# Dehydrozaluzanin C- derivative protects septic mice by alleviating over-activated inflammatory response and promoting the phagocytosis of macrophages

Ying-xiang Zou[1,2,#], Tian-nan Xiang[3,1,2,#], Li-rong Xu[1,2], Huan Zhang[1,2], Yu-he Ma[1,2], Lu Zhang[1,2], Chun-xian Zhou[1,2], Xiao Wu[1,2], Qi-lin Huang[1,2], Biao Lei[1,2], Jing-wen Mu[1,2], Xiang-yang Qin[4,*], Xin Jiang[1,2,*], Yue-juan Zheng[1,2,5,*]

[1]The Research Center for Traditional Chinese Medicine, Shanghai Institute of Infectious Diseases and Biosecurity, Shanghai University of Traditional Chinese Medicine, Shanghai, 201203, China

[2] School of Integrative Medicine, Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China

[3]Department of Chinese Medicine, Hubei College of Chinese Medicine, Jingzhou, Hubei, 434020, China

[4]Department of Chemistry, school of pharmacy, Furth Military University, Xi'an, Shaanxi, 710032, China

[5]Shanghai Key Laboratory of Health Identification and Assessment, Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China

[#]These authors contributed equally to this work.

[*] **Correspondence**: Prof. Yue-juan Zheng and Prof. Xin Jiang, School of Integrative Medicine, Shanghai University of Traditional Chinese Medicine, Shanghai, 201203, China. Tel: (+86)21-5132 2134, Fax: (+86)-21-5132 2130, Email: 13641776412@163.com (Yue-juan Zheng), and jiangxingao@163.com (Xin Jiang).

Prof. Xiang-yang Qin, Department of Chemistry, school of pharmacy, Furth Military

University, Xi'an, Shaanxi 710032, China. Tel: (+86) 029-8477 4473, Fax: (+86)

029-8477 4473, Email: qinxiangyang@fmmu.edu.cn.

**Abstract**

Host-directed therapy (HDT) is a new adjuvant strategy that interfere with host

cell factors that are required by a pathogen for replication or persistence. In this study,

we assessed the effect of dehydrozaluzanin C-derivative (DHZD), a modified

compound from dehydrozaluzanin C (DHZC), as a potential HDT agent for severe

infection. LPS-induced septic mouse model and Carbapenem resistant *Klebsiella*

*pneumoniae* (CRKP) infection mouse model was used for testing *in vivo*. RAW264.7

cells, mouse primary macrophages, and DCs were used for *in vitro* experiments.

Dexamethasone (DXM) was used as a positive control agent. DHZD ameliorated

tissue damage (lung, kidney, and liver) and excessive inflammatory response induced

by LPS or CRKP infection in mice. Also, DHZD improved the hypothermic

symptoms of acute peritonitis induced by CRKP, inhibited heat-killed CRKP

(HK-CRKP)-induced inflammatory response in macrophages, and upregulated the

proportions of phagocytic cell types in lungs. *In vitro* data suggested that DHZD

decreases LPS-stimulated expression of IL-6, TNF-α and MCP-1 via

PI3K/Akt/p70S6K signaling pathway in macrophages. Interestingly, the combined

treatment group of DXM and DHZD had a higher survival rate and lower level of

IL-6 than those of the DXM-treated group; the combination of DHZD and DXM

played a synergistic role in decreasing IL-6 secretion in sera. Moreover, the

phagocytic receptor CD36 was increased by DHZD in macrophages, which was

accompanied by increased bacterial phagocytosis in a clathrin- and actin-dependent

manner. This data suggests that DHZD may be a potential drug candidate for treating

bacterial infections.

*Keywords:* Dehydrozaluzanin C-derivative; sepsis; inflammatory response;

phagocytosis; macrophages

**Introduction**

Sepsis is a life-threatening organ dysfunction caused by a dysregulated response

to infection (1, 2). Approximately 15 to 30 million cases of sepsis are reported

annually (3, 4). Although age-standardized sepsis incidence has dropped by 37.0%

and mortality decreased by 52.8% from 1990 to 2017 (5), antimicrobial resistance still

challenges the treatment of sepsis. With the increasing use of carbapenems in

hospitals, CRKP has become a major threat to public health. CRKP accounts for about

64% of carbapenem-resistant enterobacterial infections in China (6, 7). Although a

number of studies have reported that the combined therapy with multiple active agents

may be effective for CRKP strains, the rate of clinical failure remains high (8-10).

Thus, identifying new effective treatment strategies for CRKP is crucial.

The current clinical treatment for sepsis caused by bacterial infection mainly

relies on antibiotic therapy to control the infection and the use of glucocorticoids to

inhibit the excessive inflammatory response. Glucocorticoids, such as DXM, are

effective but also accelerate specific immune cells' apoptosis. Common side effects of

long-term glucocorticoid use include osteoporosis (11) and opportunistic infections

due to its immunosuppressive effects (12). The development of new antimicrobial

drugs, especially those that target G⁻ bacteria, lags behind the emergence of antibiotic

resistance, which has become a major obstacle in the clinic.

As a major component of the cell wall of G⁻ bacteria, LPS commonly known as

endotoxin, is one of the most important pathogen-associated molecular patterns,

which is sensed by our immune system via the Toll-like receptor 4 (TLR4) (13). This

ligation activates adaptor molecules such as myeloid differentiation protein 88

(MyD88), interleukin-1 (IL-1) receptor-associated kinases (IRAKs), tumor necrosis

factor (TNF) receptor-associated factor 6 (TRAF6), transforming growth factor

(TGF)-activated kinase 1 (TAK1), and subsequent MAPKs and nuclear factor κB

(NF-κB). Yet, excessive secretion of inflammatory cytokines (such as IL-6, TNF-α

and IL-Iβ) can lead to "cytokine storm", commonly known as systemic inflammatory

response syndrome (SIRS). The pathologic progress of sepsis is usually accompanied

by SIRS. The uncontrolled inflammatory response may lead to organ dysfunction,

multiple organ failure, and, in more severe cases, even death (14).

Besides the inflammatory immune damage, the poor clearance of pathogens is

usually associated with poor prognosis during bacterial infection.

Monocytes/macrophages and dendritic cells (DCs) are the main phagocytes that

phagocytize bacteria and present their antigens to T cells (15). Neutrophil is also an

important cell type of phagocytes. Phagocytosis includes three main stages:

receptor-mediated, membrane fusion, and actin-driven (16). During this process,

phagocytic receptors, including C-type lectin receptors (CLRs), mannose receptors,

scavenger receptors (SR-A1, MARCO and CD36), complement receptors (CR3 and

CR4) and Fc receptors, have a crucial role (16-18).

HDT is a new adjuvant strategy often used to treat severe infections that can

overcome AMR. It is not mediated by killing pathogens directly; instead, it promotes

the immune clearing of pathogens through phagocytes and/or reduces tissue damage

caused by excessive inflammatory response (19).

DHZC, a sesquiterpenoid from *Magnolia officinalis*, exhibits strong

anti-inflammatory activity, but its drug toxicity is relatively high. In this study, we

assessed the effect of DHZD, a modified compound from DHZC, as a potential HDT

agent for severe infection with anti-inflammatory and immunomodulatory effects.

**Materials and Methods**

**Animals and reagents**

Six-week-old wild-type female C57BL/6J mice ($18 \pm 2$ g) were obtained from

Joint Ventures Sipper BK Experimental Animal Co. (Shanghai, China). All mice had

free access to food and water during all the experiments and were housed in a 12/12 h

light/dark cycle. The experimental procedures were approved by the Experimental

Animal Ethical Committee, Shanghai University of Traditional Chinese Medicine

(Shanghai, China) (PZSHUTCM220307018).

DHZD was modified from DHZC isolated from *Magnolia officinalis*. LPS (0111:

B4), DMSO, dexamethasone and chlorpromazine hydrochloride (CPZ) were

purchased from Sigma (St. Louis, MO, USA). Cell counting kit 8 (CCK-8) were

obtained from Meilunbio (Dalian, China). Enzyme-linked immunosorbent assay

(ELISA) kits for mouse IL-6, TNF-α, MCP-1, IL-10, IL-1β, IFN-β and

granulocyte-macrophage colony stimulating factor (mGM-CSF) were purchased from

R&D Systems (Minneapolis, MN, USA). Mouse recombinant cytokine IL-4 were

from Peprotech (USA). The detection kit of nitric oxide (NO) was obtained from

Beyotime Biotechnology (Shanghai, China). FITC anti-mouse CD11c, APC

anti-mouse CD40, APC/Fire™ 750 anti-mouse CD80, PerCP/Cyanine5.5 anti-mouse

CD86, Brilliant Violet 421™ anti-mouse I-A/I-E, FITC anti-mouse CD45, Brilliant

Violet 650 Ly6G, Brilliant Violet 421 Ly6C and PE/Cyanine7 F4/80 were purchased

from Biolegend (USA). Dexamethasone sodium phosphate injection was from

Sinopharm (China). SYBR RT-PCR Kit was purchased from Takara (Dalian, China).

PHrodo-labelled *E. coli* was purchased from ThermoFisher (USA). Cytochalasin D

was obtained from Abcam (USA). Filipin III was from MCE (USA). Levofloxacin

(LVX) and meropenem (MEM) were obtained from the National Institute for Food

and Drug Control of China (Beijing, China) CRKP (HS11286) was kindly provided

by Dr. Yi-jian Chen from Huashan Hospital affiliated to Fudan University.

**Cell culture**

Mouse macrophage-like cell line RAW264.7 were purchased from ATCC

(Manassas, VA, USA) and cultured as previously described (20).

Thioglycolate-elicited mouse primary peritoneal macrophages were isolated

from female C57BL/6J mice at 6-8 weeks of age and cultured in DMEM (GE, USA)

with 10 % FBS (Biowest, France) and 1 % Penicillin-Streptomycin, as previously

described (21).

Bone marrow-derived macrophages (BMDMs) and bone marrow-derived

dendritic cells (BMDCs) were both obtained from C57BL/6J mice (4 weeks old). The

BMDMs were cultured in DMEM containing 10 ng/mL M-CSF (Peprotech, USA)

and 10 % FBS. The BMDCs were differentiated in RPMI 1640 (GE, USA) medium

containing 10 % FBS, 50 ng/mL mGM-CSF (R&D Systems, USA) and 4 ng/mL

mIL-4 (Peprotech, USA).

**Cell viability assay**

RAW264.7 cells ($1\times10^5$ cells/well) were cultured in 96-well plates overnight and

treated with different concentrations of DHZD (0-15 μM) in 12, 24, 48 and 72 h. The

cytotoxicity of DHZD was evaluated by CCK-8 counting kit according to the

manufacturer's instructions (Meilunbio, Dalian, China).

**Preparation of HK-CRKP**

Monoclonal colonies were picked from the CRKP bacterial culture plates and

cultured in LB medium for 12 h at 37°C on a constant temperature shaker. After

heating 1 mL of the bacterial solution in a 65°C water bath for 45 min, the solution

was centrifuged (4400 rpm×15 min), resuspended in sterile PBS, centrifuged again

(4400 rpm×15 min), and resuspended in DMEM for further use (MOI=10).

**Enzyme-linked immunosorbent assay**

RAW264.7 ($2\times10^5$ cells/well) were plated as indicated and administrated with

DHZD and/or LPS (100 ng/mL) for 6 h and 18 h. The levels of cytokines (IL-6,

TNF-α, MCP-1, IFN-β, IL-1β and IL-10) in the cell culture supernatants were

detected by ELISA (R&D Systems, USA).

Mouse primary macrophages were plated ($3.5 \times 10^5$ cells/500 μL) in 24-well

plates at the day before stimulation. Cell culture supernatants were collected at 6 h

and 18 h after co-stimulation of different concentrations of DHZD and LPS (100

ng/mL), and the concentrations of IL-6, TNF-α, MCP-1, IL-1β, IFN-β and IL-10 were

measured by ELISA.

BMDCs were re-seeded on day 5 of differentiation at $3.5 \times 10^5$ cells/500 μL in

24-well plates and stimulated by DHZD (0-15 μM) with or without LPS (100 ng/mL)

for 6 h or 18 h. The secretion of IL-6, TNF-α and IL-12p70 in cell culture

supernatants were detected by ELISA.

Mouse primary macrophages were plated ($3.5 \times 10^5$ cells/500 μL) in 24-well

plates one day before and were stimulated by different concentrations of DHZD and

HK-CRKP (MOI=10). After 12 h or 24 h, cell culture supernatants were collected and

the concentrations of IL-6, TNF-α, MCP-1, IL-1β, IL-10 and MIP-2 were measured

by ELISA.

BMDMs were re-seeded on day 5 of differentiation ($3.5 \times 10^5$ cells/well) in

24-well plates and then stimulated by LPS (100 ng/mL) and DHZD for 6 h. The

expression of cytokines (IL-6, TNF-α and IL-10) was detected by ELISA.

**Assessment of NO level**

Mouse primary macrophages were plated ($3.5 \times 10^5$ cells/500 µL) in 24-well plates at the day before stimulation. Cell culture supernatants were collected at 6 h and 18 h after co-stimulation of different concentrations of DHZD and LPS (100 ng/mL). The secretion of NO was detected by corresponding NO detection kit (S0021) which was purchased from Beyotime Biotechnology (Shanghai, China).

**Detection of MHC II and co-stimulatory molecules on BMDCs**

Bone marrow stem cells were derived and differentiated into BMDCs. After 5 days, BMDCs were seeded in 12-well plates overnight. LPS (100 ng/mL) and DHZD (15 µM) were added into culture medium of BMDCs for another 24 h. Then BMDCs were harvested and labeled with FITC anti-mouse CD11c (Biolegend, 117305, USA), APC anti-mouse CD40 (Biolegend, 124611, USA), APC/Fire™ 750 anti-mouse CD80 (Biolegend, 104739, USA), PerCP/Cyanine5.5 anti-mouse CD86 (Biolegend, 105027, USA) and Brilliant Violet 421™ anti-mouse I-A/I-E (Biolegend, 107631, USA). And then cells were analyzed by Flow cytometry (Beckman, USA), and data were analyzed using FlowJo software.

**The establishment of septic mice by lethal dose of LPS challenge**

Six groups were included (n=10): LPS group, LPS+DXM group (5 mg/kg), LPS+DHZD group (10 mg/kg), LPS+DHZD group (20 mg/kg), LPS+DHZD group (40 mg/kg) and LPS+MEM (5 mg/kg) +DHZD group (10 mg/kg). The septic mouse model was established by intraperitoneal injection of LPS (12.5 mg/kg, Sigma) as described (22, 23). At the same time, DHZD and DXM were injected intraperitoneally.

194 Survival status of different groups was recorded every 2-3 h in the following

195 continuous 180 h (24).

**LPS-induced acute peritonitis**

Acute peritonitis mouse model (n=9) was carried out by intraperitoneal injection

of LPS (10 mg/kg). DXM (5 mg/kg) and/or DHZD (10 or 40 mg/kg) were

administered at the same time. After 12 h, all the mice in different groups were

sacrificed. Meanwhile, blood samples were collected and allowed to clot for 3 h

followed by centrifugation at 3500 rpm for 25 min (4 □) to prepare serum samples.

Serum levels of cytokines (IL-6, TNF-α, MCP-1 and IL-10) were measured by

corresponding ELISA kits (R&D Systems, USA) according to the manufacturer's

instructions.

Total RNA was extracted from lung tissues and reversely transcribed to cDNA.

Real-time PCR was performed with SYBR RT-PCR kit (Takara, Dalian, China). The

mRNA levels were normalized against β-Actin and presented as $2^{-\Delta\Delta Ct}$.

The lung tissues were snipped and digested with collagenase (Sigma Aldrich)

and DNase I (Sigma Aldrich) in gentleMACS$^{TM}$ (Miltenyi, Germany) at 80 rpm

(37 °C) for 30 min. After digestion, the samples were filtered through a 40 mm filter

using a syringe piston, and centrifuged at 1000 rpm (4 °C) for 5 min. RBC lysis buffer

(BD Biosciences, USA) was added to cell pellets at room temperature for 5 min, and

then PBS (8 mL) was added and centrifuged at 1400 rpm (4 °C) for 5 min. Cells were

counted and then incubated at 4 °C for 25 min with FITC anti-mouse CD45

(Biolegend, 103107, USA), Brilliant Violet 650 Ly6G (Biolegend, 127641, USA),

Brilliant Violet 421 Ly6C (Biolegend, 128031, USA) and PE/Cyanine7 F4/80

(Biolegend, 123113, USA) antibodies. Samples were run on CytoFlex LX (Beckman,

USA). Data were analyzed using FlowJo software.

Lung, liver and kidney tissues were collected to be fixed with 4% formaldehyde

and paraffin-embedded. All of the above tissues were sliced and stained with

hematoxylin and eosin (H&E). The histopathological changes were observed using a

PANNORAMIC MIDI II microscope (3DHistech, Hungary). Pulmonary edema,

alveolar infiltration, and pulmonary vasculitis were scored on a 12-point pathological

scale (25).

**RNA extraction and real-time PCR**

RAW264.7 cells were seeded ($2 \times 10^5$/well) in 24-well plates and incubated

overnight. Cells were treated with heat killed-Methicillin resistant *Staphylococcus*

*aureus* (HK-MRSA) (MOI=10) separately along with DHZD (15 μM) for 12 and 18 h.

The cells were harvested for total RNA extraction and reversely transcribed to cDNA

for further qPCR detection.

Real-time PCR was performed with SYBR RT-PCR kit (Takara, Dalian, China).

The mRNA levels were normalized against β-Actin and presented as $2^{-\Delta\Delta Ct}$. The

primer sequences for qRT-PCR amplification were as the following:

| mβ-Actin | mβ-Actin-F | 5'-AGTGTGACGTTGACATCCGT-3' |
| | mβ-Actin-R | 5'-GCAGCTCAGTAACAGTCCGC-3' |
| mIL-6 | mIL-6F | 5'-TAGTCCTTCCTACCCCAATTTCC-3' |
| | mIL-6R | 5'-TTGGTCCTTAGCCACTCCTTC-3' |

| | | |
|---|---|---|
| mTNF-α | mTNF-αF | 5'-AAGCCTGTAGCCCACGTCGTA-3' |
| | mTNF-αR | 5'-GGCACCACTAGTTGGTTGTCTTTG-3' |
| mMCP-1 | mMCP-1F | 5'- GCATCCACGTGTTGGCTCA -3' |
| | mMCP-1R | 5'- CTCCAGCCTACTCATTGGGATCA -3' |
| mMIP-2 | mMIP-2-F | 5'-CACTCTCAAGGGCGGTCAA-3' |
| | mMIP-2-R | 5'-AGGCACATCAGGTACGATCCA-3' |
| mIL-10 | mIL-10F | 5'-GCTCTTACTGACTGGCATGAG-3' |
| | mIL-10R | 5'-CGCAGCTCTAGGAGCATGTG-3' |
| mCD206 | mCD206F | 5'-CTCTGTTCAGCTATTGGACGC-3' |
| | mCD206R | 5'-TGGCACTCCCAAACATAATTTGA-3' |
| mDectin-1 | mDectin-1F | 5'-AGGCATCCCAAACTACAGGAG-3' |
| | mDectin-1R | 5'-GGAGCAGTGTCTCTTACTTCCAT-3' |
| mCR3 | mCR3F | 5'- CCATGACCTTCCAAGAGAATGC -3' |
| | mCR3R | 5'- ACCGGCTTGTGCTGTAGTC -3' |
| mCR4 | mCR4F | 5'- CTGGATAGCCTTTCTTCTGCTG -3' |
| | mCR4R | 5'- GCACACTGTGTCCGAACTCA -3' |
| mFcγR2a | mFcγR2aF | 5'-TGGACAGCCGTGCTAAATCTT-3' |
| | mFcγR2aR | 5'-GGTCCCTTCGCATGTCAGTG-3' |

**Detection of the phagocytosis of macrophages**

The phagocytosis of pHrodo labelled *E. coli* (ThermoFisher, California, USA)

was detected in RAW264.7 cells by Flow cytometry. RAW264.7 cells were plated at

$1\times10^{5}$ cells/100 µL in 96-well plates and incubated overnight. Then cells were primed

with LPS (100 ng/mL) along with DHZD (15 μM). After 24 h, macrophages were

incubated with pHrodo-labelled *E. coli* (10 μL/well) protected from light for 1 h.

Cells were washed with sterile PBS for 3 times to get rid of the bacteria outside of the

cells. Then the phagocytosis of pHrodo-labelled *E. coli* was quantified by FACS.

RAW264.7 cells were pretreated by 3 μM of Filipin III (the inhibitor of caveolin,

MCE, USA), 10 μM of chlorpromazine hydrochloride (the inhibitor of clathrin,

Sigma, USA) or 10 μM of cytochalasin D (the inhibitor of actin, Abcam, USA) for 30

min, and then pHrodo-labelled *E. coli* were added to incubation system for another 1

h. The phagocytosis of *E. coli* was investigated by BD Accuri$^{TM}$ C6 Flow cytometry

(BD, USA), and data were analyzed by FlowJo software.

**Western blot**

The phosphorylation of important signaling proteins from RAW264.7 cells and

their corresponding total proteins were determined by Western blot as previously

described (26). RAW264.7 cells (1 × 10$^6$ cells/well) were seeded in 6-well plates

overnight and stimulated with LPS (100 ng/mL) with or without DHZD (15 μM) for a

certain period of time. All the antibodies were bought from Cell Signaling Technology

(Beverly, MA, USA).

**CRKP-induced septic mouse model**

The septic mouse model was established by intraperitoneal injection of CRKP

(6×10$^7$ CFU/mouse) in 0.4 mL of PBS. The experiments were carried out in the

following groups: CRKP group, CRKP+MEM (5 mg/kg) group, CRKP+LVX (75

mg/kg) group, CRKP+DHZD group (20 mg/kg), CRKP+DHZD group (40 mg/kg)

and the combined treatment CRKP+DHZD (20 mg/kg) +MEM (5 mg/kg) group. All

medicines were administered simultaneously. The body temperature of mice was

recorded by infrared body thermometer (HJKIR, Wuhan) before and 9 hours after

injection of CRKP.

After 12 h, mice were sacrificed, and lungs and livers were collected to be fixed

with 4% formaldehyde and embedded in paraffin. All of the above tissues were sliced

and stained with H&E. The histopathological changes were observed using a

PANNORAMIC MIDI II microscope (3DHistech, Hungary). Pulmonary edema,

alveolar infiltration, and pulmonary vasculitis were scored on a 12-point pathological

scale (25).

**Statistical analysis**

Statistical analysis was performed using GraphPad Prism 9. Data were presented

as means ± standard deviation (SD). Student's *t*-test analyses was used to evaluate the

comparisons between two groups. Comparisons multiple groups were evaluated by

one-way ANOVA followed by Tukey's post hoc comparisons, respectively. *P* values

were considered as *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; ****, $P<0.0001$.

**Results**

**DHZD inhibits the secretion of LPS-induced cytokines from RAW264.7 cells**

The structure of DHZD is shown in Figure 1A. DHZD was synthesized

according to the previously reported method (27). Spectroscopic data were in

accordance with reported values (Supplementary Fig.1) (27).

CCK-8 kit was used to determine whether DHZD affected cell viability. DHZD

(0-15 μM) did not affect the proliferation of RAW264.7 cells within 72 h, but

inhibited LPS-induced secretion of IL-6, TNF-α, MCP-1, IFN-β, and IL-10 in a

dose-dependent manner (Fig. 1B-H).

**DHZD decreases the secretion of LPS-induced expression of IL-6, TNF-α,**

**MCP-1, IL-1β, IFN-β and IL-10 in mouse primary macrophages**

Next, we assessed the effect of DHZD on the secretion of LPS-induced

expression of inflammatory mediators in mouse primary macrophages and BMDMs.

Compared to macrophage-like cell lines, primary macrophages mimic the

physiological state of cells *in vivo* and are expected to maintain their *in vivo* functions

under optimal conditions for a short period (28). DHZD inhibited the secretion of

LPS-induced pro-inflammatory mediators IL-6, TNF-α, IL-1β, IFN-β, NO, MCP-1

and anti-inflammatory cytokine IL-10 in mouse primary macrophages (Figure 2A-G)

and decreased the secretion of IL-6, TNF-α and IL-10 in BMDMs stimulated with

LPS (Supplementary Fig. 2A-C).

**DHZD decreases LPS-induced secretion of inflammatory cytokines in BMDCs,**

**and downregulated the expression of MHC II and co-stimulatory molecules on**

**the surface of BMDCs**

DCs are the most important and powerful antigen-presenting cells that activate

naive T cells, bridging innate and adaptive immunity. DCs participate in the

phagocytosis and digestion of bacteria and present their antigens to evoke T-cell

immunity (29). During the transition of DCs from immature to mature status,

co-stimulatory molecules are upregulated, downregulating the phagocytosis and

promoting the antigen-presenting ability. In order to investigate whether DHZD has a

regulatory effect on the antigen presentation function of DCs, thereby affecting the

activation of T cells, flow cytometry was used to detect the expression of

costimulatory molecules (CD80, CD86, and CD40) and MHC Class II molecule (Iab)

in cells treated with or without LPS. Briefly, DHZD inhibited the expression of these

molecules compared to control cells (Figure 3D-G).

DCs secret inflammatory cytokines (IL-6, TNF-α, IL-12 and a variety of

chemokines), contributing to the formation of cytokine storm together with activated

T cells during the pathological progress of sepsis (30). This study found that DHZD

reduces the secretion of IL-6, TNF-α, and IL-12p70 (Figure 3A-C) DCs stimulated

with LPS.

**DHZD protects septic mice by inhibiting the over-activated inflammatory**

**response**

To determine the protective role of DHZD against LPS-induced (12.5 mg/kg)

septic shock, a survival analysis was carried out. DXM, a glucocorticoid used in the

clinical treatment of bacterial infection, was used as a positive control drug.

All mice in the model group died within 40 h. The mice in the septic model

group had low body temperature, shivering, standing hair and decreased activity. In

the DXM treatment group, 60 % of mice survived (10 % of mice in the low-dose

DHZD treatment group and 30 % in the medium- or high-dose DHZD treatment

groups). All mice survived in the low-dose DHZD and DXM treatment groups (Figure

4A). Compared with the model group, the mice in the DXM group, the high-dose

DHZD-treated group, and the DXM+DHZD combination group exhibited faster

recovery of body temperature, smoother hair, and were more active. The combined

treatment group (DXM+DHZD) showed the best protective effect in terms of clinical

manifestation.

The lung, liver and kidney tissue structures in the control group were intact. On

the contrary, obvious infiltration of inflammatory cells, a large amount of fluid

exudation in the alveoli, and lung interstitial thickening were seen in mice injected

with LPS. Also, the liver tissue showed obvious inflammatory cell infiltration,

swelling and necrosis of hepatocytes, fragmentation and dissolution of some nuclei,

and destruction of the hepatic plate structure, while renal tissue showed obvious

inflammatory cell infiltration, degeneration and necrosis of renal tubular epithelial

cells, and narrowing of renal cysts. DHZD improved pulmonary edema, alveolar

infiltration, and lung vasculitis caused by LPS. DXM improved lung pathological

damage (Figure 4B-C). The destruction of lung, liver and kidney tissue structure was

improved after DXM, DHZD or DXM+DHZD treatment. There was no significant

difference among different treated groups (DXM vs DHZD, $p$=0.8856, DXM vs

DXM+DHZD, $p$=0.1168, DHZD vs DXM+DHZD, $p$=0.4112) (Figure 4B).

Multiple organ failure caused by a "cytokine storm" during infection is the

leading cause of death in septic patients (31). In LPS-challenged mice, DHZD and

DXM decreased the secretion of IL-6, TNF-α, MCP-1 and IL-10. Compared with

DXM treatment, the combination treatment of DHZD+DXM could significantly

inhibit the secretion of these cytokines (Figure 4D-G).

Sepsis induced by bacterial infection can cause systemic organ dysfunction, and

lungs are often the most easily affected important organs. Therefore, sepsis is often

accompanied by acute lung injury (ALI) (32) or acute respiratory distress syndrome

(ARDS) (33), which is one of the most important prognostic factors of sepsis death. In

this study, we detected the expression of lung inflammatory factors by qPCR. DHZD

and DHZD+DXM could significantly inhibit the production of proinflammatory

cytokines IL-6, chemokines MCP-1 and MIP-2 in the lungs of mice (Supplementary

Fig. 3A-C).

**DHZD increases the numbers of lung neutrophils, monocytes, and macrophages**

**induced by LPS**

In most cases, locally recruited natural immune cells (such as neutrophils and

monocytes/macrophages) can rapidly phagocyte bacteria, control the bacterial load in

tissues, and reduce the occurrence of severe infection. However, when severe

infection occurs, many immune cells participate in the secretion of inflammatory

cytokines, forming cytokine storms, which can lead to organ damage, sepsis and even

death. Thus, in the early stage of infection, rapid phagocytosis and elimination of

bacteria, and subsequently control of the immune response of the body within

reasonable limitation, are the key aspects for treatment (34, 35).

In this study, the infiltration of neutrophils and monocytes/macrophages in the

lungs was measured by flow cytometry. As shown in Figure 5, 12 h after LPS

injection, the ratio of neutrophils (Figure 5A-C) and monocytes/macrophages (Figure

5D-H) increased in the DXM-treated group, DHZD-treated group, and DHZD+DXM

group compared to LPS challenged group, suggesting an increase in the number of

innate immune cells with the ability to phagocyte bacteria.

DHZD may have rapid phagocytic role in the early stage of bacterial infection,

reducing the bacterial load and the severity of infection in tissues, and thus exerting a

potential protective role. From our results, DXM and DHZD have similar effects in

increasing the proportions and numbers of neutrophils and monocytes. In addition,

DXM also has a broad effect of immunosuppression, which might be accompanied by

potential adverse effects (36).

**DHZD enhances the phagocytosis of pHrodo-labelled *E. coli* in macrophages in a**

**clathrin-dependent and actin-polymerized manner via CD36**

PHrodo-labeled *E. coli* is a model bacterial that emit fluorescence only in

phagocytic cells because of the acidic environment (37, 38). To clarify the regulatory

effect of DHZD on the phagocytotic ability, RAW264.7 cells were stimulated by LPS

and treated with DHZD (15 µM) for 24 h. PHrodo-labeled *E. coli* was added and

incubated for 1 h in dark. The amount of engulfed *E. coli* in the cytoplasm was

measured by flow cytometry. DHZD significantly increased the phagocytosis of

pHrodo-labeled *E. coli* in RAW264.7 (Figure 6A-C), suggesting that DHZD could

promote the ability of bacterial phagocytosis of macrophages.

The initiation of phagocytosis depends on the recognition and binding of host

cell receptors to the corresponding ligands on the surface of the pathogens. Innate

immune cells such as macrophages, neutrophils and dendritic cells can phagocyte

pathogens through caveolin-mediated endocytosis, clathrin-mediated endocytosis,

pinocytosis and other ways (39, 40). All the above-mentioned phagocytotic

mechanisms are actin-dependent. Filipin III, CPZ and cytochalasin D are inhibitors of

caveolin, clathrin and actin, respectively. Our data show that Filipin III has no effect

on DHZD-promoted phagocytosis of bacteria in RAW264.7 cells, while CPZ could

partially inhibit DHZD-promoted phagocytosis. Cytochalasin D could prevent actin

polymerization and completely prevent enhanced phagocytosis by DHZD (Figure

6D-F).

Phagocytic receptors usually mediate bacterial phagocytosis. The expression of

phagocytic receptors CD36, FcγR2a, CD206, Dectin-1, CR3 and CR4 were detected

by qRT-PCR on the surface of macrophages. From the expression screen, DHZD

could promote the expression of CD36 (Figure 6G-H), at least partially accounting for

the increased phagocytosis ability.

**DHZD inhibits LPS-stimulated activation of PI3K/Akt/p70S6K pathway**

LPS, as a ligand of TLR4, can activate extracellular signal-regulated kinase 1/2

(ERK1/2), c-Jun N-terminal kinase (JNK), and p38 MAPK pathways, NF-κB pathway

and phosphoinositide 3-kinase (PI3K)/Akt signaling pathway, leading to the

production of cytokines (41). Inhibition of PI3K/Akt activation reduces LPS-induced

release of NO, MCP-1, IL-6, and TNF-α (42). Our results demonstrated that DHZD

can effectively reduce LPS-induced inflammatory response. DHZD inhibited

LPS-induced phosphorylation of Akt at Ser473 and Thr308 and subsequent p70S6K at

Thr389, but did not affect the activation of NF-κB, ERK and p38 MAPK signaling

pathways (Figure 7A-E).

**DHZD reverses hypothermia, tissue damage, and over-activated inflammation**

**caused by CRKP**

LPS is the main component of the cell wall of G⁻ bacteria. CRKP is one of the

most troublesome G⁻ bacteria causing severe infection and sepsis in the emergency

department or intensive care unit (ICU). Fever is usually one of the most important

symptoms of infection (43). There are variable thermoregulatory responses in sepsis,

including fever and hypothermia (44). According to the analysis of clinical data, the

mortality rate of septic patients with hypothermia is much higher than in those with

fever (45).

In order to clarify the protective effects of DHZD on CRKP-infected mice, a

septic mouse model was established by the intraperitoneal injection of CRKP, which

is sensitive to LVX, used as a positive control antibiotic. Infrared thermal imaging

thermometers were also used to monitor the body temperature of mice before and

after the CRKP challenge. Compared with the model group, average body temperature

was normalized by 1.3 □ and 3.1 □ in the DHZD (20) and DHZD (40) groups,

respectively. The results suggested that DHZD had an impressive protective effect on

the recovery of body temperature (Figure 8A, B).

Also, compared with the PBS group, the lung tissue of mice in the CRKP group

showed obvious infiltration of inflammatory cells, a large amount of fluid exudation

in the alveoli, and pulmonary interstitial thickening. Inflammatory cell infiltration,

swelling and necrosis of hepatocytes, fragmentation and lysis of some nuclei, and

destruction of hepatic plate structure were observed in the liver. DHZD ameliorated

lung and liver tissue damage. A pathological score of lung tissue, as measured by

pulmonary edema, alveolar infiltration, and pulmonary vasculitis, showed a protective

effect of DHZD administration against pathological damage to the tissue (Figure

8C-D).

*In vitro* stimulation of primary peritoneal macrophages of mice with HK-CRKP

showed that DHZD could reduce the secretion of IL-6, TNF-α, MCP-1, IL-1β and

IL-10 in a dose-dependent manner but had no obvious influence on MIP-2 secretion

(Figure 8E-J).

**Discussion**

The World Health Organization has listed AMR among top 10 threats for global

health. After entering the bloodstream, invading bacteria are recognized by

pathogen-associated molecular patterns (PAMPs) via pattern recognition receptors

(PRRs). In most cases, the innate immune response eliminated the invading bacteria.

Yet, when the bacteria prevail in the host, the immune response becomes unbalanced

(46, 47), further triggering the overwhelming production of cytokines, such as TNF-α,

IL-1β, IL-6, etc. (48). The excessive inflammatory response promotes the activation

of the coagulation system and the formation of microthrombosis in blood vessels,

leading to disseminated intravascular coagulation (DIC) (49, 50).

"Cytokine storm" is the main cause of death in septic patients with multiple

organ failure. The levels of inflammatory cytokines IL-1β, IL-6, IL-12 and IL-17 are

usually high in septic patients (31, 51). In a study of community-acquired pneumonia,

the mean serum concentration of IL-6 from severe septic patients was 54.1 pg/mL,

compared with 36.5 pg /mL from patients who did not develop severe sepsis (52).

Similarly, the serum levels of IL-1β in deceased septic patients were about 2-10 times

higher than those in surviving septic patients. These clinical data suggest that high

concentrations of IL-6 or IL-1β may be closely related to poor prognosis (53). Animal

studies have suggested that administering anti-IL-6 antibodies reduces the expression

of C5a and improves survival in septic mice (54). TNF-α is not only involved in

cytokine storm but is also responsible for the induction of hypothermia and high

mortality in sepsis (55). In a study of *E. coli*-induced sepsis in baboons, treatment

with anti-TNF antibody at 2 h before infection protected vital organ damage and

reduced mortality (56). It can be concluded from the above clinical experience that the

downregulation of the excessive cytokine expression can alleviate tissue damage,

reduce the conversion rate from mild to severe diseases, and improve the survival

status of sepsis. In our study, DHZD could inhibit the over-activated inflammatory

response (IL-6, TNF-α, IL-1β, IL-12, IFN-β and IL-10) in RAW264.7 cells, mouse

primary peritoneal macrophages, and BMDMs after the stimulation of bacteria or

bacterial mimic. In the early stage of sepsis, DCs can produce inflammatory factors

(such as IL-6, TNF-α, IL-12 and chemokines) after bacterial phagocytosis (30).

Elevated IL-12 then induces the differentiation of naive T cells into helper T cells

(Th1) and the activation of NK cells, contributing to the formation of a cytokine storm

(57). We demonstrated that DHZD downregulates the production of IL-6, TNF-α and

IL12p70 in DCs, suggesting that it may also downregulate cytokine storm via

impairing subsequent T cell over-activation.

In the LPS-induced septic mouse model, DHZD showed an impressive

anti-inflammatory role and alleviated lung, liver and kidney damage, improving the

survival rate. Both of the examined concentrations of DHZD showed protective

effects *in vivo*. The survival rate of LPS+DHZD+DXM group was 100%, which was

better than that of LPS+DXM group (60%), suggesting that the combination of

DHZD and DXM may have a potential synergistic protective effect to treat sepsis.

TNF-α, interleukin, and interferonγcan induce hypothermia by regulating the

thermotaxic center of the brain (58, 59). During infection, the activity of animals is

reduced, which can also lead to the appearance of hypothermia (60). During sepsis,

central nervous system dysfunction (88% vs 60%), elevated serum bilirubin

concentration (35% vs. 15%), prolonged prothrombin time (50% vs. 23%), shock (94%

vs 61%), nonrecovery from shock (66% vs. 26%), and death (62% vs. 26%), were

found in patients with hypothermia compared to patients in the fever group (61). In

this study, DHZD protected the loss of body temperature in CRKP-induced septic

mouse model (-5.7 □ vs -2.6 □) compared to those of CRKP group mice, suggesting

that DHZD-treated infective mice have a better prognosis.

The pathogen load is usually related to the severity of infectious diseases.

Macrophage is a ubiquitous phagocytic cell type of innate immune system involved in

defensive inflammatory response and pathogen clearance during bacterial infection

(62, 63). Besides macrophages, neutrophils and DCs are also important phagocytic

cells in the body, which can phagocytize bacteria and decrease the bacterial load. Our

study used macrophages as a cell model to investigate whether DHZD can influence

cellular phagocytosis. DHZD promoted phagocytosis of bacteria in a

clathrin-and-actin-dependent manner.

CD36, a class B scavenger receptor family member, is expressed in monocytes,

macrophages, endothelial cells, and adipocytes (52). The upregulation of CD36

expression by DHZD might at least partially explain the enhanced phagocytosis of

macrophages. Not only the expression of phagocytic receptors of phagocytes but also

the proportions and numbers of neutrophils and monocytes/macrophages in lungs

were increased by DHZD after the LPS challenge, suggesting that DHZD might

contribute to a quick control of bacterial load at an early stage of infection through

phagocytosis. Accordingly, DHZD might be protective during bacterial infection and

decrease organ damage.

As a member of innate immune cells, DCs have an important role in the line of

defense against pathogens. Immature DCs have a strong phagocytic ability. As they

mature, the phagocytic ability of DC is decreased, and the expression of MHC II

molecules and co-stimulatory molecules (CD86, CD80, and CD40) are increased,

accompanied by the increased antigen presenting ability to evoke T cell activation (29,

64, 65). In our study, the downregulated expression of co-stimulatory molecules

(CD80, CD86 and CD40) and MHC II by DHZD kept DCs in an immature status,

helping to maintain their strong phagocytic ability and prevent the over-activation of

T cells.

The PI3K/Akt signaling pathway is an important regulator of cytokine

production after TLR4 ligation with LPS during G$^-$ bacterial infection (66). Previous

studies have shown that the inhibition of PI3K/Akt activation in macrophages or DCs

reduces the secretion of cytokines (IL-6, TNF-$\alpha$, IL-12, IL-10 and IL-1$\beta$) (42, 67).

Though IL-10 has a well-known role in anti-inflammation, extremely high levels of

IL-10 are harmful to patients. For example, previous studies reported 3.3 times higher

serum IL-10 levels in septic patients who died compared to survivors (68). In this

study, DHZD inhibited the activation of the PI3K/Akt/p70S6K signaling pathway but

did not influence the activation of ERK and p38 MAPK signaling pathways,

accounting for its anti-inflammatory role.

In conclusion, DHZD exerts an obvious anti-inflammatory effect *in vitro* and *in*

*vivo* by targeting the host immune response and thus may be a potential therapeutic

drug for treating sepsis caused by bacterial infection and even drug-resistant bacterial

infection. Further experiments are needed to assess whether the phagocytic ability of

bacteria besides *E. coli* could also be increased by DHZD. Interestingly, the

combination of DHZD and DXM is expected to improve the survival rate of septic

mice, hinting that DHZD might be a potential adjuvant drug candidate for the

treatment of sepsis.

**Acknowledgments**

This work was supported by grants from National Key Research and

Development Program of China (2021YFE0200900), National Natural Science

Foundation of China (82073901, 82073726), Shanghai Municipal Science and

Technology Major Project (ZD2021CY001), the Key Research and Development

Program of Shaanxi Province (2020SF-226), three-year Action Plan for Shanghai

TCM Development and Inheritance Program [ZY (2021-2023)-0401 and ZY

(2021-2023)-0103]. Traditional Chinese Medicine Research Project of Shanghai

Municipal Health Commission (2022QN043). We are thankful for Dr. Yi-jian Chen

from Huashan Hospital affiliated to Fudan University to provide

Carbapenem-resistant *Klebsiella pneumoniae* (CRKP, HS11286). Thank Ali

Mollahassani for language polishing of this paper.

**Author contribution**

Yue-juan Zheng designed the experiments. Xiang-yang Qin modified and

provided DHZD. Ying-xiang Zou, Tian-nan Xiang, Huan Zhang, Yu-he Ma, Xiao Wu,

Qi-lin Huang, Biao Lei and Jing-wen Mu performed the laboratory assays. Ying-xiang

Zou, Tian-nan Xiang, Huan Zhang and Yu-he Ma analyzed the data. Yue-juan Zheng

and Ying-xiang Zou wrote the manuscript. Tian-nan Xiang and Xiang-yang Qin

contributed to manuscript revision. All authors read and approved the manuscript.

**Reference**

1. Mu A, Klare WP, Baines SL, Ignatius Pang CN, Guérillot R, Harbison-Price N, Keller N,

Wilksch J, Nhu NTK, Phan MD, Keller B, Nijagal B, Tull D, Dayalan S, Chua HHC,

Skoneczny D, Koval J, Hachani A, Shah AD, Neha N, Jadhav S, Partridge SR, Cork

AJ, Peters K, Bertolla O, Brouwer S, Hancock SJ, Álvarez-Fraga L, De Oliveira DMP,

Forde B, Dale A, Mujchariyakul W, Walsh CJ, Monk I, Fitzgerald A, Lum M,

Correa-Ospina C, Roy Chowdhury P, Parton RG, De Voss J, Beckett J, Monty F,

McKinnon J, Song X, Stephen JR, Everest M, Bellgard MI, Tinning M, Leeming M,

Hocking D, et al. 2023. Integrative omics identifies conserved and pathogen-specific

responses of sepsis-causing bacteria. Nat Commun 14:1530.

2. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M,

Bellomo R, Bernard GR, Chiche JD, Coopersmith CM, Hotchkiss RS, Levy MM,

Marshall JC, Martin GS, Opal SM, Rubenfeld GD, van der Poll T, Vincent JL, Angus

DC. 2016. The Third International Consensus Definitions for Sepsis and Septic Shock

(Sepsis-3). Jama 315:801-10.

3. Fleischmann C, Scherag A, Adhikari NK, Hartog CS, Tsaganos T, Schlattmann P,

Angus DC, Reinhart K. 2016. Assessment of Global Incidence and Mortality of

Hospital-treated Sepsis. Current Estimates and Limitations. Am J Respir Crit Care

Med 193:259-72.

4.    Adhikari NK, Fowler RA, Bhagwanjee S, Rubenfeld GD. 2010. Critical care and the

global burden of critical illness in adults. Lancet 376:1339-46.

5.    Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, Colombara

DV, Ikuta KS, Kissoon N, Finfer S, Fleischmann-Struzek C, Machado FR, Reinhart KK,

Rowan K, Seymour CW, Watson RS, West TE, Marinho F, Hay SI, Lozano R, Lopez

AD, Angus DC, Murray CJL, Naghavi M. 2020. Global, regional, and national sepsis

incidence and mortality, 1990-2017: analysis for the Global Burden of Disease Study.

Lancet 395:200-211.

6.    Yang Y, Yang Y, Chen G, Lin M, Chen Y, He R, Galvão KN, El-Gawad El-Sayed

Ahmed MA, Roberts AP, Wu Y, Zhong LL, Liang X, Qin M, Ding X, Deng W, Huang S,

Li HY, Dai M, Chen DQ, Zhang L, Liao K, Xia Y, Tian GB. 2021. Molecular

characterization of carbapenem-resistant and virulent plasmids in Klebsiella

pneumoniae from patients with bloodstream infections in China. Emerg Microbes

Infect 10:700-709.

7.    Zhang R, Liu L, Zhou H, Chan EW, Li J, Fang Y, Li Y, Liao K, Chen S. 2017.

Nationwide Surveillance of Clinical Carbapenem-resistant Enterobacteriaceae (CRE)

Strains in China. EBioMedicine 19:98-106.

8.    Tumbarello M, Viale P, Bassetti M, De Rosa FG, Spanu T, Viscoli C. 2015. Infections

caused by KPC-producing Klebsiella pneumoniae: differences in therapy and mortality

in a multicentre study--authors' response. J Antimicrob Chemother 70:2922.

9.    Gutiérrez-Gutiérrez B, Salamanca E, de Cueto M, Hsueh PR, Viale P, Paño-Pardo JR,

Venditti M, Tumbarello M, Daikos G, Cantón R, Doi Y, Tuon FF, Karaiskos I,

Pérez-Nadales E, Schwaber MJ, Azap Ö K, Souli M, Roilides E, Pournaras S, Akova

M, Pérez F, Bermejo J, Oliver A, Almela M, Lowman W, Almirante B, Bonomo RA,

Carmeli Y, Paterson DL, Pascual A, Rodríguez-Baño J. 2017. Effect of appropriate

combination therapy on mortality of patients with bloodstream infections due to

carbapenemase-producing Enterobacteriaceae (INCREMENT): a retrospective cohort

study. Lancet Infect Dis 17:726-734.

10.    Tzouvelekis LS, Markogiannakis A, Piperaki E, Souli M, Daikos GL. 2014. Treating

infections caused by carbapenemase-producing Enterobacteriaceae. Clin Microbiol

Infect 20:862-72.

11.    Compston J. 2018. Glucocorticoid-induced osteoporosis: an update. Endocrine

61:7-16.

12.    Cutolo M, Seriolo B, Pizzorni C, Secchi ME, Soldano S, Paolino S, Montagna P, Sulli

A. 2008. Use of glucocorticoids and risk of infections. Autoimmun Rev 8:153-5.

13.    Savva A, Roger T. 2013. Targeting toll-like receptors: promising therapeutic strategies

for the management of sepsis-associated pathology and infectious diseases. Front

Immunol 4:387.

14.    O'Neill LA, Bowie AG. 2007. The family of five: TIR-domain-containing adaptors in

Toll-like receptor signalling. Nat Rev Immunol 7:353-64.

15.    Schumann J. 2016. It is all about fluidity: Fatty acids and macrophage phagocytosis.

Eur J Pharmacol 785:18-23.

16.    Groves E, Dart AE, Covarelli V, Caron E. 2008. Molecular mechanisms of phagocytic

uptake in mammalian cells. Cell Mol Life Sci 65:1957-76.

17.    Fridman WH. 1991. Fc receptors and immunoglobulin binding factors. Faseb j

5:2684-90.

18.    Ninković J, Roy S. 2012. Morphine decreases bacterial phagocytosis by inhibiting

actin polymerization through cAMP-, Rac-1-, and p38 MAPK-dependent mechanisms.

Am J Pathol 180:1068-1079.

19.    Kaufmann SHE, Dorhoi A, Hotchkiss RS, Bartenschlager R. 2018. Host-directed

therapies for bacterial and viral infections. Nat Rev Drug Discov 17:35-56.

20.    An H, Hou J, Zhou J, Zhao W, Xu H, Zheng Y, Yu Y, Liu S, Cao X. 2008. Phosphatase

SHP-1 promotes TLR- and RIG-I-activated production of type I interferon by inhibiting

the kinase IRAK1. Nat Immunol 9:542-50.

21.    Zheng Y, An H, Yao M, Hou J, Yu Y, Feng G, Cao X. 2010. Scaffolding adaptor

protein Gab1 is required for TLR3/4- and RIG-I-mediated production of

proinflammatory cytokines and type I IFN in macrophages. J Immunol 184:6447-56.

22.    Han C, Jin J, Xu S, Liu H, Li N, Cao X. 2010. Integrin CD11b negatively regulates

TLR-triggered inflammatory responses by activating Syk and promoting degradation

of MyD88 and TRIF via Cbl-b. Nat Immunol 11:734-42.

23.    Franco-Molina MA, Mendoza-Gamboa E, Castillo-León L, Tamez-Guerra RS,

Rodríguez-Padilla C. 2004. Bovine dialyzable leukocyte extract protects against

LPS-induced, murine endotoxic shock. Int Immunopharmacol 4:1577-86.

24.    Zheng Y, Guo Z, He W, Yang Y, Li Y, Zheng A, Li P, Zhang Y, Ma J, Wen M, Yang M,

An H, Ji G, Yu Y. 2012. Ephedrine hydrochloride protects mice from LPS challenge by

promoting IL-10 secretion and inhibiting proinflammatory cytokines. Int

Immunopharmacol 13:46-53.

25.   Zhang AJ, Lee AC, Chan JF, Liu F, Li C, Chen Y, Chu H, Lau SY, Wang P, Chan CC,

Poon VK, Yuan S, To KK, Chen H, Yuen KY. 2021. Coinfection by Severe Acute

Respiratory Syndrome Coronavirus 2 and Influenza A(H1N1)pdm09 Virus Enhances

the Severity of Pneumonia in Golden Syrian Hamsters. Clin Infect Dis 72:e978-e992.

26.   Jiang X, Chen Y, Liu D, Shi T, Cheng X, He W, Li Y, Ryffel B, Zheng SG, Zheng Y.

2020. Secoeudesma sesquiterpenes lactone A alleviates inflammation and offers

adjuvant protection in severe infection of carbapenem-resistant Klebsiella

pneumoniae. J Ethnopharmacol 252:112605.

27.   Ando M, Isogai K, Azami H, Hirata N, Yamazaki H, Yanagi Y. 1992. Studies on the

synthesis of sesquiterpene lactones, 14. Syntheses of (-)-arbusclin D and

(+)-4-EPI-arbusclin D: the stereochemical assignment of arbusclin D. J Nat Prod

55:1620-5.

28.   Jiang X, Wang Y, Qin Y, He W, Benlahrech A, Zhang Q, Jiang X, Lu Z, Ji G, Zheng Y.

2017. Micheliolide provides protection of mice against Staphylococcus aureus and

MRSA infection by down-regulating inflammatory response. Sci Rep 7:41964.

29.   Darkwah S, Nago N, Appiah MG, Myint PK, Kawamoto E, Shimaoka M, Park EJ. 2019.

Differential Roles of Dendritic Cells in Expanding CD4 T Cells in Sepsis. Biomedicines

7.

30.   Bouras M, Asehnoune K, Roquilly A. 2018. Contribution of Dendritic Cell Responses

to Sepsis-Induced Immunosuppression and to Susceptibility to Secondary Pneumonia.

Front Immunol 9:2590.

31.    Chousterman BG, Swirski FK, Weber GF. 2017. Cytokine storm and sepsis disease

667           pathogenesis. Semin Immunopathol 39:517-528.

668    32.    Fang J, Huang Q, Shi C, Gai L, Wang X, Yan B. 2023. Songorine inhibits oxidative

669           stress-related inflammation through PI3K/AKT/NRF2 signaling pathway to alleviate

670           lipopolysaccharide-induced septic acute lung injury. Immunopharmacol Immunotoxicol

671           doi:10.1080/08923973.2023.2281902:1-9.

672    33.    Zhang PH, Zhang WW, Wang SS, Wu CH, Ding YD, Wu XY, Smith FG, Hao Y,

673           Sheng-Wei J. 2023. Efficient pulmonary lymphatic drainage is necessary for

674           inflammation resolution in ARDS. JCI Insight doi:10.1172/jci.insight.173440.

675    34.    Kumar V, Chhibber S. 2011. Acute lung inflammation in Klebsiella pneumoniae

676           B5055-induced pneumonia and sepsis in BALB/c mice: a comparative study.

677           Inflammation 34:452-62.

678    35.    Guirgis FW, Khadpe JD, Kuntz GM, Wears RL, Kalynych CJ, Jones AE. 2014.

679           Persistent organ dysfunction after severe sepsis: a systematic review. J Crit Care

680           29:320-6.

681    36.    O'Neil JD, Bolimowska OO, Clayton SA, Tang T, Daley KK, Lara-Reyna S, Warner J,

682           Martin CS, Mahida RY, Hardy RS, Arthur JSC, Clark AR. 2023. Dexamethasone

683           impairs the expression of antimicrobial mediators in lipopolysaccharide-activated

684           primary macrophages by inhibiting both expression and function of interferon β. Front

685           Immunol 14:1190261.

686    37.    Vilekar P, King C, Lagisetty P, Awasthi V, Awasthi S. 2014. Antibacterial activity of

687           synthetic curcumin derivatives: 3,5-bis(benzylidene)-4-piperidone (EF24) and

EF24-dimer linked via diethylenetriaminepentacetic acid (EF2DTPA). Appl Biochem

Biotechnol 172:3363-73.

38.  Lai X, Ye Y, Sun C, Huang X, Tang X, Zeng X, Yin P, Zeng Y. 2013. Icaritin exhibits

anti-inflammatory effects in the mouse peritoneal macrophages and peritonitis model.

Int Immunopharmacol 16:41-9.

39.  Doherty GJ, McMahon HT. 2009. Mechanisms of endocytosis. Annu Rev Biochem

78:857-902.

40.  Ramanan V, Agrawal NJ, Liu J, Engles S, Toy R, Radhakrishnan R. 2011. Systems

biology and physical biology of clathrin-mediated endocytosis. Integr Biol (Camb)

3:803-15.

41.  Akira S, Uematsu S, Takeuchi OJC. 2006. Pathogen recognition and innate immunity.

124:783-801.

42.  Hankittichai P, Buacheen P, Pitchakarn P, Na Takuathung M, Wikan N, Smith DR,

Potikanond S, Nimlamool W. 2020. Artocarpus lakoocha Extract Inhibits LPS-Induced

Inflammatory Response in RAW 264.7 Macrophage Cells. Int J Mol Sci 21.

43.  Walter EJ, Hanna-Jumma S, Carraretto M, Forni L. 2016. The pathophysiological

basis and consequences of fever. Crit Care 20:200.

44.  Thomas-Rüddel DO, Hoffmann P, Schwarzkopf D, Scheer C, Bach F, Komann M,

Gerlach H, Weiss M, Lindner M, Rüddel H, Simon P, Kuhn SO, Wetzker R, Bauer M,

Reinhart K, Bloos F. 2021. Fever and hypothermia represent two populations of sepsis

patients and are associated with outside temperature. Crit Care 25:368.

45.  Rumbus Z, Matics R, Hegyi P, Zsiboras C, Szabo I, Illes A, Petervari E, Balasko M,

Marta K, Miko A, Parniczky A, Tenk J, Rostas I, Solymar M, Garami A. 2017. Fever Is

Associated with Reduced, Hypothermia with Increased Mortality in Septic Patients: A

Meta-Analysis of Clinical Trials. PLoS One 12:e0170152.

46.   van der Poll T, van de Veerdonk FL, Scicluna BP, Netea MG. 2017. The

immunopathology of sepsis and potential therapeutic targets. Nat Rev Immunol

17:407-420.

47.   Takeuchi O, Akira S. 2010. Pattern recognition receptors and inflammation. Cell

140:805-20.

48.   Kang R, Chen R, Zhang Q, Hou W, Wu S, Cao L, Huang J, Yu Y, Fan XG, Yan Z, Sun

X, Wang H, Wang Q, Tsung A, Billiar TR, Zeh HJ, 3rd, Lotze MT, Tang D. 2014.

HMGB1 in health and disease. Mol Aspects Med 40:1-116.

49.   Opal SM. 2011. Immunologic alterations and the pathogenesis of organ failure in the

ICU. Semin Respir Crit Care Med 32:569-80.

50.   Tang D, Wang H, Billiar TR, Kroemer G, Kang R. 2021. Emerging mechanisms of

immunocoagulation in sepsis and septic shock. Trends Immunol 42:508-522.

51.   Mera S, Tatulescu D, Cismaru C, Bondor C, Slavcovici A, Zanc V, Carstina D, Oltean

MJA. 2010. Multiplex cytokine profiling in patients with sepsis.

52.   Wu HP, Chen CK, Chung K, Tseng JC, Hua CC, Liu YC, Chuang DY, Yang CH. 2009.

Serial cytokine levels in patients with severe sepsis. Inflamm Res 58:385-93.

53.   Mera S, Tatulescu D, Cismaru C, Bondor C, Slavcovici A, Zanc V, Carstina D, Oltean

M. 2011. Multiplex cytokine profiling in patients with sepsis. Apmis 119:155-63.

54.   van der Poll T, Levi M, Hack CE, ten Cate H, van Deventer SJ, Eerenberg AJ, de

Groot ER, Jansen J, Gallati H, Büller HR, et al. 1994. Elimination of interleukin 6

attenuates coagulation activation in experimental endotoxemia in chimpanzees. J Exp

Med 179:1253-9.

55.    Leon LR, White AA, Kluger MJ. 1998. Role of IL-6 and TNF in thermoregulation and

survival during sepsis in mice. Am J Physiol 275:R269-77.

56.    Tracey KJ, Fong Y, Hesse DG, Manogue KR, Lee AT, Kuo GC, Lowry SF, Cerami A.

1987. Anti-cachectin/TNF monoclonal antibodies prevent septic shock during lethal

bacteraemia. Nature 330:662-4.

57.    Kilbourn RG, Griffith OW, Gross SS. 1993. Pathogenetic mechanisms of septic shock.

N Engl J Med 329:1427-8.

58.    Leon LR. 2004. Hypothermia in systemic inflammation: role of cytokines. Front Biosci

9:1877-88.

59.    Lang GP, Ndongson-Dongmo B, Lajqi T, Brodhun M, Han Y, Wetzker R, Frasch MG,

Bauer R. 2020. Impact of ambient temperature on inflammation-induced

encephalopathy in endotoxemic mice-role of phosphoinositide 3-kinase gamma. J

Neuroinflammation 17:292.

60.    Jhaveri KA, Trammell RA, Toth LA. 2007. Effect of environmental temperature on

sleep, locomotor activity, core body temperature and immune responses of C57BL/6J

mice. Brain Behav Immun 21:975-87.

61.    Clemmer TP, Fisher CJ, Jr., Bone RC, Slotman GJ, Metz CA, Thomas FO. 1992.

Hypothermia in the sepsis syndrome and clinical outcome. The Methylprednisolone

Severe Sepsis Study Group. Crit Care Med 20:1395-401.

62.     Gentek R, Molawi K, Sieweke MH. 2014. Tissue macrophage identity and self-renewal.

755             Immunol Rev 262:56-73.

756     63.     Yan J, Horng T. 2020. Lipid Metabolism in Regulation of Macrophage Functions.

757             Trends Cell Biol 30:979-989.

758     64.     Rescigno M, Granucci F, Ricciardi-Castagnoli P. 1999. Dendritic cells at the end of the

759             millennium. Immunol Cell Biol 77:404-10.

760     65.     Lipscomb MF, Masten BJ. 2002. Dendritic cells: immune regulators in health and

761             disease. Physiol Rev 82:97-130.

762     66.     Liu Y, Pan YF, Xue YQ, Fang LK, Guo XH, Guo X, Liu M, Mo BY, Yang MR, Liu F, Wu

763             YT, Olsen N, Zheng SG. 2018. uPAR promotes tumor-like biologic behaviors of

764             fibroblast-like synoviocytes through PI3K/Akt signaling pathway in patients with

765             rheumatoid arthritis. Cell Mol Immunol 15:171-181.

766     67.     Saraiva M, O'Garra A. 2010. The regulation of IL-10 production by immune cells. Nat

767             Rev Immunol 10:170-81.

768     68.     Surbatovic M, Popovic N, Vojvodic D, Milosevic I, Acimovic G, Stojicic M, Veljovic M,

769             Jevdjic J, Djordjevic D, Radakovic S. 2015. Cytokine profile in severe Gram-positive

770             and Gram-negative abdominal sepsis. Sci Rep 5:11355.

**Figure legends**

**Figure 1. DHZD inhibits the secretion of LPS-induced cytokines from RAW264.7 cells.**

Chemical structure of DHZD (A). DHZD had no cytotoxicity on RAW264.7 cells (B). DHZD

decreased the secretion of IL-6 (C), TNF-α (D), MCP-1 (E), IFN-β (F), IL-1β (G) and IL-10 (H) in

the cell culture supernatants induced by LPS. Values are means of three replicate determinations

(n=3) ± SD. *, $P<0.05$, **, $P <0.01$, and ***, $P <0.001$.

**Figure 2. DHZD decreases the secretion of LPS-induced expression of IL-6, TNF-α, MCP-1,**

**IL-1β, IFN-β and IL-10 in mouse primary macrophages.** DHZD decreased LPS-induced

secretion of IL-6 (A), TNF-α (B), MCP-1 (C), IL-1β (D), IFN-β (E), IL-10 (F) and NO (G). Data

are means of three replicate determinations ± SD (n=3). *, $P <0.05$, **, $P <0.01$, and ***, $P$

$<0.001$.

**Figure 3. DHZD decreases LPS-induced secretion of inflammatory cytokines in BMDCs, and**

**downregulated the expression of MHC II and co-stimulatory molecules on the surface of**

**BMDCs.** DHZD down-regulated the production of IL-6 (A), TNF-α (B) and IL-12p70 (C) in cell

culture supernatants induced by LPS. The expression of co-stimulatory molecules CD40 (D),

CD86 (E), CD80 (F) and antigen-presenting molecule MHC II (Iab) (G) was quantified by FACS

and the MFI (mean fluorescence intensity) of each sample was analyzed using FlowJo. Data were

shown as mean ± SD of three separated experiments. *, $P<0.05$, **, $P <0.01$, ***, $P <0.001$,

and ****, $P <0.0001$.

**Figure 4. DHZD protects septic mice by inhibiting the over-activated inflammatory response.**

Survival (A). Data were analyzed using Log-Rank test and survival curve was generated by

GraphPad. H&E staining of lung, kidney and liver (100×) (B). Scale bar 100 μm, and the

histological core of lung (C). The serum levels of IL-6 (D), TNF-α (E), MCP-1 (F) and IL-10 (G)

were measured by ELISA. Data were shown as mean ± SD of nine mice per group. *, $P <0.05$, **,

$P <0.01$, ***, $P <0.001$, and ****, $P <0.0001$.

**Figure 5. DHZD increases the numbers of lung neutrophils, monocytes, and macrophages**

**induced by LPS.** The proportions of lung neutrophils, monocytes and macrophages in lungs were

detected. One representative scatter plot per group and the proportions of neutrophils (A-C). One

representative scatter plot and the proportion of monocyte-macrophages (D-H). Data were shown

as means ± SD for each group (nine mice in each group). *, $P<0.05$, **, $P<0.01$, ***, $P<0.001$,

and ****, $P < 0.0001$.

**Figure 6. DHZD enhances the phagocytosis of pHrodo-labelled E. coli in macrophages in a**

**clathrin-dependent and actin-polymerized manner via CD36.** One representative scatter plot

out of three independent experiments were shown (A). The mean fluorescence intensity (MFI) of

pHrodo-labelled *E. coli* swallowed by RAW264.7 cells were analyzed (B, C). DHZD enhanced the

phagocytosis in a clathrin-dependent and actin-polymerized manner (D-F). The expression of

CD36 (G, H) were detected by PCR. Data were shown as mean ± SD from three independent

experiments. *, $P<0.05$, **, $P<0.01$, ***, $P<0.001$, and ****, $P<0.0001$.

**Figure 7. DHZD inhibits LPS-stimulated activation of PI3K/Akt/p70S6K pathway.** Total

protein of cells was detected by Western blot (A-E). Phospho-Akt (Ser 473), phospho-GSK-3β

(Ser9), phospho-ERK, phospho-p38, phospho-IκBα (Ser32/36), phospho-p65 (Ser536) and

β-Actin in RAW264.7 were verified by Western blot. Data were shown as mean ± SD from three

independent experiments; *, $P<0.05$, **, $P<0.01$, ***, $P<0.001$, and ****, $P<0.0001$.

**Figure 8. DHZD reverses hypothermia, tissue damage, and over-activated inflammation**

**caused by CRKP.** Temperature difference (A, B). H&E staining of lung and liver (200×) (C).

Scale bar 100 μm, and the histological core of lung (D). DHZD dose-independently reduced the

817 secretion of IL-6 (E), TNF-α (F), MCP-1 (G), IL-1β (H), IL-10 (I) and MIP-2 (J) induced by

818 HK-CRKP. Data were shown as mean ± SD from three independent experiments. *, $P<0.05$, **, $P$

819 $<0.01$, ***, $P<0.001$, and ****, $P<0.0001$.