Dehydrozaluzanin C- derivative protects septic mice by alleviating over-activated inflammatory response and promotes the phagocytosis of macrophages

Ying-xiang Zou[1,2,#], Tian-nan Xiang[5,1,2,#], Li-rong Xu[1,2], Huan Zhang[1,2], Yu-he Ma[1,2], Lu Zhang[1,2], Chun-xian Zhou[1,2], Xiao Wu[1,2], Qi-lin Huang[1,2], Biao Lei[1,2], Jing-wen Mu[1,2], Xiang-yang Qin[4,*], Xin Jiang[1,2,*], Yue-juan Zheng[1,2,3,*]

[1]School of Integrative Medicine, Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China

[2]The Research Center for Traditional Chinese Medicine, Shanghai Institute of Infectious Diseases and Biosecurity, Shanghai University of Traditional Chinese Medicine, Shanghai, 201203, China

[3]Shanghai Key Laboratory of Health Identification and Assessment, Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China

[4]Department of Chemistry, school of pharmacy, Furth Military University, Xi'an, Shaanxi, 710032, China

[5]Department of Chinese Medicine, Hubei College of Chinese Medicine, Jingzhou, Hubei, 434020, China

[#]These authors contributed equally to this work.

[*] **Correspondence**: Prof. Yue-juan Zheng and Prof. Xin Jiang, School of Integrative Medicine, Shanghai University of Traditional Chinese Medicine, Shanghai, 201203, China. Tel: (+86)21-5132 2134, Fax: (+86)-21-5132 2130, Email: 13641776412@163.com (Yue-juan Zheng), and jiangxingao@163.com (Xin Jiang).

Prof. Xiang-yang Qin, Department of Chemistry, school of pharmacy, Furth Military

University, Xi'an, Shaanxi 710032, China. Tel: (+86) 029-8477 4473, Fax: (+86) 029-

8477 4473, Email: qinxiangyang@fmmu.edu.cn.

Abstract

In the course of sepsis, excessive inflammatory response can lead to cytokine

storm. The severity of cytokine storm is closely related with poor prognosis.

Dehydrozaluzanin C-derivative (DHZD) is a modified compound from *Magnolia*

officinalis. Here, we demonstrated that DHZD ameliorated tissue damage (lung, kidney

and liver) and excessive inflammatory response induced by LPS challenge or the

infection of *Carbapenem-resistant Klebsiella pneumoniae* (CRKP). In LPS-induced

sepsis model, DHZD could reduce the secretion of inflammatory factors IL-6, TNF-α,

etc. and increased the survival rate. *In vitro* experiments, DHZD also decreased LPS-

stimulated expression of IL-6, TNF-α and MCP-1 via PI3K/Akt/p70S6K signaling

pathway in macrophages, etc., might accounting for the decreased tissue damage and

its protective role *in vivo*. Interestingly, the combined treatment group of DXM and

DHZD had higher survival rate and lower level of IL-6 than those of DXM-treated

group. The combination of DHZD and DXM played a synergistically role to decrease

IL-6 secretion in sera. DHZD not only improved the hypothermic symptoms of acute

peritonitis induced by CRKP, but also inhibited heat-killed CRKP-induced

inflammatory response in macrophages. Similar to that of DXM-treated group, the

proportions of phagocytic cell types, e.g., neutrophils and monocytes/macrophages in

lungs were upregulated by DHZD. What's more, the phagocytic receptor CD36 was

increased by DHZD in macrophages, which was accompanied by increased bacterial

phagocytosis in clathrin- and actin-dependent manner. In short, DHZD is a potential

drug candidate with host-directed immunomodulatory effects during bacterial infection

through increasing the numbers and the phagocytic ability of phagocytes and decreasing

their inflammatory responses together.

Key words: DHZD, sepsis, CRKP, phagocytosis, cytokine storm

Introduction

Sepsis is a life-threatening syndrome of multiple organ dysfunction, which is

usually caused by bacterial infection, etc. Gram-negative (G$^-$) bacteria account for a

large proportion of severe sepsis, which is a serious threat to global public health and

economic burden (1, 2). The incidence of sepsis is much higher than myocardial

infarction (3) or the combination of prostate, breast, and lung cancer (4). As reported,

there are 15 to 30 million cases of sepsis worldwide every year (5, 6). The mortality

rate is close to 20 %. In recent decades, antimicrobial resistance (AMR) poses an

increasing threat to global public health. CRKP is an important strain in clinical practice,

which can cause severe nosocomial infections, such as sepsis, pneumonia, urinary tract

infection, surgical site infection and soft tissue infection [11]. CRKP accounts for about

64 % of carbapenem-resistant enterobacterial infections in China [12].

The current clinical treatment for sepsis caused by bacterial infection mainly relies

on antibiotic therapy to control the infection and the use of glucocorticoid to inhibit

excessive inflammatory response. Glucocorticoids (such as dexamethasone) are

effective in treating sepsis, but they also accelerate the apoptosis of specific immune

cells. Side effects of long-term glucocorticoid use include osteoporosis (7) or

opportunistic infections due to its immunosuppressive effects (8). However, the

development of new antimicrobial drugs, especially targeting G⁻ bacteria, lags behind

the emergence of antibiotic resistance obviously, which has become a major obstacle in

the clinic. New drug candidates are eagerly wanted.

As a major component of the cell wall of G⁻ bacteria, lipopolysaccharide (LPS) is

one of the most important pathogen-associated molecular patterns, and can elicit

inflammatory immune response via Toll-like receptor 4 (TLR4) in various immune cells

(9). After ligation of TLR4, adaptor molecules such as myeloid differentiation protein

88 (MyD88), interleukin-1 (IL-1) receptor-associated kinases (IRAKs), tumor necrosis

factor (TNF) receptor-associated factor 6 (TRAF6), transforming growth factor (TGF)-

activated kinase 1 (TAK1), and subsequent MAPKs and nuclear factor κB (NF-κB) are

activated. Excessive inflammation can lead to "cytokine storm", commonly known as

systemic inflammatory response syndrome (SIRS). The pathologic progress of sepsis

is usually accompanied by SIRS. The uncontrolled inflammatory response usually

causes organ dysfunction, multiple organ failure and even death (10).

Besides the inflammatory immune damage, the poor clearance of pathogens is

usually associated with poor prognosis during bacterial infection.

Monocytes/macrophages, and dendritic cells (DCs) are the main phagocytes to

phagocytize bacteria and present their antigens to T cells (11). Neutrophil is also an

important cell type of phagocytes. The progress of phagocytosis includes 3 stages:

receptor-mediated, membrane fusion, and actin-driven (12). There are a variety of

phagocytic receptors, including C-type lectin receptors (CLRs), mannose receptors,

scavenger receptors (SR-A1, MARCO and CD36), complement receptors (CR3 and

CR4) and Fc receptors (12-14).

Host-directed therapy (HDT) is a new adjuvant therapy strategy especially for the

treatment of severe infection. It is not mediated by killing pathogens directly, but

promoting the immune clearing of pathogens through phagocytes, and/or reduces tissue

damage caused by excessive inflammatory response. Thus, HDT drug candidates could

play a protective role for the treatment of bacterial infection from the aspect of host.

In our previous work, dehydrozaluzanin C (DHZC), a sesquiterpenoid from

Magnolia officinalis, has strong anti-inflammatory activity but its drug toxicity is

relatively high. DZHD is a modified compound from DHZC by decreasing cytotoxicity

and increase the druggability.

In this study, the protective effects of DHZD on mice in lethal LPS challenge or

CRKP-infected septic mouse models and the underlying mechanisms were

demonstrated.

Materials and methods

Animals and reagents

Six-week-old wild-type female C57BL/6J mice (18 ± 2 g) were obtained from

Joint Ventures Sipper BK Experimental Animal Co. (Shanghai, China). All mice had

free access to food and water during all the experiments and were housed in a 12/12 h

light/dark cycle. The experimental procedures were approved by the Experimental

Animal Ethical Committee, Shanghai University of Traditional Chinese Medicine

(Shanghai, China) (PZSHUTCM220307018).

DHZD was modified from DHZC isolated from *Magnolia officinalis*. LPS (0111:

B4), DMSO, dexamethasone (DXM) and chlorpromazine hydrochloride (CPZ) were

purchased from Sigma (St. Louis, MO, USA). Cell counting kit 8 (CCK-8) were

obtained from Meilunbio (Dalian, China). Enzyme-linked immunosorbent assay

(ELISA) kits for mouse IL-6, TNF-α, MCP-1, IL-10 IFN-β, and granulocyte-

macrophage colony stimulating factor (mGM-CSF) were purchased from R&D

Systems (Minneapolis, MN, USA). Mouse recombinant cytokine IL-4 were from

Peprotech (USA). The detection kit of nitric oxide (NO) was obtained from Beyotime

Biotechnology (Shanghai, China). FITC anti-mouse CD11c, APC anti-mouse CD40,

APC/Fire™ 750 anti-mouse CD80, PerCP/Cyanine5.5 anti-mouse CD86, Brilliant

Violet 421™ anti-mouse I-A/I-E, FITC anti-mouse CD45, Brilliant Violet 650 Ly6G,

Brilliant Violet 421 Ly6C and PE/Cyanine7 F4/80 were purchased from Biolegend

(USA). Dexamethasone sodium phosphate injection was from Sinopharm (China).

SYBR RT-PCR Kit was purchased from Takara (Dalian, China). PHrodo-labelled

E. coli was purchased from ThermoFisher (USA). Cytochalasin D was obtained from

Abcam (USA). Filipin III was from MCE (USA). Levofloxacin (LVX) and meropenem

(MEM) were obtained from the National Institute for Food and Drug Control of China

(Beijing, China) CRKP (HS11286) was kindly provided by Dr. Yi-jian Chen from

Huashan Hospital affiliated to Fudan University.

Cell culture

Mouse macrophage-like cell line RAW264.7 were purchased from ATCC

(Manassas, VA, USA) and cultured as previously described (15).

Thioglycolate-elicited mouse primary peritoneal macrophages were isolated from

female C57BL/6J mice at 6-8 weeks of age and cultured in DMEM (GE, USA) with

10 % FBS (Biowest, France) and 1 % Penicillin-Streptomycin, as previously described

(16).

Bone marrow-derived macrophages (BMDMs) and bone marrow-derived

dendritic cells (BMDCs) were both obtained from C57BL/6J mice (4 weeks old). The

BMDMs were cultured in DMEM containing 10 ng/mL M-CSF (Peprotech, USA) and

10 % FBS. The BMDCs were differentiated in RPMI 1640 (GE, USA) medium

containing 10 % FBS, 50 ng/mL mGM-CSF (R&D Systems, USA) and 4 ng/mL mIL-

4 (Peprotech, USA).

Cell viability assay

RAW264.7 cells (1×10^5 cells/well) were cultured in 96-well plates overnight and

treated with different concentrations of DHZD (0-15 μM) in 12, 24, 48 and 72 h. The

cytotoxicity of DHZD was evaluated by CCK-8 counting kit according to the

manufacturer's instructions (Meilunbio, Dalian, China).

Enzyme-linked immunosorbent assay

The inflammatory factors (IL-6, TNF-α, MCP-1, IL-10 and IFN-β) in the cell

culture supernatants were detected by ELISA (R&D Systems, USA).

Blood samples were collected at 12 h after LPS challenge and allowed to clot for

3 h followed by centrifugation at 3500 rpm for 25 min (4 °C) to prepare serum sample.

Serum inflammatory cytokines (IL-6, TNF-α, MCP-1 and IL-10) were measured by

corresponding ELISA kit (R&D Systems, USA) according to the manufacturer's

instructions.

Assessment of nitric oxide (NO) level

158 Quantitative detection of NO was carried out by NO detection kit (S0021) which

159 was purchased from Beyotime Biotechnology (Shanghai, China).

Detection of MHC II and co-stimulatory molecules on BMDCs

Bone marrow stem cells were derived and differentiated into BMDCs. After 5 days,

BMDCs were seeded in 12-well plates overnight. LPS (100 ng/mL) and DHZD (15 μM)

were added into culture medium of BMDCs for another 24 h. Then BMDCs were

harvested and labeled with FITC anti-mouse CD11c (Biolegend, 117305, USA), APC

anti-mouse CD40 (Biolegend, 124611, USA), APC/Fire™ 750 anti-mouse CD80

(Biolegend, 104739, USA), PerCP/Cyanine5.5 anti-mouse CD86 (Biolegend, 105027,

USA) and Brilliant Violet 421™ anti-mouse I-A/I-E (Biolegend, 107631, USA). And

then cells were analyzed by Flow cytometry (Beckman, USA), and data were analyzed

using FlowJo software.

The establishment of septic mice by lethal dose of LPS challenge

The septic mouse model was established by intraperitoneal injection of LPS (12.5

mg/kg, Sigma) as described (17, 18). Six groups were included: LPS group, LPS +

DXM group (5 mg/kg), LPS + DHZD group (10 mg/kg), LPS + DHZD group (20

mg/kg), LPS + DHZD group (40 mg/kg) and LPS + MEM (5 mg/kg) + DHZD group

(10 mg/kg). The survival statuses of mice in different groups were recorded at different

intervals as described previously (19).

RNA extraction and real-time PCR

178   Total RNA was extracted from lung tissues or different types of cells and reverse-

179 transcribed into cDNA. Real-time PCR was performed with SYBR RT-PCR kit (Takara,

180 Dalian, China). The mRNA levels were normalized against β-Actin and presented as 2^{-}

181 $^{\Delta\Delta Ct}$. The primer sequences for qRT-PCR amplification were as the following:

mβ-Actin	mβ-Actin-F	5'-AGTGTGACGTTGACATCCGT-3'
	mβ-Actin-R	5'-GCAGCTCAGTAACAGTCCGC-3'
mIL-6	mIL-6F	5'-TAGTCCTTCCTACCCCAATTTCC-3'
	mIL-6R	5'-TTGGTCCTTAGCCACTCCTTC-3'
mTNF-α	mTNF-αF	5'-AAGCCTGTAGCCCACGTCGTA-3'
	mTNF-αR	5'-GGCACCACTAGTTGGTTGTCTTTG-3'
mMCP-1	mMCP-1F	5'- GCATCCACGTGTTGGCTCA -3'
	mMCP-1R	5'- CTCCAGCCTACTCATTGGGATCA -3'
mMIP-2	mMIP-2-F	5'-CACTCTCAAGGGCGGTCAA-3'
	mMIP-2-R	5'-AGGCACATCAGGTACGATCCA-3'
mIL-10	mIL-10F	5'-GCTCTTACTGACTGGCATGAG-3'
	mIL-10R	5'-CGCAGCTCTAGGAGCATGTG-3'
mCD206	mCD206F	5'-CTCTGTTCAGCTATTGGACGC-3'
	mCD206R	5'-TGGCACTCCCAAACATAATTTGA-3'
mDectin-1	mDectin-1F	5'-AGGCATCCCAAACTACAGGAG-3'
	mDectin-1R	5'-GGAGCAGTGTCTCTTACTTCCAT-3'
mCR3	mCR3F	5'- CCATGACCTTCCAAGAGAATGC -3'

	mCR3R	5'- ACCGGCTTGTGCTGTAGTC -3'
mCR4	mCR4F	5'- CTGGATAGCCTTTCTTCTGCTG -3'
	mCR4R	5'- GCACACTGTGTCCGAACTCA -3'
mFcγR2a	mFcγR2aF	5'-TGGACAGCCGTGCTAAATCTT-3'
	mFcγR2aR	5'-GGTCCCTTCGCATGTCAGTG-3'

Flow cytometry

The left lungs of mice in the different groups were collected at 12 h after LPS challenge, and the lung tissues were snipped and digested with collagenase (Sigma Aldrich) and DNase I (Sigma Aldrich) in gentleMACS™ (Miltenyi, Germany) at 80 rpm (37 °C) for 55 min. After digestion, the samples were filtered through a 40 mm filter using a syringe piston, and centrifuged at 1000 rpm (4 °C) for 5 min. RBC lysis buffer (BD Biosciences, USA) was added to cell pellets at room temperature for 5 min, and then PBS (8 mL) was added and centrifuged at 1400 rpm (4 °C) for 5 min. Cells were counted and then incubated at 4 °C for 25 min with FITC anti-mouse CD45 (Biolegend, 103107, USA), Brilliant Violet 650 Ly6G (Biolegend, 127641, USA), Brilliant Violet 421 Ly6C (Biolegend, 128031, USA) and PE/Cyanine7 F4/80 (Biolegend, 123113, USA) antibodies. Samples were run on CytoFlex LX (Beckman, USA). Data were analyzed using FlowJo software.

Detection of the phagocytosis of macrophages

The phagocytosis of pHrodo labelled *E. coli* (ThermoFisher, California, USA) was detected in RAW264.7 cells by Flow cytometry. Cells were stimulated by DHZD (15 μM) with or without LPS (100 ng/mL) for 24 h, and then co-incubated with pHrodo-

labelled *E. coli* (10 μL/well) at 37 °C for 1 h. RAW264.7 cells were pretreated by 3 μM

of Filipin III (MCE, USA), 10 μM of chlorpromazine hydrochloride (Sigma, USA) or

10 μM of cytochalasin D (Abcam, USA) for 30 min, and then pHrodo labelled *E. coli*

were added for another 1 h. The phagocytosis of *E. coli* was investigated by BD

Accuri™ C6 Flow cytometry (BD, USA), and data were analyzed by FlowJo software.

Western blot

The phosphorylation of important signaling proteins from RAW264.7 cells and

their corresponding total proteins were determined by Western blot as previously

described (20). All the antibodies were bought from Cell Signaling Technology

(Beverly, MA, USA).

CRKP-induced septic mouse model

The septic mouse model was established by intraperitoneal injection of CRKP

(9×10^7 CFU/mouse) in 0.4 mL of PBS. The experiments were carried out in the

following groups: CRKP group, CRKP + MEM (5 mg/kg) group, CRKP + LVX (75

mg/kg) group, CRKP + DHZD group (20 mg/kg), CRKP + DHZD group (40 mg/kg)

and the combined treatment CRKP + DHZD (20 mg/kg) + MEM (5 mg/kg) group. The

body temperature of mice was recorded by infrared body thermometer (HJKIR, Wuhan)

before or 9 hours after injection of CRKP.

Preparation of HK-CRKP

Monoclonal colonies were picked from the CRKP bacterial culture plates and

cultured in LB medium for 12 h at 37°C on a constant temperature shaker. After heating

1 mL of the bacterial solution in a 65°C water bath for 45 min, the solution was

centrifuged (4400 rpm×15 min), resuspended in sterile PBS, centrifuged again (4400

rpm×15 min), and resuspended in DMEM for further use (MOI=10).

Histopathology

Hematoxylin and eosin (H&E) were used to measure the pathological damage of

lungs, livers and kidneys. After 12 h of LPS challenge (10 mg/kg), mice were sacrificed

and lung, liver and kidney tissues were collected to be fixed with 4 % formaldehyde

and paraffin-embedded.

CRKP-induced (6×10^7 CFU/mouse) peritonitis mouse model was carried out with

C57BL/6J mice and different groups were treated as indicated. After 12 h, mice were

sacrificed and lungs and livers were collected to be fixed with 4 % formaldehyde and

embedded in paraffin.

All of the above tissues were sliced and stained with H&E. The histopathological

changes were observed using a PANNORAMIC MIDI II microscope (3DHistech,

Hungary). Pulmonary edema, alveolar infiltration, and pulmonary vasculitis were

scored on a 12-point pathological scale (21).

Statistical analysis

Statistical analysis was performed using GraphPad Prism 9. Data were presented

as means ± standard deviation (SD). Student's t-test analyses was used to evaluate the

comparisons between two groups. Comparisons multiple groups were evaluated by

one-way ANOVA followed by Tukey's post hoc comparisons, respectively. P values

were considered as *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; ****, $P<0.0001$.

Results

DHZD inhibited the secretion of LPS-induced cytokines in RAW264.7 cells

CCK-8 kit was used to determine whether DHZD affected cell viability. The

structure of DHZD was shown in Fig. 1A. DHZD was synthesized according to the

previously reported method (22). Spectroscopic data were in accordance with reported

values (Supplementary Fig.1)(22). DHZD (0-15 μM) did not affect the proliferation of

RAW264.7 cells within 72 h. At the same time, DHZD inhibited LPS-induced secretion

of IL-6, TNF-α, MCP-1, IFN-β, and IL-10 in a dose-dependent manner (Fig. 1B-H).

Figure 1. DHZD inhibited the secretion of LPS-induced cytokines in RAW264.7 cells.

RAW264.7 were co-cultured with different concentrations of DHZD, and the absorbance at 450 nm

at 12, 24, 48, 72 h to calculate the cell survival rate according to the formula (B). RAW264.7 (2×10^5

cells/well) were plated as indicated and administrated with DHZD and/or LPS (100 ng/mL) for 6 h

255   and 18 h. ELISA was carried out to detect the secretion of IL-6 (C), TNF-α (D), MCP-1 (E), IFN-β

256   (F), IL-1β (G) and IL-10 (H) in the cell culture supernatants. Values are means of three replicate

257   determinations (n=3) ± SD. *, $P<0.05$, **, $P<0.01$, and ***, $P<0.001$.

258   DHZD suppressed the secretion of LPS-induced expression of IL-6, TNF-α, MCP-

259   1, IL-1β, IFN-β and IL-10 in mouse primary macrophages

Compared to macrophage-like cell line, the biological activities of primary

macrophages are closer to the physiological and pathological characteristics of

macrophages *in vivo*. Therefore, mouse primary peritoneal macrophages and BMDMs

were used to validate the anti-inflammatory effects of DHZD. DHZD was also able to

inhibit the secretion of LPS-induced pro-inflammatory mediators IL-6, TNF-α, IL-1β,

IFN-β, NO, MCP-1 and anti-inflammatory cytokine IL-10 (Fig. 2A-G). Similarly,

DHZD decreased the secretion of IL-6, TNF-α and IL-10 after LPS stimulation in

BMDMs (Supplementary Fig. 2A-C).

Figure 2. DHZD suppressed the secretion of LPS- induced expression of IL-6, TNF-α, MCP-

1, IL-1β, IFN-β and IL-10 in mouse primary macrophages. Mouse primary macrophages were

plated (3.5×10^5 cells/500 μL) in 24-well plates at the day before stimulated. Cell culture

supernatants were collected at 6 h and 18 h after co-stimulation of different concentrations of DHZD

and LPS (100 ng/mL), and the concentrations of IL-6 (A), TNF-α (B), MCP-1 (C), IL-1β (D), IFN-

β (E), IL-10 (F) and NO (G) were measured by ELISA. Data are means of three replicate

determinations ± SD (n=3). *, $P<0.05$, **, $P<0.01$, and ***, $P<0.001$.

DHZD decreased LPS-induced secretion of inflammatory cytokines in BMDCs,

and DHZD also downregulated the expression of MHC II and co-stimulatory

molecules on the surface of BMDCs

DCs are the most important and powerful antigen presenting cells to activate

naive T cells, bridging innate immunity and adaptive immunity. They participate in the

phagocytosis, digestion of bacteria and presenting their antigens to evoke T cell

immunity. During the transition of DCs from immature to mature status, co-stimulatory

molecules are upregulated, downregulating the phagocytosis and promoting the antigen

presenting ability, demonstrating the biological function of DCs. Meanwhile, DCs also

secret inflammatory cytokines (IL-6, TNF-α, IL-12 and a variety of chemokines),

contributing to the formation of cytokine storm together with activated T cells during

the pathological progress of sepsis. DHZD reduced the secretion of IL-6, TNF-α, and

IL-12p70 (Fig. 3A-C) induced by LPS.

In order to investigate whether DHZD has a regulatory effect on the antigen

presentation function of DCs, thereby affecting the activation of T cells, Flow

cytometry was used to detect LPS-stimulated expression of costimulatory molecules

(CD80, CD86, and CD40) and MHC Class II molecule (Iab). DHZD inhibited the

expression of these molecules (Fig. 3D-G).

Figure 3. DHZD decreased LPS-induced secretion of inflammatory cytokines in BMDCs, and

DHZD also downregulated the expression of MHC II and co-stimulatory molecules on the

surface of BMDCs. Mouse bone marrow-derived dendritic cells were seeded at 3.5×10^5 cells/500

μL in 24-well plates along with stimulation by DHZD (0-15 μM) with or without LPS (100 ng/mL)

for 6 h or 18 h. The production of IL-6 (A), TNF-α (B) and IL-12p70 (C) in cell culture supernatants

were detected by ELISA. BMDCs were seeded in 12-well plate and treated with or without LPS

(100 ng/mL) along with DHZD (15 μM) for another 24 h. The expression of costimulatory

molecules CD40 (D), CD86 (E), CD80 (F) and antigen-presenting molecule MHC II (Iab) (G) was

quantified by FACS and the MFI (mean fluorescence intensity) of each sample was analyzed using

FlowJo. Data were show as mean ± SD of three separated experiments. *, $P<0.05$, **, $P<0.01$, ***,

$P<0.001$, and ****, $P<0.0001$.

DHZD protected septic mice by inhibiting the over-activated inflammatory response

To determine the protective role of DHZD against LPS-induced (12.5 mg/kg)

septic shock, a survival analysis was carried out. DXM, a glucocorticoid used in the

clinical treatment of bacterial infection, was used as a positive control drug.

All mice in the model group died within 40 h. 60 % of mice in DXM treatment

group survived, 10 % of mice in low dose DHZD treatment group survived, 30 % of

mice in medium or high dose DHZD treatment group survived at the end of observation.

However, 100 % of mice in low dose DHZD and DXM treatment group survived (Fig.

4A).

In the control group, the lung, liver and kidney tissue structure were intact. On the

contrary, after LPS challenged, the lung tissue pathology of mice showed obvious

infiltration of inflammatory cells, a large amount of fluid exudation in the alveoli, and

lung interstitial thickening. The liver tissue also showed obvious inflammatory cell

infiltration, swelling and necrosis of hepatocytes, fragmentation and dissolution of some nuclei, and destruction of the hepatic plate structure. The renal tissue also showed obvious inflammatory cell infiltration, degeneration and necrosis of renal tubular epithelial cells, and narrowing of renal cysts. The destruction of lung, liver and kidney tissue structure was improved after DHZD treatment (Fig. 4B). DHZD was found to improve pulmonary edema, alveolar infiltration, and pulmonary vasculitis by pathological score of lung tissue (Fig. 4C).

Multiple organ failure caused by a "cytokine storm" during infection is the leading cause of death in sepsis patients. In septic mice model of LPS-induced (10 mg/kg), DHZD and DXM suppressed LPS-induced production of IL-6, TNF-α, MCP-1 and IL-10. Compared with DXM treatment group, the combination treatment could significantly inhibit the secretion of these cytokines (Fig. 4D-G).

Sepsis induced by bacterial infection can cause systemic organ dysfunction, and lungs are often the most easily affected important organs. Therefore, Sepsis is often accompanied by acute lung injury (ALI) or acute respiratory distress syndrome (ARDS), which is one of the most important prognostic factors of sepsis death. We detected the expression of lung inflammatory factors by qPCR, and found that both DHZD administration group and combined administration group could significantly inhibit the production of proinflammatory cytokines IL-6, chemokines MCP-1 and MIP-2 in the lungs of mice (Supplementary Fig. 3A-C).

Figure 4. DHZD protected septic mice by inhibiting the over-activated inflammatory response.

C57BL/6J female mice were randomly divided into 6 groups (10 mice/group). All groups of mice

were challenged by lethal dose of LPS (12.5 mg/kg, ip) and corresponding agents, respectively.

Survival status of different groups were recorded every 2-3 h in the following continuous 180 h (A).

Data were analyzed using Log-Rank test and survival curve was generated by GraphPad Prism 9.0.

C57BL/6J mice were randomly divided into 5 groups (9 mice/group). All groups of mice were

infected with LPS (10 mg/kg, ip) group，PBS group, LPS + DXM (5 mg/kg) group, LPS + DHZD

(40 mg/kg) group and LPS + DXM (5 mg/kg) + DHZD (10 mg/kg) group. After 12 h of intervention,

blood samples were collected from eyeball. H&E staining of lung, kidney and liver (100×) (B).

Scale bar 100 μm, and the histological core of lung (C). The serum levels of IL-6 (D), TNF-α (E),

MCP-1 (F) and IL-10 (G) were measured by ELISA. Data were shown as mean ± SD of nine mice

per group. *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, and ****, $P < 0.0001$.

DHZD increased the cell numbers of lung neutrophils, monocytes and

macrophages induced by LPS challenge

In most cases, locally recruited natural immune cells (such as neutrophils and

monocytes/macrophages) can rapidly phagocyte bacteria, control the bacterial load in

tissues, and reduce the occurrence of severe infection. However, when severe infection

occurs, a large number of immune cells participate in the secretion of inflammatory

cytokines, forming cytokine storms, which can lead to organ damage, sepsis and even

death. Thus, in the early stage of infection, rapid phagocytosis and elimination of

bacteria, and subsequently control of the body's immune response within reasonable

limitation, are the key aspects for treatment.

The infiltration of neutrophils and monocytes/macrophages in the lungs were

measured by Flow cytometry. As shown in Figure 5, at 12 h after LPS injection, the

ratio of neutrophils (Fig. 5A-C), monocytes/macrophages (Fig. 5D-H) increased in

DHZD-treated group and in DHZD + DXM group compared to LPS challenged group,

suggesting an increase in the number of innate immune cells with the ability to

phagocyte bacteria. It may be beneficial to play a rapid phagocytic role in the early

stage of bacterial infection, reducing the bacterial load and the severity of infection in

tissues, and thus play a potential protective role.

Figure 5. DHZD increased the proportions of lung neutrophils, monocytes and macrophages

induced by LPS challenge. Mice were injected by PBS, LPS, DXM, or DHZD, accordingly. At 12

h after stimulation, mice were sacrificed to collect their lung tissues and single cell suspensions

were prepared for FACS analysis. The proportions of lung neutrophils, monocytes and macrophages

in lungs were detected. One representative scatter plot per group and the proportions of neutrophils

(A-C). One representative scatter plot and the proportion of monocyte-macrophages (D-H). Data

were shown as means ± SD for each group (nine mice in each group). *, $P<0.05$, **, $P <0.01$, ***,

$P <0.001$, and ****, $P < 0.0001$.

DHZD enhanced the phagocytosis of pHrodo-labelled *E. coli* via CD36 in a

clathrin-dependent and actin-polymerized manner in macrophages

PHrodo-labeled *E. coli* is a type of model bacteria that emits fluorescence only in

phagocytic cells because of the acidic environment. To clarify the regulatory effect of

DHZD on the phagocytotic ability, RAW264.7 cells were stimulated by LPS and DHZD

(15 µM) for 24 h. PHrodo-labeled *E. coli* was added and incubated for 1 h in dark. The

amount of engulfed *E. coli* in the cytoplasm was measured by Flow cytometry. DHZD

significantly increased the phagocytosis of pHrodo-labeled *E. coli* in RAW264.7 (Fig.

6A-C), suggesting that DHZD could promote the ability of bacterial phagocytosis of

macrophages.

The initiation of phagocytosis depends on the recognition and binding of host cell

receptors to the corresponding ligands on the surface of the pathogens. Innate immune

cells such as macrophages, neutrophils and dendritic cells can phagocyte pathogens

through caveolin-mediated endocytosis, clathrin-mediated endocytosis, pinocytosis

and other ways (23, 24). All the above-mentioned phagocytotic mechanisms are actin

dependent. Filipin III, CPZ and cytochalasin D are inhibitors of caveolin, clathrin and

actin, respectively. Filipin III had no effects on DHZD-promoted phagocytosis of

bacteria in RAW264.7 cells. CPZ could partially inhibit DHZD-promoted phagocytosis.

Cytochalasin D has the ability of preventing actin polymerization, and could completely

prevent the enhanced phagocytosis by DHZD (Fig. 6D-F).

Bacterial phagocytosis is usually mediated by phagocytic receptors. The

expression of phagocytic receptors CD36, FcγR2a, CD206, Dectin-1, CR3 and CR4

were detected by qRT-PCR on the surface of macrophages. From the expression screen,

DHZD could promote the expression of CD36 (Fig. 6G-H), at least partially accounting

for the increased ability of phagocytosis.

Figure 6. DHZD enhanced the phagocytosis of pHrodo-labelled *E. coli* in a clathrin-dependent

and actin-polymerized manner and also increased the expression of CD36 in RAW264.7 cells.

RAW264.7 cells were plated at 1×10^5 cells/100 µL in 96-well plates and incubated overnight. Then

cells were primed with LPS (100 ng/mL) along with DHZD (15 µM). After 24 h, macrophages were

incubated with pHrodo-labelled *E. coli* protected from light for 1 h. Cells were washed with sterile

PBS for three times to get rid of the bacteria outside of the cells. Then the phagocytosis of pHrodo-

labelled *E. coli* was quantified by FACS. One representative scatter plot out of three independent

experiments were shown (A). The mean fluorescence intensity (MFI) of pHrodo-labelled *E. coli*

swallowed by RAW264.7 cells were analyzed (B, C). RAW264.7 cells (1×10^5 cells/100 µL) were

seeded in 96-well plates and cultured overnight. Cells were pretreated with filipin III (3 µM), Cyto

D (10 µM) or CPZ (10 µM) for 30 min. Then cells were co-cultured with LPS (100 ng/mL) and

DHZD (15 µM) for 24 h (D-F). RAW264.7 cells were seeded (2×10^5/well) in 24-well plates and

incubated overnight. Cells were treated with HK-MRSA (MOI=10) separately along with DHZD

(15 µM) for 12 and 18 h. The expression of CD36 (G, H) were detected by PCR. Data were shown

as mean ± SD from three independent experiments; *, $P <0.05$, **, $P <0.01$, ***, $P <0.001$, and

****, $P <0.0001$.

DHZD inhibited LPS-stimulated activation of PI3K/Akt/p70S6K pathway

423   Our results have demonstrated that DHZD effectively reduced LPS-induced

424   inflammatory response. The results showed that DHZD inhibited LPS-induced

425   phosphorylation of Akt at Ser473 and Thr308 and subsequent p70S6K at Thr389.

426   Meanwhile, DHZD had no effects on the activation of ERK and p38 MAPK signaling

427   pathways (Fig. 7A-E).

Figure 7. DHZD inhibited LPS-stimulation of PI3K/Akt activation. RAW264.7 cells (1×10^6 cells/well) were seeded in 6-well plates overnight and stimulated with LPS (100 ng/mL) with or without DHZD (15 μM) for a certain period of time. Total protein of cells was detected by Western blot (A-E). Phospho-Akt (Ser 473), phospho-Gsk-3β (Ser9), phospho-ERK, phospho-p38, phospho-IκBα (Ser32/36), phospho-p65 (Ser536) and β-Actin in RAW264.7 were verified by Western blot. Data were shown as mean ± SD from three independent experiments; *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, and ****, $P < 0.0001$.

DHZD reversed hypothermia, tissue damage, and over-activated inflammation caused by CRKP

LPS is the main component of the cell wall of G⁻ bacteria. CRKP is one of the most troublesome G⁻ bacteria causing severe infection and sepsis in emergency department or intensive care unit (ICU). Fever is usually one of the most important symptoms of infection (25). There are variable thermoregulatory responses in sepsis, including fever and hypothermia (26). According to the analysis of clinical data, the mortality rate of septic patients with hypothermia is much higher than those with fever (27).

In order to clarify the protective effects of DHZD on CRKP-infected mice, a septic mouse model was established by the injection of CRKP intraperitoneally, which is sensitive to LVX. LVX was used as a positive control antibiotic in our mouse model. Infrared thermal imaging thermometer were also used to monitor body temperature of mice before and after CRKP challenge. Compared with the model group, DHZD-L group restored the average body temperature by 1.3 °C, and DHZD-H group restored the average body temperature by 3.1 °C. The results suggested that DHZD had an impressive protective effect on the recovery of body temperature (Fig. 8A-B). Compared with the PBS group, the lung tissue of mice in the CRKP group showed obvious infiltration of inflammatory cells, a large amount of fluid exudation in the alveoli, and pulmonary interstitial thickening. Inflammatory cell infiltration, swelling and necrosis of hepatocytes, fragmentation and lysis of some nuclei, and destruction of hepatic plate structure were observed in the liver. DHZD ameliorated lung and liver

tissue damage. A pathological score of lung tissue, as measured by pulmonary edema,

alveolar infiltration, and pulmonary vasculitis, showed a protective effect of DHZD

administration against pathological damage to the tissue (Fig. 8C-D).

In vitro stimulation of primary peritoneal macrophages of mice with HK-CRKP

showed that DHZD could reduce the secretion of IL-6, TNF-α, MCP-1, IL-1β and IL-

10 in a dose-dependent manner, but had no obvious influence on MIP-2 secretion (Fig.

8E-J).

Figure 8. DHZD reversed hypothermia, tissue damage, and inflammation caused by CRKP.

C57BL/6J female mice were randomly divided into 6 groups (10 mice/group). All groups of mice

were challenged by lethal dose of CRKP and corresponding agents, respectively. Temperature

difference (A, B). H&E staining of lung and liver (200×) (C). Scale bar 100 μm, and the histological

core of lung (D). Mouse primary macrophages were plated (3.5×10^5 cells/500 μL) in 24-well plates

at the day before stimulated. Cell culture supernatants were collected at 12 and 24 h after co-

stimulation of different concentrations of DHZD and HK-CRKP (MOI=10), and the concentrations

of IL-6 (E), TNF-α (F), MCP-1 (G), IL-1β (H), IL-10 (I) and MIP-2 (J) were measured by ELISA.

Data were shown as mean ± SD from three independent experiments. *, $P<0.05$, **, $P<0.01$, ***,

$P<0.001$, and ****, $P<0.0001$.

Discussion

AMR is an increasingly difficult problem for human beings, and it is also the main

cause accounting for the high mortality rate of septic patients. After bacterial infection,

the invading bacteria encounter host innate immune system at the beginning, and innate

cells "sense" the pathogens by recognizing pathogen-associated molecular patterns

(PAMPs) via pattern recognition receptors (PRRs). In most cases, the innate immune

response can help to eliminate the invading bacteria. When the bacteria prevail in host,

the immune response may become unbalanced (28, 29), further triggering the

overwhelming production of cytokines, such as TNF-α, IL-1β, IL-6, etc. (30). Then, the

excessive inflammatory response promotes the activation of the coagulation system and

the formation of microthrombosis in blood vessels, leading to disseminated

intravascular coagulation (DIC) (31, 32).

"Cytokine storm" is the main cause of death in septic patients with multiple organ

failure. The levels of inflammatory cytokines IL-1β, IL-6, IL-12 and IL-17 are usually

high in septic patients. In a mouse model of cecal ligation and puncture, the

administration of anti-IL-6 antibody reduced the expression of C5a and improved the

survival of the mice (33). In a study of community-acquired pneumonia, the mean

serum concentration of IL-6 from severe septic patients was 54.1 pg/mL, compared

with 36.5 pg /mL from patients who do not develop severe sepsis (34). Similarly, the

serum levels of IL-1β in deceased septic patients were about 2-10 times higher than those in surviving septic patients. These clinical data suggested that high concentrations of IL-6 or IL-1β may be closely related to poor prognosis (35). TNF-α is not only involved in cytokine storm but also responsible for the induction of hypothermia and high mortality in sepsis (36). In a study of *E. coli*-induced sepsis in baboons, treatment with anti-TNF antibody at 2 h before infection protected vital organ damage and reduced mortality (37). From the clinical experience, the downregulation of the excessive cytokine expression can alleviate tissue damage, reduce the conversion rate from mild to severe diseases, and thus improve the survival status of sepsis. DHZD could inhibit the over-activated inflammatory response (IL-6, TNF-α, IL-1β, IL-12, IFN-β and IL-10, etc.) in RAW264.7 cells, mouse primary peritoneal macrophages and BMDMs after the stimulation of bacteria or bacterial mimic. In the early stage of sepsis, DCs could also produce a series of inflammatory factors (such as IL-6, TNF-α, IL-12 and chemokines) after bacterial phagocytosis (38). Elevated IL-12 is also found during sepsis, which induces the differentiation of naive T cells into helper T cell (Th1) and the activation of NK cells, contributing to the formation of cytokine storm (39). DHZD downregulated the production of IL-6, TNF-α and IL12p70 in DCs, suggesting that it may also down-regulate cytokine storm via impairing subsequent T cell over-activation.

In LPS-induced septic mouse model, DHZD showed an impressive anti-inflammatory role and alleviated lung, liver and kidney damage, improving the survival rate. Both of the examined concentrations of DHZD showed protective effects *in vivo*. The survival rate of LPS + DHZD + DXM group was 100 %, which was better than

that of LPS + DXM group (60 %), suggesting that the combination of DHZD and DXM may have a potential synergistic protective effect to treat sepsis.

All of TNF-α, interleukin, and interferon γ can induce hypothermia via regulating thermotaxic center of the brain (40, 41). During infection, the activity of animals is reduced, which can also lead to the appearance of hypothermia (42). In sepsis, compared with patients in the fever group, patients in the hypothermia group had central nervous system dysfunction (88 % vs 60 %), elevated serum bilirubin concentration (35 % vs 15 %), prolonged prothrombin time (50 % vs 23 %), shock (94 % vs 61 %), nonrecovery from shock (66 % vs 26 %), and death (62 % vs 26 %) (43). In our current study, DHZD protected the loss of body temperature in CRKP-induced septic mouse model (-5.7 °C vs -2.6 °C) compared to those of CRKP group mice, hinting that DHZD-treated infective mice have better prognosis.

The load of pathogens is usually related to the severity of infectious diseases. Macrophage is a ubiquitous phagocytic cell type of innate immune system, and is involved in defensive inflammatory response and pathogen clearance during bacterial infection (44, 45). Besides macrophages, neutrophils and DCs are also important phagocytic cells in the body, which can phagocytize bacteria and decrease bacterial load. In our study, macrophages were used as a cell model to investigate whether DHZD has an ability to influence the cellular phagocytosis. DHZD was shown to promote the phagocytic ability of bacteria of macrophages in a clathrin-and actin-dependent manner. CD36, a member of class B scavenger receptor family, is expressed on cells such as monocytes, macrophages, endothelial cells, and adipocytes (34). The up-regulation of

CD36 expression by DHZD, might at least partially provide explanations for the

enhanced phagocytosis of macrophages. Not only the expression of phagocytic

receptors of phagocytes, but also the proportions and numbers of neutrophils and

monocytes/macrophages in lungs were increased by DHZD after LPS challenge,

suggesting that DHZD might contribute to a quick control of bacterial load at early

stage of infection through phagocytosis. Accordingly, DHZD might play a protective

role during bacterial infection and decrease organ damage.

As a member of innate immune cells, DC plays an important role on the line of

defense against pathogen. Immature DCs have a strong phagocytic ability. As it matures,

the phagocytic ability of DC is decreased and the expression of MHC II molecules and

co-stimulatory molecules (CD86, CD80 and CD40) are increased, accompanied by the

increased antigen presenting ability to evoke T cell activation (46-48). The down-

regulated expression of co-stimulatory molecules (CD80, CD86 and CD40) and MHC

II by DHZD keeps DC in an immature status, and might maintain its strong phagocytic

ability of bacteria and preventing the over-activation of T cells.

The PI3K/Akt signaling pathway also acts as an important regulator of cytokine

production after TLR4 ligation with LPS during G^{-} bacterial infection (49). Previous

studies have shown that the inhibition of PI3K/Akt activation in DCs or macrophages

reduces the secretion of pro-inflammatory factors (IL-6, TNF-α, IL-12 and IL-1β, etc.)

while promotes the secretion of the anti-inflammatory factor IL-10 in TLR2 and TLR4

signaling (50, 51). DHZD inhibited the activation of PI3K/Akt/p70S6K signaling

pathway, but had no influence on the activation of ERK and p38 MAPK signaling

pathways, accounting for its anti-inflammatory role.

In conclusion, DHZD has an obvious anti-inflammatory effect *in vitro* and *in vivo*

via targeting the host immune response. It may be a potential therapeutic drug for the

treatment of sepsis caused by bacterial infection and even drug-resistant bacterial

infection. Whether the phagocytic ability of bacteria besides *E. coli* could also be

increased by DHZD needs further experiments. Interestingly, the combination of

DHZD and DXM is expected to improve the survival rate of septic mice, hinting that

DHZD might be a potential adjuvant drug candidate for the treatment of sepsis.

Acknowledgments

This work was supported by grants from National Key Research and Development

Program of China (2021YFE0200900), National Natural Science Foundation of China

(82073901, 82073726), Shanghai Municipal Science and Technology Major Project

(ZD2021CY001), the Key Research and Development Program of Shaanxi Province

(2020SF-226), three-year Action Plan for Shanghai TCM Development and Inheritance

Program [ZY (2021-2023)-0401 and ZY (2021-2023)-0103]. Traditional Chinese

Medicine Research Project of Shanghai Municipal Health Commission (2022QN043).

We are thankful for Dr. Yi-jian Chen from Huashan Hospital affiliated to Fudan

University to provide Carbapenem-resistant *Klebsiella pneumoniae* (CRKP, HS11286).

Thank Ali Mollahassani for language polishing of this paper.

Author contribution

Yue-juan Zheng designed the experiments. Xiang-yang Qin modified and provided

DHZD. Ying-xiang Zou, Tian-nan Xiang, Huan Zhang, Yu-he Ma, Xiao Wu, Qi-lin

Huang, Biao Lei and Jing-wen Mu performed the laboratory assays. Ying-xiang Zou,

Tian-nan Xiang, Huan Zhang and Yu-he Ma analyzed the data. Yue-juan Zheng and

Ying-xiang Zou wrote the manuscript. Tian-nan Xiang and Xiang-yang Qin contributed

to manuscript revision. All authors read and approved the manuscript.

Supplementary information

Figure S1. Plot of the spectral data of DHZD.

Figure S2. DHZD decreased the secretion of LPS- induced IL-6, TNF-α, MCP-1 and IL-10 in

BMDMs. Tibias and femurs were obtained from C57BL/6 mice and differentiated into bone marrow

derived macrophages (BMDMs) by the addition of L929 conditional media for 5 constitutive days.

Cells were seeded (3.5×10^5 cells/well) in 24-well plates and then stimulated by LPS (100 ng/mL)

and DHZD for 6 h. The secretion of IL-6 (A), TNF-α (B) and IL-10 (C) were detected by ELISA.

Data were shown as mean ± SD from three independent experiments. *, P <0.05, **, P <0.01, and

***, P <0.001.

Figure S3. DHZD deceased the expression of inflammatory response in lung of LPS-induced

acute peritonitis mice. C57BL/6J mice were randomly divided into 5 groups (9 mice/group). All

groups of mice were stimulated with LPS (10 mg/kg) and interfered with DXM (5 mg/kg), DHZD

(10 mg/kg, 40 mg/kg) and their combination according to the groups. After 12 hours of intervention,

the mice were sacrificed. The mRNA expression of IL-6 (A), MCP-1(B) and MIP-2 (C) in lung

tissues were detected by qRT-PCR. Data were shown as mean ± SD of three mice per group. *, P

<0.05, and ***, P <0.001.

Reference:

1. Mu A, Klare WP, Baines SL, Ignatius Pang CN, Guérillot R, Harbison‑Price N, Keller N,
Wilksch J, Nhu NTK, Phan MD, Keller B, Nijagal B, Tull D, Dayalan S, Chua HHC, Skoneczny
D, Koval J, Hachani A, Shah AD, Neha N, Jadhav S, Partridge SR, Cork AJ, Peters K, Bertolla
O, Brouwer S, Hancock SJ, Álvarez‑Fraga L, De Oliveira DMP, Forde B, Dale A,
Mujchariyakul W, Walsh CJ, Monk I, Fitzgerald A, Lum M, Correa-Ospina C, Roy
Chowdhury P, Parton RG, De Voss J, Beckett J, Monty F, McKinnon J, Song X, Stephen JR,
Everest M, Bellgard MI, Tinning M, Leeming M, Hocking D, et al. 2023. Integrative omics
identifies conserved and pathogen-specific responses of sepsis-causing bacteria. Nat

Commun 14:1530.

2. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, Bellomo
R, Bernard GR, Chiche JD, Coopersmith CM, Hotchkiss RS, Levy MM, Marshall JC, Martin
GS, Opal SM, Rubenfeld GD, van der Poll T, Vincent JL, Angus DC. 2016. The Third
International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). Jama 315:801-
10.

3. Gersh, B. JJYoC. 2011. Population Trends in the Incidence and Outcomes of Acute
Myocardial Infarction.

4. Siegel R, Ma J, Zou Z, Jemal A. 2014. Cancer statistics, 2014. CA Cancer J Clin 64:9-29.

5. Fleischmann C, Scherag A, Adhikari NK, Hartog CS, Tsaganos T, Schlattmann P, Angus DC,
Reinhart K. 2016. Assessment of Global Incidence and Mortality of Hospital-treated Sepsis.
Current Estimates and Limitations. Am J Respir Crit Care Med 193:259-72.

6. Adhikari NK, Fowler RA, Bhagwanjee S, Rubenfeld GD. 2010. Critical care and the global
burden of critical illness in adults. Lancet 376:1339-46.

7. Compston J. 2018. Glucocorticoid-induced osteoporosis: an update. Endocrine 61:7-16.

8. Cutolo M, Seriolo B, Pizzorni C, Secchi ME, Soldano S, Paolino S, Montagna P, Sulli A. 2008.
Use of glucocorticoids and risk of infections. Autoimmun Rev 8:153-5.

9. Savva A, Roger T. 2013. Targeting toll-like receptors: promising therapeutic strategies for
the management of sepsis-associated pathology and infectious diseases. Front Immunol
4:387.

10. O'Neill LA, Bowie AG. 2007. The family of five: TIR-domain-containing adaptors in Toll-
like receptor signalling. Nat Rev Immunol 7:353-64.

11. Schumann J. 2016. It is all about fluidity: Fatty acids and macrophage phagocytosis. Eur J
Pharmacol 785:18-23.

12. Groves E, Dart AE, Covarelli V, Caron E. 2008. Molecular mechanisms of phagocytic uptake
in mammalian cells. Cell Mol Life Sci 65:1957-76.

13. Fridman WH. 1991. Fc receptors and immunoglobulin binding factors. Faseb j 5:2684-90.

14. Ninković J, Roy S. 2012. Morphine decreases bacterial phagocytosis by inhibiting actin
polymerization through cAMP-, Rac-1-, and p38 MAPK-dependent mechanisms. Am J
Pathol 180:1068-1079.

15. An H, Hou J, Zhou J, Zhao W, Xu H, Zheng Y, Yu Y, Liu S, Cao X. 2008. Phosphatase SHP-
1 promotes TLR- and RIG-I-activated production of type I interferon by inhibiting the
kinase IRAK1. Nat Immunol 9:542-50.

16. Zheng Y, An H, Yao M, Hou J, Yu Y, Feng G, Cao X. 2010. Scaffolding adaptor protein
Gab1 is required for TLR3/4- and RIG-I-mediated production of proinflammatory
cytokines and type I IFN in macrophages. J Immunol 184:6447-56.

17. Han C, Jin J, Xu S, Liu H, Li N, Cao X. 2010. Integrin CD11b negatively regulates TLR-
triggered inflammatory responses by activating Syk and promoting degradation of
MyD88 and TRIF via Cbl-b. Nat Immunol 11:734-42.

18. Franco-Molina MA, Mendoza-Gamboa E, Castillo-León L, Tamez-Guerra RS, Rodríguez-
Padilla C. 2004. Bovine dialyzable leukocyte extract protects against LPS-induced, murine
endotoxic shock. Int Immunopharmacol 4:1577-86.

19. Zheng Y, Guo Z, He W, Yang Y, Li Y, Zheng A, Li P, Zhang Y, Ma J, Wen M, Yang M, An H,
Ji G, Yu Y. 2012. Ephedrine hydrochloride protects mice from LPS challenge by promoting

IL-10 secretion and inhibiting proinflammatory cytokines. Int Immunopharmacol 13:46-
53.

20. Jiang X, Chen Y, Liu D, Shi T, Cheng X, He W, Li Y, Ryffel B, Zheng SG, Zheng Y. 2020.
Secoeudesma sesquiterpenes lactone A alleviates inflammation and offers adjuvant
protection in severe infection of carbapenem-resistant Klebsiella pneumoniae. J
Ethnopharmacol 252:112605.

21. Zhang AJ, Lee AC, Chan JF, Liu F, Li C, Chen Y, Chu H, Lau SY, Wang P, Chan CC, Poon VK,
Yuan S, To KK, Chen H, Yuen KY. 2021. Coinfection by Severe Acute Respiratory Syndrome
Coronavirus 2 and Influenza A(H1N1)pdm09 Virus Enhances the Severity of Pneumonia
in Golden Syrian Hamsters. Clin Infect Dis 72:e978-e992.

22. Ando M, Isogai K, Azami H, Hirata N, Yamazaki H, Yanagi Y. 1992. Studies on the synthesis
of sesquiterpene lactones, 14. Syntheses of (-)-arbusclin D and (+)-4-EPI-arbusclin D: the
stereochemical assignment of arbusclin D. J Nat Prod 55:1620-5.

23. Doherty GJ, McMahon HT. 2009. Mechanisms of endocytosis. Annu Rev Biochem 78:857-
902.

24. Ramanan V, Agrawal NJ, Liu J, Engles S, Toy R, Radhakrishnan R. 2011. Systems biology
and physical biology of clathrin-mediated endocytosis. Integr Biol (Camb) 3:803-15.

25. Walter EJ, Hanna-Jumma S, Carraretto M, Forni L. 2016. The pathophysiological basis and
consequences of fever. Crit Care 20:200.

26. Thomas-Rüddel DO, Hoffmann P, Schwarzkopf D, Scheer C, Bach F, Komann M, Gerlach
H, Weiss M, Lindner M, Rüddel H, Simon P, Kuhn SO, Wetzker R, Bauer M, Reinhart K,
Bloos F. 2021. Fever and hypothermia represent two populations of sepsis patients and
are associated with outside temperature. Crit Care 25:368.

27. Rumbus Z, Matics R, Hegyi P, Zsiboras C, Szabo I, Illes A, Petervari E, Balasko M, Marta K,
Miko A, Parniczky A, Tenk J, Rostas I, Solymar M, Garami A. 2017. Fever Is Associated with
Reduced, Hypothermia with Increased Mortality in Septic Patients: A Meta-Analysis of
Clinical Trials. PLoS One 12:e0170152.

28. van der Poll T, van de Veerdonk FL, Scicluna BP, Netea MG. 2017. The immunopathology
of sepsis and potential therapeutic targets. Nat Rev Immunol 17:407-420.

29. Takeuchi O, Akira S. 2010. Pattern recognition receptors and inflammation. Cell 140:805-
20.

30. Kang R, Chen R, Zhang Q, Hou W, Wu S, Cao L, Huang J, Yu Y, Fan XG, Yan Z, Sun X,
Wang H, Wang Q, Tsung A, Billiar TR, Zeh HJ, 3rd, Lotze MT, Tang D. 2014. HMGB1 in
health and disease. Mol Aspects Med 40:1-116.

31. Opal SM. 2011. Immunologic alterations and the pathogenesis of organ failure in the ICU.
Semin Respir Crit Care Med 32:569-80.

32. Tang D, Wang H, Billiar TR, Kroemer G, Kang R. 2021. Emerging mechanisms of
immunocoagulation in sepsis and septic shock. Trends Immunol 42:508-522.

33. van der Poll T, Levi M, Hack CE, ten Cate H, van Deventer SJ, Eerenberg AJ, de Groot ER,
Jansen J, Gallati H, Büller HR, et al. 1994. Elimination of interleukin 6 attenuates
coagulation activation in experimental endotoxemia in chimpanzees. J Exp Med
179:1253-9.

34. Wu HP, Chen CK, Chung K, Tseng JC, Hua CC, Liu YC, Chuang DY, Yang CH. 2009. Serial
cytokine levels in patients with severe sepsis. Inflamm Res 58:385-93.

35.　Mera S, Tatulescu D, Cismaru C, Bondor C, Slavcovici A, Zanc V, Carstina D, Oltean M.
2011. Multiplex cytokine profiling in patients with sepsis. Apmis 119:155-63.

36.　Leon LR, White AA, Kluger MJ. 1998. Role of IL-6 and TNF in thermoregulation and survival
during sepsis in mice. Am J Physiol 275:R269-77.

37.　Tracey KJ, Fong Y, Hesse DG, Manogue KR, Lee AT, Kuo GC, Lowry SF, Cerami A. 1987.
Anti-cachectin/TNF monoclonal antibodies prevent septic shock during lethal
bacteraemia. Nature 330:662-4.

38.　Bouras M, Asehnoune K, Roquilly A. 2018. Contribution of Dendritic Cell Responses to
Sepsis-Induced Immunosuppression and to Susceptibility to Secondary Pneumonia. Front
Immunol 9:2590.

39.　Kilbourn RG, Griffith OW, Gross SS. 1993. Pathogenetic mechanisms of septic shock. N
Engl J Med 329:1427-8.

40.　Leon LR. 2004. Hypothermia in systemic inflammation: role of cytokines. Front Biosci
9:1877-88.

41.　Lang GP, Ndongson-Dongmo B, Lajqi T, Brodhun M, Han Y, Wetzker R, Frasch MG, Bauer
R. 2020. Impact of ambient temperature on inflammation-induced encephalopathy in
endotoxemic mice-role of phosphoinositide 3-kinase gamma. J Neuroinflammation
17:292.

42.　Jhaveri KA, Trammell RA, Toth LA. 2007. Effect of environmental temperature on sleep,
locomotor activity, core body temperature and immune responses of C57BL/6J mice.
Brain Behav Immun 21:975-87.

43.　Clemmer TP, Fisher CJ, Jr., Bone RC, Slotman GJ, Metz CA, Thomas FO. 1992. Hypothermia
in the sepsis syndrome and clinical outcome. The Methylprednisolone Severe Sepsis Study
Group. Crit Care Med 20:1395-401.

44.　Gentek R, Molawi K, Sieweke MH. 2014. Tissue macrophage identity and self-renewal.
Immunol Rev 262:56-73.

45.　Yan J, Horng T. 2020. Lipid Metabolism in Regulation of Macrophage Functions. Trends
Cell Biol 30:979-989.

46.　Darkwah S, Nago N, Appiah MG, Myint PK, Kawamoto E, Shimaoka M, Park EJ. 2019.
Differential Roles of Dendritic Cells in Expanding CD4 T Cells in Sepsis. Biomedicines 7.

47.　Rescigno M, Granucci F, Ricciardi-Castagnoli P. 1999. Dendritic cells at the end of the
millennium. Immunol Cell Biol 77:404-10.

48.　Lipscomb MF, Masten BJ. 2002. Dendritic cells: immune regulators in health and disease.
Physiol Rev 82:97-130.

49.　Liu Y, Pan YF, Xue YQ, Fang LK, Guo XH, Guo X, Liu M, Mo BY, Yang MR, Liu F, Wu YT,
Olsen N, Zheng SG. 2018. uPAR promotes tumor-like biologic behaviors of fibroblast-like
synoviocytes through PI3K/Akt signaling pathway in patients with rheumatoid arthritis.
Cell Mol Immunol 15:171-181.

50.　Stark AK, Sriskantharajah S, Hessel EM, Okkenhaug K. 2015. PI3K inhibitors in inflammation,
autoimmunity and cancer. Curr Opin Pharmacol 23:82-91.

51.　Re F, Strominger JL. 2001. Toll-like receptor 2 (TLR2) and TLR4 differentially activate
human dendritic cells. J Biol Chem 276:37692-9.

745