Germline *cis* variant determines epigenetic regulation of the anti-cancer drug metabolism gene dihydropyrimidine dehydrogenase (*DPYD*)

Ting Zhang [1], Alisa Ambrodji [2,3], Huixing Huang [1], Kelly J. Bouchonville [1], Amy S. Etheridge [4], Remington E. Schmidt [1], Brianna M. Bembenek [1], Zoey B. Temesgen [1], Zhiquan Wang [5], Federico Innocenti [4], Deborah Stroka [6], Robert B. Diasio [1], Carlo R. Largiadèr [2], and Steven M. Offer [1,7,8,*]

Department of Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic, Rochester, MN 55905, USA.

Department of Clinical Chemistry, Inselspital, Bern University Hospital, University of Bern, CH-3010 Bern, Switzerland.

Graduate School for Cellular and Biomedical Sciences, University of Bern, Freiestrasse 1, CH-3010 Bern, Switzerland.

Eshelman School of Pharmacy, Division of Pharmacotherapy and Experimental Therapeutics, University of North Carolina, Chapel Hill, NC 27599, USA.

Division of Hematology, Department of Medicine, Mayo Clinic, Rochester, MN 55905 USA.

Department of Visceral Surgery and Medicine, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland.

Department of Pathology, Carver College of Medicine, University of Iowa, Iowa City, IA 52242, USA.

Lead contact.

* Correspondence: soffer@uiowa.edu; offer.steven1@mayo.edu

ABSTRACT

Enhancers are critical for regulating tissue-specific gene expression, and genetic

variants within enhancer regions have been suggested to contribute to various cancer-

related processes, including therapeutic resistance. However, the precise mechanisms

remain elusive. Using a well-defined drug-gene pair, we identified an enhancer region

for dihydropyrimidine dehydrogenase (DPD, *DPYD* gene) expression that is relevant to

the metabolism of the anti-cancer drug 5-fluorouracil (5-FU). Using reporter systems,

CRISPR genome edited cell models, and human liver specimens, we demonstrated *in*

vitro and *vivo* that genotype status for the common germline variant (rs4294451; 27%

global minor allele frequency) located within this novel enhancer controls *DPYD*

transcription and alters resistance to 5-FU. The variant genotype increases recruitment

of the transcription factor CEBPB to the enhancer and alters the level of direct

interactions between the enhancer and *DPYD* promoter. Our data provide insight into

the regulatory mechanisms controlling sensitivity and resistance to 5-FU.

KEYWORDS

Epigenetics, Gene Regulation, 5-Fluorouracil, Chemotherapy, CEBPB, Chemotherapy,

Biomarkers, Drug Resistance.

INTRODUCTION

Therapeutic resistance has been reported for nearly all anti-cancer drugs [1], and as many as 90% of mortalities in cancer can be linked to drug resistance [2]. Enhancer-mediated regulation of gene expression has been increasingly implicated in multiple cancer-related processes, including resistance, response, and toxicity to cancer therapies [3, 4]. However, the specific molecular mechanisms through which epigenetic changes drive these processes remain mostly elusive.

Enhancers regulate gene expression by recruiting transcription factors (TFs) and subsequently interacting with the promoter region of a target gene to drive expression. The activity of enhancers can be affected by localized epigenetic state and genetic variations that affect TF binding. More than 90% of disease-associated genetic variants identified in genome-wide association studies (GWAS) lie in the non-coding portion of the genome [5, 6]. Many of these GWAS variants map to putative enhancers [7]. Regardless, translating enhancer GWAS variants into disease mechanisms remains a largely unmet challenge [8].

In previous studies, we identified a *cis* expression quantitative trait locus (eQTL) for the chemotherapeutic metabolism gene dihydropyrimidine dehydrogenase enzyme (DPD, encoded by the *DPYD* gene) [9]. DPD is the initial and rate-defining enzyme for conversion of the commonly used chemotherapeutic 5-fluoruracil (5-FU) into inactive metabolites. Hepatic DPD eliminates approximately 85% of circulating 5-FU within

minutes of administration [10, 11], and deficiency of DPD is associated with severe

(clinical grade ≥3) toxicity to 5-FU [12]. Clinical and preclinical studies have identified

missense and splicing variants of *DPYD* that consistently correlate with reduced DPD

function and increased 5-FU toxicity risk [12-18]. However, these variants explain <20%

of reported cases of grade ≥3 5-FU toxicities [12, 19], and the mechanisms contributing

to 5-FU resistance, and therefore means to overcome resistance, remain elusive.

In the present study, we used human liver tissues and cellular models to identify and

characterize a novel *cis* enhancer element capable of modulating *DPYD* expression.

We additionally provide mechanistic data demonstrating that transcription-factor driven

expression of *DPYD* is dependent on allele status for a common germline genetic

variant located within this enhancer, suggesting that the variant could be a valuable

biomarker of 5-FU toxicity risk. Furthermore, the genotype-dependent regulation of DPD

expression offers a potential mechanism for overcoming resistance in patients carrying

the risk variant.

RESULTS

Identification of *DPYD* enhancer regions

Candidate *cis* enhancer regions for *DPYD* (NM_000110.4) were identified using

GeneHancer [20] and Ensembl Regulatory Build [21] data mapped to GRCh37/hg19.

We defined the upstream boundary of the target region using enhancer data supported

by multiple methods and gene-enhancer links supported by multiple methods as

implemented within GeneHancer (i.e., putative enhancer regions with "double-elite"

status; **Figure 1A**). We then mapped annotated transcription factor binding sites in the

target region using ChIP-seq data from the Ensembl Regulatory Build as an additional

layer of evidence (**Figure 1A**). In all, three candidate proximal upstream enhancer

regions were identified, which will be referred to herein as E9, E16, and E20 based on

the relevant location upstream of the *DPYD* transcription start site (**Figure 1A**).

CRISPR-interference (CRISPRi) and -activation (CRISPRa) were used to determine

which of these regions were capable of regulating *DPYD* expression. For CRISPRi, we

generated cell lines that stably expressed the Krüppel-associated box (KRAB) domain

of Kox1 fused to nuclease-deficient Cas9 protein (dCas9-KRAB; [22]). For CRISPRa,

we created cell lines that stably expressed dCas9 fused to the histone acetyltransferase

E1A binding protein P300 (dCas9-P300; [23]). Six guide RNAs (gRNAs) were used to

target dCas9 fusion proteins to each region of interest, and quantitative reverse

transcription PCR (qRT-PCR) was used to monitor changes in *DPYD* expression

following transfection.

Targeting of dCas9-KRAB (CRISPRi) to the E9 region significantly decreased *DPYD*

expression relative to control for four of six gRNAs (**Figure 1B**). Targeting of dCas9-

KRAB to E16 resulted in a significant decrease in *DPYD* expression for one of six

gRNAs (**Figure 1C**), and no significant reductions in *DPYD* expression were noted

following transfection of gRNAs for E20 in dCas9-expressing cells; however, two gRNAs

increased *DPYD* expression (**Figure 1D**). For reciprocal CRISPRa experiments,

targeting dCas9-P300 to E9 significantly increased *DPYD* expression for four of six

gRNAs tested (**Figure 1E**), whereas gRNAs specific to E16 and E20 did not elicit any

significant changes in *DPYD* expression (**Figures 1F** and **1G**). Consistent with the

results in HepG2 cells, CRISPRi and CRISPRa targeted to the E9 region significantly

altered *DPYD* expression in HCT116 cells (**Figures S1A** and **S1D**), whereas no

significant changes were noted when targeted to E16 or E20 (**Figures S1B**, **S1C**, **S1E**,

and **S1F**).

To confirm that the expected epigenetic changes were induced at the E9 region, we

performed chromatin immunoprecipitation (ChIP) coupled with quantitative PCR (ChIP-

qPCR) in dCas9-KRAB and -P300 expressing cells transfected with E9 gRNAs. As

expected, targeting dCas9-KRAB to E9 resulted in a localized increase in H3K9me3

compared to controls, indicating a shift from active to inactive chromatin at E9 (**Figure**

2A; the primer nearest E9 is denoted by the gray box). Targeting of dCas-P300 to E9

caused changes consistent with epigenetic activation of the region, including a localized

gain of H3K27ac-marked chromatin (**Figure 2B**). Similar changes in localized histone

modifications were noted when dCas9-KRAB and dCas9-P300 HCT116 cells were

directed to E9 (**Figure 2C–D**). Collectively, these results indicate that the E9 region

might act as a *cis* regulatory element modulating *DPYD* expression.

Identification of rs4294451 as a putative regulatory SNP within E9

In an expression quantitative trait loci (eQTL) genome-wide association study (GWAS)

conducted in human liver specimens, Etheridge *et al.* previously identified three

independent haplotype blocks that significantly associated with altered *DPYD*

expression [9]. One of the blocks identified in that study spanned the E9 region,

prompting us to investigate if genetic variants in E9 could potentially perturb *DPYD*

regulation. Based on the biological role of enhancers, we postulated that a causal

functional single nucleotide polymorphism (SNP) could interfere with transcription factor

binding. Within the Ensembl Regulatory Build [21], we identified a region of

approximately 150 bp showing evidence for binding by multiple transcription factors in

HepG2 cells, but not in other cell types tested (**Figure S2**). Within this region, we

identified the variant rs4294451, which is a tag SNP for the previously identified eQTL

block [9], prompting us to hypothesize that this variant might be the causal eQTL SNP.

To assess the contribution of rs4294451 alleles to *DPYD* regulation, we first performed

reporter assays by cloning the *DPYD* promoter and E9 enhancer region containing

either the reference or variant allele for rs4294451 (T or A, respectively) into a luciferase

reporter vector (**Figure 3A**). Reporter vectors containing the T-allele yielded significantly

higher luciferase activity compared to vectors containing only the *DPYD* promoter

(p=0.00054), whereas those containing the A allele within E9 showed a more modest

increase (p=0.070, **Figure 3B**). Directly comparing E9-containing reporters, vectors

Page 7

containing the T allele showed significantly higher activity than those with the A allele

(p=0.046, **Figure 3B**), suggesting that the variant could impact *DPYD* expression.

Impact of rs4294451 genotype on *DPYD* expression and regulation

To directly characterize the impact of rs4294451 alleles on *DPYD* expression at the

endogenous locus, we used CRISPR-mediated genome editing to create matched

isogenic HCT116 knock-in cell models for each genotype. Consistent with the results

from the reporter assay (**Figure 3B**), cells homozygous for the A allele had significantly

lower *DPYD* expression compared to cells that were homozygous for the T allele

(p=0.0011, **Figure 3C**). Heterozygous rs4294451 A/T cells displayed intermediate

expression compared to cells with homozygous A/A and T/T genotypes (p=0.00013 and

p=0.028, respectively; **Figure 3C**). At the chromatin level, cells homozygous for the A

allele showed reductions in H3K27ac and accumulation of H3K9me at E9 compared to

cells that were homozygous or heterozygous for the T allele (**Figure 3D–3E**). This

finding is consistent with the A allele being associated with a less active epigenetic state

at E9. Cells carrying a heterozygous genotype for rs4294451 displayed an intermediate

epigenetic state, which is consistent with one epigenetically active and one inactive

copy of E9 (**Figure 3D–3E**). Similar epigenetic differences between rs4294451

genotypes were noted at E9 in liver specimens obtained from human donors (**Figure**

S3A–S3B).

Allele-specific expression of *DPYD* transcripts

To precisely determine the impact of rs4294451 alleles on *DPYD* expression *in vivo*, we

measured allele-specific expression using reverse transcriptase digital droplet PCR

(RT-ddPCR) using donor liver tissues. Because rs4294451 falls outside of the coding

region, we used coding region variants as proxies to measure strand-specific

expression. These coding variants consisted of *DPYD*-c.85T>C (rs1801265) and

c.496A>G (rs2297595), both of which are in linkage disequilibrium (LD) with the

rs4294451 T allele (c.85C: D′=0.92, R^2=0.80; c.496G: D′= 0.67, R^2= 0.21). Imperfect LD

between rs4294451 and proxies allowed us to use samples that are heterozygous for

the coding region SNP, but homozygous for rs429441 genotype, as an additional level

of control. In samples measured using the c.85 genotype, the fractional abundance of

the C-allele ranged from 0.542 to 0.602 in samples heterozygous for rs4294451 (**Figure**

3F). Expressed in terms of relative expression, the C-allele was expressed at levels 18–

51% higher than the T-allele (i.e., 0.542/0.458=1.18 and 0.602/0.398=1.51). In contrast,

the sample that was homozygous for the rs4294451 T allele showed allelic expression

at the c.85 locus that was indistinguishable from the DNA control (**Figure 3F**), indicating

equal expression of both transcripts. In measurements using c.496, a significant allelic-

imbalance of 0.565 (CI: 0.546–0.584) in favor of the variant c.496 G-allele compared to

the DNA control (CI: 0.488–0.511) was observed (**Figure 3G**). This again indicated

higher expression of the linked rs4294451 T allele. Notably, liver specimens that were

homozygous at the rs4294451 locus again showed c.496 allelic expression that was

indistinguishable from the DNA control (**Figure 3G**). These data indicate that rs4294451

significantly affects the expression of *DPYD* in the liver.

Page 9

Rs4294451 genotype affects interactions between the *DPYD* promoter and E9

We next investigated the impact of rs4294451 genotype on intra-chromatin interactions

between the E9 region and the *DPYD* promoter using quantitative analysis of chromatin

conformation capture (3C-qPCR) in knock-in cells. The schematic in **Figure 4A** shows

the location of primers used for 3C-qPCR relative to the *DPYD* transcription start site

(TSS) and E9 region, as well as the location of *Hin*dIII digestion sites. When using

primers anchored at the *DPYD* promoter, the E9 region showed a significantly stronger

interaction with the promoter in cells carrying the T allele compared to those

homozygous for the A allele ($p=8.9 \times 10^{-4}$ comparing TT to AA; ANOVA $p=3.0 \times 10^{-4}$

across all three genotypes; **Figure 4C**). With primers anchored to the E9 region, cells

homozygous for the rs4294451 T allele showed stronger promoter interaction than

those carrying the A allele ($p=4.9 \times 10^{-4}$ comparing TT to AA; ANOVA $p=1.2 \times 10^{-4}$

across all three genotypes; **Figure 4D**).

Rs4294451 genotype confers differential sensitivity to 5-FU

Having demonstrated that the rs4294451 genotype impacts *DPYD* expression, we next

evaluated the impact on 5-FU sensitivity using real-time cellular analysis (RTCA). RTCA

provides a measure of the number of viable cells present in a culture. We have

previously demonstrated the utility of this technology for measuring differences in drug-

sensitivity to 5-FU and that those differences directly correlate with DPD enzyme

function within cells [14, 17]. Consistent with lower expression of *DPYD*, knock-in cells

homozygous for the rs4294451 A allele were significantly more sensitive to 5-FU than

cells homozygous for the T allele (IC_{50} concentrations were 9.1 and 95.0 µM 5-FU,

respectively; p<0.0001; **Figure 4E**). Heterozygous T/A cells showed an intermediate

IC_{50} value of 29.9 μM 5-FU (**Figure 4E**).

Transcription factor binding within E9 is affected by rs4294451 status

Based on ENCODE data, rs4294451 is localized to a region that previously has been

shown to be bound by multiple transcription factors (**Figure S2**). To determine if known

transcription factor binding sites could be affected by allele status at rs4294451, we

computed binding scores for JUND, MYBL2, HNF4A, CEBPB, FOXA1, and FOXA2 for

sequences comprising the E9 region containing the T and A alleles at rs4294451.

CEBPB, FOXA1, and FOXA2 showed differential predicted binding scores between the

rs4294451 A- and T-containing query sequences (data not shown). For CEBPB, the

differential binding site had a higher score than other sites in the queried region that did

not overlap with rs4294451, suggesting that the SNP could affect CEBPB binding in the

region. While predicted binding in the region varied for FOXA1 and FOXA2 depending

on rs4294451 genotype, other binding sites that did not overlap with the variant site

showed stronger predicted binding, indicating that the SNP was likely not affecting the

critical binding site in the region. Binding scores did not differ by genotype for JUND or

HNF4A, and no binding site above the threshold score was detected in the region for

MYBL2.

Page 11

Based on the above, we sought to determine if CEBPB could regulate *DPYD*

expression through the E9 region and if regulation was affected by rs4294451

genotype. We first used reporter assays in conjunction with CEBPB. CEBPB expression

significantly increased reporter activity relative to GFP control for luciferase vectors

containing the *DPYD* promoter, indicating that the promoter contains CEBPB

recognition sites (p=9.1×10^{-4}; **Figure 5A**). In cells overexpressing CEBPB, plasmids

containing the promoter and E9 region resulted in higher luciferase activity compared to

those containing only the *DPYD* promoter, regardless of the presence of the A or T

allele (p=0.0036 and p=0.0017, respectively), suggesting that rs4294451 genotype does

not completely disrupt CEBPB regulation through E9 (**Figure 5A**). Comparing results for

the A and T alleles at rs4294451 in the presence of overexpressed CEBPB indicates

that the T allele is likely more responsive to CEBPB (**Figure 5A**; p=0.036).

To directly examine CEBPB binding to the E9 region and to determine if binding affinity

differs between the rs4294451 A and T alleles, we performed ChIP with CEBPB

antibodies in isogenic knock-in HCT116 cells for the rs4294451 A/A, A/T, and T/T

genotypes. Cells homozygous for the reference T allele showed significantly higher

CEBPB occupancy at both E9 and the *DPYD* promoter than A/T or A/A cells (**Figure**

5B–5C). These results suggest that rs4294451 genotype determined the binding

potential for CEBPB at the E9 enhancer region, which, in turn, affects CEBPB-driven

expression of *DPYD* from the promoter.

Upregulation of *DPYD* expression by CEBPB is dependent on the rs4294451-T allele

To further characterize the role of CEBPB in regulating *DPYD*, we disrupted CEBPB expression using two independent shRNAs (denoted as "sh1" and "sh2") in HCT116 knock-in cells carrying rs4294451 A/A, A/T, and T/T genotypes. Knockdown of CEBPB was confirmed at the protein (**Figure 6A**) and mRNA (**Figures 6B–6D**) levels. Knockdown of CEBPB significantly reduced *DPYD* expression in rs4294451 T/A and T/T cells, but not in rs4294451 A/A cells (**Figures 6E–6F**). CEBPB knockdown also reduced occupancy of the transcription factor at both E9 and the *DPYD* TSS in rs4294451 A/T and T/T cells, but not A/A cells, and the level of CEBPB occupancy at both regions under CEBPB knockdown conditions is similar to that in control shRNA-treated A/A cells (**Figures S4A–S4B**).

We next determined the extent to which CEBPB contributed to 5-FU chemoresistance in cells with the T allele of rs4294451 (**Figure 4E**). In rs4294451 A/A cells, knockdown of CEBPB did not affect 5-FU sensitivity (**Figure 6H**). In contrast, CEBPB knockdown significantly reduced the IC_{50} for 5-FU in rs4294451 T/T cells (p<0.0001; **Figure 6I**).

The rs4294451-T allele is enriched in individuals of African ancestry

Data from the Genome Aggregation Database (gnomAD [24]) were used to estimate the frequency of the rs4294451-T allele in global populations (**Figure S5A**). The highest minor allele frequency (MAF) was noted for African/African American individuals (40%

Page 13

MAF), where the MAF was lowest in East Asian individuals (7%). For comparison,

individuals of European (Non-Finnish) ancestry, the population with the highest number

of individuals reported in gnomAD, had a MAF of 23%. Within Latino-Admixed American

gnomAD subjects, similar differences in MAFs were noted in local ancestry-informed

frequency data (**Figure S5B**).

DISCUSSION

The antitumor efficacy and risk of severe adverse events associated with 5-FU are

determined by the overall bioavailability of the drug in plasma. As the critical

determinant of 5-FU pharmacokinetics, liver DPD expression is pivotal to both the risk of

severe adverse events and therapeutic resistance in 5-FU chemotherapy at opposite

ends of the exposure spectrum. This is underscored by the narrow therapeutic window

for 5-FU, with toxicity and efficacy occurring at partially overlapping drug exposure

levels [25]. Deleterious germline coding-region *DPYD* variants have been linked to

severe 5-FU toxicity [12, 13]; however, these variants are responsible for only a small

fraction of severe adverse events in 5-FU use and are unlikely to contribute to drug

resistance [26]. Elevated expression of DPD in tumor cells is known to confer 5-FU

resistance [27, 28], and upregulation of hepatic DPD expression has been shown to

reduce drug exposures and promote the development of 5-FU–resistant tumors [29].

However, the mechanisms regulating DPD expression are not well characterized, nor is

it understood how the regulatory processes can be altered to support the development

of 5-FU resistance.

In the current study, we identified a novel *cis* enhancer region for *DPYD* that is located

approximately 9 kb upstream from the gene's transcription start site. We demonstrated

that CEBPB is a critical transcription factor for *DPYD* that binds to this enhancer region,

termed E9, promoting enhancer–promoter interactions and increasing *DPYD*

expression. We also showed that the allele status of the germline variant rs4294451,

located within the E9 region, can affect CEBPB-driven *DPYD* expression and

sensitivity/resistance to 5-FU, making it a strong candidate biomarker for 5-FU toxicity

risk and potentially tumor resistance to 5-FU–based cancer therapy. These findings are

consistent with the recent identification of a haplotype block linked to the rs4294451 T

allele that was significantly associated with elevated *DPYD* expression in human liver

tissues [9]. In the present manuscript, we show that the rs4294451 T allele is enriched

for active chromatin marks in both human liver (**Figure S3**) and in cellular knock-in

models (**Figure 3D–E**). Furthermore, using an innovative digital droplet RT-qPCR–

based approach, we demonstrate that the rs4294451 T allele is associated with

elevated expression of the *cis DPYD* transcript in human liver (**Figure 3F–G**).

Allele frequency data retrieved from gnomAD suggest that a majority (65-70%,

estimated from allele frequencies) of individuals of African ancestry carry the

rs4294451-T allele (**Figure S5**), whereas only about 35% of individuals of European

Page 15

ancestry are predicted to be carriers of the T allele. African American patients have

worse overall survival in colorectal cancer compared to white patients, owing to

biological and non-biological factors. While differential access to healthcare, treatment

bias, and socioeconomic factors have been shown to contribute to the poorer prognosis

[30], other unrecognized factors also contribute to this difference [31, 32]. Our data

support a hypothesis that higher systemic 5-FU catabolism to due elevated liver DPD

expression in carriers of the rs4294451-T allele results in lower exposure to active anti-

tumor metabolites of 5-FU. Additionally, we demonstrate that colorectal cancer cell lines

likely retain the enhancer functions associated with the rs4294451-T allele, suggesting

that tumor cells carrying this variant could more readily inactive 5-FU via increased DPD

expression. The higher likelihood of carrying the T-allele in individuals of African

ancestry would therefore place them at greater risk. Additional studies will be needed to

investigate the degree to which rs4294451-T contributes to survival and progression in

colorectal cancer and in other solid cancers frequently treated with 5-FU.

The transcription factor CEBPB is a member of the CCAAT Enhancer Binding Protein

family, a group of transcription factors that contain basic leucine zipper (bZIP) domains

and is highly expressed in liver [33, 34]. Multiple CEBPB isoforms have been detected,

with some acting as transcriptional activators and others as inhibitors. The data

presented herein indicate that active isoforms of CEBPB are up-regulating *DPYD*

through binding to the E9 enhancer region. Inhibitory isoforms of CEBPB have also

been shown to be important for certain physiological processes including tumorigenesis

and liver regeneration [35, 36]. Our over-expression studies used the full-length active

isoform. Therefore, we cannot rule out a role for the inactive isoform participating in the

regulation of *DPYD* expression. In addition to acting as a homodimer, CEBPB can also

heterodimerize with other CEBP family proteins and interact with other transcription

factors, including P300/CBP, CREB, NFKB, AP1, and NFAT, to co-regulate gene

expression [37, 38]. Additional studies are underway to characterize the role of

additional regulatory factors within the *DPYD* enhancer region identified in this

manuscript.

In silico analyses suggested that the rs4294451 A allele created a stronger binding site

for CEBPB within the E9 enhancer region. However, our data demonstrate that the T

allele is associated with higher reporter activity (**Figure 3B**), higher DPD expression in

both knock-in cells (**Figure 3C**) and human liver tissues (**Figure 3F–G**), and localized

epigenetic activation (**Figure 3D–E**). Enrichment of CEBPB to the E9 region was

likewise shown to be higher in cells with the rs4294451 T allele (**Figure 5**), and higher

levels of interaction were also noted between the E9 region with the T allele and the

DPYD promoter (**Figure 4**). These findings are also consistent with previously reported

eQTL results for the haplotype block linked to rs4294451, where the T allele was

associated with higher levels of *DPYD* expression in human liver specimens [9], and our

observation that the T allele is associated with cellular resistance to 5-FU (**Figure 4E**).

The binding of CEBPB to specific motifs has recently been shown to be cell-type

specific and to rely on non-consensus binding motifs with binding strengths that can be

modulated by the sequence and structure of surrounding DNA regions [39, 40],

providing a possible explanation for the discrepancy between predicted and observed

results.

To our knowledge, this is the first report directly linking CEBPB to 5-FU metabolism and

the first mechanistic data demonstrating *cis* effects of a regulatory variant on *DPYD*

expression. The role of CEBPB in modulating 5-FU resistance and toxicity is not without

precedent. For example, mir-191 is abnormally expressed in several cancers and has

been associated with both 5-FU resistance and the regulation of CEBPB expression

[41]; however, the CEBPB–*DPYD* regulatory axis has not previously been recognized.

Furthermore, CEBPB signaling has been shown to be activated in colorectal cancer

cells following treatment with 5-FU [42], suggesting that CEBPB-mediated activation of

DPYD expression might represent a dynamic response to therapy.

While the contributions of regulatory variants to 5-FU metabolism have not been widely

studied to date, previous studies have explored DPD regulation. Our laboratories

previously characterized a *trans*-acting regulatory variant for DPD located within the

microRNA mir-27a that was subsequently shown to further increase 5-FU toxicity risk in

individuals that carried deleterious nonsynonymous *DPYD* variants [43, 44]. The variant

at rs4294451 is in LD with the *DPYD* variants c.85T>C and c.496G>A, which served as

coding region proxies for allele-specific expression in our present study (**Figure 3F–G**).

The haplotypes defined by these two coding-region variants together with a third variant

(c.1129-5923C>G/rs75017182) previously associated with varied levels of systemic

DPD activity [45]. A subsequent retrospective analysis indicated that these haplotype

differences translate to differential risk of severe 5-FU toxicity [46]; however, additional

studies are needed that are powered to evaluate more than the most common

haplotypes. Taken together, these results suggest that the differential regulatory effects

of rs4294451 alleles could further impact overall DPD activity and, by extension,

modulate the risk of severe 5-FU–related toxicity conferred by coding or splice-variants

in *DPYD*. Further work is also needed to identify interactions between the linked

variants that impact DPD enzyme activity and the regulatory variant to define systemic

DPD function and 5-FU toxicity risk.

METHODS

Cells

HEK293T, HCT116, and HepG2 cell lines were obtained from the American Tissue

Collection Center (ATCC, Manassas, VA). All cell lines were cultured in Dulbecco's

modified Eagle's medium (DMEM; Corning, Corning, NY) supplemented with 10% fetal

bovine serum (FBS; MilliporeSigma, Billerica, MA), 2 mM L-glutamate (Corning), and 1x

penicillin/streptomycin solution (MilliporeSigma). Cells were maintained at 37°C with 5%

CO_2. To support cell attachment and expansion, HepG2 cells were grown on plates

coated with 5% Matrigel (Corning).

Liver tissues

Human liver tissues used for ChIP analyses were processed through Dr. Mary Relling's

laboratory at St. Jude Children's Research Hospital, part of the Pharmacogenetics of

Anticancer Agents Research (PAAR) Group and were provided by the Liver Tissue Cell

Distribution System funded by NIH Contract #N01-DK-7-0004/HHSN267200700004C

and by the Cooperative Human Tissue Network. The acquisition and use of specimens

for this manuscript was conducted with the approval of the University of North Carolina

at Chapel Hill IRB (study number 10-2253), which has designated the use of these

livers for the current analyses as nonhuman subject research and the need for direct

consent for use in this study was waived. Human liver tissues used for allele-specific

expression were obtained from the University Clinic of Visceral Surgery and Medicine,

Inselspital, Bern, Switzerland. Specimens were from donated material from patients who

had undergone liver surgery at Inselspital and signed a written consent form for remnant

tissues to be used in research (KEK-BE:2016-02202). Patients with impaired liver

function due to cirrhosis or other conditions were excluded from analyses.

Quantitative RT-PCR (RT-qPCR)

Total RNA was extracted from cells using the Direct-zol RNA Kit (Zymo Research,

Tustin, CA) following the manufacturer's protocol. Reverse transcription into cDNA was

performed using the Transcriptor Reverse Transcriptase kit (Roche, Indianapolis, IN)

and random hexamer primers (Roche) according to the manufacturer's instructions.

Quantitative PCR (qPCR) was performed on a LightCycler 480 System (Roche) using

LightCycler 480 SYBR Green I Master Mix reagents (Roche). Primers used for RT-

qPCR are listed in **Supplementary Table S1**. RNA expression was normalized to the

reference gene L32, and relative gene expression was calculated using the $2^{-\Delta\Delta CT}$

method. For all analyses, three independent experiments were performed; for each

experiment, gene expression was assessed in triplicate.

Western blot analysis

Whole-cell lysates were extracted using the RIPA lysis buffer system (Santa Cruz

Biotechnology, Dallas, TX), separated by SDS-PAGE, and transferred to PVDF

membrane (MilliporeSigma). Blots were blocked using Odyssey Blocking Buffer (LI-

COR Biosciences, Lincoln, NE) and incubated with primary antibody at 4°C overnight.

Primary antibodies consisted of anti-CEBPB (PA5-27244; 1:1,000 dilution; Thermo

Fisher Scientific, Waltham, MA), anti-alpha Tubulin (ab4074; 1:7,500 dilution; Abcam,

Waltham, MA), and anti-cas9 (sc-517386; 1:1,000 dilution; Santa Cruz Biotechnology,

Dallas, TX). Membranes were washed and incubated with anti-mouse and anti-rabbit

secondary antibodies (#926-32212 and #926-68073; both 1:5,000 dilution; LI-COR

Biosciences) for 1 hour at room temperature. Blots were imaged using the Odyssey

infrared imaging system (LI-COR Biosciences).

Plasmids

The gRNA expression empty vector lentiGuide-Puro was a gift from Feng Zhang

(Addgene_52963). pCMV-FLAG LAP2 was a gift from Joan Massague

(Addgene_15738). The oligonucleotides targeting enhancer regions (E9, E16, and E20)

were designed using GuideScan [47], hybridized, phosphorylated, and cloned into

lentiGuide-Puro via *Bsm*BI sites. The luciferase expression vector pGL4.10 was

purchased from Promega (Madison, WI). Lentivirus vectors expressing shRNAs

targeting CEBPB were obtained from the University of Minnesota Genomics Center

(Sh1: TRCN0000007440 and Sh2: TRCN0000007442).

CRISPR inactivation (CRISPRi) and activation (CRISPRa)

HepG2 cell lines that overexpress dCas9-KRAB and dCas9-P300 were generated by

lentiviral transduction using lenti-EF1a-dCas9-KRAB-Puro and pLV-dCas9-p300-P2A-

PuroR, respectively. To generate lentiviral particles, HEK293T cells were co-

transfected with lenti-EF1a-dCas9-KRAB-Puro plasmid (a gift from Kristen Brennand;

Addgene_99372) or pLV-dCas9-p300-P2A-PuroR plasmid (a gift from Charles

Gersbach; Addgene_83889), psPAX2 (a gift from Didier Trono; Addgene_12260), and

pMD2.G (a gift from Didier Trono; Addgene_12259) using TransIT-Lenti Transfection

Reagent (Mirus Bio, Madison, WI). A 3:1 ratio of transfection reagent to total plasmid

was used for all transfections. For all transfections, medium was changed 14 hours after

transfection, and viral supernatants were collected 34 hours later. Supernatants were

filtered using 0.45 μm PVDF filters (MilliporeSigma) to remove debris/cells and used

directly for transductions. For transductions, HCT116 cells or HepG2 cells were seeded

at a density of 4×10^5 cells per well in 6-well plates and incubated with 500 µL virus-

containing supernatant, 12.5 µg/mL polybrene (MilliporeSigma), and 1.5 mL fresh

DMEM culture medium. Medium was changed after 24 hours. Cells were treated with 1

µg/mL puromycin to initiate selection for transduced cells 48 hours after transduction.

Expression of dCas9-KRAB and dCas9-P300 was confirmed by western blotting. Guide

RNAs (gRNAs) for each target region were identified and designed using GuideScan

[47]. Oligonucleotides corresponding to each gRNA were obtained from IDT (Coralville,

IA), hybridized, phosphorylated using T4 polynucleotide kinase (New England Biolabs,

Ipswich, MA), and ligated into digested BsmBI-digested (New England Biolabs)

lentiGuide-Puro vector (a gift from Feng Zhang; Addgene_52963). Cell lines stably

expressing dCas9-KRAB or dCas9-P300 were transfected with plasmids encoding

gRNAs or lentiGuide empty vector using TransIT-X2 (Mirus Bio). RNA was extracted 2

days after transfection, and *DPYD* expression was measured by RT-qPCR.

Chromatin immunoprecipitation coupled with quantitative PCR (ChIP-qPCR)

ChIP assays were performed using the ChIP-IT Express Enzymatic Kit (Active Motif,

Carlsbad, CA) following manufacturer's directions. One million cells were harvested,

washed, and crossed linked using 1% formaldehyde (Thermo Fisher Scientific) in

serum-free medium for 10 minutes followed by quenching with 125 mM glycine for 5

minutes at room temperature. The chromatin was digested using enzymatic shearing

cocktail provided by the kit to an average size of 200–1000 bp. Two percent of the

sheared chromatin was retained as input control. Approximately 25 µg sheared

chromatin was incubated with 2 μg H3K27ac antibody (ab4729; Abcam), 2 μg

H3K9me3 antibody (ab8898; Abcam), 2 μg CEBPB antibody (PA5-27244; Thermo

Fisher Scientific), or 2 μg control normal Rabbit IgG antibody (antibody 2729; Cell

Signaling Technology, Danvers, MA), in the presence of protein G magnetic beads,

ChIP buffer, and protease inhibitor cocktail (all Active Motif) for 4 hours at 4°C. Magnetic

beads were washed, chromatin was eluted, cross-linking was reversed, and proteinase

K treatment performed using reagents provided in the kit following manufacturer's

directions. DNA was purified using the QIAquick PCR Purification Kit (Qiagen,

Germantown, MD). Purified DNA was used for subsequent qPCR reactions using SYBR

Green I Master Mix on a LightCycler 480 System (Roche). Enrichment was calculated

using the following formula: (a) % ChIP = $2^{(Input\ Ct\ -\ ChIP\ Ct)}$ * (dilution factor) (100); (b) %

IgG = $2^{(Input\ Ct\ -\ IgG\ Ct)}$ * (dilution factor) (100); (c) Fold Enrichment = % ChIP ÷ % IgG.

Primers used for ChIP-qPCR are listed in **Supplementary Table S1**.

Luciferase reporter assays

The *DPYD* promoter region, consisting of the 1154 bp of genomic DNA directly

upstream of the *DPYD* TSS, was amplified by PCR, digested with EcoRV and HindIII

(New England Biolabs), and cloned into compatible sites on the pGL4.10 vector

(Promega). The 1392-bp region comprising the E9 region was PCR amplified from

genomic DNA and cloned upstream of the *DPYD* promoter using KpnI and SacI sites

(New England Biolabs). A vector containing the A allele of rs4294451 within E9 was

confirmed by Sanger sequencing. The vector containing the rs4294451 T allele was

generated by site-directed mutagenesis and confirmed by sequencing. All primers used

in vector construction are listed in **Supplementary Table S1**. For reporter assays, 10^5

HEK293T cells were seeded into 24-well plates and co-transfected with pGL4.10-based

plasmids and pRL-SV40 Renilla luciferase plasmid (Promega). After 48 hours,

luciferase activity was measured using the Dual-Glo Luciferase Assay (Promega)

following manufacturer's recommendations on a Synergy HTX Multimode Plate Reader

(Agilent Technologies, Santa Clara, CA).

Knock-in cell lines for rs4294451 genotypes

Knock-in cell lines were generated using CRISPR/Cas9 gene editing. Homology-

directed repair (HDR) donor templates and target-specific Alt-R crRNA were designed

using the Alt-R HDR Design Tool (IDT). Equimolar amounts of crRNA (IDT) and

common Alt-R tracrRNA (IDT) were annealed to form the gRNA duplex. RNP

complexes were formed by combining gRNA with Alt-R S.p. Cas9 Nuclease V3 (IDT) to

a final Cas9:gRNA ratio of 4:4.8. RNA complex and Alt-R HDR Donor Oligos were

transfected into HCT116 cells by electroporation (Lonza Nucleofector 96-well Shuttle

System; Lonza, Bend, OR) using parameters provided by the manufacturer. Seventy-

two hours after transfection, serial dilutions were performed to obtain single-cell clones.

Clones were expanded, genomic DNA was isolated, and rs42944551 genotype was

determined by rhAmp Genotyping (assay ID: Hs.GT.rs4294451.A.1; IDT) using rhAmp

Genotyping Master Mix and universal probe Reporter Mix (both IDT).

Allele-specific gene expression

Allele-specific expression of *DPYD* was measured using reverse transcriptase droplet

digital PCR (RT-ddPCR) by targeting variants in the coding region of *DPYD* (c.85T>C

and c.496A>G). Linkage disequilibrium (LD) between variants was calculated using

LDpair implemented within LDlink [48]. DNA was extracted from donor liver tissues

using the QIAamp DNA Mini Kit (Qiagen). For RNA extraction, tissues were lysed in

QIAzol (Qiagen), and RNA was extracted using the miRneasy Kit (Qiagen) with on-

column DNA digestion using RNase-free DNase (Qiagen). RNA quality was assessed

using an Agilent 2100 Bioanalyzer running 2100 Expert Software v.B.02.10 using

Agilent RNA 6000 Nano kits.

DNA samples were genotyped for c.85, c.496, and rs4294451 loci using TaqMan SNP

Genotyping assays (Applied Biosystems, Waltham, MA). Tissues that were

heterozygous for at least one of the coding region SNPs (i.e., c.85T>C and/or

c.496A>G) were suitable for allele-specific expression analysis because expression

from both alleles could be discriminated using the coding-region markers in mRNA.

Allele-specific expression was measured using RT-ddPCR with the One-Step RT-

ddPCR Advanced Kit for Probes (Qiagen) on a QX200 ddPCR Droplet Reader (BioRad,

Hercules, CA). Fractional abundance was calculated using QuantaSoft software

(BioRad). Poisson distributions were determined using Quantasoft and were used to

define 95% confidence intervals. To address possible biases associated with differing

probe efficacies caused by differential probe binding affinities or specificities, fractional

abundances are also reported relative to those measured in DNA.

Quantitative analysis of chromatin conformation capture (3C-qPCR)

Quantitative analysis of chromatin conformation capture (3C-qPCR) was performed as

described by Hagère *et al.* [49] with minor modifications. Cells were trypsinized using

0.25% w/v trypsin-EDTA (Thermo Fisher Scientific) and resuspended in 1% FBS

(MilliporeSigma) in DMEM (Corning) for counting by flow cytometry using a NovoCyte

3000 RYB system (Agilent Technologies). Cells (10^7) were pelleted by centrifugation at

$300 \times g$ at 22°C for 5 minutes. Supernatant was discarded, and cell pellets were

resuspended in 500 μL of 10% FBS in DMEM. Single-cell suspensions were obtained

by filtration through a 40-μm cell strainer (Corning). Crosslinking was performed by

adding 9.5 mL of 1% formaldehyde in 10% FBS in PBS per 10^7 cells. Reactions were

incubated at 22°C while tumbling for 10 minutes. Reactions were transferred to ice and

crosslinking was quenched by the addition of ice-cold glycine (MilliporeSigma) to

achieve a final concentration of 0.125M. Samples were centrifuged at $300 \times g$ at 4°C for

5 minutes. Supernatant was removed and discarded. Crosslinked cell pellets were lysed

in 5 mL cold lysis buffer (10 mM Tris-HCl, pH7.5; 10 mM NaCl; 5 mM MgCl; 0.1 mM

EGTA; 1x Roche Complete protease inhibitor cocktail). Lysis was allowed to proceed for

10 minutes on ice with intermittent gentle pipetting to obtain homogeneous suspensions

of nuclei. Nuclei were pelleted at $500 \times g$ at 4°C for 5 minutes and resuspended in 0.5

mL of 1.2x NEBuffer r2.1 (New England Biolabs) containing 3% sodium dodecyl sulfate

(SDS, MilliporeSigma). Samples were incubated at 37°C for 1 hour with shaking.

Following incubation, Triton X-100 (MilliporeSigma) was added to a final concentration

of 2%. Reactions were incubated at 37°C for 1 hour with shaking. To digest DNA, 600 U

of HindIII (New England Biolabs) was added, and reactions were incubated for 16 hours

at 37°C with shaking. To inactivate digestion, SDS was added to a final concentration of

1.6%, and reactions were incubated at 65°C for 25 minutes with shaking. Excess SDS

was sequestered by the addition of Triton X-100 to a final concentration of 1%. Samples

were divided into two aliquots, one for ligation and the other for non-ligation control.

Ligation was performed on a 7-fold dilution of HindIII-digested chromatin using 100 units

of Quick T4 DNA ligase (New England Biolabs) at 16°C for 16 hours, followed by 1 hour

at 22°C. Proteinase K (300 μg; New England Biolabs) was added to ligation mixtures

and non-ligated controls, and samples were incubated at 65°C for 16 hours to reverse

crosslinking. DNA was subsequently purified by adding 300 μg RNase A (Thermo-

Fisher) followed by a 30-minute incubation at 37°C and subsequent phenol-chloroform

extraction as described [49]. DNA pellets were washed with 70% ethanol and dissolved

in 10 mM Tris pH 7.5.

For qPCR, directional primers were designed within each fragment as depicted in

Figure 4A. Two anchors, one localized to the fragment containing the *DPYD* promoter

and the other to the E9 region, were selected and paired with primers designed to

"walk" across the length of the surrounding region. Reactions were carried out in 10 μL

reaction volumes, consisting of 5 pmol each primer and 1 μL of a 1:50 dilution of each

3C sample. Amplification was performed using LightCycler 480 SYBR Green I Master

Mix (Roche) on a LightCycler 480 thermocycler (Roche). PCR conditions were 95°C for

10 minutes, and 40 cycles of 95°C for 15 seconds, 65°C for 1 minute, and 72°C for 15

seconds. Enrichment was calculated as $2^{-(cp\ (ligated\ DNA)\ -\ cp\ (non-ligated\ DNA))}$. Enrichment with

the fragment containing the nearest primer was used as the control for interaction

frequency for further normalization between replicate experiments. Specifically, when

primer 5 was used as the anchor, data were normalized to the average of values

obtained from using primers 5 and 4. When primer 7 was used as the anchor, data were

normalized to the average of values from primers 7 and 8. All experiments were

performed in triplicate. Primer sequences and positions relative to the *DPYD*

transcription start site (TSS) for qPCR are listed in **Supplementary Table S2**.

Cellular sensitivity to 5-FU

Cell viability was monitored using the xCELLigence MP Real-Time Cell Analysis (RTCA)

system (Agilent) as previously described [14]. Background impedance values for RTCA

E-View plates (Agilent) were obtained using complete DMEM prior to plating cells. Cells

were seeded at a density of 5,000 cells per well and incubated for 20 hours, at which

time medium was removed and replaced with medium containing serial dilutions of 5-FU

ranging from 1.25 µM to 100 µM. Impedance values (expressed as cell index, CI, units)

were recorded every 15 minutes over the course of the experiment to monitor

proliferation. Results represent the average of three independent cultures. To account

for minute differences in plating and potential cell loss during drug addition, relative CI

Page 29

was calculated as the CI measured 48 hours after 5-FU divided by the CI recorded

immediately after treatment. Four parameter logistic non-linear regression analysis was

used to determine IC_{50} concentrations (GraphPad Prism version 9, GraphPad Software,

San Diego, CA).

CEBPB knockdown cells

Lentiviral particles for CEBPB knockdown were generated by transfecting shRNA

plasmids (TRCN0000007440 and TRCN0000007442) or scramble shRNA control (a gift

from David Sabatini; Addgene_1864 [50]), psPAX2 (a gift from Didier Trono;

Addgene_12260), and pMD2.G (a gift from Didier Trono; Addgene_12259) into

HEK293T cells using TransIT-Lenti (Mirus Bio, Madison, WI). A 3:1 ratio of transfection

reagent to total plasmid was used for all transfections. For all transfections, medium

was changed 14 hours after transfection, and viral supernatants were collected 34

hours later. Supernatants were filtered using 0.45 μm PVDF filters (Millipore) to remove

debris/cells and used directly for transductions. For transductions, target cells were

seeded at a density of 4×10^5 cells per well in 6-well plates and incubated with 500 µL

virus-containing supernatant, 12.5 µg/mL polybrene (MilliporeSigma), and 1.5 mL fresh

DMEM culture medium.

Analysis of transcription factor binding motifs

Potential transcription factor binding sites within the E9 region were determined using

TFBSTools [51]. DNA sequence corresponding to the 101-nucleotide region centered

on rs4294451 was retrieved from genome build GRCh38.p13 (NC_000001.11). A

second DNA string was created to mimic the sequence corresponding to the rs4294451

A allele. Position frequency matrices for each transcription factor were retrieved from

JASPAR CORE 2022 [52] and converted to log-scale position weight matrices using the

toPWM method implemented in TFBSTools. JASPAR includes binding site information

from multiple sources. ENCODE data [53] were available for CEBPB (matrix ID

MA04661) and JUND (MA0491.1). In absence of ENCODE data, alternatives including

PAZAR [54] (HNF4A, MA0114.2; FOXA1, MA0148.3), REMAP [55] (FOXA2,

MA0047.3), and data from an individual publication [56] (MYBL2, MA0777.1) were used.

EP300 is a cofactor that does not recognize a specific DNA motif on its own; instead, it

interacts with various DNA-binding factors to modify chromatin and facilitate activation

of target genes. As such, P300 does not have a DNA-binding motif and was not

included in analysis. Nucleotide sequences were scanned using the patterns presented

in the position weight matrices to identify putative transcription factor binding sites.

Forward and reverse strands were searched, and the 70th percentile between the

minimum and maximum possible value for a matrix was used as the minimum threshold

score. Empirical p-values for each score were calculated by an exact method using the

TFMPvalue R package. R version 4.2.2 was used for analyses

Statistical analyses

Significance was defined as $p < 0.05$ unless otherwise noted in the text. Pairwise

comparisons were performed using unpaired two-tailed Student's t tests calculated

using GraphPad Prism version 9. One-way ANOVA statistics were calculated using

GraphPad Prism. Summary statistics pertaining to allele-specific expression were

calculated using Quantasoft software as described in the Allele-specific gene

expression section. Transcription factor binding predictions and associated analyses

were performed in R version 4.2.2 as described above.

Availability of data and materials

All data generated or analyzed during this study are included in this published article

and its supplementary information files.

Acknowledgements

This work was supported by the National Cancer Institute of the National Institutes of

Health under award number R01CA251065 (SMO, P.I.). Allele-specific expression

experiments were supported by the Swiss National Science Foundation under award

number 320030_212583 (CRL, P.I.). The funding bodies did not contribute to the design

of the study, the collection, analysis, and interpretation of data, or in writing the

manuscript.

REFERENCES

1. Ramos, A., S. Sadeghi, and H. Tabatabaeian, *Battling Chemoresistance in Cancer: Root Causes and Strategies to Uproot Them.* Int J Mol Sci, 2021. **22**(17).

2. Mansoori, B., A. Mohammadi, S. Davudian, S. Shirjang, and B. Baradaran, *The Different Mechanisms of Cancer Drug Resistance: A Brief Review.* Adv Pharm Bull, 2017. **7**(3): p. 339-348.

3. Koutsi, M.A., M. Pouliou, L. Champezou, G. Vatsellas, A.I. Giannopoulou, C. Piperi, and M. Agelopoulos, *Typical Enhancers, Super-Enhancers, and Cancers.* Cancers (Basel), 2022. **14**(18).

4. Lauschke, V.M., Y. Zhou, and M. Ingelman-Sundberg, *Novel genetic and epigenetic factors of importance for inter-individual differences in drug disposition, response and toxicity.* Pharmacol Ther, 2019. **197**: p. 122-152.

5. Manolio, T.A., F.S. Collins, N.J. Cox, D.B. Goldstein, L.A. Hindorff, D.J. Hunter, M.I. McCarthy, E.M. Ramos, L.R. Cardon, A. Chakravarti, J.H. Cho, A.E. Guttmacher, A. Kong, L. Kruglyak, E. Mardis, C.N. Rotimi, M. Slatkin, D. Valle, A.S. Whittemore, M. Boehnke, A.G. Clark, E.E. Eichler, G. Gibson, J.L. Haines, T.F. Mackay, S.A. McCarroll, and P.M. Visscher, *Finding the missing heritability of complex diseases.* Nature, 2009. **461**(7265): p. 747-53.

6. Maurano, M.T., R. Humbert, E. Rynes, R.E. Thurman, E. Haugen, H. Wang, A.P. Reynolds, R. Sandstrom, H. Qu, J. Brody, A. Shafer, F. Neri, K. Lee, T. Kutyavin, S. Stehling-Sun, A.K. Johnson, T.K. Canfield, E. Giste, M. Diegel, D. Bates, R.S. Hansen, S. Neph, P.J. Sabo, S. Heimfeld, A. Raubitschek, S. Ziegler, C. Cotsapas, N. Sotoodehnia, I. Glass, S.R. Sunyaev, R. Kaul, and J.A. Stamatoyannopoulos, *Systematic localization of common disease-associated variation in regulatory DNA.* Science, 2012. **337**(6099): p. 1190-5.

7. Boix, C.A., B.T. James, Y.P. Park, W. Meuleman, and M. Kellis, *Regulatory genomic circuitry of human disease loci by integrative epigenomics.* Nature, 2021. **590**(7845): p. 300-307.

8. Claringbould, A. and J.B. Zaugg, *Enhancers in disease: molecular basis and emerging treatment strategies.* Trends Mol Med, 2021. **27**(11): p. 1060-1073.

9. Etheridge, A.S., P.J. Gallins, D. Jima, K.A. Broadaway, M.J. Ratain, E. Schuetz, E. Schadt, A. Schroder, C. Molony, Y. Zhou, K.L. Mohlke, F.A. Wright, and F. Innocenti, *A New Liver Expression Quantitative Trait Locus Map From 1,183 Individuals Provides Evidence for Novel Expression Quantitative Trait Loci of*

Drug Response, Metabolic, and Sex-Biased Phenotypes. Clin Pharmacol Ther,
2019.

10. Sommadossi, J.P., D.A. Gewirtz, R.B. Diasio, C. Aubert, J.P. Cano, and I.D.
Goldman, *Rapid catabolism of 5-fluorouracil in freshly isolated rat hepatocytes as*
analyzed by high performance liquid chromatography. J Biol Chem, 1982.
257(14): p. 8171-6.

11. Heggie, G.D., J.P. Sommadossi, D.S. Cross, W.J. Huster, and R.B. Diasio,
Clinical pharmacokinetics of 5-fluorouracil and its metabolites in plasma, urine,
and bile. Cancer Res, 1987. **47**(8): p. 2203-6.

12. Amstutz, U., L.M. Henricks, S.M. Offer, J. Barbarino, J.H.M. Schellens, J.J.
Swen, T.E. Klein, H.L. McLeod, K.E. Caudle, R.B. Diasio, and M. Schwab,
Clinical Pharmacogenetics Implementation Consortium (CPIC) Guideline for
Dihydropyrimidine Dehydrogenase Genotype and Fluoropyrimidine Dosing: 2017
Update. Clin Pharmacol Ther, 2018. **103**(2): p. 210-216.

13. Lee, A.M., Q. Shi, E. Pavey, S.R. Alberts, D.J. Sargent, F.A. Sinicrope, J.L.
Berenberg, R.M. Goldberg, and R.B. Diasio, *DPYD variants as predictors of 5-*
fluorouracil toxicity in adjuvant colon cancer treatment (NCCTG N0147). J Natl
Cancer Inst, 2014. **106**(12).

14. Shrestha, S., C. Zhang, C.R. Jerde, Q. Nie, H. Li, S.M. Offer, and R.B. Diasio,
Gene-Specific Variant Classifier (DPYD-Varifier) to Identify Deleterious Alleles of
Dihydropyrimidine Dehydrogenase. Clin Pharmacol Ther, 2018. **104**(4): p. 709-
718.

15. Offer, S.M., C.C. Fossum, N.J. Wegner, A.J. Stuflesser, G.L. Butterfield, and
R.B. Diasio, *Comparative functional analysis of DPYD variants of potential*
clinical relevance to dihydropyrimidine dehydrogenase activity. Cancer Res,
2014. **74**(9): p. 2545-54.

16. Offer, S.M., A.M. Lee, L.K. Mattison, C. Fossum, N.J. Wegner, and R.B. Diasio,
A DPYD variant (Y186C) in individuals of african ancestry is associated with
reduced DPD enzyme activity. Clin Pharmacol Ther, 2013. **94**(1): p. 158-66.

17. Offer, S.M., N.J. Wegner, C. Fossum, K. Wang, and R.B. Diasio, *Phenotypic*
profiling of DPYD variations relevant to 5-fluorouracil sensitivity using real-time
cellular analysis and in vitro measurement of enzyme activity. Cancer Res, 2013.
73(6): p. 1958-68.

18. Nie, Q., S. Shrestha, E.E. Tapper, C.S. Trogstad-Isaacson, K.J. Bouchonville,
A.M. Lee, R. Wu, C.R. Jerde, Z. Wang, P.A. Kubica, S.M. Offer, and R.B. Diasio,
Quantitative Contribution of rs75017182 to Dihydropyrimidine Dehydrogenase

mRNA Splicing and Enzyme Activity. Clin Pharmacol Ther, 2017. **102**(4): p. 662-
670.

19.   Meulendijks, D., L.M. Henricks, G.S. Sonke, M.J. Deenen, T.K. Froehlich, U.
Amstutz, C.R. Largiader, B.A. Jennings, A.M. Marinaki, J.D. Sanderson, Z. Kleibl,
P. Kleiblova, M. Schwab, U.M. Zanger, C. Palles, I. Tomlinson, E. Gross, A.B.
van Kuilenburg, C.J. Punt, M. Koopman, J.H. Beijnen, A. Cats, and J.H.
Schellens, *Clinical relevance of DPYD variants c.1679T>G, c.1236G>A/HapB3,*
and c.1601G>A as predictors of severe fluoropyrimidine-associated toxicity: a
systematic review and meta-analysis of individual patient data. Lancet Oncol,
2015. **16**(16): p. 1639-50.

20.   Fishilevich, S., R. Nudel, N. Rappaport, R. Hadar, I. Plaschkes, T. Iny Stein, N.
Rosen, A. Kohn, M. Twik, M. Safran, D. Lancet, and D. Cohen, *GeneHancer:*
genome-wide integration of enhancers and target genes in GeneCards.
Database (Oxford), 2017. **2017**.

21.   Zerbino, D.R., S.P. Wilder, N. Johnson, T. Juettemann, and P.R. Flicek, *The*
ensembl regulatory build. Genome Biol, 2015. **16**: p. 56.

22.   Yeo, N.C., A. Chavez, A. Lance-Byrne, Y. Chan, D. Menn, D. Milanova, C.C.
Kuo, X. Guo, S. Sharma, A. Tung, R.J. Cecchi, M. Tuttle, S. Pradhan, E.T. Lim,
N. Davidsohn, M.R. Ebrahimkhani, J.J. Collins, N.E. Lewis, S. Kiani, and G.M.
Church, *An enhanced CRISPR repressor for targeted mammalian gene*
regulation. Nat Methods, 2018. **15**(8): p. 611-616.

23.   Hilton, I.B., A.M. D'Ippolito, C.M. Vockley, P.I. Thakore, G.E. Crawford, T.E.
Reddy, and C.A. Gersbach, *Epigenome editing by a CRISPR-Cas9-based*
acetyltransferase activates genes from promoters and enhancers. Nat
Biotechnol, 2015. **33**(5): p. 510-7.

24.   Chen, S., L.C. Francioli, J.K. Goodrich, R.L. Collins, M. Kanai, Q. Wang, J.
Alföldi, N.A. Watts, C. Vittal, L.D. Gauthier, T. Poterba, M.W. Wilson, Y.
Tarasova, W. Phu, M.T. Yohannes, Z. Koenig, Y. Farjoun, E. Banks, S. Donnelly,
S. Gabriel, N. Gupta, S. Ferriera, C. Tolonen, S. Novod, L. Bergelson, D.
Roazen, V. Ruano-Rubio, M. Covarrubias, C. Llanwarne, N. Petrillo, G. Wade, T.
Jeandet, R. Munshi, K. Tibbetts, g.P. Consortium, A. O'Donnell-Luria, M.
Solomonson, C. Seed, A.R. Martin, M.E. Talkowski, H.L. Rehm, M.J. Daly, G.
Tiao, B.M. Neale, D.G. MacArthur, and K.J. Karczewski, *A genome-wide*
mutational constraint map quantified from variation in 76,156 human genomes.
bioRxiv, 2022: p. 2022.03.20.485034.

25.   Beumer, J.H., E. Chu, C. Allegra, Y. Tanigawara, G. Milano, R. Diasio, T.W. Kim,
R.H. Mathijssen, L. Zhang, D. Arnold, K. Muneoka, N. Boku, and M. Joerger,
Therapeutic Drug Monitoring in Oncology: International Association of

Therapeutic Drug Monitoring and Clinical Toxicology Recommendations for 5-
Fluorouracil Therapy. Clin Pharmacol Ther, 2019. **105**(3): p. 598-613.

26.   Henricks, L.M., C. Lunenburg, F.M. de Man, D. Meulendijks, G.W.J. Frederix, E.
818   Kienhuis, G.J. Creemers, A. Baars, V.O. Dezentje, A.L.T. Imholz, F.J.F.
819   Jeurissen, J.E.A. Portielje, R.L.H. Jansen, P. Hamberg, A.J. Ten Tije, H.J.
820   Droogendijk, M. Koopman, P. Nieboer, M.H.W. van de Poel, C. Mandigers, H.
821   Rosing, J.H. Beijnen, E.V. Werkhoven, A.B.P. van Kuilenburg, R.H.N. van
822   Schaik, R.H.J. Mathijssen, J.J. Swen, H. Gelderblom, A. Cats, H.J. Guchelaar,
823   and J.H.M. Schellens, *DPYD genotype-guided dose individualisation of*
824   *fluoropyrimidine therapy in patients with cancer: a prospective safety analysis.*
825   Lancet Oncol, 2018. **19**(11): p. 1459-1467.

826   27.   Jiang, W., Z. Lu, Y. He, and R.B. Diasio, *Dihydropyrimidine dehydrogenase*
827   *activity in hepatocellular carcinoma: implication in 5-fluorouracil-based*
828   *chemotherapy.* Clin Cancer Res, 1997. **3**(3): p. 395-9.

829   28.   Kikuchi, O., S. Ohashi, Y. Nakai, S. Nakagawa, K. Matsuoka, T. Kobunai, T.
830   Takechi, Y. Amanuma, M. Yoshioka, T. Ida, Y. Yamamoto, Y. Okuno, S.
831   Miyamoto, H. Nakagawa, K. Matsubara, T. Chiba, and M. Muto, *Novel 5-*
832   *fluorouracil-resistant human esophageal squamous cell carcinoma cells with*
833   *dihydropyrimidine dehydrogenase overexpression.* Am J Cancer Res, 2015. **5**(8):
834   p. 2431-40.

835   29.   Li, L.H., H. Dong, F. Zhao, J. Tang, X. Chen, J. Ding, H.T. Men, W.X. Luo, Y. Du,
836   J. Ge, B.X. Tan, D. Cao, and J.Y. Liu, *The upregulation of dihydropyrimidine*
837   *dehydrogenase in liver is involved in acquired resistance to 5-fluorouracil.* Eur J
838   Cancer, 2013. **49**(7): p. 1752-60.

839   30.   Mayberry, R.M., R.J. Coates, H.A. Hill, L.A. Click, V.W. Chen, D.F. Austin, C.K.
840   Redmond, C.M. Fenoglio-Preiser, C.P. Hunter, M.A. Haynes, and et al.,
841   *Determinants of black/white differences in colon cancer survival.* J Natl Cancer
842   Inst, 1995. **87**(22): p. 1686-93.

843   31.   Govindarajan, R., R.V. Shah, L.G. Erkman, and L.F. Hutchins, *Racial differences*
844   *in the outcome of patients with colorectal carcinoma.* Cancer, 2003. **97**(2): p.
845   493-8.

846   32.   Alexander, D.D., J. Waterbor, T. Hughes, E. Funkhouser, W. Grizzle, and U.
847   Manne, *African-American and Caucasian disparities in colorectal cancer mortality*
848   *and survival by data source: an epidemiologic review.* Cancer Biomark, 2007.
849   **3**(6): p. 301-13.

850   33.   Akira, S., H. Isshiki, T. Sugita, O. Tanabe, S. Kinoshita, Y. Nishio, T. Nakajima,
851   T. Hirano, and T. Kishimoto, *A nuclear factor for IL-6 expression (NF-IL6) is a*
852   *member of a C/EBP family.* EMBO J, 1990. **9**(6): p. 1897-906.

34. Jakobsen, J.S., J. Waage, N. Rapin, H.C. Bisgaard, F.S. Larsen, and B.T. Porse,
Temporal mapping of CEBPA and CEBPB binding during liver regeneration
reveals dynamic occupancy and specific regulatory codes for homeostatic and
cell cycle gene batteries. Genome Res, 2013. **23**(4): p. 592-603.

35. Begay, V., J.J. Smink, C. Loddenkemper, K. Zimmermann, C. Rudolph, M.
Scheller, D. Steinemann, U. Leser, B. Schlegelberger, H. Stein, and A. Leutz,
Deregulation of the endogenous C/EBPbeta LIP isoform predisposes to
tumorigenesis. J Mol Med (Berl), 2015. **93**(1): p. 39-49.

36. Li, Y., E. Bevilacqua, C.B. Chiribau, M. Majumder, C. Wang, C.M. Croniger, M.D.
Snider, P.F. Johnson, and M. Hatzoglou, *Differential control of the*
CCAAT/enhancer-binding protein beta (C/EBPbeta) products liver-enriched
transcriptional activating protein (LAP) and liver-enriched transcriptional inhibitory
protein (LIP) and the regulation of gene expression during the response to
endoplasmic reticulum stress. J Biol Chem, 2008. **283**(33): p. 22443-56.

37. Miller, M., *Interactions of CCAAT/enhancer-binding protein beta with*
transcriptional coregulators. Postepy Biochem, 2016. **62**(3): p. 343-348.

38. Seo, J., D.D. Kocak, L.C. Bartelt, C.A. Williams, A. Barrera, C.A. Gersbach, and
T.E. Reddy, *AP-1 subunits converge promiscuously at enhancers to potentiate*
transcription. Genome Res, 2021. **31**(4): p. 538-550.

39. Cohen, D.M., H.W. Lim, K.J. Won, and D.J. Steger, *Shared nucleotide flanks*
confer transcriptional competency to bZip core motifs. Nucleic Acids Res, 2018.
46(16): p. 8371-8384.

40. Lountos, G.T., S. Cherry, J.E. Tropea, A. Wlodawer, and M. Miller, *Structural*
basis for cell type specific DNA binding of C/EBPbeta: The case of cell cycle
inhibitor p15INK4b promoter. J Struct Biol, 2022. **214**(4): p. 107918.

41. Zhang, X.F., K.K. Li, L. Gao, S.Z. Li, K. Chen, J.B. Zhang, D. Wang, R.F. Tu, J.X.
Zhang, K.X. Tao, G. Wang, and X.D. Zhang, *miR-191 promotes tumorigenesis of*
human colorectal cancer through targeting C/EBPbeta. Oncotarget, 2015. **6**(6): p.
4144-58.

42. Wang, D., L. Yang, W. Yu, Q. Wu, J. Lian, F. Li, S. Liu, A. Li, Z. He, J. Liu, Z.
Sun, W. Yuan, and Y. Zhang, *Colorectal cancer cell-derived CCL20 recruits*
regulatory T cells to promote chemoresistance via FOXO1/CEBPB/NF-kappaB
signaling. J Immunother Cancer, 2019. **7**(1): p. 215.

43. Offer, S.M., G.L. Butterfield, C.R. Jerde, C.C. Fossum, N.J. Wegner, and R.B.
Diasio, *microRNAs miR-27a and miR-27b directly regulate liver*
dihydropyrimidine dehydrogenase expression through two conserved binding
sites. Mol Cancer Ther, 2014. **13**(3): p. 742-51.

44.   Amstutz, U., S.M. Offer, J. Sistonen, M. Joerger, R.B. Diasio, and C.R. Largiader,
891   *Polymorphisms in MIR27A Associated with Early-Onset Toxicity in*
892   *Fluoropyrimidine-Based Chemotherapy.* Clin Cancer Res, 2015. **21**(9): p. 2038-
893   44.

894   45.   Hamzic, S., D. Scharer, S.M. Offer, D. Meulendijks, C. Nakas, R.B. Diasio, S.
895   Fontana, M. Wehrli, S. Schurch, U. Amstutz, and C.R. Largiader, *Haplotype*
896   *structure defines effects of common DPYD variants c.85T > C (rs1801265) and*
897   *c.496A > G (rs2297595) on dihydropyrimidine dehydrogenase activity:*
898   *Implication for 5-fluorouracil toxicity.* Br J Clin Pharmacol, 2021. **87**(8): p. 3234-
899   3243.

900   46.   Medwid, S., T.J. Wigle, and R.B. Kim, *Fluoropyrimidine-associated toxicity and*
901   *DPYD variants c.85T>C, c.496A>G, and c.1236G>A: impact of haplotype.*
902   Cancer Chemother Pharmacol, 2023. **91**(1): p. 97-102.

903   47.   Perez, A.R., Y. Pritykin, J.A. Vidigal, S. Chhangawala, L. Zamparo, C.S. Leslie,
904   and A. Ventura, *GuideScan software for improved single and paired CRISPR*
905   *guide RNA design.* Nat Biotechnol, 2017. **35**(4): p. 347-349.

906   48.   Machiela, M.J. and S.J. Chanock, *LDlink: a web-based application for exploring*
907   *population-specific haplotype structure and linking correlated alleles of possible*
908   *functional variants.* Bioinformatics, 2015. **31**(21): p. 3555-7.

909   49.   Hagege, H., P. Klous, C. Braem, E. Splinter, J. Dekker, G. Cathala, W. de Laat,
910   and T. Forne, *Quantitative analysis of chromosome conformation capture assays*
911   *(3C-qPCR).* Nat Protoc, 2007. **2**(7): p. 1722-33.

912   50.   Sarbassov, D.D., D.A. Guertin, S.M. Ali, and D.M. Sabatini, *Phosphorylation and*
913   *regulation of Akt/PKB by the rictor-mTOR complex.* Science, 2005. **307**(5712): p.
914   1098-101.

915   51.   Tan, G. and B. Lenhard, *TFBSTools: an R/bioconductor package for transcription*
916   *factor binding site analysis.* Bioinformatics, 2016. **32**(10): p. 1555-6.

917   52.   Castro-Mondragon, J.A., R. Riudavets-Puig, I. Rauluseviciute, R.B. Lemma, L.
918   Turchi, R. Blanc-Mathieu, J. Lucas, P. Boddie, A. Khan, N. Manosalva Perez, O.
919   Fornes, T.Y. Leung, A. Aguirre, F. Hammal, D. Schmelter, D. Baranasic, B.
920   Ballester, A. Sandelin, B. Lenhard, K. Vandepoele, W.W. Wasserman, F. Parcy,
921   and A. Mathelier, *JASPAR 2022: the 9th release of the open-access database of*
922   *transcription factor binding profiles.* Nucleic Acids Res, 2022. **50**(D1): p. D165-
923   D173.

924   53.   Consortium, E.P., *An integrated encyclopedia of DNA elements in the human*
925   *genome.* Nature, 2012. **489**(7414): p. 57-74.

54.   Portales-Casamar, E., S. Kirov, J. Lim, S. Lithwick, M.I. Swanson, A. Ticoll, J.
Snoddy, and W.W. Wasserman, *PAZAR: a framework for collection and*
dissemination of cis-regulatory sequence annotation. Genome Biol, 2007. **8**(10):
p. R207.

55.   Hammal, F., P. de Langen, A. Bergon, F. Lopez, and B. Ballester, *ReMap 2022:*
a database of Human, Mouse, Drosophila and Arabidopsis regulatory regions
from an integrative analysis of DNA-binding sequencing experiments. Nucleic
Acids Res, 2022. **50**(D1): p. D316-D325.

56.   Jolma, A., J. Yan, T. Whitington, J. Toivonen, K.R. Nitta, P. Rastas, E.
Morgunova, M. Enge, M. Taipale, G. Wei, K. Palin, J.M. Vaquerizas, R.
Vincentelli, N.M. Luscombe, T.R. Hughes, P. Lemaire, E. Ukkonen, T. Kivioja,
and J. Taipale, *DNA-binding specificities of human transcription factors.* Cell,
2013. **152**(1-2): p. 327-39.
939

 FIGURES AND FIGURE LEGENDS

Zhang et al., Figure 1

Figure 1. Identification of a novel *DPYD* enhancer. (A) Candidate *DPYD* enhancer

regions were selected for further study using data from GeneHancer and Ensembl

Regulatory Build. Coordinates are based on GRCh37/hg19. Regions are termed E9,

E16, and E20 based on approximate distance upstream of the *DPYD* transcription start

site. For CRISPR inhibition (CRISPRi), *DPYD* expression was measured in HepG2 cells

expressing dCas9-KRAB following transfection with guide-RNAs specific to the E9 (B),

E16 (C), and E20 (D) regions. For CRISPR activation (CRISPRa), *DPYD* expression

was measured in HepG2 cells expressing dCas9-P300 following transfection with guide-

950   RNAs specific to E9 (E), E16 (F), and E20 (G). Data represent the mean of three

951   independent biological replicates ± SD. *, $p<0.05$; **, $p<0.005$. P-values were calculated

952   using two-tailed Student's t-test comparing results to those from lentiguide controls.

Figure 2. Epigenetic changes at the E9 region induced by CRISPRi/CRISPRa.

Lentiguide vectors encoding guide-RNAs targeting the E9 region (E9 gRNAs) or empty

vector control (Lentiguide) were transduced into HepG2 cells expressing dCas9-KRAB

(A) or dcas9-P300 (B) and HCT116 cells expressing dCas9-KRAB (C) or dcas9-P300

(D). Chromatin immunoprecipitation (ChIP) was performed using antibodies specific to

Page 43

961   H3K9me3 (A, C) or H3K27ac (B, D). Quantitative PCR using primers centered at the

962   indicated regions (E) was used to measure the relative abundance of H3K9me3 and

963   H3K27ac. Data are presented relative to input DNA control and are further normalized

964   to IgG control. Error bars represent the SD of three independent biological replicate

965   experiments.

Figure 3. The rs4294451 T allele is associated with elevated *DPYD* expression. (A)

Luciferase reporter constructs were generated by cloning the E9 region containing the

reference rs4294451 T allele or the variant rs4294451 A allele into reporter vectors

containing the *DPYD* promoter. (B) Luciferase reporter activity was measured for

vectors shown in panel A. Error bars represent the standard deviation of three

independent biological replicates. (C) Relative expression of endogenous *DPYD*

measured via RT-qPCR in HCT116 cells engineered using CRISPR-mediated genome

editing to contain the depicted genotypes for rs4294451. Chromatin enrichment of

Page 45

H3K27ac (D) and H3K9me3 (E) was measured using ChIP-qPCR in HCT116 cells

engineered to contain the indicated genotypes at rs4294451. (F) *DPYD* allele-specific

expression was measured in human liver tissues using the C and T alleles at position

c.85. (G) Allele-specific expression was measured using the c.496-A and G alleles. All

panels: *, $p<0.05$; **, $p<0.005$; ***, $p<0.0005$. P-values were calculated as pairwise

comparisons between the indicated groups using two-tailed Student's t-tests.

Figure 4. The rs4294451 T allele is associated with increased interaction between E9 and the *DPYD* promoter. (A) Schematic of *Hind*III restriction enzyme sites (vertical bars) and primers (arrows) used for chromatin conformation capture (3C) relative to the *DPYD* transcription start site (TSS) and E9 region. (B) Legend for panels C–E. (C) 3C of chromatin interactions in rs4294451 knock-in HCT116 cells using anchor primer positioned within the digestion fragment containing the *DPYD* promoter. A, location of anchor primer; N, location of primer used for data normalization. (D) 3C of knock-in cells using anchor primer positioned within the fragment containing the E9 region. For panels C–D: one-way ANOVA p: *, p<0.01; **, p<0.001; all other data points, p>0.01. (E) Knock-in cells were treated with dilutions of 5-fluorouracil (5-FU) and viability assessed using real-time cell analysis (RTCA). Cell index is a measure of impedance between electrodes that are arrayed at the bottom of the RTCA plate and is representative of the number of live cells attached to the culture plate. Data are from 48 hours of 5-FU treatment. For all plotted data, the mean ± SD of three independent replicates is presented.

Page 47

1000

Figure 5. Rs4294451 T allele is associated with higher occupancy of CEBPB at E9

and the *DPYD* promoter. (A) Expression plasmids for CEBPB or GFP (control) were

co-transfected into HEK293T cells with the luciferase reporter plasmids depicted in

Figure 3A. (B) ChIP-qPCR was performed to examine the relative CEBPB enrichment at

the DNA fragments within the E9 region in knock-in HCT116 cells for the rs4294451

genotypes indicated. Primers used for E9 are centered on the position at 9201

nucleotides upstream of the *DPYD* TSS. (C) ChIP-qPCR was used to measure CEBPB

occupancy surrounding the *DPYD* promoter region. *, $p<0.05$; **, $p<0.005$; ***,

$p<0.0005$. P-values were calculated as pairwise comparisons between the indicated

groups using two-tailed Student's t-tests. For all panels, error bars represent the SD of

three independent replicates.

1012

Zhang et al., Figure 6

Figure 6. CEBPB-mediated upregulation of *DPYD* is dependent on rs4294451 T

allele. (A) Immunoblot showing knockdown of CEBPB expression in HCT116 cells

carrying different rs4294451 genotypes transduced with lentiviral particles encoding two

independent shRNAs against CEBPB (sh1 and sh2) or a scrambled control shRNA

(scr). *CEBPB* expression was measured by RT-qPCR in HCT116 A/A (B), T/A (C) and

T/T cells (D) transduced with the indicated shRNA lentiviral particles. *DPYD* expression

was measured in shCEBPB and scramble control HCT116 A/A (E), T/A (F) and T/T (G)

cells. The effect of CEBPB knockdown on cell viability in HCT116 A/A cells (H) and

HCT116 T/T cells (I) was measured by RTCA. Data shown are from 48 hours of 5-FU

treatment at the indicated concentrations. *, $p < 0.05$, calculated as a pairwise two-sided

Student's t-test comparing the indicated data to that of the associated scr control. Error

bars represent the SD from three independent replicates.

1026

 SUPPLEMENTARY TABLES

1028

Supplementary Table S1: Primers used for ChIP, cloning, qPCR, and site-directed mutagenesis.

Target	Forward Primer (5'-3')	Reverse Primer (5'-3')
E9 ChIP-1	CCTACCTACCACCCCCAAGA	TGCCCAGGACATTACACATGA
E9 ChIP-2	CATCAGGTTGCTTTTTGCAGC	TTAGCGTGGACTACCAGGGA
E9 ChIP-3	TGGGACTCAAAAAGCGGTTCA	GGTGTTGCGGGTGCTGATTA
E9 ChIP-4	GGTTCAACAGAGGATGCAACAC	AGGGAAAGATTTTCCTGGCCC
E9 ChIP-5	TAAAGATGCACCGAGGTGGG	AGTGTCTGGATTTAAGTAGATGTGC
E9 ChIP-6	AGCAGGAATGAGAAGGAGAGAAG	TCAGTCTCTCACTCCAAACCC
E9 ChIP-7	GCTCTTTTCATTGAAGCCTAAAACA	CCATCATACTTTTTCCAATTGTTGC
E9 ChIP-8	AATGATGAGAGGAAGATGACAAAGT	TGCCTACGCGATGAGTTGTA
DPYD promoter	AGTCGATATCCACAGTGTCTGTGTCTGGC	AGTCAAGCTTGCTCGATGTCCGCCGAG
E9 region	AGTCGGTACCGAATAAAACCAAAATAAAATCCATTTGGACGTTT	AGTCGAGCTCTTTGTGCAAAGGACCTTGGTATTTCC
Rs4294451 A>T	AAAGAAAAATAAATAAAAAAAGGAAAAATCTATAAGC	TTCTGGGGGTTGGTGTTG
Cas9	AACAGCCGCGAGAGAATGAA	CACGGGGTGTTCTTTCAGGA
DPYD	GTAAGGACTCGGCGGACATC	GCCGAAGTGGAACACAGAGT
L32	CCTTGTGAAGCCCAAGATCG	TGCCGGATGAACTTCTTGGT
CEBPB	CGCCGCCTGCCTTTAAATC	AAGCAGTCCGCCTCGTAGTA

1029

Page 51

Supplementary Table S2. Sequences and positions of the primers used for 3C analysis.

Primer position (TSS: +1)	Sequences (5'-3')
-22822	GGGAAGTTGAGAGAGCTAGGC
-19322	TGCTCTGTCAGCTGAGAAGACCTAGA
-16420	GTCACTACTGGGACTCTGAGAAA
-15616	AAAAGAAATTGCAACCTCTGGCA
-11097	GTTGCTTTTTGCAGCTGGGAT
-8087 (also used as anchor for E9 region)	AGTGCTTGAAGCTGATGAAGGG
-5884	CTGCAGAACAAGAACAGCACAT
1280 (also used as anchor for promoter)	TTAGGGTAGTCTATTCCTTTTTGGT
2826	TGCTTTGTGAGTGTACTGTTTGG
3076	CCTCCACCGGCAAGGATAAT

1030

1031

 SUPPLEMENTARY FIGURES AND FIGURE LEGENDS

1033

Zhang et al., Figure S1

1034

Figure S1. CRISPRi and CRISPRa screen to identify *DPYD cis* regulatory elements

in HCT116 cells. For CRISPRi, *DPYD* expression was measured in HCT116 cells

expressing dCas9-KRAB following transfection with guide-RNAs specific to the E9 (A),

E16 (B), and E20 (C) regions. For CRISPRa, *DPYD* expression was measured in

HCT116 cells expressing dCas9-P300 following transfection with guide-RNAs specific

to E9 (D), E16 (E), and E20 (F). Data represent the mean of three independent

biological replicates ± SD. *, $p<0.05$. P-values were calculated using two-tailed

Student's t-test comparing results to those from lentiguide controls.

Zhang et al., Figure S2

Figure S2. Genomic context of rs4294451. Rs4294451 is located within a putative

enhancer region showing evidence for regulatory activity in Ensembl Regulatory Build

data and within transcription factor binding sites in ENCODE Factorbook data.

Page 54

1047

Figure S3. Rs4294451 A allele is associated with epigenetic repression at the E9

region in human liver specimens. Chromatin enrichment of H3K27ac (A) and

H3K9me3 (B) was measured using ChIP-qPCR of liver specimens obtained from

human donors carrying different rs4294451 genotypes. Data represent three

independent measurements from a single liver specimen with each indicated genotype

± SD.

1054

Figure S4. Disruption of CEBPB reduces enrichment at the E9 region and *DPYD*

promoter in rs4294451 T/T and A/T cells, but not in T/T cells. CEBPB enrichment at

the E9 region (A) and the *DPYD* promoter (B) was measured by ChIP-qPCR in CEBPB

knockdown and scramble (scr) control knock-in HCT116 cells containing the indicated

rs4294451 genotype. Data represent the mean of three independent replicates ± SD. *,

p<0.05. P-values were calculated using two-tailed Student's t-test.

1061

1062

1063 **Figure S5. Allele frequency for rs4294451-T allele in global populations.** (A) Allele

1064 frequencies in various populations for the rs4294451-T allele was retrieved from the

1065 gnomAD browser v3.1.2. (B) Local ancestry-informed frequency data was retrieved for

1066 the rs4294451-T allele within Latino-Admixed American samples of gnomAD v3.1

Page 57