1   ***Candida albicans*** **suppression of neutrophil reactive nitrogen species is rescued by**

2   **host Hif-1α** ***in vivo***

Thomas B. Burgess[1], Ffion R. Hammond[1], Piotr T. Szkuta[1], Amy Lewis[1], Stella Christou[2],

Kathryn R. Ayscough[2], Simon A. Johnston[1], Alison M. Condliffe[1] and Philip M. Elks[1]

[1]The Bateson Centre, School of Medicine and Population Health, University of Sheffield,

Beech Hill Road, Sheffield, S10 2RX, UK.

[2]The School of Biosciences, University of Sheffield, Firth Court, Western Bank, Sheffield,

S10 2TN, UK.

26 **Corresponding Author:**

27 Philip M. Elks (p.elks@sheffield.ac.uk)

Abstract

Candida spp. cause 750,000 cases per annum of invasive disease worldwide, with up to a 50% mortality rate. Poor efficacy of current antifungals, lack of vaccines and rising antifungal resistance rates point towards an urgent need to develop new therapies. *Candida albicans* is a human commensal fungus that can cause life-threatening invasive infection in immunocompromised individuals. *C. albicans* is able to manipulate host macrophages and neutrophils to escape phagosomal killing and has previously been shown to suppress reactive nitrogen species (RNS) production *in vitro*. However, the effects of *C. albicans* on RNS *in vivo* and the molecular and cellular mechanisms involved remain unclear. Using a zebrafish model, we aimed to characterise RNS suppression by *C. albicans in vivo*. We demonstrate that *C. albicans* suppressed neutrophil RNS both proximally and distally to the infection site in a partially active process, with heat-killed *C. albicans* not reducing RNS to the same extent as live fungi. Using a *car1Δ* mutant, we show that fungal arginase is partially responsible for the reduction in neutrophil RNS. Stabilisation of Hif-1α, a transcription factor with a key role in immune regulation, rescued neutrophil RNS production during *C. albicans* infection, leading to improved infection outcomes. The protective effect of Hif-1α stabilisation was neutrophil- and nitric oxide synthase-dependent. Together, these data demonstrate that Hif-1α stabilisation can restore the neutrophil RNS response in *C. albicans* infection, leading to improved infection outcomes, highlighting the potential of targeting Hif-1α and RNS in host directed therapies against fungal infections.

Introduction

Candida albicans is a commensal fungus, colonising 70% of healthy individuals (Witherden *et al*., 2017), which can cause superficial mucosal infections with negligible mortality. However, *C. albicans* can also act as an opportunistic pathogen capable of causing life-threatening, invasive disease in immunocompromised individuals and neonates (Witherden *et al*., 2017; Flores-Maldonado *et al*., 2019; Pellon, Sadeghi Nasab and Moyes, 2020). *Candida* species are the 4[th] most common nosocomial infection in the USA and

account for up to 22% of health-care-associated bloodstream infections (Pfaller and

Diekema, 2010; Magill *et al.*, 2014). There are approximately 750,000 cases of severe

candidiasis worldwide per year and the estimated mortality rate is around 50% (Bongomin *et*

al., 2017; WHO, 2022). This high mortality, combined with the lack of fungal vaccines and

rising rates of antifungal resistance, has resulted in *C. albicans* being designated as a critical

priority fungal pathogen, with urgent research into new alternative therapies required (WHO,

2022).

C. albicans infections are initially detected by pattern recognition receptors, including

Toll-like receptors, C-type lectin receptors and the Rig-I like receptor, MDA-5, resulting in a

downstream pro-inflammatory response and innate immune cell recruitment (reviewed in

Burgess, Condliffe and Elks, 2022). Although many immune cell types are recruited to the

sites of *C. albicans* infection, neutrophils are one of the most potent killers of *C. albicans* and

neutropenic patients are particularly at risk of invasive disease (Uzun *et al.*, 2001; Brown,

2011). Candidalysin, produced by hyphal *C. albicans*, is a particularly strong stimulator of

CXCL1-mediated neutrophil recruitment (Drummond *et al.*, 2019). Following recruitment,

neutrophils have multiple mechanisms of eliminating *C. albicans*, including the production of

reactive oxygen (ROS) and nitrogen (RNS) species (Winterbourn *et al.*, 2006). Inhibition of

inducible nitric oxide synthase (iNOS) leads to increased mortality in mouse models of *C.*

albicans infections, showing the importance of RNS mediated fungicidal activity (Tillmann,

Gow and Brown, 2011; Wagener *et al.*, 2017; Navarathna, Lionakis and Roberts, 2019).

C. albicans has a variety of mechanisms that can subvert the host innate immune

response (Luo *et al.*, 2013). Additionally, *C. albicans'* ability to rapidly adapt to changes in

environmental pH, metabolic flexibility and a robust stress response allows colonisation of a

wide range of niches within the host (Mayer, Wilson and Hube, 2013). The morphotype

yeast-to-hyphal transition is essential for host tissue invasion and limits innate immune

sensing of *C. albicans* via masking of β-glucan (Gantner, Simmons and Underhill, 2005; van

der Graaf *et al.*, 2005). *C. albicans* can suppress ROS and RNS production by macrophages

in vitro (Wellington, Dolan and Krysan, 2009) with an, as yet, unidentified *C. albicans*

secreted compound responsible for RNS inhibition in murine macrophages (Collette, Zhou

and Lorenz, 2014). However, there is currently a lack of *in vivo* observations on the effects of

C. albicans on RNS and little is known about the molecular and cellular mechanisms

underlying RNS suppression and whether it is possible to rescue this with potential

therapeutic application.

Hypoxia inducible factor (HIF) proteins are important regulators of cellular responses

to hypoxia, innate immunity and inflammation (McGettrick and O'Neill, 2020). In hypoxic

conditions, HIF-1α proteins are stabilised, allowing entry into the nucleus, where they form

heterodimers with HIF-1β. Consequent transcription of target genes regulate cellular

metabolism, proliferation, migration, differentiation, angiogenesis and pro-inflammatory

mediators (Krzywinska and Stockmann, 2018; Hammond, Lewis and Elks, 2020). HIF-1α

stabilisation is associated with increased neutrophil survival, via NF-κB and promotes

neutrophil extracellular trap formation (Walmsley *et al.*, 2005; McInturff *et al.*, 2012). We

have previously shown that stabilisation of Hif-1α increases neutrophil RNS output, which is

protective against *Mycobacterium marinum* infection *in vivo* (Elks *et al.*, 2013). Therefore,

Hif-1α stabilisation represents a potential host directed therapy to target neutrophil RNS

during fungal infections, such as *C. albicans*.

Zebrafish are a highly tractable *in vivo* model system to investigate the behaviours of

innate immune cells during infection (Torraca and Mostowy, 2018). Zebrafish larvae are

transparent, allowing the use of innate immune transgenic lines to directly observe neutrophil

behaviours and interactions with pathogens (Lam *et al.*, 2004; Renshaw *et al.*, 2006).

Adaptive immunity does not mature until 4 to 6 weeks post fertilisation, whereas innate

immune cells are present from early larval stages, allowing examination of innate immune

responses to infections without the presence of adaptive immunity (Lam *et al.*, 2004).

Several infection sites have been used in zebrafish to model *C. albicans* infection, including

localised infection into the hindbrain ventricle and systemic infection via injection into the

caudal vein, allowing neutrophil behaviours to be observed to better our understanding of

host-*C. albicans* interactions in an *in vivo* context (Bergeron *et al.*, 2017; Gratacap *et al.*,

2017; Seman *et al.*, 2018).

Here, we show that *C. albicans* infection suppressed neutrophil RNS in zebrafish.

RNS suppression by *C. albicans* was shown to occur both locally and distally to the infection

site, and was a partially active process, with heat-killed *C. albicans* having a reduced effect

on neutrophil RNS. Fungal arginase (*car1*) plays a role in RNS reduction, with a *car1Δ*

mutant having a reduced effect on neutrophil RNS. Furthermore, we show that it is possible

to restore host RNS production via Hif-1α stabilisation, enhancing host control of *C. albicans*

infection. Together, this work highlights Hif-1α and RNS as potential targets for host directed

therapies against invasive *C. albicans* infections.

Methods

Ethics Statement

Zebrafish were handled in accordance with Animals (Scientific Procedures) Act 1986,

under Project Licenses P1A4A7A5E and PP7684817. Ethical approval was granted by the

University of Sheffield Local Ethical Review Panel.

Zebrafish

Zebrafish were maintained in Home Office approved facilities in Biological Services

Unit (BSU) aquaria at the University of Sheffield, in accordance with standard protocols and

local animal welfare regulations. Adult fish were maintained at 28°C with a 14/10 hour

light/dark cycle. Embryos were maintained at 28 °C with a 14/10 hour light/dark cycle in 1X

E3 medium (E3 60X stock: 5.0 mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$;

diluted to 1X in distilled water) with 10-5% Methylene Blue (Sigma-Aldrich). Methylene Blue

was added to prevent fungal growth. The following zebrafish strains were used: nacre (wild

type), *Tg(mpx:GFP)i114* (Renshaw *et al.*, 2006), *Tg(lyz:nfsB.mCherry)sh260* (Buchan *et al.*,

2019), Tg(*mpeg:nlsClover*)sh436 (Bernut *et al.*, 2019).

Fungal Strains

Several *C. albicans* strains were used in this study (Table 1). *C. albicans* was initially grown on YPD (4001022, MP Bio) plates at 28°C for 48 hours, then transferred to overnight liquid culture in YPD broth in a shaking incubator at 30°C at 200 rpm.

Table 1: List of *C. albicans* strains used in this study

Strain	Genotype	Reference
C. albicans TT21-dTomato	*ade2::hisG/ade2::hisG ura3::imm434/ura3::imm434::URA3-tetO ENO1/eno1::ENO1 tetR-ScHAP4AD-3XHA-ADE2* pENO1-dTomato-NATR	(Seman *et al.*, 2018)
C. albicans SN148 GFP	SN148 *ENO1/ENO1-GFP::LEU2*	(Christou, Ayscough and Johnston, 2023)
C. albicans SC5314	wt; clinical blood isolate	(Gillum, Tsay and Kirsch, 1984)
C. albicans car1Δ	As SC5314 but *car1Δcar1Δ*	(Schaefer *et al.*, 2020)

RNA Injections

Embryos were injected with dominant active *hif-1αb* variant RNA (DA1) or dominant negative *hif-1αb* variant RNA at the one-cell stage, as previously described (Elks *et al.*, 2011). Phenol Red (PR; Sigma-Aldrich) was used as a vehicle control.

Morpholino Injections

A *csf3r* morpholino (Gene Tools) was used to deplete neutrophil populations. A standard control morpholino (Gene Tools) used as a negative control. The antisense morpholino oligonucleotide sequences were: *csf3r* – 5'-GAAGCACAAGCGAGACGGATGCCAT-3'; control – 5'-CCTCTTACCTCAGTTACAATTTATA-3'.

F0 CRISPR-Cas9 Knockdown (CRISPant)

Two sgRNAs were designed to target the ATG/first exon of *nos2a* and *nos2b* to knockdown iNOS and RNS production. Efficacy of knockdown of RNS production was functionally validated by anti-nitrotyrosine staining. Tyrosinase (*tyr*) sgRNA was used as a negative control (Isles *et al.*, 2019). The target sequences were: *nos2a* – 5'-TTTCTCATTTTCAATGATAG-3'; *nos2b* – 5'-GTTCGCTCTTGTGAGTGACC-3'; *tyr* – 5'-CCTGACCTCCTGAAGACCCC-3'. 25µM sgRNA was co-injected with tracrRNA, 1:2 diluted Cas9 and phenol red.

C. albicans Infections

Overnight *C. albicans* cultures were washed 3 times in phosphate buffered saline (PBS) and counted on a haemocytometer. Washed cultures were resuspended in 10% PVP (polyvinylpyrrolidone; Calbiochem) to achieve the desired inoculation. Infection dose for systemic caudal vein infections were 100 colony forming units (cfu) or 200 cfu. Infection dose for localised hindbrain infections was 20 cfu. *C. albicans* was heat-killed by incubation in a heat block at 65°C for 1 hour (Panpetch *et al.*, 2017). To denature fungal proteins, *C. albicans* was boiled at 105°C for 5 minutes.

For survival curves, zebrafish survival was monitored daily and dead zebrafish were removed from the Petri dish. Death was determined by a lack of heartbeat/circulation or tissue degradation post mortem. At 4 days post infection (dpi; equivalent to 5 days post fertilisation, dpf), all surviving zebrafish larvae were culled.

Anti-nitrotyrosine Staining

1 dpi embryos were fixed in 4% paraformaldehyde in PBS overnight in 4°C. Nitrotyrosine levels were labelled using a rabbit polyclonal anti-nitrotyrosine antibody (06-284; Merck Millipore) and were detected using Alexa-633 conjugated goat-anti-rabbit secondary antibody (Invitrogen Life Technologies) imaged using line-scanning confocal microscopy, as previously described (Elks *et al.*, 2013, 2014).

Microscopy and Quantification of Anti-nitrotyrosine

For *C. albicans* SC5314 and *car1Δ*, brightfield images were taken using an inverted Leica DMi8 with a 40x lens and a Hamamatsu OrcaV4 camera. Stained embryos were imaged on a Leica DMi8 inverted microscope with a Leica TCS-SPE line-scanning confocal for imaging with a 40x lens. For quantification purposes, acquisition settings were kept the same across the groups. Corrected fluorescence intensity was calculated using Image J measurements assessing the cell fluorescence of individual cells corrected for cell size and background fluorescence of the image (Elks *et al.*, 2013, 2014).

Statistical Analysis

All data were analysed using GraphPad Prism 9.5.1 1 (GraphPad Software, La Jolla, California, USA, www.graphpad.com). Nitrotyrosine staining data was analysed with Kruskal-Wallis test, with Dunn's multiple comparisons test. Nitrotyrosine staining data for proximal and distal RNS levels was analysed with 2-way ANOVA, with Sidák's multiple comparison test. Survival curves were analysed with Gehan-Breslow-Wilcoxon test, with Bonferroni correction. The p values shown are: ns=not significant, $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$

Results

C. albicans infection suppresses neutrophil RNS production

C. albicans has previously been demonstrated to modulate host innate immune

responses, by suppressing production of ROS and RNS by macrophages *in vitro* (Wellington,

Dolan and Krysan, 2009; Collette, Zhou and Lorenz, 2014). In zebrafish, we have previously

shown that RNS, identified using an anti-nitrotyrosine antibody, is predominantly produced in

neutrophils, both in resting/mock infected individuals (Elks *et al.*, 2014) and to an elevated

level after bacterial infection with *Mycobacterium marinum* (Elks *et al.*, 2013). We aimed to

establish if *C. albicans* is capable of modulating neutrophil RNS in an *in vivo* zebrafish model.

1 day post fertilisation (dpf) zebrafish larvae were infected in the caudal vein with either live

C. albicans, heat killed *C. albicans* or a mock infected (PVP) control. At 24 hours post

infection (hpi), zebrafish larvae were fixed and stained with an anti-nitrotyrosine antibody as

a measure of host neutrophil RNS production. In mock infected (PVP) controls there was a

basal level of anti-nitrotyrosine levels in neutrophils (Elks *et al.*, 2013, 2014). Compared to

the PVP control, *C. albicans* TT21-dTomato caused a 90.0% (p<0.0001) reduction in the

corrected fluorescence intensity of nitrotyrosine staining (Figure 1A-B). Heat-killed *C.*

albicans TT21-dTomato had an intermediate level of RNS suppression, between PVP

(PVP=5.17, heat-killed *C. albicans* TT21-dTomato=2.47; p<0.0001) and live *C. albicans*

infection (*C. albicans* TT21-dTomato=0.52, heat-killed *C. albicans* TT21-dTomato=2.47;

p<0.0001). Similar phenotypes of RNS suppression were observed in infection with live and

heat-killed *C. albicans* SN148 GFP, suggesting the effect is not strain specific (Figure S1).

Hence, we used these fluorescence strains interchangeably, depending on the colour

combinations required for each experiment. Together these data show that *C. albicans*

suppresses neutrophil RNS production *in vivo*, while heat-killed *C. albicans* is capable of

generating an intermediate level of RNS suppression.

Suppression of neutrophil RNS by *C. albicans* occurs both proximally and distally to the infection site

To determine whether neutrophils with reduced RNS need to be proximal to *C. albicans* infection, a local hindbrain ventricle *C. albicans* infection was performed, with RNS production measured both at the site of infection in the hindbrain and a distal, uninfected site in the caudal haematopoietic tissue (CHT). *C. albicans* caused a significant decrease in nitrotyrosine levels in both proximal neutrophils in the hindbrain (PVP=2.38, *C. albicans*=0.52; p<0.0001) and distal neutrophils in the CHT (PVP=2.59, *C. albicans*=0.71; p<.0001) compared to PVP control (Figure 2A-C). Hence, live *C. albicans* is capable of causing both a proximal and distal suppression of neutrophil RNS, suggestive of the influence of a secreted factor. Heat-killed *C. albicans* caused an intermediate RNS reduction in the hindbrain but had no effect on RNS in distal neutrophils in the CHT region, suggesting that multiple mechanisms may reduce neutrophil RNS that differ between proximal and distal regulation. Boiled *C. albicans* had no effect on neutrophil RNS at either location, implying that a secreted factor regulating neutrophil RNS may be a protein (that does not function once denatured). Together, these results suggest that *C. albicans* may have multiple mechanisms of neutrophil RNS suppression, some which act locally and others which act more distally.

C. albicans **arginase, CAR1, plays a role in the suppression of neutrophil RNS**

Host RNS levels are controlled by a balance of two enzymes that share the substrate L-arginine, iNOS and arginase. iNOS is the enzyme responsible for RNS production and arginase, shunts the metabolism of L-arginine away from RNS towards L-ornithine production (Bansal and Ochoa, 2003). Fungi are known to produce enzymes with similar functions in nitric oxide production, for example C. albicans produces its own version of arginase, *CAR1* (Schaefer *et al.*, 2020). We therefore hypothesised that *C. albicans* arginase interferes with the host arginase-iNOS axis, resulting in neutrophil RNS suppression. 1 dpf zebrafish were infected with *C. albicans car1Δ* or its parental strain *C. albicans* SC5314 and

RNS production was measured at 24 hpi by anti-nitrotyrosine staining. *C. albicans car1Δ*

infection decreased neutrophil RNS production compared to PVP ($p < 0.0001$) (Figure 3A-B).

The RNS produced after live *C. albicans car1Δ* infection was significantly different from live

C. albicans SC5314 ($p < 0.05$), heat-killed *C. albicans* SC5314 ($p < 0.0001$) and heat-killed *C.*

albicans car1Δ ($p < 0.0001$), Heat-killed *C. abicans car1Δ* was not significantly different from

heat-killed *C. albicans* SC5314 ($p > 0.9999$), suggesting an active component is responsible

for differences observed in live *C. albicans*. *C. albicans car1Δ* caused an intermediate level

of RNS suppression between live *C. albicans* SC5314 and heat-killed *C. albicans* (Figure

3A-B), indicating that *CAR1* plays a partial role in suppression of neutrophil RNS.

Hif-1α is protective against *C. albicans* infection in a neutrophil-dependent manner

Hif-1α is a transcription factor with known roles in regulation of innate immunity

(McGettrick and O'Neill, 2020) and has previously been shown to be protective against

Mycobacterium marinum infection *in vivo* by increasing neutrophil RNS levels (Elks *et al.*,

2013; Hammond, Lewis and Elks, 2020). We aimed to stabilise Hif-1α in *C. albicans* infected

zebrafish embryos, in order to restore innate immune responses and increase host survival

in *C. albicans* infection.

1 cell stage zebrafish embryos were injected with phenol red (solvent control, PR),

dominant negative *hif-1α* (DN1) or dominant active *hif-1α* (DA1) RNA, then infected

systemically with *C. albicans* TT21-dTomato infection at 30 hpf. Mortality was measured

over the following 4 days. DA1 injected embryos had significantly greater survival across 4

days than both PR ($p < 0.01$) and DN1 ($p < 0.001$) injected larvae (Figure 4A-B), suggesting

Hif-1α stabilisation is host protective in *C. albicans* infection.

Neutrophils play an essential role in innate immune control of invasive candidiasis

(Desai and Lionakis, 2018). Brothers *et al.* showed neutrophils are responsible for killing of *C.*

albicans, whereas macrophages internalised, but did not kill, *C. albicans in vivo* (Brothers *et*

al., 2013). Therefore, we hypothesised that the protective effect of Hif-1α stabilisation was

mediated by neutrophils. To assess the impact of Hif-1α stabilisation on macrophage-

mediated immunity, we depleted the macrophage population by injection of clodronate

liposomes (van Rooijen and Hendrikx, 2010) at 30 hpf or PBS liposomes as a negative

control, leading to a significant depletion of the macrophage population (Figure S3). 1 cell

stage embryos were injected with DA1 or PR control. At 30 hpf, embryos were injected into

the caudal vein with clodronate liposomes or PBS liposomes followed by infection with

100cfu *C. albicans* TT21-dTomato at 2 dpf. Both groups of PBS liposome injected larvae had

greater survival than clodronate liposome, macrophage depleted, siblings. DA1 was

protective in both the presence or depletion of macrophages, with clodronate liposome

larvae having greater survival than PR clodronate liposome embryos (Figure 4C; p<0.0001),

indicating that the host-protective effect of Hif-1α stabilisation is not dependent on the

presence of macrophages.

Next, we aimed to establish if neutrophils were responsible for the protective effect of

Hif-1α stabilisation in *C. albicans* infection. We injected 1 cell stage zebrafish embryos with a

csf3r morpholino (Ellett *et al*., 2011) to deplete neutrophil numbers in 2 dpf zebrafish larvae

(Figure S4). *Csf3r* morpholino injection led to a significant decrease in survival of 100cfu *C.*

albicans infected larvae, in both PR and DA1 injected fish (Fig 4D). DA1 + *csf3r* larvae had

lower survival compared to PR + *csf3r* larvae, showing a loss of the protective effect of Hif-

1α stabilisation in neutrophil ablated larvae (Figure 4D) and demonstrating that the

protective effect of Hif-1α stabilisation is neutrophil dependent.

Host protective effect of Hif-1α is RNS-dependent

Hif-1α stabilisation has previously been shown to increase RNS production in

zebrafish neutrophils (Elks *et al*., 2013). We hypothesised that the host protective effect of

Hif-1α stabilisation in *C. albicans* infection was due to increased neutrophil RNS production.

Therefore, we investigated the effect of dominant active *hif-1α* (DA1) on neutrophil

behaviours during *C. albicans* infection. DA1 did not alter the recruitment of neutrophils to

the site of a local *C. albicans* infection compared to phenol red (PR) injected controls (Figure

S5), implying a change in neutrophil behaviour, not neutrophil numbers, is responsible for

the protective effect of Hif-1α stabilisation. In anti-nitrotyrosine stained larvae, DA1/PVP

caused a doubling in nitrotyrosine levels compared to PR/PVP, mock injected larvae (Figure

5A-B; PR/PVP=16.36, DA1/PVP=33.25; p<0.0001) as previously described (Elks *et al.*,

2013). While *C. albicans* reduced nitrotyrosine levels in PR control injected embryos, DA1

injection restored nitrotyrosine levels in *C. albicans* infected larvae to comparable levels of

uninfected larvae (Fig 5A-B).

323   To demonstrate that this restoration of RNS in Hif-1α stabilisation is responsible for

the host protective effect in *C. albicans* infection, we inhibited production of RNS

pharmacologically, using L-NIL, and genetically, using *nos2a+b* (zebrafish iNOS) double

CRISPR-Cas9 knockdown ("CRISPants"; Moore *et al.*, 1994; Isles *et al.*, 2019). L-NIL

treatment did not impact the growth of *C. albicans* outside of the host *in vitro* (Figure S6). PR

+ L-NIL larvae had reduced survival compared to PR + dH$_2$O (drug solvent control) treated

larvae (Figure 5C), demonstrating the importance of RNS in *C. albicans* control. DA1 + L-NIL

treated larvae did not have a significant survival advantage compared to either PR + dH$_2$O or

PR + L-NIL groups (Figure 5C), indicative of iNOS inhibition removing the host protective

effect of Hif-1α stabilisation. *Nos2ab* double CRISPants were shown to inhibit nitrotyrosine

production in PVP injected zebrafish (Figure S7). DA1 with *nos2ab* CRISPants no longer

had increased survival compared to PR + *tyr* sgRNA (*tyrosinase* CRISPant control; Isles *et*

al., 2019) larvae, supporting that RNS production by *nos2* is responsible for the host

protective effect of Hif-1α stabilisation (Figure 5D). Together these data demonstrate that the

Hif-1α stabilisation is host-protective via a neutrophil- (Fig 4) and RNS-dependent (Fig 5)

mechanism.

Discussion

341   The increasing prevalence of antifungal resistance worldwide highlights the

importance of discovering new antifungal targets or host directed therapies capable of

improving disease outcomes. Here, we have shown that Hif-1α stabilisation represents a

potential host directed therapy capable of restoring the neutrophil RNS response to *C.*

albicans infection.

We first demonstrated that *C. albicans* can suppress neutrophil RNS production *in*

vivo. Collette *et al.* previously showed *C. albicans* suppresses RNS production in murine

bone marrow derived macrophages *in vitro* (Collette, Zhou and Lorenz, 2014), which is

supported by a variety of sources showing suppression of ROS by *C. albicans in vitro*

(Frohner *et al.*, 2009; Wellington, Dolan and Krysan, 2009). While previous publications have

demonstrated RNS suppression in macrophages, our data demonstrates an effect in

neutrophils. Human neutrophils produce higher levels of RNS than macrophages (Muijsers

et al., 2001), as is the case in zebrafish larvae, suggesting our findings may reflect the

human RNS response to *C. albicans* making zebrafish a useful *in vivo* model to investigate

the immune response in these difficult to study *ex vivo*, short-lived cells.

We showed *C. albicans* suppression of neutrophil RNS occurs both proximally at the

site of infection and also at distal sites. Investigations using *in vitro* models have been

unable to fully answer questions about host-*C. albicans* interactions in the context of a whole

organism. Previous work has suggested direct contact between *C. albicans* and

macrophages is required for RNS suppression and that an unidentified secreted factor plays

a role (Collette, Zhou and Lorenz, 2014). Our findings do not preclude a suppression effect

from direct contact between *C. albicans* and neutrophils, but suggest that direct contact is

not a requirement and that suppression involves the secretion of a factor, allowing

suppression of distal neutrophil RNS. Our *in vivo* model suggests that *C. albicans* has a

secreted mechanism of RNS suppression which is capable of having effects large distances

(millimetres) across an organism. This raises new questions about the role of *C. albicans* in

the development of polymicrobial coinfections in immune compromised patients. This

secreted factor is likely to be a protein as boiling *C. albicans*, resulting in denaturation of

proteins, inhibits the distal suppression of neutrophil RNS. Pathogen-specific secreted

compounds are a rich source for potential biomarkers of infection (Stastna and Van Eyk,

2012). Further research into the specific molecule responsible for suppressing neutrophil

RNS distally is required to determine its suitability to act as a biomarker of *C. albicans*

infection. The intermediate phenotype of RNS suppression by heat-killed *C. albicans* hints at

multiple mechanisms of neutrophil RNS suppression by *C. albicans*. Current literature

supports the idea of multiple mechanisms of suppression. *C. albicans* SOD5 has been

associated with degradation of antimicrobial ROS *in vitro* (Frohner *et al.*, 2009), *C. albicans*

chitin induces host arginase-1 activity, decreasing RNS production (Wagener *et al.*, 2017),

YHB1 is able to detoxify RNS (Hromatka, Noble and Johnson, 2005) and Collette *et al.*

identified an unknown, small secreted molecule capable of suppressing RNS (Collette, Zhou

and Lorenz, 2014). The diversity of proteins and genes associated with decreased RNS

production correlates with our hypothesis that multiple mechanisms of RNS suppression

exist. Further investigation could aim to establish which, if any, mechanism contributes the

most to host RNS suppression, and if any mechanisms can be targeted to downregulate the

suppressive effect. Alternative explanations may be that heat-killing is partially denaturing

fungal proteins, reducing activity of the hypothesised secreted protein, or that heat-killing is

damaging the fungal cell wall, resulting in exposure of chitin, which is known to decrease

RNS production (Wagener *et al.*, 2017).

Here, we identify that *C. albicans* arginase is one such mechanism involved in the

suppression of host RNS production. *CAR1* is the only known arginase gene in the *C.*

albicans genome, encoding a cytosolic arginase (Schaefer *et al.*, 2020). Fungal arginase

may be interfering with the host arginase-iNOS balance, resulting in reduced iNOS activity

and RNS suppression. The modest decrease in neutrophil RNS production after infection

with the *CAR1* mutant suggests that *C. albicans* arginase production may not be the

predominant RNS reducing mechanism, but additive strategies of neutrophil RNS

suppression by *C. albicans* may underlie their success at the large overall suppression of

neutrophil RNS observed locally and distally.

We have demonstrated that Hif-1α stabilisation has a protective effect in *C. albicans*

infection in zebrafish embryos, via a neutrophil-mediated, RNS-dependent mechanism.

Neutrophils are the primary innate immune cell responsible for killing *Candida spp.* (Brown,

2011), supporting our conclusion that neutrophil suppression is an important strategy of *C.*

albicans during infection that, if rescued, could represent an exciting therapeutic opportunity.

RNS has well characterised, highly candidacidal effects (Rementería, García-Tobalina and

Sevilla, 1995; Vazquez-Torres *et al.*, 1995; Tillmann, Gow and Brown, 2011; Navarathna,

Lionakis and Roberts, 2019). Similarly, Macherla *et al.* revealed NO-releasing nanoparticles

inhibit the growth of *C. albicans in vitro* and *in vivo* (Macherla *et al.*, 2012). Hence,

upregulation of neutrophil RNS production, following Hif-1α stabilisation, is a potential

mechanism that could be targeted therapeutically to improve immune candidacidal activity.

We have previously demonstrated Hif-1α stabilisation protects against *M. marinum* infection

in a similar manner (Elks *et al.*, 2013), suggesting Hif-1α stabilisation could offer protective

effect against multiple and different groups of pathogens (fungi and bacteria) where

neutrophil-mediated immune responses are subverted.

Our data indicates that Hif-1α stabilisation is able to overcome *C. albicans*

suppression of neutrophil RNS, conferring a neutrophil-mediated, nitric oxide-dependent

protective effect *in vivo*. Hence, Hif-1α stabilisation has potential as a host directed

therapeutic strategy to aid in the treatment of *C. albicans* infections, following further

validation in human systems. We also begin to determine the mechanisms underlying RNS

suppression by *C. albicans*. Identification of specific proteins, genes or compounds

responsible for RNS suppression could point towards new potential antifungal targets or

prospective biomarkers to aid the diagnosis and treatment of *C. albicans* infections.

    Figures and Legends

Figure 1: *C. albicans* suppressed neutrophil RNS production *in vivo*.

(A) Anti-nitrotyrosine fluorescence at 24 hpi following injection of PVP, live *C. albicans* TT21-dTomato or heat-killed *C. albicans* TT21-dTomato into the caudal vein at 30 hpf. N=96-108 neutrophils from 16-18 fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance determined by Kruskal-Wallis test, then Dunn's multiple comparisons test. ****p<0.0001.

(B) Representative images of PVP, *C. albicans* TT21-dTomato and heat-killed *C. albicans* TT21-dTomato infected zebrafish embryos at 24 hpi. Scale bars = 50 μm.

Figure 2: *C. albicans* suppression of RNS occurs proximally and distally to site of infection.

(A) 1 dpf zebrafish were injected with PVP, *C. albicans,* heat-killed *C. albicans* or boiled *C. albicans*, into the hindbrain ventricle. At 24 hpi, embryos were fixed and stained with anti-nitrotyrosine primary antibody and goat-anti-rabbit Alexa-633 secondary antibody. Zebrafish were then imaged at the hindbrain and CHT regions and anti-nitrotyrosine fluorescence was quantified.

(B) Anti-nitrotyrosine fluorescence at 24 hpi following injection of PVP, *C. albicans,* heat-killed *C. albicans* or boiled *C. albicans* into the hindbrain ventricle. N=90 neutrophils from 15 fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance determined by 2-way ANOVA, then Sidák's multiple comparisons test. ns=not significant, *$p<0.05$, ****$p<0.0001$

(C) Representative images of anti-nitrotyrosine stained 24 hpi zebrafish embryos at proximal

and distal sites.

Figure 3: *C. albicans car1Δ* has a role in suppression of neutrophil RNS.

(A) Anti-nitrotyrosine fluorescence at 24 hpi following injection of PVP, *C. albicans* SC5314*,*

heat-killed *C. albicans* SC5314*, C. albicans car1Δ* or heat-killed *C. albicans car1Δ* into the

caudal vein. N=144 neutrophils from 24 fish, obtained from 3 independent experiments.

Error bars show SEM. Statistical significance determined by Kruskal-Wallis test, with Dunn's

multiple comparisons tests. *p<0.05, ****p<0.0001.

(B) Representative images of anti-nitrotyrosine stained embryos. Dashed lines show *C.*

albicans. Scale bars = 50 µm.

Figure 4: Hif-1α stabilisation is protective against *C. albicans* infection.

(A) Survival of *C. albicans* TT21-dTomato infected zebrafish embryos following injection with dominant active *hif-1α* (DA1), dominant negative *hif-1α* (DN1) RNA or phenol red (PR) solvent control. Mortality was measured daily. N=102-103 fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance determined by Gehan-Breslow-Wilcoxon test with Bonferroni correction. **p<0.01, ****p<0.0001.

(B) Representative image of 200 cfu *C. albicans* TT21-dTomato infected zebrafish embryo at 24 hpi. Scale bar = 1000 µm.

(C) Survival curve of 100 cfu *C. albicans* TT21-dTomato infected zebrafish embryos following injection with DA1 or PR, then treatment with PBS or clodronate liposomes. Mortality was measured daily. N=102-103 fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance determined by Gehan-Breslow-Wilcoxon test with Bonferroni correction. ****p<0.0001.

(D) Survival curve of 100 cfu *C. albicans* TT21-dTomato infected zebrafish embryos

following injection with PR or DA1 and control or *csf3r* morpholino. Mortality was measured

daily. N=135 fish, obtained from 3 independent experiments. Error bars show SEM.

Statistical significance determined by Gehan-Breslow-Wilcoxon test with Bonferroni

correction. *p<0.05, **p<0.01.

Figure 5: The host protective effect of Hif-1α stabilisation is RNS-dependent.

(A) Anti-nitrotyrosine fluorescence at 24 hpi in PR and DA1 injected fish, following injection

of PVP or 500 cfu *C. albicans* SN148 GFP. N=108 neutrophils from 18 fish, obtained from 3

independent experiments. Error bars show SEM. Statistical significance determined by

Kruskal-Wallis test, with Dunn's multiple comparisons test. ns=not significant, ***p<0.001,

****p<0.0001

(B) Representative images of 1 dpi Tg(*lyz:NTRmCherry*) zebrafish embryos injected with PR

or DA1 and infected with 500 cfu *C. albicans* SN148 GFP or PVP. Scale bars = 50 μm.

(C) Survival curve of 200 cfu *C. albicans* TT21-dTomato infected zebrafish embryos

following injection with PR or DA1, then treatment with dH$_2$O or L-NIL. N=135 fish, obtained

from 3 independent experiments. Error bars show SEM. Statistical significance determined

by Gehan-Breslow-Wilcoxon test, with Bonferroni correction. **p<0.01, ***p<0.001.

(D) Survival curve of 200 cfu *C. albicans* TT21-dTomato infected zebrafish embryos

following injection with PR or DA1 and *tyr* or *nos2ab* sgRNA. N=135 fish, obtained from 3

independent experiments. Error bars show SEM. Statistical significance determined by

Gehan-Breslow-Wilcoxon test, with Bonferroni correction. **p<0.01, ***p<0.001.

Acknowledgements

498   The authors would like to thank The BSU Aquarium Team for fish care and the IICD

499 Technical Team for practical assistance (University of Sheffield). We would like to thank

500 Professor Neil Gow (University of Exeter) and Dr Robert Wheeler (University of Maine) for

501 kindly providing Candida strains.

Author Contributions

504   Conceived and designed the experiments: TBB, FRH, PTS, AL, SC, AMC, PME.

505 Performed the experiments: TBB, FRH, PTS, AL. Generation of Candida strains: SC, KRA,

506 SAJ. Analysed the data: TBB, FRH, PTS, AL, AMC, PME. Wrote and drafted the manuscript:

507 TBB, FRH, PTS, AL, AMC, PME.

Funding

510   A.L. and P.M.E. are funded by a Sir Henry Dale Fellowship, jointly funded by the

511 Wellcome Trust and the Royal Society (Grant Number 105570/Z/14/A) held by P.M.E. T.B.B

512 is supported by a studentship from the MRC Discovery Medicine North (DiMeN) Doctoral

513 Training Partnership (MR/N013840/1). F.R.H. and P.T.S. were funded by a University of

514 Sheffield PhD scholarship. S.C. was supported by a BBSRC White Rose DTP studentship

515 (BB/M011151/1). S.A.J. was supported by Medical Research Council and Department for

516 International Development Career Development Award Fellowship (MR/J009156/1).

517 Zebrafish infection work was performed in The Wolfson Laboratories for Zebrafish Models of

518 Infection (The Wolfson Foundation/Royal Society grant number WLR\R1\170024) at the

519 University of Sheffield.

521 **Competing Interests**

522 The authors declare that they have no conflict of interest.

524 **References**

525 Bansal, V. and Ochoa, J.B. (2003) 'Arginine availability, arginase, and the immune response',

526 *Current Opinion in Clinical Nutrition and Metabolic Care*, 6(2), pp. 223–228. Available at:

527 https://doi.org/10.1097/00075197-200303000-00012.

528 Bergeron, A.C. *et al.* (2017) 'Polyclonal anti-Candida antibody improves phagocytosis and

529 overall outcome in zebrafish model of disseminated candidiasis', *Developmental &*

530 *Comparative Immunology*, 68, pp. 69–78. Available at:

531 https://doi.org/10.1016/j.dci.2016.11.017.

532 Bernut, A. *et al.* (2019) 'CFTR Protects against Mycobacterium abscessus Infection by Fine-

533 Tuning Host Oxidative Defenses', *Cell Reports*, 26(7), pp. 1828-1840.e4. Available at:

534 https://doi.org/10.1016/j.celrep.2019.01.071.

535 Bongomin, F. *et al.* (2017) 'Global and Multi-National Prevalence of Fungal Diseases—

536 Estimate Precision', *Journal of Fungi*, 3(4). Available at: https://doi.org/10.3390/jof3040057.

537 Brothers, K.M. *et al.* (2013) 'NADPH Oxidase-Driven Phagocyte Recruitment Controls

538 Candida albicans Filamentous Growth and Prevents Mortality', *PLOS Pathogens*, 9(10), p.

539 e1003634. Available at: https://doi.org/10.1371/journal.ppat.1003634.

Brown, G.D. (2011) 'INNATE ANTIFUNGAL IMMUNITY: THE KEY ROLE OF

PHAGOCYTES', *Annual review of immunology*, 29, pp. 1–21. Available at:

https://doi.org/10.1146/annurev-immunol-030409-101229.

Buchan, K.D. *et al.* (2019) 'A transgenic zebrafish line for in vivo visualisation of neutrophil

myeloperoxidase', *PLOS ONE*, 14(4), p. e0215592. Available at:

https://doi.org/10.1371/journal.pone.0215592.

Burgess, T.B., Condliffe, A.M. and Elks, P.M. (2022) 'A Fun-Guide to Innate Immune

Responses to Fungal Infections', *Journal of Fungi*, 8(8), p. 805. Available at:

https://doi.org/10.3390/jof8080805.

Christou, S., Ayscough, K.R. and Johnston, S.A. (2023) 'Candida albicans cells lacking AP-2

have defective hyphae and are avirulent despite increased host uptake and intracellular

proliferation in macrophages'. bioRxiv, p. 2023.08.03.551781. Available at:

https://doi.org/10.1101/2023.08.03.551781.

Collette, J.R., Zhou, H. and Lorenz, M.C. (2014) 'Candida albicans Suppresses Nitric Oxide

Generation from Macrophages via a Secreted Molecule', *PLoS ONE*, 9(4), p. e96203.

Available at: https://doi.org/10.1371/journal.pone.0096203.

Desai, J.V. and Lionakis, M.S. (2018) 'The role of neutrophils in host defense against

invasive fungal infections', *Current clinical microbiology reports*, 5(3), pp. 181–189. Available

at: https://doi.org/10.1007/s40588-018-0098-6.

Drummond, R.A. *et al.* (2019) 'CARD9+ microglia promote antifungal immunity via IL-1β-

and CXCL1-mediated neutrophil recruitment', *Nature Immunology*, 20(5), pp. 559–570.

Available at: http://dx.doi.org/10.1038/s41590-019-0377-2.

Elks, P.M. *et al.* (2011) 'Activation of hypoxia-inducible factor-1α (Hif-1α) delays

inflammation resolution by reducing neutrophil apoptosis and reverse migration in a

zebrafish inflammation model', *Blood*, 118(3), pp. 712–722. Available at:

https://doi.org/10.1182/blood-2010-12-324186.

Elks, P.M. *et al.* (2013) 'Hypoxia Inducible Factor Signaling Modulates Susceptibility to

Mycobacterial Infection via a Nitric Oxide Dependent Mechanism', *PLOS Pathogens*, 9(12),

p. e1003789. Available at: https://doi.org/10.1371/journal.ppat.1003789.

Elks, P.M. *et al.* (2014) 'Mycobacteria counteract a TLR-mediated nitrosative defense

mechanism in a zebrafish infection model', *PloS One*, 9(6), p. e100928. Available at:

https://doi.org/10.1371/journal.pone.0100928.

Ellett, F. *et al.* (2011) 'mpeg1 promoter transgenes direct macrophage-lineage expression in

zebrafish', *Blood*, 117(4), pp. e49–e56. Available at: https://doi.org/10.1182/blood-2010-10-

314120.

Flores-Maldonado, O. *et al.* (2019) 'Neonatal invasive candidiasis: A poorly understood

entity', *Revista Medicina Universitaria*, 21. Available at:

https://doi.org/10.24875/RMU.18000031.

Frohner, I.E. *et al.* (2009) 'Candida albicans cell surface superoxide dismutases degrade

host-derived reactive oxygen species to escape innate immune surveillance', *Molecular

Microbiology*, 71(1), pp. 240–252. Available at: https://doi.org/10.1111/j.1365-

2958.2008.06528.x.

Gantner, B.N., Simmons, R.M. and Underhill, D.M. (2005) 'Dectin-1 mediates macrophage

recognition of Candida albicans yeast but not filaments', *The EMBO journal*, 24(6), pp.

1277–1286. Available at: https://doi.org/10.1038/sj.emboj.7600594.

Gillum, A.M., Tsay, E.Y. and Kirsch, D.R. (1984) 'Isolation of the Candida albicans gene for

orotidine-5'-phosphate decarboxylase by complementation of S. cerevisiae ura3 and E. coli

pyrF mutations', *Molecular & general genetics: MGG*, 198(2), pp. 179–182. Available at:

https://doi.org/10.1007/BF00328721.

van der Graaf, C.A.A. *et al.* (2005) 'Differential Cytokine Production and Toll-Like Receptor

Signaling Pathways by Candida albicans Blastoconidia and Hyphae', *Infection and Immunity*,

73(11), pp. 7458–7464. Available at: https://doi.org/10.1128/iai.73.11.7458-7464.2005.

Gratacap, R.L. *et al.* (2017) 'Control of Mucosal Candidiasis in the Zebrafish Swim Bladder

Depends on Neutrophils That Block Filament Invasion and Drive Extracellular-Trap

Production', *Infection and Immunity*. Edited by G.S. Deepe, 85(9), pp. e00276-17. Available at: https://doi.org/10.1128/IAI.00276-17.

Hammond, F.R., Lewis, A. and Elks, P.M. (2020) 'If it's not one thing, HIF's another: immunoregulation by hypoxia inducible factors in disease', *The FEBS Journal*, 287(18). Available at: https://doi.org/10.1111/febs.15476.

Hromatka, B.S., Noble, S.M. and Johnson, A.D. (2005) 'Transcriptional Response of Candida albicans to Nitric Oxide and the Role of the YHB1 Gene in Nitrosative Stress and Virulence', *Molecular Biology of the Cell*, 16(10), pp. 4814–4826. Available at: https://doi.org/10.1091/mbc.E05-05-0435.

Isles, H.M. *et al.* (2019) 'The CXCL12/CXCR4 Signaling Axis Retains Neutrophils at Inflammatory Sites in Zebrafish', *Frontiers in Immunology*, 10. Available at: https://www.frontiersin.org/articles/10.3389/fimmu.2019.01784 (Accessed: 30 May 2023).

Krzywinska, E. and Stockmann, C. (2018) 'Hypoxia, Metabolism and Immune Cell Function', *Biomedicines*, 6(2), p. E56. Available at: https://doi.org/10.3390/biomedicines6020056.

Lam, S. *et al.* (2004) 'Development and maturation of the immune system in zebrafish, Danio rerio: a gene expression profiling, in situ hybridization and immunological study', *Developmental & Comparative Immunology*, 28(1), pp. 9–28. Available at: https://doi.org/10.1016/S0145-305X(03)00103-4.

Luo, S. *et al.* (2013) 'Complement and innate immune evasion strategies of the human pathogenic fungus Candida albicans', *Molecular Immunology*, 56(3), pp. 161–169. Available at: https://doi.org/10.1016/j.molimm.2013.05.218.

Macherla, C. *et al.* (2012) 'Nitric Oxide Releasing Nanoparticles for Treatment of Candida Albicans Burn Infections', *Frontiers in Microbiology*, 3. Available at: https://www.frontiersin.org/article/10.3389/fmicb.2012.00193 (Accessed: 23 June 2022).

Magill, S.S. *et al.* (2014) 'Multistate Point-Prevalence Survey of Health Care–Associated Infections', *New England Journal of Medicine*, 370(13), pp. 1198–1208. Available at: https://doi.org/10.1056/NEJMoa1306801.

Mayer, F.L., Wilson, D. and Hube, B. (2013) 'Candida albicans pathogenicity mechanisms', *Virulence*, 4(2), pp. 119–128. Available at: https://doi.org/10.4161/viru.22913.

McGettrick, A.F. and O'Neill, L.A.J. (2020) 'The Role of HIF in Immunity and Inflammation', *Cell Metabolism*, 32(4), pp. 524–536. Available at: https://doi.org/10.1016/j.cmet.2020.08.002.

McInturff, A.M. *et al.* (2012) 'Mammalian target of rapamycin regulates neutrophil extracellular trap formation via induction of hypoxia-inducible factor 1 α', *Blood*, 120(15), pp. 3118–3125. Available at: https://doi.org/10.1182/blood-2012-01-405993.

Moore, W.M. *et al.* (1994) 'L-N6-(1-iminoethyl)lysine: a selective inhibitor of inducible nitric oxide synthase', *Journal of Medicinal Chemistry*, 37(23), pp. 3886–3888. Available at: https://doi.org/10.1021/jm00049a007.

Muijsers, R.B.R. *et al.* (2001) 'l-Arginine is not the limiting factor for nitric oxide synthesis by human alveolar macrophages in vitro', *European Respiratory Journal*, 18(4), pp. 667–671. Available at: https://doi.org/10.1183/09031936.01.00101301.

Navarathna, D.H., Lionakis, M.S. and Roberts, D.D. (2019) 'Endothelial nitric oxide synthase limits host immunity to control disseminated Candida albicans infections in mice', *PLOS ONE*, 14(10), p. e0223919. Available at: https://doi.org/10.1371/journal.pone.0223919.

Panpetch, W. *et al.* (2017) 'Oral administration of live- or heat-killed Candida albicans worsened cecal ligation and puncture sepsis in a murine model possibly due to an increased serum (1→3)-β-D-glucan', *PLoS ONE*, 12(7), p. e0181439. Available at: https://doi.org/10.1371/journal.pone.0181439.

Pellon, A., Sadeghi Nasab, S.D. and Moyes, D.L. (2020) 'New Insights in Candida albicans Innate Immunity at the Mucosa: Toxins, Epithelium, Metabolism, and Beyond', *Frontiers in Cellular and Infection Microbiology*, 10. Available at: https://www.frontiersin.org/articles/10.3389/fcimb.2020.00081 (Accessed: 22 July 2022).

Pfaller, M.A. and Diekema, D.J. (2010) 'Epidemiology of invasive mycoses in North America', *Critical Reviews in Microbiology*, 36(1), pp. 1–53. Available at: https://doi.org/10.3109/10408410903241444.

Rementería, A., García-Tobalina, R. and Sevilla, M.J. (1995) 'Nitric oxide-dependent killing

of *Candida albicans* by murine peritoneal cells during an experimental infection', *FEMS*

Immunology & Medical Microbiology, 11(3), pp. 157–162. Available at:

https://doi.org/10.1111/j.1574-695X.1995.tb00112.x.

Renshaw, S.A. *et al.* (2006) 'A transgenic zebrafish model of neutrophilic inflammation',

Blood, 108(13), pp. 3976–3978. Available at: https://doi.org/10.1182/blood-2006-05-024075.

van Rooijen, N. and Hendrikx, E. (2010) 'Liposomes for specific depletion of macrophages

from organs and tissues', *Methods in Molecular Biology (Clifton, N.J.)*, 605, pp. 189–203.

Available at: https://doi.org/10.1007/978-1-60327-360-2_13.

Schaefer, K. *et al.* (2020) 'Three Related Enzymes in Candida albicans Achieve Arginine-

and Agmatine-Dependent Metabolism That Is Essential for Growth and Fungal Virulence',

mBio, 11(4), pp. e01845-20. Available at: https://doi.org/10.1128/mBio.01845-20.

Seman, B.G. *et al.* (2018) 'Yeast and Filaments Have Specialized, Independent Activities in

a Zebrafish Model of Candida albicans Infection', *Infection and Immunity*, 86(10), pp.

e00415-18. Available at: https://doi.org/10.1128/IAI.00415-18.

Stastna, M. and Van Eyk, J.E. (2012) 'Secreted proteins as a fundamental source for

biomarker discovery', *Proteomics*, 12(4–5), pp. 722–735. Available at:

https://doi.org/10.1002/pmic.201100346.

Tillmann, A., Gow, N.A.R. and Brown, A.J.P. (2011) 'Nitric oxide and nitrosative stress

tolerance in yeast', *Biochemical Society Transactions*, 39(1), pp. 219–223. Available at:

https://doi.org/10.1042/BST0390219.

Torraca, V. and Mostowy, S. (2018) 'Zebrafish Infection: From Pathogenesis to Cell Biology',

Trends in Cell Biology, 28(2), pp. 143–156. Available at:

https://doi.org/10.1016/j.tcb.2017.10.002.

Uzun, O. *et al.* (2001) 'Risk factors and predictors of outcome in patients with cancer and

breakthrough candidemia', *Clinical Infectious Diseases: An Official Publication of the*

Infectious Diseases Society of America, 32(12), pp. 1713–1717. Available at:

https://doi.org/10.1086/320757.

Vazquez-Torres, A. *et al.* (1995) 'Nitric Oxide Enhances Resistance of SCID Mice to

Mucosal Candidiasis', *The Journal of Infectious Diseases*, 172(1), pp. 192–198. Available at:

https://doi.org/10.1093/infdis/172.1.192.

Wagener, J. *et al.* (2017) 'Candida albicans Chitin Increases Arginase-1 Activity in Human

Macrophages, with an Impact on Macrophage Antimicrobial Functions', *mBio*, 8(1), pp.

e01820-16. Available at: https://doi.org/10.1128/mBio.01820-16.

Walmsley, S.R. *et al.* (2005) 'Hypoxia-induced neutrophil survival is mediated by HIF-1α–

dependent NF-κB activity', *The Journal of Experimental Medicine*, 201(1), pp. 105–115.

Available at: https://doi.org/10.1084/jem.20040624.

Wellington, M., Dolan, K. and Krysan, D.J. (2009) 'Live Candida albicans Suppresses

Production of Reactive Oxygen Species in Phagocytes', *Infection and Immunity*, 77(1), pp.

405–413. Available at: https://doi.org/10.1128/IAI.00860-08.

WHO (2022) *WHO fungal priority pathogens list to guide research, development and public*

health action. Geneva: World Health Organization. Available at:

https://www.who.int/publications-detail-redirect/9789240060241 (Accessed: 11 May 2023).

Winterbourn, C.C. *et al.* (2006) 'Modeling the reactions of superoxide and myeloperoxidase

in the neutrophil phagosome: implications for microbial killing', *The Journal of Biological*

Chemistry, 281(52), pp. 39860–39869. Available at: https://doi.org/10.1074/jbc.M605898200.

Witherden, E.A. *et al.* (2017) 'The Human Mucosal Mycobiome and Fungal Community

Interactions', *Journal of Fungi*, 3(4), p. 56. Available at: https://doi.org/10.3390/jof3040056.