***Candida spp.* suppress neutrophil reactive nitrogen species to evade killing**

Thomas B. Burgess[1], Ffion R. Hammond[1], Piotr T. Szkuta[1], Amy Lewis[1], Stella Christou[2],

Kieran A Bowden[1], Tihana Bicanic[4], Lynne R Prince[1], Kathryn R. Ayscough[2], David G.

Partridge[3], Simon A. Johnston[1], Alison M. Condliffe[1] and Philip M. Elks[1]

[1]Bateson Centre for Disease Mechanisms and Florey Institute of Infection, School of Medicine

and Population Health, University of Sheffield, Beech Hill Road, Sheffield, S10 2RX, UK.

[2]The School of Biosciences, University of Sheffield, Firth Court, Western Bank, Sheffield, S10

2TN, UK.

[3]Department of Microbiology, Laboratory Medicine, Northern General Hospital, Sheffield

Teaching Hospital NHS Foundation Trust, Sheffield S5 7AU, UK

[4]Institute of Infection and Immunity, City St George's University of London, London SW17 0RE,

UK

Corresponding Author:

Philip M. Elks (p.elks@sheffield.ac.uk)

Abstract

Candida albicans is a human commensal that can cause life-threatening invasive

infection in immunocompromised individuals. Human immunity to *C. albicans* infection is

thought to be largely dependent on neutrophil reactive oxygen and nitrogen species

(ROS/RNS) generation by neutrophils. Despite this, our understanding of innate immune

killing and escape by *C. albicans* is primarily studied in macrophages and the precise

mechanisms of evasion are unclear in neutrophils. Here we sought to determine the

importance of neutrophil reactive nitrogen species (RNS) production during *C. albicans*

infection *in vivo*. Using a zebrafish model, we found that *C. albicans* rapidly downregulated

neutrophil RNS below basal levels during the first day post infection, a time at which neutrophil

RNS is upregulated in bacterial infections as an important host-defense mechanism, indicating

fungal evasion of host neutrophils. We confirmed downregulation of RNS in human primary

neutrophils and with clinical *Candida* spp. isolates, including emerging human pathogens *C.*

auris and *C. glabrata*. Using a zebrafish *arginase 2* transgenic line and a *C. albicans car1Δ*

mutant, we show that both host and fungal arginase contribute the reduction in neutrophil RNS.

Despite pathogen downregulation, upregulation of neutrophil RNS via Hypoxia inducible factor

(Hif)-1α stabilisation, was sufficient to improve *C. albicans* infection survival, dependent on the

presence of neutrophils and Nitric oxide synthase 2 (Nos2). Finally, restoration of neutrophil

RNS, via Hif-1α stabilisation, was synergistic with clinically relevant antifungal treatment,

increasing survival and clearance of *C. albicans* infections. Together, these data demonstrate

that restoration of the neutrophil RNS response in *C. albicans* infection improves infection

outcomes, highlighting the potential of targeting Hif-1α and RNS in host directed therapies

against fungal infections.

Introduction

Candida albicans (*C. albicans*) is a commensal fungus, colonising 70% of healthy individuals (Witherden *et al.*, 2017) and can cause superficial mucosal infections with negligible mortality. However, *C. albicans* can act as an opportunistic pathogen capable of causing life-threatening, invasive disease in immunocompromised individuals (Witherden *et al.*, 2017; Flores-Maldonado *et al.*, 2019; Pellon, Sadeghi Nasab and Moyes, 2020). There are approximately 750,000 cases of severe candidiasis worldwide per year and the estimated mortality rate is high, estimated to be around 50% (Bongomin *et al.*, 2017; WHO, 2022), likely an underestimation due to poor surveillance mechanisms worldwide (Lamoth *et al.*, 2018; Almeida, Rodrigues and Coelho, 2019). This high mortality, combined with a lack of fungal vaccines and rising rates of antifungal resistance, has resulted in *C. albicans* being designated as one of four critical priority fungal pathogens by the World Health Organization, with another *Candida* spp., the emerging human pathogen *Candida auris*, also a critical priority pathogen (WHO, 2022). Hence, urgent research into new therapies is required.

C. albicans is detected by pattern recognition receptors, including Toll-like receptors, C-type lectin receptors and the Rig-I like receptor MDA-5, resulting in a downstream pro-inflammatory response and innate immune cell recruitment (Burgess, Condliffe and Elks, 2022). Although many immune cell types are recruited to the sites of *C. albicans* infection, neutrophils are one of the most potent killers of *C. albicans*, with neutropenic patients being particularly at risk of invasive disease (Uzun *et al.*, 2001; Brown, 2011). Following recruitment, neutrophils have multiple mechanisms of eliminating *C. albicans*, including the production of reactive oxygen (ROS) and nitrogen (RNS) species (Winterbourn *et al.*, 2006). Pharmacological inhibition of nitric oxide synthase (NOS) leads to increased mortality in *C. albicans* infections in mice *in vivo* and reduced candidacidal activity in murine macrophages and peritoneal cells *in vitro*, showing the importance of RNS-mediated fungicidal activity (Rementería, García-Tobalina and Sevilla, 1995; Vazquez-Torres *et al.*, 1995; Vazquez-Torres, Jones-Carson and Balish, 1996; Navarathna, Lionakis and Roberts, 2019). *C. albicans'* ability to rapidly adapt to changes in environmental pH, metabolic flexibility and a robust stress

response allows colonisation of a wide range of niches within the host (Mayer, Wilson and

Hube, 2013). As part of these mechanisms, *C. albicans* is able to subvert the host innate

immune response, including ROS and RNS (Luo *et al.*, 2013). Despite *Candida* spp. being

primarily controlled by neutrophils in human disease, most immune evasion studies have used

murine macrophage cells. This is, in part, because neutrophils are a more challenging cell

type to study *in vitro* due to their short lifespan and ease of accidental activation. *C. albicans*

can suppress ROS and RNS production by macrophages *in vitro* (Wellington, Dolan and

Krysan, 2009; Collette, Zhou and Lorenz, 2014)*.* Culture supernatant also suppressed

macrophage RNS to a lower degree implying secreted compounds are partially involved in

RNS suppression (Collette, Zhou and Lorenz, 2014). How *Candida* spp. evade neutrophil RNS

is a major knowledge gap, critical to understand the pathogenesis of human disease. However,

to overcome challenges associated with studying neutrophils *in vitro*, we require *in vivo* animal

models to understand the neutrophil response in intact tissues.

Zebrafish are a highly tractable *in vivo* model system to investigate the behaviours of

innate immune cells during infection (Henry *et al.*, 2013; Speirs *et al.*, 2024; Lam *et al.*, 2004;

Renshaw *et al.*, 2006). Adaptive immunity does not mature until 4 to 6 weeks post fertilisation,

whereas innate immune cells are present from early larval stages, allowing examination of

innate immune responses to infections without the presence of adaptive immunity (Lam *et al.*,

2004). Zebrafish models of *C. albicans* infection are well established, with a dose-dependent

effect on zebrafish mortality (Chao *et al.*, 2010), and have been used to uncover important

disease mechanisms (Rosowski *et al.*, 2018; Archambault *et al.*, 2019; Blair *et al.*, 2025).

Furthermore, *C. albicans* is able to undergo the yeast-to-hyphae transition in zebrafish larvae,

demonstrating a comparable pathogenesis to human infection (Chao *et al.*, 2010; Gratacap *et*

al., 2017; Seman *et al.*, 2018; Christou *et al.*, 2025).

Stimulation of RNS has the potential to be a host directed therapy to *C. albicans*

infection that would complement existing antifungals. We have previously shown that Hypoxia

inducible factor (Hif)-1α increases neutrophil RNS output, which is protective against the

bacteria *Mycobacterium marinum* infection in zebrafish (Elks *et al.*, 2013). HIF proteins are

important regulators of cellular responses to hypoxia, innate immunity and inflammation

(McGettrick and O'Neill, 2020). HIF-1α proteins are stabilised in hypoxic conditions, or by TLR

signalling in normoxia, triggering transcription of target genes that regulate cellular metabolism,

proliferation, migration, differentiation, angiogenesis and pro-inflammatory mediators

(Peyssonnaux *et al.*, 2005; Jantsch *et al.*, 2011; Krzywinska and Stockmann, 2018; Hammond,

Lewis and Elks, 2020). Therefore, we set out to address whether Hif-1α stabilisation may be

a potential therapeutic mechanism that can enhance neutrophil RNS during *C. albicans*

infections.

Here, we show that *C. albicans* infection suppressed neutrophil RNS in zebrafish

larvae. Importantly, a similar dampening effect on RNS by *C. albicans* was also observed in

human primary neutrophils. RNS suppression was not limited to laboratory strains of *C.*

albicans; it was found during infection with *C. albicans* clinical isolates from non-invasive and

invasive disease and with other *Candida* species. Heatkilled *C. albicans* was only partially

able to decrease neutrophil RNS. Significantly, the ability of strains to reduce neutrophil RNS

positively correlated with virulence *in vivo*. Both fungal arginase (car1) and host arginase

induction were found to contribute to RNS reduction, with a *car1Δ* mutant having a reduced

effect on neutrophil RNS and host arginase induced in zebrafish. Upregulation of host RNS

production via Hif-1α stabilisation enhanced host control of *C. albicans* infection. Restoration

of neutrophil RNS had an additive effect on host survival and clearance of *C. albicans* infection

when combined with antifungals. Together, these data highlight RNS and Hif-1α as potential

targets for host directed therapies against *C. albicans* infections.

Methods

Ethics Statement

Zebrafish were handled in accordance with Animals (Scientific Procedures) Act 1986,

under Project Licenses P1A4A7A5E and PP7684817. Ethical approval was granted by the

University of Sheffield Local Ethical Review Panel. Human blood was collected from healthy

individuals following informed consent. Ethical approval was granted by the University of

Sheffield Research Ethics Committee (study reference number 031773).

Zebrafish Husbandry

Zebrafish were maintained in Home Office approved facilities in Biological Services

Unit (BSU) aquaria at the University of Sheffield, in accordance with standard protocols and

local animal welfare regulations. Adult fish were maintained at 28°C with a 14/10 hour

light/dark cycle. Larvae were maintained at 28°C with a 14/10 hour light/dark cycle in 1X E3

medium (E3 60X stock: 5.0 mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$; diluted

to 1X in distilled water) with 0.001% Methylene Blue (Sigma-Aldrich) to prevent fungal growth

in the first 24 hours. From 24hpf, larvae were kept in E3 without Methylene Blue so that

Candida spp. infection was not impacted. The following zebrafish strains were used: nacre

(wild type), *Tg(mpx:GFP)i114* (Renshaw *et al.*, 2006), *Tg(lyz:nfsB.mCherry)sh260* (Buchan *et*

al., 2019), *Tg(mpeg:nlsClover)*sh436 (Bernut *et al.*, 2019), *TgBAC(arg2:GFP)sh571*

(Hammond *et al.*, 2023).

Candida **spp.** **Culture**

C. albicans strains (Table 1) and *Candida* spp. clinical isolates (Table 2) were used in

this study. *C. albicans* was initially grown on YPD (4001022, MP Bio) plates at 28°C for 48

hours, then transferred to overnight liquid culture in YPD broth in a shaking incubator at 30°C

at 200 rpm. *C. albicans* was heat-killed by incubation in a heat block at 65°C for 1 hour

(Panpetch *et al.*, 2017). To denature fungal proteins, *C. albicans* was boiled at 100°C for 5

minutes.

Table 1: List of *C. albicans* strains

Strain	Genotype	Reference
C. albicans **TT21-dTomato**	*ade2::hisG/ade2::hisG* *ura3::imm434/ura3::imm434::URA3-tetO ENO1/eno1*::ENO1 tetR-ScHAP4AD-3XHA-ADE2 pENO1-dTomato-NATR	(Seman *et al.*, 2018)
C. albicans **SN148 GFP**	SN148 *ENO1/ENO1-GFP::LEU2*	(Christou *et al.*, 2025)
C. albicans **SC5314**	wt; clinical blood isolate	(Gillum, Tsay and Kirsch, 1984)
C. albicans **car1Δ**	As SC5314 but *car1Δcar1Δ*	(Schaefer *et al.*, 2020)
C. albicans **NRG1^{OEX}-dTomato**	e2::hisG/ade2::hisG *ura3::imm434/ura3::imm434::URA3-tetO-NRG1 ENO1/eno1*::ENO1 tetR-ScHAP4AD-3XHA-ADE2 pENO1-dTomato-NATR	(Seman *et al.*, 2018)

163   **Table 2: List of *Candida* spp. clinical isolates**

Species	Name	Invasive/non-invasive	Sample Site	Source
Candida albicans	AJP4	Non-invasive	Vaginal sample	Provided by Dr David Partridge
Candida albicans	AJP5	Invasive	Blood sample	Collected at Sheffield Children's Hospital
Candida albicans	AJP9	Non-invasive	Tongue swab	Collected at Sheffield Children's Hospital
Candida albicans	AJP25	Invasive	Line tip sample	Collected at Sheffield Children's Hospital

Candida glabrata	AJP12	Invasive	Urine sample	Collected at Sheffield Children's Hospital
Candida parapsilosis	AJP22	Invasive	Urine sample	Collected at Sheffield Children's Hospital
Candida guilliermondii	AJP24	Non-invasive	Tracheal swab	Collected at Sheffield Children's Hospital
Candida auris	StG2	Invasive	Bloodstream	Provided by Dr Tihana Bicanic

C. albicans Infections

Overnight liquid *C. albicans* cultures were washed 3 times in phosphate buffered saline

(PBS) and counted on a haemocytometer. Washed cultures were resuspended in 10% PVP

(polyvinylpyrrolidone; Calbiochem) to achieve the desired inoculation. Infection dose for

systemic caudal vein infections were 200 colony forming units (cfu) in a 1nl injection volume,

100 cfu in 1nl for experiments with immune cell ablation or 500 cfu in 1nl in experiments with

antifungals.

For survival curves, zebrafish survival was monitored daily and dead zebrafish were

removed. Death was determined by a cessation of heartbeat/circulation or tissue degradation

post-mortem. At 4 days post infection (dpi; equivalent to 5 days post fertilisation, dpf), all

surviving zebrafish larvae were culled.

RNA Injections

178   Larvae were injected with dominant active *hif-1ab* variant RNA (DA1) or dominant

179 negative *hif-1ab* variant RNA in dH_2O with 10% phenol red (Sigma-Aldrich, to visualise

180 successful injection) at the one-cell stage, as previously described (Elks *et al.*, 2011). 10%

181 (v/v) phenol red in dH_2O (PR; Sigma-Aldrich) was used as a solvent control.

Morpholino Injections

A *csf3r* morpholino (Gene Tools) was used to deplete neutrophil populations (Ellett *et al.*, 2011). A standard control morpholino (Gene Tools) used as a negative control. The antisense morpholino oligonucleotide sequences were: *csf3r* – 5'-GAAGCACAAGCGAGACGGATGCCAT-3'; control – 5'-CCTCTTACCTCAGTTACAATTTATA-3'. *Csf3r* morpholino was functionally validated by whole body neutrophil count at 2 dpf.

F0 CRISPR-Cas9 Knockdown

Two sgRNAs were designed to target the ATG/first exon of *nos2a* and *nos2b* to knockdown zebrafish *nos2* genes by CRISPR-Cas9. Efficacy of knockdown of RNS production was functionally validated by whole-body anti-nitrotyrosine staining. Tyrosinase (*tyr*) sgRNA, targeting a pigment gene that plays no role in neutrophil RNS production, was used as a negative control and to demonstrate Cas9 function (Isles *et al.*, 2019). The target sequences were: *nos2a* – 5'-TTTCTCATTTTCAATGATAG-3'; *nos2b* – 5'-GTTCGCTCTTGTGAGTGACC-3'; *tyr* – 5'-CCTGACCTCCTGAAGACCCC-3'. 25µM sgRNA was co-injected with tracrRNA, 1 µl Cas9 and phenol red.

Pharmacological Treatment of Zebrafish Larvae

For pharmacological inhibition of nitric oxide synthase, larvae were treated with 200 µM N6-(1-Iminoethyl)-lysine (L-NIL) by immersion. dH_2O was used as a solvent control.

For antifungal survival curves and clearance assays, larvae were treated with either 5.0 µg/ml fluconazole in DMSO or 0.5 µg/ml caspofungin in dH_2O by immersion. DMSO or dH_2O was used as a solvent control. At 2 dpi, larvae were transferred into fresh E3 and treatment was re-administered.

Anti-nitrotyrosine Staining

1 dpi larvae (2 dpf) were fixed in 4% formaldehyde in PBS overnight at 4°C. Whole-body nitrotyrosine levels were labelled using a rabbit polyclonal anti-nitrotyrosine antibody (06-284; Merck Millipore) and were detected using Alexa-633 conjugated goat-anti-rabbit secondary antibody (Invitrogen Life Technologies), as previously described (Elks *et al.*, 2013, 2014) in a T*g(mpx:GFP)i114* background to assess for neutrophil specific RNS.

Microscopy and Quantification of Anti-nitrotyrosine Staining

For *C. albicans* SC5314 and *car1Δ,* brightfield images were taken using an inverted Leica DMi8 with a 40x lens and a Hamamatsu OrcaV4 camera. Stained larvae were imaged on a Leica DMi8 inverted microscope with a Leica TCS-SPE line-scanning confocal for imaging with a 40x 1.1NA water immersion lens. For quantification purposes, acquisition settings were kept the same across the groups. Corrected fluorescence intensity was calculated using FIJI measurements assessing the cell fluorescence of individual cells corrected for cell size and background fluorescence of the image (Burgess *et al.*, 2010; Elks *et al.*, 2013, 2014).

Hyphal staging

For assessment of fungal growth of *Candida* spp clinical isolates, which lack fluorescent proteins, brightfield images were taken using an inverted Leica DMi8 with a 40x lens and a Hamamatsu OrcaV4 camera. Classification of hyphal stages was done manually, based on a previously described descriptive criteria (Thrikawala *et al.*, 2022). Criteria for hyphal staging were:

1. Yeast: only spherical, yeast cells were observed. No signs of any hyphal growth.
2. Germinating: ovoid cells, resembling pseudohyphae or very short hyphae, were observed.
3. Hyphal extension: a small number of short or medium length hyphae were observed.

236   4. Destructive hyphae: multiple invasive hyphae were observed. Majority of observed *C.*
237   *albicans* was in hyphal morphotype.

Human neutrophils

Human neutrophils were isolated by Plasma-Percoll density gradient centrifugation of
whole blood from healthy donors as previously described (Herman, Rahman and Prince, 2020).
0.5×10^6 cells were added to individual wells in 96-well plates in 100µl volumes. Cells were
primed with 1 µg/ml lipopolysaccharides from *Escherichia coli* (LPS—Sigma, L8274).
Neutrophils were infected with an MOI of 1.0 or 0.5 *Candida albicans*, MOI 1.0 heat-killed
Candida albicans or uninfected (sterile RPMI media) and were incubated at 37°C, 5% CO_2 for
3 hours. Cells were cytocentrifuged onto microscope slides and fixed in 4% formaldehyde for
staining. Immunostaining was performed as described above.

Microscopy of *C. albicans* Infection clearance

250   Brightfield images were taken using an inverted Leica DMi8 with a 2.5x lens and a
251 Hamamatsu OrcaV4 camera. Surviving fish were classified as 'infection cleared' only if no
252 signs of remaining fluorescent *C. albicans* infection could be observed at 2.5x magnification.

Statistical Analysis

255   All data were analysed using GraphPad Prism 10.5.0 1 (GraphPad Software, La Jolla,
256 California, USA, www.graphpad.com). Nitrotyrosine staining data was analysed with two-tailed
257 Mann-Whitney test or Kruskal-Wallis test, with Dunn's multiple comparisons test. Survival
258 curves were analysed with Gehan-Breslow-Wilcoxon test, with Bonferroni correction. The p
259 values shown are: ns=not significant, *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$

Results

C. albicans infection suppresses neutrophil RNS production below basal levels

Live *C. albicans* caused a 90.0% reduction in the levels of RNS (visualised using an anti-nitrotyrosine antibody) compared to mock infection (PVP) controls (Figure 1A-B), indicating not only a lack of induction of RNS, but also a downregulation of basal RNS. We have previously shown that RNS is predominantly found in neutrophils both in unchallenged and mock infected (PVP) individuals (Elks *et al.*, 2014) and is upregulated after bacterial challenge with *Mycobacterium marinum* (Elks *et al.*, 2013). This established that *C. albicans* supressed neutrophil RNS *in vivo* in contrast to bacterial challenge. In mock infected (PVP) controls there was a basal level of nitrotyrosine levels in neutrophils, as observed previously (Elks *et al.*, 2013, 2014). We tested another bacterium, *Staphylococcus aureus* to determine if induction of neutrophil RNS was limited to mycobacteria. *S. aureus* (SH1000) infection also caused a significant increase in neutrophil nitrotyrosine fluorescence, corroborating previous observations of increased RNS levels in response to mycobacterial infection (Figure S1; Elks *et al.*, 2013). RNS reduction by *C. albicans* was not limited to a single strain as we infected with a different *C. albicans* strain, SN148 GFP, and observed similar RNS suppression (Figure S2). This allowed the use of TT21-dTomato and SN148 GFP interchangeably, depending on the colour combinations required for each experiment. Zebrafish larvae infected with heat-killed C. albicans TT21-dTomato had an intermediate level of RNS suppression, between the PVP control and live C. albicans infection (Figure 1A-B). Hence, the full effect of neutrophil RNS suppression is dependent on live C. albicans suggestive of partially active and passive mechanisms.

To confirm our finding in humans, we performed anti-nitrotyrosine staining on human primary polymorphonuclear neutrophils (PMNs) after *C. albicans* infection *in vitro*. Detectable and robust RNS induction in human neutrophils was induced by priming with LPS which led to a robust nitrotyrosine response after 3 hours in the absence of infection (PVP mock infected) (Figure 1C-D). When *C. albicans* was added at an MOI of 1.0 nitrotyrosine was suppressed to baseline unstimulated levels, despite the presence of LPS (Figure 1C-D). A lower MOI of 0.5

live *C. albicans* showed less of an RNS suppression compared to an MOI of 1.0 of live *C. albicans*, indicating a dose dependent effect (Figure 1C-D). Similarly, an MOI of 1.0 of heat-killed *C. albicans* exhibited lower RNS suppression compared to the same MOI of live *C. albicans*. Together these data show that *C. albicans* suppresses RNS production in human neutrophils as in zebrafish.

The laboratory reference strain SC5314 and strains derived from this (including SN148 and TT21) have inactivating missense mutations in RNAi component Argonaute and so may behave differently to *C. albicans* strains found in the clinic (Iracane *et al.*, 2024). Therefore, clinical isolates from invasive (AJP5 and AJP25) and non-invasive (AJP4 and AJP9) infections were tested for neutrophil RNS suppression in zebrafish. Our clinical isolates lacked a fluorescent protein marker, so brightfield microscopy was used to identify sites of infection in anti-nitrotyrosine stained zebrafish larvae. Both invasive (AJP5 and AJP25) and non-invasive (AJP4 and AJP9) isolates were able to significantly reduce neutrophil RNS compared to mock infection (PVP) controls (Figure 1E-F). These observations demonstrate neutrophil RNS suppression by *C. albicans* is conserved in clinical isolates, implying RNS suppression in *C. albicans* infections may be a clinically relevant phenomenon.

Full neutrophil RNS suppression is dependent on hypha-forming *C. albicans*

C. albicans morphological switch from yeast to hyphae is associated with increased virulence and shifts in the transcriptome and secretome (Nantel *et al.*, 2002; Höfs, Mogavero and Hube, 2016; Vaz *et al.*, 2021). Classification of hyphal growth by *C. albicans* TT21 and clinical isolates in zebrafish revealed the isolates that caused greater neutrophil RNS suppression (AJP9 and AJP25) (Figure 1E-F) exhibited greater hyphal growth *in vivo* compared to those that reduced RNS to a lesser extent (AJP4 and AJP5) (Figure 2A-B). Based on this, we hypothesised hyphal switching may be partially responsible for the suppression of neutrophil RNS. To investigate this, 1 dpf zebrafish were infected with *C. albicans* TT21-dTomato or yeast-locked strain *C. albicans* NRG1^{OEX}-dTomato (Seman *et al.*, 2018). Both *C.*

albicans NRG1^{OEX}-dTomato and heat-killed *C. albicans* NRG1^{OEX}-dTomato caused a modest

decrease in RNS compared to wildtype strains, with reductions comparable to heat-killed *C.*

albicans (Figure 2C-D). These data suggest that an important component of full RNS

suppression by *C. albicans* is hyphal switch-dependent, but that hyphal switch is not essential

for neutrophil RNS suppression.

Host arginase is induced by *C. albicans*

Inducible nitric oxide synthase (iNOS; Nos2 in zebrafish) is the enzyme responsible for

RNS production. iNOS competes with the arginase enzyme for a shared substrate, L-arginine,

and arginase can competitively inhibit RNS production by iNOS (Bansal and Ochoa, 2003).

We investigated the impact of *C. albicans* infection on host arginase levels, as a potential

mechanism by which neutrophil RNS is reduced. PVP injected larvae had a low basal level of

arg2:GFP expression (Figure 3A-B). *Mycobacterium marinum* (used as a control for arginase

induction) caused a significant upregulation of *arg2:GFP* compared to PVP, as previously

described (Hammond et al., 2023). *C. albicans* TT21-dTomato caused a greater upregulation

in *arg2:GFP* expression compared to *M. marinum* indicating a robust host arginase

upregulation. *C. albicans* NRG1^{OEX}-dTomato also caused upregulation of *arg2:GFP* compared

to both PVP and *M. marinum* indicating this is a hyphal-independent process. Examination of

arg2:GFP expression in neutrophils (labelled by *mpx:GFP*) revealed *C. albicans* TT21-

dTomato caused an increase in the percentage of *arg2:GFP*+ neutrophils (41.59% of the

neutrophil population in field of view; Figure 4C-D) compared to PVP (2.01%) and *M. marinum*

(10.97%). Together, these results reveal *C. albicans* infection upregulates host arginase

expression in a subpopulation of neutrophils to a greater level than the professional bacterial

pathogen *M. marinum*.

***C. albicans* derived arginase contributes to the suppression of neutrophil RNS**

Fungi are known to produce enzymes with similar structure and functions to vertebrate

enzymes. *C. albicans* produces its own version of arginase, *CAR1* (Schaefer *et al.*, 2020). We

hypothesised that *C. albicans* arginase interfered with the host RNS production, resulting in neutrophil RNS suppression. 1 dpf zebrafish were infected with *C. albicans car1Δ* or its parental strain *C. albicans* SC5314 and RNS production was measured at 1 dpi by anti-nitrotyrosine staining. *C. albicans car1Δ* infection decreased neutrophil RNS production compared to PVP (Figure 4A-B). *C. albicans car1Δ* caused an intermediate level of RNS suppression between live *C. albicans* SC5314 and heat-killed *C. albicans* SC5314 (Figure 4A-B), indicating that CAR1 contributes to the suppression of neutrophil RNS. Heat-killed *C. albicans car1Δ* was not significantly different from heat-killed *C. albicans* SC5314. Infection with *C. albicans car1Δ* resulted in significantly greater zebrafish larvae survival than infection with *C. albicans* SC5314 (Figure 4C), suggesting RNS suppression by car1 is important for *C. albicans* virulence *in vivo*.

The level of neutrophil RNS suppression by *Candida* spp. correlates with virulence

  C. albicans is the leading causative agent of candidiasis, but there are many other clinically important *Candida* spp. We investigated the effect of a range of *Candida* spp. clinical isolates on neutrophil RNS levels *in vivo*. Clinical isolates of *C. parapsilosis* (AJP22), *C. guilliermondii* (AJP24) and *C. auris* (StG2) all robustly reduced neutrophil RNS compared to PVP controls, but to a lesser extent than a laboratory strain of *C. albicans* (Figure 5A-B). *C. glabrata* AJP12 caused a more modest decrease in neutrophil RNS compared to *C. albicans*. (Figure 5A-B). Together, these results show that neutrophil RNS suppression is conserved across *Candida* spp. clinical isolates, but that levels of RNS suppression can vary between species.

  When percentage zebrafish survival at 4 dpi were plotted against the percentage reduction of RNS levels by *C. albicans* strains/isolates compared to PVP control, survival negatively correlated with reduction in neutrophil RNS levels (Figure 5C. Full survival curves for *Candida* spp. are in Figure S3 and S4.) (Pearson's correlation coefficient: -0.718, 95% confidence interval: -0.208 – -0.921; $p < 0.05$, R^2 value: 0.516). Therefore, a greater

suppression of host RNS levels is correlated with increased virulence in zebrafish infection *in*

vivo, indicating that suppression of neutrophil RNS is a potential virulence mechanism in

human disease-relevant *Candida* spp..

Increased neutrophil RNS, via Hif-1α stabilisation, is protective against *C. albicans*

infection

Hif-1α is a transcription factor with known roles in regulation of innate immunity

(McGettrick and O'Neill, 2020). We have previously shown stabilised Hif-1α (using injection of

a dominant active variant RNA) is protective against *M. marinum* infection *in vivo* by increasing

neutrophil RNS levels (Elks *et al.*, 2013; Hammond, Lewis and Elks, 2020). We therefore

hypothesised that stabilisation of Hif-1α would restore neutrophil RNS in *C. albicans* infection

and increase host survival.

1 cell stage zebrafish larvae were injected with water/10 % phenol red (solvent) control

(PR), dominant negative *hif-1α* (DN *hif-1α*) or dominant active *hif-1α* (DA *hif-1α*) RNA (as

previously described; (Elks *et al.*, 2011)), and subsequently infected systemically with *C.*

albicans TT21-dTomato infection at 30 hpf. DA *hif-1α* larvae had significantly greater survival

over the first 4 days post infection than both PR and DN *hif-1α* larvae (Figure 6A),

demonstrating that Hif-1α stabilisation is host-protective in *C. albicans* infection.

Neutrophils play an essential role in innate immune control of invasive candidiasis

(Desai and Lionakis, 2018). In zebrafish neutrophils are responsible for the killing of *C.*

albicans, whereas macrophages internalise but do not kill (Brothers *et al.*, 2013). Therefore,

we hypothesised that the protective effect of Hif-1α stabilisation would be mediated by

neutrophils rather than macrophages. Firstly, to assess the impact of Hif-1α stabilisation on

macrophage-mediated immunity, we depleted the macrophage population by injection of

clodronate liposomes (van Rooijen and Hendrikx, 2010) at 30 hpf (or PBS liposomes as a

negative control) leading to a significant depletion of the macrophage population (Figure S5).

At 30 hpf, PR or DA *hif-1α* larvae were injected into the caudal vein with clodronate liposomes

or PBS liposomes followed by infection with 100cfu *C. albicans* TT21-dTomato at 2 dpf. Both

groups of PBS liposome-injected larvae had greater survival than their clodronate liposome,

macrophage-depleted, siblings. DA *hif-1α* was host protective both with or without depletion

of macrophages, with DA *hif-1α* + clodronate liposome larvae having greater survival than PR

+ clodronate liposome larvae (Figure 6B; p<0.0001), indicating that the host-protective effect

of Hif-1α stabilisation is not dependent on the presence of macrophages. Next, we injected 1

cell stage zebrafish larvae with a *csf3r* morpholino (Ellett *et al.*, 2011) to deplete neutrophil

numbers by over 60% in 2 dpf zebrafish larvae (Figure S6). *csf3r* morpholino injection led to

a significant decrease in survival of 100 cfu *C. albicans* infected larvae in both PR and DA *hif-*

1α larvae (Figure 6C). DA *hif-1α* + *csf3r* larvae had lower survival compared to PR + *csf3r*

larvae, showing a loss of the protective effect of Hif-1α stabilisation in neutrophil ablated larvae

(Figure 6C), indicating that the protective effect of Hif-1α stabilisation is neutrophil-dependent.

DA *hif-1α* did not alter the recruitment of neutrophils to the site of a local hindbrain *C.*

albicans infection compared to PR injected controls (Figure S7), therefore we hypothesised

that the host protective effect of Hif-1α stabilisation in *C. albicans* infection was due to

increased neutrophil RNS production. DA *hif-1α* doubled the levels of nitrotyrosine compared

to PR in mock infected (PVP) larvae (Figure 7A-B) similar to previous observations (Elks *et*

al., 2013). While *C. albicans* reduced nitrotyrosine levels in PR control injected larvae, DA *hif-*

1α C. albicans infected larvae had high levels of neutrophil nitrotyrosine compared to mock

infected (PVP) DA1 controls (Figure 7A-B).

To demonstrate that the increase in RNS after Hif-1α stabilisation is responsible for

the host protective effect in *C. albicans* infection, we inhibited production of RNS both

pharmacologically, using L-NIL, and genetically, using *nos2a+b* double CRISPR-Cas9

knockdown ("CRISPants"; Moore *et al.*, 1994; Isles *et al.*, 2019). L-NIL treatment did not

impact the growth of *C. albicans in vitro* (Figure S8). Following infection with *C. albicans*, PR

+ L-NIL-treated larvae had reduced survival compared to PR + dH_2O (drug solvent control)-

treated larvae (Figure 7C), indicating a role of RNS in *C. albicans* control. DA *hif-1α* + L-NIL-

treated larvae did not have a significant survival advantage compared to either PR + dH_2O or

PR + L-NIL groups (Figure 7C), demonstrating that iNOS inhibition is sufficient to remove the

host protective effect of Hif-1α stabilisation. *Nos2a+b* double CRISPants were shown to inhibit

nitrotyrosine production in mock infected zebrafish (Figure S9). DA *hif-1α* with *nos2a+b*

CRISPants no longer had increased survival following *C. albicans* infection compared to PR

+ *tyr* sgRNA (*tyrosinase* CRISPant control; Isles *et al.*, 2019) larvae, supporting that Nos2 is

required for Hif-1α stabilisation dependent host protection (Figure 7D). Together these data

demonstrate that the Hif-1α stabilisation is host-protective in *C. albicans* infection via a

neutrophil- (Figure 6) and RNS-dependent (Figure 7) mechanism.

Increased neutrophil RNS has an additive host protective effect when used alongside

antifungals

We next investigated the potential of Hif-1α stabilisation to act as an adjunctive therapy

alongside established antifungals. At 1 dpf, PR/DA *hif-1α* larvae were injected into the caudal

vein with 500 cfu *C. albicans* TT21-dTomato then treated by immersion with 5.0 μg/ml

fluconazole or 1.0 μg/ml caspofungin. Both fluconazole (Figure S10) and caspofungin (Figure

S11) treatment had dose dependent effects on zebrafish larval survival post *C. albicans*

infection. Fluconazole significantly increased host survival in both PR and DA *hif-1α* larvae

(Figure 8A). DA *hif-1α* + fluconazole had increased survival compared to PR + fluconazole

and DA *hif-1α* + DMSO. Combination of DA *hif-1α* + caspofungin similarly increased survival

compared to PR + caspofungin (Figure 8B) and DA *hif-1α* + dH$_2$O (Figure 8B), demonstrating

that combination of Hif-1α stabilisation with fluconazole or caspofungin has an additive effect

on survival.

Surviving larvae were imaged at 4 dpi to quantify clearance of *C. albicans* infection

(Figure 8C-F). 27% of DA *hif-1α* + fluconazole larvae had completely cleared infection at 4 dpi

(Figure 8C-D), greater than both PR alone (4%) and PR + fluconazole (8%). 22% of larvae

treated with DA *hif-1α* + caspofungin cleared infection (Figure 8E-F) compared to 3% of PR-

treated larvae and 4% of larvae treated with PR and caspofungin. Hence, Hif-1α stabilisation

also has an additive effect on clearance when combined with fluconazole or caspofungin.

Discussion

The increasing prevalence of antifungal resistance worldwide highlights the

importance of discovering new antifungal targets or host-directed therapies capable of

improving disease outcomes. Here, we have shown that *Candida* spp*.* suppression of

neutrophil RNS is an important factor in their virulence *in vivo* and that restoration of neutrophil

RNS represents a potential host directed therapy for difficult-to-treat *C. albicans* infection.

We found that *C. albicans* can suppress neutrophil RNS production in zebrafish *in vivo*

and in human primary neutrophils *in vitro*. *C. albicans* has been shown to suppress RNS

production in murine bone marrow derived macrophages *in vitro* (Collette, Zhou and Lorenz,

2014), which is supported by a variety of sources demonstrating suppression of ROS by *C.*

albicans in vitro (Frohner *et al.*, 2009; Wellington, Dolan and Krysan, 2009). While previous

publications have demonstrated RNS suppression in macrophages, our data demonstrates an

effect in neutrophils in a live tissue-setting. Human alveolar macrophages are poor RNS

producers (Muijsers *et al.*, 2001) and *C. albicans* is primarily controlled by neutrophils in

human disease (Gazendam *et al.*, 2016), therefore we chose to dissect the mechanisms

behind RNS suppression in neutrophils in the tractable *in vivo* zebrafish model*.*

Here, we identify that RNS suppression is partially dependent on live, hyphae-forming

C. albicans. The hyphal morphotype is highly associated with a shift in transcription and

increased virulence (Sudbery, 2011; Desai, 2018), which may trigger expression of protein(s)

necessary for RNS suppression by *C. albicans*. Proteomic analysis of supernatant from hyphal

C. albicans in vitro identified 301 secreted proteins specific to hyphae (Vaz *et al.*, 2021).

Further examination and characterisation of these proteins could point towards candidates

involved in active RNS suppression. However, RNS suppression was still observed in non-

hyphae-forming, non-albicans *Candida* spp. isolates (*C. glabrata, C. parapsilosis, C.*

guilliermondii), suggesting that the presence of hyphae themselves is not essential for RNS

suppression.

Host and *C. albicans* derived arginase also contribute to the suppression of neutrophil

RNS production. CAR1 is the only known arginase gene in the *C. albicans* genome, encoding

a cytosolic arginase (Schaefer *et al.*, 2020). Fungal arginase may be interfering with the host

arginase-iNOS balance, resulting in reduced iNOS activity and RNS suppression. The modest

decrease in neutrophil RNS production after infection with the CAR1 mutant suggests that *C.*

albicans arginase production is not the sole RNS reducing mechanism. Our findings suggest

that multiple, additive mechanisms of neutrophil RNS suppression by *C. albicans* may underlie

their success at the large overall suppression of neutrophil RNS observed. *C. albicans* SOD5

has been associated with degradation of antimicrobial ROS *in vitro* (Frohner *et al.*, 2009), *C.*

albicans chitin induces host arginase-1 activity, decreasing RNS production (Wagener *et al.*,

2017), *YHB1* is able to detoxify RNS (Hromatka, Noble and Johnson, 2005) and Collette *et al.*

identified an unknown, small secreted molecule capable of suppressing RNS (Collette, Zhou

and Lorenz, 2014). The presence of chitin may account for the partial suppression of RNS by

heat-killed *C. albicans* that we observed. Further investigation could aim to establish which

are the most important mechanisms and effector proteins involved in RNS suppression in

neutrophils.

Neutrophil RNS suppression was observed in infection with various *C. albicans*

laboratory strains and clinical isolates alongside non-albicans *Candida* spp. clinical isolates.

This suggests that RNS suppression could be conserved widely across clinically relevant

Candida spp. Importantly, *Candida* spp. virulence correlated with neutrophil RNS suppression

in vivo indicating that neutrophil RNS suppression may be a virulence factor. Virulence is

determined by a wide range of factors, not solely neutrophil RNS suppression (Ghannoum

and Abu-Elteen, 1990), any of which could be differentially regulated in these *Candida* spp.

isolates. It was interesting to note that the emerging human pathogen *C. auris* was able to

efficiently suppress neutrophil RNS in zebrafish, despite only been identified as a human

pathogen since 2009 (Lockhart *et al.*, 2017), suggesting that this immune evasion mechanism

may have evolved in the environment (Casadevall, Kontoyiannis and Robert, 2019).

Suppression of neutrophil RNS by environmental strains may act as a useful proxy for

determining whether *Candida* spp. or strains have the potential to be virulent in humans.

We found that upregulation of RNS, via Hif-1α stabilisation, has a protective effect in

C. albicans infection in zebrafish larvae, via a neutrophil-mediated, RNS-dependent

mechanism. Neutrophils are the primary innate immune cell responsible for killing *Candida*

spp. (Brown, 2011). Neutrophil suppression is therefore an important mechanism of *C.*

albicans pathogenesis that, if restored, could represent an exciting therapeutic opportunity.

RNS is well characterised as being candidacidal (Rementería, García-Tobalina and Sevilla,

1995; Vazquez-Torres *et al.*, 1995; Tillmann, Gow and Brown, 2011; Navarathna, Lionakis

and Roberts, 2019) and NO-releasing nanoparticles inhibited the growth of *C. albicans in vitro*

and *in vivo* (Macherla *et al.*, 2012). Hence, upregulation of neutrophil RNS production is a

potential mechanism that could be targeted therapeutically to improve immune candidacidal

activity. We have previously demonstrated Hif-1α stabilisation protects against *M. marinum*

infection in a similar manner (Elks *et al.*, 2013), suggesting Hif-1α stabilisation could offer

protective effect against multiple and different groups of pathogens (bacteria and fungi) where

neutrophil-mediated immune responses are dampened during pathogenesis. Further to this,

we showed restoration of neutrophil RNS had an additive effect on survival and clearance

when combined with traditional antifungals. Early fungicidal activity by neutrophils prior to

hyphal growth is vital for subsequent clearance and resolution of *C. albicans* infection, with

early phagocytosis a strong prognostic indicator of survival (Bergeron *et al.*, 2017; Zhu *et al.*,

2023). As a primarily fungistatic agent (Vasicek *et al.*, 2014), fluconazole may help to restrict

initial growth of *C. albicans*, allowing neutrophils with increased RNS production to kill *C.*

albicans, resulting in greater survival and clearance. Conversely, caspofungin predominantly

has a fungicidal effect (Letscher-Bru and Herbrecht, 2003; Szymański *et al.*, 2022), though

increased neutrophil RNS production may also facilitate earlier killing of *C. albicans* in

combination and allow lower doses of these poorly tolerated drugs for shorter treatment

periods.

Our data shows that *Candida* spp. can efficiently suppress neutrophil RNS as a

virulence mechanism and that Hif-1α stabilisation is able to overcome this, conferring a

neutrophil-mediated, RNS-dependent protective effect *in vivo*. Neutrophil RNS restoration has

the potential as an adjunctive host-directed therapeutic strategy to aid in the treatment of

Candida spp. infections.

Acknowledgements

The authors would like to thank The BSU Aquarium Team for fish care and the SMPH

Technical Team for practical assistance (University of Sheffield). We would like to thank

Professor Neil Gow (University of Exeter), Dr Robert Wheeler (University of Maine) and Dr

Ben Caswall (St George's, University of London) for kindly providing *Candida* spp. strains. For

the purpose of open access, the author has applied a Creative Commons Attribution (CC BY)

licence to any Author Accepted Manuscript version arising

Author Contributions

Conceived and designed the experiments: TBB, FRH, PTS, AL, SC, SAJ, AMC, PME.

Performed the experiments: TBB, FRH, PTS, AL. Generation/provision of *Candida spp*. strains:

SC, TB KRA, DGP. Analysed the data: TBB, FRH, PTS, AL, AMC, PME. Wrote and drafted

the manuscript: TBB, FRH, PTS, AL, SAJ, AMC, PME.

Funding

TBB was supported by a studentship from the MRC Discovery Medicine North (DiMeN)

Doctoral Training Partnership (MR/N013840/1). PME and AL were funded by a Sir Henry Dale

Fellowship, jointly funded by the Wellcome Trust and the Royal Society (Grant Number

105570/Z/14/A) held by PME. This research was funded in whole, or in part, by the Wellcome

Trust (105570/Z/14/A). For the purpose of Open Access, the author has applied a CC BY

public copyright licence to any Author Accepted Manuscript version arising from this

submission. FRH and PTS were funded by a University of Sheffield PhD scholarship. SC was

supported by a BBSRC White Rose DTP studentship (BB/M011151/1). SAJ was supported

by Medical Research Council and Department for International Development Career

Development Award Fellowship (MR/J009156/1) and MRC grant MR/Z505948/1. Zebrafish

infection work was performed in The Wolfson Laboratories for Zebrafish Models of Infection

(The Wolfson Foundation/Royal Society grant number WLR\R1\170024) at the University of

Sheffield.

Competing Interests

571   The authors declare that they have no conflict of interest.

References

Almeida, F., Rodrigues, M.L. and Coelho, C. (2019) 'The Still Underestimated Problem of
Fungal Diseases Worldwide', *Frontiers in Microbiology*, 10, p. 214. Available at:
https://doi.org/10.3389/fmicb.2019.00214.

Archambault, L.S. *et al.* (2019) 'Intravital Imaging Reveals Divergent Cytokine and Cellular
Immune Responses to Candida albicans and Candida parapsilosis', *mBio*, 10(3), pp. e00266-
19. Available at: https://doi.org/10.1128/mBio.00266-19.

Bansal, V. and Ochoa, J.B. (2003) 'Arginine availability, arginase, and the immune response',
Current Opinion in Clinical Nutrition and Metabolic Care, 6(2), pp. 223–228. Available at:
https://doi.org/10.1097/00075197-200303000-00012.

Bergeron, A.C. *et al.* (2017) 'Polyclonal anti-Candida antibody improves phagocytosis and
overall outcome in zebrafish model of disseminated candidiasis', *Developmental &
Comparative Immunology*, 68, pp. 69–78. Available at:
https://doi.org/10.1016/j.dci.2016.11.017.

Bernut, A. *et al.* (2019) 'CFTR Protects against Mycobacterium abscessus Infection by Fine-
Tuning Host Oxidative Defenses', *Cell Reports*, 26(7), pp. 1828-1840.e4. Available at:
https://doi.org/10.1016/j.celrep.2019.01.071.

Blair, B.A. *et al.* (2025) 'Forward genetic screen in zebrafish identifies new fungal regulators
that limit host-protective Candida-innate immune interaction', *mBio*, 16(5), p. e0052925.
Available at: https://doi.org/10.1128/mbio.00529-25.

Bongomin, F. *et al.* (2017) 'Global and Multi-National Prevalence of Fungal Diseases—
Estimate Precision', *Journal of Fungi*, 3(4). Available at: https://doi.org/10.3390/jof3040057.

Brothers, K.M. *et al.* (2013) 'NADPH Oxidase-Driven Phagocyte Recruitment Controls
Candida albicans Filamentous Growth and Prevents Mortality', *PLOS Pathogens*, 9(10), p.
e1003634. Available at: https://doi.org/10.1371/journal.ppat.1003634.

Brown, G.D. (2011) 'INNATE ANTIFUNGAL IMMUNITY: THE KEY ROLE OF PHAGOCYTES',
Annual review of immunology, 29, pp. 1–21. Available at: https://doi.org/10.1146/annurev-
immunol-030409-101229.

Buchan, K.D. *et al.* (2019) 'A transgenic zebrafish line for in vivo visualisation of neutrophil
myeloperoxidase', *PLOS ONE*, 14(4), p. e0215592. Available at:
https://doi.org/10.1371/journal.pone.0215592.

Burgess, A. *et al.* (2010) 'Loss of human Greatwall results in G2 arrest and multiple mitotic
defects due to deregulation of the cyclin B-Cdc2/PP2A balance', *Proceedings of the National
Academy of Sciences*, 107(28), pp. 12564–12569. Available at:
https://doi.org/10.1073/pnas.0914191107.

Burgess, T.B., Condliffe, A.M. and Elks, P.M. (2022) 'A Fun-Guide to Innate Immune
Responses to Fungal Infections', *Journal of Fungi*, 8(8), p. 805. Available at:
https://doi.org/10.3390/jof8080805.

Casadevall, A., Kontoyiannis, D.P. and Robert, V. (2019) 'On the Emergence of Candida auris:
Climate Change, Azoles, Swamps, and Birds', *mBio*, 10(4), p. 10.1128/mbio.01397-19.
Available at: https://doi.org/10.1128/mbio.01397-19.

Chao, C.-C. *et al.* (2010) 'Zebrafish as a Model Host for Candida albicans Infection', *Infection
and Immunity*, 78(6), pp. 2512–2521. Available at: https://doi.org/10.1128/IAI.01293-09.

Christou, S. *et al.* (2025) 'Candida albicans Cells Lacking AP-2 Have Defective Hyphae and
Are Avirulent Despite Increased Host Uptake and Intracellular Proliferation in Macrophages',
Molecular Microbiology [Preprint]. Available at: https://doi.org/10.1111/mmi.70032.

Collette, J.R., Zhou, H. and Lorenz, M.C. (2014) 'Candida albicans Suppresses Nitric Oxide
Generation from Macrophages via a Secreted Molecule', *PLoS ONE*, 9(4), p. e96203.
Available at: https://doi.org/10.1371/journal.pone.0096203.

Desai, J.V. (2018) 'Candida albicans Hyphae: From Growth Initiation to Invasion', *Journal of
Fungi*, 4(1), p. 10. Available at: https://doi.org/10.3390/jof4010010.

Desai, J.V. and Lionakis, M.S. (2018) 'The role of neutrophils in host defense against invasive
fungal infections', *Current clinical microbiology reports*, 5(3), pp. 181–189. Available at:
https://doi.org/10.1007/s40588-018-0098-6.

Elks, P.M. *et al.* (2011) 'Activation of hypoxia-inducible factor-1α (Hif-1α) delays inflammation
resolution by reducing neutrophil apoptosis and reverse migration in a zebrafish inflammation
model', *Blood*, 118(3), pp. 712–722. Available at: https://doi.org/10.1182/blood-2010-12-
324186.

Elks, P.M. *et al.* (2013) 'Hypoxia Inducible Factor Signaling Modulates Susceptibility to
Mycobacterial Infection via a Nitric Oxide Dependent Mechanism', *PLOS Pathogens*, 9(12), p.
e1003789. Available at: https://doi.org/10.1371/journal.ppat.1003789.

Elks, P.M. *et al.* (2014) 'Mycobacteria counteract a TLR-mediated nitrosative defense
mechanism in a zebrafish infection model', *PloS One*, 9(6), p. e100928. Available at:
https://doi.org/10.1371/journal.pone.0100928.

Ellett, F. *et al.* (2011) 'mpeg1 promoter transgenes direct macrophage-lineage expression in
zebrafish', *Blood*, 117(4), pp. e49–e56. Available at: https://doi.org/10.1182/blood-2010-10-
314120.

Flores-Maldonado, O. *et al.* (2019) 'Neonatal invasive candidiasis: A poorly understood entity', *Revista Medicina Universitaria*, 21. Available at: https://doi.org/10.24875/RMU.18000031.

Frohner, I.E. *et al.* (2009) 'Candida albicans cell surface superoxide dismutases degrade host-derived reactive oxygen species to escape innate immune surveillance', *Molecular Microbiology*, 71(1), pp. 240–252. Available at: https://doi.org/10.1111/j.1365-2958.2008.06528.x.

Gazendam, R.P. *et al.* (2016) 'How neutrophils kill fungi', *Immunological Reviews*, 273(1), pp. 299–311. Available at: https://doi.org/10.1111/imr.12454.

Gillum, A.M., Tsay, E.Y. and Kirsch, D.R. (1984) 'Isolation of the Candida albicans gene for orotidine-5'-phosphate decarboxylase by complementation of S. cerevisiae ura3 and E. coli pyrF mutations', *Molecular & general genetics: MGG*, 198(2), pp. 179–182. Available at: https://doi.org/10.1007/BF00328721.

Gratacap, R.L. *et al.* (2017) 'Control of Mucosal Candidiasis in the Zebrafish Swim Bladder Depends on Neutrophils That Block Filament Invasion and Drive Extracellular-Trap Production', *Infection and Immunity*. Edited by G.S. Deepe, 85(9), pp. e00276-17. Available at: https://doi.org/10.1128/IAI.00276-17.

Hammond, F.R. *et al.* (2023) 'An arginase 2 promoter transgenic line illuminates immune cell polarisation in zebrafish', *Disease Models & Mechanisms*, 16(6), p. dmm049966. Available at: https://doi.org/10.1242/dmm.049966.

Hammond, F.R., Lewis, A. and Elks, P.M. (2020) 'If it's not one thing, HIF's another: immunoregulation by hypoxia inducible factors in disease', *The FEBS Journal*, 287(18). Available at: https://doi.org/10.1111/febs.15476.

Henry, K.M. *et al.* (2013) 'Zebrafish as a model for the study of neutrophil biology', *Journal of Leukocyte Biology*, 94(4), pp. 633–642. Available at: https://doi.org/10.1189/jlb.1112594.

Herman, K.D., Rahman, A. and Prince, L.R. (2020) 'Isolation and High Throughput Flow Cytometric Apoptosis Assay of Human Neutrophils to Enable Compound Library Screening', *Bio-protocol*, 10(11). Available at: https://doi.org/10.21769/BioProtoc.3640.

Höfs, S., Mogavero, S. and Hube, B. (2016) 'Interaction of Candida albicans with host cells: virulence factors, host defense, escape strategies, and the microbiota', *Journal of Microbiology*, 54(3), pp. 149–169. Available at: https://doi.org/10.1007/s12275-016-5514-0.

Hromatka, B.S., Noble, S.M. and Johnson, A.D. (2005) 'Transcriptional Response of Candida albicans to Nitric Oxide and the Role of the YHB1 Gene in Nitrosative Stress and Virulence', *Molecular Biology of the Cell*, 16(10), pp. 4814–4826. Available at: https://doi.org/10.1091/mbc.E05-05-0435.

Iracane, E. *et al.* (2024) 'Identification of an active RNAi pathway in Candida albicans', *Proceedings of the National Academy of Sciences*, 121(17), p. e2315926121. Available at: https://doi.org/10.1073/pnas.2315926121.

Isles, H.M. *et al.* (2019) 'The CXCL12/CXCR4 Signaling Axis Retains Neutrophils at Inflammatory Sites in Zebrafish', *Frontiers in Immunology*, 10. Available at: https://www.frontiersin.org/articles/10.3389/fimmu.2019.01784 (Accessed: 30 May 2023).

Jantsch, J. *et al.* (2011) 'Toll-like receptor activation and hypoxia use distinct signaling pathways to stabilize hypoxia-inducible factor 1α (HIF1A) and result in differential HIF1A-

dependent gene expression', *Journal of Leukocyte Biology*, 90(3), pp. 551–562. Available at: https://doi.org/10.1189/jlb.1210683.

Krzywinska, E. and Stockmann, C. (2018) 'Hypoxia, Metabolism and Immune Cell Function', *Biomedicines*, 6(2), p. E56. Available at: https://doi.org/10.3390/biomedicines6020056.

Lam, S. *et al.* (2004) 'Development and maturation of the immune system in zebrafish, Danio rerio: a gene expression profiling, in situ hybridization and immunological study', *Developmental & Comparative Immunology*, 28(1), pp. 9–28. Available at: https://doi.org/10.1016/S0145-305X(03)00103-4.

Lamoth, F. *et al.* (2018) 'Changes in the epidemiological landscape of invasive candidiasis', *Journal of Antimicrobial Chemotherapy*, 73(suppl_1), pp. i4–i13. Available at: https://doi.org/10.1093/jac/dkx444.

Letscher-Bru, V. and Herbrecht, R. (2003) 'Caspofungin: the first representative of a new antifungal class', *Journal of Antimicrobial Chemotherapy*, 51(3), pp. 513–521. Available at: https://doi.org/10.1093/jac/dkg117.

Lockhart, S.R. *et al.* (2017) 'Simultaneous Emergence of Multidrug-Resistant Candida auris on 3 Continents Confirmed by Whole-Genome Sequencing and Epidemiological Analyses', *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*, 64(2), pp. 134–140. Available at: https://doi.org/10.1093/cid/ciw691.

Luo, S. *et al.* (2013) 'Complement and innate immune evasion strategies of the human pathogenic fungus Candida albicans', *Molecular Immunology*, 56(3), pp. 161–169. Available at: https://doi.org/10.1016/j.molimm.2013.05.218.

Macherla, C. *et al.* (2012) 'Nitric Oxide Releasing Nanoparticles for Treatment of Candida Albicans Burn Infections', *Frontiers in Microbiology*, 3. Available at: https://www.frontiersin.org/article/10.3389/fmicb.2012.00193 (Accessed: 23 June 2022).

Mayer, F.L., Wilson, D. and Hube, B. (2013) 'Candida albicans pathogenicity mechanisms', *Virulence*, 4(2), pp. 119–128. Available at: https://doi.org/10.4161/viru.22913.

McGettrick, A.F. and O'Neill, L.A.J. (2020) 'The Role of HIF in Immunity and Inflammation', *Cell Metabolism*, 32(4), pp. 524–536. Available at: https://doi.org/10.1016/j.cmet.2020.08.002.

Moore, W.M. *et al.* (1994) 'L-N6-(1-iminoethyl)lysine: a selective inhibitor of inducible nitric oxide synthase', *Journal of Medicinal Chemistry*, 37(23), pp. 3886–3888. Available at: https://doi.org/10.1021/jm00049a007.

Muijsers, R.B.R. *et al.* (2001) 'l-Arginine is not the limiting factor for nitric oxide synthesis by human alveolar macrophages in vitro', *European Respiratory Journal*, 18(4), pp. 667–671. Available at: https://doi.org/10.1183/09031936.01.00101301.

Nantel, A. *et al.* (2002) 'Transcription Profiling of Candida albicans Cells Undergoing the Yeast-to-Hyphal Transition', *Molecular Biology of the Cell*, 13(10), pp. 3452–3465. Available at: https://doi.org/10.1091/mbc.E02-05-0272.

Navarathna, D.H., Lionakis, M.S. and Roberts, D.D. (2019) 'Endothelial nitric oxide synthase limits host immunity to control disseminated Candida albicans infections in mice', *PLOS ONE*, 14(10), p. e0223919. Available at: https://doi.org/10.1371/journal.pone.0223919.

Panpetch, W. *et al.* (2017) 'Oral administration of live- or heat-killed Candida albicans
worsened cecal ligation and puncture sepsis in a murine model possibly due to an increased
serum (1→3)-β-D-glucan', *PLoS ONE*, 12(7), p. e0181439. Available at:
https://doi.org/10.1371/journal.pone.0181439.

Pellon, A., Sadeghi Nasab, S.D. and Moyes, D.L. (2020) 'New Insights in Candida albicans
Innate Immunity at the Mucosa: Toxins, Epithelium, Metabolism, and Beyond', *Frontiers in*
Cellular and Infection Microbiology, 10. Available at:
https://www.frontiersin.org/articles/10.3389/fcimb.2020.00081 (Accessed: 22 July 2022).

Peyssonnaux, C. *et al.* (2005) 'HIF-1alpha expression regulates the bactericidal capacity of
phagocytes', *The Journal of Clinical Investigation*, 115(7), pp. 1806–1815. Available at:
https://doi.org/10.1172/JCI23865.

Rementería, A., García-Tobalina, R. and Sevilla, M.J. (1995) 'Nitric oxide-dependent killing of
Candida albicans by murine peritoneal cells during an experimental infection', *FEMS*
Immunology & Medical Microbiology, 11(3), pp. 157–162. Available at:
https://doi.org/10.1111/j.1574-695X.1995.tb00112.x.

Renshaw, S.A. *et al.* (2006) 'A transgenic zebrafish model of neutrophilic inflammation', *Blood*,
108(13), pp. 3976–3978. Available at: https://doi.org/10.1182/blood-2006-05-024075.

van Rooijen, N. and Hendrikx, E. (2010) 'Liposomes for specific depletion of macrophages
from organs and tissues', *Methods in Molecular Biology (Clifton, N.J.)*, 605, pp. 189–203.
Available at: https://doi.org/10.1007/978-1-60327-360-2_13.

Rosowski, E.E. *et al.* (2018) 'The Zebrafish as a Model Host for Invasive Fungal Infections',
Journal of Fungi (Basel, Switzerland), 4(4), p. 136. Available at:
https://doi.org/10.3390/jof4040136.

Schaefer, K. *et al.* (2020) 'Three Related Enzymes in Candida albicans Achieve Arginine- and
Agmatine-Dependent Metabolism That Is Essential for Growth and Fungal Virulence', *mBio*,
11(4), pp. e01845-20. Available at: https://doi.org/10.1128/mBio.01845-20.

Seman, B.G. *et al.* (2018) 'Yeast and Filaments Have Specialized, Independent Activities in a
Zebrafish Model of Candida albicans Infection', *Infection and Immunity*, 86(10), pp. e00415-
18. Available at: https://doi.org/10.1128/IAI.00415-18.

Speirs, Z.C. *et al.* (2024) 'What can we learn about fish neutrophil and macrophage response
to immune challenge from studies in zebrafish', *Fish & Shellfish Immunology*, 148, p. 109490.
Available at: https://doi.org/10.1016/j.fsi.2024.109490.

Sudbery, P.E. (2011) 'Growth of Candida albicans hyphae', *Nature Reviews Microbiology*,
9(10), pp. 737–748. Available at: https://doi.org/10.1038/nrmicro2636.

Szymański, M. *et al.* (2022) 'Echinocandins – structure, mechanism of action and use in
antifungal therapy', *Journal of Enzyme Inhibition and Medicinal Chemistry*, 37(1), pp. 876–894.
Available at: https://doi.org/10.1080/14756366.2022.2050224.

Thrikawala, S. *et al.* (2022) 'Cyclooxygenase production of PGE2 promotes phagocyte control
of A. fumigatus hyphal growth in larval zebrafish | PLOS Pathogens', *PLOS Pathogens*, 18(3),
p. e1010040. Available at: https://doi.org/10.1371/journal.ppat.1010040.

Tillmann, A., Gow, N.A.R. and Brown, A.J.P. (2011) 'Nitric oxide and nitrosative stress
tolerance in yeast', *Biochemical Society Transactions*, 39(1), pp. 219–223. Available at:
https://doi.org/10.1042/BST0390219.

Uzun, O. *et al.* (2001) 'Risk factors and predictors of outcome in patients with cancer and
breakthrough candidemia', *Clinical Infectious Diseases: An Official Publication of the*
Infectious Diseases Society of America, 32(12), pp. 1713–1717. Available at:
https://doi.org/10.1086/320757.

Vasicek, E.M. *et al.* (2014) 'Disruption of the Transcriptional Regulator Cas5 Results in
Enhanced Killing of Candida albicans by Fluconazole', *Antimicrobial Agents and*
Chemotherapy, 58(11), pp. 6807–6818. Available at: https://doi.org/10.1128/aac.00064-14.

Vaz, C. *et al.* (2021) 'Mass Spectrometry-Based Proteomic and Immunoproteomic Analyses
of the Candida albicans Hyphal Secretome Reveal Diagnostic Biomarker Candidates for
Invasive Candidiasis', *Journal of Fungi*, 7(7), p. 501. Available at:
https://doi.org/10.3390/jof7070501.

Vazquez-Torres, A. *et al.* (1995) 'Nitric Oxide Enhances Resistance of SCID Mice to Mucosal
Candidiasis', *The Journal of Infectious Diseases*, 172(1), pp. 192–198. Available at:
https://doi.org/10.1093/infdis/172.1.192.

Vazquez-Torres, A., Jones-Carson, J. and Balish, E. (1996) 'Peroxynitrite contributes to the
candidacidal activity of nitric oxide-producing macrophages', *Infection and Immunity*, 64(8),
pp. 3127–3133. Available at: https://doi.org/10.1128/iai.64.8.3127-3133.1996.

Wagener, J. *et al.* (2017) 'Candida albicans Chitin Increases Arginase-1 Activity in Human
Macrophages, with an Impact on Macrophage Antimicrobial Functions', *mBio*, 8(1), pp.
e01820-16. Available at: https://doi.org/10.1128/mBio.01820-16.

Wellington, M., Dolan, K. and Krysan, D.J. (2009) 'Live Candida albicans Suppresses
Production of Reactive Oxygen Species in Phagocytes', *Infection and Immunity*, 77(1), pp.
405–413. Available at: https://doi.org/10.1128/IAI.00860-08.

WHO (2022) *WHO fungal priority pathogens list to guide research, development and public*
health action. Geneva: World Health Organization. Available at:
https://www.who.int/publications-detail-redirect/9789240060241 (Accessed: 11 May 2023).

Winterbourn, C.C. *et al.* (2006) 'Modeling the reactions of superoxide and myeloperoxidase in
the neutrophil phagosome: implications for microbial killing', *The Journal of Biological*
Chemistry, 281(52), pp. 39860–39869. Available at: https://doi.org/10.1074/jbc.M605898200.

Witherden, E.A. *et al.* (2017) 'The Human Mucosal Mycobiome and Fungal Community
Interactions', *Journal of Fungi*, 3(4), p. 56. Available at: https://doi.org/10.3390/jof3040056.

Zhu, W. *et al.* (2023) 'Dual wave of neutrophil recruitment determines the outcome of C.
albicans infection', *Frontiers in Cellular and Infection Microbiology*, 13. Available at:
https://doi.org/10.3389/fcimb.2023.1239593.

Figure Legends:

Figure 1: *Candida albicans* **suppresses neutrophil reactive nitrogen species (RNS) production below resting levels.**

(A) Schematic of experiment: 1 dpf zebrafish were injected with PVP or *C. albicans* into the caudal vein. At 1 dpi, larvae were fixed and stained with anti-nitrotyrosine primary antibody and goat-anti-rabbit Alexa-633 secondary antibody. Zebrafish were then imaged at the site of infection in the caudal haematopoietic tissue (CHT) and anti-nitrotyrosine fluorescence was quantified. Graph shows anti-nitrotyrosine fluorescence at 1 dpi following injection of PVP, live *C. albicans* TT21-dTomato or heat-killed *C. albicans* TT21-dTomato into the caudal vein at 30 hpf. N=96-108 neutrophils from 16-18 fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance determined by Kruskal-Wallis test, then Dunn's multiple comparisons test. P values shown: ****$p < 0.0001$.

(B) Representative images of PVP, *C. albicans* TT21-dTomato and heat-killed *C. albicans* TT21-dTomato infected zebrafish larvae at 1 dpi. Scale bars = 50 µm.

(C) Corrected fluorescence intensity of PMNs fixed at 3 hours post LPS treatment and stained with anti-nitrotyrosine antibody (in arbitrary units, AU). Non-LPS stimulated resting neutrophils, LPS stimulated neutrophils and a secondary antibody only groups were included as controls. Neutrophils infected with an MOI of 0.5 and 1.0 live *C. albicans* and heat-killed *C. albicans* all show a decrease in nitrotyrosine in the presence of LPS compared to LPS-only controls. N=42-48 neutrophils from 2 independent experiments and donors.

(D) Representative images of human PMNs stained with anti-NT. Scale bars = 50 µm. *$p < 0.05$, ****$p < 0.0001$.

(E) Anti-nitrotyrosine fluorescence at 1 dpi following injection of PVP, *C. albicans* TT21-dTomato or *C. albicans* clinical isolates into the caudal vein at 30 hpf. N=144 neutrophils from 24 fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance determined by Kruskal-Wallis test, then Dunn's multiple comparisons test. P values shown: **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

(F) Representative images of PVP, *C. albicans* TT21-dTomato, *C. albicans* AJP4, *C. albicans*

AJP5, *C. albicans* AJP9 and *C. albicans* AJP25 infected zebrafish larvae at 1 dpi. White dotted

circles show *C. albicans.* Scale bars = 50 µm.

Figure 2: Robust RNS suppression is associated with hyphae-forming *C. albicans*

(A) Hyphal classification of zebrafish larvae infected with range of *C. albicans* isolates at 1 dpi.

N=24 fish, obtained from 3 independent experiments.

(B) Representative images of hyphal classification in 1 dpi zebrafish larvae infected with *C.*

albicans TT21-dTomato or *C. albicans* clinical isolate. Images are labelled with the *C. albicans*

isolate. White dotted circles show regions of fungal growth. Scale bar = 50 µm.

(C) Anti-nitrotyrosine fluorescence at 1 dpi following injection of PVP, *C. albicans* TT21-

dTomato, heat-killed *C. albicans* TT21-dTomato*, C. albicans* NRG1^{OEX}-dTomato or heat-killed

C. albicans NRG1^{OEX}-dTomato into the caudal vein. N=120 neutrophils from 20 fish, obtained

from 3 independent experiments. Error bars show SEM. Statistical significance determined by

Kruskal-Wallis test, with Dunn's multiple comparisons tests. $*p<0.05$, $****p<0.0001$.

(D) Representative images of PVP, *C. albicans* TT21-dTomato, heat-killed *C. albicans* TT21-

dTomato*, C. albicans* NRG1^{OEX}-dTomato or heat-killed *C. albicans* NRG1^{OEX}-dTomato

infected larvae at 1 dpi. Scale bars = 50 µm.

Figure 3: *C. albicans* infection stimulates host *arginase 2* expression.

(A) Corrected fluorescence intensity of *Tg(arg2:GFP)* zebrafish larvae at 1 dpi following

injection of PVP, *C. albicans* TT21-dTomato, *C. albicans* NRG1^{OEX}-dTomato or

M. marinum Crimson into the caudal vein at 30 hpf. For PVP, *C. albicans* TT21-dTomato and

C. albicans NRG1^{OEX}-dTomato: n=84-108 cells from 14-18 fish, obtained from 3 independent

experiments. For *M. marinum*: n=48 cells from 8 fish, obtained from 2 independent

experiments. Error bars show SEM. Statistical significance determined by Kruskal-Wallis test,

then Dunn's multiple comparisons test. P values shown: **p<0.01, ****p<0.0001

(B) Representative images of *Tg(arg2:GFP)* zebrafish embryos injected with PVP, *C. albicans*

TT21-dTomato, *C. albicans* NRG1[OEX]-dTomato or *M. marinum* Crimson at 24hpi. White arrow

heads point towards neutrophils. Yellow arrow heads point towards *C. albicans*. Scale bars =

50 µm.

(C) Number of *lyz:nfsB.mCherry*-expressing cells that are *arg2:GFP+* and *arg2:GFP-,*

following infection with *C. albicans* TT21-dTomato, *C. albicans* NRG1[OEX]-dTomato, *M.*

marinum Crimson or PVP. N=113-199 cells from 12 zebrafish, obtained from 2 independent

experiments.

(D) Percentage of *lyz:nfsB.mCherry*-expressing neutrophils that are *arg2:GFP+* and

arg2:GFP-, following infection with *C. albicans* TT21-dTomato, *C. albicans* NRG1[OEX]-dTomato,

M. marinum Crimson or PVP. N=113-199 cells from 12 zebrafish, obtained from 2 independent

experiments.

Figure 4: *C. albicans car1* contributes to suppression of neutrophil RNS.

(A) Anti-nitrotyrosine fluorescence at 1 dpi following injection of PVP, *C. albicans* SC5314*,*

heat-killed *C. albicans* SC5314*, C. albicans car1Δ* or heat-killed *C. albicans car1Δ* into the

caudal vein. N=144 neutrophils from 24 fish, obtained from 3 independent experiments. Error

bars show SEM. Statistical significance determined by Kruskal-Wallis test, with Dunn's

multiple comparisons tests. *p<0.05, ****p<0.0001.

(B) Representative images of anti-nitrotyrosine stained larvae. Dashed lines show *C. albicans*.

Scale bars = 50 µm.

(C) 1 dpf nacre larvae were injected into the caudal vein with 200 cfu *C. albicans* SC5314 or

C. albicans car1Δ. Larval survival was measured daily up to 4 dpi. n=180 fish, obtained from

3 independent experiments. Error bars show SEM. Statistical significance determined by

Gehan-Breslow-Wilcoxon test, with Bonferroni correction. P values shown: ****p<0.0001

Figure 5: Non-albicans *Candida* spp*. suppress neutrophil RNS.

(A) Anti-nitrotyrosine fluorescence at 1 dpi following injection of PVP, *C. albicans* TT21-dTomato, *C. glabrata* AJP12, *C. parapsilosis* AJP22, *C. guilliermondii* AJP24 or *C. auris* StG2 into the caudal vein. N=144 neutrophils from 24 fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance determined by Kruskal-Wallis test, with Dunn's multiple comparisons tests. * indicates difference compared to PVP. † indicates difference compared to *C. albicans* TT21-dTomato. ‡ indicates difference compared to *C. glabrata* AJP12. P values shown: **p<0.01, ***p<0.001, ****p<0.0001.

(B) Representative images of PVP, *C. albicans* TT21-dTomato, *C. glabrata* AJP12, *C. parapsilosis* AJP22, *C. guilliermondii* AJP24 or *C. auris* StG2 infected larvae at 1 dpi. Scale bars = 50 μm.

(C) Reduction in RNS levels compared to PVP control plotted against zebrafish survival at 4dpi, following systemic infection with 200 cfu *Candida* spp*. Full survival curves for *Candida* spp. are in Figures S4 and S5. Each plotted point is based on 3 independent repeats of relevant experiment. Line of best fit and 95% confidence error lines are shown. Pearson's correlation coefficient and R^2 value were calculated.

Figure 6: Hif-1α stabilisation is protective against *C. albicans* infection.

(A) Survival of *C. albicans* TT21-dTomato infected zebrafish larvae following injection with dominant active *hif-1α* (DA1), dominant negative *hif-1α* (DN1) RNA or phenol red (PR) control. Mortality was measured daily. N=102-103 fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance determined by Gehan-Breslow-Wilcoxon test with Bonferroni correction. **p<0.01, ****p<0.0001.

(B) Survival curve of 100 cfu *C. albicans* TT21-dTomato infected zebrafish larvae following injection with DA1 or PR, then treatment with PBS or clodronate liposomes. Mortality was

measured daily. N=102-103 fish, obtained from 3 independent experiments. Error bars show

SEM. Statistical significance determined by Gehan-Breslow-Wilcoxon test with Bonferroni

correction. ****p<0.0001.

(C) Survival curve of 100 cfu *C. albicans* TT21-dTomato infected zebrafish larvae following

injection with PR or DA1 and control or *csf3r* morpholino. Mortality was measured daily. N=135

fish, obtained from 3 independent experiments. Error bars show SEM. Statistical significance

determined by Gehan-Breslow-Wilcoxon test with Bonferroni correction. *p<0.05, **p<0.01.

Figure 7: The host protective effect of Hif-1α stabilisation is RNS-dependent.

(A) Anti-nitrotyrosine fluorescence at 1 dpi in PR and DA1 injected fish, following injection of

PVP or 500 cfu *C. albicans* SN148 GFP. N=108 neutrophils from 18 fish, obtained from 3

independent experiments. Error bars show SEM. Statistical significance determined by

Kruskal-Wallis test, with Dunn's multiple comparisons test. ***p<0.001, ****p<0.0001

(B) Representative images of 1 dpi Tg(*lyz:NTRmCherry*) zebrafish larvae injected with PR or

DA1 and infected with 500 cfu *C. albicans* SN148 GFP or PVP. Scale bars = 50 μm.

(C) Survival curve of 200 cfu *C. albicans* TT21-dTomato infected zebrafish larvae following

injection with PR or DA1, then treatment with dH_2O or L-NIL. N=135 fish, obtained from 3

independent experiments. Error bars show SEM. Statistical significance determined by

Gehan-Breslow-Wilcoxon test, with Bonferroni correction. **p<0.01, ***p<0.001.

(D) Survival curve of 200 cfu *C. albicans* TT21-dTomato infected zebrafish larvae following

injection with PR or DA1 and *tyr* or *nos2a+b* sgRNA. N=135 fish, obtained from 3 independent

experiments. Error bars show SEM. Statistical significance determined by Gehan-Breslow-

Wilcoxon test, with Bonferroni correction. **p<0.01, ***p<0.001.

Figure 8: Hif-1α stabilisation has an additive effect with antifungals.

(A) Survival of *C. albicans* TT21-dTomato infected zebrafish larvae following injection with

dominant active *hif-1α* (DA1) or phenol red (PR) control. Larvae were treated with 5.0 μg/ml

fluconazole or DMSO solvent control. Mortality was measured daily. N=135 fish, obtained from

3 independent experiments. Error bars show SEM. Statistical significance determined by

Gehan-Breslow-Wilcoxon test with Bonferroni correction. **$p<0.01$, ****$p<0.0001$

(B) Survival of *C. albicans* TT21-dTomato infected zebrafish larvae following injection with

DA1or PR control. Larvae were treated with 1.0 µg/ml caspofungin or dH$_2$O solvent control.

Mortality was measured daily. N=135 fish, obtained from 3 independent experiments. Error

bars show SEM. Statistical significance determined by Gehan-Breslow-Wilcoxon test with

Bonferroni correction. **$p<0.01$, ****$p<0.0001$

(C) Clearance of *C. albicans* infection at 4 dpi by zebrafish larvae following injection with DA1

or PR control and treatment with 5.0 µg/ml fluconazole or DMSO solvent control. N=3

independent experiments, 135 fish per group. Error bars show Standard Deviation. Statistical

significance determined by one-way ANOVA, with Tukey's multiple comparisons test. *$p<0.05$,

**$p<0.01$

(D) Infection outcomes of *C. albicans* TT21-dTomato infected zebrafish larvae at 4 dpi,

following injection with DA1 or PR and treatment with 5.0 µg/ml fluconazole or DMSO solvent

control. N=3 independent experiments, 135 fish per group.

(E) Clearance of *C. albicans* infection at 4 dpi by zebrafish larvae following injection with DA1

or PR control and treatment with 1.0 µg/ml caspofungin or dH$_2$O solvent control. N=3

independent experiments, 135 fish per group. Error bars show Standard Deviation. Statistical

significance determined by one-way ANOVA, with Tukey's multiple comparisons test.

(F) Infection outcomes of *C. albicans* TT21-dTomato infected zebrafish larvae at 4 dpi,

following injection with DA1 or PR and treatment with 1.0 µg/ml caspofungin or dH$_2$O solvent

control. N=3 independent experiments, 135 fish per group.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8