Chemical inhibition of stomatal differentiation by perturbation of the master-regulatory

bHLH heterodimer via an ACT-Like domain

Ayami Nakagawa[1]*, Krishna Mohan Sepuru[2,3]*, Shu Jan Alicia Yip[1]*, Hyemin Seo[2,3], Calvin M.

Coffin[2,3], Yasutomo Segawa[4], Rie Iwasaki[1], Hiroe Kato[1], Stephanie Kim[3], Yusuke Aihara[1,5],

Toshinori Kinoshita[1], Kenichiro Itami[1], Soon-Ki Han[1,6], Kei Murakami[1,5,7+], and Keiko U. Torii[1,2,3+]

[1]Institute of Transformative Bio-Molecules, Nagoya University, Nagoya, Aichi, 464-8601, Japan

[2]Howard Hughes Medical Institute, The University of Texas at Austin, Austin, TX 78712 USA

[3]Department of Molecular Biosciences, The University of Texas at Austin, Austin, TX 78712 USA

[4]Institute for Molecular Science and SOKENDAI, Myodaiji, Okazaki, 444-8585, Japan
12
[5]PRESTO, Japan Science and Technology Agency (JST), Chiyoda, Tokyo 102-0076, Japan

[6]Institute for Advanced Research (IAR), Nagoya University, Nagoya, Aichi, 464-8601, Japan

[7]Department of Chemistry, Kwansei Gakuin University, Sanda, Hyogo 669-1330, Japan

*These authors contributed equally

+Corresponding authors

kei.murakami@kwansei.ac.jp

ktorii@utexas.edu

Selective perturbation of protein interactions with chemical compounds enables dissection and control of developmental processes. Differentiation of stomata, cellular valves vital for plant growth and survival, is specified by the basic-helix-loop-helix (bHLH) heterodimers. Harnessing a new amination reaction, we here report a synthesis, target identification, and mode of action of an atypical doubly-sulfonylated imidazolone, Stomidazolone, which triggers stomatal stem cell arrest. Our forward chemical genetics followed by biophysical analyses elucidated that Stomidazolone directly binds to the C-terminal ACT-Like (ACTL) domain of MUTE, a master regulator of stomatal differentiation, and perturbs its heterodimerization with a partner bHLH, SCREAM. Guided by structural docking modeling, we rationally designed MUTE with reduced Stomidazolone binding. These engineered MUTE proteins are fully functional and confer Stomidazolone resistance *in vivo*. Our study identifies doubly-sulfonylated imidazolone as a direct inhibitor of the stomatal master regulator, further expanding the chemical space for perturbing bHLH-ACTL proteins to manipulate plant development.

INTRODUCTION

Plant growth and survival rely on stomata ~ adjustable vales on the shoot epidermis of land plants

facilitating efficient gas exchange while minimizing water loss. Each stoma consists of paired

guard cells that can adjust the aperture in response to the environment, such as light, drought,

and CO_2 levels [1]. A wealth of studies has elucidated the molecular basis of signaling mechanisms

leading to stomatal opening and closure [2,3]. In addition to stomatal movement, stomatal

development is also influenced by environmental conditions and in turn impacts plant's water-use

efficiency [4,5]. During leaf development, stomata are generated through a stereotypical sequence

of cell division and differentiation events [6,7]. In dicotyledonous plants, including Arabidopsis, the

stomatal-cell lineage is initiated via *de novo* asymmetric cell divisions in the protoderm, which

create meristemoids, proliferating stomatal precursor cells with a stem-cell-like property. After a

few rounds of asymmetric divisions, the meristemoid differentiates into a guard mother cell (GMC).

The GMC then undergoes a single symmetric division to terminally differentiate a stoma [8].

The cell-state transitional steps during stomatal development are governed by the

sequential actions of basic-helix-loop-helix (bHLH) transcription factors (TFs), SPEECHLESS

(SPCH), MUTE, and FAMA, which form heterodimers with their partner bHLHs, SCREAM (SCRM,

also known as ICE1) and SCRM2 [9-12]. The activities of these stomatal bHLH modules are enforced

by the inhibitory peptide-receptor kinase signaling pathways. Briefly, secreted peptide ligands

EPIDERMAL PATTERNING FACTORs (EPFs) are perceived by their receptors, ERECTA-family

receptor kinases and TOO MANY MOUTHS (TMM) [13-18]. This activates a downstream Mitogen

Activated Protein Kinase (MAPK) cascade, which in turn inhibits the bHLH heterodimers via direct

phosphorylation and degradation events [19-21].

Interestingly, these stomatal bHLH TFs possess a C-terminal ACT-Like (ACTL) domain, a

small structurally defined module resembling the ACT domain found in diverse metabolic

enzymes in bacteria and eukaryotes (named after <u>a</u>spartate kinase-<u>c</u>horismate mutase-<u>t</u>yrA

prephenate dehydrogenase) [21,22]. Recently, we revealed that the SCRM ACTL domain serves as

a heterodimer selectivity interface with SPCH, MUTE, and FAMA [23]. This rather surprising finding

offers a potential for the ACTL domain of bHLH TFs as a drug target for selective inhibition of

heterodimerization, hence stomatal differentiation. Whereas the ACTL domain is predominant

amongst plant bHLH proteins [24,25] no chemical inhibitors or binders targeting this domain, or more

broadly, bHLH-ACTL TFs are available.

Chemical compounds that selectively perturb functions and interactions of key regulatory

proteins can be a powerful tool to dissect and manipulate the development and physiology of cells

to whole organisms [26,27]. Stomata are not the exception [28]. For example, previous chemical
screens identified compounds that suppress stomatal movement [29,30]. Likewise, small compounds
have been utilized to examine the roles of signaling kinases and phosphatases in stomatal
development [31-33]. Chemical library screens identified bubblin, which severely disrupts proper cell
polarity in dividing meristemoids [34], as well as those that increase the number of stomata [35].
However, in all the above cases, the chemical compounds impact multiple processes, thus
causing pleiotropic effects and/or their targets remain unidentified.

To expand a chemical space to control stomatal function and development, we have
produced a collection of nitrogen-containing chemical compounds by utilizing a new amination
reactions [36]. Nitrogen-containing molecules are widely recognized for their potent biological
activities [37]. Previous chemical screens and subsequent structure-activity-relationship studies
have shown that sulfonamidated compounds SIM1 (Stomata Influencing Molecule) and its
derivative SIM3* inhibit stomatal movement [38]. Here, we report a complex doubly-sulfonylated
imidazolone compound named Stomidazolone as a potent inhibitor of stomatal differentiation. We
discover that Stomidazolone tightly binds to the MUTE ACTL domain and perturbs the MUTE-
SCRM heterodimerization. We rationally designed Stomidazolone-resistant versions of MUTE
and demonstrated their efficacy *in vivo*. Together with our chemical synthesis pipelines, our work
will expand the chemical and pharmacological space for bioactive compounds and open the future
directions for selective perturbation of prominent ACTL-domain containing plant bHLH TFs for
transient manipulation of plant growth and development.

100
101

RESULTS

Doubly-sulfonylated imidazolone inhibits stomatal differentiation
In our initial attempt to derivatize SIM1 [38], we synthesized SIM1* from 2,4-diphenylimidazole
through C–H imidation followed by monodesulfonylation, we serendipitously found the first one-
shot synthesis of doubly-sulfonylated imidazolone (Fig. 1a; See Methods). Treatment of 2,4-
diphenyloxazole with chloramine B in the presence of CuI afforded this doubly-sulfonylated,
AYSJ929 (later named as Stomidazolone: Fig. 1a, Document S1 for synthesis and NMR spectra
of the synthesized compounds). Further optimization produced Stomidazolone in 42% yield. Our
new reaction enabled the conversion of diphenyloxazole into substituted imidazolone with two
equivalents of chloramine (see Document S1). Even after searching the Reaxys database, we
failed to uncover any records of a reaction that transforming diphenyloxazole into imidazolones.

The structure was unambiguously confirmed by X-ray crystallographic analysis, which clearly
shows the acute angular shape of this doubly-sulfonylated imidazolone (Fig. 1b, Table S1,
Supplementary Document S1). Unlike SIM1 and its derivative SIM3* [38], this compound did not
exhibit any discernable effects on stomatal movement using our standardized assays (Fig. S1).

The synthesized compound features numerous aryl groups, oxygen and nitrogen atoms
as hydrogen bond acceptors, as well as hydrogen atoms serving as hydrogen bond donors (see
Fig. 1b; Document S1), suggestive of potential bioactivity. We thus subjected it to our chemical
library screening pipeline for stomatal development [35]. Intriguingly, its application to Arabidopsis
seedlings conferred reduced number of stomata but concomitant increase in the number of
meristemoids (Fig. 1c). These increased meristemoids properly express both early and late
meristemoid reporter markers, *TMMpro::GUS-GFP* and *MUTEpro::nucYFP*, respectively (Fig. 1d).
Further quantitative analysis of stomata and meristemoid densities confirmed the dose-
dependency of the compound, now named Stomidazolone, to decrease the number of stomata
(24.45 %, 59.50 %, and 69.53 % decrease from mock to 20, 50, and 100 μM Stomidazolone,
respectively) while dramatically increasing the number of meristemoids (315.8 %, 801.5 %, and
647.6 % increase from mock to 20, 50, and 100 μM Stomidazolone, respectively) (Fig. 1e). On
the other hand, total number of stomata and meristemoids are only slightly reduced by the
Stomidazolone treatment (14.46 % and 25.03% at 20 and 50 μM) (Fig. 1e). These results suggest
that Stomidazolone inhibits stomatal differentiation at the transition from proliferating
meristemoids to differentiation. Stomidazolone does not affect overall seedling growth including
root elongation and cotyledon expansion, even at 100 μM (Fig. 1f). It is thus highly unlikely that
the observed inhibition of meristemoid differentiation is due to general seedling growth retardation.
We therefore conclude that Stomidazolone potently triggers the developmental arrest of the
meristemoids, and in a lesser extent, inhibits the stomatal-lineage initiation.
138

Chemical genetics delineate Stomidazolone as a potential inhibitor of MUTE

To unravel which pathways of stomatal development are impacted by Stomidazolone, we took a
chemical-genetic approach utilizing a series of mutants known to regulate stomatal development
(Fig. 2a). SPCH, MUTE, and FAMA consecutively drive stomatal-lineage progression [9-11] (Fig.
2a). To address whether Stomidazolone interferes with each cell-transitional step specified by
these stomatal bHLH proteins, we applied Stomidazolone to *spch*, *mute*, and *fama* mutant
seedlings. As shown in Fig. 2b and c, Stomidazolone exerted no effects on the pavement cell-
only epidermis of *spch*. Likewise, the arrested meristemoid phenotype of *mute* was not enhanced
by Stomidazolone. On the other hand, Stomidazolone significantly reduced the number of *fama*

tumors whereas increased the arrested meristemoids (Fig. 2c,d). Thus, *spch* and *mute* mutations

are epistatic to Stomidazolone, suggesting that this compound inhibits the MUTE-step of stomatal

fate commitment.

It is established that *ERECTA*-family genes interact synergistically to enforce stomatal

development [14]. Among them, *ERECTA* primarily inhibits the entry into stomatal cell lineages, a

process regulated by *SPCH*, whereas *ERECTA-LIKE1* (*ERL1*) restricts the differentiation

potential of the meristemoids, a process orchestrated by *MUTE* [39-41](Figs. 2a). The loss of function

in these receptor genes compromise the MAPK activation that inhibits SPCH and MUTE protein

accumulation [19,20,40]. Compared to wild type, *erecta* null allele (*er*: *er-105*) overly produces

stomatal-lineage cells [14] (Fig. 3a-d). Stomidazolone treatment exaggerated the *er-105* phenotype:

the number of stomata drastically reduced and the meristemoids vastly increased (Fig. 3a-d, k).

In *er erl1*, Stomidazolone treatment did not decrease the number of stomata, although it increased

the number of meristemoids (Fig. 3e, f, k). The result supports the previous finding that the loss

of *ERL1* intensifies stomatal differentiation by over-accumulation of MUTE [40], thereby conferring

partial resistance to Stomidazolone. The number of arrested meristemoids are even more

increased in *er erl2* (Fig. 3g, h), consistent with the loss of *ERL1* in this genetic background.

Finally, *er erl1 erl2* triple mutant seedlings appeared insensitive to Stomidazolone (Fig. 3i, j).

Combined, these rather complex results further support that Stomidazolone primarily interferes

with the MUTE activity for commitment to stomatal differentiation. Our results also suggest that if

ERECTA-family mediated signaling pathways are absent, Stomidazolone cannot effectively

interfere with the overly activated stomatal differentiation pathway.

It is known that HD-ZIP IV-family gene *HDG2* positively regulates stomatal differentiation

by promoting *MUTE* expression and that its paralog *AtML1* enhances the *hdg2* mutant phenotype

[42] (Fig. 2a). We next tested whether loss-of-function mutations in these genes enhance the effects

of Stomidazolone to trigger meristemoid arrests. As shown in Fig. 3l, m, Stomidazolone increased

the meristemoid numbers of *hdg2* epidermis while the total numbers of stomata and meristemoids

were unaffected. Strikingly, *hdg2 atml1* double mutant exhibited hypersensitivity to

Stomidazolone, and its application exaggerated *mute*-like epidermal phenotype (Fig. 3n, o).

Taken together, a series of chemical-genetic dissections indicates that Stomidazolone likely

inhibits the activity of MUTE.

***In vivo* potency and selectivity of Stomidazolone relies on SCREAM**

SCRM is a heterodimeric partner of MUTE as well as SPCH and FAMA, and its activity is required

for stomatal cell-state transitions [12] (Fig. 2a). We investigated whether the activity of SCRM

influences the effects of Stomidazolone on stomatal differentiation. Strikingly, Stomidazolone
treatment to *scrm* loss-of-function mutant (also known as *ice1-2*) severely inhibited stomatal
lineage initiation, resulting in the *spch*-like epidermis vastly composed of pavement cells (Fig. 4a,
b). Conversely, Stomidazolone failed to inhibit the 'stomata-only' epidermis of *scrm-D*, a
constitutively active *SCRM* mutation that breaks the molecular bridge between MAPK cascade
and SPCH [12,21](Fig. 4c, d). Therefore, Stomidazolone's potency as an inhibitor of stomatal
differentiation relies on the activity of SCRM: In the absence of SCRM, Stomidazolone can
effectively inhibit the SPCH-regulated step of stomatal-lineage initiation. Under normal levels of
SCRM (i.e. in wild type), Stomidazolone predominantly triggers meristemoid arrest. Lastly, the
constitutively active form of SCRM is insensitive to Stomidazolone.
Next, we sought to delineate if Stomidazolone interferes with the *in vivo* heterodimerization
potential of SCRM. Recently, we have shown that the SCRM C-terminal ACTL domain serves as
a heterodimerization interface and its dysfunctions selectively abrogate heterodimerization of
SCRM with MUTE, but not SPCH [23]. We therefore used our sensitized mutants, *scrm-D_S343*
and *scrm-D_S423*, which encode scrm-D$_{L484F}$ and scrm-D$_{\Delta ACTL}$, respectively [23]. Whereas both
scrm-D_S343 and *scrm-D_S423* exhibit nearly wild-type epidermis, Stomidazolone triggered
excessive formation of arrested meristemoids [23] (Fig. 4 e-h). This phenotype resembles that of
scrm-D mute [43]. Taken together, our intricate *in vivo* phenotypic evidence highlights that
Stomidazolone inhibits meristemoid differentiation, most likely by disrupting the
heterodimerization of MUTE and SCRM through the C-terminal ACTL domain.
202

Stomidazolone directly targets MUTE and interferes with its heterodimerization
To address the molecular action of Stomidazolone, we investigated whether Stomidazolone
directly disrupts the heterodimerization of MUTE (as well as other stomatal-lineage bHLHs) with
SCRM. First, we performed yeast two-hybrid (Y2H) assays. The growth assays show that
Stomidazolone diminishes the association of SCRM with MUTE as well as with SPCH and FAMA
(note that the N-terminal activation domain of SPCH and FAMA have been removed to prevent
their autoactivation)[21] (Fig. 5a). Further quantitative α-galactosidase assays show that
Stomidazolone significantly inhibit these stomatal bHLH heterodimers, with slightly less impact on
the SPCH-SCRM heterodimers (Fig. 5b). A precursor analog of Stomidazolone, SIM3* acts as an
inhibitor of stomatal movement [38] (Fig. S2a). Unlike Stomidazolone, SIM3* has no effects on
stomatal development (Fig. S2b, c). Likewise, SIM3* does not affect heterodimerization of
stomatal bHLH proteins (Fig. S2d, e).

To quantitatively assess the direct inhibition of MUTE-SCRM heterodimers by
Stomidazolone, we next performed two *in vitro* biophysical protein-protein interaction kinetics
assays using biolayer interferometry (BLI) and isothermal titration calorimetry (ITC) (Fig. 5c, d;
Fig. S3, Table S2). For this purpose, recombinant full-length SCRM and MUTE proteins were
expressed and purified. Preincubation of Stomidazolone and MUTE decreased the binding affinity
of MUTE and SCRM by two orders of magnitude for both BLI and ITC methods (Kd values
increased from 7.5 ± 1.2 nM to 856.5 ± 15.1 nM in BLI and from 10.3 ± 2.1 nM and 1400 ± 40 nM
in ITC for MUTE-SCRM binding with or without pre-incubation with Stomidazolone, respectively)
(Figs. 5c, d, Fig. S3). These results demonstrate that Stomidazolone directly interferes with the
MUTE-SCRM heterodimerization.

We next tested whether Stomidazolone directly binds to MUTE. Indeed, the BLI assay
showed Stomidazolone binding to MUTE (Kd = 8.9 ± 1.2 μM) (Fig. 5e). In addition, we found that
Stomidazolone binds with recombinant SPCH, FAMA, and SCRM proteins, albeit with lower
affinity (Fig. 5e, Fig. S3, Table S3). Finally, to identify the binding domain of Stomidazolone on
MUTE, we performed BLI assays with the purified recombinant bHLH domain and ACTL domain
of MUTE. As shown in Fig. 5f, Stomidazolone specifically binds to the ACTL domain (Kd = 13.1 ±
1.3 μM). Again, Stomidazolone showed a lower affinity with the ACTL domains of SCRM, SPCH,
and FAMA (Fig. S3, Table S3). Taken together, a series of yeast and *in vitro* studies demonstrates
that Stomidazolone directly and preferentially targets the ACTL domain of MUTE and interferes
with the MUTE-SCRM heterodimerization. Our results reinforce the previous finding that the ACTL
domain integrity is strictly required for the heterodimerization of MUTE-SCRM, but not SPCH-
SCRM or FAMA-SCRM [23], and further provide the molecular, mechanistic basis of Stomidazolone
as an inhibitor of meristemoid differentiation.

Docking modeling identifies the Stomidazolone-MUTE binding interface
We demonstrated that the ACTL domain of MUTE is targeted by Stomidazolone (Fig. 5). To
delineate the specific binding interface, we performed computational docking modeling (see
Methods). Briefly, we used HADDOCK-based calculations to drive the docking process, using
mutagenesis data and binding site information from Swiss Dock as ambiguous interaction
restraints. Structural models showed that different racemic mixtures of Stomidazolone adopted
the same geometry in all clusters, with ionic interactions mediated by charged residues within the
MUTE ACTL domain, Arg-133 (R133), Arg-134 (R134), Glu-162 (E162), and Glu-163 (E163), and
H-bonding and packing interactions by polar and hydrophobic residues Ile135, Val136, Thr164

and Val165 (Fig. 6a, b). The binding free energies of both racemic mixtures were similar (Table

S4).

Amino-acid sequence alignments of MUTE and its orthologs as well as their sister bHLHs

(SPCH and FAMA) show that R133 and R134 residues are highly conserved among the MUTE

and FAMA clades, whereas substitutions to His (H) and Lys (K) are prevalent in the SPCH clade

(Fig. S4). The E162 and E163 residues are replaced by Glu (E) and Gln (Q) and Asp (D) and D/E

in FAMA and SPCH clades, respectively (Fig. S4). Overall, the Stomidazolone-

SPCH/MUTE/FAMA ACTL domain binding interfaces possess positively and negatively charged

residues with clade-specific sequences, which might reflect the differences in physical binding

affinities (Fig. 5e, Fig. S2).

To confirm the predicted binding residues for Stomidazolone-MUTE interaction, we further

performed Alanine substitutions. For this purpose, we generated MUTE$_{R133A\_R134A}$,

MUTE$_{E162A\_E163A}$, and MUTE$_{R133A\_R134A\_E162A\_E163A}$ mutants and then quantitatively analyzed

Stomidazolone binding affinity on these mutant versions of MUTE ACTL domain by using

biophysical assays, BLI and ITC (Fig. 6c, Fig. S2 and Table S3). In both assays, the binding

affinity was significantly reduced; The Kd values obtained for BLI assays are 85.6 ± 5.5 μM and

125.0 ± 12.0 μM for double mutants, and no binding was detected for the quadruple mutant (Fig.

6c). These results provide the experimental evidence for the predicted Stomidazolone-binding

interface of specific MUTE residues within its ACTL domain.

Engineering Stomidazolone-resistance via a molecular docking modeling

Our molecular docking modeling and experimental verification deciphered the Stomidazolone

binding interface at the MUTE ACTL domain (Fig. 6a, b). To the best of our knowledge, nothing

is known about the bioactivity of doubly-sulfonylated imidazolones nor drugs that target the ACTL

domains of bHLH TFs. We thus first explored the functional importance of the Stomidazolone-

binding sites within the MUTE ACTL domain. We expressed two different *MUTE* mutant versions

that are diminished in Stomidazolone binding (Fig. 6c) into *mute* knockout (*mute-2*) plants driven

by its own promoter (*MUTEpro::MUTE$_{E162A\_E163A}$-GFP* and *MUTEpro:: MUTE$_{R133A\_R134A}$-GFP*). We

in addition generated a control *MUTEpro::MUTE-GFP* construct using the identical molecular

cloning strategy, which was also transformed into *mute-2* in parallel (see Methods). Like the wild-

type MUTE-GFP, the engineered versions of MUTE-GFP are detected in the nucleus of a

differentiating meristemoid, and their expression fully rescued the *mute* arrested meristemoid

phenotype (Fig. 6d, e, Fig. S4a). These transgenic *mute* null lines grow into healthy, fertile plants

indistinguishable from the wild type. Thus, the site-directed mutations within the predicted

Stomidazolone-MUTE interaction interface do not impact the expression patterns or
functionalities of MUTE.

Next, we tested the effects of Stomidazolone. As expected, control *MUTEpro::MUTE-GFP*
mute-2 seedlings significantly reduced the number of stomata upon Stomidazolone treatment (Fig.
6e, f, Fig. S5b, c). In contrast, independent lines of *MUTEpro::MUTE$_{E162A\_E163A}$-GFP* and
MUTEpro:: MUTE$_{R133A\_R134A}$-GFP exhibit Stomidazolone resistance, because Stomidazolone
treatment did not significantly affect the numbers of stomata and meristemoids (Fig. 6e, f, Fig.
S5b, c). These results validate the structurally modelled Stomidazolone-MUTE binding interface
and unambiguously demonstrate that the *in vivo* binding of Stomidazolone to the MUTE ACTL
domain is the cause of the inhibition of stomatal differentiation. Furthermore, our structure-guided
engineering of Stomidazolone-resistant plants suggests that the Stomidazolone-binding interface
within the ACTL domain can be uncoupled from the biological activity of the bHLH-ACTL TFs.

294
295

DISCUSSION
In this study, we developed a new catalytic amination and materialized a synthesis of
Stomidazolone, a complex, doubly-sulfonylated imidazolone that selectively inhibits stomatal
differentiation. Leveraged by the wealth of stomatal development mutant resources, our forward
chemical genetics pinpointed the exact *in vivo* target of Stomidazolone: the master regulator
MUTE. Further biochemical, biophysical, and structural modeling studies elucidated that
Stomidazolone perturbs the MUTE-SCRM heterodimerization through interacting with the ACTL-
domain of MUTE. Chemical synthesis of TF dimerization inhibitors is an under/unexplored field in
plant biology. As such, our work expands a chemical space for manipulation of developmental
processes.

Stomidazolone binds strongly to MUTE ACTL domain, but also exhibits weak *in vitro*
binding to other bHLH-ACTL proteins, including SPCH and FAMA, and interferes with their
heterodimerization *in vitro* and in yeast (Fig. 5e, Fig. S2). Nevertheless, selective inhibition of
meristemoid-to-GMC transition indicates that the MUTE-SCRM heterodimer is the primary *in vivo*
Stomidazolone target. It has been shown that heterodimerization of MUTE-SCRM, but not SPCH-
SCRM or FAMA-SCRM, relies on the presence or integrity of the SCRM ACTL domain [23]. Our
findings support the notion that the MUTE-SCRM pair is the most labile, hence druggable,
amongst the stomatal bHLH proteins. SPCH, MUTE, and FAMA are closely related paralogs. Yet,
they possess distinct domains, which may influence the efficacy of Stomidazolone. For instance,
SPCH has a MAPK-substrate domain, and its phosphorylation leads to SPCH protein

destabilization [19]. Interestingly, we found that mutants that decouple SPCH from MAPK-mediated

negative regulation are insensitive to Stomidazolone: they are complete loss of upstream receptor

kinases (*er erl1 erl2*)[14] and constitutively-active *scrm-D* that is unable to recruit MAPK to SPCH,

both causing exaggerated stomatal differentiation [12,21] (Figs 3i, j, 4b). It has been shown that, in

a rare instance, *er erl1 erl2* mutations can partially rescue the arrested meristemoid phenotype of

mute [44]. As such, the Stomidazolone-insensitivity phenotype of *er erl1 erl2* and *scrm-D* can be

explained by the overly-active and stable SPCH-SCRM heterodimers compensating for MUTE-

SCRM. This accords with the recent view hinting a hidden functional overlap between SPCH and

MUTE [45].

Our discovery that Stomidazolone targets functionally-dispensable residues within the

MUTE ACTL domain brings several implications. First, Stomidazolone (and its derivatives) could

be optimized in the future to map the functional significance of ACTL domains among the bHLH

TFs. The ACTL domain-containing bHLH proteins are prominent within the land plant lineages [24].

Whereas studies indicated the roles of the ACTL domains in dimerization [25], their functional

significance or mode of action remains largely unclear. In the well-studied case of Maize R, its

ACTL domain functions as auto-inhibitory homodimerization module that prevent the DNA binding

[46,47], a mechanism different from the stomatal bHLH-ACTL proteins [23]. Second, the finding hints

a hitherto unidentified small ligand-binding property of the ACTL-domain-containing bHLH

proteins. The prototypical ACT domain are found in biosynthetic enzymes and functions as a

sensor domain for small metabolic ligands, including amino acids and thiamine [22]. On the contrary,

it is unknown whether bHLH-ACTL TFs bind any endogenous small molecules. If they do, how

Stomidazolone interferes with such endogenous ligands would be interesting future direction.

Third, as we successfully achieved for MUTE (Fig. 6, Fig. S4), other bHLH-ACTL TFs could be

engineered to generate drug-resistant versions to selectively modulate plant phenotypes without

compromising their biological functions.

Harnessing synthetic chemistry to disrupt TF dimerization - while it seems like an obvious

means of manipulating growth and development, this approach is surprisingly unexplored in

plants. Studies identified small proteins that can competitively interfere with TF dimers and

prevent their DNA binding [48]. For instance, LONG HYPOCOTYL IN FAR-RED1 (HFR1) protein

titrates the bHLH TF, PHYTOCHROME INTERACTING FACTORs (PIFs), from forming functional

dimers [49]. In mammalian system, natural compound library screens have identified small

molecules that disrupt the homodimerization of the Hes1, an bHLH TF regulating neural cell

proliferation and differentiation [50,51]. In all these cases, unlike Stomidazolone, the inhibitors target

the core dimerization HLH modules that are essential for the TF function, thereby leaving little
room for designing and engineering the drug resistant versions of TFs.

In the synthetic chemistry point of view, the serendipitous discoveries of Stomidazolone
and subsequent chemical synthesis proffered a new catalytic transformation of oxazoles [37,52]. Our
newly discovered reaction can convert diphenyloxazole into imidazolone to create a highly
angular compounds (see Document S1), which are unexplored in a pharmacology field. Now that
we demonstrated this high degree of three-dimensionality of Stomidazolone interacts with a
discrete structural module of the ACTL domain and selectively inhibits the key step of stomatal
differentiation, further elaboration of our new amination reaction may expand the chemical space
for manipulating protein functions in plants and beyond.

359
360

Materials and Methods

Materials availability

Materials generated and used in this study will be available from Keiko Torii (ktorii@utexas.edu).
The synthesized imidazolones (Stomidazolone; ID-AYSJ929) may be limited in quantity. Inquiries
for availability could be directed to Kei Murakami (kei.murakami@kwansei.ac.jp).

Plant growth condition and chemical treatment

Arabidopsis accession Columbia (Col-0) was used as wild-type. The following mutants/transgenic
lines are reported elsewhere: *TMMpro::GUS-GFP* [13]; *spch-3/+, mute/+,* and *fama-1/+* [11]; *mute-*
2/+ [44]; *MUTEpro::nucYFP* [40]; *ice1-2/scrm* (SALK_003155), *scrm2-1* (SAIL_808_B10) [12]; *er-105*
erl1-2 erl2-1 [14]; *hdg2-2* (SALK_127828C) and *hdg2-2 atml1-3* (SALK_128172)[42], *scrm-D_S343*
and *scrm-D_S423* [23]. Their genotype and transcript reduction were confirmed. The presence of
transgenes/mutant alleles were confirmed by genotyping. Screening of small molecules that
reduce stomatal numbers was performed as previously described [35], in which all molecules were
dissolved in dimethyl sulfoxide (DMSO, Molecular biology grade, Fujifilm Wako, Japan 046-
21981), Col-0 seeds were sown in 96-well plates (TL5003; True Line) containing 95 µL of 1/2 MS
medium with rotary shaking at 140 rpm under continuous light at 22°C. Five µL of compounds
were dissolved in DMSO at 10 mM, and diluted with liquid 1/2 MS media to 1 mM and added on
1-day-old seedlings (final concentration, 50 µM). Seedlings were incubated for 8 days, and at day
9 seedling abaxial cotyledons were observed through confocal microscopy.

Molecular cloning and generation of transgenic plants

The following constructs for transgenic plants were generated in this study: *MUTEpro::MUTE-GFP* (pAH11), *MUTEpro::MUTE$_{E162AE163A}$-GFP* (pHS133), *MUTEpro::MUTE$_{R133AR134A}$-GFP* (pHS134). The original *MUTEpro::MUTE-GFP* construct (pLJP155) was generated via a different cloning strategy with different antibiotics selection [11]. We therefore re-generate the wild-type Three-way Gateway version of *MUTEpro::MUTE-GFP* (pAH11) as a control. *MUTE* genomic coding construct without stop codon (pAR205) and *MUTE* promoter construct (pAR202) were subjected to the Gateway reaction with R4pGWB504 [53]. The wild-type MUTE-GFP lines with the identical R4pGWB504 backbone enable a proper comparison of phenotypic complementation. The mutant versions of *MUTE* constructs were generated in the same way after the site-directed mutagenesis using Q5 Site-directed mutagenesis kit (NEB) and subjected to the Three-Way Gateway reaction after sequence confirmation. The constructs were then transformed into the *Agrobacterium* strain GV3101 and floral dipped into *mute-2/+* plants. For detailed lists of oligo DNA and plasmid information, see Tables S5 and S6, respectively.

Confocal microscopy

For confocal microscopy, cell peripheries of seedlings were visualized with propidium iodide (Sigma, P4170). Images were acquired using LSM800 or LSM700 (Zeiss) using a 20x lens. The GFP, YFP reporter and PI signals were detected with excitation at 488 nm and an emission filter of 410 to 546 nm, at 488 nm and an emission filter of 410 to 546 nm, and with excitation at 561 nm and 582-617 nm emission range, respectively. Raw data were collected with 1024 x 1024 pixel images. For qualitative image presentation, Adobe Photoshop CS6 was used to trim and uniformly adjust the contrast/brightness.

Quantitative analysis of epidermal phenotype

For quantitative analysis of the abaxial leaf epidermis, confocal images were taken on the days after germination as indicated in the Figure legends. Preparation of images was done as described previously [32,54]. For counting epidermal cell types, stomatal density, stomatal precursor cells (meristemoids), total stomata-lineage cells (stomata and meristemoids) were calculated by counting cell types in an area of 0.4 mm^2 (0.63 mm x 0.63 mm) and plotted as per mm^2. All raw data are presented in Table S7.

Yeast two hybrid assay

Y2H assays were performed using the Matchmaker[TM] Two-Hybrid System (Clontech). Bait

(pGBKT7) and prey (pGADT7) constructs were co-transformed into the yeast strain AH109 using

Frozen-EZ Yeast Transformation II Kit (Zymo Research, T2001). The resulting transformants

were spotted on SD (−Leu, −Trp) and SD (−Trp, −Leu, −His) selection media containing different

concentrations of 3-amino-1,2,4-triazole (Sigma, A8056) as previously reported [21]. Quantitative

α-galactosidase assay was also performed according to manufactural instruction (Clontech).

Briefly, the transformants were incubated in 2 ml SD medium with 0, 10, 50 100 µM Stomidazolone

or SIM3* with rotary shake (220 rpm) at $30^{\circ}C$ overnight. The cultured media containing yeast were

transferred to microtubes and centrifuged to obtain the supernatant. The colorimetric assay was

performed by incubating 8 µL of supernatant with 24 µL Assay Buffer that contains 0.5 M sodium

acetate (pH 4.5) (Nakalai tesque, 31137-25): 100 mM *p*-nitrophenyl α-D-Galactopyranoside

Solution (Sigma N0877) = 2:1 (v/v) at $30^{\circ}C$ for 60 min, then terminated by adding 960 µL 1X Stop

Solution (0.1 M sodium carbonate, Fujifilm Wako 195-01582). After OD_{410} was recorded, α-

galactosidase units were calculated according to the protocol (Clontech, P3024-1).

Chemical synthesis

2,4-diphenyloxazole   Chloramine B   **Stomidazolone**
                    2.0 equiv   **(AYSJ929)**

Synthesis of Stomidazolone (AYSJ929): 4 Å Molecular sieve powder (500 mg) was added into a

large screw-capped tube and heat-dried in vacuo for 1 h. 2,4-Diphenyloxazole (111 mg, 0.50

mmol), Chloramine B dihydrate (250 mg, 1.0 mmol, 2.0 equiv) and CuI (9.5 mg, 50 µmol, 10

mol%) were added into the tube with a magnetic stirring bar in open air. The tube was evacuated

and refilled with N_2 gas following the usual Schlenk technique. Anhydrous 1,2-dichloroethane (5.0

mL, 0.10 M) was added to the tube via the rubber top, the tube capped and its cap wrapped with

parafilm. The reaction mixture was stirred and heated at 40 ºC for 12 h. The mixture was then

cooled to room temperature, and the crude mixture was filtered through a pad of silica gel topped

with Na_2SO_4 in a short column and concentrated *in vacuo*. Purification by flash chromatography

on silica gel (hexane/ethyl acetate = 4:1) followed by recrystallization from CH_2Cl_2/methanol

provided Stomidazolone (AYSJ929) in 42% yield (113 mg, 0.21 mmol) as a colorless crystal. For

details of chemical synthesis see Document S1.

NMR and mass-spectrometry

The high-resolution mass spectra were recorded on Thermo Fisher Scientific Exactive Plus (ESI)

with Thermo Scientific Dionex UltiMate 3000 Series UHPLC or Agilent Technologies 6130

Quadrupole Liquid Chromatography/Mass Spectrometry (LC-MS). Nuclear magnetic resonance

(NMR) spectra were recorded on JEOL JNM-ECA-600 (^{1}H 600 MHz, ^{13}C 150 MHz) spectrometer.

Chemical shifts for ^{1}H NMR are expressed in parts per million (ppm) relative to residual CHCl$_3$ in

CDCl$_3$ (δ 7.26 ppm), and for ^{13}C NMR in ppm relative to CDCl$_3$ (δ 77.2 ppm). Data are reported

as follows: chemical shift, multiplicity (s = singlet, d = doublet, dd = doublet of doublets, t = triplets,

dt = doublet of triplets, td = triplet of doublets, tt = triplet of triplets, q = quartets, m = multiplets,

brs = broad singlet), coupling constant (Hz), and integration. For the original NMR and mass-

spectrometry data, see Dataset S1.

X-ray crystallography

A suitable crystal was mounted with mineral oil on a MiTeGen MicroMeshes and transferred to

the goniometer of the kappa goniometer of a RIGAKU XtaLAB Synergy-S system with 1.2 kW

MicroMax-007HF microfocus rotating anode (Graphite-monochromated Mo Kα radiation (λ =

0.71073 Å)) and HyPix6000HE hybrid photon-counting detector. Cell parameters were

determined and refined, and raw frame data were integrated using CrysAlis[Pro] (Agilent

Technologies, 2010). The structures were solved by direct methods with SHELXT [55] and refined

by full-matrix least-squares techniques against F^2 (SHELXL-2018/3) by using Olex2 software

package [56]. The intensities were corrected for Lorentz and polarization effects. The non-hydrogen

atoms were refined anisotropically. Hydrogen atoms were placed using AFIX instructions. CCDC

2089650 contains the supplementary crystallographic data for this paper. These data can be

obtained free of charge from The Cambridge Crystallographic Data Centre via

www.ccdc.cam.ac.uk/data_request/cif. See Table S1 for X-ray crystallography data.

Recombinant protein expression and purification

A. thaliana SCRM1 (ICE1) (1-494), $_{\Delta N}$SPCH (98-364), MUTE (1-202), $_{\Delta N}$FAMA (190-414),

MUTE$_{\Delta C}$ (1-114), SCRM-ACT (405-494), MUTE-ACT (114-202), SPCH-ACT (285-364), FAMA-

ACT (306-414), MUTE$_{R133A\_R134A}$, MUTE$_{E162A\_E163A}$, MUTE$_{R133A\_R134A\_E162A\_E163A}$ were cloned into

pGEX-4T-1 vector with an N-terminal GST tag and a thrombin cleavage sequence. For protein

expression, the constructs were transformed into *E. coli* strain BL21. For each transformant, a

single colony was selected and incubated in 10 ml LB liquid medium. The overnight-incubated *E.*

coli suspensions were transferred to 1L LB medium and incubated at 37 °C for around 2 h until

the OD_{600} reached 0.4–0.6. The isopropyl β-D-1-thiogalactopyranoside (0.25 µM of final

concentration) was added to the cultures and the strains were incubated at 22 °C for a further

18 h. GST-fused proteins were purified using glutathione agarose resin. The soluble portion of the

cell lysate was loaded onto a GST-Sepharose column. Nonspecifically bound proteins were

removed by washing the column with 20 mM Tris pH 8.0, 200 mM NaCl. The bound GST-fused

protein was eluted with 10 mm glutathione and 20 mM Tris pH 8.0, 200 mM NaCl. The GST-fused

proteins were exchanged with phosphate-buffered saline buffer, and then the solution was treated

with 50 µg of thrombin for 10–12 h at 16 °C. The GST portion of the protein was cleaved during

thrombin digestion, and then the whole solution was reloaded onto the glutathione *S*-transferase

column to obtain pure protein. The purified proteins were further purified by gel filtration on a

Superdex-200 (GE) column using fast protein liquid chromatography and phosphate buffer (pH

7.2) as the eluent. The purity of the protein was checked by SDS-PAGE. For detailed lists of oligo

DNA and plasmid information, see Tables S5 and S6, respectively.

Isothermal titration calorimetry (ITC)

Binding of the small molecule Stomidazolone to SCRM, MUTE, SPCH, FAMA, MUTEΔ_C,

MUTE_ACTL, SCRM_ACTL, SPCH_ACTL and FAMA_ACTL, MUTE$_{R133A\_134A}$, MUTE$_{E162A\_163A}$,

MUTE$_{R133A\_R134A\_E162A\_E163A}$, as well as SCRM binding to MUTE in the presence and absence of

Stomidazolone was characterized at 25 °C using a Malvern PEAQ-ITC microcalorimeter. All the

protein samples were dialyzed overnight using PBS buffer containing 2% DMSO. Stomidazolone

was dissolved in PBS buffer containing 2% DMSO. Titrations were performed by injecting 1 × 0.5-

µL and 12 × 3-µL aliquots of 1 mM Stomidazolone to 30 mM of protein in PBS buffer pH 7.4,

containing 2% DMSO. All titrations were carried out at least twice. The raw data were corrected

using buffer and protein controls and analyzed using the software supplied by the manufacturer.

See Table S3 for details.

Biolayer interferometry (BLI)

The binding affinities of the Stomidazolone with GST-tagged SCRM, MUTE, SPCH, FAMA, MUTE

ΔACTL, MUTE_ACTL, SCRM_ACTL, SPCH_ACTL and FAMA_ACTL, MUTE$_{R133A\_134A}$,

MUTE$_{E162A\_E163A}$, MUTE$_{R133A\_R134A\_E162A\_E163A}$ were measured using the Octet Red96 system

(ForteBio, Pall Life Sciences) following the manufacturer's protocols. The optical probes coated

with anti-GST were first loaded with 500 nM GST tagged proteins before kinetic binding analyses.

The experiment was performed in 96-well plates maintained at 30 °C. Each well was loaded with

200 µL reaction volume for the experiment. The binding buffer used in these experiments
contained 1× PBS supplemented with 2% DMSO, 0.02% Tween20. The concentrations of the
Stomidazolone as the analyte in the binding buffer were 100, 50, 25, 12.5, 6.25, 3.12, and 1.56 µM.
There was no binding of the analytes to the unloaded probes as shown by the control wells.
Binding kinetics to all seven concentrations of the analytes were measured simultaneously using
default parameters on the instrument. The data were analyzed using the Octet data analysis
software. The association and dissociation curves were fit with the 1:1 homogeneous ligand
model. The k_{obs} (observed rate constant) values were used to calculate K_d, with steady-state
analysis of the direct binding. See Table S3 for details.

Protein structural modelling
MUTE and MUTE ACTL domain homology models were used from Alpha fold program
(https://alphafold.ebi.ac.uk/entry/Q9M8K6) The MUTE ACTL domain was subjected to
constrained energy minimization to allow the global energy minimization and structural analysis
using the AMBER 12 suite and VADAR [57,58]. Molecular docking of both racemic forms of
Stomidzolone to the MUTE, MUTE ACTL domain monomers was carried out using the high
ambiguity-driven biomolecular docking (HADDOCK) approach [59,60]. Ambiguous interaction
restraints (AIRs) were selected based on mutagenesis data and preliminary AUTODOCK model
[61]. The pairwise ligand interface RMSD matrix over all structures was calculated, and the final
structures were clustered using an RMSD cut-off value of 3.5 Å for both ligand and protein. The
clusters were then prioritized using RMSD and the HADDOCK score (weighted sum of a
combination of energy terms). See Table S4 for binding energies for Stomidazolone-MUTE
docking simulation.

Measurement of stomatal opening in *Commelina benghalensis*
The experiments were performed as described previously [30,62]. Briefly, Stomidazolone or ABA
were added to a multi-well plate with basal buffer [5 mM MES/bistrispropane (pH 6.5), 50 mM KCl,
and 0.1 mM CaCl₂] to be diluted to a final concentration of 50 or 20 µM, respectively. *C.*
benghalensis plants were incubated in the dark overnight; then, 4 mm diameter leaf discs were
excised from fully expanded leaves using a hole punch (Biopsy Punch, Kai Medical). Samples
were immersed in a basal buffer containing compounds for 30 min before being exposed to light
(150 µmol m^{-2} s^{-1} red light and 50 µmol m^{-2} s^{-1} blue light) for 3 h. The stomatal apertures were
observed using a stereoscopic microscope (BX43; Olympus). The microscopic images were
acquired with a charge-coupled device (CCD) camera (DP27; Olympus), followed by autofocus

image processing using cellSens standard software (Olympus). Measurement of stomatal

apertures of *C. benghalensis* was performed by using an automation program, *DeepStomata* [63].

Quantification and statistical analysis

A series of Z-stack confocal images was obtained to obtain images covering the epidermis and

capturing GFP/YFP signal from the reporter lines. The area and the number of epidermal cells

were quantified by using FIJI-ImageJ. Statistical analyses were performed using R ver. 4.0.2.

Sample sizes (n) are indicated and defined in each legend. For the multiple sample comparison,

1-way ANOVA was performed followed by a post hoc Tukey HSD test. For the two-sample

comparison, Student's T-test was performed. Graphs were generated using R ggplot2 package.

For all boxplots, each box represents IQR (interquartile range), in which top, middle line, and

bottom indicating 75, 50, and 25 percentiles, respectively, and the bar representing maximum to

minimal values. Each dot represents jittered individual data point. The datapoints that go beyond

the bar are outliers. The color of Boxplot graph was based on ColorBrewer.org. The value of n,

the number of each experiment or samples, the means of error bars, and how statistical

significance was defined are indicated in a relevant figure legend.

ACKNOWLEDGEMENTS

We thank Amanda Rychel and Alex Hofstetter for constructing the 3-Way Gateway version of

MUTEpro::MUTE-GFP, Zixuan Li for assisting with chemical synthesis, Nagoya ITbM Molecular

Structure Center for assistance in NMR and mass-spectrometry, Jason McLellan for the access

to ITC, and Kosuke Watanabe for assistance with X-ray crystal structure analysis. We especially

thank Yuichiro Tsuchiya for the insightful discussion and generously offering the lab space/facility

for A.N., and Eundeok Kim for comments. A part of this work was conducted in Institute for

Molecular Science supported by ARIM (JPMXP1223MS5012). This work was supported by

MEXT/JSPS KAKENHI (16H01237, 17H06476 and 19H00990) to K.U.T.; Japan Science and

Technology Agency (JST) PRESTO (JPMJPR20D8), TOBE MAKI Scholarship Foundation, and

Tatematsu Foundation to K.M.; and ITbM Research Award to S.J.A.Y., A.N., and S.-K.H. A.N.

was in part supported by the JST CREST Award (JPMJCR1924). S.J.A.Y. was supported by the

JSPS Research Fellowship for Young Scientists DC1. S.-K.H. was supported by the Young

Leader Cultivation Program from the Institute of Advanced Research, Nagoya University. K.U.T.

acknowledges the support from the Howard Hughes Medical Institute and Johnson & Johnson

Centennial Chair of Plant Cell Biology at UT Austin.

CONTRIBUTIONS

A.N., K.M., S.J.A.Y, S.-K.H and K.U.T. conceived the study. K.U.T., K.M., S.J.A.Y, S.K-H., A.N., and K.M.S. designed the experiments. A.N., K.M.S., S.J.A.Y, H.S., C.M.C., T.S., H.K., R.I, S.K., Y.A., K.U.T. performed the experiments (forward chemical genetics/transgenic/phenotypic analysis A.N., S.J.A.Y, H.S., S.-K.H., C.M.C., H.K., K.U.T.; yeast/biochemical/biophysical experiments, K.M.S., R.I.; chemical synthesis, K.M.; structural analyses/modeling, K.M., K.S.M., Y.S.; stomatal movement assays, Y.A., T.K.). A.N., K.S.M., K.M., Y.A. and K.U.T. performed data analysis and visualization. K.U.T. wrote the manuscript draft. All authors contributed to the editing of the manuscript. K.M. and K.U.T. are responsible for project administration and funding acquisition.

COMPETING INTERESTS

The authors declare no competing interests.

600
601
Figure 1. Stomidazolone is a potent chemical for meristemoid arrest
(a) Schematics of chemical synthesis. Supplementary Document S1 for the details of the synthesis. Bottom right, structure of Stomidazolone (ID: AYSJ929).
(b) X-ray crystal structure of Stomidazolone shown as a stick model (White, H; Beige, C; Blue, N; Yellow, S; Red, O).

(c) Confocal microscopy images of representative cotyledon abaxial epidermis from Arabidopsis
wild-type (WT) seedlings grown in the presence of 0 (mock), 20, 50, and 100 μM Stomidazolone.
Orange brackets, stomatal-lineage cells with an arrested meristemoid in the center. Scale bars,
50 μm.
(d) Arrested meristemoids by Stomidazolone express proper meristemoid markers. Shown are
confocal microscope images of representative abaxial cotyledons epidermis expressing
TMMpro::GUS-GFP (top) and *MUTEpro::nucYFP* (bottom) grown in the presence of 0 (mock:
Left) and 50 μM Stomidazolone (Right). Scale bars, 50 μm
(e) Stomidazolone treatment reduces the numbers of stomata but increases that of meristemoids.
Quantitative analysis of stomatal (s; green), meristemoid (m; cyan), and stomata+meristemoid (t;
gray) density per mm^2 of cotyledon abaxial epidermis from seedlings grown in the presence of 0
(mock), 20, 50, and 100 μM Stomidazolone. One-way ANOVA followed by Tukey's HSD analysis
were performed for each cell type (stomata and meristemoids). Letters (a,b,c) indicate groups
that are statistically different from other groups within each cell type. n = 8
(f) Stomidazolone does not impact seedling growth. Shown are two representative 9-day-old WT
seedlings, grown in the presence of 0 (mock), 20, 50, and 100 μM Stomidazolone. Scale bars, 10
mm.
624

625
626
Figure 2. *spch* and *mute*, but not *fama*, are epistatic to Stomidazolone
(a) Schematic diagram showing the genetic control of stomatal cell-state transitions. Key genes/gene products and their roles are indicated in the cartoon. See main text for details of each step. Arrows, positive regulation; T-bars, negative regulation. Upregulation of *MUTE* by *HDG2* and inhibition of *MUTE* by *ERL1* are highlighted in red. MMC, meristemoid mother cell; GMC, guard mother cell; GC, guard cell.

(b) Representative cotyledon abaxial epidermis of mock and 50 μM Stomidazolone-treated WT,

spch (spch-3), *mute (mute-1)*, and *fama*. Scale bars, 50 μm.

(c, d) Quantitative analysis of meristemoids density per 1 mm^2 (c) and *fama* tumors (d) of

cotyledon abaxial epidermis from seedlings grown in the presence of 0 (mock) and 50 μM

Stomidazolone. Student's T test was performed for each plant. p values and NS (not significant)

are indicated above each boxplot. n = 8.

Figure 3. Stomidazolone is a potent inhibitor of cell-state transition at the MUTE-step

(a-j) Effects of Stomidazolone on the epidermis of wild-type (WT: a, b), *er* (c, d), *er erl1* (e, f), *er erl2* (g, h), and *er erl1 erl2* (i, j). Representative cotyledon abaxial epidermis of mock- (left) and 50 μM Stomidazolone (right). Orange brackets, clusters of arrested meristemoids. Scale bars, 50 μm. (b, d, f, h, j) Quantitative analysis of stomatal (s; green), meristemoid (m; cyan), and stomata+meristemoid (t; gray) density per mm^2 of cotyledon abaxial epidermis from seedlings grown in the presence of 0 (mock) and 50 μM Stomidazolone. One-way ANOVA followed by Tukey's HSD analysis were performed for each cell type (stomata and meristemoids). Letters (a,b,c) indicate groups that are statistically different from other groups within each cell type. n = 8.

(k) Number of meristemods, data replotted from other panels.

(l-o) Effects of Stomidazolone on the epidermis of *hdg2* (*hdg2-2*: l, m) and *hdg2 atml1* (*hdg2-2 atml1-3*: n, o). All experiments and analyses were performed as described above. Scale bars, 50 μm. n = 8.

657
658
Figure 4. SCRM activity dictates *in vivo* potency and selectivity of Stomidazolone

(a) Abaxial epidermis of cotyledons in *scrm* (*ice1-2*) mutants treated with mock or 50 µM Stomidazolone on 1-day-old seedlings and further incubated for 8 days. *scrm* loss-of-function mutants occasionally exhibit parallelly paired-stomata, so called 'four-lips' phenotype (orange plus) as well as stoma made with extra guard cell (orange asterisk) likely due to FAMA mis-regulation. Scale bars, 50 µm.

(b) Stomatal index (SI, %) for Mock- or Stomidazolone-treated *scrm* mutant. n=8.

(c) Abaxial epidermis of cotyledons in *scrm-D* mutants treated with mock or 50 µM Stomidazolone on 1-day-old seedlings and further incubated for 8 days. Scale bars, 50 µm.

(d) Stomatal index (SI, %) for Mock- or Stomidazolone-treated *scrm-D* mutant. Student's T test was performed, and p values are shown above each boxplot. n=8.

(e-h) Effect of Stomidazolone on the epidermis of *scrm-D_S343* and *scrm_D_S423*. (e, g) Abaxial epidermis of cotyledons in *scrm-D_S343 and scrm_D_S423* mutants treated with mock or 50 µM Stomidazolone for 8 days. Clustered meristemoids and their sister stomatal-lineage ground cells (SLGCs: orange brackets) become exaggerated. Scale bars, 50 µm. (f, h) The cumulative bar graphs indicate the densities of stomata (green), small cells (meristemoids and SLGCs, cyan) and pavement cells (PC; dark brown) per 0.41 mm^2 with mock or 50 µM Stomidazolone.

Figure 5. Stomidazolone binds to the ACTL domain and disrupts bHLH heterodimers

(a) Y2H analysis. Yeasts expressing control and pairs of stomatal bHLH proteins were spotted in 10-fold serial dilutions on appropriate selection media with or without 50 μM Stomidazolone. 1 mM 3-AT treatment shows that Stomidazolone reduces the interactions of SCRM with stomatal bHLH proteins. 3-AT, 3-amino-1,2,4-triazole.

(b) Quantitative α-galactosidase assays. Yeasts expressing control and pairs of stomatal bHLH proteins were cultured in 0, 10, 50, and 100 μM Stomidazolone and enzyme assays were performed. One-way ANOVA followed by Tukey's HSD analysis were performed. Letters (a,b,c) indicate that the enzyme activities are statistically different within the group. Experiments were repeated at least 3 times.

(c, d) Quantitative Analysis of SCRM-MUTE Interactions in the absence (c), and presence (d) of
Stomidazolone. *In vitro* binding response curves are provided for SCRM with MUTE and
with/without Stomidazolone. The data represent the mean ± SD and are representative of two
independent experiments.
(e) Quantitative analysis of Stomidazolone binding with SCRM (blue), MUTE (cyan), FAMA
(green), and SPCH (lavender). The MUTE protein exhibit higher affinity to Stomidazolone. *In vitro*
binding response curves are provided Stomidazolone with SCRM/MUTE/SPCH/FAMA/MUTE.
The data represent the mean ± SD and are representative of two independent experiments.
(f) Quantitative analysis of Stomidazolone binding with MUTE bHLH (green) and ACTL (brown)
domains. *In vitro* binding response curves are provided for Stomidazolone with $MUTE_{bHLH}$ and
$MUTE_{ACTL}$. The data represent the mean ± SD and are representative of two independent
experiments.
700
701

Figure 6. Engineering Stomidazolone-resistant plants based on the MUTE-Stomidazolone docking model

(a, b) MUTE-Stomidazolone binding models. (a) Alphafold2 predicted full-length MUTE protein is shown as ribbon representation. MUTE bHLH domain, green; ACTL domain, peach; Linker region, red. Stomidazolone is shown as stick model (gray). (b) Magnified view of MUTE-Stomidazolone binding interface. The key binding residues (R133, R134, E162 and E163) sidechains are highlighted in magenta.

(c) Quantitative analysis of Stomidazolone binding to wild-type MUTE and MUTE site-directed mutants ($MUTE_{R133A\_R134A}$, $MUTE_{E162A\_E163A}$, $MUTE_{R133A\_R134AE162A\_E163A}$) by BLI. *In vitro* binding response curves are presented for Stomidazolone with MUTE and its mutants. Stomidazolone compound was tested at seven different concentrations (200, 100, 50, 25, 12.5, 6.25, and 3.125 μM). The data represent the mean ± SD and are representative of two independent experiments.

(d) Site-directed mutant versions of MUTE that disrupt Stomidazolone binding are expressed normally and biologically functional. Representative cotyledon abaxial epidermis of 5-day-old *mute-2* seedlings expressing *MUTEpro::MUTE-GFP* (left), *MUTEpro::MUTE$_{E162A\_E163A}$ -GFP* (center), and *MUTEpro::MUTE$_{R133A\_R134A}$-GFP* (right). Insets, representative late meristemoids from each panel with MUTE-GFP signals in nucleus. Confocal images are inverted for clear cellular outlines. Scale bars, 20 μm.

(e-f) MUTE mutant versions that disrupt Stomidazolone binding confer drug resistance *in vivo*. (e) Representative cotyledon abaxial epidermis of five-day-old *mute-2* seedlings expressing *MUTEpro::MUTE-GFP* (left), *MUTEpro::MUTE$_{E162A\_E163A}$ -GFP* (center), and

MUTEpro::MUTE$_{R133A\_R134A}$ -GFP (right) with mock- (top) and 20 μM Stomidazolone for eight days

(bottom). Scale bars, 50 μm.

(f) Quantitative analysis of total (t: stomata and meristemoids), stomatal (s: green) or meristemoid

(m: light cyan) density per mm^2 of cotyledon abaxial epidermis from seedlings grown in the

presence of 0 (mock) and 20 μM Stomidazolone. Student's T test was performed were performed

for each cell type (total, stomata and meristemoids) and p values are indicated. *, p<0.05, ***,

p<0.0005. Experiments were repeated twice. n = 8.

REFERENCES

Hetherington, A. M. & Woodward, F. I. The role of stomata in sensing and driving environmental change. *Nature* **424**, 901-908 (2003).

Hsu, P. K., Dubeaux, G., Takahashi, Y. & Schroeder, J. I. Signaling mechanisms in abscisic acid-mediated stomatal closure. *Plant J* **105**, 307-321 (2021).

Kim, T. H., Bohmer, M., Hu, H., Nishimura, N. & Schroeder, J. I. Guard cell signal transduction network: advances in understanding abscisic acid, CO2, and Ca2+ signaling. *Annu Rev Plant Biol* **61**, 561-591 (2010).

Qi, X. & Torii, K. U. Hormonal and environmental signals guiding stomatal development. *BMC Biol* **16**, 21 (2018).

Bertolino, L. T., Caine, R. S. & Gray, J. E. Impact of Stomatal Density and Morphology on Water-Use Efficiency in a Changing World. *Front Plant Sci* **10**, 225 (2019).

Han, S. K. & Torii, K. U. Lineage-specific stem cells, signals and asymmetries during stomatal development. *Development* **143**, 1259-1270 (2016).

Lau, O. S. & Bergmann, D. C. Stomatal development: a plant's perspective on cell polarity, cell fate transitions and intercellular communication. *Development* **139**, 3683-3692 (2012).

Nadeau, J. A. & Sack, F. D. Stomatal development in Arabidopsis. *Arabidopsis Book* **1**, e0066 (2002).

MacAlister, C. A., Ohashi-Ito, K. & Bergmann, D. C. Transcription factor control of asymmetric cell divisions that establish the stomatal lineage. *Nature* **445**, 537-540 (2007).

Ohashi-Ito, K. & Bergmann, D. C. Arabidopsis FAMA controls the final proliferation/differentiation switch during stomatal development. *Plant Cell* **18**, 2493-2505 (2006).

Pillitteri, L. J., Sloan, D. B., Bogenschutz, N. L. & Torii, K. U. Termination of asymmetric cell division and differentiation of stomata. *Nature* **445**, 501-505 (2007).

Kanaoka, M. M., Pillitteri, L. J., Fujii, H., Yoshida, Y., Bogenschutz, N. L., Takabayashi, J., Zhu, J. K. & Torii, K. U. SCREAM/ICE1 and SCREAM2 specify three cell-state transitional steps leading to arabidopsis stomatal differentiation. *Plant Cell* **20**, 1775-1785 (2008).

Nadeau, J. A. & Sack, F. D. Control of stomatal distribution on the Arabidopsis leaf surface. *Science* **296**, 1697-1700 (2002).

Shpak, E. D., McAbee, J. M., Pillitteri, L. J. & Torii, K. U. Stomatal patterning and differentiation by synergistic interactions of receptor kinases. *Science* **309**, 290-293 (2005).

Hara, K., Kajita, R., Torii, K. U., Bergmann, D. C. & Kakimoto, T. The secretory peptide gene EPF1 enforces the stomatal one-cell-spacing rule. *Genes Dev* **21**, 1720-1725 (2007).

Hara, K., Yokoo, T., Kajita, R., Onishi, T., Yahata, S., Peterson, K. M., Torii, K. U. & Kakimoto, T. Epidermal cell density is autoregulated via a secretory peptide, EPIDERMAL PATTERNING FACTOR 2 in Arabidopsis leaves. *Plant Cell Physiol* **50**, 1019-1031 (2009).

Hunt, L. & Gray, J. E. The signaling peptide EPF2 controls asymmetric cell divisions during stomatal development. *Curr Biol* **19**, 864-869 (2009).

Lee, J. S., Kuroha, T., Hnilova, M., Khatayevich, D., Kanaoka, M. M., McAbee, J. M., Sarikaya, M., Tamerler, C. & Torii, K. U. Direct interaction of ligand-receptor pairs specifying stomatal patterning. *Genes Dev* **26**, 126-136 (2012).

Lampard, G. R., Macalister, C. A. & Bergmann, D. C. Arabidopsis stomatal initiation is controlled by MAPK-mediated regulation of the bHLH SPEECHLESS. *Science* **322**, 1113-1116 (2008).

Lampard, G. R., Lukowitz, W., Ellis, B. E. & Bergmann, D. C. Novel and expanded roles for MAPK signaling in Arabidopsis stomatal cell fate revealed by cell type-specific manipulations. *Plant Cell* **21**, 3506-3517 (2009).

Putarjunan, A., Ruble, J., Srivastava, A., Zhao, C., Rychel, A. L., Hofstetter, A. K., Tang, X., Zhu, J. K., Tama, F., Zheng, N. & Torii, K. U. Bipartite anchoring of SCREAM enforces

stomatal initiation by coupling MAP kinases to SPEECHLESS. *Nat Plants* **5**, 742-754
(2019).
22   Grant, G. A. The ACT domain: a small molecule binding domain and its role as a common
regulatory element. *J Biol Chem* **281**, 33825-33829 (2006).
23   Seo, H., Sepuru, K. M., Putarjunan, A., Aguirre, L., Burrows, B. A. & Torii, K. U. Intragenic
suppressors unravel the role of the SCREAM ACT-like domain for bHLH partner selectivity
in stomatal development. *Proc Natl Acad Sci U S A* **119** (2022).
24   Lee, Y. S., Shiu, S. H. & Grotewold, E. Evolution and diversification of the ACT-like domain
associated with plant basic helix-loop-helix transcription factors. *Proc Natl Acad Sci U S*
A **120**, e2219469120 (2023).
25   Lee, Y. S., Herrera-Tequia, A., Silwal, J., Geiger, J. H. & Grotewold, E. A hydrophobic
residue stabilizes dimers of regulatory ACT-like domains in plant basic helix-loop-helix
transcription factors. *J Biol Chem*, 100708 (2021).
26   Spring, D. R. Chemical genetics to chemical genomics: small molecules offer big insights.
Chem Soc Rev **34**, 472-482 (2005).
27   McCourt, P. & Desveaux, D. Plant chemical genetics. *New Phytol* **185**, 15-26 (2010).
28   Kinoshita, T., Toh, S. & Torii, K. U. Chemical control of stomatal function and development.
Curr Opin Plant Biol **60**, 102010 (2021).
29   Kim, T. H., Hauser, F., Ha, T., Xue, S., Bohmer, M., Nishimura, N., Munemasa, S.,
Hubbard, K., Peine, N., Lee, B. H., Lee, S., Robert, N., Parker, J. E. & Schroeder, J. I.
Chemical genetics reveals negative regulation of abscisic acid signaling by a plant
immune response pathway. *Curr Biol* **21**, 990-997 (2011).
30   Toh, S., Inoue, S., Toda, Y., Yuki, T., Suzuki, K., Hamamoto, S., Fukatsu, K., Aoki, S.,
Uchida, M., Asai, E., Uozumi, N., Sato, A. & Kinoshita, T. Identification and
Characterization of Compounds that Affect Stomatal Movements. *Plant Cell Physiol* **59**,
1568-1580 (2018).
31   De Rybel, B., Audenaert, D., Vert, G., Rozhon, W., Mayerhofer, J., Peelman, F., Coutuer,
S., Denayer, T., Jansen, L., Nguyen, L., Vanhoutte, I., Beemster, G. T., Vleminckx, K.,
Jonak, C., Chory, J., Inze, D., Russinova, E. & Beeckman, T. Chemical inhibition of a
subset of Arabidopsis thaliana GSK3-like kinases activates brassinosteroid signaling.
Chem Biol **16**, 594-604 (2009).
32   Houbaert, A., Zhang, C., Tiwari, M., Wang, K., de Marcos Serrano, A., Savatin, D. V., Urs,
M. J., Zhiponova, M. K., Gudesblat, G. E., Vanhoutte, I., Eeckhout, D., Boeren, S., Karimi,
M., Betti, C., Jacobs, T., Fenoll, C., Mena, M., de Vries, S., De Jaeger, G. & Russinova,
E. POLAR-guided signalling complex assembly and localization drive asymmetric cell
division. *Nature* **563**, 574-578 (2018).
33   Bian, C., Guo, X., Zhang, Y., Wang, L., Xu, T., DeLong, A. & Dong, J. Protein phosphatase
2A promotes stomatal development by stabilizing SPEECHLESS in Arabidopsis. *Proc Natl*
Acad Sci U S A **117**, 13127-13137 (2020).
34   Sakai, Y., Sugano, S. S., Kawase, T., Shirakawa, M., Imai, Y., Kawamoto, Y., Sugiyama,
H., Nakagawa, T., Hara-Nishimura, I. & Shimada, T. The chemical compound bubblin
induces stomatal mispatterning in Arabidopsis by disrupting the intrinsic polarity of
stomatal lineage cells. *Development* **144**, 499-506 (2017).
35   Ziadi, A., Uchida, N., Kato, H., Hisamatsu, R., Sato, A., Hagihara, S., Itami, K. & Torii, K.
U. Discovery of synthetic small molecules that enhance the number of stomata: C-H
functionalization chemistry for plant biology. *Chem Commun (Camb)* **53**, 9632-9635
(2017).
36   Ito, E., Fukushima, T., Kawakami, T., Murakami, K. & Itami, K. Catalytic Dehydrogenative
C-H Imidation of Arenes Enabled by Photo-generated Hole Donation to Sulfonimide.
Chem **2**, 383-392 (2017).

37 Hili, R. & Yudin, A. K. Making carbon-nitrogen bonds in biological and chemical synthesis.
Nat Chem Biol **2**, 284-287 (2006).
38 Toda, Y., Perry, G. J. P., Inoue, S., Ito, E., Kawakami, T., Narouz, M. R., Takahashi, K.,
Aihara, Y., Maeda, B., Kinoshita, T., Itami, K. & Murakami, K. Identification of stomatal-
regulating molecules from de novo arylamine collection through aromatic C-H amination.
Sci Rep **12**, 949 (2022).
39 Horst, R. J., Fujita, H., Lee, J. S., Rychel, A. L., Garrick, J. M., Kawaguchi, M., Peterson,
K. M. & Torii, K. U. Molecular Framework of a Regulatory Circuit Initiating Two-
Dimensional Spatial Patterning of Stomatal Lineage. *PLoS Genet* **11**, e1005374 (2015).
40 Qi, X., Han, S. K., Dang, J. H., Garrick, J. M., Ito, M., Hofstetter, A. K. & Torii, K. U.
Autocrine regulation of stomatal differentiation potential by EPF1 and ERECTA-LIKE1
ligand-receptor signaling. *Elife* **6**, e24102 (2017).
41 Qi, X., Yoshinari, A., Bai, P., Maes, M., Zeng, S. M. & Torii, K. U. The manifold actions of
signaling peptides on subcellular dynamics of a receptor specify stomatal cell fate. *Elife* **9**
(2020).
42 Peterson, K. M., Shyu, C., Burr, C. A., Horst, R. J., Kanaoka, M. M., Omae, M., Sato, Y.
& Torii, K. U. Arabidopsis homeodomain-leucine zipper IV proteins promote stomatal
development and ectopically induce stomata beyond the epidermis. *Development* **140**,
1924-1935 (2013).
43 Pillitteri, L. J., Peterson, K. M., Horst, R. J. & Torii, K. U. Molecular profiling of stomatal
meristemoids reveals new component of asymmetric cell division and commonalities
among stem cell populations in Arabidopsis. *Plant Cell* **23**, 3260-3275 (2011).
44 Pillitteri, L. J., Bogenschutz, N. L. & Torii, K. U. The bHLH protein, MUTE, controls
differentiation of stomata and the hydathode pore in Arabidopsis. *Plant Cell Physiol* **49**,
934-943 (2008).
45 Smit, M. E. & Bergmann, D. C. The stomatal fates: Understanding initiation and
enforcement of stomatal cell fate transitions. *Curr Opin Plant Biol*, 102449 (2023).
46 Kong, Q., Pattanaik, S., Feller, A., Werkman, J. R., Chai, C., Wang, Y., Grotewold, E. &
Yuan, L. Regulatory switch enforced by basic helix-loop-helix and ACT-domain mediated
dimerizations of the maize transcription factor R. *Proc Natl Acad Sci U S A* **109**, E2091-
2097 (2012).
47 Feller, A., Hernandez, J. M. & Grotewold, E. An ACT-like domain participates in the
dimerization of several plant basic-helix-loop-helix transcription factors. *J Biol Chem* **281**,
28964-28974 (2006).
48 Seo, P. J., Hong, S. Y., Kim, S. G. & Park, C. M. Competitive inhibition of transcription
factors by small interfering peptides. *Trends Plant Sci* **16**, 541-549 (2011).
49 Hornitschek, P., Lorrain, S., Zoete, V., Michielin, O. & Fankhauser, C. Inhibition of the
shade avoidance response by formation of non-DNA binding bHLH heterodimers. *EMBO*
J **28**, 3893-3902 (2009).
50 Arai, M. A., Masada, A., Ohtsuka, T., Kageyama, R. & Ishibashi, M. The first Hes1 dimer
inhibitors from natural products. *Bioorg Med Chem Lett* **19**, 5778-5781 (2009).
51 Arai, M. A., Makita, Y., Yamaguchi, Y., Kawano, H., Suganami, A., Tamura, Y. & Ishibashi,
M. Total synthesis of lindbladione, a Hes1 dimerization inhibitor and neural stem cell
activator isolated from Lindbladia tubulina. *Sci Rep* **10**, 21433 (2020).
52 Jiao, J., Murakami, K. & Itami, K. Catalytic methods for aromatic C–H amination: an ideal
strategy for nitrogen-based functional molecules. *ACS Catalysis* **6**, 610-633 (2016).
53 Nakagawa, T., Nakamura, S., Tanaka, K., Kawamukai, M., Suzuki, T., Nakamura, K.,
Kimura, T. & Ishiguro, S. Development of R4 gateway binary vectors (R4pGWB) enabling
high-throughput promoter swapping for plant research. *Biosci Biotechnol Biochem* **72**,
624-629 (2008).

54   Han, S. K., Herrmann, A., Yang, J., Iwasaki, R., Sakamoto, T., Desvoyes, B., Kimura, S.,
Gutierrez, C., Kim, E. D. & Torii, K. U. Deceleration of the cell cycle underpins a switch
from proliferative to terminal divisions in plant stomatal lineage. *Dev Cell* **57**, 569-582 e566
(2022).
55   Sheldrick, G. M. SHELXT - integrated space-group and crystal-structure determination.
Acta Crystallogr A Found Adv **71**, 3-8 (2015).
56   Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. OLEX2:
a complete structure solution, refinement and analysis program. *J Appl Cryst* **42**, 339-341
(2009).
57   Case, D. A., Darden, T. A., Cheatham III, T. E., Simmerling, C. L., Wang, J., Duke, R. E.,
Luo, R., Merz, K. M., Wang, B., Pearlman, D. A., Crowley, M., Brozell, S., Tsui, V., Gohlke,
H., Mongan, J., Hrnak, V., Cui, G., Beroza, P., Schafmeister, C., Caldwell, J. W., Ross, W.
S. & Kollman, P. A. Amber 9. (2004).
58   Willard, L., Ranjan, A., Zhang, H., Monzavi, H., Boyko, R. F., Sykes, B. D. & Wishart, D.
S. VADAR: a web server for quantitative evaluation of protein structure quality. *Nucleic*
Acids Res **31**, 3316-3319 (2003).
59   Dominguez, C., Boelens, R. & Bonvin, A. M. HADDOCK: a protein-protein docking
approach based on biochemical or biophysical information. *J Am Chem Soc* **125**, 1731-
1737 (2003).
60   de Vries, S. J., van Dijk, A. D., Krzeminski, M., van Dijk, M., Thureau, A., Hsu, V.,
Wassenaar, T. & Bonvin, A. M. HADDOCK versus HADDOCK: new features and
performance of HADDOCK2.0 on the CAPRI targets. *Proteins* **69**, 726-733 (2007).
61   Morris, G. M., Huey, R., Lindstrom, W., Sanner, M. F., Belew, R. K., Goodsell, D. S. &
Olson, A. J. AutoDock4 and AutoDockTools4: Automated docking with selective receptor
flexibility. *J Comput Chem* **30**, 2785-2791 (2009).
62   Aihara, Y., Maeda, B., Goto, K., Takahashi, K., Nomoto, M., Toh, S., Ye, W., Toda, Y.,
Uchida, M., Asai, E., Tada, Y., Itami, K., Sato, A., Murakami, K. & Kinoshita, T.
Identification and improvement of isothiocyanate-based inhibitors on stomatal opening to
act as drought tolerance-conferring agrochemicals. *Nat Commun* **14**, 2665 (2023).
63   Toda, Y., Toh, S., Bourdais, G., Robatzek, S., MacLean, D. & Kinoshita, T. DeepStomata:
Facial Recognition Technology for Automated Stomatal Aperture Measurement. *bioRxiv*
p 365098. (2018).
918