Chemical inhibition of stomatal differentiation by perturbation of the master-regulatory

bHLH heterodimer via an ACT-Like domain

Ayami Nakagawa[1]*, Krishna Mohan Sepuru[2,3]*, Shu Jan Yip[1]*, Hyemin Seo[2,3], Calvin M.

Coffin[2,3], Kota Hashimoto[4], Zixuan Li[4], Yasutomo Segawa[5], Rie Iwasaki[1], Hiroe Kato[1], Daisuke

Kurihara[1,7], Yusuke Aihara[1,6], Stephanie Kim[3], Toshinori Kinoshita[1], Kenichiro Itami[1], Soon-Ki

Han[1,7], Kei Murakami[1,4,6, +], and Keiko U. Torii[1,2,3+]

[1]Institute of Transformative Bio-Molecules (WPI-ITbM), Nagoya University, Nagoya, Aichi, 464-

8601, Japan

[2]Howard Hughes Medical Institute, The University of Texas at Austin, Austin, TX 78712 USA

[3]Department of Molecular Biosciences, The University of Texas at Austin, Austin, TX 78712

USA

[4]Department of Chemistry, Kwansei Gakuin University, Sanda, Hyogo 669-1330, Japan

[5]Institute for Molecular Science and SOKENDAI, Myodaiji, Okazaki, 444-8585, Japan

16
[6]PRESTO, Japan Science and Technology Agency (JST), Chiyoda, Tokyo 102-0076, Japan

[7]Institute for Advanced Research (IAR), Nagoya University, Nagoya, Aichi, 464-8601, Japan

*These authors contributed equally

+Corresponding authors

kei.murakami@kwansei.ac.jp

ktorii@utexas.edu

23
24
25
26

Selective perturbation of protein interactions with chemical compounds enables dissection and control of developmental processes. Differentiation of stomata, cellular valves vital for plant growth and survival, is specified by the basic-helix-loop-helix (bHLH) heterodimers. Harnessing a new amination reaction, we here report a synthesis, derivatization, target identification, and mode of action of an atypical doubly-sulfonylated imidazolone, Stomidazolone, which triggers stomatal stem cell arrest. Our forward chemical genetics followed by biophysical analyses elucidated that Stomidazolone directly binds to the C-terminal ACT-Like (ACTL) domain of MUTE, a master regulator of stomatal differentiation, and perturbs its heterodimerization with a partner bHLH, SCREAM *in vitro* and in plant cells. On the other hand, Stomidazolone analogs that are biologically inactive did not bind to MUTE or disrupt the SCREAM-MUTE heterodimers. Guided by structural docking modeling, we rationally designed MUTE with reduced Stomidazolone binding. These engineered MUTE proteins are fully functional and confer Stomidazolone resistance *in vivo*. Our study identifies doubly-sulfonylated imidazolone as a direct inhibitor of the stomatal master regulator, further expanding the chemical space for perturbing bHLH-ACTL proteins to manipulate plant development.

INTRODUCTION

Plant growth and survival rely on stomata ~ adjustable valves on the shoot epidermis of land
plants facilitating efficient gas exchange while minimizing water loss. Each stoma consists of
paired guard cells that can adjust the aperture in response to the environment, such as light,
drought, and CO_2 levels [1]. A wealth of studies has elucidated the molecular basis of signaling
mechanisms leading to stomatal opening and closure [2,3]. In addition to stomatal movement,
stomatal development is also influenced by environmental conditions and in turn impacts plants'
water-use efficiency [4,5]. During leaf development, stomata are generated through a stereotypical
sequence of cell division and differentiation events [6,7]. In dicotyledonous plants, including
Arabidopsis, the stomatal-cell lineage is initiated via *de novo* asymmetric cell divisions in the
protoderm, which create meristemoids, proliferating stomatal precursor cells with a stem-cell-
like property. After a few rounds of asymmetric divisions, the meristemoid differentiates into a
guard mother cell (GMC). The GMC then undergoes a single symmetric division to terminally
differentiate a stoma [8].

The cell-state transitional steps during stomatal development are governed by the
sequential actions of basic-helix-loop-helix (bHLH) transcription factors (TFs), SPEECHLESS
(SPCH), MUTE, and FAMA, which form heterodimers with their partner bHLHs, SCREAM
(SCRM, also known as ICE1) and SCRM2 [9-12]. The activities of these stomatal bHLH modules
are enforced by the inhibitory peptide-receptor kinase signaling pathways. Briefly, secreted
peptide ligands EPIDERMAL PATTERNING FACTORs (EPFs) are perceived by their receptors,
ERECTA-family receptor kinases and TOO MANY MOUTHS (TMM) [13-18]. This activates a
downstream Mitogen Activated Protein Kinase (MAPK) cascade, which in turn inhibits the bHLH
heterodimers via direct phosphorylation and degradation events [19-21].

Interestingly, these stomatal bHLH TFs possess a C-terminal ACT-Like (ACTL) domain,
a small structurally defined module resembling the ACT domain found in diverse metabolic
enzymes in bacteria and eukaryotes (named after <u>a</u>spartate kinase-<u>c</u>horismate mutase-<u>t</u>yrA
prephenate dehydrogenase) [21,22]. Recently, we revealed that the SCRM ACTL domain serves
as a heterodimer selectivity interface with SPCH, MUTE, and FAMA [23]. This rather surprising
finding offers a potential for the ACTL domain of bHLH TFs as a drug target for selective
inhibition of heterodimerization, hence stomatal differentiation. Whereas the ACTL domain is
predominant amongst plant bHLH proteins [24,25] no chemical inhibitors or binders targeting this
domain, or more broadly, bHLH-ACTL TFs are available.

Chemical compounds that selectively perturb functions and interactions of key regulatory
proteins can be a powerful tool to dissect and manipulate the development and physiology of
cells to whole organisms [26,27]. Stomata are no exception [28]. For example, previous chemical
screens identified compounds that suppress stomatal movement [29,30]. Likewise, small
compounds have been utilized to examine the roles of signaling kinases and phosphatases in
stomatal development [31-33]. Chemical library screens identified bubblin, which severely disrupts
proper cell polarity in dividing meristemoids [34], as well as those that increase the number of
stomata [35]. However, in all the above cases, the chemical compounds impact multiple
processes, thus causing pleiotropic effects and/or their targets remain unidentified.

In an effort to expand the chemical space for controlling stomatal function and
development, we have created a collection of nitrogen-containing chemical compounds by
utilizing a newly developed amination reaction [36]. Nitrogen-containing molecules are widely
recognized for their potent biological activities [37]. Previous chemical screens and subsequent
structure-activity-relationship studies have shown that sulfonamidated compounds SIM1
(Stomata Influencing Molecule) and its derivative SIM3* inhibit stomatal movement [38]. Here, we
report a complex doubly-sulfonylated imidazolone compound named Stomidazolone as a potent
inhibitor of stomatal differentiation. We discover that Stomidazolone tightly binds to the MUTE
ACTL domain and perturbs the SCRM-MUTE heterodimerization. Additionally, we synthesized
and identified biologically-inactive Stomidazolone analogs that severely compromise MUTE
binding and its potency to interfere with SCRM-MUTE. Finally, we rationally designed
Stomidazolone-resistant versions of MUTE and demonstrated their efficacy *in vivo*. Together
with our chemical synthesis pipelines, our work will expand the chemical and pharmacological
space for bioactive compounds and open the future directions for selective perturbation of
prominent ACTL-domain-containing plant bHLH TFs for transient manipulation of plant growth
and development.

RESULTS

Doubly-sulfonylated imidazolone inhibits stomatal differentiation
In our initial attempt to derivatize SIM1 [38], we synthesized SIM1* from 2,4-diphenylimidazole
through C–H imidation followed by monodesulfonylation, and serendipitously found the first one-
shot synthesis of doubly-sulfonylated imidazolone (Fig. 1a; See Methods). Treatment of 2,4-
diphenyloxazole with chloramine B in the presence of CuI afforded this doubly-sulfonylated

AYSJ929 (later named as Stomidazolone: Fig. 1a, Document S1 for synthesis and NMR spectra

of the synthesized compounds). Further optimization produced Stomidazolone in 42% yield. Our

new reaction enabled the conversion of diphenyloxazole into substituted imidazolone with two

equivalents of chloramine (see Document S1). Even after searching the Reaxys database, we

failed to uncover any records of a reaction that transforming diphenyloxazole into imidazolones.

The structure was unambiguously confirmed by X-ray crystallographic analysis, which clearly

shows the acute angular shape of this doubly-sulfonylated imidazolone (Fig. 1b, Table S1,

Supplementary Document S1). Unlike SIM1 and its derivative SIM3* [38], this compound did not

exhibit any discernable effects on stomatal movement using our standardized assays (Fig. S1).

The synthesized compound features numerous aryl groups, oxygen and nitrogen atoms

as hydrogen bond acceptors, as well as hydrogen atoms serving as hydrogen bond donors (see

Fig. 1b; Document S1), suggestive of potential bioactivity. We thus subjected it to our chemical

library screening pipeline for stomatal development [35]. Intriguingly, its application to Arabidopsis

seedlings conferred reduced number of stomata but concomitant increase in the number of

meristemoids (Fig. 1c). These increased meristemoids properly express both early and late

meristemoid reporter markers, *TMMpro::GUS-GFP* and *MUTEpro::nucYFP*, respectively (Fig.

1d). Further quantitative analysis of stomata and meristemoid densities confirmed the dose-

dependency of the compound, now named Stomidazolone, to decrease the number of stomata

(24.45 %, 59.50 %, and 69.53 % decrease from mock to 20, 50, and 100 µM Stomidazolone,

respectively) while dramatically increasing the number of meristemoids (315.8 %, 801.5 %, and

647.6 % increase from mock to 20, 50, and 100 µM Stomidazolone, respectively) (Fig. 1e). On

the other hand, total number of stomata and meristemoids are only slightly reduced by the

Stomidazolone treatment (14.46 % and 25.03% at 20 and 50 µM) (Fig. 1e). These results

suggest that Stomidazolone inhibits stomatal differentiation at the transition from proliferating

meristemoids to differentiation.

Our bioassay pipelines utilizes a liquid culture system [35], which may impose some

limitation to the utility of Stomidazolone. We therefore investigated whether Stomidazolone

supplemented to solid media can be taken up to influence stomatal differentiation. Like in liquid

culture, Stomidazolone in the solid media inhibited stomatal differentiation in a dose-dependent

manner, with concomitant increase in arrested meristemoids (Fig. S2a-c). These results suggest

that Stomidazolone can be taken up by the seedling roots to impact stomatal development.

Stomidazolone does not affect overall seedling growth including root elongation and

cotyledon expansion, even at 100 µM in liquid culture (Fig. 1f). Likewise, Stomidazolone in the

solid media does not impact seedling morphology at 20 and 50 µM (Fig. S2d). It is thus highly

unlikely that the observed inhibition of meristemoid differentiation is due to general seedling
growth retardation. We therefore conclude that Stomidazolone potently triggers the
developmental arrest of the meristemoids, and in a lesser extent, inhibits the stomatal-lineage
initiation.
156
Stability of Stomidazolone and identification of inactive Stomidazolone analogs
To examine the stability of Stomidazolone and identify any degraded products (if unstable), we
treated Stomidazolone with an acid or a base. As shown in Document S1 we confirmed that
Stomidazolone is stable under acidic conditions (with 3 equiv. of acetic acid) but completely
degraded under basic conditions (with 3 equiv. of sodium hydroxide). The major degraded
product under basic conditions was isolated with gel-permeation chromatography (GPC) in 48%
yield. The product, which we designated as lzx103, was identified as
(phenylsulfonyl)benzimidamide [39] through ^{1}H and ^{13}C NMR analyses (Fig. S3a, Document S1).
Next, to directly address the stability of Stomidazolone in the aqueous condition we used
for our bioassays, we performed liquid chromatography-mass spectrometry (LC/MS) (Fig. S4).
Stomidazolone was incubated in the 1/2 MS liquid medium (pH 5.7) at room temperature for 48
hours. Stomidazolone peak was detected at the predicted elution time (2.47 min) whereas no
lzx103 peak (2.13 min) was detected (Fig. S4a, c, d). Additionally, we tested Stomidazolone's
stability at phosphate buffer (pH 7.0). Again, our LC/MS analysis did not detect the degradation
of Stomidazolone (Fig. S4b,c,d). Based on these results, we conclude that Stomidazolone is a
stable compound under biological conditions.
Subsequently, we performed bioassays of lzx103 on developing Arabidopsis seedlings.
Unlike Stomidazolone, the application of 50 µM lzx103 did not cause any effects on stomatal
development (Fig. S3b,c,d). Our results rule out the possibility that the inhibition of stomatal
differentiation is attributable to the degradation product rather than Stomidazolone itself. These
results establish Stomidazolone as a chemical compound with novel inhibitory activity on
stomatal differentiation.
Stomidazolone possesses a complex angular structure not shared by its precursor SIM1
or its degradation product lzx103 (see Figs. 1 and S3, Document S1). To further investigate the
structure-activity relationship of Stomidazolone as an inhibitor of stomatal differentiation, we
next synthesized two Stomidazolone analogs (Document S1). Specifically, the Stomidazolone
analogs, designated as AYSJ1059 and AYSJ1061 were synthesized from methylphenyl-
substituted oxazole and bromophenyl-substituted oxazole using the copper-catalyzed conditions
(Fig. 2a, Document S1). Unlike Stomidazolone, the application of AYSJ1059 as well as

AYSJ1061 at 50 µM to Arabidopsis seedlings did not affect the number of stomata (Fig. 2b, c),
indicating that these two Stomidazolone analogs do not possess bioactivity to inhibit stomatal
differentiation. Our results imply that the additional substituent (either methyl or bromo moieties)
may perturb Stomidazolone's ability to bind to its target(s).
190
Mechanistic genetic analyses delineate Stomidazolone as a potential inhibitor of MUTE
To unravel which pathways of stomatal development are impacted by Stomidazolone, we
conducted mechanistic studies leveraged by the availability of a series of mutants known to
regulate stomatal development (Fig. 3a). SPCH, MUTE, and FAMA consecutively drive
stomatal-lineage progression [9-11] (Fig. 3a). To address whether Stomidazolone interferes with
each cell-transitional step specified by these stomatal bHLH proteins, we applied Stomidazolone
to *spch*, *mute*, and *fama* mutant seedlings. As shown in Fig. 3b, c, Stomidazolone exerted no
effects on the pavement cell-only epidermis of *spch*. Likewise, the arrested meristemoid
phenotype of *mute* was not enhanced by Stomidazolone. On the other hand, Stomidazolone
significantly reduced the number of *fama* tumors whereas increased the arrested meristemoids
(Fig. 3c, d). Thus, *spch* and *mute* mutations are epistatic to Stomidazolone, suggesting that this
compound inhibits the MUTE-step of stomatal fate commitment.
It is established that *ERECTA*-family genes interact synergistically to enforce stomatal
development [14]. Among them, *ERECTA* primarily inhibits the entry into stomatal cell lineages, a
process regulated by *SPCH*, whereas *ERECTA-LIKE1* (*ERL1*) restricts the differentiation
potential of the meristemoids, a process orchestrated by *MUTE* [40-42] (Fig. 3a). The loss of
function in these receptor genes compromise the MAPK activation that inhibits SPCH and
MUTE protein accumulation [19,20,41]. Compared to wild type, *erecta* null allele (*er*: *er-105*) overly
produces stomatal-lineage cells [14] (Fig. 4a-d). Stomidazolone treatment exaggerated the *er-105*
phenotype: the number of stomata drastically decreased and the meristemoids vastly increased
(Fig. 4a-d, k). In *er erl1*, Stomidazolone treatment did not reduce the number of stomata,
although it increased the number of meristemoids (Fig. 4e, f, k). The result supports the
previous finding that the loss of *ERL1* intensifies stomatal differentiation by over-accumulation of
MUTE [41], thereby conferring partial resistance to Stomidazolone. The number of arrested
meristemoids is even more increased in *er erl2* (Fig. 4g, h), consistent with the loss of *ERL1* in
this genetic background. Finally, *er erl1 erl2* triple mutant seedlings appeared insensitive to
Stomidazolone (Fig. 4i, j). Combined, these rather complex results further support that
Stomidazolone primarily interferes with the MUTE activity for commitment to stomatal
differentiation. Our results also suggest that if ERECTA-family mediated signaling pathways are

absent, Stomidazolone cannot effectively interfere with the overly activated stomatal
differentiation pathway.

It is known that HD-ZIP IV-family gene *HDG2* positively regulates stomatal differentiation
by promoting *MUTE* expression and that its paralog *AtML1* enhances the *hdg2* mutant
phenotype [43] (Fig. 3a). We next tested whether loss-of-function mutations in these genes
enhance the effects of Stomidazolone to trigger meristemoid arrests. As shown in Fig. 4l, m,
Stomidazolone increased the meristemoid numbers of *hdg2* epidermis while the total numbers
of stomata and meristemoids were unaffected. Strikingly, *hdg2 atml1* double mutant exhibited
hypersensitivity to Stomidazolone, and its application exaggerated *mute*-like epidermal
phenotype (Fig. 4n, o). Taken together, a series of chemical-genetic dissections indicates that
Stomidazolone likely inhibits the activity of MUTE.

***In vivo* potency and selectivity of Stomidazolone relies on SCREAM**
SCRM and its partially redundant paralog, SCRM2, serve as a heterodimeric partner of MUTE
as well as SPCH and FAMA, and their activity is unequally required for stomatal cell-state
transitions [12] (Fig. 3a). We investigated whether the activity of SCRMs influences the effects of
Stomidazolone on stomatal differentiation (Fig. 5). Stomidazolone treatment to *scrm* loss-of-
function mutant (also known as *ice1-2*) severely reduced stomatal lineage initiation, resulting in
a *spch*-like epidermis vastly composed of pavement cells (Fig. 5a-d). This result is different from
other core stomatal development mutants, whereby Stomidazolone triggered a *mute*-like
phenotype (Figs. 3, 4). Unlike *SCRM*, *SCRM2* knockout mutant *scrm2-1* does not confer any
discernable phenotype [12]. The *scrm2-1* seedlings responded to Stomidazolone treatment just
like wild-type seedlings: reduction of the number of stomata with meristemoid arrests (Fig. 5e, f).
The complete loss of *SCRMs*, *scrm scrm2-1* double mutant, develops pavement cell-only
epidermis identical to *spch* [12]. Stomidazolone has no effects on *scrm scrm2-1* epidermis (Fig. 5g,
h). Conversely, a constitutively active form of SCRM, scrm-D confers stomata-only epidermis
[12,21]. This *scrm-D* phenotype was not affected by Stomidazolone (Fig. 5i, j). These results
suggest that Stomidazolone's potency as an inhibitor of stomatal differentiation relies on the
activity of SCRM. In the absence of SCRM, Stomidazolone can additionally inhibit the SPCH-
regulated step of stomatal-lineage initiation. Here, SCRM2 is the only available partner for
SPCH/MUTE/FAMA and as such, Stomidazolone might be able to target SCRM2 and SPCH.

Next, we sought to delineate if Stomidazolone interferes with the *in vivo*
heterodimerization potential of SCRM. Recently, we have shown that the SCRM C-terminal
ACTL domain serves as a heterodimerization interface and its dysfunctions selectively abrogate

heterodimerization of SCRM with MUTE, but not SPCH [23]. We therefore used our *scrm-D*

intragenic suppressor mutants, *scrm-D_S343* and *scrm-D_S423*, which encode scrm-D_{L484F} and

scrm-$D_{\Delta ACTL}$, respectively [23]. As reported previously [23], both *scrm-D_S343* and *scrm-D_S423*

exhibit nearly wild-type epidermis. Stomidazolone triggered excessive formation of arrested

meristemoids in these *scrm-D* intragenic suppressors (Fig. 5k-p). The phenotype is slightly

weaker yet resembling to that of *scrm-D mute* [44], further supporting the hypothesis that

Stomidazolone perturbs the activity of MUTE.

In the absence of *SCRM2*, both *scrm-D_S343* and *scrm-D_S423* confers massive

clusters of arrested meristemoids, identical to the phenotype of *scrm-D mute* [23]. This occurs

because scrm-D ACTL-domain mutant proteins (scrm-D_{L484F} and scrm-$D_{\Delta ACTL}$) cannot stably

heterodimerize with MUTE while still heterodimerizing with SPCH [23]. We thus predicted that if

Stomidazolone mainly targets the SCRM-MUTE (but not SCRM-SPCH) heterodimer, its

application will not affect the massive meristemoid cluster phenotype of *scrm-D_S343* and

scrm-D_S423 in the *scrm2* null mutant background. Consistent with this hypothesis, both *scrm-*

D_S343 scrm2-1 and *scrm-D_S423 scrm2-1* double mutant showed insensitivity to

Stomidazolone (Fig. 5 q,r). Taken together, our intricate *in vivo* phenotypic evidence highlights

that Stomidazolone inhibits meristemoid differentiation, most likely by disrupting the

heterodimerization of MUTE and SCRM through the C-terminal ACTL domain.

Stomidazolone directly targets MUTE and interferes with its heterodimerization

To address the molecular action of Stomidazolone, we investigated whether Stomidazolone

directly disrupts the heterodimerization of SCRM with MUTE (as well as other stomatal-lineage

bHLHs). First, we performed yeast two-hybrid (Y2H) assays. The growth assays show that

Stomidazolone diminishes the association of SCRM with MUTE as well as with SPCH and

FAMA (note that the N-terminal activation domain of SPCH and FAMA have been removed to

prevent their autoactivation) [21] (Fig. 6a). Further quantitative α-galactosidase assays show that

Stomidazolone inhibits these stomatal bHLH heterodimers in a dose-dependent manner, with

slightly less impact on the SPCH-SCRM heterodimers (Fig. 6b). Unlike Stomidazolone, its

precursor analog SIM3* has no effects on stomatal development nor heterodimerization of

stomatal bHLH proteins (Fig. S5).

To examine the effects of Stomidazolone on the heterodimerization of SCRM with full-

length SPCH, MUTE, or FAMA in the context of the plant cells, we next performed ratiometric

bimolecular fluorescent complementation (BiFC) assays. Briefly, *Nicotiana benthamiana* leaves

are transfected with a combination of SCRM and SPCH/MUTE/FAMA, each fused with a

complementary half YFP, along with a full-length RFP (mScarlet-I3) fused with Histone H2B.
Subsequently, leaf disks were prepared and treated with mock- or Stomidazolone (see Materials
and Methods for details). In the absence of Stomidazolone, strong YFP signals were detected in
the nuclei expressing the combination of SCRM-cYFP with SPCH/MUTE/FAMA-nYFP along
with RFP signals (Fig. 6c). Strikingly, however, Stomidazolone treatment severely diminished
the YFP signals in SCRM-cYFP + MUTE-nYFP pairwise combination (Fig. 6c). On the other
hand, its effects on SCRM-cYFP + SPCH-nYFP or SCRM-cYFP + FAMA-nYFP were unclear
(Fig. 6c). Subsequent quantitative analyses of YFP/RFP signal ratio confirm that Stomidazolone
severely perturbs the heterodimerization of SCRM-MUTE while weakly affecting SCRM-SPCH
but not affecting SCRM-FAMA (Fig. 6d).

We observed that Stomidazolone can confer a weak *spch*-like phenotype when applied
to *scrm* mutant seedlings (Fig. 5c, d). This rather unexpected phenotype can be explained if
Stomidazolone effectively blocks the heterodimerization of SPCH with SCRM2. To further
address this possibility, we conducted rBiFC assays with SCRM2 and SPCH/MUTE/FAMA. Like
SCRM, SCRM2 heterodimerizes with SPCH, MUTE, and FAMA, as detected by YFP signal
reconstitutions in the nuclei (Fig. S6a, b). Unlike the case of SCRM, however, Stomidazolone
treatment abolished the heterodimerization of SCRM2 with all of SPCH/MUTE/FAMA (Fig. S6a,
b). The findings that Stomidazolone disrupts SCRM2-SPCH, but not SCRM-SPCH heterodimers
fully explain the observed *in vivo* effects on Stomidazolone on *scrm* mutant (Fig. 5c, d).
Additionally, perturbation of SCRM2-MUTE heterodimers by Stomidazolone is consistent with
the exaggerated meristemoid arrest phenotypes by Stomidazolone treatment on *scrm-D_S343*
and *scrm-D_S423* seedlings (Fig. 5k-p).

To quantitatively assess the direct inhibition of SCRM-MUTE heterodimers by
Stomidazolone, we further performed two *in vitro* biophysical protein-protein interaction kinetics
assays using biolayer interferometry (BLI) and isothermal titration calorimetry (ITC) (Fig. 6e, f;
Figs. S7-S9, Tables S2, S3). For this purpose, recombinant full-length SCRM and MUTE
proteins were expressed and purified. Preincubation of Stomidazolone and MUTE decreased
the binding affinity of MUTE and SCRM by two orders of magnitude for both BLI and ITC
methods (Kd values increased from 7.5 ± 1.2 nM to 856.5 ± 15.1 nM in BLI and from 9.5 ± 1.6
nM to 900 ± 34.5 nM in ITC for SCRM-MUTE binding without or with pre-incubation with
Stomidazolone, respectively) (Figs. 6e, f, S7). Consistent with the lack of biological activity to
inhibit stomatal differentiation, the Stomidazolone analogs AYSJ1059 and AYSJ1061 showed
minimal effects on the SCRM-MUTE heterodimerization (Fig. S9, Tables S2, S3). These results
demonstrate that Stomidazolone directly interferes with the SCRM-MUTE heterodimerization.

We next tested whether Stomidazolone directly binds to MUTE. Indeed, the BLI assay

showed Stomidazolone binding to MUTE (Kd = 8.9 ± 1.2 μM) (Fig. 6g). Again, biologically

inactive Stomidazolone analogs, AYSJ1059 and AYSJ1061 (Fig. 2), exhibit no binding to MUTE

when analyzed by ITC and substantially weak binding by BLI (Fig. S8, Tables S2, S3). In

addition, we found that Stomidazolone binds with recombinant SPCH, FAMA, and SCRM

proteins, albeit with lower affinity (Fig. 6g, Fig. S8a, b, d, Table S3). These results support that

inhibition of stomatal differentiation by Stomidazolone primarily occurs via its direct binding to

MUTE.

Finally, to identify the binding domain of Stomidazolone on MUTE, we performed BLI

assays with the purified recombinant bHLH domain and ACTL domain of MUTE. As shown in

Fig. 6h, Stomidazolone specifically binds to the ACTL domain (Kd = 13.1 ± 1.3 μM). Based on

the ITC data, the binding is primarily driven by enthalpic interactions (Table S4). Again,

Stomidazolone showed a lower affinity with the ACTL domains of SCRM, SPCH, and FAMA

(Fig. S8e, f, h, Table S3). Together with a series of *in planta* and *in vitro* studies, the results

indicate that Stomidazolone directly and primarily targets the ACTL domain of MUTE and

interferes with the SCRM-MUTE heterodimerization. Furthermore, our results reinforce the

previous finding that the ACTL domain integrity is strictly required for the heterodimerization of

SCRM-MUTE [23], and convincingly provide the molecular, mechanistic basis of Stomidazolone

as an inhibitor of meristemoid differentiation.

Docking modeling identifies the Stomidazolone-MUTE binding interface

We demonstrated that the ACTL domain of MUTE is targeted by Stomidazolone (Fig. 6). To

delineate the specific binding interface, we performed computational docking modeling (Fig.

S10). Briefly, we used HADDOCK-based calculations to drive the docking process, using

mutagenesis data and binding site information from Swiss Dock as ambiguous interaction

restraints. Structural models showed that different racemic mixtures of Stomidazolone adopted

the same geometry in all clusters, with ionic interactions mediated by charged residues within

the MUTE ACTL domain, Arg-133 (R133), Arg-134 (R134), Glu-162 (E162), and Glu-163

(E163), and H-bonding and packing interactions by polar and hydrophobic residues Ile135,

Val136, Thr164 and Val165 (Fig. 7a, b, Fig. S10a). The binding free energies of both racemic

mixtures were similar (Fig. S10, Table S4). In contrast, the biologically inactive Stomidazolone

analogs AYSJ1059 and AYSJ1061, in both enantiomers, failed to converge into a discrete

binding model (Fig. S10b, c), consistent with its lack of direct binding to the MUTE ACTL

domain detected with ITC (Figs. S8l, m).

Amino-acid sequence alignments of MUTE and its orthologs as well as their sister

bHLHs (SPCH and FAMA) show that R133 and R134 residues are highly conserved among the

MUTE and FAMA clades, whereas substitutions to His (H) and Lys (K) are prevalent in the

SPCH clade (Fig. S11). The E162 and E163 residues are replaced by Glu (E) and Gln (Q) and

Asp (D) and D/E in FAMA and SPCH clades, respectively (Fig. S11). Overall, the

Stomidazolone-SPCH/MUTE/FAMA ACTL domain binding interfaces possess positively and

negatively charged residues with clade-specific sequences, which might reflect the differences

in physical binding affinities (Figs. 7b, S11, Tables S2, S3).

To confirm the predicted binding residues for Stomidazolone-MUTE interaction, we

further performed Alanine substitutions. For this purpose, we generated $MUTE_{R133A\_R134A}$,

$MUTE_{E162A\_E163A}$, and $MUTE_{R133A\_R134A\_E162A\_E163A}$ mutants and then quantitatively analyzed

Stomidazolone binding affinity on these mutant versions of MUTE ACTL domain by using

biophysical assays, BLI and ITC (Fig. 7c, Fig. S8i, j, k and Table S3). In both assays, the

binding affinity was significantly reduced; The Kd values obtained for BLI assays are 85.6 ± 5.5

μM and 125.0 ± 12.0 μM for double mutants, and no binding was detected for the quadruple

mutant (Fig. 7c, S8i, j, k, Tables S2, S3). These results provide the experimental evidence for

the predicted Stomidazolone-binding interface of specific MUTE residues within its ACTL

domain.

Engineering Stomidazolone-resistance via a molecular docking modeling

Our molecular docking modeling and experimental verification deciphered the Stomidazolone

binding interface at the MUTE ACTL domain (Fig. 7a, b). To the best of our knowledge, nothing

is known about the bioactivity of doubly-sulfonylated imidazolones nor drugs that target the

ACTL domains of bHLH TFs. We thus first explored the functional importance of the

Stomidazolone-binding sites within the MUTE ACTL domain. We expressed two different *MUTE*

mutant versions that are diminished in Stomidazolone binding (Fig. 7c) into *mute* knockout

(*mute-2*) plants driven by its own promoter (*MUTEpro::MUTE$_{E162A\_E163A}$-GFP* and *MUTEpro::*

MUTE$_{R133A\_R134A}$-GFP). We in addition generated a control *MUTEpro::MUTE-GFP* construct

using the identical molecular cloning strategy, which was also transformed into *mute-2* in

parallel (see Methods). Like the wild-type MUTE-GFP, the engineered versions of MUTE-GFP

are detected in the nucleus of a differentiating meristemoid, and their expression fully rescued

the *mute* arrested meristemoid phenotype (Fig. 7d, e, Fig. S12a). These transgenic *mute* null

lines grow into healthy, fertile plants indistinguishable from the wild type. Thus, the site-directed

mutations within the predicted Stomidazolone-MUTE interaction interface do not impact the

expression patterns or functionalities of MUTE.

Next, we tested the effects of Stomidazolone. As expected, control *MUTEpro::MUTE-*

GFP mute-2 seedlings significantly reduced the number of stomata upon Stomidazolone

treatment (Fig. 7e, f, Fig. S12b, c). In contrast, independent lines of $MUTEpro::MUTE_{E162A\_E163A}$-

GFP and $MUTEpro:: MUTE_{R133A\_R134A}$-*GFP* exhibit Stomidazolone resistance, because

Stomidazolone treatment did not significantly affect the numbers of stomata and meristemoids

(Fig. 7e, f, Fig. S12b, c). These results validate the structurally modelled Stomidazolone-MUTE

binding interface and unambiguously demonstrate that the *in vivo* binding of Stomidazolone to

the MUTE ACTL domain is the cause of the inhibition of stomatal differentiation. Furthermore,

our structure-guided engineering of Stomidazolone-resistant plants suggests that the

Stomidazolone-binding interface within the ACTL domain can be uncoupled from the biological

activity of the bHLH-ACTL TFs.

DISCUSSION

In this study, we developed a new catalytic amination and materialized a synthesis of

Stomidazolone, a complex, doubly-sulfonylated imidazolone that selectively inhibits stomatal

differentiation. Leveraged by the wealth of stomatal development mutant resources, our forward

chemical genetics pinpointed the exact *in vivo* target of Stomidazolone: the master regulator

MUTE. Further biochemical, biophysical, and structural modeling studies elucidated that

Stomidazolone primarily perturbs the SCRM-MUTE heterodimerization through interacting with

the ACTL-domain of MUTE. Chemical synthesis of TF dimerization inhibitors is an

under/unexplored field in plant biology. As such, our work may expand a chemical space for

manipulation of developmental processes.

In yeast, Stomidazolone interferes with the heterodimerization of not only SCRM-MUTE

but also SCRM-SPCH and SCRM-FAMA (Fig. 6a, b). Remarkably, in the context of *in vivo* plant

cells, however, Stomidazolone exhibits selective disruption of the SCRM-MUTE heterodimers

(Fig. 6c, d) and, consistently, confers meristemoids arrest in developing seedling epidermis

(Figs. 1, 3). It is known that SPCH, MUTE, and FAMA are closely related paralogs. Yet, they

possess distinct domains, with MUTE being the smallest without the extended N-terminal

domain [11]. SPCH, MUTE, and FAMA are known to associate with different protein complexes,

including those mediating transcription and epigenetic regulations [45-47]. It is therefore plausible

that association with distinct protein complexes may contribute to the *in vivo* selectivity of

Stomidazolone toward SCRM-MUTE, which cannot be recapitulated in the heterologous yeast

system (note that truncated SPCH and FAMA proteins lacking their N-terminal domain were

used in yeast in order to avoid transcriptional auto-activation) (Fig. 6a, b). Moreover, prevalent

effects of Stomidazolone to perturb SCRM-MUTE heterodimers may also reflect the inherent

property of bHLH-ACTL heterodimerization. It has been shown that heterodimerization of

SCRM-MUTE, but not SCRM-SPCH or SCRM-FAMA, relies on the presence or integrity of the

SCRM ACTL domain [23]. Thus, our findings support the notion that the SCRM-MUTE pair is the

most labile, hence druggable, amongst the stomatal bHLH heterodimers.

432   The *in vitro* Stomidazolone binding assays show that Stomidazolone most strongly

associates with the ACTL domain of MUTE but also weakly associate with that of SPCH, FAMA,

and SCRM (Fig. 6). Thus, whereas the specificity for perturbing the bHLH-ACTL

heterodimerization may be impacted by the *in vivo* protein complex status, the specificity of

direct binding of Stomidazolone to the ACTL domain lies in local protein sequence. In support of

this idea, the key Stomidazolone-interacting amino-acid residues within the MUTE ACTL

domain are not fully conserved in SPCH, FAMA, and SCRM (Fig. S11). Future structural

analysis of Stomidazolone-ACTL binding interface may clarify these points.

440   Our exhaustive mechanistic genetic analyses with Stomidazolone revealed that, in the

absence of SCRM, Stomidazolone vastly reduces the number of stomata, thus phenocopying

the weak *spch* mutant [9] (Fig. 5c, d). This unexpected phenotype underscores the unequal

redundancy between *SCRM* and its lesser-biologically potent paralog *SCRM2* [12]. In this study,

we showed that SCRM2-SPCH heterodimers, but not SCRM-SPCH heterodimers, are targeted

in vivo by Stomidazolone (Fig. S6). The high efficacy of Stomidazolone on perturbing SCRM2-

SPCH/MUTE/FAMA heterodimerization may likely stem from the less stable, labile nature of

SCRM2, which may result in weaker heterodimerization affinity. Consistently, it has been shown

that the constitutively active version of SCRM2, $SCRM2_{R203H}$, which possesses the Arginine-to-

Histidine substitution within its KRAAM motif identical to scrm-D, does not confer excessive

stomatal differentiation phenotype as severe as *scrm-D* [12]. In any event, the presence of SCRM,

including those with ACTL-domain mutations, can mask the effects of Stomidazolone on

SCRM2 (Fig. 5). In contrast, the mutations that stabilize SCRM confer Stomidazolone

insensitivity (Fig. 4i-k, 5e,f, m-o). They include loss of upstream receptor kinases (*er erl1 erl2*)[14],

constitutively active *scrm-D* [12,21], as well as *scrm-D* ACTL domain mutants in the absence of

SCRM2 [23]. Because these mutations prevent SPCH from undergoing MAPK-mediated,

phosphorylation-dependent degradation [19,21,23], the observed Stomidazolone insensitivity can be

explained if these stabilized versions of SCRM-SPCH heterodimers can compensate for the
reduction of SCRM-MUTE.

The discovery that Stomidazolone targets functionally-dispensable residues within the
MUTE ACTL domain brings several implications. First, Stomidazolone (and its derivatives) could
be optimized in the future to map the functional significance of ACTL domains among the bHLH
TFs. The ACTL domain-containing bHLH proteins are prominent within the land plant lineages [24].
Whereas studies indicated the roles of the ACTL domains in dimerization [25], their functional
significance or mode of action remains largely unclear. In the well-studied case of Maize R, its
ACTL domain functions as an auto-inhibitory homodimerization module that prevents the DNA
binding [48,49], a mechanism different from the stomatal bHLH-ACTL proteins [23]. Second, the
finding hints at a hitherto unidentified small ligand-binding property of the ACTL-domain-
containing bHLH proteins. The prototypical ACT domain is found in biosynthetic enzymes and
functions as a sensor domain for small metabolic ligands, including amino acids and thiamine [22].
On the contrary, it is unknown whether bHLH-ACTL TFs bind any endogenous small molecules.
If they do, how Stomidazolone interferes with such endogenous ligands would be an interesting
future direction. Third, as we successfully achieved for MUTE (Fig. 7, Fig. S12), other bHLH-
ACTL TFs could be engineered to generate drug-resistant versions to selectively modulate plant
phenotypes without compromising their biological functions. From a practical point of view,
Stomidazolone can be applied to both liquid and solid culture media to target specific bHLH-
ACTL heterodimers (Figs. S2) and it is quite stable in the MS media (Fig. S4). Moreover, our
streamlined synthesis of Stomidazolone analogs (Fig. 2, Document S1) may be expanded to
generate a library of potential ACTL-targeting imidazolones. In any event, further structural
analyses would help elucidate the exact biological significance and applicability of
Stomidazolone to delineate and manipulate bHLH-ACTL proteins.

Harnessing synthetic chemistry to disrupt TF dimerization - while it seems like an
obvious means of manipulating growth and development, this approach is surprisingly
unexplored in plants. Studies identified small proteins that can competitively interfere with TF
dimers and prevent their DNA binding [50]. For instance, LONG HYPOCOTYL IN FAR-RED1
(HFR1) protein titrates the bHLH TF, PHYTOCHROME INTERACTING FACTORs (PIFs), from
forming functional dimers [51]. In mammalian systems, natural compound library screens have
identified small molecules that disrupt the homodimerization of Hes1, an bHLH TF regulating
neural cell proliferation and differentiation [52,53]. In all these cases, unlike Stomidazolone, the
inhibitors target the core dimerization HLH modules that are essential for the TF function,
thereby leaving little room for designing and engineering the drug resistant versions of said TFs.

In the synthetic chemistry point of view, the serendipitous discoveries of Stomidazolone

and subsequent chemical synthesis proffered a new catalytic transformation of oxazoles [37,54].

Our newly discovered reaction can convert diphenyloxazole into imidazolone to create highly

angular compounds (see Document S1), which are unexplored in a pharmacology field. Now

that we demonstrated the interaction of this high degree of three-dimensionality of

Stomidazolone interacts with a discrete structural module of the ACTL domain and selectively

inhibits the key step of stomatal differentiation, further elaboration of our new amination reaction

may expand the chemical space for manipulating protein functions in plants and beyond.

Materials and Methods

Materials availability

Materials generated and used in this study will be available from Keiko Torii (ktorii@utexas.edu).

The synthesized imidazolones (Stomidazolone; ID AYSJ929) and biologically inactive analogs

(AYSJ1059 and AYSJ1061) may be limited in quantity. Inquiries for availability could be directed

to Kei Murakami (kei.murakami@kwansei.ac.jp).

Plant growth condition and chemical treatment

Arabidopsis accession Columbia (Col-0) was used as wild-type. The following

mutants/transgenic lines are reported elsewhere: *TMMpro::GUS-GFP* [13]; *spch-3/+, mute/+,* and

fama-1/+ [11]; *mute-2/+* [55]; *MUTEpro::nucYFP* [41]; *ice1-2/scrm* (SALK_003155), *scrm2-1*

(SAIL_808_B10), and *ice1-2/scrm scrm2-1*[12]; *er-105 erl1-2 erl2-1* [14]; *hdg2-2* (SALK_127828C)

and *hdg2-2 atml1-3* (SALK_128172)[43]; *scrm-D_S343, scrm-D_S423, scrm-D_S343 scrm2-1,*

and *scrm-D_S423 scrm2-1* [23]. Their genotype and transcript reduction were confirmed. The

presence of transgenes/mutant alleles were confirmed by genotyping. Liquid culture with

Stomidazolone, the analogs (AYSJ1059 and AYSJ1061), and degraded compound lzx103 was

performed as previously described [35], in which all molecules were dissolved in dimethyl

sulfoxide (DMSO, Molecular biology grade, Fujifilm Wako, Japan 046-21981), Col-0 seeds were

sown in 96-well plates (TL5003; True Line) containing 95 µL of 1/2 MS media containing 0.5%

sucrose with rotary shaking at 140 rpm under continuous light at 22 $^{\circ}$C. Five µL of compounds

were dissolved in DMSO at 10 mM and diluted with liquid 1/2 MS media to 1 mM and added on

1-day-old seedlings (final concentration, 50 µM). Seedlings were incubated for 8 days, and at

day 9 seedling abaxial cotyledons were observed through confocal microscopy. For liquid

culture of *scrm-D_S343*, *scrm-D_S423*, 1/2 MS liquid media without sucrose was used. For the

solid culture, 0, 20, 50 or 100 µM Stomidazolone were solidified with low-melting agarose

(5805S, PrimeGelTM Agarose LMT 1-20K, TAKARA Bio). Briefly, 1/2 MS salt, 0.5% sucrose,

and 0.8% low melting agarose were mixed in water and autoclaved. After cooling the medium at

around 30 $^{\circ}$C, Stomidazolone (500x, dissolved in DMSO) was added and poured into 90 mm

petri dishes (Asnol Petri Dish JP φ90 x 15mm, 3-1491-01, AS ONE). Col-0 seeds were sown on

Stomidazolone-containing 1/2 MS agarose media and grown under continuous light for 9 days.

Molecular cloning and generation of transgenic plants

The following constructs for transgenic plants were generated in this study: *MUTEpro::MUTE-*

GFP (pAH11), *MUTEpro::MUTE$_{E162AE163A}$-GFP* (pHS133), *MUTEpro::MUTE$_{R133AR134A}$-GFP*

(pHS134). The original *MUTEpro::MUTE-GFP* construct (pLJP155) was generated via a

different cloning strategy with different antibiotics selection [11]. We therefore re-generated the

wild-type Three-way Gateway version of *MUTEpro::MUTE-GFP* (pAH11) as a control. *MUTE*

genomic coding construct without stop codon (pAR205) and *MUTE* promoter construct

(pAR202) were subjected to the Gateway reaction with R4pGWB504 [56]. The wild-type MUTE-

GFP lines with the identical R4pGWB504 backbone enable a proper comparison of phenotypic

complementation. The mutant versions of *MUTE* constructs were generated in the same way

after site-directed mutagenesis using Q5 Site-directed mutagenesis kit (NEB) and subjected to

the Three-Way Gateway reaction after sequence confirmation. The constructs were then

transformed into the *Agrobacterium* strain GV3101 and floral dipped into *mute-2/+* plants. To

construct *CaMV35S::H2B-mScarlet-I3/pPZP211* (DKv1400), 35S promoter (*35Sp::RFP-CenH3*

with *DKp665-DKp1254*) [57], H2B-(SGGGG)2 linker (from DKv1200 [58] with DKp2412-DKp2599),

mScarlet-I3 (from codon optimized synthetic DNA for *A. thaliana*), Nos terminator (from

DKv1200 with DKp1251-DKp2598) were amplified using KOD one polymerase (TOYOBO). The

PCR products were assembled into BamHI/EcoRI-digested pPZP211 [59] using the NEBuilder

HiFi DNA Assembly (NEB). For detailed lists of oligo DNA and plasmid information, see Tables

S5 and S6, respectively.

Confocal microscopy

For confocal microscopy, cell peripheries of seedlings were visualized with propidium iodide

(Sigma, P4170). Images were acquired using LSM800 or LSM700 (Zeiss) using a 20x lens. The

GFP, YFP reporter and PI signals were detected with excitation at 488 nm and an emission filter

of 410 to 546 nm, at 488 nm and an emission filter of 410 to 546 nm, and with excitation at 561

nm and 582-617 nm emission range, respectively. Raw data were collected with 1024 x 1024

pixel images. For qualitative image presentation, Adobe Photoshop CC was used to trim and

uniformly adjust the contrast/brightness.

Yeast two hybrid assay

Y2H assays were performed using the Matchmaker^{TM} Two-Hybrid System (Clontech). Bait

(pGBKT7) and prey (pGADT7) constructs were co-transformed into the yeast strain AH109

using Frozen-EZ Yeast Transformation II Kit (Zymo Research, T2001). The resulting

transformants were spotted on SD (–Leu, –Trp) and SD (–Trp, –Leu, –His) selection media

containing different concentrations of 3-amino-1,2,4-triazole (Sigma, A8056) as previously

reported [21]. Quantitative α-galactosidase assay was also performed according to manufactural

instruction (Clontech). Briefly, the transformants were incubated in 2 ml SD medium with 0, 10,

50 100 μM Stomidazolone or SIM3* with rotary shake (220 rpm) at 30 ^{o}C overnight. The

cultured media containing yeast were transferred to microtubes and centrifuged to obtain the

supernatant. The colorimetric assay was performed by incubating 8 μL of supernatant with 24

μL Assay Buffer that contains 0.5 M sodium acetate (pH 4.5) (Nakalai tesque, 31137-25): 100

mM *p*-nitrophenyl α-D-Galactopyranoside Solution (Sigma N0877) = 2:1 (v/v) at 30 ^{o}C for 60 min,

then terminated by adding 960 μL 1X Stop Solution (0.1 M sodium carbonate, Fujifilm Wako

195-01582). After OD_{410} was recorded, α-galactosidase units were calculated according to the

protocol (Clontech, P3024-1).

Biomolecular Fluorescence Complementation (BiFC)

For ratiometric BiFC assays [60], the cDNAs of SPCH, MUTE, and FAMA were cloned into the

35S CaMV promoter-driven destination vector pSPYNE, which contains the N terminus of the

EYFP protein (nYFP-174 amino acids), while the cDNA of SCRM and SCRM2 was inserted into

pSPYCE, which contains the C terminus of the EYFP protein (cYFP-64 amino acids) (see Table

S5 for detailed information of the constructs). The constructs were then transformed into

Agrobacterium tumefaciens strain GV3101. The Agrobacterium was resuspended in infiltration

buffer (10 mM $MgCl_2$, 10 mM 2-(N-morpholino)ethanesulfonic acid [pH 5.6], and 150 μM

acetosyringone). Bacterial culture densities were adjusted to a final optical density (OD600) of

1.0, and the cell suspensions were incubated at room temperature for 4 hours prior to

infiltration. Equal volumes of cultures carrying the corresponding complementary pair of BiFC

constructs (YFPn and YFPc), along with *CaMV35S::Histone H2B-mScarlet-I3* and the silencing

suppressor plasmid p19 (a gift from Dr. Sir David Baulcombe) were mixed and then co-infiltrated

into 4-week-old *N. benthamiana* leaves. The infiltrated *N. benthamiana* leaves were incubated
at 25°C for 2 days. Subsequently, leaf discs were cut out and cultured in mock or in liquid 1/2
MS media in mock (DMSO) or 100 µM Stomidazolone for 1 day. Confocal imaging of *N.*
benthamiana leaves was performed using a Leica SP8 confocal microscope using a 20X
objective lens (Leica Microsystems), simultaneously capturing YFP (excitation at 518 nm and
emission at 530 nm) and mScarlet-I3 (excitation at 569 nm and emission at 600 nm) differential
interference contrast channels.

600
601 **Chemical synthesis**

2,4-diphenyloxazole   Chloramine B   Stomidazolone
                              2.0 equiv   (AYSJ929)

Synthesis of Stomidazolone (AYSJ929): 4 Å Molecular sieve powder (500 mg) was added into a
large screw-capped tube and heat-dried in vacuo for 1 h. 2,4-Diphenyloxazole (111 mg, 0.50
mmol), Chloramine B dihydrate (250 mg, 1.0 mmol, 2.0 equiv) and CuI (9.5 mg, 50 µmol, 10
mol%) were added into the tube with a magnetic stirring bar in open air. The tube was
evacuated and refilled with N_2 gas following the usual Schlenk technique. Anhydrous 1,2-
dichloroethane (5.0 mL, 0.10 M) was added to the tube via the rubber top, the tube capped and
its cap wrapped with parafilm. The reaction mixture was stirred and heated at 40 ºC for 12 h.
The mixture was then cooled to room temperature, and the crude mixture was filtered through a
pad of silica gel topped with Na_2SO_4 in a short column and concentrated *in vacuo*. Purification
by flash chromatography on silica gel (hexane/ethyl acetate = 4:1) followed by recrystallization
from CH_2Cl_2/methanol provided Stomidazolone (AYSJ929) in 42% yield (113 mg, 0.21 mmol) as
a colorless crystal. For details of chemical synthesis see Document S1. For synthesis of
biologically inactive Stomidazolone analogs AYSJ1059 and AYSJ1061 as well as the major
inactive degradation product lzx103 under alkaline condition, see Document S1.

NMR and mass-spectrometry
The high-resolution mass spectra were recorded on Thermo Fisher Scientific Exactive Plus
(ESI) with Thermo Scientific Dionex UltiMate 3000 Series UHPLC or Agilent Technologies 6130
Quadrupole Liquid Chromatography/Mass Spectrometry (LC-MS). Nuclear magnetic resonance

(NMR) spectra were recorded on JEOL JNM-ECA-600 (^{1}H 600 MHz, ^{13}C 150 MHz)
spectrometer. Chemical shifts for ^{1}H NMR are expressed in parts per million (ppm) relative to
residual CHCl$_3$ in CDCl$_3$ (δ 7.26 ppm), and for ^{13}C NMR in ppm relative to CDCl$_3$ (δ 77.2 ppm).
Data are reported as follows: chemical shift, multiplicity (s = singlet, d = doublet, dd = doublet of
doublets, t = triplets, dt = doublet of triplets, td = triplet of doublets, tt = triplet of triplets, q =
quartets, m = multiplets, brs = broad singlet), coupling constant (Hz), and integration. For the
original NMR and mass-spectrometry data, see Dataset S1.

X-ray crystallography

A suitable crystal was mounted with mineral oil on a MiTeGen MicroMeshes and transferred to
the goniometer of the kappa goniometer of a RIGAKU XtaLAB Synergy-S system with 1.2 kW
MicroMax-007HF microfocus rotating anode (Graphite-monochromated Mo Kα radiation (λ =
0.71073 Å)) and HyPix6000HE hybrid photon-counting detector. Cell parameters were
determined and refined, and raw frame data were integrated using CrysAlis^{Pro} (Agilent
Technologies, 2010). The structures were solved by direct methods with SHELXT [61] and refined
by full-matrix least-squares techniques against F^2 (SHELXL-2018/3) by using Olex2 software
package [62]. The intensities were corrected for Lorentz and polarization effects. The non-
hydrogen atoms were refined anisotropically. Hydrogen atoms were placed using AFIX
instructions. CCDC 2089650 contains the supplementary crystallographic data for this paper.
These data can be obtained free of charge from The Cambridge Crystallographic Data Centre
via www.ccdc.cam.ac.uk/data_request/cif. See Table S1 for X-ray crystallography data.

Liquid chromatography-mass spectroscopy (LC/MS)

Stomidazolone in 1/2 MS liquid media (pH 5.7) or phosphate buffer (pH 7.0) was detected by LC
(Agilent 1200, Agilent Technologies, Santa Clara, USA) coupled to a single quadrupole mass
spectrometer (Agilent 6130) equipped with an APCI-ES source. For the stability assay of
Stomidazolone, the analytical samples were prepared as follows: 100 µL 5 mM Stomidazolone
in DMSO was diluted with 900 µL 1/2MS (pH5.7) or 50 mM Phosphate buffer (pH 7.0). The
Stomidazolone solutions were incubated at room temperature with rotating. At 0 or 48 hr after
incubation, the 10 µL of solution were collected, diluted with water : acetonitrile = 50:50, and
applied to LC/MS. Stomidazolone and the degraded compound lzx103 were separated by
Agilent Poroshell 120 EC-C18 (2.1 × 100 mm, 2.7) column. The mobile phase consisted of 5
mM ammonium formate : acetonitrile = 80:20 at a flow rate of 0.6 mL/min. in isocratic mode, and
detected by mass spectrometry (MS). For the detection of these compounds, 0.5 µL of the

sample were injected, and the total run time was 7 min. Parameters were set as follows: drying

gas flow 13.0 L/min, nebulizer pressure 3100 Torr, drying gas temperature 350 °C, and capillary

voltage +4000 V (positive ionization) and 3500V (negative ionization). The electrospray

ionization mass spectra (ESI-MS) recorded in the mass-to-charge ratio range from m/z 80–1000

was collected. For identification of Stomidazolone and Izx103, selected ion monitoring (SIM)

mode was used, and the following m/z values were used to detect the different protonated

molecules $[M+H]^+$ at m/z 532 for Stomidazolone, and m/z 261 for Izx103. Identification of

Stomidazolone and Izx103 was exclusively based on LC retention time and high-resolution

mass spectra.

Recombinant protein expression and purification

A. thaliana SCRM (ICE1) (1-494), $_{\Delta N}$SPCH (98-364), MUTE (1-202), $_{\Delta N}$FAMA (190-414),

MUTE$_{\Delta C}$ (1-114), MUTE$_{R133A\_R134A}$, MUTE$_{E162A\_E163A}$, MUTE$_{R133A\_R134A\_E162A\_E163A}$, SCRM-ACTL

(405-494), MUTE-ACTL (114-202), SPCH-ACTL (285-364), FAMA-ACTL (306-414),

MUTE_ACTL$_{R133A\_R134A}$, MUTE_ACTL$_{E162A\_E163A}$, MUTE_ACTL$_{R133A\_R134A\_E162A\_E163A}$ were cloned

into pGEX-4T-1 vector with an N-terminal GST tag and a thrombin cleavage sequence. For

protein expression, the constructs were transformed into *E. coli* strain BL21. For each

transformant, a single colony was selected and incubated in 10 ml LB liquid media. The

overnight-incubated *E. coli* suspensions were transferred to 1L LB medium and incubated at

37 °C for around 2 h until the OD_{600} reached 0.4–0.6. The isopropyl β-D-1-

thiogalactopyranoside (0.25 µM of final concentration) was added to the cultures and the

strains were incubated at 22 °C for a further 18 h. GST-fused proteins were purified using

glutathione agarose resin. The soluble portion of the cell lysate was loaded onto a GST-

Sepharose column. Nonspecifically bound proteins were removed by washing the column with

20 mM Tris pH 8.0, 200 mM NaCl. The bound GST-fused protein was eluted with 10

mm glutathione and 20 mM Tris pH 8.0, 200 mM NaCl. The GST-fused proteins were

exchanged with phosphate-buffered saline buffer, and then the solution was treated with 50 µg

of thrombin for 10–12 h at 16 °C. The GST portion of the protein was cleaved during thrombin

digestion, and then the whole solution was reloaded onto the glutathione *S*-transferase column

to obtain pure protein. The purified proteins were further purified by gel filtration on a Superdex-

200 (GE) column using fast protein liquid chromatography and phosphate buffer (pH 7.2) as the

eluent. The purity of the protein was checked by SDS-PAGE. For detailed lists of oligo DNA and

plasmid information, see Tables S5 and S6, respectively.

Isothermal titration calorimetry (ITC)

Binding of the small molecule Stomidazolone to SCRM, MUTE, SPCH, FAMA, MUTEΔ_C,

MUTE_ACTL, SCRM_ACTL, SPCH_ACTL and FAMA_ACTL, MUTE$_{R133A\_134A}$, MUTE$_{E162A\_163A}$,

MUTE$_{R133A\_R134A\_E162A\_E163A}$, as well as SCRM binding to MUTE in the presence and absence of

Stomidazolone / AYSJ1059 /AYSJ1061 was characterized at 30 °C using a Malvern PEAQ-ITC

microcalorimeter. All the protein samples were dialyzed overnight using PBS buffer containing

2% DMSO and 1mM DTT. Stomidazolone was dissolved in PBS buffer containing 2% DMSO

and 1mM DTT. Titrations were performed by injecting 1 × 0.5-µL and 12 × 3-µL or 1 × 0.75-µL

and 18 × 2-µL or 1 × 0.2-µL and 25 × 1.3-µL aliquots of 1 mM Stomidazolone / AYSJ1059 /

AYSJ1061 to 5-30 µM of protein in PBS buffer pH 7.4, containing 2% DMSO and 1mM DTT. All

titrations were carried out at least twice. The raw data were corrected using buffer and protein

controls and analyzed using the software supplied by the manufacturer. See Table S3 for details.

Biolayer interferometry (BLI)

The binding affinities of SCRM to MUTE / MUTE$_{R133A\_134A}$ / MUTE$_{E162A\_E163A}$ /

MUTE$_{R133A\_R134A\_E162A\_E163A}$ in presence and absence of Stomidazolone and its analogues

(AYSJ1059 and AYSJ1061), and Stomidazolone / AYSJ1059 / AYSJ1061 interactions with

GST-tagged SCRM, MUTE, SPCH, FAMA, MUTE ΔACTL, MUTE_ACTL, SCRM_ACTL,

SPCH_ACTL and FAMA_ACTL, MUTE$_{R133A\_134A}$, MUTE$_{E162A\_E163A}$, MUTE$_{R133A\_R134A\_E162A\_E163A}$

were measured using the Octet Red96 system (ForteBio, Pall Life Sciences) following the

manufacturer's protocols. The optical probes coated with anti-GST were first loaded with

500 nM GST tagged proteins before kinetic binding analyses. The experiment was performed

in 96-well plates maintained at 30 °C. Each well was loaded with 200 µL reaction volume for

the experiment. The binding buffer used in these experiments contained 1× PBS supplemented

with 2% DMSO, 0.02% Tween20. The concentrations of the Stomidazolone as the analyte in

the binding buffer were 100, 50, 25, 12.5, 6.25, 3.12, and 1.56 µM. There was no binding of the

analytes to the unloaded probes as shown by the control wells. Binding kinetics to all seven

concentrations of the analytes were measured simultaneously using default parameters on the

instrument. The data were analyzed using the Octet data analysis software. The association

and dissociation curves were fit with the 1:1 homogeneous ligand model. The k_{obs} (observed

rate constant) values were used to calculate K_d, with steady-state analysis of the direct binding.

See Table S3 for details.

Protein structural modelling

MUTE and MUTE ACTL domain homology models were used from Alpha fold program
(https://alphafold.ebi.ac.uk/entry/Q9M8K6) The MUTE ACTL domain was subjected to
constrained energy minimization to allow the global energy minimization and structural analysis
using the AMBER 12 suite and VADAR [63,64]. Molecular docking of both racemic forms of
Stomidazolone, AYSJ1059, and AYSJ1061 to the MUTE ACTL domain monomers was carried
out using the high ambiguity-driven biomolecular docking (HADDOCK) approach [65,66].
Ambiguous interaction restraints (AIRs) were selected based on mutagenesis data and
preliminary AUTODOCK model [67]. The pairwise ligand interface RMSD matrix over all structures
was calculated, and the final structures were clustered using an RMSD cut-off value of 3.5 Å for
both ligand and protein. The clusters were then prioritized using RMSD and the HADDOCK
score (weighted sum of a combination of energy terms). See Table S4 for binding energies for
Stomidazolone-MUTE docking simulation.

Measurement of stomatal opening in *Commelina benghalensis*
The experiments were performed as described previously [30,68]. Briefly, Stomidazolone or ABA
were added to a multi-well plate with basal buffer [5 mM MES/bistrispropane (pH 6.5), 50 mM
KCl, and 0.1 mM $CaCl_2$] to be diluted to a final concentration of 50 or 20 μM, respectively. *C.*
benghalensis plants were incubated in the dark overnight; then, 4 mm diameter leaf discs were
excised from fully expanded leaves using a hole punch (Biopsy Punch, Kai Medical). Samples
were immersed in a basal buffer containing compounds for 30 min before being exposed to light
(150 μmol m^{-2} s^{-1} red light and 50 μmol m^{-2} s^{-1} blue light) for 3 h. The stomatal
apertures were observed using a stereoscopic microscope (BX43; Olympus). The microscopic
images were acquired with a charge-coupled device (CCD) camera (DP27; Olympus), followed
by autofocus image processing using cellSens standard software (Olympus). Measurement of
stomatal apertures of *C. benghalensis* was performed by using an automation program,
DeepStomata [69].

Quantification and statistical analysis
For quantitative analysis of the abaxial leaf epidermis, confocal images were taken on the days
after germination as indicated in the Figure legends. Preparation of images was done as
described previously [32,70]. For counting epidermal cell types, stomatal density, stomatal
precursor cells (meristemoids), total stomata-lineage cells (stomata and meristemoids) were
calculated by counting cell types in an area of 0.4 mm^2 (0.63 mm x 0.63 mm) and plotted as per
mm^2. A series of Z-stack confocal images was obtained to obtain images covering the epidermis

and capturing GFP/YFP signal from the reporter lines. The area and the number of epidermal cells were quantified by using FIJI-ImageJ. Statistical analyses were performed using R ver. 4.0.2. Sample sizes (n) are indicated and defined in each legend. For the multiple sample comparison, 1-way ANOVA was performed followed by a post hoc Tukey HSD test. For the two-sample comparison, Student's T-test was performed. Graphs were generated using R ggplot2 package. For all boxplots, each box represents IQR (interquartile range), in which top, middle line, and bottom indicating 75, 50, and 25 percentiles, respectively, and the bar representing maximum to minimal values. Each magenta dot represents jittered individual data point. The datapoints that go beyond the bar are outliers, which are indicated as black dots. If the data point is greater than Q3 + 1.5*IQR (above the third quartile is greater than 1.5 times of the interquartile range) or less than Q1 - 1.5*IQR, that data point is classified as outlier. The color of Boxplot graph was based on ColorBrewer.org. The value of n, the number of each experiment or samples, the means of error bars, and how statistical significance was defined are indicated in a relevant figure legend.

ACKNOWLEDGEMENTS

We thank Amanda Rychel and Alex Hofstetter for constructing the 3-Way Gateway version of *MUTEpro::MUTE-GFP*, Nagoya ITbM Molecular Structure Center for assistance in NMR and mass-spectrometry, Miki Muranaka for plant care, Jason McLellan for the access to ITC, and Kosuke Watanabe for assistance with X-ray crystal structure analysis. We appreciate Satoshi Matsubara and Tokihiro Furuhashi for their advice on the identification of lzx103. We especially thank Yuichiro Tsuchiya for the insightful discussion and generously offering the lab space/facility for A.N., and Eundeok Kim for comments. A part of this work was conducted in Institute for Molecular Science supported by ARIM (JPMXP1223MS5012). This work was supported by MEXT/JSPS KAKENHI (JP16H01237, JP17H06476 and JP19H00990) to K.U.T.; Japan Science and Technology Agency (JST) PRESTO (JPMJPR20D8), TOBE MAKI Scholarship Foundation, and Tatematsu Foundation to K.M.; and ITbM Research Award to S.J.Y., A.N., and S.-K.H. A.N. was in part supported by the JST CREST Award (JPMJCR1924). S.J.Y. was supported by the JSPS Research Fellowship for Young Scientists DC1. S.-K.H. was supported by the Young Leader Cultivation Program from the Institute of Advanced Research, Nagoya University. K.U.T. acknowledges the support from the Howard Hughes Medical Institute and Johnson & Johnson Centennial Chair of Plant Cell Biology at UT Austin.

CONTRIBUTIONS

A.N., K.M., S.J.Y., S.-K.H and K.U.T. conceived the study. K.U.T., K.M., S.J.Y., S.K-H., A.N., and K.M.S. designed the experiments. A.N., K.M.S., S.J.Y., H.S., C.M.C., T.S., H.K., R.I, S.K., Y.A., K.U.T. performed the experiments (forward chemical genetics/transgenic/phenotypic analysis A.N., S.J.Y., H.S., S.-K.H., C.M.C., H.K., K.U.T.; yeast/biochemical/biophysical experiments, K.M.S., R.I.; BiFC, H.S.; chemical synthesis, S.J.Y., K.H., Z.L., K.M.; structural analyses/modeling, K.M., K.S.M., Y.S.; stomatal movement assays, Y.A., T.K.). D.K. developed a codon-optimized plasmid; A.N., K.S.M., H.S., K.M., Y.A. and K.U.T. performed data analysis and visualization. K.U.T. wrote the manuscript draft. All authors contributed to the editing of the manuscript. K.M. and K.U.T. are responsible for project administration and funding acquisition.

COMPETING INTERESTS

The authors declare no competing interests.

DATA AVAILABILITY

Crystallographic data for the structure reported in this article have been deposited at the Cambridge Crystallographic Data Centre, under deposition numbers CCDC 2089650. Primer oligo DNA information and plasmid information are provided in Tables S5 and S6, respectively. Raw data and original spreadsheet of cell counting are in the process of deposition to the Texas Data Repository (https://dataverse.tdl.org/) and also will be provided by the corresponding authors upon request.

817
Figure 1. Stomidazolone is a potent chemical for meristemoid arrest
(a) Schematics of chemical synthesis. Supplementary Document S1 for the details of the
synthesis. Bottom right, structure of Stomidazolone (ID: AYSJ929).
(b) X-ray crystal structure of Stomidazolone shown as a stick model (White, H; Beige, C; Blue,
N; Yellow, S; Red, O).
(c) Confocal microscopy images of representative cotyledon abaxial epidermis from Arabidopsis
wild-type (WT) seedlings 9-days-after germination (dag) grown in the presence of 0 (mock), 20,

50, and 100 µM Stomidazolone. Orange brackets, stomatal-lineage cells with an arrested
meristemoid in the center. Scale bars, 50 µm.
(d) Arrested meristemoids by Stomidazolone express proper meristemoid markers. Shown are
confocal microscope images of 9-dag representative abaxial cotyledons epidermis expressing
TMMpro::GUS-GFP (top) and *MUTEpro::nucYFP* (bottom) grown in the presence of 0 (mock:
Left) and 50 µM Stomidazolone (Right). Scale bars, 50 µm
(e) Stomidazolone treatment reduces the number of stomata but increases that of meristemoids.
Quantitative analysis of stomatal (s; green), meristemoid (m; cyan), and stomata+meristemoid
(t; gray) density per mm^2 of cotyledon abaxial epidermis from seedlings grown in the presence
of 0 (mock), 20, 50, and 100 µM Stomidazolone. One-way ANOVA followed by Tukey's HSD
analysis were performed for each cell type (stomata and meristemoids). Letters (a,b,c) indicate
groups that are statistically different from other groups within each cell type. n = 8
(f) Stomidazolone does not impact seedling growth. Shown are two representative 9-dag WT
seedlings, grown in the presence of 0 (mock), 20, 50, and 100 µM Stomidazolone. Scale bars,
10 mm.
840

841
842

843
844
845

Figure 2. Identification of biologically inactive Stomidazlone analogs

(a) Chemical structures of Stomidazolone and its derivatives AYSJ1059 and AYSJ1061. See Document S1 for detailed procedures of synthesis and analysis.

(b) Representative confocal images of cotyledon abaxial epidermis from seedlings 9-days-after-germination grown with mock, or 50 µM Stomidazolone, or its derivatives grown on liquid culture medium. Scale bars, 50 µm.

(c) Quantitative analysis. Number of stomata per 1 mm^2 of cotyledon abaxial epidermis from seedlings grown in the presence of mock or 50 µM Stomidazolone and its derivatives. Student's T test was performed and p values are indicated. ***, $p<0.0005$. n = 8.

856
857
Figure 3. *spch* and *mute*, but not *fama*, are epistatic to Stomidazolone
(a) Schematic diagram showing the genetic control of stomatal cell-state transitions. Key
genes/gene products and their roles are indicated in the cartoon. See main text for details of
each step. Arrows, positive regulation; T-bars, negative regulation. Upregulation of *MUTE* by
HDG2 and inhibition of *MUTE* by *ERL1* are highlighted in red. MMC, meristemoid mother cell;
GMC, guard mother cell; GC, guard cell.

(b) Representative cotyledon abaxial epidermis of mock and 50 µM Stomidazolone-treated WT,

spch (spch-3), *mute (mute-1)*, and *fama*. Scale bars, 50 µm.

(c, d) Quantitative analysis of meristemoids density per 1 mm^2 (c) and *fama* tumors (d) of

cotyledon abaxial epidermis from seedlings grown in the presence of 0 (mock) and 50 µM

Stomidazolone. Student's T test was performed for each plant. p values and NS (not significant)

are indicated above each boxplot. n = 8.

Figure 4. Stomidazolone is a potent inhibitor of cell-state transition at the MUTE-step

(a-j) Effects of Stomidazolone on the epidermis of wild-type (WT: a, b), *er* (c, d), *er erl1* (e, f), *er erl2* (g, h), and *er erl1 erl2* (i, j). Representative cotyledon abaxial epidermis of mock- (left) and 50 µM Stomidazolone (right). Orange brackets, clusters of arrested meristemoids. Scale bars, 50 µm.

(b, d, f, h, j) Quantitative analysis of stomatal (s; green), meristemoid (m; cyan), and stomata+meristemoid (t; gray) density per mm^2 of cotyledon abaxial epidermis from seedlings grown in the presence of 0 (mock) and 50 µM Stomidazolone (Stomid). Student's T test was performed for each plant. p values are indicated above each boxplot. n = 8.

(k) Number of meristemods, data replotted from other panels.

(l-o) Effects of Stomidazolone on the epidermis of *hdg2* (*hdg2-2*: l, m) and *hdg2 atml1* (*hdg2-2 atml1-3*: n, o). All experiments and analyses were performed as described above. Scale bars, 50 µm. n = 8.

887
888
Figure 5. SCRM activity dictates *in vivo* potency and selectivity of Stomidazolone
(a, c, e, g, i) Abaxial epidermis of cotyledons in wild-type (WT: a), *scrm* (*ice1-2:* c), *scrm2-1* (e),
scrm scrm2-1 (g), and *scrm-D* (e) 9-dag seedlings treated with mock or 50 µM Stomidazolone.
scrm loss-of-function mutants occasionally exhibit parallelly paired-stomata, so called 'four-lips'
phenotype (orange plus) as well as stomata made with an extra guard cell (orange asterisk)
likely due to *FAMA* mis-regulation. Scale bars, 50 µm.
(b, d, f, h, j) Stomatal index (SI, %) for Mock- or Stomidazolone-treated (Stomid) WT (b), *scrm*
(d), *scrm2-1* (f), *scrm scrm2-1* (h), and *scrm-D* (j). n=8. Student's T-test was performed for
pairwise comparison.
(k, m, o) Abaxial epidermis of cotyledons in WT (k) and the intragenic *scrm-D* suppressors:
scrm-D_S343 (m) and *scrm_D_S423* (o) 9-dag seedlings treated with mock or 50 µM
Stomidazolone. Clustered meristemoids and their sister stomatal-lineage ground cells (SLGCs:
orange brackets) become exaggerated. Scale bars, 50 µm.
(l, n, p) Stomatal and small cell (SC) index (%) for Mock- or Stomidazolone-treated (Stomid) WT
(h), *scrm-D_S343* (j) and *scrm_D_S423* (l). n=8. Student's T-test was performed for pairwise
comparisons.

(q, r) Abaxial epidermis of cotyledons in the intragenic *scrm-D* suppressors in the absence of
SCRM2: *scrm-D_S343 scrm2-1* (q) and *scrm_D_S423 scrm2-1* (r) 9-dag seedlings treated with
mock or 50 µM Stomidazolone. Scale bars, 20 µm.
908
909

911
912

Figure 6. Stomidazolone primarily disrupts SCRM-MUTE heterodimers through direct binding

(a) Y2H analysis. Yeasts expressing control and pairs of stomatal bHLH proteins were spotted in 10-fold serial dilutions on appropriate selection media with or without 50 µM Stomidazolone. 1 mM 3-AT treatment shows that Stomidazolone reduces the interactions of SCRM with stomatal bHLH proteins. 3-AT, 3-amino-1,2,4-triazole.

(b) Quantitative α-galactosidase assays. Yeasts expressing control and pairs of stomatal bHLH proteins were cultured in 0, 10, 50, and 100 µM Stomidazolone and enzyme assays were performed. One-way ANOVA followed by Tukey's HSD analysis were performed. Letters (a,b,c) indicate that the enzyme activities are statistically different within the group. Experiments were repeated at least 3 times.

(c) Ratiometric BiFC analysis. *N. benthamiana* leaves are infiltrated with Histone H2B (H2B)-mScarlet-I3 and pairs of stomatal bHLH proteins. Subsequently, leaf disks were treated either with mock (top) or 100 µM Stomidazolone (bottom). Shown are representative nucleus from each combination imaged simultaneously for mScarlet-I3 (control) and YFP (interaction) signals. Scale bars, 5 µm.

(d) Quantitative analysis of BiFC YFP signal intensity ratio normalized by the signal intensity of
H2B-mScarlet-I3. Student T-test was performed for each pairwise comparison. NS, not
significant. n=6.
(e, f) Quantitative Analysis of SCRM-MUTE Interactions in the absence (e), and presence (f) of
Stomidazolone. *In vitro* binding response curves are provided for SCRM with MUTE and
with/without Stomidazolone. The data represent the mean ± SD and are representative of two
independent experiments.
(g) Quantitative analysis of Stomidazolone binding with SCRM (blue), MUTE (cyan), FAMA
(green), and SPCH (lavender). The MUTE protein exhibit higher affinity to Stomidazolone. *In*
vitro   binding   response   curves   are   provided   Stomidazolone   with
SCRM/MUTE/SPCH/MUTE/FAMA. The data represent the mean ± SD and are representative
of two independent experiments.
(h) Quantitative analysis of Stomidazolone binding with MUTE bHLH (green) and ACTL (brown)
domains. *In vitro* binding response curves are provided for Stomidazolone with MUTE$_{bHLH}$ and
MUTE$_{ACTL}$. The data represent the mean ± SD and are representative of two independent
experiments.
946
947

Figure 7. Engineering Stomidazolone-resistant plants based on the MUTE-Stomidazolone docking model

(a, b) MUTE-Stomidazolone binding models. (a) Alphafold2 predicted full-length MUTE protein is shown as ribbon representation. MUTE bHLH domain, green; ACTL domain, peach; Linker region, red. Stomidazolone is shown as stick model (gray). (b) Magnified view of MUTE-Stomidazolone binding interface. The key binding residues (R133, R134, E162 and E163) sidechains are highlighted in magenta.

(c) Quantitative analysis of Stomidazolone binding to wild-type MUTE and MUTE site-directed mutants (MUTE$_{R133A\_R134A}$, MUTE$_{E162A\_E163A}$, MUTE$_{R133A\_R134AE162A\_E163A}$) by BLI. *In vitro* binding response curves are presented for Stomidazolone with MUTE and its mutants. Stomidazolone compound was tested at seven different concentrations (200, 100, 50, 25, 12.5, 6.25, and 3.125 μM). The data represent the mean□±□SD and are representative of two independent experiments.

(d) Site-directed mutant versions of MUTE that disrupt Stomidazolone binding are expressed normally and biologically functional. Representative cotyledon abaxial epidermis of 5-day-old *mute-2* seedlings expressing *MUTEpro::MUTE-GFP* (left), *MUTEpro::MUTE$_{E162A\_E163A}$-GFP* (center), and *MUTEpro::MUTE$_{R133A\_R134A}$-GFP* (right). Insets, representative late meristemoids from each panel with MUTE-GFP signals in nucleus. Confocal images are inverted for clear cellular outlines. Scale bars, 20 μm.

(e-f) MUTE mutant versions that disrupt Stomidazolone binding confer drug resistance *in vivo*.

(e) Representative cotyledon abaxial epidermis of 9-day-old *mute-2* seedlings expressing

MUTEpro::MUTE-GFP (left), *MUTEpro::MUTE$_{E162A\_E163A}$ -GFP* (center), and
MUTEpro::MUTE$_{R133A\_R134A}$ -GFP (right) with mock- (top) and 20 μM Stomidazolone for eight
days (bottom). Scale bars, 50 μm.
(f) Quantitative analysis of total (t: stomata and meristemoids), stomatal (s: green) or
meristemoid (m: light cyan) density per mm^2 of cotyledon abaxial epidermis from seedlings
grown in the presence of 0 (mock) and 20 μM Stomidazolone. Student's T test was performed
were performed for each cell type (total, stomata and meristemoids) and p values are indicated.
*, $p<0.05$, ***, $p<0.0005$. Experiments were repeated twice. n = 8.
979
980
981
982

REFERENCES

Hetherington, A. M. & Woodward, F. I. (2003) The role of stomata in sensing and driving environmental change. *Nature* **424**, 901-908 (2003).

Hsu, P. K., Dubeaux, G., Takahashi, Y. & Schroeder, J. I. (2021) Signaling mechanisms in abscisic acid-mediated stomatal closure. *Plant J* **105**, 307-321 (2021).

Kim, T. H., Bohmer, M., Hu, H., Nishimura, N. & Schroeder, J. I. (2010) Guard cell signal transduction network: advances in understanding abscisic acid, CO2, and Ca2+ signaling. *Annu Rev Plant Biol* **61**, 561-591 (2010).

Qi, X. & Torii, K. U. (2018) Hormonal and environmental signals guiding stomatal development. *BMC Biol* **16**, 21 (2018).

Bertolino, L. T., Caine, R. S. & Gray, J. E. (2019) Impact of Stomatal Density and Morphology on Water-Use Efficiency in a Changing World. *Front Plant Sci* **10**, 225 (2019).

Han, S. K. & Torii, K. U. (2016) Lineage-specific stem cells, signals and asymmetries during stomatal development. *Development* **143**, 1259-1270 (2016).

Lau, O. S. & Bergmann, D. C. (2012) Stomatal development: a plant's perspective on cell polarity, cell fate transitions and intercellular communication. *Development* **139**, 3683-3692 (2012).

Nadeau, J. A. & Sack, F. D. (2002) Stomatal development in Arabidopsis. *Arabidopsis Book* **1**, e0066 (2002).

MacAlister, C. A., Ohashi-Ito, K. & Bergmann, D. C. (2007) Transcription factor control of asymmetric cell divisions that establish the stomatal lineage. *Nature* **445**, 537-540 (2007).

Ohashi-Ito, K. & Bergmann, D. C. (2006) Arabidopsis FAMA controls the final proliferation/differentiation switch during stomatal development. *Plant Cell* **18**, 2493-2505 (2006).

Pillitteri, L. J., Sloan, D. B., Bogenschutz, N. L. & Torii, K. U. (2007) Termination of asymmetric cell division and differentiation of stomata. *Nature* **445**, 501-505 (2007).

Kanaoka, M. M., Pillitteri, L. J., Fujii, H., Yoshida, Y., Bogenschutz, N. L., Takabayashi, J., Zhu, J. K. & Torii, K. U. (2008) SCREAM/ICE1 and SCREAM2 specify three cell-state transitional steps leading to arabidopsis stomatal differentiation. *Plant Cell* **20**, 1775-1785 (2008).

Nadeau, J. A. & Sack, F. D. (2002) Control of stomatal distribution on the Arabidopsis leaf surface. *Science* **296**, 1697-1700 (2002).

Shpak, E. D., McAbee, J. M., Pillitteri, L. J. & Torii, K. U. (2005) Stomatal patterning and differentiation by synergistic interactions of receptor kinases. *Science* **309**, 290-293 (2005).

Hara, K., Kajita, R., Torii, K. U., Bergmann, D. C. & Kakimoto, T. (2007) The secretory peptide gene EPF1 enforces the stomatal one-cell-spacing rule. *Genes Dev* **21**, 1720-1725 (2007).

Hara, K., Yokoo, T., Kajita, R., Onishi, T., Yahata, S., Peterson, K. M., Torii, K. U. & Kakimoto, T. (2009) Epidermal cell density is autoregulated via a secretory peptide, EPIDERMAL PATTERNING FACTOR 2 in Arabidopsis leaves. *Plant Cell Physiol* **50**, 1019-1031 (2009).

Hunt, L. & Gray, J. E. (2009) The signaling peptide EPF2 controls asymmetric cell divisions during stomatal development. *Curr Biol* **19**, 864-869 (2009).

Lee, J. S., Kuroha, T., Hnilova, M., Khatayevich, D., Kanaoka, M. M., McAbee, J. M., Sarikaya, M., Tamerler, C. & Torii, K. U. (2012) Direct interaction of ligand-receptor pairs specifying stomatal patterning. *Genes Dev* **26**, 126-136 (2012).

1032	19	Lampard, G. R., Macalister, C. A. & Bergmann, D. C. (2008) Arabidopsis stomatal
1033		initiation is controlled by MAPK-mediated regulation of the bHLH SPEECHLESS.
1034		*Science* **322**, 1113-1116 (2008).
1035	20	Lampard, G. R., Lukowitz, W., Ellis, B. E. & Bergmann, D. C. (2009) Novel and
1036		expanded roles for MAPK signaling in Arabidopsis stomatal cell fate revealed by cell
1037		type-specific manipulations. *Plant Cell* **21**, 3506-3517 (2009).
1038	21	Putarjunan, A., Ruble, J., Srivastava, A., Zhao, C., Rychel, A. L., Hofstetter, A. K., Tang,
1039		X., Zhu, J. K., Tama, F., Zheng, N. & Torii, K. U. (2019) Bipartite anchoring of SCREAM
1040		enforces stomatal initiation by coupling MAP kinases to SPEECHLESS. *Nat Plants* **5**,
1041		742-754 (2019).
1042	22	Grant, G. A. (2006) The ACT domain: a small molecule binding domain and its role as a
1043		common regulatory element. *J Biol Chem* **281**, 33825-33829 (2006).
1044	23	Seo, H., Sepuru, K. M., Putarjunan, A., Aguirre, L., Burrows, B. A. & Torii, K. U. (2022)
1045		Intragenic suppressors unravel the role of the SCREAM ACT-like domain for bHLH
1046		partner selectivity in stomatal development. *Proc Natl Acad Sci U S A* **119:** (2022).
1047	24	Lee, Y. S., Shiu, S. H. & Grotewold, E. (2023) Evolution and diversification of the ACT-
1048		like domain associated with plant basic helix-loop-helix transcription factors. *Proc Natl*
1049		*Acad Sci U S A* **120**, e2219469120 (2023).
1050	25	Lee, Y. S., Herrera-Tequia, A., Silwal, J., Geiger, J. H. & Grotewold, E. (2021) A
1051		hydrophobic residue stabilizes dimers of regulatory ACT-like domains in plant basic
1052		helix-loop-helix transcription factors. *J Biol Chem*, 100708 (2021).
1053	26	Spring, D. R. (2005) Chemical genetics to chemical genomics: small molecules offer big
1054		insights. *Chem Soc Rev* **34**, 472-482 (2005).
1055	27	McCourt, P. & Desveaux, D. (2010) Plant chemical genetics. *New Phytol* **185**, 15-26
1056		(2010).
1057	28	Kinoshita, T., Toh, S. & Torii, K. U. (2021) Chemical control of stomatal function and
1058		development. *Curr Opin Plant Biol* **60**, 102010 (2021).
1059	29	Kim, T. H., Hauser, F., Ha, T., Xue, S., Bohmer, M., Nishimura, N., Munemasa, S.,
1060		Hubbard, K., Peine, N., Lee, B. H., Lee, S., Robert, N., Parker, J. E. & Schroeder, J. I.
1061		(2011) Chemical genetics reveals negative regulation of abscisic acid signaling by a
1062		plant immune response pathway. *Curr Biol* **21**, 990-997 (2011).
1063	30	Toh, S., Inoue, S., Toda, Y., Yuki, T., Suzuki, K., Hamamoto, S., Fukatsu, K., Aoki, S.,
1064		Uchida, M., Asai, E., Uozumi, N., Sato, A. & Kinoshita, T. (2018) Identification and
1065		Characterization of Compounds that Affect Stomatal Movements. *Plant Cell Physiol* **59**,
1066		1568-1580 (2018).
1067	31	De Rybel, B., Audenaert, D., Vert, G., Rozhon, W., Mayerhofer, J., Peelman, F., Coutuer,
1068		S., Denayer, T., Jansen, L., Nguyen, L., Vanhoutte, I., Beemster, G. T., Vleminckx, K.,
1069		Jonak, C., Chory, J., Inze, D., Russinova, E. & Beeckman, T. (2009) Chemical inhibition
1070		of a subset of Arabidopsis thaliana GSK3-like kinases activates brassinosteroid signaling.
1071		*Chem Biol* **16**, 594-604 (2009).
1072	32	Houbaert, A., Zhang, C., Tiwari, M., Wang, K., de Marcos Serrano, A., Savatin, D. V.,
1073		Urs, M. J., Zhiponova, M. K., Gudesblat, G. E., Vanhoutte, I., Eeckhout, D., Boeren, S.,
1074		Karimi, M., Betti, C., Jacobs, T., Fenoll, C., Mena, M., de Vries, S., De Jaeger, G. &
1075		Russinova, E. (2018) POLAR-guided signalling complex assembly and localization drive
1076		asymmetric cell division. *Nature* **563**, 574-578 (2018).
1077	33	Bian, C., Guo, X., Zhang, Y., Wang, L., Xu, T., DeLong, A. & Dong, J. (2020) Protein
1078		phosphatase 2A promotes stomatal development by stabilizing SPEECHLESS in
1079		Arabidopsis. *Proc Natl Acad Sci U S A* **117**, 13127-13137 (2020).
1080	34	Sakai, Y., Sugano, S. S., Kawase, T., Shirakawa, M., Imai, Y., Kawamoto, Y., Sugiyama,
1081		H., Nakagawa, T., Hara-Nishimura, I. & Shimada, T. (2017) The chemical compound

1082   bubblin induces stomatal mispatterning in Arabidopsis by disrupting the intrinsic polarity
1083   of stomatal lineage cells. *Development* **144**, 499-506 (2017).

1084   35   Ziadi, A., Uchida, N., Kato, H., Hisamatsu, R., Sato, A., Hagihara, S., Itami, K. & Torii, K.
1085   U. (2017) Discovery of synthetic small molecules that enhance the number of stomata:
1086   C-H functionalization chemistry for plant biology. *Chem Commun (Camb)* **53**, 9632-
1087   9635 (2017).

1088   36   Ito, E., Fukushima, T., Kawakami, T., Murakami, K. & Itami, K. (2017) Catalytic
1089   Dehydrogenative C-H Imidation of Arenes Enabled by Photo-generated Hole Donation to
1090   Sulfonimide. *Chem* **2**, 383-392 (2017).

1091   37   Hili, R. & Yudin, A. K. (2006) Making carbon-nitrogen bonds in biological and chemical
1092   synthesis. *Nat Chem Biol* **2**, 284-287 (2006).

1093   38   Toda, Y., Perry, G. J. P., Inoue, S., Ito, E., Kawakami, T., Narouz, M. R., Takahashi, K.,
1094   Aihara, Y., Maeda, B., Kinoshita, T., Itami, K. & Murakami, K. (2022) Identification of
1095   stomatal-regulating molecules from de novo arylamine collection through aromatic C-H
1096   amination. *Sci Rep* **12**, 949 (2022).

1097   39   Yi, F., Sun, Q., Sun, J., Fu, C. & Yi, W. (2019) Terminal Alkyne-Assisted One-Pot
1098   Synthesis of Arylamidines: Carbon Source of the Amidine Group from Oxime Chlorides.
1099   *J Org Chem* **84**, 6780-6787 (2019).

1100   40   Horst, R. J., Fujita, H., Lee, J. S., Rychel, A. L., Garrick, J. M., Kawaguchi, M., Peterson,
1101   K. M. & Torii, K. U. (2015) Molecular Framework of a Regulatory Circuit Initiating Two-
1102   Dimensional Spatial Patterning of Stomatal Lineage. *PLoS Genet* **11**, e1005374 (2015).

1103   41   Qi, X., Han, S. K., Dang, J. H., Garrick, J. M., Ito, M., Hofstetter, A. K. & Torii, K. U.
1104   (2017) Autocrine regulation of stomatal differentiation potential by EPF1 and ERECTA-
1105   LIKE1 ligand-receptor signaling. *Elife* **6**, e24102 (2017).

1106   42   Qi, X., Yoshinari, A., Bai, P., Maes, M., Zeng, S. M. & Torii, K. U. (2020) The manifold
1107   actions of signaling peptides on subcellular dynamics of a receptor specify stomatal cell
1108   fate. *Elife* **9:** (2020).

1109   43   Peterson, K. M., Shyu, C., Burr, C. A., Horst, R. J., Kanaoka, M. M., Omae, M., Sato, Y.
1110   & Torii, K. U. (2013) Arabidopsis homeodomain-leucine zipper IV proteins promote
1111   stomatal development and ectopically induce stomata beyond the epidermis.
1112   *Development* **140**, 1924-1935 (2013).

1113   44   Pillitteri, L. J., Peterson, K. M., Horst, R. J. & Torii, K. U. (2011) Molecular profiling of
1114   stomatal meristemoids reveals new component of asymmetric cell division and
1115   commonalities among stem cell populations in Arabidopsis. *Plant Cell* **23**, 3260-3275
1116   (2011).

1117   45   Kim, E. D., Dorrity, M. W., Fitzgerald, B. A., Seo, H., Sepuru, K. M., Queitsch, C.,
1118   Mitsuda, N., Han, S. K. & Torii, K. U. (2022) Dynamic chromatin accessibility deploys
1119   heterotypic cis/trans-acting factors driving stomatal cell-fate commitment. *Nat Plants* **8**,
1120   1453-1466 (2022).

1121   46   Matos, J. L., Lau, O. S., Hachez, C., Cruz-Ramirez, A., Scheres, B. & Bergmann, D. C.
1122   (2014) Irreversible fate commitment in the Arabidopsis stomatal lineage requires a
1123   FAMA and RETINOBLASTOMA-RELATED module. *Elife* **3:** (2014).

1124   47   Chen, L., Guan, L., Qian, P., Xu, F., Wu, Z., Wu, Y., He, K., Gou, X., Li, J. & Hou, S.
1125   (2016) NRPB3, the third largest subunit of RNA polymerase II, is essential for stomatal
1126   patterning and differentiation in Arabidopsis. *Development* **143**, 1600-1611 (2016).

1127   48   Kong, Q., Pattanaik, S., Feller, A., Werkman, J. R., Chai, C., Wang, Y., Grotewold, E. &
1128   Yuan, L. (2012) Regulatory switch enforced by basic helix-loop-helix and ACT-domain
1129   mediated dimerizations of the maize transcription factor R. *Proc Natl Acad Sci U S A*
1130   **109**, E2091-E2097 (2012).

49 Feller, A., Hernandez, J. M. & Grotewold, E. (2006) An ACT-like domain participates in
the dimerization of several plant basic-helix-loop-helix transcription factors. *J Biol Chem*
281, 28964-28974 (2006).
50 Seo, P. J., Hong, S. Y., Kim, S. G. & Park, C. M. (2011) Competitive inhibition of
transcription factors by small interfering peptides. *Trends Plant Sci* **16**, 541-549 (2011).
51 Hornitschek, P., Lorrain, S., Zoete, V., Michielin, O. & Fankhauser, C. (2009) Inhibition of
the shade avoidance response by formation of non-DNA binding bHLH heterodimers.
EMBO J **28**, 3893-3902 (2009).
52 Arai, M. A., Masada, A., Ohtsuka, T., Kageyama, R. & Ishibashi, M. (2009) The first
Hes1 dimer inhibitors from natural products. *Bioorg Med Chem Lett* **19**, 5778-5781
(2009).
53 Arai, M. A., Makita, Y., Yamaguchi, Y., Kawano, H., Suganami, A., Tamura, Y. &
Ishibashi, M. (2020) Total synthesis of lindbladione, a Hes1 dimerization inhibitor and
neural stem cell activator isolated from Lindbladia tubulina. *Sci Rep* **10**, 21433 (2020).
54 Jiao, J., Murakami, K. & Itami, K. (2016) Catalytic methods for aromatic C–H amination:
an ideal strategy for nitrogen-based functional molecules. *ACS Catalysis* **6**, 610-633
(2016).
55 Pillitteri, L. J., Bogenschutz, N. L. & Torii, K. U. (2008) The bHLH protein, MUTE,
controls differentiation of stomata and the hydathode pore in Arabidopsis. *Plant Cell*
Physiol **49**, 934-943 (2008).
56 Nakagawa, T., Nakamura, S., Tanaka, K., Kawamukai, M., Suzuki, T., Nakamura, K.,
Kimura, T. & Ishiguro, S. (2008) Development of R4 gateway binary vectors (R4pGWB)
enabling high-throughput promoter swapping for plant research. *Biosci Biotechnol*
Biochem **72**, 624-629 (2008).
57 Kurihara, D., Matsunaga, S., Uchiyama, S. & Fukui, K. (2008) Live cell imaging reveals
plant aurora kinase has dual roles during mitosis. *Plant Cell Physiol* **49**, 1256-1261
(2008).
58 Susaki, D., Suzuki, T., Maruyama, D., Ueda, M., Higashiyama, T. & Kurihara, D. (2021)
Dynamics of the cell fate specifications during female gametophyte development in
Arabidopsis. *PLoS Biol* **19**, e3001123 (2021).
59 Hajdukiewicz, P., Svab, Z. & Maliga, P. (1994) The small, versatile pPZP family of
Agrobacterium binary vectors for plant transformation. *Plant Mol Biol* **25**, 989-994
(1994).
60 Walter, M., Chaban, C., Schutze, K., Batistic, O., Weckermann, K., Nake, C., Blazevic,
D., Grefen, C., Schumacher, K., Oecking, C., Harter, K. & Kudla, J. (2004) Visualization
of protein interactions in living plant cells using bimolecular fluorescence
complementation. *Plant J* **40**, 428-438 (2004).
61 Sheldrick, G. M. (2015) SHELXT - integrated space-group and crystal-structure
determination. *Acta Crystallogr A Found Adv* **71**, 3-8 (2015).
62 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H.
(2009) OLEX2: a complete structure solution, refinement and analysis program. *J Appl*
Cryst **42**, 339-341 (2009).
63 Case, D. A., Darden, T. A., Cheatham III, T. E., Simmerling, C. L., Wang, J., Duke, R. E.,
Luo, R., Merz, K. M., Wang, B., Pearlman, D. A., Crowley, M., Brozell, S., Tsui, V.,
Gohlke, H., Mongan, J., Hrnak, V., Cui, G., Beroza, P., Schafmeister, C., Caldwell, J. W.,
Ross, W. S. & Kollman, P. A. Amber 9. (2004).
64 Willard, L., Ranjan, A., Zhang, H., Monzavi, H., Boyko, R. F., Sykes, B. D. & Wishart, D.
S. (2003) VADAR: a web server for quantitative evaluation of protein structure quality.
Nucleic Acids Res **31**, 3316-3319 (2003).

65   Dominguez, C., Boelens, R. & Bonvin, A. M. (2003) HADDOCK: a protein-protein
docking approach based on biochemical or biophysical information. ***J Am Chem Soc***
125, 1731-1737 (2003).
66   de Vries, S. J., van Dijk, A. D., Krzeminski, M., van Dijk, M., Thureau, A., Hsu, V.,
Wassenaar, T. & Bonvin, A. M. (2007) HADDOCK versus HADDOCK: new features and
performance of HADDOCK2.0 on the CAPRI targets. ***Proteins*** **69**, 726-733 (2007).
67   Morris, G. M., Huey, R., Lindstrom, W., Sanner, M. F., Belew, R. K., Goodsell, D. S. &
Olson, A. J. (2009) AutoDock4 and AutoDockTools4: Automated docking with selective
receptor flexibility. ***J Comput Chem*** **30**, 2785-2791 (2009).
68   Aihara, Y., Maeda, B., Goto, K., Takahashi, K., Nomoto, M., Toh, S., Ye, W., Toda, Y.,
Uchida, M., Asai, E., Tada, Y., Itami, K., Sato, A., Murakami, K. & Kinoshita, T. (2023)
Identification and improvement of isothiocyanate-based inhibitors on stomatal opening to
act as drought tolerance-conferring agrochemicals. ***Nat Commun*** **14**, 2665 (2023).
69   Toda, Y., Toh, S., Bourdais, G., Robatzek, S., MacLean, D. & Kinoshita, T. (2018)
DeepStomata: Facial Recognition Technology for Automated Stomatal Aperture
Measurement. ***bioRxiv*** p 365098. (2018).
70   Han, S. K., Herrmann, A., Yang, J., Iwasaki, R., Sakamoto, T., Desvoyes, B., Kimura, S.,
Gutierrez, C., Kim, E. D. & Torii, K. U. (2022) Deceleration of the cell cycle underpins a
switch from proliferative to terminal divisions in plant stomatal lineage. ***Dev Cell*** **57**, 569-
582 e566 (2022).
1200