Paradoxical dominant negative activity of an immunodeficiency-associated activating
***PIK3R1* variant**

Patsy R. Tomlinson[1,2], Rachel Knox[1,2,3], Olga Perisic[4], Helen C. Su[5], Gemma V. Brierley[6,1],
Roger L. Williams[4], Robert K. Semple[1,7,8,3]

[1] The University of Cambridge Metabolic Research Laboratories, Wellcome Trust-MRC
Institute of Metabolic Science, Cambridge, UK
[2] MRC Metabolic Diseases Unit, Wellcome Trust-MRC Institute of Metabolic Science,
Cambridge, UK
[3] The National Institute for Health Research Cambridge Biomedical Research Centre,
Cambridge, UK
[4] MRC Laboratory of Molecular Biology, Cambridge, UK
[5] Laboratory of Clinical Immunology & Microbiology, Intramural Research Program, National
Institute of Allergy and Infectious Disease, National Institutes of Health, Bethesda, MD,
USA
[6] Department of Comparative Biomedical Science, The Royal Veterinary College, London
NW1 0TU, UK
[7] Centre for Cardiovascular Science, University of Edinburgh, Edinburgh, UK
[8] MRC Human Genetics Unit, Institute of Genetics and Cancer, University of Edinburgh,
Edinburgh, UK

Correspondence to:
Prof. Robert K. Semple, Centre for Cardiovascular Science, University of Edinburgh,
Queen's Medical Research Institute, Little France Crescent, Edinburgh EH16 4TJ, UK. Tel:
+44 131 242 6051/ Email: rsemple@ed.ac.uk

Character Count: 50,222 (to end of references; with spaces)

Running Title: Dominant negative effect of an activating PIK3R1 variant

Abstract

PIK3R1 encodes three regulatory subunits of class IA phosphoinositide 3-kinase (PI3K), each associating with any of three catalytic subunits, namely p110α, p110β or p110δ. Constitutional *PIK3R1* mutations cause diseases with a genotype-phenotype relationship not yet fully explained: heterozygous loss-of-function mutations cause SHORT syndrome, featuring insulin resistance and short stature attributed to reduced p110α function, while heterozygous activating mutations cause immunodeficiency, attributed to p110δ activation and known as APDS2. Surprisingly, APDS2 patients do not show features of p110α hyperactivation, but do commonly have SHORT syndrome-like features, suggesting p110α hypofunction. We sought to investigate this. In dermal fibroblasts from an APDS2 patient, we found no increased PI3K signalling, with p110δ expression markedly reduced. In preadipocytes, the APDS2 variant was potently dominant negative, associating with Irs1 and Irs2 but failing to heterodimerise with p110α. This attenuation of p110α signalling by a p110δ-activating PIK3R1 variant potentially explains co-incidence of gain-of-function and loss-of-function *PIK3R1* phenotypes.

Keywords: APDS2, cancer, immunodeficiency, insulin resistance, phosphoinositide 3-kinase, PIK3R1, SHORT syndrome

Introduction

PIK3R1 gene products (p85α, p55α and p50α) are essential for class IA phosphoinositide 3-kinase (PI3K) signalling. Each of the three protein products can bind any of three catalytic subunits, p110α, β and δ, encoded by *PIK3CA*, *PIK3CB*, and *PIK3CD* respectively. This stabilises and inhibits the catalytic subunits in the basal state and confers responsiveness to upstream stimuli including receptor tyrosine kinase activation, enabled by two phosphotyrosine-binding SH2 domains shared by all PIK3R1 products (Fruman *et al*, 2017).

PI3K signalling mediates responses to a myriad of cues, including growth factors, hormones such as insulin, and processed antigens, and so recent discovery that genetic perturbation of PIK3R1 in humans disrupts growth, insulin action and immunity is no surprise. Beyond this headline observation lies considerable complexity, however, and important questions about genotype-phenotype correlations in PIK3R1-related disorders are unresolved.

Some PIK3R1 mutations reduce basal inhibition of catalytic subunits, usually due to disruption of the inhibitory inter-SH2 domain, and are found in cancers (Philp *et al*, 2001) and vascular malformations with overgrowth(Cottrell *et al*, 2021). In both diseases, hyperactivated PI3Kα, composed of heterodimers of *PIK3R1* products and p110α, drives pathological growth. Distinct inter-SH2 domain *PIK3R1* mutations, mostly causing skipping of exon 11 and deletion of residues 434-475, hyperactivate PI3Kδ in immune cells, causing highly penetrant monogenic immunodeficiency (Deau *et al*, 2014; Lucas *et al*, 2014b). This phenocopies the immunodeficiency caused by genetic activation of p110δ itself, which is named Activated PI3K Delta Syndrome 1 (APDS1) (Angulo *et al*, 2013; Lucas *et al*, 2014a). The PIK3R1-related syndrome, discovered shortly afterwards, is thus named APDS2.

Despite ubiquitous *PIK3R1* expression, features attributable to PI3Kα activation have been described in only a single case of APDS2 to date (Wentink *et al*, 2017). In that case, the causal variant (N564K) was a constitutional activating mutation associated with cancer rather than one of the common APDS2 variants. Biochemical studies have suggested that

apparently selective p110δ activation occurs because APDS2 mutations derepress p110δ,

the predominant immune system catalytic subunit, more than p110α, due to differences in

the inhibitory contacts between *PIK3R1* products and different catalytic subunits (Dornan *et*

al, 2017).

More surprisingly, phenotypic overlap is reported between APDS2 and SHORT syndrome.

SHORT syndrome, named for the characteristic developmental features (S̲hort stature,

H̲yperextensibility, H̲ernia, O̲cular depression, R̲ieger anomaly, and T̲eething delay) is

caused by loss of PI3Kα function due to disruption of the phosphotyrosine-binding C-

terminal SH2 domain (Chudasama *et al*, 2013; Dyment *et al*, 2013; Thauvin-Robinet *et al*,

2013). Like APDS2 it is stereotyped and highly penetrant. It features short stature, insulin

resistance, and dysmorphic features (Avila *et al*, 2016). In recent years, both individual case

reports (Bravo Garcia-Morato *et al*, 2017; Petrovski *et al*, 2016; Ramirez *et al*, 2020;

Szczawinska-Poplonyk *et al*, 2022) and larger case series (Elkaim *et al*, 2016; Jamee *et al*,

2020; Maccari *et al*, 2023; Nguyen *et al*, 2023; Olbrich *et al*, 2016; Petrovski *et al*., 2016)

have established that many people with APDS2 have overt features of SHORT syndrome,

while, more generally, linear growth impairment is common in APDS2, but not in APDS1.

These clinical observations are bolstered by the impaired linear growth and increased *in*

utero mortality reported in mice with knock in of the common causal APDS2 mutation in

Pik3r1(Nguyen *et al*., 2023). All people with SHORT-APDS2 overlap syndromes have well

established activating PIK3R1 mutations in the inter-SH2 domain implicated in APDS2, but

none have characteristic SHORT syndrome mutations, which are usually in the C-terminal

SH2 domain. Conversely, no patient with a classical SHORT syndrome mutation has been

shown to have immunodeficiency. There is thus now convincing evidence of a syndrome

with features of both gain and loss of PI3K function. The mechanistic basis of this is

unexplained.

PI3K activity is determined not just by activity of individual PI3K holoenzymes, but also by

subunit stoichiometry. Regulatory and catalytic subunits stabilise each other upon binding

(Brachmann *et al*, 2005; Yu *et al*, 1998), and only regulatory subunit monomers are stable

enough to be detected in mammalian cells (Yu *et al.,* 1998). Molar excess of regulatory

subunits over catalytic subunits gives rise to free regulatory subunits that compete with

holoenzyme for phosphotyrosine binding (Ueki *et al*, 2002). This has been invoked to explain

increased insulin sensitivity on knockout or reduction of p85α alone (Mauvais-Jarvis *et al*,

2002)*,* of p55α and p50α (Chen *et al*, 2004) or of all *Pik3r1* products (Fruman *et al*, 2000).

The "free-p85" model has been contested by some reports suggesting no excess p85

(Geering *et al*, 2007; Zhao *et al*, 2006), however most observations, including some made

recently using *in vivo* tagging and pulldown of PI3K components (Tsolakos *et al*, 2018)

suggest that monomeric p85α can be seen *in vivo* in some tissues.

We now address mechanisms underlying human mixed gain- and loss-of function

phenotypes observed in association with PIK3R1 mutations using primary cells, cell models

and in vitro enzyme assay and suggest that they are explained by competing effects of

APDS2 PIK3R1 mutations on PI3K activity and stability.

Results

PI3K subunit expression and signalling in APDS2 fibroblasts

Disorders in the *PIK3CA*-related overgrowth spectrum (PROS) feature soft tissue

overgrowth and are caused by heterozygous, mosaic activating mutations in *PIK3CA*,

encoding the p110α catalytic subunit of PI3Kα(Madsen *et al*, 2018). In this group, basal

hyperactivation of PI3Kα signalling can be easily discerned in dermal fibroblasts from

affected areas of the body(Lindhurst *et al*, 2012). *PIK3R1*, like *PIK3CA*, is ubiquitously

expressed, yet overgrowth is not reported in APDS2 caused by heterozygous, constitutional

activating mutations in *PIK3R1*. Dermal fibroblasts strongly express *PIK3R1*, but no studies

to date have evaluated whether PI3K activity is increased in APDS2.

We began by assessing dermal fibroblasts cultured from a previously described woman with

APDS2 due to the common causal PIK3R1 mutation. This affects a splice donor site that

causes skipping of exon 11, leading to in-frame deletion of 42 amino acids (434-475

inclusive) in the inter-SH2 domain, which is shared by all PIK3R1 isoforms (Patient A.1 in

(Lucas *et al.*, 2014b))(**Fig 1 figure supplement 1**). These cells were compared to cells from

healthy control volunteers, or from people with PROS. Confirmation of expression of all

pathogenic mutations was undertaken by cDNA sequencing prior to further study (**Fig 1**

figure supplement 2A). We found that truncated p85α was expressed, but at a much lower

protein level than full length wild-type p85α (**Fig 1A, Fig 1 figure supplement 2B,C**).

Despite this, no increase in basal or insulin-stimulated AKT phosphorylation was seen in

APDS2 cells compared to cells from wild-type volunteers or from people with PROS and

activating *PIK3CA* mutations H1047L or H1047R (**Fig 1A-C, Fig 1 figure supplement**

3A,B). Although insulin-induced AKT phosphorylation was lower in fibroblasts from the one

APDS2 patient studied compared to controls, we have previously reported extensive

variability in insulin-responsiveness of primary dermal fibroblasts. Moreover even primary

cells from a patient expressing high levels of the SHORT syndrome-associated p85α Y657X

did not show attenuated insulin action. The reduced insulin action in APDS2 cells in the

current study thus should not be overinterpreted until reproduced in further APDS2 cells.

Previous studies have suggested that the truncated, APDS2-causal p85α variant exerts a

much greater activating effect on p110δ, which has a more restricted tissue expression,

including immune cells, than p110α, which is ubiquitous (Dornan *et al.*, 2017). We found that

both p110α and p110δ were easily detectable in control dermal fibroblasts, however p110δ

was almost absent in APDS2 fibroblasts compared to controls, with lower levels also seen in

PROS cells (**Fig 1A,D, Fig 1 figure supplement 3D**). p110α was unchanged in APDS2

cells but modestly increased in PROS cells (**Fig 1A,E, Fig 1 figure supplement 3C**).

Collectively these findings suggest that any ability of the APDS2 PIK3R1 variant in skin cells

to activate PI3K may be overcome by reduced protein levels of p110δ, likely through

reduced binding and/or reduced stabilisation of p110δ by the mutant regulatory subunit.

Overexpressed PIK3R1 ΔExon11 is potently dominant negative in 3T3-L1 preadipocytes

We next turned to a well-established cellular system allowing conditional overexpression of

PK3R1 alleles of interest. We have previously shown that overexpression of two SHORT

syndrome PIK3R1 variants – R649W and Y657X – impair insulin signalling and adipocyte

differentiation of murine 3T3-L1 preadipocytes, consistent with impaired PI3Kα function and

with the lipodystrophy and insulin resistance seen in SHORT syndrome (Huang-Doran *et al*,

2016). To assess the effect of the APDS2 ΔEx11 in the same, non-immunological, cell

context, we used lentiviral vectors, as previously described (Huang-Doran *et al.*, 2016), to

generate clonal 3T3-L1 cell lines allowing conditional, tuneable overexpression of PIK3R1

variants in response to doxycycline (**Figure 2 figure supplement 1)**. For signalling studies,

we also generated cells conditionally overexpressing the PROS- and cancer-associated

PIK3CA H1047R mutation as a positive control for increased PI3Kα signalling, while for

differentiation studies we used previously reported lines conditionally expressing PIK3R1

R649W or Y657X (Huang-Doran *et al.*, 2016) (**Fig 1 figure supplement 1**). In

undifferentiated cells, we first confirmed doxycycline-dependent overexpression of p85α or

p110α transgenes (**Fig 2A**), before assessing basal and insulin-stimulated Akt

phosphorylation. As expected, overexpression of oncogenic H1047R p110α strongly

increased basal Akt phosphorylation (**Fig 2A-C, Fig 2 figure supplement 1A,C**) with no

additional increase on insulin stimulation (**Fig 2A-C, Fig 2 figure supplement 1B,D**).

Surprisingly, however, not only did overexpression of the APDS2 ΔEx11 p85α not increase

basal PI3Kα signalling, but it potently inhibited insulin-induced Akt phosphorylation,

consistent with a strong dominant negative action on pathway activation (**Fig 2A-C, Fig 2**

figure supplement 1). Overexpressing wild-type p85α mildly reduced insulin-stimulated Akt

phosphorylation (**Fig 2A-C, Fig 2 figure supplement 1B,C**), although WT p85α was not

expressed to as high a level as ΔEx11 p85α. In keeping with impaired PI3Kα activity,

overexpression of the ΔEx11 variant also severely impaired adipocyte differentiation,

assessed by triglyceride accumulation in response to a standard differentiation protocol (**Fig**

2D). A similar effect was seen on overexpressing SHORT syndrome variants R649W and

Y657X (**Fig 2D**).

To assess whether the inhibitory effect of ΔEx11 p85α might be an artefact caused by strong

overexpression, doxycycline titration was used to assess whether a low level of

overexpression might unmask signalling hyperactivation. However, no such hyperactivation

was seen, and instead, a graded diminution of insulin-induced AKT phosphorylation was

observed from the lowest to highest levels of p85α ΔEx11 overexpression (**Fig 2 figure**

supplement 3).

Effect of PIK3R1 mutations on Phosphoinositide 3-kinase activity in vitro

Given this evidence that APDS2-associated PIK3R1 ΔEx11 potently inhibits PI3Kα signalling

when overexpressed in 3T3-L1 preadipocytes, we next sought to investigate the biochemical

basis of this observation. First, we assessed the effect of disease-causing PIK3R1 mutations

on basal and phosphotyrosine-stimulated activity of purified PI3Kα, β and δ holoenzyme in a

previously described reconstituted *in vitro* system(Burke *et al*, 2012). As well as the APDS2

ΔEx11 mutation, we selected three SHORT syndrome-associated mutations to study. These

were the most common causal mutation, R649W, which abolishes phosphotyrosine binding

by the C-terminal SH2 (cSH2) domain (Chudasama *et al*., 2013), Y657X, which truncates

the cSH2 domain (Huang-Doran *et al*., 2016; Kwok *et al*, 2020), and E489K which,

atypically, lies in the inter-SH2 domain where most cancer, overgrowth and APDS2-

associated mutations lie (Thauvin-Robinet *et al*., 2013). PIK3R1 E489K-containing primary

cells were previously suggested to show basal hyperactivation (Thauvin-Robinet *et al*.,

2013).

Wild-type and SHORT syndrome mutant holoenzymes were successfully purified for *in vitro*

assay, but despite multiple attempts, ΔEx11 holoenzyme could only be made in small

amounts under identical conditions, and moreover was unstable on storage, precluding

further study. Such instability of *in vitro* synthesised ΔEx11 holoenzyme was previously

reported (Dornan *et al.*, 2017). Also in keeping with previous reports (Chudasama *et al.*,

2013; Dornan *et al.*, 2017; Dornan *et al*, 2020), p85α R649W showed severely reduced

phosphotyrosine-stimulated activity in complex with p110α, with highly significant but lesser

loss of function seen for Y657X and E489K (**Fig 3A, Fig 3 figure supplement 1A**). No

increase in basal activity was seen for any variant.

We also assessed whether any of the SHORT syndrome variants affect function of PI3Kβ

and found that all variants impaired phosphotyrosine-stimulated activity of PI3Kβ. Again, this

impairment was less for p85α Y657X than for R649W, and in this case only very mild for

E489K (**Fig 3B, Fig 3 figure supplement 1B**). For PI3Kδ, p85α R649W conferred severe

loss of function, as for other isoforms (**Fig 3C, Fig 3 figure supplement 1C**), suggesting

that the absence of immunodeficiency in SHORT syndrome is not accounted for by selective

inhibition of PI3Kα function by causal mutations.

Binding of p110α by mutant p85α

Given the potent dominant negativity of APDS2-related p85α ΔEx11 in cells, and the

instability of p85α ΔEx11-containing PI3K holoenzyme *in vitro*, we next used

immunoprecipitation to assess binding of p110α by this and other p85α variants in cells.

p85α was easily detected in anti-p110α immunoprecipitates from all cell lines at baseline

(**Fig 4**). No increase in p110α levels was seen on conditional overexpression of wild-type or

R649W p85α. Given the known instability of monomeric p110α, this suggests that all p110α

is bound to p85α before overexpression. Although wild-type endogenous p85α may have

been replaced by heterologously overexpressed p85α in these cells, this could not be

detected without a size difference of the variant from wild-type. For cells overexpressing the

SHORT syndrome-associated p85α Y657X, the truncated variant was strongly co-

immunoprecipitated, accounting for nearly all of the p85α signal in anti-p110α

immunoprecipitate. This demonstrates preserved binding of p110α by mutant p85α (**Fig 4**).

In sharp contrast, although truncated p85α ΔEx11 was easily detected in cell lysates before

immunoprecipitation and in supernatant after immunoprecipitation (arrowed in **Fig 4**), no

truncated p85α ΔEx11 was seen in p110α immunoprecipitates, and no change in p110α

expression was detected (**Fig 4**). This suggests that this truncated APDS2 causal variant

does not supplant endogenous, full-length p85α binding to p110α, despite overexpression.

This argues against destabilisation of p110α as the mechanism of the observed dominant

negative activity.

Effect of PIK3R1 mutations on insulin-induced PI3K recruitment to IRS1/2

As APDS2 p85α ΔEx11 does not appear to displace wild-type p85α from p110α, despite

strong overexpression, it is likely that there are high levels of truncated p85α unbound to

p110α in the cell. This may be important, as p85α mediates recruitment of PI3K to activated

tyrosine kinase receptors and their tyrosine phosphorylated substrates, including the insulin-

receptor substrate proteins Irs1 (Myers *et al*, 1992) and Irs2(Sun *et al*, 1995). Excess free

regulatory subunits compete with heterodimeric PI3K holoenzyme for binding to these

phosphotyrosines (Ueki *et al.*, 2002), raising the possibility that excess free, truncated

APDS2 p85α ΔEx11 may exert its inhibitory action similarly by outcompeting PI3K

holoenzyme for phosphotyrosine binding.

To assess this possibility, we again used the 3T3-L1 cellular model to determine whether

overexpression of disease-causing p85α variants impairs recruitment of p110α to Irs1 and

Irs2. Irs1 was immunoprecipitated with or without conditional p85α overexpression and with

or without insulin stimulation. Overexpression of wild-type p85α had no effect on basal or

insulin-induced p110α recruitment to Irs1 (**Fig 5A,B, Fig 5 supplement 1A,B**). In contrast,

overexpression of either p85α R649W or Y657X sharply reduced insulin-stimulated p110α

recruitment to Irs1 on insulin stimulation (**Fig 5A,B, Fig 5 supplement 1A,B**). In keeping

with the inability of the R649W cSH2 domain to bind phosphotyrosines, p85α R649W was

also not recruited to Irs1, while overexpression of Y657X increased stimulated but not basal

p85α recruitment (**Fig 5A,C, Fig 5 supplement 1C,D**). This suggests a pure defect in PI3K

holoenzyme recruitment for R649W. For Y657X the signalling defect may have mixed

mechanisms, with reduced activation by phosphotyrosine seen in *in vitro* studies coupled to

increased abundance of monomeric mutant p85α, leading to recruitment of non p110α-

bound mutant p85α to Irs1.

In keeping with our finding of severely attenuated insulin signalling upon p85α ΔEx11

overexpression (**Fig 2**), overexpression of this mutant p85α abolished p110α recruitment to

Irs1 (**Fig 5A,B, Fig 5 supplement 1A,B**). However, non p110α-bound p85α ΔEx11 was

strongly recruited to Irs1 even in the absence of insulin stimulation (**Fig 5A,C, Fig 5**

supplement 1C,D). This suggests that although p85α ΔEx11 does not effectively compete

with wild-type p85α for binding to p110α, it has preserved or possibly enhanced ability to

bind Irs1. This gives the mutant p85α properties that render it a more potent endogenous

inhibitor of PI3K signalling than free wild-type p85α. Experiments conducted with

immunoprecipitation of Irs2 instead of Irs1 yielded closely similar findings for all p85α

species (**Fig 6 and Fig 6 supplement 1**).

Discussion

Murine genetic studies of Pik3r1 have proved complicated to interpret, due in part to

functional redundancy among PI3K regulatory subunits, and in part to unanticipated effects

of perturbing PI3K subunit stoichiometry. This has left several important questions about *in*

vivo functions of Pik3r1 unresolved. Identification of a series of human genetic disorders

caused by constitutional PIK3R1 mutations over the past 10 years has given fresh impetus

to the field and has been mechanistically illuminating.

Homozygous truncating PIK3R1 mutations abolishing p85α expression while preserving

p55α and p50α produce agammaglobulinaemia (Conley *et al*, 2012; Tang *et al*, 2018)

(**Figure 1 figure supplement 1**). This resembles the immunodeficiency reported in *Pik3r1*

knockout mice (Fruman *et al*, 1999; Suzuki *et al*, 1999) and suggests an essential, non-

redundant function of p85α in B cell development in humans. Thereafter, human genetics

has provided more novel insights. *PIK3R1* mutations were identified in SHORT syndrome in

2013 (Chudasama *et al.*, 2013; Dyment *et al.*, 2013; Thauvin-Robinet *et al.*, 2013), nearly all

in the C-terminal SH2 domain, which together with the N-terminal SH2 domain enables PI3K

recruitment to activated RTKs (Rordorf-Nikolic *et al*, 1995). SHORT syndrome features short

stature and insulin resistance consistent with impaired ligand-induced p110α action, a

phenotype distinct from the enhanced insulin sensitivity produced by genetic ablation of one

or more *Pik3r1* products in mice (Chen *et al.*, 2004; Fruman *et al.*, 2000; Mauvais-Jarvis *et*

al., 2002). Mice with SHORT syndrome mutations knocked in were generated after human

findings, and faithfully reproduce the human phenotype (Kwok *et al.*, 2020; Solheim *et al*,

2018; Winnay *et al*, 2016), confirming that expression of a signalling-impaired PIK3R1 has

different consequences to Pik3r1 knockout. No immunodeficiency has been described

associated with classic SHORT syndrome mutations, however, suggesting a selective effect

on PI3Kα, but the basis of this selectivity has not previously been investigated.

In contrast to SHORT syndrome, mutations in the inter SH2 domain of PIK3R1, mostly

leading to skipping of exon 11, were shown in 2014 to activate PI3K *in vitro* and to cause

immunodeficiency (APDS2) (Deau *et al.*, 2014; Lucas *et al.*, 2014b) similar to that caused by

activating mutations in p110δ (APDS1) (Angulo *et al.*, 2013; Lucas *et al.*, 2014a). Neither

overgrowth nor metabolic features of APDS2 have been described to suggest p110α

hyperactivation, and indeed short stature is common in APDS2 (Elkaim *et al.*, 2016; Jamee

et al., 2020; Maccari *et al.*, 2023; Olbrich *et al.*, 2016; Petrovski *et al.*, 2016), with a growing

number of APDS2 patients described with features of SHORT syndrome (Bravo Garcia-

Morato *et al.*, 2017; Maccari *et al.*, 2023; Nguyen *et al.*, 2023; Petrovski *et al.*, 2016;

Ramirez *et al.*, 2020; Szczawinska-Poplonyk *et al.*, 2022). Moreover mice with the common

APDS2 causal Pik3r1 variant knocked in show impaired growth and *in utero* survival, unlike

APDS2 murine models(Nguyen *et al.*, 2023).

Thus, study of distinct PIK3R1-related syndromes shows that established loss-of-function

PIK3R1 mutations produce phenotypes attributable to selectively impaired PI3Kα

hypofunction, while activating mutations produce phenotypes attributable to selectively

increased PI3Kδ signalling. Indeed, not only do such activating mutations not produce

phenotypes attributable to PI3Kα activation, but they surprisingly have features

characteristic of impaired PI3Kα function.

Lack of overgrowth in APDS2 has been attributed to greater ability of APDS2 PIK3R1

variants to activate p110δ than p110α (Dornan *et al.*, 2017)*,* but the co-occurence of gain-

and loss- of function phenotypes has not been explained to date. Our findings suggest that

the explanation may lie in competing effects of APDS2 PIK3R1 variants to activate PI3Kδ on

one hand, as previously shown (Angulo *et al.*, 2013; Dornan *et al.*, 2017; Lucas *et al.*,

2014b), while on the other hand interfering with PI3Kα through the dominant negative effect

of non p110α-bound mutant p85α.

The low expression of truncated p85α ΔEx11 we described in dermal fibroblasts is similar to

observations made in lymphocytes, however, in lymphocytes this is associated with

increased basal AKT phosphorylation (Deau *et al.*, 2014; Lucas *et al.*, 2014b) that is

abolished by p110δ inhibition (Lucas *et al.*, 2014b). P110δ is also expressed in fibroblasts,

but protein levels were reduced in the cells studied, consistent with the previously reported

observation that when expressed in insect cells the PI3K holoenzyme containing

p85α ΔEx11 has a low yield and is unstable (Dornan *et al.*, 2017). We speculate that in cells

with low endogenous p110δ protein expression, the destabilising effect of mutant PIK3R1

predominates over its activating effect.

The balance between expression and signalling in different cells may be a fine one,

however, as transient overexpression of FLAG-tagged p85α ΔEx11 did increase AKT

phosphorylation in 3T3 fibroblasts in a previous study, although expression of p110α, β and

δ was not determined (Deau *et al.*, 2014). We find, in contrast, that overexpression of

untagged p85α ΔEx11 has a strong inhibitory effect on insulin signalling, and we were

unable to identify a window of overexpression where increased AKT phosphorylation could

be observed. We further demonstrated that most or all of the mutant p85α expressed is not

bound to p110α, while unbound mutant p85α still binds to Irs1/2, effectively competing with

PI3K holoenzyme. The observation that p85α ΔEx11 can associate with Irs1/2 is in

agreement with reports that p85α ΔEx11-containing PI3Kα and PI3Kδ can be stimulated by

pY2-peptides (Dornan *et al.*, 2017; Dornan *et al.*, 2020) and that p85α ΔEx11 is recruited to

tyrosine-phosphorylated LAT in T cells(Lucas *et al.*, 2014b). The competition we suggest

between unbound mutant p85α and PI3K holoenzyme for binding to the activated RTKs is in

keeping with longstanding evidence that free p85 downregulates PI3K signalling through the

same competitive mechanism (Thorpe *et al*, 2017; Ueki *et al.*, 2002), with a single study

suggesting in addition that overexpression of tagged p85α leads to Irs1 sequestration with

free p85α in cytosolic foci where PI(3,4,5)P3 production does not occur (Luo *et al*, 2005).

The current study has limitations. We have studied primary cells from only a single APDS2

patient, and in the 3T3-L1 cell model, we did not determine whether p110δ protein could be

detected. If not, this could explain the lack of detectable AKT phosphorylation with induction

of Pik3r1 ΔEx11. Indeed, previous pharmacological studies in 3T3-L1 adipocytes has shown

that selective inhibition of p110δ or p110β does not alter insulin-induced phosphorylation of

any protein studied in the PI3-K pathway, attesting to the dominance of p110α in insulin

action in this cell model (Knight *et al*, 2006). Our study moreover raises further questions.

Full length p85α subunits can homodimerize (Cheung *et al*, 2015; Harpur *et al*, 1999;

LoPiccolo *et al*, 2015), and it is speculated that homodimers may outcompete p85α/p110

heterodimers for binding to activated Irs1 due to configuration of the four SH2 domains. In

the 3T3-L1 preadipocyte APDS2 models, p85α ΔEx11 expression was high despite impaired

p110α association, but whether this is composed of monomeric or homodimeric p85α, and

whether mutant p85α expression leads to sequestration of Irs1 remote from the insulin

receptor, as previously suggested for tagged wild-type p85α(Luo *et al.*, 2005), is

undetermined.

In summary, it is already established that: **A.** genetic activation of PIK3CD causes

immunodeficiency without disordered growth, while **B.** inhibition of PIK3R1 recruitment to

RTKs and their substrates impairs growth and insulin action, without immunodeficiency,

despite all catalytic subunits being affected and **C.** loss of p85α alone causes

immunodeficiency. The current study, coupled with prior reports, suggests that the common

APDS2 mutation in PIK3R1 has mixed consequences, producing greater hyperactivation of

p110δ than p110α, based on subtle differences in the inhibitory interactions of regulatory

and catalytic subunits, while also destabilising PI3K holoenzyme and exerting dominant

negative activity on wild-type PIK3R1 function. We suggest that these competing activating

and inhibitory consequences are finely balanced, potentially differing among tissues, leading

to mixed clinical profiles of gain- and loss-of function features.

Materials and Methods

Ethics

The patient studied gave written informed consent for NIH IRB-approved research protocols

06-I-0015 and 09-I-0133 and was reported previously(Lucas *et al.*, 2014b). Dermal

fibroblasts were isolated from a skin punch biopsy and cultured as previously

described(Huang-Doran *et al.*, 2016).

Baculovirus Generation

p85α point mutation expression constructs were created by site-directed mutagenesis of a

Hsp85α_pACEBac1 plasmid using a QuikChange II XL Site-Directed Mutagenesis Kit

(Agilent, 200521). In-Fusion cloning (Takara, 638909) was used to generate

Hsp85α_pACEBac1_p85α ΔEx11. In brief, Hsp85α_pACEBac1 backbone was digested

(BamHI/NotI) and purified using Zymoclean Gel DNA Recovery Kit (Zymo Research,

D4001), while the cDNA insert was purified using NucleoSpin® Clean-up Kit (Takara,

740609.10). Insert generation was performed by High-Fidelity PCR using primers designed

to yield fragments containing exons 1-10 or 12-16 with the necessary overlap, with inserts

verified by electrophoresis and purified using NucleoSpin® Clean-up kit. In-Fusion reactions

were performed according to manufacturer's guideline using 100ng each of linearised

plasmid and both inserts in 10µL. Products were transformed into Stellar Competent cells

(Takara, 636763). Purified plasmid inserts were sequenced and verified by restriction

enzyme digest, exploiting loss of a DpnI site within excised exon 11.

Hsp85α_pACEBac1,   and   Hsp110α_pACEBac1,   Hsp110β_pACEBac1   and

Hsp110δ_pFastBac™HT B plasmids encoding N terminally tandem His-tagged p110

subunits were used to generate bacmid DNA. MAX Efficiency DH10Bac™ Competent

(Invitrogen, 10361-012) or EMBacY cells (Thermofisher, 10361012) were transformed with

40ng plasmid DNA and turbid cultures plated onto agar containing 50µg/mL kanamycin,

10µg/mL gentamicin, 10µg/mL tetracycline, 100µg/mL X-Gal and 40µg/mL IPTG. Single

colonies were picked, expanded and purified at 2 days and bacmid concentrations quantified

by Nanodrop 1000 Spectrophotometer (Thermo Scientific). 2-4µg bacmid was transfected

into Sf9 cells in 6 well plates using Insect-XPRESS (Promega E2311). Cells were incubated

at 2°C for 5 days and bacmid YFP expression assessed by Leica DM IL LED Fluo

microscope using a green fluorescent protein filter cube. Pooled virus from 2 wells of

supernatant (P1 stock) was used to produce high titre P2 stock by transfection of 1.5x106

Sf9 cells/mL (Thermofisher, 11496015) in 450mL using Insect-XPRESS with L-Glutamine

(Lonza LZBE12-730Q). For catalytic subunits, 1-2mL P2 stock was used for a further round

of Sf9 transfection and expansion to generate P3 stock.

Purification of PI3K holoenzymes

1.5-2L of 1.5x106 Sf9 cells/mL were co-infected with 18mL P3 catalytic subunit and 4mL P2

regulatory subunit baculovirus. Non infected Sf9 cells were negative controls, and a prior

protein preparation of 320 kDa was the positive control. Cells were cultured at 27°C,

harvested 48h post-infection, pelleted, washed and stored at -80°C. Pellets were later lysed

by sonication in buffer containing 20 mM TrisHCl (pH 8.0), 300 mM NaCl, 20 mM imidazole

(pH 8.0) and 1 mM TCEP (pH 7.0) at 4 °C. Universal Nuclease (Thermofisher, 88702) was

added to lysates before ultracentrifugation at 35,000 rpm for 35min at 4°C. p85α/p110

heterodimers were pulled down via 6 tandem p110 N-terminal His tags, preventing

purification of monomeric p85α, using tandem Ni2+HisTrap Fast Flow columns (GE

Healthcare) (equilibration buffer 20 mM TrisHCl (pH 8.0), 100 mM NaCl, 20 mM imidazole

and 1 mM TCEP (pH 7.0); elution buffer 20 mM TrisHCl (pH 8.0), 100 mM NaCl, 200 mM

imidazole and 1 mM TCEP (pH 7.0)). Further purification utilised a heparin HiTrap Q HP

column (GE Healthcare) equilibrated with 20mM TrisHCl (pH 8.0) and 1mM TCEP (pH 7.0),

with proteins eluted in 20mM TrisHCl (pH 8.0), 1 M NaCl and 1mM TCEP (pH 7.0). Eluted

fractions were concentrated to ≥1mg/mL using Millipore Amicon Ultra-15 Centrifugal Filter

Units with Ultracel-50 membrane, and 1mL concentrated fractions were gel filtered on

Superdex 200 16/60 columns (GE Healthcare) equilibrated in 20mM HEPES (pH 7.5),

100mM NaCl and 2mM TCEP (pH 7.0). The p110 6-His tag was retained for proteins used in

functional analyses. KTA Protein Purification Systems (GE Healthcare) and UNICORN

Control Software version 5.11 (GE Healthcare) were used for all purifications. Purity of

eluted complexes was verified by SDS-PAGE, and purified proteins were quantified using a

Nanodrop 1000 Spectrophotometer (Thermofisher). Protein concentration was calculated

using the molecular extinction coefficient (assuming full cysteine reduction) determined by

heterodimer sequence input to the ProtParam tool (ExPASy). Purified proteins were stored

at −80°C in single-use aliquots.

Total injection volume onto gel filtration columns was kept at 1 mL. Yields of PI3K

complexes were determined using the area under the curve (280nm mAU per mL eluted

protein) from gel filtration chromatograms, and normalised to the volume of Sf9 cells

infected.

Liposome preparation

Chloroform/methanol solutions of sphingomyelin, cholesterol, porcine brain

phosphatidylcholine, porcine brain phosphatidylethanolamine, porcine brain

phosphatidylserine and porcine brain PI(4,5)P2 (Avanti Polar Lipids) were mixed to generate

a preparation containing 5/10/15/45/20/5% (wt/vol) of each lipid, with a total lipid

concentration of 5 mg/mL and final PI(4,5)P2 concentration 250 µg/mL. Lipid preparations

were dessicated under argon and then in a vacuum desiccator. Lipids were rehydrated in

buffer containing 20mM HEPES (pH 7.5), 100 mM KCl and 1 mM EGTA (pH 8.0) using

vortexing for 3 min, waterbath sonication for 15 min, and 10 cycles between liquid nitrogen

and a 43°C water bath. Unilamellar vesicles were generated by extrusion through

polycarbonate filters with 100nm pores, using a glass-tight syringe. Single use aliquots were

stored at -80 °C.

Fluorescence Polarisation Assay

PI(3,4,5)P3 production was measured by modified PI3-Kinase activity fluorescence

polarisation assay (Echelon Biosciences, Salt Lake City, UT, USA). 10µL reactions in 384-

well black microtitre plates used 1mM liposomes containing 50µM PI(4,5)P2, optimised

concentrations of purified PI3K proteins, 100µM ATP, 2mM MgCl2, with or without 1µM

tyrosine bisphosphorylated 33-mer peptide derived from mouse PDGFRβ residues 735-767,

including phosphotyrosine at positions 740 and 751 ("pY2"; 735-

ESDGG**Y**MDMSKDESID**Y**VPMLDMKGDIKYADIE-767; Cambridge peptides). Reactions

were quenched with 5µL of solution containing 20mM HEPES (pH 7.5), 150mM NaCl, 30mM

EDTA (pH 7.4) and 400nM GST–Grp1-PH, followed by addition of 5µL 40nM TAMRA Red

Fluorescent Probe in HNT buffer. Identity of the lipid group coupled to the TAMRA probe

was not disclosed by Echelon Biosciences, but as Grp1 recognises specific lipid head

groups in competition with PI(3,4,5)P3, it is likely to be a lipid with a similar head group such

as inositol 1,3,4,5-tetrakisphosphate. After 1h equilibration, fluorescence polarisation was

measured using a PHERAstar spectrofluorometer (BMG Labtech, Ortenberg, Germany) with

the FP/540-20/590-20/590-20 optical module. The concentrations of wild-type protein

complexes used in the assay were: 20-40nM p85α/p110α, 80-160nM p85α/p110β and 80-

160nM p85α/p110δ for basal activities, and 1-2nM p85α/p110α, 2-4 nM p85α/p110β and 2-

6nM p85α/p110δ for pY2-stimulated activities. Concentrations of thawed protein aliquots

were assessed as 280 nm absorbance by Nanodrop 1000 Spectrophotometer, and samples

of preparations used in assays were resolved by SDS-PAGE to confirm equal amounts of

each complex. Three technical replicates were used per assay on at least three different

occasions. Relevant wild-type p85α/p110 catalytic isoforms were included in each assay as

controls. Other controls were 1. Stop mix + probe without lipids, to assess maximum

fluorescence polarisation. 2. Stop mix + probe with lipids, to determine the effect of lipids on

maximum fluorescence polarisation. 3. Probe without stop mix or lipids to determine

minimum fluorescence polarisation and 4. Probe with lipids but no stop mix to determine the

effect of lipids on minimum fluorescence polarisation. Standard curves of diC8-PI(3,4,5)P3 in

the presence of liposomes were included in every experiment and used to infer PIP3

generated by purified PI3K. Graphpad Prism v6.0 was used to generate sigmoidal standard

curves by plotting log transformed diC8-PI(3,4,5)P3 concentrations against fluorescence

polarisation. Standard curves for each experiment provided the linear range within which

PIP3 could be accurately determined. Preliminary studies found only trivial differences in

measured fluorescence polarisation in the presence of pY2, which was thus omitted from

subsequent standard curves. Quadruplicate interpolated PIP3 concentrations were averaged

and normalised by reaction time (30 min) and enzyme concentration in pM to yield PI3K

activity in pmol PIP3/min/pM enzyme. Specific activities were then further normalised to

activity of p85α wild-type controls in the presence of pY2 in each experiment.

Generation of 3T3-L1 cells conditionally expressing p85α or p110α

3T3-L1 preadipocytes (Zenbio, lot 3T3062104, passage 8) were cultured, differentiated and

stained for neutral lipid with Oil Red O as previously described(Huang-Doran *et al*, 2011).

Generation of 3T3-L1 sublines inducibly expressing WT, Y657X, or R649W p85α was also

previously reported(Huang-Doran *et al.*, 2016; Hussain *et al*, 2011). The 3T3-L1 lines

inducibly expressing p85α ΔEx11 or p110α H1047R were generated using essentially the

same procedure, starting with In-Fusion subcloning of the p85α ΔEx11 cDNA insert from the

pACEBac1 p85α ΔEx11 plasmid described above, and of PIK3CA H1047R cDNA (Based on

Uniprot P42336) derived by PCR of a pre-existing expression vector, into the pEN_Tmcs

entry vector. After sequence verification these inserts were integrated into a pSLIK-Hygro

lentiviral vector, packaged into VSV-G–pseudotyped lentiviral particles, using packaging

plasmids pMDLg/pRRE and pRSVREV, and VSV-G envelope plasmid pVSV, and used to

infect 3T3-L1 preadipocytes (Plasmids listed in **Table 2**). Transgene expression was

induced with 1 µg/mL doxycycline for 72 hours, or with variable concentrations of

doxycycline as indicated.

Insulin signalling studies

3T3-L1 cells serum starved in DMEM containing 0.5 % BSA for 16h were stimulated with

100nM Actrapid insulin (Novo Nordisk) for 10min. Monolayers were snap frozen in liquid N2

and stored at -80°C. 200-600µL co-IP lysis (20mM HEPES, 150mM NaCl, 1.5mM MgCl2,

10% (v/v) glycerol, 1% (v/v) Triton X-100, 1mM EGTA pH 7.4, 1mM PMSF, 2mM activated

sodium orthovanadate, Complete Mini EDTA-free protease inhibitor cocktail and PhosSTOP

(Roche 04906837001), in Milli-Q Ultrapure water (Millipore)) or RIPA buffer (50mM Tris HCl

pH 8.0, 150mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS with added

Complete Mini EDTA-free protease inhibitor cocktail and PhosSTOP, in Milli-Q Ultrapure

Water (Millipore)) was added to frozen cells before scraping of lysate into pre-chilled tubes,

incubation on ice for 30min and clearing by centrifugation. Protein was quantified using the

Bio-Rad DC assay.

Lysates were mixed with NuPAGE® LDS Loading buffer (Life Technologies, NP0008)

supplemented with 5% β-mercaptoethanol and boiled before SDS-PAGE. For co-

immunoprecipitation (Co-IP) lysates (150-300µg protein) were incubated in 500µL Co-IP

buffer with immunoprecipitating antibody overnight at 4°C. 10µg lysate mixed with

NuPAGE® SDS Loading buffer and 5% β-mercaptoethanol was stored to represent Co-IP

input. Co-IP samples were incubated for 2h with 1.5mg PBST-washed Protein G

Dynabeads® (Life Technologies, 10003D) at 4°C before centrifugation and bead removal

using a DynaMag™-2 Magnet (Invitrogen). Supernatants mixed 1:1 with Co-IP elution buffer

(2X NuPAGE® LDS, 100 mM NuPAGE® Sample Reducing Agent (Invitrogen, NP0009), in

Co-IP lysis buffer) were boiled for 10 min. Beads were washed 5 times in 50μL Co-IP lysis

buffer, before protein elution with 25μL Co-IP Elution buffer and boiling for 10min. Input,

supernatant and co-IP samples underwent SDS-PAGE using NuPAGE® 4-12% gradient

Bis-Tris gels (ThermoFisher, NP0321BOX) in NuPAGE® MOPS SDS Running Buffer (Life

Technologies, NP0001). Transfer to nitrocellulose was performed using the iBlot™ Dry

Blotting System (Invitrogen), with preincubation in Equilibration Transfer Buffer for proteins

above 150kDa.

For immunoblotting, blocked membranes were incubated overnight at 4°C in primary

antibody, washed and incubated with HRP-linked anti-rabbit or anti-mouse IgG secondary

antibody diluted 1:5000 in blocking buffer. Proteins were visualized using the ChemiDoc™

MP System (Bio-Rad) and band intensities quantified using Image Lab software (Bio-Rad).

Statistical Analysis

For quantitative data, all biological replicates (i.e. results of independent experiments on

different days) are represented in figures, with paired points (e.g. with/without insulin in the

same experiment) connected by lines. For fluorescence polarisation assays, each biological

replicate shown is the mean of 3 technical replicates. Sample size for individual experiments

was not pre-determined. To avoid the pitfalls of dichotomous significance testing on low-

throughput biological datasets, we used estimation statistics (Data Analysis using Bootstrap-

Coupled ESTimation) with default settings (5000 resamples, BCa boostrap)(Ho *et al*, 2019).

This focuses on effect sizes and derives confidence intervals derived from bootstrapping for

differences in means; the small (but typical) sample size of the experiments analysed limits

reliable bootstrapping, but it offers additional confidence to the consistent patterns seen in

independent replicates. We have indicated significance in main text figures with an asterisk,

and show the 95% confidence intervals for mean differences in extended view figures.

Data and Resource Availability

This study includes no data deposited in external repositories. Uncut Western blot and other

source images are available on the Open Science Framework project webpage:

https://osf.io/eyc4h/?view_only=9af8645c8fe5408790ce19920d0084c6. All antibodies used

are listed in **Table 1**, and plasmids in **Table 2**.

Acknowledgements

This work was supported by the Wellcome Trust through a grant to RKS [210752/Z/18/Z]

and a studentship to PRT [102356/Z/13/Z]. Additional support was from the UK Medical

Research Council (MRC) [MC_UU_12012/5 and MC_U105184308 (to RLW)] and the

Intramural Research Program of the National Institute of Allergy and Infectious Diseases,

National Institutes of Health (to HCS). We thank Dr. Koneti Rao, Debra Long-Priel, and

Angela Wang for clinical, technical, and regulatory assistance.

Author Contributions

PRT: conceptualisation, methodology, formal analysis, investigation, visualization, writing -

original draft. **RK**: formal analysis, investigation, visualization. **OP**: supervision,

methodology, investigation, writing - review & editing. **HCS:** resources, funding acquisition,

writing - review & editing. **GVB:** supervision, methodology, writing - review & editing. **RLW**:

supervision, methodology, investigation, funding acquisition, writing - review & editing. **RKS:**

conceptualisation, supervision, methodology, funding acquisition, writing - original draft,

writing - review & editing, funding acquisition.

Disclosure and competing interests statement

RKS has consulted for Novartis on clinical aspects of PIK3CA-related overgrowth, and for

Alnylam, Amryt and AstraZeneca on clinical aspects of monogenic insulin resistance and

lipodystrophy

References

Angulo I, Vadas O, Garcon F, Banham-Hall E, Plagnol V, Leahy TR, Baxendale H, Coulter T,
Curtis J, Wu C *et al* (2013) Phosphoinositide 3-kinase delta gene mutation predisposes to
respiratory infection and airway damage. *Science* 342: 866-871
Avila M, Dyment DA, Sagen JV, St-Onge J, Moog U, Chung BHY, Mo S, Mansour S, Albanese
A, Garcia S *et al* (2016) Clinical reappraisal of SHORT syndrome with PIK3R1 mutations:
toward recommendation for molecular testing and management. *Clin Genet* 89: 501-506
Brachmann SM, Ueki K, Engelman JA, Kahn RC, Cantley LC (2005) Phosphoinositide 3-kinase
catalytic subunit deletion and regulatory subunit deletion have opposite effects on insulin
sensitivity in mice. *Mol Cell Biol* 25: 1596-1607
Bravo Garcia-Morato M, Garcia-Minaur S, Molina Garicano J, Santos Simarro F, Del Pino
Molina L, Lopez-Granados E, Ferreira Cerdan A, Rodriguez Pena R (2017) Mutations in
PIK3R1 can lead to APDS2, SHORT syndrome or a combination of the two. *Clin Immunol* 179:
77-80
Burke JE, Perisic O, Masson GR, Vadas O, Williams RL (2012) Oncogenic mutations mimic
and enhance dynamic events in the natural activation of phosphoinositide 3-kinase
p110alpha (PIK3CA). *Proc Natl Acad Sci U S A* 109: 15259-15264
Chen D, Mauvais-Jarvis F, Bluher M, Fisher SJ, Jozsi A, Goodyear LJ, Ueki K, Kahn CR (2004)
p50alpha/p55alpha phosphoinositide 3-kinase knockout mice exhibit enhanced insulin
sensitivity. *Mol Cell Biol* 24: 320-329
Cheung LW, Walkiewicz KW, Besong TM, Guo H, Hawke DH, Arold ST, Mills GB (2015)
Regulation of the PI3K pathway through a p85alpha monomer-homodimer equilibrium. *Elife*
4: e06866
Chudasama KK, Winnay J, Johansson S, Claudi T, Konig R, Haldorsen I, Johansson B, Woo JR,
Aarskog D, Sagen JV *et al* (2013) SHORT syndrome with partial lipodystrophy due to
impaired phosphatidylinositol 3 kinase signaling. *Am J Hum Genet* 93: 150-157
Conley ME, Dobbs AK, Quintana AM, Bosompem A, Wang YD, Coustan-Smith E, Smith AM,
Perez EE, Murray PJ (2012) Agammaglobulinemia and absent B lineage cells in a patient
lacking the p85alpha subunit of PI3K. *J Exp Med* 209: 463-470
Cottrell CE, Bender NR, Zimmermann MT, Heusel JW, Corliss M, Evenson MJ, Magrini V,
Corsmeier DJ, Avenarius M, Dudley JN *et al* (2021) Somatic PIK3R1 variation as a cause of
vascular malformations and overgrowth. *Genet Med*
Deau MC, Heurtier L, Frange P, Suarez F, Bole-Feysot C, Nitschke P, Cavazzana M, Picard C,
Durandy A, Fischer A *et al* (2014) A human immunodeficiency caused by mutations in the
PIK3R1 gene. *J Clin Invest* 124: 3923-3928
Dornan GL, Siempelkamp BD, Jenkins ML, Vadas O, Lucas CL, Burke JE (2017)
Conformational disruption of PI3Kdelta regulation by immunodeficiency mutations in
PIK3CD and PIK3R1. *Proc Natl Acad Sci U S A* 114: 1982-1987
Dornan GL, Stariha JTB, Rathinaswamy MK, Powell CJ, Boulanger MJ, Burke JE (2020)
Defining How Oncogenic and Developmental Mutations of PIK3R1 Alter the Regulation of
Class IA Phosphoinositide 3-Kinases. *Structure* 28: 145-156 e145
Dyment DA, Smith AC, Alcantara D, Schwartzentruber JA, Basel-Vanagaite L, Curry CJ,
Temple IK, Reardon W, Mansour S, Haq MR *et al* (2013) Mutations in PIK3R1 cause SHORT
syndrome. *Am J Hum Genet* 93: 158-166
Elkaim E, Neven B, Bruneau J, Mitsui-Sekinaka K, Stanislas A, Heurtier L, Lucas CL, Matthews
H, Deau MC, Sharapova S *et al* (2016) Clinical and immunologic phenotype associated with
activated phosphoinositide 3-kinase delta syndrome 2: A cohort study. *J Allergy Clin*
Immunol 138: 210-218 e219

Fruman DA, Chiu H, Hopkins BD, Bagrodia S, Cantley LC, Abraham RT (2017) The PI3K
Pathway in Human Disease. *Cell* 170: 605-635
Fruman DA, Mauvais-Jarvis F, Pollard DA, Yballe CM, Brazil D, Bronson RT, Kahn CR, Cantley
LC (2000) Hypoglycaemia, liver necrosis and perinatal death in mice lacking all isoforms of
phosphoinositide 3-kinase p85 alpha. *Nat Genet* 26: 379-382
Fruman DA, Snapper SB, Yballe CM, Davidson L, Yu JY, Alt FW, Cantley LC (1999) Impaired B
cell development and proliferation in absence of phosphoinositide 3-kinase p85alpha.
Science 283: 393-397
Geering B, Cutillas PR, Nock G, Gharbi SI, Vanhaesebroeck B (2007) Class IA
phosphoinositide 3-kinases are obligate p85-p110 heterodimers. *Proc Natl Acad Sci U S A*
104: 7809-7814
Harpur AG, Layton MJ, Das P, Bottomley MJ, Panayotou G, Driscoll PC, Waterfield MD (1999)
Intermolecular interactions of the p85alpha regulatory subunit of phosphatidylinositol 3-
kinase. *J Biol Chem* 274: 12323-12332
Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A (2019) Moving beyond P values: data
analysis with estimation graphics. *Nat Methods* 16: 565-566
Huang-Doran I, Bicknell LS, Finucane FM, Rocha N, Porter KM, Tung YC, Szekeres F, Krook A,
Nolan JJ, O'Driscoll M *et al* (2011) Genetic defects in human pericentrin are associated with
severe insulin resistance and diabetes. *Diabetes* 60: 925-935
Huang-Doran I, Tomlinson P, Payne F, Gast A, Sleigh A, Bottomley W, Harris J, Daly A, Rocha
N, Rudge S *et al* (2016) Insulin resistance uncoupled from dyslipidemia due to C-terminal
PIK3R1 mutations. *JCI Insight* 1: e88766
Hussain K, Challis B, Rocha N, Payne F, Minic M, Thompson A, Daly A, Scott C, Harris J,
Smillie BJ *et al* (2011) An activating mutation of AKT2 and human hypoglycemia. *Science*
334: 474
Jamee M, Moniri S, Zaki-Dizaji M, Olbrich P, Yazdani R, Jadidi-Niaragh F, Aghamahdi F,
Abolhassani H, Condliffe AM, Aghamohammadi A *et al* (2020) Clinical, Immunological, and
Genetic Features in Patients with Activated PI3Kdelta Syndrome (APDS): a Systematic
Review. *Clin Rev Allergy Immunol* 59: 323-333
Knight ZA, Gonzalez B, Feldman ME, Zunder ER, Goldenberg DD, Williams O, Loewith R,
Stokoe D, Balla A, Toth B *et al* (2006) A pharmacological map of the PI3-K family defines a
role for p110alpha in insulin signaling. *Cell* 125: 733-747
Kwok A, Zvetkova I, Virtue S, Luijten I, Huang-Doran I, Tomlinson P, Bulger DA, West J,
Murfitt S, Griffin J *et al* (2020) Truncation of Pik3r1 causes severe insulin resistance
uncoupled from obesity and dyslipidaemia by increased energy expenditure. *Mol Metab* 40:
101020
Lindhurst MJ, Parker VE, Payne F, Sapp JC, Rudge S, Harris J, Witkowski AM, Zhang Q,
Groeneveld MP, Scott CE *et al* (2012) Mosaic overgrowth with fibroadipose hyperplasia is
caused by somatic activating mutations in PIK3CA. *Nat Genet* 44: 928-933
LoPiccolo J, Kim SJ, Shi Y, Wu B, Wu H, Chait BT, Singer RH, Sali A, Brenowitz M, Bresnick AR
et al (2015) Assembly and Molecular Architecture of the Phosphoinositide 3-Kinase
p85alpha Homodimer. *J Biol Chem* 290: 30390-30405
Lucas CL, Kuehn HS, Zhao F, Niemela JE, Deenick EK, Palendira U, Avery DT, Moens L,
Cannons JL, Biancalana M *et al* (2014a) Dominant-activating germline mutations in the gene
encoding the PI(3)K catalytic subunit p110delta result in T cell senescence and human
immunodeficiency. *Nat Immunol* 15: 88-97

Lucas CL, Zhang Y, Venida A, Wang Y, Hughes J, McElwee J, Butrick M, Matthews H, Price S,
Biancalana M *et al* (2014b) Heterozygous splice mutation in PIK3R1 causes human
immunodeficiency with lymphoproliferation due to dominant activation of PI3K. *J Exp Med*
211: 2537-2547
Luo J, Field SJ, Lee JY, Engelman JA, Cantley LC (2005) The p85 regulatory subunit of
phosphoinositide 3-kinase down-regulates IRS-1 signaling via the formation of a
sequestration complex. *J Cell Biol* 170: 455-464
Maccari ME, Wolkewitz M, Schwab C, Lorenzini T, Leiding JW, Aladjdi N, Abolhassani H,
Abou-Chahla W, Aiuti A, Azarnoush S *et al* (2023) Activated phosphoinositide 3-kinase delta
syndrome: Update from the ESID Registry and comparison with other autoimmune-
lymphoproliferative inborn errors of immunity. *J Allergy Clin Immunol* 152: 984-996 e910
Madsen RR, Vanhaesebroeck B, Semple RK (2018) Cancer-Associated PIK3CA Mutations in
Overgrowth Disorders. *Trends Mol Med* 24: 856-870
Mauvais-Jarvis F, Ueki K, Fruman DA, Hirshman MF, Sakamoto K, Goodyear LJ, Iannacone M,
Accili D, Cantley LC, Kahn CR (2002) Reduced expression of the murine p85alpha subunit of
phosphoinositide 3-kinase improves insulin signaling and ameliorates diabetes. *J Clin Invest*
109: 141-149
Myers MG, Jr., Backer JM, Sun XJ, Shoelson S, Hu P, Schlessinger J, Yoakim M, Schaffhausen
B, White MF (1992) IRS-1 activates phosphatidylinositol 3'-kinase by associating with src
homology 2 domains of p85. *Proc Natl Acad Sci U S A* 89: 10350-10354
Nguyen T, Lau A, Bier J, Cooke KC, Lenthall H, Ruiz-Diaz S, Avery DT, Brigden H, Zahra D,
Sewell WA *et al* (2023) Human PIK3R1 mutations disrupt lymphocyte differentiation to
cause activated PI3Kdelta syndrome 2. *J Exp Med* 220
Olbrich P, Lorenz M, Cura Daball P, Lucena JM, Rensing-Ehl A, Sanchez B, Fuhrer M,
Camacho-Lovillo M, Melon M, Schwarz K *et al* (2016) Activated PI3Kdelta syndrome type 2:
Two patients, a novel mutation, and review of the literature. *Pediatr Allergy Immunol* 27:
640-644
Petrovski S, Parrott RE, Roberts JL, Huang H, Yang J, Gorentla B, Mousallem T, Wang E,
Armstrong M, McHale D *et al* (2016) Dominant Splice Site Mutations in PIK3R1 Cause Hyper
IgM Syndrome, Lymphadenopathy and Short Stature. *J Clin Immunol* 36: 462-471
Philp AJ, Campbell IG, Leet C, Vincan E, Rockman SP, Whitehead RH, Thomas RJ, Phillips WA
(2001) The phosphatidylinositol 3'-kinase p85alpha gene is an oncogene in human ovarian
and colon tumors. *Cancer Res* 61: 7426-7429
Ramirez L, Tamayo W, Ale H (2020) APDS2 and SHORT Syndrome in a Teenager with PIK3R1
Pathogenic Variant. *J Clin Immunol* 40: 1020-1025
Rordorf-Nikolic T, Van Horn DJ, Chen D, White MF, Backer JM (1995) Regulation of
phosphatidylinositol 3'-kinase by tyrosyl phosphoproteins. Full activation requires
occupancy of both SH2 domains in the 85-kDa regulatory subunit. *J Biol Chem* 270: 3662-
3666
Solheim MH, Winnay JN, Batista TM, Molven A, Njolstad PR, Kahn CR (2018) Mice Carrying a
Dominant-Negative Human PI3K Mutation Are Protected From Obesity and Hepatic
Steatosis but Not Diabetes. *Diabetes* 67: 1297-1309
Sun XJ, Wang LM, Zhang Y, Yenush L, Myers MG, Jr., Glasheen E, Lane WS, Pierce JH, White
MF (1995) Role of IRS-2 in insulin and cytokine signalling. *Nature* 377: 173-177
Suzuki H, Terauchi Y, Fujiwara M, Aizawa S, Yazaki Y, Kadowaki T, Koyasu S (1999) Xid-like
immunodeficiency in mice with disruption of the p85alpha subunit of phosphoinositide 3-
kinase. *Science* 283: 390-392

Szczawinska-Poplonyk A, Bernat-Sitarz K, Schwartzmann E, Piechota M, Badura-Stronka M
(2022) Clinical and immunological assessment of APDS2 with features of the SHORT
syndrome related to a novel mutation in PIK3R1 with reduced penetrance. *Allergol*
Immunopathol (Madr) 50: 1-9
Tang P, Upton JEM, Barton-Forbes MA, Salvadori MI, Clynick MP, Price AK, Goobie SL (2018)
Autosomal Recessive Agammaglobulinemia Due to a Homozygous Mutation in PIK3R1. *J Clin*
Immunol 38: 88-95
Thauvin-Robinet C, Auclair M, Duplomb L, Caron-Debarle M, Avila M, St-Onge J, Le Merrer
M, Le Luyer B, Heron D, Mathieu-Dramard M *et al* (2013) PIK3R1 mutations cause syndromic
insulin resistance with lipoatrophy. *Am J Hum Genet* 93: 141-149
Thorpe LM, Spangle JM, Ohlson CE, Cheng H, Roberts TM, Cantley LC, Zhao JJ (2017) PI3K-
p110alpha mediates the oncogenic activity induced by loss of the novel tumor suppressor
PI3K-p85alpha. *Proc Natl Acad Sci U S A* 114: 7095-7100
Tsolakos N, Durrant TN, Chessa T, Suire SM, Oxley D, Kulkarni S, Downward J, Perisic O,
Williams RL, Stephens L *et al* (2018) Quantitation of class IA PI3Ks in mice reveals p110-free-
p85s and isoform-selective subunit associations and recruitment to receptors. *Proc Natl*
Acad Sci U S A 115: 12176-12181
Ueki K, Fruman DA, Brachmann SM, Tseng YH, Cantley LC, Kahn CR (2002) Molecular balance
between the regulatory and catalytic subunits of phosphoinositide 3-kinase regulates cell
signaling and survival. *Mol Cell Biol* 22: 965-977
Wentink M, Dalm V, Lankester AC, van Schouwenburg PA, Scholvinck L, Kalina T, Zachova R,
Sediva A, Lambeck A, Pico-Knijnenburg I *et al* (2017) Genetic defects in PI3Kdelta affect B-
cell differentiation and maturation leading to hypogammaglobulineamia and recurrent
infections. *Clin Immunol* 176: 77-86
Winnay JN, Solheim MH, Dirice E, Sakaguchi M, Noh HL, Kang HJ, Takahashi H, Chudasama
KK, Kim JK, Molven A *et al* (2016) PI3-kinase mutation linked to insulin and growth factor
resistance in vivo. *J Clin Invest* 126: 1401-1412
Yu J, Zhang Y, McIlroy J, Rordorf-Nikolic T, Orr GA, Backer JM (1998) Regulation of the
p85/p110 phosphatidylinositol 3'-kinase: stabilization and inhibition of the p110alpha
catalytic subunit by the p85 regulatory subunit. *Mol Cell Biol* 18: 1379-1387
Zhao JJ, Cheng H, Jia S, Wang L, Gjoerup OV, Mikami A, Roberts TM (2006) The p110alpha
isoform of PI3K is essential for proper growth factor signaling and oncogenic
transformation. *Proc Natl Acad Sci U S A* 103: 16296-16300
764
765

 Figures

Figure 1. PI3K subunit expression and signalling in primary dermal fibroblasts.
Immunoblotting of AKT, AKT phosphorylated at threonine 308 (T308) or serine 473 (S473),
p85α, p110δ and p110α and are shown with and without stimulation by 100 nM insulin (Ins)
for 10 minutes. β-actin is shown as a loading control, with different amounts of pooled lysate
(Pool) used to demonstrate signal intensity in the linear range. Results from 4 healthy controls
(WT; 1-4), 1 patient with Activating p110 Delta Syndrome 2 (APDS2) due to the p85α Δexon11
variant, and 3 patients with PIK3CA-Related Overgrowth Spectrum (PROS) caused by the
activating *PIK3CA* mutations indicated. (A) Immunoblots, with the truncated p85α Δexon11
variant arrowed; (B)-(E) quantification of immunoblot bands from 3 independent experiments
shown for phosphoAKT-S473, phosphoAKT-T308, p110δ and p110α respectively. Each point
represents data from one of the patient cell lines in the immunoblots. Paired datapoints +/-
insulin are shown in (B) and (C), and dotted lines mark means. Asterisks indicate a significant
difference. More detailed statistical analysis including 95% confidence intervals for the paired
mean differences for these comparisons are shown in Figure 1 figure supplement 2.

Figure 2. Blunted insulin signaling in 3T3-L1 preadipocyte models of APDS2 and
SHORT syndrome. Immunoblotting of Akt, Akt phosphorylated at threonine 308 (T308) or
serine 473 (S473), p85α, and p110α and are shown with and without stimulation with 100 nM
insulin (Ins) for 10 minutes. Cells were treated with doxycycline (Dox) 1μg/mL for 72 hours
prior to insulin stimulation as indicated. (A) One immunoblot representing 3 experiments is
shown. (B)-(C) Quantification of immunoblot bands from all 3 independent experiments shown
for phosphoAkt-S473 and phosphoAkt-T308 respectively. Paired datapoints +/- insulin are
shown, and dotted lines mark means. Asterisks indicate a significant difference. More detailed
statistical analysis including 95% confidence intervals for the paired mean differences for
these comparisons are shown in Figure 2 figure supplement 2. (D) Staining for neutral lipid
with Oil Red O of 3T3-L1 cells at day 10 of adipocyte differentiation. Induction of transgene
expression by 1μg/mL doxycycline throughout differentiation is shown. Images of entire plates
are shown above, with representative bright field microscopy images below.

Figure 3. SHORT syndrome p85α mutations impair phosphotyrosine-stimulated phosphoinositide 3-kinase activity. Lipid kinase activity of purified recombinant PI3K complexes generated using baculoviral expression in *Sf9* cells was measured using a modified fluorescence polarisation assay. WT p85α or p85α SHORT syndrome mutations, E489K, R649W or Y657X bound to either (A) p110α (B) p110β or (C) p110δ were assayed for basal and bisphosphotyrosine (pY2)-stimulated lipid kinase activity. Dotted lines mark means, and asterisks indicate a significant difference between the bisphosphotyrosine (pY2)-stimulated state for WT and comparator mutant p85α. More detailed statistical analysis including 95% confidence intervals for the paired mean differences for these comparisons are shown in Figure 3 figure supplement 1.

Figure 4. Ability of pathogenic p85α variants to bind p110α, assessed by co-immunoprecipitation. Results of immunoblotting of anti-p110α immunoprecipitates from 3T3-L1 cells expressing wild-type, APDS2-associated or SHORT syndrome-associated mutant p85α under the control of doxycycline (Dox) are shown. (A) One representative immunoblot of immunoprecipitate, cell lysate prior to immunoprecipitation, and post immunoprecipitation supernatant is shown. (B) Quantification of immunoblot bands from immunoprecipitates from 3 independent experiments, expressed as a percentage relative to the intensity of the band in WT cells without doxycycline exposure. Co-immunoprecipitated p85α is shown normalized to immunoprecipitated p110α from all 3 independent experiments. Datapoints from the same experiment +/- doxycline are connected by lines. No significant differences were found among conditions.

Figure 5. Attenuated insulin-induced association of p110α with Irs1 in the presence of APDS2 and SHORT syndrome mutant p85α. Results of immunoblotting of anti-Irs1 immunoprecipitates from 3T3-L1 cells expressing wild-type, APDS2-associated or SHORT syndrome-associated mutant p85α under the control of doxycycline (Dox) are shown.

Treatment with 100nM insulin (Ins) is indicated. (A) One representative immunoblot of immunoprecipitate, cell lysate prior to immunoprecipitation, and post immunoprecipitation supernatant is shown. Two separate sets of gels, including independent wild-type controls are shown on left and right. (B-C) Quantification of immunoblot bands from immunoprecipitates from 3 independent experiments. Immunoprecipitated p110α is shown normalized to immunoprecipitated Irs1 from all 3 independent experiments in (B), and immunoprecipitated p85α similarly in (C). Datapoints from the same experiment +/- insulin are connected by lines. Asterisks indicate significant differences induced by transgene overexpression (i.e. plus versus minus doxycycline). More detailed statistical analysis including 95% confidence intervals for the paired mean differences for these comparisons are shown in Figure 5 figure supplement 1.

Figure 6. Attenuated insulin-induced association of p110α with Irs2 in the presence of APDS2 and SHORT syndrome mutant p85α. Results of immunoblotting of anti-Irs2 immunoprecipitates from 3T3-L1 cells expressing wild-type, APDS2-associated or SHORT syndrome-associated mutant p85α under the control of doxycycline (Dox) are shown.

Treatment with 100nM insulin (Ins) is indicated. (A) One representative immunoblot of
immunoprecipitate, cell lysate prior to immunoprecipitation, and post immunoprecipitation
supernatant is shown. Two separate sets of gels, including independent wild-type controls are
shown on left and right. (B-C) Quantification of immunoblot bands from immunoprecipitates
from 3 independent experiments. Immunoprecipitated p110α is shown normalized to
immunoprecipitated Irs2 from all 3 independent experiments in (B), and immunoprecipitated
p85α similarly in (C). Datapoints from the same experiment +/- insulin are connected by lines.
92

93   **Tables**

Antibody Target	Company	Catalogue No.	Species	Dilution used (immunoblot/IP)
p85α	CST	4257	Rabbit	1:1000
Phospho-AKT/Akt (T308)	CST	9275	Rabbit	1:1000
Phospho-AKT/Akt (S473)	CST	9271	Rabbit	1:1000
AKT/Akt (1,2,3)	CST	2920	Mouse	1:1000
p110α	CST	4249	Rabbit	1:1000/1:50
p110δ	CST	34050	Rabbit	1:1000
Irs1	Millipore	06-248	Rabbit	1:500
Irs1	CST	2382	Rabbit	1:1000/1:50
Irs2 clone 9.52	Millipore	MABS15	Mouse	1:1000/1:50
β-actin	CST	4967	Rabbit	1:5000
Anti-rabbit IgG; HRP-linked	CST	7074	Goat	1:5000
Anti-mouse IgG; HRP-linked	CST	7076	Horse	1:5000

101   **Table 1: Antibodies Used**

Plasmid Name	Insert	Comment
Hsp85a_pACEBac1	Human p85α (UniProtKB P27986.2)	Generated in Williams Lab, MRC-LMB
Hsp85a-Y657*_pACEBac1	Human p85α-Y657*	Based on Hsp85a_pACEBac1
Hsp85a-R649W_pACEBac1	Human p85α-R649W	Based on Hsp85a_pACEBac1
Hsp85a-E489K_pACEBac1	Human p85α-E489K	Based on Hsp85a_pACEBac1
Hsp85a-dEx11_pACEBac1	Human p85α-dEx11	Based on Hsp85a_pACEBac1
Hsp110a_pFastBacHT B	Human p110α (UniProtKB P42336.2)	Generated in Williams Lab, MRC-LMB
Hsp110b_pACEBac1	Human p110β (UniProtKB P42338.1)	Generated in Williams Lab, MRC-LMB
Hsp110d_pFastBacHT B	Human p110δ (UniProtKB O00329.2)	Generated in Williams Lab, MRC-LMB
pEN_Tmcs		Addgene #25751
pSLIK-Hygro		Addgene #25737
Hsp85a-dEx11_pSLIK-Hygro	Human p85α-dEx11	
pMDLg/pRRE		Addgene #12251
pRSV-Rev		Addgene #12253
pMD2.G		Addgene #12259

104 **Table 1: Plasmids Used**