Paradoxical dominant negative activity of an immunodeficiency-associated activating *PIK3R1* variant

Patsy R. Tomlinson[1,2,3], Rachel Knox[1,2,3], Olga Perisic[4], Helen C. Su[6], Gemma V. Brierley[1,2,3], Roger L. Williams[4], Robert K. Semple[3,5,7]

[1] The University of Cambridge Metabolic Research Laboratories, Wellcome Trust-MRC Institute of Metabolic Science, Cambridge, UK
[2] The National Institute for Health Research Cambridge Biomedical Research Centre, Cambridge, UK
[3] MRC Metabolic Diseases Unit, Wellcome Trust-MRC Institute of Metabolic Science, Cambridge, UK
[4] MRC Laboratory of Molecular Biology, Cambridge, UK
[5] Centre for Cardiovascular Science, University of Edinburgh, Edinburgh, UK
[6] Laboratory of Clinical Immunology & Microbiology, Intramural Research Program, National Institute of Allergy and Infectious Disease, National Institutes of Health, Bethesda, MD, USA
[7] MRC Human Genetics Unit, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh, UK

Correspondence to:

Prof. Robert K. Semple, Centre for Cardiovascular Science, University of Edinburgh, Queen's Medical Research Institute, Little France Crescent, Edinburgh EH16 4TJ, UK. Tel: +44 131 242 6051/ Email: rsemple@ed.ac.uk

Character Count: 50,222 (to end of references; with spaces)

Running Title: Dominant negative effect of an activating PIK3R1 variant

Abstract

PIK3R1 encodes three regulatory subunits of class IA phosphoinositide 3-kinase (PI3K),

each associating with any of three catalytic subunits, namely p110α, p110β or p110δ.

Constitutional *PIK3R1* mutations cause diseases with a genotype-phenotype relationship not

yet fully explained: heterozygous loss-of-function mutations cause SHORT syndrome,

featuring insulin resistance and short stature attributed to reduced p110α function, while

heterozygous activating mutations cause immunodeficiency, attributed to p110δ activation

and known as APDS2. Surprisingly, APDS2 patients do not show features of p110α

hyperactivation, but do commonly have short stature or SHORT syndrome, suggesting

p110α hypofunction. We sought to investigate this. In dermal fibroblasts from an APDS2

patient, we found no increased PI3K signalling, with p110δ expression markedly reduced. In

preadipocytes, the APDS2 variant was potently dominant negative, associating with Irs1 and

Irs2 but failing to heterodimerise with p110α. This attenuation of p110α signalling by a

p110δ-activating PIK3R1 variant potentially explains co-incidence of gain-of-function and

loss-of-function *PIK3R1* phenotypes.

Keywords: APDS2, cancer, immunodeficiency, insulin resistance, phosphoinositide 3-

kinase, PIK3R1, SHORT syndrome

Introduction

PIK3R1 gene products (p85α, p55α and p50α) are essential for class IA phosphoinositide 3-kinase (PI3K) signalling. Each of the three protein products can bind any of three catalytic subunits, p110α, β and δ, encoded by *PIK3CA*, *PIK3CB*, and *PIK3CD* respectively. This stabilises and inhibits the catalytic subunits in the basal state and confers responsiveness to upstream stimuli including receptor tyrosine kinase activation, enabled by two phosphotyrosine-binding SH2 domains shared by all PIK3R1 products (Fruman *et al*, 2017).

PI3K signalling mediates responses to a myriad of cues, including growth factors, hormones such as insulin, and processed antigens, and so recent discovery that genetic perturbation of PIK3R1 in humans disrupts growth, insulin action and immunity is no surprise. Beyond this headline observation lies considerable complexity, however, and important questions about genotype-phenotype correlations in PIK3R1-related disorders are unresolved.

Some PIK3R1 mutations reduce basal inhibition of catalytic subunits, usually due to disruption of the inhibitory inter-SH2 domain, and are found in cancers (Philp *et al*, 2001) and vascular malformations with overgrowth(Cottrell *et al*, 2021). In both diseases, hyperactivated PI3Kα, composed of heterodimers of *PIK3R1* products and p110α, drives pathological growth. Distinct inter-SH2 domain *PIK3R1* mutations, mostly causing skipping of exon 11 and deletion of residues 434-475, hyperactivate PI3Kδ in immune cells, causing immunodeficiency (Deau *et al*, 2014; Lucas *et al*, 2014b). This phenocopies the immunodeficiency caused by genetic activation of p110δ itself, called Activated PI3K Delta Syndrome 1 (APDS1) (Angulo *et al*, 2013; Lucas *et al*, 2014a), and is thus named APDS2.

Despite ubiquitous *PIK3R1* expression, features attributable to PI3Kα activation have been described in only a single case of APDS2 to date (Wentink *et al*, 2017). In that case, the causal variant was a constitutional activating mutation associated with cancer rather than one of the common APDS2 variants. Biochemical studies have suggested that apparently selective p110δ activation occurs because APDS2 mutations derepress p110δ, the

predominant immune system catalytic subunit, more than p110α, due to differences in the

inhibitory contacts between *PIK3R1* products and different catalytic subunits (Dornan *et al*,

2017).

More surprisingly, phenotypic overlap is reported between APDS2 and SHORT syndrome.

SHORT syndrome is caused by loss of PI3Kα function due to disruption of the

phosphotyrosine-binding C-terminal SH2 domain (Chudasama *et al*, 2013; Dyment *et al*,

2013; Thauvin-Robinet *et al*, 2013). It features reduced linear growth, insulin resistance, and

dysmorphic features (Avila *et al*, 2016). In recent years, both case reports (Bravo Garcia-

Morato *et al*, 2017; Petrovski *et al*, 2016; Ramirez *et al*, 2020; Szczawinska-Poplonyk *et al*,

2022) and larger series (Elkaim *et al*, 2016; Jamee *et al*, 2020; Maccari *et al*, 2023; Nguyen

et al, 2023; Olbrich *et al*, 2016; Petrovski *et al.*, 2016) have established that many people

with APDS2 have overt features of SHORT syndrome, while, more generally, linear growth

impairment is common in APDS2, but not in APDS1. These clinical observations are

bolstered by the impaired linear growth and increased *in utero* mortality reported in mice with

knock in of the common causal APDS2 mutation in Pik3r1(Nguyen *et al.*, 2023). All people

with SHORT-APDS2 overlap syndromes have well established activating PIK3R1 mutations

in the inter-SH2 domain implicated in APDS2, but none have characteristic SHORT

syndrome mutations, which are usually in the C-terminal SH2 domain. Conversely, no

patient with a classical SHORT syndrome mutation has been shown to have

immunodeficiency. There is thus now convincing evidence of a syndrome with features of

both gain and loss of PI3K function. The mechanistic basis of this is unexplained.

PI3K activity is determined not just by activity of individual PI3K holoenzymes, but also by

subunit stoichiometry. Regulatory and catalytic subunits stabilise each other upon binding

(Brachmann *et al*, 2005; Yu *et al*, 1998), and only regulatory subunit monomers are stable

enough to be detected in mammalian cells (Yu *et al.*, 1998). Molar excess of regulatory

subunits over catalytic subunits gives rise to free regulatory subunits that compete with

holoenzyme for phosphotyrosine binding (Ueki *et al*, 2002). This has been invoked to explain

increased insulin sensitivity on knockout or reduction of p85α alone (Mauvais-Jarvis *et al*,

2002), of p55α and p50α (Chen *et al*, 2004) or of all *Pik3r1* products (Fruman *et al*, 2000).

The "free-p85" model has been contested by some reports suggesting no excess p85

(Geering *et al*, 2007; Zhao *et al*, 2006), however most observations, including some made

recently using *in vivo* tagging and pulldown of PI3K components (Tsolakos *et al*, 2018)

suggest that monomeric p85α can be seen *in vivo* in some tissues.

We now address mechanisms underlying human mixed gain- and loss-of function

phenotypes observed in association with PIK3R1 mutations using primary cells, cell models

and in vitro enzyme assay and suggest that they are explained by competing effects of

APDS2 PIK3R1 mutations on PI3K activity and stability.

Results

PI3K subunit expression and signalling in APDS2 fibroblasts

Disorders in the *PIK3CA*-related overgrowth spectrum (PROS) feature soft tissue

overgrowth and are caused by mosaic activating mutations in *PIK3CA*, encoding the

p110α catalytic subunit of PI3Kα(Madsen *et al*, 2018). In this group, basal hyperactivation of

PI3Kα signalling can be easily discerned in dermal fibroblasts from affected areas of the

body(Lindhurst *et al*, 2012). *PIK3R1*, like *PIK3CA*, is ubiquitously expressed, yet overgrowth

is not reported in APDS2 caused by constitutional activating mutations in *PIK3R1*. Dermal

fibroblasts strongly express *PIK3R1*, but no studies to date have evaluated whether PI3K

activity is increased in APDS2.

We began by assessing dermal fibroblasts cultured from a previously described woman with

APDS2 due to the common causal PIK3R1 mutation in the splice donor site that causes

skipping of exon 11 and an in-frame deletion in the inter-SH2 domain shared by all PIK3R1

isoforms (Patient A.1 in (Lucas *et al.*, 2014b)). These cells were compared to cells from

healthy control volunteers, or from people with PROS. Confirmation of expression of all

pathogenic mutations was undertaken by cDNA sequencing prior to further study (**Fig**

EV1A). We found that truncated p85α was expressed, but at a much lower level than full

length wild-type p85α (**Fig 1A, FigEV1B,C**). Despite this, no increase in basal or insulin-

stimulated AKT phosphorylation was seen in APDS2 cells compared to cells from wild-type

volunteers or from people with PROS and activating *PIK3CA* mutations H1047L or H1047R

(**Fig 1A-C, Fig EV2A,B**). Indeed, insulin-induced phosphorylation of AKT was lower in

fibroblasts from this APDS2 patient compared to controls.

Previous studies have suggested that the truncated, APDS2-causal p85α variant exerts a

much greater activating effect on p110δ, which has a more restricted tissue expression,

including immune cells, than p110α, which is ubiquitous (Dornan *et al.*, 2017). We found that

both p110α and p110δ were easily detectable in control dermal fibroblasts, however p110δ

was almost absent in APDS2 fibroblasts compared to controls, with lower levels also seen in

PROS cells (**Fig 1A,D, Fig EV2D**). p110α was unchanged in APDS2 cells but modestly

increased in PROS cells (**Fig 1A,E, Fig EV2C**). Collectively these findings suggest that any

ability of the APDS2 PIK3R1 variant in skin cells to activate PI3K may be overcome by

reduced protein levels of p110δ, likely through reduced binding and/or reduced stabilisation

of p110δ by the mutant regulatory subunit.

Overexpressed PIK3R1 ΔExon11 is potently dominant negative in 3T3-L1 preadipocytes

We next turned to a well-established cellular system allowing conditional overexpression of

PK3R1 alleles of interest. We have previously shown that overexpression of two SHORT

syndrome PIK3R1 variants – R649W and Y657X – impair insulin signalling and adipocyte

differentiation of murine 3T3-L1 preadipocytes, consistent with impaired PI3Kα function and

with the lipodystrophy and insulin resistance seen in SHORT syndrome (Huang-Doran *et al*,

2016). To assess the effect of the APDS2 ΔEx11 in the same, non-immunological, cell

context, we used lentiviral vectors, as previously described (Huang-Doran *et al.*, 2016), to

generate clonal 3T3-L1 cell lines allowing conditional, tuneable overexpression of PIK3R1

variants in response to doxycycline. For signalling studies, we also generated cells

conditionally overexpressing the PROS- and cancer-associated PIK3CA H1047R mutation

as a positive control for increased PI3Kα signalling, while for differentiation studies we used

previously reported lines conditionally expressing PIK3R1 R649W or Y657X (Huang-Doran

et al., 2016).

In undifferentiated cells, we first confirmed doxycycline-dependent overexpression of p85α

or p110α transgenes (**Fig 2A**), before assessing basal and insulin-stimulated Akt

phosphorylation. As expected, overexpression of oncogenic H1047R p110α strongly

increased basal Akt phosphorylation (**Fig 2A-C, Fig EV3A,C**) with no additional increase on

insulin stimulation (**Fig 2A-C, Fig EV3B,D**). Surprisingly, however, not only did

overexpression of the APDS2 ΔEx11 p85α not increase basal PI3Kα signalling, but it

potently inhibited insulin-induced Akt phosphorylation, consistent with a strong dominant

negative action on pathway activation (**Fig 2A-C, Fig EV3**). Overexpressing wild-type p85α

mildly reduced insulin-stimulated Akt phosphorylation (**Fig 2A-C, Fig EV3B,C**), although WT

p85α was not expressed to as high a level as ΔEx11 p85α. In keeping with impaired PI3Kα

activity, overexpression of the ΔEx11 variant also severely impaired adipocyte differentiation,

assessed by triglyceride accumulation in response to a standard differentiation protocol (**Fig**

2D). A similar effect was seen on overexpressing SHORT syndrome variants R649W and

Y657X (**Fig 2D**).

To assess whether the inhibitory effect of ΔEx11 p85α might be an artefact caused by strong

overexpression, doxycycline titration was used to assess whether a low level of

overexpression might unmask signalling hyperactivation. However, no such hyperactivation

was seen, and instead, a graded diminution of insulin-induced AKT phosphorylation was

observed from the lowest to highest levels of p85α ΔEx11 overexpression (**Fig EV4**).

Effect of PIK3R1 mutations on Phosphoinositide 3-kinase activity in vitro

Given this evidence that APDS2-associated PIK3R1 ΔEx11 potently inhibits PI3Kα when

overexpressed in 3T3-L1 preadipocytes, we next sought to investigate the biochemical basis

of this observation. First, we assessed the effect of disease-causing PIK3R1 mutations on

basal and phosphotyrosine-stimulated activity of purified PI3Kα, β and δ holoenzyme in a

previously described reconstituted *in vitro* system(Burke *et al*, 2012). As well as the APDS2

ΔEx11 mutation, we selected three SHORT syndrome-associated mutations to study. These

were the most common causal mutation, R649W, which abolishes phosphotyrosine binding

by the C-terminal SH2 (cSH2) domain (Chudasama *et al.*, 2013), Y657X, which truncates

the cSH2 domain (Huang-Doran *et al.*, 2016; Kwok *et al*, 2020), and E489K which,

atypically, lies in the inter-SH2 domain where most cancer, overgrowth and APDS2-

associated mutations lie (Thauvin-Robinet *et al.*, 2013). PIK3R1 E489K-containing primary

cells were previously suggested to show basal hyperactivation (Thauvin-Robinet *et al.*,

2013).

Wild-type and SHORT syndrome mutant holoenzymes were successfully purified for *in vitro*

assay, but despite multiple attempts, ΔEx11 holoenzyme could only be made in small

amounts under identical conditions, and moreover was unstable on storage, precluding

further study. In keeping with previous reports (Chudasama *et al.*, 2013; Dornan *et al.*, 2017;

Dornan *et al*, 2020), p85α R649W showed severely reduced phosphotyrosine-stimulated

activity in complex with p110α, with highly significant but lesser loss of function seen for

Y657X and E489K (**Fig 3A, Fig EV5A**). No increase in basal activity was seen for any

variant.

We also assessed whether any of the SHORT syndrome variants affect function of PI3Kβ

and found that all variants impaired phosphotyrosine-stimulated activity of PI3Kβ. Again, this

impairment was less for p85α Y657X than for R649W, and in this case only very mild for

E489K (**Fig 3B, Fig EV5B**). For PI3Kδ, p85α R649W conferred severe loss of function, as

for other isoforms (**Fig 3C, Fig EV5C**), suggesting that the absence of immunodeficiency in

SHORT syndrome is not accounted for by selective inhibition of PI3Kα function by causal

mutations.

Binding of p110α by mutant p85α

Given the potent dominant negativity of APDS2-related p85α ΔEx11 in cells, and the

instability of p85α ΔEx11-containing PI3K holoenzyme *in vitro*, we next used

immunoprecipitation to assess binding of p110α by this and other p85α variants in cells.

p85α was easily detected in anti-p110α immunoprecipitates from all cell lines at baseline

(**Fig 4**). No increase was seen on conditional overexpression of wild-type or R649W p85α.

Given the known instability of monomeric p110α, this suggests that all p110α is bound to

p85α before overexpression. Although wild-type endogenous p85α may have been replaced

by heterologously overexpressed p85α in these cells, this could not be detected without a

size difference of the variant from wild-type. For cells overexpressing the SHORT syndrome-

associated p85α Y657X, the truncated variant was strongly co-immunoprecipitated,

accounting for nearly all of the p85α signal in anti-p110α immunoprecipitate. This

demonstrates preserved binding of p110α by mutant p85α (**Fig 4**). In sharp contrast,

although truncated p85α ΔEx11 was easily detected in cell lysates before

immunoprecipitation and in supernatant after immunoprecipitation (arrowed in **Fig 4**), no

truncated p85α ΔEx11 was seen in p110α immunoprecipitates, and no change in p110α

expression was detected (**Fig 4**). This suggests that this truncated APDS2 causal variant

does not supplant endogenous, full-length p85α binding to p110α, despite overexpression.

This argues against destabilisation of p110α as the mechanism of the observed dominant

negative activity.

Effect of PIK3R1 mutations on insulin-induced PI3K recruitment to IRS1/2

As APDS2 p85α ΔEx11 does not appear to displace wild-type p85α from p110α, despite

strong overexpression, it is likely that there are high levels of the monomeric truncated p85α

in the cell. This may be important as it is p85α that mediates recruitment of PI3K to activated

tyrosine kinase receptors and their tyrosine phosphorylated substrates, including the insulin-

receptor substrate proteins Irs1 and Irs2. Excess free regulatory subunits compete with

heterodimeric PI3K holenzyme (Ueki *et al.*, 2002), raising the possibility that excess free,

truncated APDS2 p85α ΔEx11 may exert its inhibitory action by competing with PI3K

holenzyme.

To assess this possibility, we again used the 3T3-L1 cellular model to determine whether

overexpression of disease-causing p85α variants impairs recruitment of p110α to Irs1 and

Irs2. Irs1 was immunoprecipitated with or without conditional p85α overexpression and with

or without insulin stimulation. Overexpression of wild-type p85α had no effect on basal or

insulin-induced p110α recruitment to Irs1 (**Fig 5A,B, Fig EV6A,B**). In contrast,

overexpression of either p85α R649W or Y657X sharply reduced insulin-stimulated p110α

recruitment to Irs1 on insulin stimulation (**Fig 5A,B, Fig EV6A,B**). In keeping with the

inability of the R649W cSH2 domain to bind phosphotyrosines, p85α R649W was also not

recruited to Irs1, while overexpression of Y657X increased stimulated but not basal p85α

recruitment (**Fig 5A,C, Fig EV6C,D**). This suggests a pure defect in PI3K holoenzyme

recruitment for R649W. For Y657X the signalling defect may have mixed mechanisms, with

reduced activation by phosphotyrosine seen in *in vitro* studies coupled to increased

abundance of monomeric mutant p85α, leading to recruitment of non p110α-bound mutant

p85α to Irs1.

In keeping with our finding of severely attenuated insulin signalling upon p85α ΔEx11

overexpression (**Fig 2**), overexpression of this mutant p85α abolished p110α recruitment to

Irs1 (**Fig 5A,B, Fig EV6A,B**). However, non p110α-bound p85α ΔEx11 was strongly

recruited to Irs1 even in the absence of insulin stimulation (**Fig 5A,C, Fig EV6C,D**). This

suggests that although p85α ΔEx11 does not effectively compete with wild-type p85α for

binding to p110α, it has preserved or possibly enhanced ability to bind Irs1. This gives the

mutant p85α properties that render it a more potent endogenous inhibitor of PI3K signalling

than free wild-type p85α. Experiments conducted with immunoprecipitation of Irs2 instead of

Irs1 yielded closely similar findings for all p85α species (**Fig EV6 and EV7**).

Discussion

Murine genetic studies of Pik3r1 have proved complicated to interpret, due in part to

functional redundancy among PI3K regulatory subunits, and in part to unanticipated effects

of perturbing PI3K subunit stoichiometry. This has left several important questions about *in*

vivo functions of Pik3r1 unresolved. Identification of a series of human genetic disorders

caused by constitutional PIK3R1 mutations over the past 10 years has given fresh impetus

to the field and has been mechanistically illuminating.

Homozygous truncating PIK3R1 mutations abolishing p85α expression while preserving

p55α and p50α produce agammaglobulinaemia (Conley *et al*, 2012; Tang *et al*, 2018). This

resembles the immunodeficiency reported in *Pik3r1* knockout mice (Fruman *et al*, 1999;

Suzuki *et al*, 1999) and suggests an essential, non-redundant function of p85α in B cell

development in humans. Thereafter, human genetics has provided more novel insights.

PIK3R1 mutations were identified in SHORT syndrome in 2013 (Chudasama *et al.*, 2013;

Dyment *et al.*, 2013; Thauvin-Robinet *et al.*, 2013), nearly all in the C-terminal SH2 domain,

which together with the N-terminal SH2 domain enables PI3K recruitment to activated RTKs

(Rordorf-Nikolic *et al*, 1995). SHORT syndrome features short stature and insulin resistance

consistent with impaired ligand-induced p110α action, a phenotype distinct from the

enhanced insulin sensitivity produced by genetic ablation of one or more *Pik3r1* products in

mice (Chen *et al.*, 2004; Fruman *et al.*, 2000; Mauvais-Jarvis *et al.*, 2002). Mice with SHORT

syndrome mutations knocked in were generated after human findings, and faithfully

reproduce the human phenotype (Kwok *et al.*, 2020; Solheim *et al*, 2018; Winnay *et al*,

2016), confirming that expression of a signalling-impaired PIK3R1 has different

consequences to Pik3r1 knockout. No immunodeficiency has been described associated

with classic SHORT syndrome mutations, however, suggesting a selective effect on PI3Kα,

but the basis of this selectivity has not previously been investigated.

In contrast to SHORT syndrome, mutations in the inter SH2 domain of PIK3R1, mostly

leading to skipping of exon 11, were shown in 2014 to activate PI3K *in vitro* and to cause

immunodeficiency (APDS2) (Deau *et al.*, 2014; Lucas *et al.*, 2014b) similar to that caused by

activating mutations in p110δ (APDS1) (Angulo *et al.*, 2013; Lucas *et al.*, 2014a). Neither

overgrowth nor metabolic features of APDS2 have been described to suggest p110α

hyperactivation, and indeed short stature is common in APDS2 (Elkaim *et al.*, 2016; Jamee

et al., 2020; Maccari *et al.*, 2023; Olbrich *et al.*, 2016; Petrovski *et al.*, 2016), with a growing

number of APDS2 patients described with features of SHORT syndrome (Bravo Garcia-

Morato *et al.*, 2017; Maccari *et al.*, 2023; Nguyen *et al.*, 2023; Petrovski *et al.*, 2016;

Ramirez *et al.*, 2020; Szczawinska-Poplonyk *et al.*, 2022). Moreover mice with the common

APDS2 causal Pik3r1 variant knocked in show impaired growth and *in utero* survival, unlike

APDS2 murine models(Nguyen *et al.*, 2023).

Thus, study of distinct PIK3R1-related syndromes shows that impaired signalling caused by

PIK3R1 mutations produces phenotypes related to impaired PI3Kα function, without

affecting PI3Kδ. Conversely, mutations of PIK3R1 that produce phenotypes related to

increased PI3Kδ signalling do not increase PI3Kα signalling. Indeed, they surprisingly have

features characteristic of impaired PI3Kα function.

Lack of overgrowth in APDS2 has been attributed to greater ability of APDS2 PIK3R1

variants to activate p110δ than p110α (Dornan *et al.*, 2017)*,* but the co-ocurrence of gain-

and loss- of function phenotypes has not been explained to date. Our findings suggest that

the explanation may lie in competing effects of APDS2 PIK3R1 variants to activate PI3Kδ on

one hand, as previously shown (Angulo *et al.*, 2013; Dornan *et al.*, 2017; Lucas *et al.*,

2014b), while on the other hand interfering with PI3Kα through the dominant negative effect

of non p110α-bound mutant p85α.

The low expression of truncated p85α ΔEx11 we described in dermal fibroblasts is similar to

observations made in lymphocytes, however, in lymphocytes this is associated with

increased basal AKT phosphorylation (Deau *et al.*, 2014; Lucas *et al.*, 2014b) that is

abolished by p110δ inhibition (Lucas *et al.*, 2014b). P110δ is also expressed in fibroblasts,

but protein levels were reduced in the cells studied, consistent with the previously reported

observation that when expressed in insect cells the PI3K holoenzyme containing

p85α ΔEx11 has a low yield and is unstable (Dornan *et al.*, 2017). We speculate that in cells

with low endogenous p110δ protein expression, the destabilising effect of mutant PIK3R1

predominates over its activating effect.

The balance between expression and signalling may be a fine one, however, as transient

overexpression of FLAG-tagged p85α ΔEx11 did increase AKT phosphorylation in 3T3

fibroblasts in a previous study, although expression of p110α, β and δ was not determined

(Deau *et al.*, 2014). We find, in contrast, that overexpression of untagged p85α ΔEx11 has a

strong inhibitory effect on insulin signalling, and we were unable to identify a window of

overexpression where increased AKT phosphorylation could be observed. We further

demonstrated that most or all of the mutant p85α expressed is not bound to p110α, while

unbound mutant p85α still binds to Irs1/2, effectively competing with PI3K holoenzyme. The

observation that p85α ΔEx11 can associate with Irs1/2 is in agreement with reports that

p85α ΔEx11-containing PI3Kα and PI3Kδ can be stimulated by pY2-peptides (Dornan *et al.*,

2017; Dornan *et al.*, 2020) and that p85α ΔEx11 is recruited to tyrosine-phosphorylated LAT

in T cells(Lucas *et al.*, 2014b). The competition we suggest between unbound mutant p85α

and PI3K holoenzyme for binding to the activated RTKs is in keeping with longstanding

evidence that free p85 downregulates PI3K signalling through the same competitive

mechanism (Thorpe *et al*, 2017; Ueki *et al.*, 2002), with a single study suggesting in addition

that overexpression of tagged p85α leads to Irs1 sequestration with free p85α in cytosolic

foci where PI(3,4,5)P3 production does not occur (Luo *et al*, 2005).

The current study has limitations. We have studied primary cells from only a single APDS2

patient, and in the 3T3-L1 cell model, we did not determine whether p110δ protein could be

detected. If not, this could explain the lack of detectable AKT phosphorylation with induction

of Pik3r1 ΔEx11. Our study moreover raises further questions. Full length p85α subunits can

homodimerize (Cheung *et al*, 2015; Harpur *et al*, 1999; LoPiccolo *et al*, 2015), and it is

speculated that homodimers may outcompete p85α/p110 heterodimers for binding to

activated Irs1 due to configuration of the four SH2 domains. In the 3T3-L1 preadipocyte

APDS2 models, p85α ΔEx11 expression was high despite impaired p110α association, but

whether this is composed of monomeric or homodimeric p85α, and whether mutant p85α

expression leads to sequestration of Irs1 remote from the insulin receptor, as previously

suggested for tagged wild-type p85α, is undetermined.

In summary, it is already established that: A. genetic activation of PIK3CD causes

immunodeficiency without disordered growth, while B. inhibition of PIK3R1 recruitment to

RTKs and their substrates impairs growth and insulin action, without immunodeficiency,

despite all catalytic subunits being affected and C. loss of p85α alone causes

immunodeficiency. The current study, coupled with prior reports, suggests that the common

APDS2 mutation in PIK3R1 has mixed consequences, producing greater hyperactivation of

p110δ than p110α, based on subtle differences in the inhibitory interactions of regulatory

and catalytic subunits, while also destabilising PI3K holoenzyme and exerting dominant

negative activity on wild-type PIK3R1 function. We suggest that these competing activating

and inhibitory consequences are finely balanced, potentially differing among tissues, leading

to mixed clinical profiles of gain- and loss-of function features.

Materials and Methods

Ethics

The patient studied gave written informed consent for NIH IRB-approved research protocols

06-I-0015 and 09-I-0133 and was reported previously(Lucas *et al.*, 2014b). Dermal

fibroblasts were isolated from a skin punch biopsy and cultured as previously

described(Huang-Doran *et al.*, 2016).

Baculovirus Generation

p85α point mutation expression constructs were created by site-directed mutagenesis of a

pACEBac1_Homo sapiens p85α plasmid using a QuikChange II XL Site-Directed

Mutagenesis Kit. In-Fusion cloning was used to generate pACEBac1_p85α ΔEx11. Purified

plasmid inserts were sequenced and verified by restriction enzyme digest, exploiting loss of

a DpnI site within excised exon 11.

pACEBac1_p85α and pFastBac™HT B plasmids encoding N terminally tandem His-tagged

p110 subunits were used to generate bacmid DNA. Bacmids were expanded using MAX

Efficiency DH10Bac™ Competent or EMBacY cells and 2-4μg was transfected into Sf9 cells

using Insect-XPRESS (Promega E2311). Pooled virus from low titre supernatant (P1 stock)

was used to produce high titre P2 stock by transfection of 1.5×10^6 Sf9 cells/mL in 450mL

using. For catalytic subunits, 1-2mL P2 stock was used for a further round of Sf9 transfection

and expansion to generate P3 stock.

Purification of PI3K holoenzymes

1.5-2L of 1.5×10^6 Sf9 cells/mL were co-infected with 18mL P3 catalytic subunit and 4mL P2

regulatory subunit baculovirus. Cells were cultured at 27°C, harvested 48 h post-infection,

pelleted, washed and stored at -80°C. Pellets were later lysed by sonication before

ultracentrifugation. p85α/p110 heterodimers were pulled down via the tandem p110 N-

terminal His tags, preventing purification of monomeric p85α, using tandem Ni^{2+}HisTrap Fast

Flow columns. Further purification utilised a heparin HiTrap Q HP column. Eluted fractions

were concentrated to ≥1mg/mL and 1mL concentrated fractions were gel filtered on

Superdex 200 16/60 columns. The p110 tandem His tag was retained for proteins used in

functional analyses. Purity of eluted complexes was verified by SDS-PAGE, and purified

proteins were quantified using a Nanodrop 1000 Spectrophotometer (Thermo Scientific),

with concentrations calculated using the molecular extinction coefficient (assuming full

cysteine reduction) determined by sequence input to the ProtParam tool (ExPASy). Total

injection volume onto gel filtration columns was kept at 1 mL. Yields of PI3K complexes were

determined using the area under the curve (280nm mAU per mL eluted protein) from gel

filtration chromatograms, and normalised to the volume of Sf9 cells infected.

Liposome preparation

Chloroform/methanol solutions of sphingomyelin, cholesterol, porcine brain

phosphatidylcholine, porcine brain phosphatidylethanolamine, porcine brain

phosphatidylserine and porcine brain $PI(4,5)P_2$ were mixed to generate a preparation

containing 5/10/15/45/20/5% of each lipid, with a total lipid concentration of 5 mg/mL and

final $PI(4,5)P_2$ concentration 250µg/mL. Lipid preparations were dessicated under argon and

then in a vacuum desiccator before rehydrated in buffer containing 20mM HEPES (pH 7.5),

100 mM KCl and 1 mM EGTA (pH 8.0). Unilamellar vesicles were generated by extrusion

through polycarbonate filters with 100nm pores, using a glass-tight syringe.

Fluorescence Polarisation Assay

$PI(3,4,5)P_3$ production was measured by modified PI3-Kinase activity fluorescence

polarisation assay (Echelon Biosciences, Salt Lake City, UT, USA) in 384-well black

microtitre plates using 1mM liposomes containing 50µM $PI(4,5)P_2$, optimised concentrations

of purified PI3K proteins, 100µM ATP, 2mM $MgCl_2$, with or without 1µM tyrosine

bisphosphorylated peptide derived from PDGFRβ ("pY2"). Reactions were quenched before

addition of TAMRA Red Fluorescent Probe in HNT buffer. Identity of the lipid group coupled

to the TAMRA probe was not disclosed by Echelon Biosciences, but as Grp1 recognises

specific lipid head groups in competition with $PI(3,4,5)P_3$, it is likely to be a lipid with a similar

head group such as inositol 1,3,4,5-tetrakisphosphate. After 1h fluorescence polarisation

was measured using a PHERAstar spectrofluorometer (BMG Labtech, Ortenberg,

Germany). Concentrations of wild-type protein complexes used in the assay were: 20-40nM

p85α/p110α, 80-160nM p85α/p110β and 80-160nM p85α/p110δ for basal activities, and 1-

2nM p85α/p110α, 2-4 nM p85α/p110β and 2-6nM p85α/p110δ for pY2-stimulated activities.

Concentrations of thawed protein aliquots were assessed as 280 nm absorbance by

Nanodrop 1000 Spectrophotometer (Thermo Scientific), and samples of preparations used

in assays were resolved by SDS-PAGE to confirm equal amounts of each complex. Three

technical replicates were used per assay on at least three different occasions. Relevant wild-

type p85α/p110 catalytic isoforms were included in each assay as controls. Other controls

are detailed in the Appendix. Standard curves of diC8-PI(3,4,5)P_3 in the presence of

liposomes were included in every experiment and used to infer PIP_3 generated by purified

PI3K. Standard curves for each experiment provided the linear range within which PIP_3

could be accurately determined. Quadruplicate interpolated PIP_3 concentrations were

averaged and normalised by reaction time (30 min) and enzyme concentration in pM to yield

PI3K activity in pmol PIP_3/min/pM enzyme. Specific activities were then further normalised to

activity of p85α wild-type controls in the presence of pY2 for each experiment.

Generation of 3T3-L1 cells conditionally expressing p85α or p110α

3T3-L1 preadipocytes from Zenbio were cultured, differentiated and stained for neutral lipid

with Oil Red O as previously described(Huang-Doran *et al*, 2011). Generation of 3T3-L1

sublines inducibly expressing WT, Y657X, or R649W p85α was also previously

reported(Huang-Doran *et al.*, 2016; Hussain *et al*, 2011). The 3T3-L1 lines inducibly

expressing p85α ΔEx11 or p110α H1047R were generated using essentially the same

procedure, starting with In-Fusion subcloning of the p85α ΔEx11 cDNA insert from the

pACEBac1 p85α ΔEx11 plasmid described above, and of a PIK3CA H1047R cDNA insert

derived by PCR from a pcDNA5.FRT.Teton_FLAG_PIK3CA-H1047R plasmid, into the

pEN_Tmcs entry vector. Transgene expression was induced with 1 μg/mL doxycycline for 72

hours.

Insulin signalling studies

3T3-L1 cells serum starved in DMEM containing 0.5 % BSA for 16h were stimulated with

100nM Actrapid insulin (Novo Nordisk) for 10min. Monolayers were snap frozen in liquid N2

and stored at -80°C. 200-600μL co-IP lysis or RIPA buffer was added to frozen before

scraping of lysate into pre-chilled tubes, incubation on ice for 30min and clearing by

centrifugation. Protein was quantified using the Bio-Rad DC assay.

Lysates mixed with NuPAGE® SDS Loading bufferwith 5% β-mercaptoethanol were boiled

before SDS-PAGE, with 10μg boiled lysate used to represent co-immunoprecipitation (Co-

IP) input. For Co-IP, lysates were incubated with immunoprecipitating antibody overnight at

4°C. before incubation for 2h with Protein G Dynabeads®, followed by centrifugation and

bead removal using a DynaMag™-2 Magnet. Supernatants mixed 1:1 with Co-IP Elution

Buffer were boiled for 10 min. Protein was eluted from washed beads and boiled for 10min.

Input, supernatant and co-IP samples underwent SDS-PAGE using NuPAGE® 4-12%

gradient Bis-Tris gels in NuPAGE® MOPS SDS Running Buffer. Transfer to nitrocellulose

was performed using the iBlot™ Dry Blotting System (Invitrogen), with preincubation in

Equilibration Transfer Buffer for proteins above 150kDa.

For immunoblotting, blocked membranes were incubated overnight at 4°C in primary

antibody, washed and incubated with HRP-linked anti-rabbit or anti-mouse IgG secondary

antibody diluted 1:5000 in blocking buffer. Proteins were visualized using the ChemiDoc™

MP System (Bio-Rad) and band intensities quantified using Image Lab software (Bio-Rad).

Statistical Analysis

For quantitative data, all biological replicates (i.e. results of independent experiments on

different days) are represented in figures, with paired points (e.g. with/without insulin in the

same experiment) connected by lines. For fluorescence polarisation assays, each biological

replicate shown is the mean of 3 technical replicates. Sample size for individual experiments

was not pre-determined. To avoid the pitfalls of dichotomous significance testing on low-

throughput biological datasets, we used estimation statistics (Data Analysis using Bootstrap-

Coupled ESTimation) with default settings (5000 resamples, BCa boostrap)(Ho *et al*, 2019).

This focuses on effect sizes and derives confidence intervals derived from bootstrapping for

differences in means; the small (but typical) sample size of the experiments analysed limits

reliable bootstrapping, but it offers additional confidence to the consistent patterns seen in

independent replicates. We have indicated significance in main text figures with an asterisk,

and show the 95% confidence intervals for mean differences in extended view figures.

Expanded details of methods, including sources and concentrations of antibodies, and buffer

compositions, are listed in the Appendix.

Data Availability

This study includes no data deposited in external repositories. Uncut Western blot and other

source images are available on the Open Science Framework project webpage:

https://osf.io/eyc4h/?view_only=9af8645c8fe5408790ce19920d0084c6.

Acknowledgements

This work was supported by the Wellcome Trust through a grant to RKS [210752/Z/18/Z]

and a studentship to PRT [102356/Z/13/Z]. Additional support was from the UK Medical

Research Council (MRC) [MC_UU_12012/5 and MC_U105184308 (to RLW)] and the

Intramural Research Program of the National Institute of Allergy and Infectious Diseases,

National Institutes of Health (to HCS). We thank Dr. Koneti Rao, Debra Long-Priel, and

Angela Wang for clinical, technical, and regulatory assistance.

Author Contributions

PRT: conceptualisation, methodology, formal analysis, investigation, visualization, writing -

original draft. **RK**: formal analysis, investigation, visualization. **OP**: supervision,

methodology, investigation, writing - review & editing. **HCS:** resources, funding acquisition,

writing - review & editing. **GVB:** supervision, methodology, writing - review & editing. **RLW**:

supervision, methodology, investigation, funding acquisition, writing - review & editing. **RKS:**

conceptualisation, supervision, methodology, funding acquisition, writing - original draft,

writing - review & editing, funding acquisition.

Disclosure and competing interests statement

RKS has consulted for Novartis on clinical aspects of PIK3CA-related overgrowth, and for

Alnylam, Amryt and AstraZeneca on clinical aspects of monogenic insulin resistance and

lipodystrophy

The Paper Explained

PROBLEM: The *PIK3R1* gene encodes a key component of phosphoinositide 3-kinase

(PI3K), which is crucial for actions of insulin and growth factors on many cells. Mutations in

PIK3R1 cause several different genetic diseases. One of these is SHORT syndrome, which

includes short stature and insulin resistance. This is caused by PIK3R1 mutations that impair

PI3K signalling, especially through the PI3Kα form of PI3K. Another PIK3R1-related disease

is Activated PI3K δ syndrome 2 (APDS2), a form of immunodeficiency caused by PIK3R1

mutations that activate PI3K signalling, especially involving the PI3Kδ form of PI3K.

Surprisingly, many patients with APDS2 have short stature or frank SHORT syndrome,

however. How these signs of loss of PI3K function are caused by an activating PIK3R1

mutation is unknown. We set out to investigate this.

RESULTS: We discovered that skin cells from a person with APDS2 had very low levels of

the protein that binds to PIK3R1 to make PI3Kδ, and low levels of insulin-triggered PI3K

activity. The PIK3R1 variant that causes APDS2, previously convincingly shown to activate

PI3Kδ, made only unstable PI3K. When we forced expression of disease-causing PIK3R1

variants in mouse fat cell precursors, we found that the APDS2 variant inhibited insulin-

induced PI3K even more than variants that cause SHORT syndrome. This was because the

APDS2 variant did not make stable PI3Kα, and outcompeted normal PI3Kα for binding to

insulin receptor target proteins.

IMPACT: These findings demonstrate that a PIK3R1 mutation that was previously shown to

cause immunodeficiency by activating PI3Kδ also ablates insulin signalling by indirectly

inhibiting PI3Kα activity. We suggest that this explains the mixed PI3K gain- and loss-of

function features that co-exist in APDS2. Our findings imply that PI3Kδ inhibition alone may

not be sufficient to reverse all components of APDS2, and that additional therapeutic

strategies targeting the mutant PI3KR1 products may be warranted.

For More Information

SHORT syndrome (OMIM): https://omim.org/entry/269880

APDS2 (OMIM): https://omim.org/entry/616005

Semple Lab: https://www.ed.ac.uk/mrc-human-genetics-unit/research/robert-semple-group

References

Angulo I, Vadas O, Garcon F, Banham-Hall E, Plagnol V, Leahy TR, Baxendale H, Coulter T,
Curtis J, Wu C *et al* (2013) Phosphoinositide 3-kinase delta gene mutation predisposes to
respiratory infection and airway damage. *Science* 342: 866-871
Avila M, Dyment DA, Sagen JV, St-Onge J, Moog U, Chung BHY, Mo S, Mansour S, Albanese
A, Garcia S *et al* (2016) Clinical reappraisal of SHORT syndrome with PIK3R1 mutations:
toward recommendation for molecular testing and management. *Clin Genet* 89: 501-506
Brachmann SM, Ueki K, Engelman JA, Kahn RC, Cantley LC (2005) Phosphoinositide 3-kinase
catalytic subunit deletion and regulatory subunit deletion have opposite effects on insulin
sensitivity in mice. *Mol Cell Biol* 25: 1596-1607
Bravo Garcia-Morato M, Garcia-Minaur S, Molina Garicano J, Santos Simarro F, Del Pino
Molina L, Lopez-Granados E, Ferreira Cerdan A, Rodriguez Pena R (2017) Mutations in
PIK3R1 can lead to APDS2, SHORT syndrome or a combination of the two. *Clin Immunol* 179:
77-80
Burke JE, Perisic O, Masson GR, Vadas O, Williams RL (2012) Oncogenic mutations mimic
and enhance dynamic events in the natural activation of phosphoinositide 3-kinase
p110alpha (PIK3CA). *Proc Natl Acad Sci U S A* 109: 15259-15264
Chen D, Mauvais-Jarvis F, Bluher M, Fisher SJ, Jozsi A, Goodyear LJ, Ueki K, Kahn CR (2004)
p50alpha/p55alpha phosphoinositide 3-kinase knockout mice exhibit enhanced insulin
sensitivity. *Mol Cell Biol* 24: 320-329
Cheung LW, Walkiewicz KW, Besong TM, Guo H, Hawke DH, Arold ST, Mills GB (2015)
Regulation of the PI3K pathway through a p85alpha monomer-homodimer equilibrium. *Elife*
4: e06866
Chudasama KK, Winnay J, Johansson S, Claudi T, Konig R, Haldorsen I, Johansson B, Woo JR,
Aarskog D, Sagen JV *et al* (2013) SHORT syndrome with partial lipodystrophy due to
impaired phosphatidylinositol 3 kinase signaling. *Am J Hum Genet* 93: 150-157
Conley ME, Dobbs AK, Quintana AM, Bosompem A, Wang YD, Coustan-Smith E, Smith AM,
Perez EE, Murray PJ (2012) Agammaglobulinemia and absent B lineage cells in a patient
lacking the p85alpha subunit of PI3K. *J Exp Med* 209: 463-470

Cottrell CE, Bender NR, Zimmermann MT, Heusel JW, Corliss M, Evenson MJ, Magrini V,
Corsmeier DJ, Avenarius M, Dudley JN *et al* (2021) Somatic PIK3R1 variation as a cause of
vascular malformations and overgrowth. *Genet Med*
Deau MC, Heurtier L, Frange P, Suarez F, Bole-Feysot C, Nitschke P, Cavazzana M, Picard C,
Durandy A, Fischer A *et al* (2014) A human immunodeficiency caused by mutations in the
PIK3R1 gene. *J Clin Invest* 124: 3923-3928
Dornan GL, Siempelkamp BD, Jenkins ML, Vadas O, Lucas CL, Burke JE (2017)
Conformational disruption of PI3Kdelta regulation by immunodeficiency mutations in
PIK3CD and PIK3R1. *Proc Natl Acad Sci U S A* 114: 1982-1987
Dornan GL, Stariha JTB, Rathinaswamy MK, Powell CJ, Boulanger MJ, Burke JE (2020)
Defining How Oncogenic and Developmental Mutations of PIK3R1 Alter the Regulation of
Class IA Phosphoinositide 3-Kinases. *Structure* 28: 145-156 e145
Dyment DA, Smith AC, Alcantara D, Schwartzentruber JA, Basel-Vanagaite L, Curry CJ,
Temple IK, Reardon W, Mansour S, Haq MR *et al* (2013) Mutations in PIK3R1 cause SHORT
syndrome. *Am J Hum Genet* 93: 158-166
Elkaim E, Neven B, Bruneau J, Mitsui-Sekinaka K, Stanislas A, Heurtier L, Lucas CL, Matthews
H, Deau MC, Sharapova S *et al* (2016) Clinical and immunologic phenotype associated with
activated phosphoinositide 3-kinase delta syndrome 2: A cohort study. *J Allergy Clin*
Immunol 138: 210-218 e219
Fruman DA, Chiu H, Hopkins BD, Bagrodia S, Cantley LC, Abraham RT (2017) The PI3K
Pathway in Human Disease. *Cell* 170: 605-635
Fruman DA, Mauvais-Jarvis F, Pollard DA, Yballe CM, Brazil D, Bronson RT, Kahn CR, Cantley
LC (2000) Hypoglycaemia, liver necrosis and perinatal death in mice lacking all isoforms of
phosphoinositide 3-kinase p85 alpha. *Nat Genet* 26: 379-382
Fruman DA, Snapper SB, Yballe CM, Davidson L, Yu JY, Alt FW, Cantley LC (1999) Impaired B
cell development and proliferation in absence of phosphoinositide 3-kinase p85alpha.
Science 283: 393-397
Geering B, Cutillas PR, Nock G, Gharbi SI, Vanhaesebroeck B (2007) Class IA
phosphoinositide 3-kinases are obligate p85-p110 heterodimers. *Proc Natl Acad Sci U S A*
104: 7809-7814
Harpur AG, Layton MJ, Das P, Bottomley MJ, Panayotou G, Driscoll PC, Waterfield MD (1999)
Intermolecular interactions of the p85alpha regulatory subunit of phosphatidylinositol 3-
kinase. *J Biol Chem* 274: 12323-12332
Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A (2019) Moving beyond P values: data
analysis with estimation graphics. *Nat Methods* 16: 565-566
Huang-Doran I, Bicknell LS, Finucane FM, Rocha N, Porter KM, Tung YC, Szekeres F, Krook A,
Nolan JJ, O'Driscoll M *et al* (2011) Genetic defects in human pericentrin are associated with
severe insulin resistance and diabetes. *Diabetes* 60: 925-935
Huang-Doran I, Tomlinson P, Payne F, Gast A, Sleigh A, Bottomley W, Harris J, Daly A, Rocha
N, Rudge S *et al* (2016) Insulin resistance uncoupled from dyslipidemia due to C-terminal
PIK3R1 mutations. *JCI Insight* 1: e88766
Hussain K, Challis B, Rocha N, Payne F, Minic M, Thompson A, Daly A, Scott C, Harris J,
Smillie BJ *et al* (2011) An activating mutation of AKT2 and human hypoglycemia. *Science*
334: 474
Jamee M, Moniri S, Zaki-Dizaji M, Olbrich P, Yazdani R, Jadidi-Niaragh F, Aghamahdi F,
Abolhassani H, Condliffe AM, Aghamohammadi A *et al* (2020) Clinical, Immunological, and

Genetic Features in Patients with Activated PI3Kdelta Syndrome (APDS): a Systematic Review. *Clin Rev Allergy Immunol* 59: 323-333

Kwok A, Zvetkova I, Virtue S, Luijten I, Huang-Doran I, Tomlinson P, Bulger DA, West J, Murfitt S, Griffin J *et al* (2020) Truncation of Pik3r1 causes severe insulin resistance uncoupled from obesity and dyslipidaemia by increased energy expenditure. *Mol Metab* 40: 101020

Lindhurst MJ, Parker VE, Payne F, Sapp JC, Rudge S, Harris J, Witkowski AM, Zhang Q, Groeneveld MP, Scott CE *et al* (2012) Mosaic overgrowth with fibroadipose hyperplasia is caused by somatic activating mutations in PIK3CA. *Nat Genet* 44: 928-933

LoPiccolo J, Kim SJ, Shi Y, Wu B, Wu H, Chait BT, Singer RH, Sali A, Brenowitz M, Bresnick AR *et al* (2015) Assembly and Molecular Architecture of the Phosphoinositide 3-Kinase p85alpha Homodimer. *J Biol Chem* 290: 30390-30405

Lucas CL, Kuehn HS, Zhao F, Niemela JE, Deenick EK, Palendira U, Avery DT, Moens L, Cannons JL, Biancalana M *et al* (2014a) Dominant-activating germline mutations in the gene encoding the PI(3)K catalytic subunit p110delta result in T cell senescence and human immunodeficiency. *Nat Immunol* 15: 88-97

Lucas CL, Zhang Y, Venida A, Wang Y, Hughes J, McElwee J, Butrick M, Matthews H, Price S, Biancalana M *et al* (2014b) Heterozygous splice mutation in PIK3R1 causes human immunodeficiency with lymphoproliferation due to dominant activation of PI3K. *J Exp Med* 211: 2537-2547

Luo J, Field SJ, Lee JY, Engelman JA, Cantley LC (2005) The p85 regulatory subunit of phosphoinositide 3-kinase down-regulates IRS-1 signaling via the formation of a sequestration complex. *J Cell Biol* 170: 455-464

Maccari ME, Wolkewitz M, Schwab C, Lorenzini T, Leiding JW, Aladjdi N, Abolhassani H, Abou-Chahla W, Aiuti A, Azarnoush S *et al* (2023) Activated phosphoinositide 3-kinase delta syndrome: Update from the ESID Registry and comparison with other autoimmune-lymphoproliferative inborn errors of immunity. *J Allergy Clin Immunol* 152: 984-996 e910

Madsen RR, Vanhaesebroeck B, Semple RK (2018) Cancer-Associated PIK3CA Mutations in Overgrowth Disorders. *Trends Mol Med* 24: 856-870

Mauvais-Jarvis F, Ueki K, Fruman DA, Hirshman MF, Sakamoto K, Goodyear LJ, Iannacone M, Accili D, Cantley LC, Kahn CR (2002) Reduced expression of the murine p85alpha subunit of phosphoinositide 3-kinase improves insulin signaling and ameliorates diabetes. *J Clin Invest* 109: 141-149

Nguyen T, Lau A, Bier J, Cooke KC, Lenthall H, Ruiz-Diaz S, Avery DT, Brigden H, Zahra D, Sewell WA *et al* (2023) Human PIK3R1 mutations disrupt lymphocyte differentiation to cause activated PI3Kdelta syndrome 2. *J Exp Med* 220

Olbrich P, Lorenz M, Cura Daball P, Lucena JM, Rensing-Ehl A, Sanchez B, Fuhrer M, Camacho-Lovillo M, Melon M, Schwarz K *et al* (2016) Activated PI3Kdelta syndrome type 2: Two patients, a novel mutation, and review of the literature. *Pediatr Allergy Immunol* 27: 640-644

Petrovski S, Parrott RE, Roberts JL, Huang H, Yang J, Gorentla B, Mousallem T, Wang E, Armstrong M, McHale D *et al* (2016) Dominant Splice Site Mutations in PIK3R1 Cause Hyper IgM Syndrome, Lymphadenopathy and Short Stature. *J Clin Immunol* 36: 462-471

Philp AJ, Campbell IG, Leet C, Vincan E, Rockman SP, Whitehead RH, Thomas RJ, Phillips WA (2001) The phosphatidylinositol 3'-kinase p85alpha gene is an oncogene in human ovarian and colon tumors. *Cancer Res* 61: 7426-7429

Ramirez L, Tamayo W, Ale H (2020) APDS2 and SHORT Syndrome in a Teenager with PIK3R1
Pathogenic Variant. *J Clin Immunol* 40: 1020-1025
Rordorf-Nikolic T, Van Horn DJ, Chen D, White MF, Backer JM (1995) Regulation of
phosphatidylinositol 3'-kinase by tyrosyl phosphoproteins. Full activation requires
occupancy of both SH2 domains in the 85-kDa regulatory subunit. *J Biol Chem* 270: 3662-
3666
Solheim MH, Winnay JN, Batista TM, Molven A, Njolstad PR, Kahn CR (2018) Mice Carrying a
Dominant-Negative Human PI3K Mutation Are Protected From Obesity and Hepatic
Steatosis but Not Diabetes. *Diabetes* 67: 1297-1309
Suzuki H, Terauchi Y, Fujiwara M, Aizawa S, Yazaki Y, Kadowaki T, Koyasu S (1999) Xid-like
immunodeficiency in mice with disruption of the p85alpha subunit of phosphoinositide 3-
kinase. *Science* 283: 390-392
Szczawinska-Poplonyk A, Bernat-Sitarz K, Schwartzmann E, Piechota M, Badura-Stronka M
(2022) Clinical and immunological assessment of APDS2 with features of the SHORT
syndrome related to a novel mutation in PIK3R1 with reduced penetrance. *Allergol*
Immunopathol (Madr) 50: 1-9
Tang P, Upton JEM, Barton-Forbes MA, Salvadori MI, Clynick MP, Price AK, Goobie SL (2018)
Autosomal Recessive Agammaglobulinemia Due to a Homozygous Mutation in PIK3R1. *J Clin*
Immunol 38: 88-95
Thauvin-Robinet C, Auclair M, Duplomb L, Caron-Debarle M, Avila M, St-Onge J, Le Merrer
M, Le Luyer B, Heron D, Mathieu-Dramard M *et al* (2013) PIK3R1 mutations cause syndromic
insulin resistance with lipoatrophy. *Am J Hum Genet* 93: 141-149
Thorpe LM, Spangle JM, Ohlson CE, Cheng H, Roberts TM, Cantley LC, Zhao JJ (2017) PI3K-
p110alpha mediates the oncogenic activity induced by loss of the novel tumor suppressor
PI3K-p85alpha. *Proc Natl Acad Sci U S A* 114: 7095-7100
Tsolakos N, Durrant TN, Chessa T, Suire SM, Oxley D, Kulkarni S, Downward J, Perisic O,
Williams RL, Stephens L *et al* (2018) Quantitation of class IA PI3Ks in mice reveals p110-free-
p85s and isoform-selective subunit associations and recruitment to receptors. *Proc Natl*
Acad Sci U S A 115: 12176-12181
Ueki K, Fruman DA, Brachmann SM, Tseng YH, Cantley LC, Kahn CR (2002) Molecular balance
between the regulatory and catalytic subunits of phosphoinositide 3-kinase regulates cell
signaling and survival. *Mol Cell Biol* 22: 965-977
Wentink M, Dalm V, Lankester AC, van Schouwenburg PA, Scholvinck L, Kalina T, Zachova R,
Sediva A, Lambeck A, Pico-Knijnenburg I *et al* (2017) Genetic defects in PI3Kdelta affect B-
cell differentiation and maturation leading to hypogammaglobulineamia and recurrent
infections. *Clin Immunol* 176: 77-86
Winnay JN, Solheim MH, Dirice E, Sakaguchi M, Noh HL, Kang HJ, Takahashi H, Chudasama
KK, Kim JK, Molven A *et al* (2016) PI3-kinase mutation linked to insulin and growth factor
resistance in vivo. *J Clin Invest* 126: 1401-1412
Yu J, Zhang Y, McIlroy J, Rordorf-Nikolic T, Orr GA, Backer JM (1998) Regulation of the
p85/p110 phosphatidylinositol 3'-kinase: stabilization and inhibition of the p110alpha
catalytic subunit by the p85 regulatory subunit. *Mol Cell Biol* 18: 1379-1387
Zhao JJ, Cheng H, Jia S, Wang L, Gjoerup OV, Mikami A, Roberts TM (2006) The p110alpha
isoform of PI3K is essential for proper growth factor signaling and oncogenic
transformation. *Proc Natl Acad Sci U S A* 103: 16296-16300
706

 Figure Legends

Figure 1. PI3K subunit expression and signalling in primary dermal fibroblasts.

Immunoblotting of AKT, AKT phosphorylated at threonine 308 (T308) or serine 473 (S473),

p85α, p110δ and p110α and are shown with and without stimulation by 100 nM insulin (Ins)

for 10 minutes. β-actin is shown as a loading control, with different amounts of pooled lysate

(Pool) used to demonstrate signal intensity in the linear range. Results from 4 healthy

controls (WT; 1-4), 1 patient with Activating p110 Delta Syndrome 2 (APDS2) due to the

p85α Δexon11 variant, and 3 patients with PIK3CA-Related Overgrowth Spectrum (PROS)

caused by the activating *PIK3CA* mutations indicated. (A) Immunoblots, with the truncated

p85α Δexon11 variant arrowed; (B)-(E) quantification of immunoblot bands from 3

independent experiments shown for phosphoAKT-S473, phosphoAKT-T308, p110δ and

p110α respectively. Paired datapoints +/- insulin are shown in (B) and (C), and dotted lines

mark means. Asterisks indicate a significant difference. More detailed statistical analysis

including 95% confidence intervals for the paired mean differences for these comparisons

are shown in Figure EV2.

Figure 2. Blunted insulin signaling in 3T3-L1 preadipocyte models of APDS2 and

SHORT syndrome. Immunoblotting of Akt, Akt phosphorylated at threonine 308 (T308) or

serine 473 (S473), p85α, and p110α and are shown with and without stimulation with 100

nM insulin (Ins) for 10 minutes. Cells were treated with doxycycline (Dox) 1μg/mL for 72

hours prior to insulin stimulation as indicated. (A) One immunoblot representing 3

experiments is shown. (B)-(C) Quantification of immunoblot bands from all 3 independent

experiments shown for phosphoAkt-S473 and phosphoAkt-T308 respectively. Paired

datapoints +/- insulin are shown, and dotted lines mark means. Asterisks indicate a

significant difference. More detailed statistical analysis including 95% confidence intervals

for the paired mean differences for these comparisons are shown in Figure EV3. (D)

Staining for neutral lipid with Oil Red O of 3T3-L1 cells at day 10 of adipocyte differentiation.

Induction of transgene expression by 1μg/mL doxycycline throughout differentiation is

shown. Images of entire plates are shown above, with representative bright field microscopy

images below.

Figure 3. SHORT syndrome p85α mutations impair phosphotyrosine-stimulated

phosphoinositide 3-kinase activity. Lipid kinase activity of purified recombinant PI3K

complexes generated using baculoviral expression in *Sf9* cells was measured using a

modified fluorescence polarisation assay. WT p85α or p85α SHORT syndrome mutations,

E489K, R649W or Y657X bound to either (A) p110α (B) p110β or (C) p110δ were assayed

for basal and bisphosphotyrosine (pY2)-stimulated lipid kinase activity. Dotted lines mark

means, and asterisks indicate a significant difference between the bisphosphotyrosine

(pY2)-stimulated state for WT and comparator mutant p85α. More detailed statistical

analysis including 95% confidence intervals for the paired mean differences for these

comparisons are shown in Figure EV5.

Figure 4. Ability of pathogenic p85α variants to bind p110α, assessed by co-

immunoprecipitation. Results of immunoblotting of anti-p110α immunoprecipitates from

3T3-L1 cells expressing wild-type, APDS2-associated or SHORT syndrome-associated

mutant p85α under the control of doxycycline (Dox) are shown. (A) One representative

immunoblot of immunoprecipitate, cell lysate prior to immunoprecipitation, and post

immunoprecipitation supernatant is shown. (B) Quantification of immunoblot bands from

immunoprecipitates from 3 independent experiments. Co-immunoprecipitated p85α is shown

normalized to immunoprecipitated p110α from all 3 independent experiments. Datapoints

from the same experiment +/- doxycline are connected by lines. No significant differences

were found among conditions.

Figure 5. Attenuated insulin-induced association of p110α with Irs1 in the presence of

APDS2 and SHORT syndrome mutant p85α. Results of immunoblotting of anti-Irs1

immunoprecipitates from 3T3-L1 cells expressing wild-type, APDS2-associated or SHORT

syndrome-associated mutant p85α under the control of doxycycline (Dox) are shown.

Treatment with 100nM insulin (Ins) is indicated. (A) One representative immunoblot of

immunoprecipitate, cell lysate prior to immunoprecipitation, and post immunoprecipitation

supernatant is shown. Two separate sets of gels, including independent wild-type controls

are shown on left and right. (B-C) Quantification of immunoblot bands from

immunoprecipitates from 3 independent experiments. Immunoprecipitated p110α is shown

normalized to immunoprecipitated Irs1 from all 3 independent experiments in (B), and

immunoprecipitated p85α similarly in (C). Datapoints from the same experiment +/- insulin

are connected by lines. Asterisks indicate significant differences induced by transgene

overexpression (i.e. plus versus minus doxycycline). More detailed statistical analysis

including 95% confidence intervals for the paired mean differences for these comparisons

are shown in Figure EV6.

                                    Expanded View Figure Legends

Figure EV1. Further characterisation of primary dermal fibroblasts studied. (A) Details

of cDNA sequence for *PIK3CA* and *PIK3R1* from cells derived from healthy controls (WT),

patients with APDS2 (p85α ΔEx11) or *PIK3CA*-related overgrowth syndrome (PROS),

confirming expected expression of mutant alleles. (B) Higher magnification detail of

immunoblot from wild-type and APDS2 fibroblasts showing truncated p85α Δex11. (C) All 3

immunoblot replicates for p110δ blots quantified in Figure 1, showing severely reduced

p110δ expression in the APDS2 cell line

Figure EV2. Full statistical analysis of data presented in main Figure 1 Analysis of

insulin-induced increase in (A) AKT S473/4 and (B) T308/9 phosphorylation. The paired

mean difference for 3-4 comparisons are shown in Cumming estimation plots. The raw data,

as presented in Figure 1, are re-plotted on the upper axes with paired observations

connected by a line. On the lower axes, paired mean differences are plotted as a bootstrap

sampling distribution. Mean differences are depicted as dots; 95% confidence intervals are

indicated by the ends of the vertical error bars. (C-D) Analysis of differences in (C) p110α

and (D) p110δ protein expression between healthy control cells and cells from PROS

patients harbouring activating PIK3CA mutations. Mean differences are shown in Gardner-

Altman estimation plots, with expression data plotted on the left axes and mean difference

on floating axes on the right, again as a bootstrap sampling distribution with mean difference

depicted as a dot and 95% confidence intervals by the ends of the vertical bar.

Figure EV3. Full statistical analysis of data presented in main Figure 2. Analysis of the

effects of doxycycline-induced expression of wild-type (WT) or ΔEx11 (APDS2) p85α, or of

p110α H1047R (PROS) on Akt S473/4 (A,B)   and T308/9 (C,D) phosphorylation.

Comparisons are made in both the basal, non-insulin stimulated state (A,C) and after

stimulation with 10 nmol/L insulin (B,D). Paired mean differences for 3 comparisons are

shown in Cumming estimation plots. Raw data, as presented in Figure 2, are re-plotted on

the upper axes with paired observations connected by a line. On the lower axes, paired

mean differences are plotted as a bootstrap sampling distribution. Mean differences are

depicted as dots; 95% confidence intervals are indicated by the ends of the vertical error

bars.

Figure EV4. The effect of graded expression of wild-type or disease-associated p85α

on 3T3-L1 preadipocytes. Immunoblots of p85α, phosphoAkt (S473) and total Akt are

shown for control 3T3-L1 cells and 3T3-L1 cells conditionally expressing wild-type (WT), or

APDS2-associated mutant p85α under the control of doxycycline (Dox), with and without 10

minutes of exposure to insulin as indicated. The filled black triangles indicate increasing

concentrations of doxycycline (from left to right: 0, 0.02, 0.03, 0.045, 0.065, or 0.1 µg/mL).

Exposure was for 72 hours in all cases. The truncated p85α variant can be seen below the

WT p85α for the APDS2 ΔEx11 mutant.

Figure EV5. Full statistical analysis of data presented in main Figure 3. Analysis of

fluorescence polarisation assay of phosphoinositide 3-kinase (PI3K) activity of *in vitro*

synthesised wild-type (WT) or mutant (E489K, R649W or Y657X) p85α. Results for p110α-

and p110β-containing PI3K are shown in (A) and (B) respectively. All data were acquired in

the presence of phosphotyrosine peptide. Paired mean differences for 3 comparisons are

shown in Cumming estimation plots. Raw data, as presented in Figure 3, are re-plotted on

the upper axes with paired observations connected by a line. On the lower axes, paired

mean differences are plotted as a bootstrap sampling distribution. Mean differences are

depicted as dots; 95% confidence intervals are indicated by the ends of the vertical error

bars. Results for the R649W p85α mutation only are shown with p110δ in (C). In this case

raw data are re-plotted on the left hand axes with paired observations connected by 3 nearly

superimposed lines. On the right hand axes, paired mean differences are plotted as a

bootstrap sampling distribution.

Figure EV6. Full statistical analysis of data presented in main Figure 5. Analysis of the

effects of doxycycline (dox)-induced expression of wild-type (WT), Y657X, R649W or ΔEx11

p85α on association of p110α (A,B)   and p85α (C,D) with Irs1. Results of co-

immunoprecipitation with (B,D) and without exposure to 10 nmol/L insulin Comparisons are

made in both the basal, non-insulin stimulated state (A,C) and after stimulation with 10

nmol/L insulin (B,D) are shown. Paired mean differences for 3 comparisons are shown in

Cumming estimation plots. Raw data, as presented in Figure 5, are re-plotted on the upper

axes with paired observations connected by a line. On the lower axes, paired mean

differences are plotted as a bootstrap sampling distribution. Mean differences are depicted

as dots; 95% confidence intervals are indicated by the ends of the vertical error bars.

Figure EV7. Attenuated insulin-induced association of p110α with Irs2 in the presence

of APDS2 and SHORT syndrome mutant p85α. Results of immunoblotting of anti-Irs2

immunoprecipitates from 3T3-L1 cells expressing wild-type, APDS2-associated or SHORT

syndrome-associated mutant p85α under the control of doxycycline (Dox) are shown.

Treatment with 100nM insulin (Ins) is indicated. (A) One representative immunoblot of

immunoprecipitate, cell lysate prior to immunoprecipitation, and post immunoprecipitation

supernatant is shown. Two separate sets of gels, including independent wild-type controls

are shown on left and right. (B-C) Quantification of immunoblot bands from

immunoprecipitates from 3 independent experiments. Immunoprecipitated p110α is shown

normalized to immunoprecipitated Irs2 from all 3 independent experiments in (B), and

immunoprecipitated p85α similarly in (C). Datapoints from the same experiment +/- insulin

are connected by lines.

Figure EV8. Full statistical analysis of data presented in Figure EV7. Analysis of the

effects of doxycycline (dox)-induced expression of wild-type (WT), Y657X, R649W or ΔEx11

p85α on association of p110α (A,B)   and p85α (C,D) with Irs2. Results of co-

immunoprecipitation with (B,D) and without exposure to 10 nmol/L insulin Comparisons are

made in both the basal, non-insulin stimulated state (A,C) and after stimulation with 10

nmol/L insulin (B,D). Paired mean differences for 3 comparisons are shown in Cumming

estimation plots. Raw data, as presented in Figure EV7, are re-plotted on the upper axes

with paired observations connected by a line. On the lower axes, paired mean differences

are plotted as a bootstrap sampling distribution. Mean differences are depicted as dots; 95%

confidence intervals are indicated by the ends of the vertical error bars.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5