A mitochondrial carrier transports glycolytic intermediates to link cytosolic and mitochondrial glycolysis in the human gut parasite *Blastocystis*

Eva Pyrihová[1,2], Martin S. King[1], Alannah C. King[1], M. Rey Toleco[2], Mark van der Giezen[2]* and Edmund R.S. Kunji[1]*

[1]Medical Research Council Mitochondrial Biology Unit, The Keith Peters Building, Cambridge Biomedical Campus, Hills Road, Cambridge, CB2 0XY, United Kingdom.

[2]University of Stavanger, Department of Chemistry, Bioscience, and Environmental Engineering, Richard Johnsens gate 4, N-4021 Stavanger, Norway.

*Authors to whom correspondence should be addressed
Tel.: +47 5183 4493, e-mail: mark.vandergiezen@uis.no
Tel.: + 44 1223 252850, e-mail: ek@mrc-mbu.cam.ac.uk

Abstract

Stramenopiles form a clade of diverse eukaryotic organisms, including multicellular

algae, the fish and plant pathogenic oomycetes, such as the potato blight

Phytophthora, and the human intestinal protozoan *Blastocystis*. In most eukaryotes,

glycolysis is a strictly cytosolic metabolic pathway that converts glucose to pyruvate,

resulting in the production of NADH and ATP. In contrast, stramenopiles have a

branched glycolysis in which the enzymes of the pay-off phase are located in both

the cytosol and the mitochondrial matrix. Here, we identify a mitochondrial carrier in

Blastocystis that can transport glycolytic intermediates, such as dihydroxyacetone

phosphate and glyceraldehyde-3-phosphate, across the mitochondrial inner

membrane, linking the cytosolic and mitochondrial branches of glycolysis.

Comparative analyses with the phylogenetically-related human mitochondrial

oxoglutarate carrier (SLC25A11) and dicarboxylate carrier (SLC25A10) show that the

glycolytic intermediate carrier has lost its ability to transport the canonical substrates

malate and oxoglutarate. *Blastocystis* lacks several key components of oxidative

phosphorylation required for the generation of mitochondrial ATP, such as complexes

III and IV, ATP synthase and ADP/ATP carriers. The presence of the glycolytic pay-

off phase in the mitochondrial matrix generates ATP, which powers energy-requiring

processes, such as macromolecular synthesis, as well as NADH, used by

mitochondrial complex I to generate a proton motive force to drive the import of

proteins and molecules. Given its unique substrate specificity and central role in

carbon and energy metabolism, the carrier for glycolytic intermediates identified here

represents a specific drug and pesticide target against stramenopile pathogens,

which are of great economic importance.

Introduction

Glycolysis is a crucial metabolic pathway that most organisms use to obtain energy from the oxidation of sugars, which consists of ten enzymatic reactions in total, divided into two phases. The first 'preparatory' phase consists of five reactions, the products of which are the triose phosphates dihydroxyacetone phosphate and glyceraldehyde-3-phosphate. The second 'pay-off' phase leads to the production of ATP, NADH and pyruvate, which is transported into the mitochondrial matrix by the mitochondrial pyruvate carrier (Bricker et al., 2012; Herzig et al., 2012). In most eukaryotes, glycolysis is a cytosolic process with the exception of trypanosomes, where it is localised in the glycosomes (Opperdoes and Borst, 1977). However, it was recently discovered that a large group of eukaryotes, the stramenopiles, have a branched glycolytic pathway in which the pay-off phase is duplicated and also localised in mitochondria (Abrahamian et al., 2017; Bártulos et al., 2018). The stramenopiles are an extremely diverse eukaryotic group consisting mainly of protist algae, which are among the most important primary oxygen producers in the oceans (Burki et al., 2020). The group also includes multicellular algae (kelps) and the economically-devastating oomycetes, such as *Phytophthora infestans* (Jiang and Tyler, 2012), which causes potato blight, one of the major contributing factors of the Great Famine in Ireland in the late 19th century.

The human gut protozoan *Blastocystis* also belongs to the stramenopiles. Despite more than one billion infected individuals worldwide (Andersen and Stensvold, 2016), the public health significance of *Blastocystis* is debated. *Blastocystis* is often linked to patients with diarrhoea, irritable bowel syndrome (Poirier et al., 2012), or inflammatory bowel disease (Dogruman-Al et al., 2009). However, it is also frequently found in healthy individuals, and a few microbiome studies associated *Blastocystis* with higher microbial diversity (Andersen et al., 2015; Audebert et al., 2016; Stensvold et al., 2022). The controversy may arise from the extreme genetic diversity of *Blastocystis*, which is classified into over 40 subtypes (Andersen et al., 2015; Stensvold et al., 2023), and thus its influence on gut microbiota is likely to be subtype-specific (Tito et al., 2019). Nevertheless, *Blastocystis* ST7-B, which was used in this study, is most frequently associated with pathogenic phenotypes (Ajjampur et al., 2016; Deng et al., 2023; Stensvold et al., 2009; Wu et al., 2014; Yason et al., 2019) and was recently linked to a decreased microbial diversity in

diarrheal patients (Deng et al., 2022). The treatment of choice is metronidazole
(Stenzel and Boreham, 1996), but there have been reports of resistance (Haresh et
al., 1999; Mirza et al., 2011; Moghaddam et al., 2005), highlighting the need for new
therapeutic interventions.

Unlike other stramenopiles, *Blastocystis* has lost the cytosolic pay-off phase of
glycolysis, consistent with the absence of a gene encoding a mitochondrial pyruvate
carrier (Bártulos et al., 2018). The only exception is the last enzyme of the pathway,
pyruvate kinase, which was also identified as a cytosolic protein (Lantsman et al.,
2008). *Blastocystis* is an anaerobe (Zierdt, 1991) with highly divergent mitochondria
(Stechmann et al., 2008). It has respiratory complexes I and II, but lacks the rest of
the electron transport chain, as well as ATP synthase and mitochondrial ADP/ATP
carriers (Gentekaki et al., 2017; Stechmann et al., 2008; Wawrzyniak et al., 2008).
Therefore, *Blastocystis* relies mainly on glycolysis and fermentation for its metabolic
energy requirements (Müller et al., 2012). The transport of glycolytic intermediates
across the mitochondrial inner membrane is necessary for the completion of
glycolysis and is critical for the survival of the parasite. Glycolytic enzymes are
powerful drug targets in parasites that are strictly dependent on glycolysis for ATP
production. The structural differences between host and parasitic glyceraldehyde-3-
phosphate dehydrogenase and triosephosphate isomerase in *Plasmodium*
falciparum have been studied with the aim of inhibiting glycolysis (Bruno et al., 2014;
Penna-Coutinho et al., 2011; Ravindra and Balaram, 2005). Similarly, inhibitors
against phosphofructokinase in *Trypanosoma brucei* have a rapid effect on the
parasite viability without inhibiting the host enzyme (McNae et al., 2021).
Understanding the unique aspects of parasite metabolism is crucial for the discovery
of new drug targets.

It is clear that the branched stramenopile glycolysis needs to be linked by a specific
transporter in the mitochondrial inner membrane, as the solute exchange between
the cytosol and mitochondrial matrix is tightly controlled (Kunji et al., 2020; Ruprecht
and Kunji, 2020). The mitochondrial carrier family (SLC25) is the largest solute
transporter family in humans (Kunji et al., 2020). Its members facilitate the transport
of nucleotides, amino acids, inorganic ions, keto acids, vitamins and other nutrients
across the mitochondrial inner membrane. All SLC25 members have a three-fold

pseudo-symmetric structure, evident at the sequence level by the presence of three
conserved homologous repeats (Saraste and Walker, 1982) and specific motifs
important for their transport mechanism (Ruprecht et al., 2019; Ruprecht and Kunji,
2020, 2021).

Here, we report a stramenopile-specific group of transport proteins closely related to
mitochondrial carboxylate carriers of the SLC25 family. We characterised one of
these unique *Blastocystis* carriers, which has lost its ability to transport the canonical
substrates malate and oxoglutarate. Instead, this carrier can transport several
glycolytic intermediates, including dihydroxyacetone phosphate, glyceraldehyde-3-
phosphate, 3-phosphoglycerate and phosphoenolpyruvate, thus providing a
previously undiscovered transport link between cytosolic and mitochondrial
glycolysis. Since *Blastocystis* relies solely on mitochondrial glycolysis, the inhibition
of this carrier might represent a promising way to treat this parasite.

Results

A group of mitochondrial carboxylate carriers that is unique to stramenopiles

The last enzyme of the cytosolic phase of glycolysis in *Blastocystis* is
triosephosphate isomerase (Bártulos et al., 2018). Consequently, dihydroxyacetone
phosphate and glyceraldehyde-3-phosphate need to be transported into the
mitochondrial matrix for the pay-off phase of glycolysis. Given the presence of
mitochondrial glycolysis in stramenopiles, we searched for a transporter that is not
only ubiquitous in stramenopiles, but also absent in other eukaryotes. As glycolysis in
Blastocystis would be critically dependent on this transporter, we used this parasite
as a model organism. We focused on mitochondrial carboxylate carriers because of
their ability to transport phosphorylated three-carbon substrates (Castegna et al.,
2010; Fiermonte et al., 2001; Palmieri et al., 1972) and compared the sequences of
carboxylate carriers from all major eukaryotic supergroups. Our analyses revealed
four well-defined groups of carriers: dicarboxylate carriers (DIC, SLC25A10),
oxoglutarate/malate carriers (OGC, SLC25A11), di/tricarboxylate carriers (DTC), and
a new stramenopile-specific group, the putative glycolytic intermediate carriers (GIC)
(**Figure 1**). At least one copy of the putative GIC carrier was identified in each
stramenopile species. In *Blastocystis* ST7-B, there are four copies of the putative
GIC protein, but two of them are truncated and lack crucial mitochondrial carrier

elements (Ruprecht et al., 2019; Ruprecht and Kunji, 2020, 2021), possibly rendering these proteins non-functional (XP_012899519.1 has an N-terminal truncation; XP_012896318.1 has a C-terminal truncation) (Ruprecht and Kunji, 2020).

The unidentified *Blastocystis* carriers localise to mitochondria

Another important criterion is that the putative glycolytic intermediate carriers localise to mitochondria in *Blastocystis*. Specific (peptide) rabbit antibodies were raised against the two putative glycolytic intermediate carriers bGIC-1 and bGIC-2 as well as an ortholog of the *Blastocystis* oxoglutarate carrier bOGC (**Figure 2B**). Mitochondria were visualized using antibodies raised against a triosephosphate isomerase/glyceraldehyde-3-phosphate dehydrogenase mitochondrial fusion protein (TPI-GAPDH) (Bártulos et al., 2018). Immunoconfocal microscopy clearly show that bOGC, bGIC-1 and bGIC-2 localise to mitochondria in *Blastocystis*, showing that they are potential candidates for the missing transport links (**Figure 2A**).

Human and *Blastocystis* carriers purified from yeast mitochondria are folded

All three *Blastocystis* proteins were expressed in and purified from *Saccharomyces cerevisiae* mitochondria using the detergent lauryl maltose neopentyl glycol (**Figure 3A**). They were compared to the human oxoglutarate carrier SLC25A11 (hOGC) and dicarboxylate carrier SLC25A10 (hDIC), prepared in the same way. Next, thermostability analysis was used to assess whether the purified proteins are correctly folded. Spectral properties of tryptophan/tyrosine residues change as their local environment changes during protein denaturation in a temperature ramp (Alexander et al., 2014). The assay produces an apparent melting temperature (Tm) at which the rate of protein unfolding is the highest. All three proteins produced unfolding curves, showing that they are folded. The apparent melting temperatures for bOGC (52.6 ± 0.5 °C), bGIC-1 (60.7 ± 0.4 °C) and bGIC-2 (59.8 ± 0.4 °C) (**Figure 3B** and **C**) are similar to those observed for hOGC (51.0 °C), hDIC (54.1 °C) and other mitochondrial carriers (Crichton et al., 2015; Jaiquel Baron et al., 2021; Majd et al., 2018).

Two mitochondrial carriers of *Blastocystis* bind glycolytic intermediates

We have previously shown that inhibitors (Crichton et al., 2015; Jaiquel Baron et al., 2021; Tavoulari et al., 2022) and substrates (Majd et al., 2018; Mavridou et al., 2022)

can specifically stabilize a population of transport proteins in thermal denaturation
assays. This technique provides a simple, high-throughput method for screening
libraries of compounds to identify potential protein binders that can then be confirmed
in low-throughput transport assays with radiolabelled substrates (Majd et al., 2018).
We used detergent-solubilized, purified protein to screen a library of 39 commercially
available compounds, which included mono-, di- and tricarboxylates, acids, inorganic
ions, and several glycolytic intermediates (with the exception of 1,3-
bisphosphoglycerate and 2-phosphoglycerate, which were not available).

In order to quantify the binding effect of a compound, we calculated a thermostability
shift (ΔTm) by subtracting the apparent melting temperature in the absence of a
compound from that in its presence. A positive shift suggests that the compound is
stabilizing, indicating a putative substrate (Majd et al., 2018). We first tested the
library on purified hOGC (SLC25A11) and hDIC (SLC25A10). A range of compounds
had stabilizing effects on hOGC in thermostability shift assays (**Figure 4A**).
Oxoglutarate (3.8 ± 0.4 °C), malate (3.6 ± 0.3 °C), maleate (3.1 ± 0.1 °C), succinate
(1.9 ± 0.3 °C) and malonate (1.3 ± 0.2 °C) all produced positive shifts. For hDIC,
larger shifts than for hOGC were observed for malate (4.8 ± 0.2 °C), maleate (8.0 ±
0.1 °C), succinate (3.0 ± 0.3 °C) and malonate (7.3 ± 0.1 °C) (**Figure 4B**). Shifts were
also observed for sulphate (3.0 ± 0.2 °C), thiosulphate (4.0 ± 0.4 °C), phosphate (3.2
± 0.2 °C), dihydroxyacetone phosphate (2.5 ± 0.4 °C) and glyceraldehyde-3-
phosphate (5.6 ± 0.6 °C) (**Figure 4B**), which were not observed for hOGC and may
reflect a difference in substrate specificity (Crompton et al., 1974).

The thermostability shifts for the *Blastocystis* ortholog bOGC were similar to those of
hOGC (SLC25A11), although some compounds, most notably citrate and isocitrate,
produced small stabilizations that were not observed for the human ortholog (**Figure
4C**). A wider range of compounds caused an increase in stability of the two candidate
carriers bGIC-1 and bGIC-2 compared to bOGC (**Figure 4D** and **E**). These
compounds included glycolytic intermediates, such as glyceraldehyde-3-phosphate
(1.0 ± 0.1 °C and 2.1 ± 0.1 °C for bGIC-1 and bGIC-2, respectively) and
dihydroxyacetone phosphate (3.6 ± 0.2 °C and 1.6 ± 0.1 °C) and inorganic ions, such
as phosphate (1.7 ± 0.1 °C and 5.0 ± 0.1 °C) and sulphate (2.3 ± 0.2 °C and 2.6 ± 0.3

°C). A significant shift was observed for phosphoenolpyruvate (PEP) in the case of bGIC-2 (4.3 ± 0.2 °C). In comparison, hOGC showed no shift for any of these compounds (**Figure 4A**), whereas bOGC showed only a minor shift in the presence of sulphate (0.5 ± 0.1 °C) (**Figure 4C**).

Human OGC and DIC have distinct and overlapping substrate specificities

In addition to the glycolytic intermediates that were the focus of this study, we selected all compounds that stabilized the proteins in thermostability assays and performed transport assays. To test whether a compound was transported, we loaded proteoliposomes with substrate by freeze-thaw-extrusion (Jaiquel Baron et al., 2021), and initiated exchange with the addition of radiolabelled [^{14}C]-malate (hOGC and bOGC, **Figure 3D** and **E**) or [^{35}S]-sulphate (hDIC). Only when the internalized compound is a transported substrate of the carrier will the uptake of radiolabelled substrate occur. For hOGC (SLC25A11), uptake was measured for oxoglutarate, oxaloacetate, malate, succinate, maleate, malonate, 2-oxoadipate (**Figure 5A**). Surprisingly, a low uptake rate was measured for pyruvate, which did not induce a significant shift (0.1 ± 0.3 °C). Pyruvate is a small molecule and therefore probably does not introduce enough additional bonds to stabilize the protein. The substrate specificity of hOGC is similar to that of the rat and bovine oxoglutarate carrier (Bisaccia et al., 1988; Fiermonte et al., 1993; Palmieri et al., 1972). For hDIC (SLC25A10), uptake was observed for malate, succinate, maleate, malonate, sulphate, thiosulphate, phosphate and, to a lesser extent, oxalic acid and dihydroxyacetone phosphate (**Figure 5B**). This result is consistent with previous data that have shown that the rat dicarboxylate carrier also transports sulphate and thiosulphate in addition to dicarboxylates (Crompton et al., 1974; Saris, 1980). Although we measured transport for dihydroxyacetone phosphate, we have not found any evidence for the presence of dihydroxyacetone phosphate inside mitochondria in the literature. It is possible that it is not transported under physiological conditions in competition with dicarboxylates or other substrates. The substrate specificity of *Blastocystis* OGC (**Figure 5C**) is very similar to human OGC (**Figure 5A**), consistent with their close phylogenetic relationship.

***Blastocystis* GIC-2 links cytosolic and mitochondrial glycolysis**

We first tested $[^{14}C]$-malate/malate homo-exchange for bGIC-1 and bGIC-2 (**Figure**

6A and **B**, respectively). No transport was measured, showing that these proteins do

not have the ability to transport malate. We next aimed to find a radiolabelled

substrate that is transported by bGIC-1 and bGIC-2. We tested $[^{33}P]$-

phosphate/phosphate and $[^{35}S]$-sulphate/sulphate homo-exchange, as both of these

compounds induce relatively large shifts in thermostability assays: bGIC-1 (1.7 ± 0.1

°C and 2.3 ± 0.2 °C) and bGIC-2 (5.0 ± 0.1 °C and 2.6 ± 0.3 °C), for phosphate and

sulphate, respectively (**Figure 4D** and **E**). No transport was detected for either

phosphate (**Figure 6C**) or sulphate (**Figure 6E**) for bGIC-1. In contrast, bGIC-2

showed transport activity for both: the rate of $[^{33}P]$-phosphate/phosphate homo-

exchange was 4.5 times above the background (**Figure 6D**), and the rate of $[^{35}S]$-

sulphate/sulphate homo-exchange was 22 times above background (**Figure 6F**). We,

therefore, used $[^{35}S]$-sulphate as a counter-substrate and tested hetero-exchange to

determine the substrate specificity of bGIC-2 (**Figure 5D**). No transport was

observed for the canonical oxoglutarate carrier substrates, including oxaloacetate,

malate, succinate and 2-oxoadipate; oxoglutarate, maleate and malonate displayed

only low transport rates. Interestingly, bGIC-2 displayed high transport rates for

phosphate, sulphate, and thiosulphate and for the glycolytic intermediates

dihydroxyacetone phosphate, glyceraldehyde-3-phosphate, 3-phosphoglycerate and

phosphoenolpyruvate (**Figure 5D**).

Discussion

Glycolysis is a textbook example of a well-conserved cytosolic pathway. However,

the discovery that glycolysis is branched and also mitochondrial in stramenopiles

(Abrahamian et al., 2017; Bártulos et al., 2018; Nakayama et al., 2012) raises

questions about how glycolytic intermediates are transported across the tightly

controlled mitochondrial inner membrane. Here, we report the discovery and

characterisation of a mitochondrial carrier protein unique to stramenopiles that

transports the glycolytic intermediates dihydroxyacetone phosphate, glyceraldehyde-

3-phosphate and phosphoenolpyruvate, providing the heretofore missing transport

link.

Since mitochondrial glycolysis is found exclusively in stramenopiles, we searched for
a transporter that is ubiquitous in these species, but not in other eukaryotes.
Phylogenetic analyses identified a new class of mitochondrial carboxylate carriers in
stramenopiles that are most closely related to plant di/tricarboxylate carriers (**Figure**
1). The plant carriers catalyse the transport of tricarboxylates (citrate, isocitrate, and
aconitate) in exchange for dicarboxylates (oxoglutarate, oxaloacetate, malate,
maleate, succinate, and malonate) (Picault et al., 2002) and are also present in
Alveolata, Rhizaria, and Haptophyta (Toleco et al., 2020), which share a common
evolutionary ancestor with stramenopiles (Burki et al., 2020). Therefore, the glycolytic
intermediate carrier of stramenopiles is probably derived from a common ancestor
with di/tricarboxylate carriers and does not exist in any other eukaryote outside of the
stramenopile clade.

Blastocystis was used in this study as a stramenopile model organism, as it relies
solely on glycolysis for its energy demands and does not have a second branch of
glycolysis in the cytosol. We characterised human and *Blastocystis* carboxylate
carriers and demonstrated that the stramenopile GIC proteins have a unique
substrate specificity that is distinct from the human carriers (**Figure 5**). We also
characterised the substrate specificity of *Blastocystis* OGC, which has the same
specificity as human OGC (SLC25A11), and we demonstrated that it is unable to
transport glycolytic intermediates and thus could not be the missing transport step in
the branched glycolytic pathway.

Blastocystis GIC-1 did not transport any of the tested substrates (**Figure 6A**, **C** and
E), but it is expressed and localised to mitochondria (**Figure 2**), demonstrating that it
is not a pseudogene. We did not test radiolabelled glycolytic intermediates directly in
transport, as phosphoenolpyruvate and dihydroxyacetone phosphate are not
available, and radiolabelled glyceraldehyde-3-phosphate is prohibitively expensive.
We cannot, therefore, exclude the possibility that GIC-1 catalyses glycolytic
intermediate exchange, but not in hetero-exchange reactions with malate, phosphate
and sulphate. It is more likely that bGIC-1 has lost its transport activity and could
possibly fulfil a different role. For example, in *Tetrahymena thermophila*, a non-
functional OGC-like carrier is incorporated into a respiratory supercomplex (Zhou et
al., 2022). In contrast to bGIC-1, bGIC-2 transports several inorganic ions and

glycolytic intermediates, including dihydroxyacetone phosphate and glyceraldehyde-
3-phosphate (**Figure 5D**), but not the canonical substrates of *Blastocystis* OGC, such
as malate, maleate, malonate, and succinate. Most mitochondrial carriers operate as
strict antiporters, where the chemical gradient of one substrate drives the uptake of
the counter-substrate (Ruprecht and Kunji, 2020). A mitochondrial phosphate carrier
homolog was identified in the phylogenetic analysis of *Blastocystis* (**Figure 1 - figure**
supplement 1), which is driven by the proton motive force to maintain high
concentrations of phosphate in the mitochondrial matrix, which in turn could drive the
uptake of glycolytic intermediates. *Blastocystis* lacks many of the conserved carrier
proteins found in the mitochondria of aerobes. In total, we have only identified 18
mitochondrial carriers in *Blastocystis* (**Figure 1 - figure supplement 1**). However, we
show that bGIC-2 seems to be able to compensate for the loss of some mitochondrial
carriers, including the transport of inorganic ions in the absence of a DIC homolog.
Additionally, the distantly related citrate carrier, which transports dicarboxylates,
tricarboxylates and PEP, was also lost in *Blastocystis*, possibly due to an incomplete
TCA cycle devoid of citrate synthase, aconitase, isocitrate dehydrogenase and α-
ketoglutarate dehydrogenase (Stechmann et al., 2008; Zierdt et al., 1988). In
Blastocystis, PEP can be synthesised from mitochondrial pyruvate via oxaloacetate
(Gentekaki et al., 2017; Stechmann et al., 2008), and here we show that PEP
transport is catalysed by bGIC-2. When PEP is exported by bGIC-2, it could be
converted into pyruvate and lactate by cytosolic pyruvate kinase and lactate
dehydrogenase, respectively (Lantsman et al., 2008), which would lead to the
synthesis of ATP and the conversion of NADH to NAD$^+$ in the cytosol. This reaction
may play a significant role in cytosolic ATP generation, as there is no mitochondrial
ADP/ATP carrier in *Blastocystis* for adenine nucleotide exchange (**Figure 1 - figure**
supplement 1). *Blastocystis* also contains a glycerol-3-phosphate dehydrogenase
(Gentekaki et al., 2017), which oxidises NADH to convert glycerol-3-phosphate to
dihydroxyacetone phosphate, which in turn can be transported into mitochondria by
bGIC-2 to generate ATP and reducing equivalents. In most other eukaryotes,
cytosolic NADH is oxidised by the malate-aspartate shuttle, but this pathway requires
an ortholog of the mitochondrial aspartate/glutamate carrier, which could not be
identified in *Blastocystis*, even though it might have a mitochondrial glutamate carrier
(**Figure 1 - figure supplement 1**).
345

What selective advantage could there be to have the pay-off phase of glycolysis in

the mitochondrial matrix? *Blastocystis* is well adapted to an anoxic environment with

its anaerobic metabolism but has a limited ability to detoxify oxygen. In its natural

environment, *Blastocystis* likely uses protons as terminal electron acceptors by using

pyruvate:ferredoxin oxidoreductase, which would result in the production of hydrogen

and a range of fermentative products, such as lactate, acetate, and propionate

(Gentekaki et al., 2017; Lantsman et al., 2008; Stechmann et al., 2008). Under

aerobic stress, such as in a dysbiotic gut (Stensvold and van der Giezen, 2018), it

possibly uses oxygen as the terminal electron acceptor by using alternative oxidase,

which produces water (Tsaousis et al., 2018). However, as a likely adaptation to the

anoxic environment, *Blastocystis* has lost some of the canonical components of

oxidative phosphorylation, including complexes III and IV of the respiratory chain,

ATP-synthase and mitochondrial ADP/ATP carriers (Stechmann et al., 2008).

Consequently, it has lost its ability to synthesize mitochondrial ATP via this route.

This role was taken over by transfer of the pay-off phase of glycolysis to the

mitochondrial matrix in *Blastocystis*, which has several advantages. The degradation

of one glucose molecule in the cytosol requires one ATP and one pyrophosphate

molecule, but will generate four ATP molecules in the mitochondrial matrix, which

can be used for ATP-requiring processes, such as protein import, mitochondrial DNA

replication and transcription, and mitochondrial protein synthesis (Denoeud et al.,

2011; Gentekaki et al., 2017). In addition, two NAD^+ molecules are reduced to

NADH, which can be used by complex I to generate a proton motive force to drive

the import of proteins and the uptake of ions and molecules into mitochondria, such

as phosphate. Cytosolic ATP can be supplied either through the export of PEP by

bGIC-2, as explained, or via a mitochondrial carrier that shares properties with the

mitochondrial Mg-ATP/Pi carrier, which is generally used to equilibrate adenine

nucleotide pools between the matrix and cytosol.

Our results show that the canonical metabolic pathways studied in common model

organisms can be altered in non-model eukaryotic species (**Figure 7**). Moreover, the

newly discovered glycolytic intermediates carrier demonstrates the plasticity of

mitochondrial carriers in terms of their substrate specificity. Through the discovery of

this carrier, we have linked the two halves of glycolysis and provided a potential

target for specific treatment not only for *Blastocystis* but also for other stramenopile
pathogens that threaten global food security.

381
Materials and Methods
***In-silico* analysis**
Bioinformatics-based approaches were used to identify potential candidates for a
glycolytic intermediate transporter. Any potential transporter should be present in all
stramenopiles, but absent in other eukaryotes, and could plausibly transport
phosphorylated three carbon substrates. Most of the transport proteins in
mitochondria belong to the mitochondrial carrier family (SLC25) (Kunji et al., 2020;
Ruprecht and Kunji, 2020), and thus the initial search focused on them.

Mitochondrial carrier sequences were gathered using 53 human mitochondrial
carriers as a BLAST query against the genomes of selected stramenopiles:
Blastocystis ST7-B*, Aureococcus anophagefferens, Ectocarpus sp.*, *Saprolegnia*
parasitica, Phaeodactylum tricornutum and *Phytopthora infestans*. InterProScan
(www.ebi.ac.uk/interpro/search/sequence/) was then used to discard non-carrier hits
(Jones et al., 2014), and partial sequences were removed (sequences with fewer
than three signature repeats or 250 amino acids). Reciprocal BLAST was then used
to eliminate incorrectly assigned sequences. The stramenopile sequences were
aligned to *Homo sapiens, Saccharomyces cerevisiae* and *Arabidopsis thaliana*
carriers. To prepare the alignment for phylogenetic analysis, the structures of the
ADP/ATP carrier (PDB: 1OKC and 6GCI) were used to identify and remove the
hypervariable loop regions in each sequence.

IQ-TREE was used to calculate a phylogenetic tree using an LG + F + R9 model
suggested by ModelFinder and branch supports were generated by ultrafast
bootstrap with 1000 replicates (Hoang et al., 2018; Minh et al., 2020). A clade of
stramenopile-specific transporters was identified as a sister group to the *A. thaliana*
di/tricarboxylate carrier homologs (DTC) (**Figure 1 - figure supplement 1**). Within
this clade, all selected stramenopile species were represented at least once.
Interestingly, *Blastocystis,* which has no homolog of the mitochondrial pyruvate
carrier, and would have to rely solely on the import of glycolytic intermediates, has
four copies of this carrier. Given that DTC substrates are three-carbon dicarboxylates

and tricarboxylates, it is possible that a closely related carrier could transport
phosphorylated three-carbon substrates.

To investigate whether this clade is not only present in stramenopiles, but also
unique to them, BLAST searches were performed using the unique stramenopile
sequences against genomes from an expanded range of eukaryotes using NCBI and
EukProt databases (Priyam et al., 2019). The collected sequences were screened to
remove partial and duplicate sequences and sequences that shared contact points
with carboxylate carriers (Kunji and Robinson, 2006; Robinson and Kunji, 2006) were
selected. They were added to a multiple sequence alignment of the identified
stramenopile carriers and other closely related transporters from the first tree. The
maximum likelihood tree was generated by IQ-TREE with the LG + F + R7 model
suggested by ModelFinder (Minh et al., 2020). Branch supports were generated by
ultrafast bootstrap with 1000 replicates (Hoang et al., 2018).

To address the absence of GIC in other species and exclude the possibility of its
presence outside of stramenopiles, we performed additional analyses. The
stramenopile GIC sequences were blasted against NCBI datasets encompassing
major eukaryotic groups, and the top hits were selected and aligned with GIC
sequences. Phylogenetic analysis of this expanded dataset revealed that the non-
stramenopile carriers are more closely related to other carboxylate carriers, indicating
that even the carriers most similar to GIC do not fall within the GIC clade. These
analyses confirmed that the putative GIC carriers are universal in stramenopiles and
also unique to them.

Protein production in mitochondria of *Saccharomyces cerevisiae*
Genes (SLC25A11, the human oxoglutarate carrier (hOGC), UniProt accession code:
Q02978; SLC25A10, the human dicarboxylate carrier (hDIC), UniProt accession
code: Q9UBX3; *Blastocystis* ST7-B oxoglutarate carrier (bOGC) UniProt accession
code: D8M7L1; *Blastocystis* ST7-B potential glycolytic intermediate carrier 1 (bGIC-
1), UniProt accession code: D8LYZ6; *Blastocystis* ST7-B potential glycolytic
intermediate carrier 2 (bGIC-2), UniProt accession code: D8LWW2) were codon-
optimized with N-terminal eight histidine tag and a Xa protease cleavage site to aid
purification (hOGC, Δ2-13; hDIC, GKP added (positions 2-4); bGIC-1, Δ2-3; bGIC-2,

Δ2). The modified genes were cloned into a pYES2/CT vector (Invitrogen) under the
control of an inducible galactose promoter, as previously described (King and Kunji,
2020), and expressed in *Saccharomyces cerevisiae* (GenScript) strain W303.1B
(hOGC and bGIC-2) or the protease-deficient strain BJ2186 (hDIC, bOGC and bGIC-
1). Transformants were selected on Sc-Ura + 2 % (w/v) glucose plates. For large-
scale expression, a pre-culture of cells grown in Sc-Ura + 2% (w/v) glucose was
inoculated into 50 L of YPG plus 0.1% glucose medium in an Applikon Pilot Plant 140
L bioreactor. Cells were grown at 30 °C for 20 hours, induced with a final
concentration of either 0.4% (hOGC, hDIC and bGIC-1) or 2.0% (bOGC and bGIC-2)
galactose, grown for a further 4 (hOGC) or 8 hours (hDIC, bOGC, bGIC-1 and bGIC-
2), and harvested by centrifugation (4,000 g, 20 minutes, 4 °C). Alternatively, 15 L of
culture were grown in shaker flasks and induced as above. Crude mitochondria were
prepared using a bead mill (Dyno-Mill Multilab, Willy A. Bachofen AG) by established
methods (King and Kunji, 2020).
461
462 **Preparation of lipid for protein purification**
463 Tetraoleoyl cardiolipin (18:1) powder was purchased from Avanti Polar Lipids. Lipids
464 were solubilized in 10% (w/v) lauryl maltose neopentyl glycol (Anatrace) by vortexing
465 for 4 hours at room temperature to give 10 mg mL^{-1} lipid in a 10% (w/v) detergent
466 stock. The stocks were stored in liquid nitrogen.
467
Protein purification by nickel affinity chromatography
Protein was purified as described previously for the human mitochondrial ADP/ATP
carrier (Jaiquel Baron et al., 2021). In total, 0.5 g of mitochondria were solubilized in
1.5% (w/v) lauryl maltose neopentyl glycol, protease inhibitor tablet (Roche), 20 mM
imidazole, and 150 mM NaCl for one hour by rotation at 4 °C. The insoluble material
was separated from the soluble fraction by centrifugation (200,000 g, 45 minutes, 4
°C). Nickel Sepharose slurry (0.5 mL resin; GE healthcare) was added to the soluble
fraction; the mixture was stirred at 4 °C for one hour. The nickel resin was harvested
by centrifugation (100 g, 10 minutes, 4 °C), transferred to a Proteus 1-step batch midi
spin column (Generon), and washed with 40 column volumes of buffer A (20 mM
HEPES pH 7.0, 150 mM NaCl, 40 mM imidazole, 0.1 mg mL^{-1} tetraoleoyl cardiolipin /
0.1% (w/v) lauryl maltose neopentyl glycol) (100 g, 5 minutes, 4 °C); followed by 10
column volumes of buffer B (20 mM HEPES pH 7.0, 50 mM NaCl, 0.1 mg mL^{-1}

tetraoleoyl cardiolipin / 0.1% (w/v) lauryl maltose neopentyl glycol) (100 g, 5 minutes,

4 °C). The nickel resin was resuspended in buffer B and incubated with 10 mM

imidazole, 20 µg factor Xa protease (NEB) and 5 mM $CaCl_2$ with inversion at 10 °C

overnight. Protein was eluted by centrifugation on a spin column (100 g, 2 minutes, 4

°C). Imidazole and NaCl were removed using a PD-10 desalting column according to

the manufacturer's instructions (GE Healthcare) at 4 °C and the concentration was

determined by spectrometry (NanoDrop Technologies) at 280 nm using calculated

extinction coefficients (hOGC: 29,000 M^{-1} cm^{-1}; hDIC: 20,900 M^{-1} cm^{-1}; bOGC:

50,350 M^{-1} cm^{-1}; bGIC-1: 48,735 M^{-1} cm^{-1}; bGIC-2: 41,620 M^{-1} cm^{-1}).

Generation and testing of antibodies

Specific peptides (C+LSGEGTSEKLYSSSF for bOGC, C+IVAPGEARLGSIKMA for

bGIC-1 and C+TAAEGRISGMAIAKS for bGIC-2) were generated as a N-terminal

Keyhole limpet haemocyanin fusion to raise the antibodies in rabbits (Eurogentec).

For Western blots, an overnight incubation with 1:2,000 α-bOGC, 1:6,000 α-bGIC1,

and 1:1,000 α-bGIC2 and one hour incubation with secondary 1:20,000 goat α-

rabbit-HRP (Millipore) against 100 ng of purified protein was used. The membranes

were developed with ECL Western Blotting Detection Reagents (Amersham).

Cell culture

Axenic cultures of *Blastocystis* ST7-B were a generous gift from Professor Kevin S.

W. Tan (National University of Singapore) and were grown as described previously

(Leonardi et al., 2021).

Confocal microscopy

Blastocystis ST7-B cells were harvested by centrifugation (1,000 g, 10 minutes) at

ambient temperature. Cells were washed with pre-reduced PBS, and fixed with 4%

methanol-free formaldehyde (Thermo Fisher Scientific), incubated at 37 °C for 30

minutes. Fixed parasites were permeabilized with 0.1% Triton X-100, and blocked

with 3% BSA prior to antibody staining. Rabbit antibodies against bGIC-1 , bGIC-2 ,

and bOGC (Eurogentec) were diluted with 3% BSA in PBS with 0.01% Triton X-100

at 1:5000, 1:1000, and 1:1500, respectively, and incubated overnight at 4 °C. Guinea

pig anti-TPI-GAPDH (1:1000; Eurogentec), previously shown to localise in the

mitochondria (Bártulos et al., 2018) was used as an mitochondrial marker. Alexa

Fluor™ 488 conjugated goat anti-rabbit (A-21428, Invitrogen) and Alexa Fluor™ 555

conjugated goat anti-guinea pig□(A-11073, Invitrogen), used as secondary

antibodies, were diluted with 3% BSA in PBS with 0.01% Triton X-100 at 0.5 µg mL^{-1}

and 2 µg mL^{-1}, respectively. Hoechst 33342 (5 µg mL^{-1}, Thermo Fisher Scientific)

was used to stain DNA. All immunofluorescence assays were carried out in No. 1.5

high precision glass cover slips (Marienfeld) coated with 20 µg Cell-Tak (Corning).

Cells were mounted using ProLong™ Gold Antifade Mountant (Molecular Probes)

and cured overnight. Images were taken using a Leica SP8 laser scanning confocal

microscope (Leica Microsystems) equipped with oil immersion objective lens (Leica

Microsystems, HC PL APO CS2 100x/1.40 NA). Z-stacks were collected with a

system-optimized sampling compliant to the Nyquest-Shannon sampling theorem

and with LAS X LIGHTNING deconvolution-optimized settings (Leica Microsystems).

Optical sections from the image stacks were further processed using Fiji Version

1.54d.

Thermostability analysis

Thermal unfolding analyses were performed with dye-free differential scanning

fluorimetry (nanoDSF), which monitors the changes in the spectral properties of

tryptophan and tyrosine residues due to changes in their environment caused by

unfolding (Alexander et al., 2014). The proteins have the following number of

tryptophans: hOGC (two); hDIC (one); bOGC (five); bGIC-1 (five); bGIC-2 (four).

Approximately 0.6 µg of protein and 10 mM substrate were added into a final volume

of 10 µL purification buffer B, and the samples were loaded into nanoDSF-grade

standard glass capillaries. The temperature was increased by 4 °C every minute from

25 to 95 °C, the intrinsic fluorescence was measured in a Prometheus NT.48

nanoDSF device, and the apparent melting temperature (Tm) was calculated with the

PR.ThermControl software (NanoTemper Technologies).

Reconstitution of carrier protein into liposomes

A mix containing *Escherichia coli* polar lipid extract (Avanti Polar Lipids order

number: 100600C), egg L-α-phosphatidylcholine (Avanti Polar Lipids order number:

840051C) and tetraoleoyl cardiolipin (Avanti Polar Lipids order number: 710335C) in

a 15:5:1 (w/w) ratio was dried under a stream of nitrogen: 6.3 mg of total lipids were

used per sample (0.625 mL). Lipids were re-hydrated in 20 mM HEPES pH 7.0, 50

mM NaCl and 5 mM MgCl$_2$. The detergent pentaethylene glycol monodecyl ether

(C$_{10}$E$_5$) was added to a final concentration of 2.5% (v/v), and the lipids were

solubilized by vortexing and incubated on ice: the equivalent 30 µg protein was

added per sample. The pentaethylene glycol monodecyl was removed by SM-2 bio-

beads (Bio-Rad): five additions of bio-beads were made to the master mix every 20

minutes with inversion at 4 °C: four additions of 30 mg, and the final addition of 240

mg per sample. The samples were incubated overnight at 4 °C with rotation. Bio-

beads were removed by passage through empty micro-bio spin columns (Bio-Rad).

The compound was internalized by freeze-thaw-extrusion (Jaiquel Baron et al.,

2021). Compound, to a final concentration of 5 mM (malate and sulphate assays) or

10 mM (phosphate assays) was added, and the sample was subjected to three

cycles of freezing in liquid nitrogen for two minutes followed by thawing in a room

temperature water bath for ten minutes. Proteoliposomes were subsequently

extruded by 21 passages through a 0.4-µm filter (Millipore), and the external

substrate was removed and exchanged into buffer (20 mM HEPES, pH 7.0 and 50

mM NaCl) using a PD10 desalting column (GE Healthcare) according to

manufacturer's instructions.

Transport assays

Transport assays were performed using the Hamilton MicroLab Star robot (Hamilton

Robotics Ltd). Proteoliposomes (100 µL) were loaded into the wells of a

MultiScreenHTS + HA 96-well filter plate (pore size 0.45-µm, Millipore). Uptake of the

radiolabelled substrate was initiated by the addition of 100 µL buffer containing 2.5

µM [^{14}C]-malate (American Radiolabelled Chemicals), 25 µM [^{35}S]-sodium sulphate

(Hartmann Analytic), or 200 µM [^{33}P]-orthophosphoric acid (Hartmann Analytic).

Uptake was stopped after 0, 10, 20, 30, 45, 60, 150, 300, 450 and 600 seconds by

filtration and washing three times with 200 µL ice-cold buffer (20 mM HEPES, pH 7.0,

50 mM NaCl). Plates were dried overnight, 200 µL MicroScint-20 (Perkin Elmer) were

added, and levels of radioactivity were determined using a TopCount scintillation

counter (Perkin Elmer). Initial rates were determined from the linear part of the

uptake curve (60 seconds).

SDS-PAGE analysis and protein quantification

582 Protein was visualized by SDS-PAGE analysis using 4-12% bis-tris mPAGE gels and
583 stained using InstantBlue (Abcam), according to the manufacturer's instructions.
584 Reconstitution efficiency was determined by comparing the amount of protein
585 incorporated into proteoliposome samples to a standard curve of known amounts of
586 purified protein.
587

Acknowledgements
MSK and ACK were supported by grant MC_UU_00028/2 of the UK Medical
Research Council to ERSK, and EP and MRT were supported by Norwegian
Research Council grant 301170 to MvdG. We thank Dr. Shane Palmer (MRC
Mitochondrial Biology Unit) for carrying out large-scale fermentations in this project,
Drs. Denis Lacabanne, Sotiria Tavoulari, Camila Cimadamore-Werthein and Pavel
Dolezal for constructive comments on the manuscript, and Associate Professor Kevin
S. W. Tan for the ST7-B *Blastocystis* culture.

Author contributions
Conception and design of the research: E.P., M.S.K., M.v.d.G., and E.R.S.K.
Bioinformatic analyses: A.C.K. and E.P.
Molecular biology: M.S.K.
Microscopy: M.R.T.
Protein purification and thermostability analyses: E.P.
Transport assays: E.P., and M.S.K.
Data analysis: E.P. and M.S.K.
Data interpretation: E.P., M.S.K., M.v.d.G., and E.R.S.K.
Writing of the paper: E.P., M.S.K., A.C.K., M.R.T., M.v.d.G., and E.R.S.K
All authors read and revised the paper.

Competing financial interests
The authors declare no competing interests.

References

Abrahamian, M., Kagda, M., Ah-Fong, A.M.V., and Judelson, H.S. (2017). Rethinking the

evolution of eukaryotic metabolism: novel cellular partitioning of enzymes in stramenopiles

links serine biosynthesis to glycolysis in mitochondria. BMC Evol Biol *17*, 241.

Ajjampur, S.S., Png, C.W., Chia, W.N., Zhang, Y., and Tan, K.S. (2016). Ex Vivo and In Vivo

Mice Models to Study Blastocystis spp. Adhesion, Colonization and Pathology: Closer to

Proving Koch's Postulates. PLoS One *11*, e0160458.

Alexander, C.G., Wanner, R., Johnson, C.M., Breitsprecher, D., Winter, G., Duhr, S., Baaske,

P., and Ferguson, N. (2014). Novel microscale approaches for easy, rapid determination of

protein stability in academic and commercial settings. Biochim Biophys Acta *1844*, 2241-

2250.

Andersen, L.O., Bonde, I., Nielsen, H.B., and Stensvold, C.R. (2015). A retrospective

metagenomics approach to studying Blastocystis. FEMS Microbiol Ecol *91*.

Andersen, L.O., and Stensvold, C.R. (2016). Blastocystis in Health and Disease: Are We

Moving from a Clinical to a Public Health Perspective? J Clin Microbiol *54*, 524-528.

Audebert, C., Even, G., Cian, A., Blastocystis Investigation, G., Loywick, A., Merlin, S.,

Viscogliosi, E., and Chabe, M. (2016). Colonization with the enteric protozoa Blastocystis is

associated with increased diversity of human gut bacterial microbiota. Sci Rep *6*, 25255.

Bártulos, R.C., Rogers, M.B., Williams, T.A., Gentekaki, E., Brinkmann, H., Cerff, R., Liaud,

M.F., Hehl, A.B., Yarlett, N.R., Gruber, A.*, et al.* (2018). Mitochondrial Glycolysis in a Major

Lineage of Eukaryotes. Genome Biol Evol *10*, 2310-2325.

Bisaccia, F., Indiveri, C., and Palmieri, F. (1988). Purification and reconstitution of two anion

carriers from rat liver mitochondria: the dicarboxylate and the 2-oxoglutarate carrier.

Biochim Biophys Acta *933*, 229-240.

Bricker, D.K., Taylor, E.B., Schell, J.C., Orsak, T., Boutron, A., Chen, Y.C., Cox, J.E., Cardon,

C.M., Van Vranken, J.G., Dephoure, N.*, et al.* (2012). A mitochondrial pyruvate carrier

required for pyruvate uptake in yeast, Drosophila, and humans. Science *337*, 96-100.

Bruno, S., Pinto, A., Paredi, G., Tamborini, L., De Micheli, C., La Pietra, V., Marinelli, L.,

Novellino, E., Conti, P., and Mozzarelli, A. (2014). Discovery of covalent inhibitors of

glyceraldehyde-3-phosphate dehydrogenase, a target for the treatment of malaria. J Med

Chem *57*, 7465-7471.

Burki, F., Roger, A.J., Brown, M.W., and Simpson, A.G.B. (2020). The New Tree of Eukaryotes.
Trends Ecol Evol *35*, 43-55.

Castegna, A., Scarcia, P., Agrimi, G., Palmieri, L., Rottensteiner, H., Spera, I., Germinario, L.,
and Palmieri, F. (2010). Identification and functional characterization of a novel
mitochondrial carrier for citrate and oxoglutarate in *Saccharomyces cerevisiae*. J Biol Chem
285, 17359-17370.

Crichton, P.G., Lee, Y., Ruprecht, J.J., Cerson, E., Thangaratnarajah, C., King, M.S., and Kunji,
E.R. (2015). Trends in thermostability provide information on the nature of substrate,
inhibitor, and lipid interactions with mitochondrial carriers. J Biol Chem *290*, 8206-8217.

Crompton, M., Palmieri, F., Capano, M., and Quagliariello, E. (1974). The transport of
thiosulphate in rat liver mitochondria. FEBS Lett *46*, 247-250.

Deng, L., Lee, J.W.J., and Tan, K.S.W. (2022). Infection with pathogenic Blastocystis ST7 is
associated with decreased bacterial diversity and altered gut microbiome profiles in
diarrheal patients. Parasit Vectors *15*, 312.

Deng, L., Wojciech, L., Png, C.W., Kioh, D.Y.Q., Gu, Y., Aung, T.T., Malleret, B., Chan, E.C.Y.,
Peng, G., Zhang, Y.*, et al.* (2023). Colonization with two different Blastocystis subtypes in
DSS-induced colitis mice is associated with strikingly different microbiome and pathological
features. Theranostics *13*, 1165-1179.

Denoeud, F., Roussel, M., Noel, B., Wawrzyniak, I., Da Silva, C., Diogon, M., Viscogliosi, E.,
Brochier-Armanet, C., Couloux, A., Poulain, J.*, et al.* (2011). Genome sequence of the
stramenopile Blastocystis, a human anaerobic parasite. Genome Biol *12*, R29.

Dogruman-Al, F., Kustimur, S., Yoshikawa, H., Tuncer, C., Simsek, Z., Tanyuksel, M., Araz, E.,
and Boorom, K. (2009). Blastocystis subtypes in irritable bowel syndrome and inflammatory
bowel disease in Ankara, Turkey. Mem Inst Oswaldo Cruz *104*, 724-727.

Fiermonte, G., Dolce, V., Palmieri, L., Ventura, M., Runswick, M.J., Palmieri, F., and Walker,
J.E. (2001). Identification of the human mitochondrial oxodicarboxylate carrier. Bacterial
expression, reconstitution, functional characterization, tissue distribution, and chromosomal
location. J Biol Chem *276*, 8225-8230.

Fiermonte, G., Walker, J.E., and Palmieri, F. (1993). Abundant bacterial expression and
reconstitution of an intrinsic membrane-transport protein from bovine mitochondria.
Biochem J *294*, 293-299.

Gentekaki, E., Curtis, B.A., Stairs, C.W., Klimes, V., Elias, M., Salas-Leiva, D.E., Herman, E.K.,

Eme, L., Arias, M.C., Henrissat, B.*, et al.* (2017). Extreme genome diversity in the hyper-

prevalent parasitic eukaryote Blastocystis. PLoS Biol *15*, e2003769.

Haresh, K., Suresh, K., Khairul Anus, A., and Saminathan, S. (1999). Isolate resistance of

Blastocystis hominis to metronidazole. Trop Med Int Health *4*, 274-277.

Herzig, S., Raemy, E., Montessuit, S., Veuthey, J.L., Zamboni, N., Westermann, B., Kunji, E.R.,

and Martinou, J.C. (2012). Identification and functional expression of the mitochondrial

pyruvate carrier. Science *337*, 93-96.

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., and Vinh, L.S. (2018). UFBoot2:

Improving the Ultrafast Bootstrap Approximation. Mol Biol Evol *35*, 518-522.

Jaiquel Baron, S., King, M.S., Kunji, E.R.S., and Schirris, T.J.J. (2021). Characterization of drug-

induced human mitochondrial ADP/ATP carrier inhibition. Theranostics *11*, 5077-5091.

Jiang, R.H., and Tyler, B.M. (2012). Mechanisms and evolution of virulence in oomycetes.

Annu Rev Phytopathol *50*, 295-318.

Jones, P., Binns, D., Chang, H.Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,

Mitchell, A., Nuka, G.*, et al.* (2014). InterProScan 5: genome-scale protein function

classification. Bioinformatics *30*, 1236-1240.

King, M.S., and Kunji, E.R.S. (2020). Expression and purification of membrane proteins in

Saccharomyces cerevisiae. Methods Mol Biol *2127*, 47-61.

Kunji, E.R.S., King, M.S., Ruprecht, J.J., and Thangaratnarajah, C. (2020). The SLC25 Carrier

Family: Important Transport Proteins in Mitochondrial Physiology and Pathology. Physiology

(Bethesda) *35*, 302-327.

Kunji, E.R.S., and Robinson, A.J. (2006). The conserved substrate binding site of

mitochondrial carriers. Biochim Biophys Acta *1757*, 1237-1248.

Lantsman, Y., Tan, K.S.W., Morada, M., and Yarlett, N. (2008). Biochemical characterization

of a mitochondrial-like organelle from Blastocystis sp. subtype 7. Microbiology (Reading)

154, 2757-2766.

Leonardi, S.S., Li, F.J., Chee, M.S., Yason, J.A., Tay, H.Y., Chen, J.Y., Koh, E.Y., He, C.Y., and

Tan, K.S. (2021). Characterisation of novel functionality within the Blastocystis

tryptophanase gene. PLoS Negl Trop Dis *15*, e0009730.

Majd, H., King, M.S., Palmer, S.M., Smith, A.C., Elbourne, L.D., Paulsen, I.T., Sharples, D.,
Henderson, P.J., and Kunji, E.R. (2018). Screening of candidate substrates and coupling ions
of transporters by thermostability shift assays. Elife *7*, e38821.

Mavridou, V., King, M.S., Tavoulari, S., Ruprecht, J.J., Palmer, S.M., and Kunji, E.R.S. (2022).
Substrate binding in the mitochondrial ADP/ATP carrier is a step-wise process guiding the
structural changes in the transport cycle. Nat Commun *13*, 3585.

McNae, I.W., Kinkead, J., Malik, D., Yen, L.H., Walker, M.K., Swain, C., Webster, S.P., Gray, N.,
Fernandes, P.M., Myburgh, E., *et al.* (2021). Fast acting allosteric phosphofructokinase
inhibitors block trypanosome glycolysis and cure acute African trypanosomiasis in mice. Nat
Commun *12*, 1052.

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A.,
and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. Mol Biol Evol *37*, 1530-1534.

Mirza, H., Teo, J.D., Upcroft, J., and Tan, K.S. (2011). A rapid, high-throughput viability assay
for Blastocystis spp. reveals metronidazole resistance and extensive subtype-dependent
variations in drug susceptibilities. Antimicrob Agents Chemother *55*, 637-648.

Moghaddam, D.D., Ghadirian, E., and Azami, M. (2005). Blastocystis hominis and the
evaluation of efficacy of metronidazole and trimethoprim/sulfamethoxazole. Parasitol Res
96, 273-275.

Müller, M., Mentel, M., van Hellemond, J.J., Henze, K., Woehle, C., Gould, S.B., Yu, R.Y., van
der Giezen, M., Tielens, A.G., and Martin, W.F. (2012). Biochemistry and evolution of
anaerobic energy metabolism in eukaryotes. Microbiol Mol Biol Rev *76*, 444-495.

Nakayama, T., Ishida, K., and Archibald, J.M. (2012). Broad distribution of TPI-GAPDH fusion
proteins among eukaryotes: evidence for glycolytic reactions in the mitochondrion? PLoS
One *7*, e52340.

Opperdoes, F.R., and Borst, P. (1977). Localization of nine glycolytic enzymes in a microbody-
like organelle in Trypanosoma brucei: the glycosome. FEBS Lett *80*, 360-364.

Palmieri, F., Quagliariello, E., and Klingenberger, M. (1972). Kinetics and specificity of the
oxoglutarate carrier in rat-liver mitochondria. Eur J Biochem *29*, 408-416.

Penna-Coutinho, J., Cortopassi, W.A., Oliveira, A.A., Franca, T.C., and Krettli, A.U. (2011).
Antimalarial activity of potential inhibitors of Plasmodium falciparum lactate dehydrogenase
enzyme selected by docking studies. PLoS One *6*, e21237.

Picault, N., Palmieri, L., Pisano, I., Hodges, M., and Palmieri, F. (2002). Identification of a
novel transporter for dicarboxylates and tricarboxylates in plant mitochondria. Bacterial
expression, reconstitution, functional characterization, and tissue distribution. J Biol Chem
277, 24204-24211.

Poirier, P., Wawrzyniak, I., Vivares, C.P., Delbac, F., and El Alaoui, H. (2012). New insights into
Blastocystis spp.: a potential link with irritable bowel syndrome. PLoS Pathog *8*, e1002545.

Priyam, A., Woodcroft, B.J., Rai, V., Moghul, I., Munagala, A., Ter, F., Chowdhary, H., Pieniak,
I., Maynard, L.J., Gibbins, M.A.*, et al.* (2019). Sequenceserver: A Modern Graphical User
Interface for Custom BLAST Databases. Mol Biol Evol *36*, 2922-2924.

Ravindra, G., and Balaram, P. (2005). Plasmodium falciparum triosephosphate isomerase:
New insights into an old enzyme. Pure Appl Chem *77*, 281-289.

Robinson, A.J., and Kunji, E.R.S. (2006). Mitochondrial carriers in the cytoplasmic state have
a common substrate binding site. Proc Natl Acad Sci USA *103*, 2617-2622.

Ruprecht, J.J., King, M.S., Zogg, T., Aleksandrova, A.A., Pardon, E., Crichton, P.G., Steyaert, J.,
and Kunji, E.R.S. (2019). The molecular mechanism of transport by the mitochondrial
ADP/ATP carrier. Cell *176*, 435-447.

Ruprecht, J.J., and Kunji, E.R.S. (2020). The SLC25 mitochondrial carrier family: structure and
mechanism. Trends Biochem Sci *45*, 244-258.

Ruprecht, J.J., and Kunji, E.R.S. (2021). Structural Mechanism of Transport of Mitochondrial
Carriers. Annu Rev Biochem *90*, 535-558.

Saraste, M., and Walker, J.E. (1982). Internal sequence repeats and the path of polypeptide
in mitochondrial ADP/ATP translocase. FEBS Lett *144*, 250-254.

Saris, N.E.L. (1980). Sulfate transport by H+ symport and by the dicarboxylate carrier in
mitochondria. Biochem J *192*, 911-917.

Stechmann, A., Hamblin, K., Perez-Brocal, V., Gaston, D., Richmond, G.S., van der Giezen, M.,
Clark, C.G., and Roger, A.J. (2008). Organelles in Blastocystis that blur the distinction
between mitochondria and hydrogenosomes. Curr Biol *18*, 580-585.

Stensvold, C.R., Berg, R., Maloney, J.G., Molokin, A., and Santin, M. (2023). Molecular
characterization of *Blastocystis* and *Entamoeba* of muskoxen and sheep in Greenland. Int J
Parasitol *53*, 673-685.

Stensvold, C.R., Lewis, H.C., Hammerum, A.M., Porsbo, L.J., Nielsen, S.S., Olsen, K.E.,
Arendrup, M.C., Nielsen, H.V., and Molbak, K. (2009). Blastocystis: unravelling potential risk

factors and clinical significance of a common but neglected parasite. Epidemiol Infect *137*,

1655-1663.

Stensvold, C.R., Sorland, B.A., Berg, R., Andersen, L.O., van der Giezen, M., Bowtell, J.L., El-

Badry, A.A., Belkessa, S., Kurt, O., and Nielsen, H.V. (2022). Stool Microbiota Diversity

Analysis of Blastocystis-Positive and Blastocystis-Negative Individuals. Microorganisms *10*.

Stensvold, C.R., and van der Giezen, M. (2018). Associations between Gut Microbiota and

Common Luminal Intestinal Parasites. Trends Parasitol *34*, 369-377.

Stenzel, D.J., and Boreham, P.F. (1996). Blastocystis hominis revisited. Clin Microbiol Rev *9*,

563-584.

Tavoulari, S., Schirris, T.J.J., Mavridou, V., Thangaratnarajah, C., King, M.S., Jones, D.T.D.,

Ding, S., Fearnley, I.M., and Kunji, E.R.S. (2022). Key features of inhibitor binding to the

human mitochondrial pyruvate carrier hetero-dimer. Mol Metab, 101469.

Tito, R.Y., Chaffron, S., Caenepeel, C., Lima-Mendez, G., Wang, J., Vieira-Silva, S., Falony, G.,

Hildebrand, F., Darzi, Y., Rymenans, L.*, et al.* (2019). Population-level analysis of Blastocystis

subtype prevalence and variation in the human gut microbiota. Gut *68*, 1180-1189.

Toleco, M.R., Naake, T., Zhang, Y., Heazlewood, J.L., and Fernie, A.R. (2020). Plant

Mitochondrial Carriers: Molecular Gatekeepers That Help to Regulate Plant Central Carbon

Metabolism. Plants (Basel) *9*.

Tsaousis, A.D., Hamblin, K.A., Elliott, C.R., Young, L., Rosell-Hidalgo, A., Gourlay, C.W.,

Moore, A.L., and van der Giezen, M. (2018). The Human Gut Colonizer Blastocystis Respires

Using Complex II and Alternative Oxidase to Buffer Transient Oxygen Fluctuations in the Gut.

Front Cell Infect Microbiol *8*, 371.

Wawrzyniak, I., Roussel, M., Diogon, M., Couloux, A., Texier, C., Tan, K.S., Vivares, C.P.,

Delbac, F., Wincker, P., and El Alaoui, H. (2008). Complete circular DNA in the mitochondria-

like organelles of Blastocystis hominis. Int J Parasitol *38*, 1377-1382.

Wu, Z., Mirza, H., and Tan, K.S. (2014). Intra-subtype variation in enteroadhesion accounts

for differences in epithelial barrier disruption and is associated with metronidazole

resistance in Blastocystis subtype-7. PLoS Negl Trop Dis *8*, e2885.

Yason, J.A., Liang, Y.R., Png, C.W., Zhang, Y., and Tan, K.S.W. (2019). Interactions between a

pathogenic Blastocystis subtype and gut microbiota: in vitro and in vivo studies. Microbiome

7, 30.

Zhou, L., Maldonado, M., Padavannil, A., Guo, F., and Letts, J.A. (2022). Structures of

Tetrahymena's respiratory chain reveal the diversity of eukaryotic core metabolism. Science

376, 831-839.

Zierdt, C.H. (1991). Blastocystis hominis--past and future. Clin Microbiol Rev *4*, 61-79.

Zierdt, C.H., Donnolley, C.T., Muller, J., and Constantopoulos, G. (1988). Biochemical and

ultrastructural study of Blastocystis hominis. J Clin Microbiol *26*, 965-970.

Figures and legends

Figure 1: Phylogenetic analysis of carboxylate carriers. The maximum likelihood tree was generated by IQ-TREE with the LG + F + R7 model suggested by ModelFinder. The support was calculated using ultrafast bootstrap analysis. The unrooted phylogenetic tree shows the uniqueness of the identified putative stramenopile glycolytic intermediate carriers amongst eukaryotes. The stramenopile-specific glycolytic intermediate carriers (GIC - pink) are most closely related to di/tricarboxylate carriers (DTC – green), and to a lesser extend to oxoglutarate carriers (OGC - blue) and dicarboxylate carriers (DIC - yellow). Stramenopile proteins are in red, proteins included in this study are in bold. Note the absence of any non-

818 stramenopile sequences in this clade. Bootstrap values less than 90 and a value for

819 the GIC group are indicated.

820
821
822

Figure 2: A) Localisation of GIC carriers in *Blastocystis* ST7-B cells using specific

antibodies (orange). The signal colocalises with the mitochondrial protein TPI-

GAPDH (green) (Bártulos et al., 2018), demonstrating its mitochondrial localisation.

DNA is stained with Hoechst 33342 (blue). The scale bar is 5 μm. **B**) Western blots of

Blastocystis OGC, GIC-1, and GIC-2 antibodies against purified protein confirm that

there is no cross reactivity.

Figure 3: Expression, purification and biophysical characterization of *Blastocystis* carriers. **A**) Instant-blue stained SDS-PAGE gel of purified bOGC, bGIC-1 and bGIC-2. **B**) Typical unfolding curves of bOGC (black trace), bGIC-1 (red trace), and bGIC-2 (blue trace) using nanoDSF. **C**) The peak in the derivative of the unfolding curve (dR/dT) is the apparent melting temperature (Tm). **D**) [^{14}C]-malate uptake curve of bOGC reconstituted into proteoliposomes loaded with (white circles) or without (black circles) 5 mM malate. Transport was initiated by the external addition of 2.5 μM [^{14}C]-malate. **E**) Initial transport rates, calculated from panel **D**) from the linear part of the uptake curve (first 60 seconds). The data represent the mean and standard deviation is of n=4 (two biological repeats, each with two technical repeats).

Figure 4: Thermostability screening of detergent-solubilized, purified protein using nanoDSF (Prometheus, Nanotemper). **A)** hOGC, **B)** hDIC, **C)** bOGC, **D)** bGIC-1, and **E)** bGIC-2 against 10 mM di- and tricarboxylates, glycolytic intermediates and inorganic ions. The temperature shift (ΔTm) is the apparent melting temperature in the presence of compound minus the apparent melting temperature in the absence of compound. The data are represented by the mean and standard deviation of two biological repeats, each in triplicate (n=6 in total). G6P, glucose-6-phosphate; F6P, fructose-6-phosphate; F16BP, fructose 1-6 bisphosphate; DHAP, dihydroxyacetone phosphate; G3P, glyceraldehyde-3-phosphate; 3PG, 3-phosphoglycerate; PEP, phosphoenolpyruvate.

Figure 5: Hetero-exchange reactions catalysed by reconstituted carriers. **A**) Proteoliposomes containing human oxoglutarate carrier (hOGC, SLC25A11), **B**) human dicarboxylate carrier (hDIC, SLC25A10), **C**) *Blastocystis* oxoglutarate carrier (bOGC) and **D**) *Blastocystis* putative glycolytic intermediate carrier 2 (bGIC-2) were loaded with 5 mM compound, and transport initiated by the external addition of 2.5 μM [^{14}C]-malate (hOGC and bOGC) or 25 μM [^{35}S]-sulphate (hDIC and bGIC-2). Initial transport rates were calculated from the linear part of the uptake curve (60 seconds). The data represent the mean and standard deviation of two biological repeats, each with two technical repeats (n=4 in total). G6P, glucose-6-phosphate; F6P, fructose-6-phosphate; F16BP, fructose 1-6 bisphosphate; DHAP, dihydroxyacetone phosphate; G3P, glyceraldehyde-3-phosphate; 3PG, 3-phosphoglycerate; PEP, phosphoenolpyruvate.

Figure 6: Proteoliposomes containing *Blastocystis* putative glycolytic intermediate
carriers 1 (bGIC-1; **A**, **C** and **E**) and 2 (bGIC-2; **B**, **D** and **F**) were loaded with either
(**A** and **B**) 5 mM malate, (**C** and **D**) 10 mM phosphate or (**E** and **F**) 5 mM sodium
sulphate, and transport initiated by the external addition of (**A** and **B**) 2.5 µM [^{14}C]-
malate, (**C** and **D**) 200 µM [^{33}P]-phosphate or (**E** and **F**) 25 µM [^{35}S]-sulphate.
Controls with no internalized substrate were also tested (black circles). Initial
transport rates were calculated from the linear part of the uptake curve (60 seconds).
The data represent the mean and standard deviation of two biological repeats, each
with two technical repeats (n=4 in total).

Figure 7: Comparing glycolysis in humans and *Blastocystis*.