High temporal frequency light response in mouse retina is mediated by ON and OFF bipolar cells and requires FAT3 signaling

Evelyn C. Avilés[1,2], Sean K. Wang[3,4], Sarina Patel[1], Shuxiang Shi[5,6], Lucas Lin[3], Vladimir J. Kefalov[7], Lisa V. Goodrich[1*], Constance L. Cepko[3,4*], Yunlu Xue[3,5,8*]

1. Department of Neurobiology, Blavatnik Institute, Harvard Medical School, Boston, MA 02115
2. Present address: Facultad de Ciencias Biologicas, Pontificia Universidad Catolica de Chile, Santiago, Chile
3. Departments of Genetics and Ophthalmology, Harvard Medical School, Boston, MA 02115
4. Howard Hughes Medical Institute, Boston, MA 02115
5. Lingang Laboratory, Shanghai, China, 200031
6. School of Life Science and Technology, ShanghaiTech University, Shanghai, China, 201210.
7. Gavin Herbert Eye Institute & Center for Translational Vision Research, University of California, Irvine, CA 92697
8. Lead contact

*Correspondence: ylxue@lglab.ac.cn, cepko@genetics.med.harvard.edu, lisa_goodrich@hms.harvard.edu

Abstract

Vision is initiated by the reception of light by photoreceptors and subsequent processing via parallel retinal circuits. Proper circuit organization depends on the multi-functional tissue polarity protein FAT3, which is required for amacrine cell connectivity and retinal lamination. Here we investigated the retinal function of *Fat3* mutant mice and found decreases in physiological and perceptual responses to high frequency flashes. These defects did not correlate with abnormal amacrine cell wiring, pointing instead to a role in bipolar cell subtypes that also express FAT3. Indeed, similar deficits were observed in mice lacking the bipolar cell glutamate receptors GRIK1 (OFF-bipolar cells) and GRM6 (ON-bipolar cells). Mechanistically, FAT3 binds to the synaptic protein PTPσ and is required to localize GRIK1 to OFF-cone bipolar cell synapses with cone photoreceptors. How FAT3 impacts ON-cone bipolar cell function at high temporal frequency remains to be uncovered. These findings expand the repertoire of FAT3's functions and reveal the importance of both ON- and OFF-bipolar cells for high frequency light response.

Keywords: high frequency vision, retinal physiology, FAT cadherins, bipolar cells, GRIK1, GRM6

Introduction

The remarkable ability to detect light over a wide range of intensities and frequencies is accomplished by highly specialized cell types and circuits within the retina. These circuits transform signals originating from the photoreceptors following their initial synapse with bipolar cells (BCs), which are classified as ON or OFF depending upon their response to photoreceptor signals, setting the stage for downstream processing events. Critical to these transformations are the >80 types of retinal interneurons[1–4], which are organized into laminae with the cell bodies of the horizontal, BCs, and amacrine cells (ACs) located in the inner nuclear layer (INL), while the photoreceptor cell bodies reside in the outer nuclear layer (ONL). Retinal interneurons form synaptic connections in two layers: in the outer plexiform layer (OPL) among BCs, horizontal cells and photoreceptors, and in the inner plexiform layer (IPL) among BCs, ACs, and the output neurons of the retina, the retinal ganglion cells (RGCs) (**Figure 1a**). Despite the stereotyped and conserved organization of retinal neurons and their connections, it is unclear how important the lamination is for the sense of vision. Also unknown is how the organization of most retinal circuits mediate specific visual responses, e.g. which cell types and connections are needed to process and transmit high frequency signals[5].

Several features of retinal circuit assembly depend upon FAT3, a tissue polarity protein[6–8]. FAT cadherins are transmembrane receptors that can sense cell position in the environment *via* their huge extracellular domains. They induce appropriate changes in cell morphology *via* their intracellular domains, thereby creating cellular asymmetries that are aligned across the tissue. In *Fat3* mutant retinas, many ACs migrate to ectopic locations in the IPL and the ganglion cell layer (GCL). They also fail to retract their trailing neurites, which go on to form ectopic synapses in two misplaced plexiform layers, one in the INL (the outer misplaced plexiform layer, OMPL) and one below the GCL (the inner misplaced plexiform layer, IMPL)[6–8]. These ectopic layers contain synapses between ACs and between ACs and rod BCs, which do not express FAT3[2], but seem to "follow" their AC partners to abnormal locations[7]. FAT3, which is localized to AC dendrites in the IPL, mediates these effects by localizing cytoskeletal effectors needed for migration and retraction[6]. The FAT3 intracellular domain also binds a variety of

synaptic proteins[6] and is therefore poised to control synapse localization or function

independent of its effects on cell morphology. A direct functional role at the synapse has

not been described, and it was not clear how the loss of FAT3 impacts vision.

Here, we used retinal physiology and behavioral analyses to investigate the

effects of FAT3 disruptions on retinal function. We demonstrate that FAT3 ensures

formation of synapses between cones and BC dendrites that are required for proper

transmission of high temporal frequency signals. Basic light responses of retinal

neurons were found to be preserved in *Fat3* mutant mice despite the highly abnormal

ectopic plexiform layers. However, the overall retinal response to 30 Hz flashes were

severely reduced in amplitude and *Fat3* mutants behaved as if they see constant

illumination. Analysis of a variety of conditional *Fat3* knock-out mice revealed that this

phenotype is not due to altered AC lamination. Instead, we found that the OFF-cone

pathway is abnormal, as revealed by an *in vivo* electroretinogram (ERG) protocol that

we developed. Loss of *Grik1*, which encodes an ionotropic glutamate receptor

specifically localized to OFF-Cone BC (CBC) dendrites[9,10], showed a reduction in the d-

wave, but had little effect on the amplitude of the 30 Hz ERG. Indeed, only the loss of

Grik1 along with *Grm6,* which encodes a metabotropic glutamate receptor in ON-CBCs,

resulted in reductions in the amplitude of the ERG response to high frequency flickering

lights, indicating that both CBC pathways are required for high temporal frequency light

responses. Further, we show that the FAT3 intracellular domain binds to the synaptic

protein PTPσ, and that in *Fat3* mutants, both PTPσ and GRIK1 are present at reduced

levels in the ribbon synapses that link cones to OFF-CBCs. Thus, FAT3 is required to

set up and/or maintain a properly organized and functional synapse between BCs and

photoreceptors and the transmission of high frequency signals, highlighting its multiple

and versatile roles in the development and function of retinal circuitry.

Results

Global loss of *Fat3* affects the response of the cone pathway to high temporal

frequency light

In *Fat3*$^{\Delta TM/\Delta TM}$ mutant mice, which lack a membrane localized form of FAT3,

retinal lamination is strongly disrupted, with changes in the position of ACs and their

synapses with other ACs, as well as with other retinal cell types[7]. The abnormal

lamination can also be seen by optical coherence tomography (OCT) imaging in

animals *in vivo* (**Figure S1a**). To assess the functional consequences of this change in

circuit organization*,* we used ERG, a common way to measure electrical changes in the

retina in response to light. By altering the stimulus, it is possible to reveal the

contributions of specific cell types. For instance, signaling through the rod pathway is

assayed by performing the ERG after dark-adaptation and under scotopic conditions,

using dim flashes that elicit little response from the cone pathway. Conversely, photopic

ERG tests isolate the cone pathway by light-adapting the eyes and using a background

light that saturates rod phototransduction. In an ERG waveform, the a-wave originates

from photoreceptors and is followed by the b-wave, which reflects the activity of rod

bipolar cells (RBC) and/or ON-types of CBCs. By flashing the stimulus on and off with

increasing frequency, it is possible to determine how well photoreceptors and BCs are

able to resolve temporal differences in the visual stimulus. As the frequency of the flash

increases, OFF-CBCs are hypothesized to dominate the response based on evidence

from a few genetically modified strains[11]. Additionally, the ERG d-wave, which emerges

when turning off a light step that lasts for a few seconds[12], is thought to represent

mainly OFF-CBC activity[13]. However, definitive *in vivo* evidence is lacking, and d-wave

measurements are not usually collected in the lab or the clinic.

Conventional ERG assays under scotopic and photopic conditions revealed no

obvious difference in *Fat3*$^{\Delta TM/\Delta TM}$ vs. *Fat3*$^{\Delta TM/+}$ littermates, indicating that overall

signaling through rod and cone pathways was preserved (**Figure S1b-e**). Likewise, the

oscillatory potentials between the a- and b-waves, which are thought to reflect the

activity of ACs and/or RGCs[14], showed no significant change in the number, amplitude

or timing of peaks (**Figure S1f**). Thus, *Fat3*$^{\Delta TM/\Delta TM}$ mutant mice can detect standard light

stimuli despite changes in retinal lamination.

Although previous work showed its role in ACs[6,7], *Fat3* is also expressed in

RGCs and some subtypes of BCs, especially OFF-CBCs[2] (**Figure 1b,c, Figure S2**).

Consistent with this expression pattern, FAT3 protein localizes not only to the IPL but

also to the OPL, where BC dendrites form synapses with photoreceptors (**Figure 1d**),

raising the possibility of additional effects on retinal function. Indeed, *in situ* hybridization

confirmed co-expression of *Fat3* with the OFF-CBC marker *Grik1,* which encodes an

ionotropic glutamate receptor[2,15](**Figure 1f-i**. *Fat3* is also expressed in some *Grik1*-

negative CBCs, which are positive for the ON-CBC marker *Grm6*, though to a less

degree (**Figure 1i, S2**). This suggested that FAT3 might be required for retinal functions

that would not be detected using standard ERG, as conventional photopic and scotopic

ERG measures overall activities from populations of photoreceptors and the ON-

pathway, but not the OFF-pathway or an individual ON-CBC subtype.

To determine whether *Fat3* is required for OFF-CBC function *in vivo*, we

recorded ERGs in response to lights that flicker in the high frequency range (i.e. over 15

Hz, as defined by Seeliger and colleagues[11]). Flicker ERG responses in this range are

hypothesized to be dominated by the OFF-pathway but direct evidence has been

lacking[11]. We observed a significant decrease in the amplitude of responses to lights

flickering at 30 Hz in *Fat3*$^{\Delta TM/\Delta TM}$ compared to their *Fat3*$^{\Delta TM/+}$ littermates (**Figure 2a,b**).

Additionally, the timing of the first peak in response to 20 Hz stimulation ("the implicit

time") was delayed (**Figure 2c**). 30 Hz implicit time was not measured, as many *Fat3*

mutant mice presented no response at this frequency. These results suggested that the

transmission of high temporal frequency light responses was impaired, and, more

specifically, in BCs, where the signal to the flicker ERG orignates[11]. To determine if this

physiological deficit in the retina had perceptual consequences, behavioral assays were

conducted, using contextual and vision-cued fear conditioning tests[16,17]. Normally, fear

conditioned animals increase the time of "freezing" if they have been conditioned to

associate an unpleasant stimulus with an environmental cue, in this case, 33 Hz light

(**Figure 2d**, **Supplementary movies**). In contrast to *Fat3*$^{\Delta TM/+}$ mice (n=8 animals),

some *Fat3*$^{\Delta TM/\Delta TM}$ mice (5 out of 9 animals) failed to show a freezing response when

switching from a static light to a 33 Hz flicker (**Figure 2e**). The lack of response was not

due to an inability to form fear memories, as all *Fat3*$^{\Delta TM/\Delta TM}$ and *Fat3*$^{\Delta TM/+}$ animals froze

less when switching from an unpleasant olfactory and tactile context (group "context") to

a novel and safe context within the group with static light (group "static light") (**Figure**

2e). Mutant mice also performed like the WT controls in an optomotor behavioral assay,

which measures spatial discrimination (**Figure 2f**).

Responses to different frequencies of light are used to study the temporal

properties of vision at photoreceptor, brain or psychophysical levels[18]. The flicker ERG

at high temporal frequency was proposed, but had not been established, as a test for

OFF-CBC function. Usually, OFF-CBC function is probed using patch-clamping of single

cells, or assessed at the population level by examination of the d-wave of the ERG. To

assay overall function of the OFF-CBC population, we designed an *in vivo* ERG protocol

for mice based on a study of the *ex vivo* ERG response of amphibian retinas. This

assay measures the retinal voltage change, the d-wave, after turning off a long step of

light[12], which is called a "step ERG". Using this newly developed *in vivo* protocol on

mice, we found that WT mice had a strong d-wave with robust oscillatory potentials at

the end of a three-second exposure to a 1,000 cd/m^2 step of light (**Figure 2g**). By

contrast, in *Fat3*$^{\Delta TM/\Delta TM}$ mice, the d-wave amplitude was decreased, and the d-wave

related oscillatory potentials were absent (**Figure 2g,h**). Thus, responses to both high

frequency flickering light and the light being turned off showed that *Fat3*$^{\Delta TM/\Delta TM}$ mice

have deficits in processing specific types of visual stimuli.

Retinal lamination defects do not disrupt responses to flickering stimuli

We next asked which changes to retinal organization and function are

responsible for the impaired high frequency light response in *Fat3*$^{\Delta TM/\Delta TM}$ mutant mice.

To test the impact of an ectopic plexiform layer, *Fat3* was deleted specifically from ACs

using *Ptf1a*^{CRE} [7,19] and a *Fat3* floxed allele (**Figure 3a,b**). In these conditional knock-out

(cKO) mutants, ACs migrate normally, but do not retract their trailing processes, leading

to formation of an OMPL[7]. We found that the 30 Hz flicker ERG amplitude and 20 Hz

implicit time were normal in *Ptf1a*^{CRE} *Fat3*cKO mice compared to littermate *Ptf1a*^{CRE}

control mice (**Figure 3d-f**), indicating that the high frequency light response defects

were not secondary either to the presence of *Fat3* mutant ACs or the lamination defects

they caused. To survey other FAT3+ cells for possible effects on high frequency light

response, we used the *Isl1*^{CRE} mouse line to drive recombination in all ON-CBCs,

starburst ACs, and RGCs, but not in OFF-CBCs[20] (**Figure 3c,g,h**). Although *Isl1*^{CRE}

*Fat3*cKO mice exhibited all of the cellular phenotypes previously described in the retina

of *Fat3*$^{\Delta TM/\Delta TM}$ mice (**Figure 3g-m**), the 30 Hz flicker ERG amplitude and 20 Hz implicit

time were normal compared to littermate $Isl1^{CRE}$ control mice (**Figure 3n-p**). Thus, the

high frequency light response defects are not due to the gross disorganization of retinal

lamination and do not reflect a role for FAT3 in RGCs or ON-CBCs alone. The 30 Hz

amplitude and 20 Hz implicit timing were also unaffected by removal of *Fat3* from

GABAergic ACs and the type 2 OFF-CBC subset using $Bhlhe22^{CRE}$ (**Figure S3a-k**) or

loss of non-type 2 OFF-CBCs, as occurs in $Fezf2^{-/-}$ mutants[2,21] (**Figure S3l-q**). We

were not able to directly test whether high frequency light response requires FAT3 in

OFF-CBCs, as there is no Cre line that is active in all OFF-CBCs. Nonetheless, these

experiments showed that the high frequency light response defects are not caused by

retinal lamination defects and most likely involve the loss of FAT3 from OFF-CBCs,

potentially in combination with loss from other cell types.

FAT3 intracellular signaling is critical for high frequency light response and the

step ERG d-wave

FAT3 is a versatile protein that can mediate non-autonomous interactions via its

extracellular domain and autonomous effects by recruiting different sets of cytoskeletal

effectors to its ICD[6]. To better understand how FAT3 signaling supports high frequency

visual signal transmission and contributes to the step ERG d-wave, we analyzed

$Fat3^{\Delta ICD\text{-}GFP}$ animals, in which most of the ICD is replaced with GFP while keeping the

extracellular and transmembrane domains anchored to the cell membrane[6] (**Figure 4a**).

These animals showed expression of the FAT3-GFP fusion protein in the OPL (**Figure**

4b,c), consistent with FAT3 protein localization (**Figure 1d**). In $Fat3^{\Delta ICD\text{-}GFP/\Delta ICD\text{-}GFP}$

mutant mice, ACs migrate to abnormal cell layers and fail to retract their neurites, but do

not form ectopic plexiform layers, as shown previously[6]. Therefore, analyzing $Fat3^{\Delta ICD\text{-}}$

$^{GFP/\Delta ICD\text{-}GFP}$ mutants can reveal whether high frequency light response depends on

intracellular signaling and/or the ability to form synapses.

Consistent with previous histological analysis[6], no ectopic plexiform layers were

detected by OCT *in vivo* imaging of $Fat3^{\Delta ICD\text{-}GFP/\Delta ICD\text{-}GFP}$ eyes (**Figure 4d**). Nonetheless,

the amplitude of the 30 Hz flicker ERG response was decreased in $Fat3^{\Delta ICD\text{-}GFP/\Delta ICD\text{-}GFP}$

mice, as in $Fat3^{\Delta TM/\Delta TM}$ mice (**Figure 4e-f**). Likewise, the 20 Hz implicit time was

delayed and the amplitude of the d-wave in step ERG responses was decreased

(**Figure 4g-i**). The loss of high frequency flicker ERG responses even in the absence of ectopic plexiform layers further underscores that changes in AC wiring do not contribute to altered visual function in *Fat3* mutant mice. Rather, these results suggest a potential role for FAT3 signaling in OFF-CBCs, in keeping with the results of the genetic analyses described above.

Disruption of glutamate receptor signaling in CBCs alters ERG responses

The above results suggest that the OFF-CBC pathway is likely to be essential for retinal responses to high frequency stimuli. To test this idea independently, we asked whether ERG responses also are altered upon loss of *Grik1*, which is the only subunit of ionotropic glutamate receptors (iGluR) enriched specifically in OFF-CBCs [2,15,22,23] (**Figure S2, Figure 5a,c,d**). To probe OFF-CBC morphology, we created a novel AAV vector (AAV8-Grik1-GFP) using our previously described *Grik1* enhancer/promoter element[15]. Viral delivery of the Grik1-GFP reporter resulted in the expression of GFP in OFF-CBCs, consistent with the *Grik1* expression profile, and revealed normal cellular morphologies in *Grik1*$^{-/-}$ compared to WT littermates (**Figure S4a,b**). The 0.5 Hz and 10 Hz flicker ERG responses of *Grik1*$^{-/-}$ were similar to WT littermates (**Figure S4e-h**). The amplitude of the response to 30 Hz flickering stimuli was decreased in *Grik1*$^{-/-}$ mice (21.34 ± 1.92 µV, n=10 eyes vs. 32.23 ± 2.65 µV in WT, n=13 eyes, **Figure 5g-i**, **S4i-k**), but more modestly than in *Fat3* mutants (12.65 ± 3.29 µV vs. 43.43 ± 1.16 µV in WT; n=10 eyes each, **Figure 2b**). Additionally, the 20 Hz implicit time was earlier in *Grik1* mutants (35.60 ± 1.13 msec, n=10 eyes vs. 39.54 ± 0.45 msec in WT, n=13 eyes, **Figure 5g,i**, **S4i,k**), whereas this response was delayed in *Fat3* mutants (46.20 ± 1.00 msec, n=10 eyes vs. 36.50 ± 0.50 msec in WT, n=10 eyes; **Figure 2c**). Like *Fat3* mutants, *Grik1*$^{-/-}$ mice showed decreased d-wave amplitudes and no d-wave oscillatory potentials in response to step stimuli when compared to WT littermates (**Figure 5j,k,S4c,d**).

As GRIK1 is the only iGluR subunit that is specifically expressed in OFF-CBCs and *Grik1*$^{-/-}$ eyes did not completely lose the flicker ERG response or the d-wave, we wondered if glutamate receptors specific to ON-CBCs also contribute to the high frequency ERG response. Since the mGluR subunit, GRM6, is predominantly

expressed by ON-CBCs and the loss of GRM6 is sufficient to remove all the transient

ON-BC responses such as the ERG b-wave[2,24] (**Figure S2, Figure 5b**), we generated

Grik1[-/-]; *Grm6*[-/-] double homozygous knockout mice and confirmed the deletion of both

proteins with antibody staining (**Figure 5c-f**)[25,26]. The effect on the ERG response to 30

Hz flickering stimuli was greater in *Grm6*[-/-];*Grik1*[-/-] than *Grik1*[-/-] mice, with strongly

reduced amplitudes (3.63 ± 1.03 µV, n=10 eyes vs. 21.34 ± 1.92 µV in *Grik1*[-/-], n=10

eyes, **Figure 5h**), similar to *Fat3* mutants (12.65 ± 3.29 µV in *Fat3*[ΔTM/ΔTM], n=10 eyes,

Figure 2b). Additionally, the 20 Hz implicit time was significantly delayed in *Grm6*[-/-] mice

(44.08 ± 1.22 msec, n=12 eyes vs. 39.54 ± 0.45 msec in WT, n=13 eyes, **Figure 5i**),

similar to *Fat3* mutants (46.20 ± 1.00 msec, in *Fat3*[ΔTM/ΔTM], n=10 eyes, **Figure 2c**). Thus,

both GRIK1+ OFF-CBCs and GRM6+ ON-BCs contribute to retinal responses to high

frequency light, despite the observation that *Fat3* mRNA was expressed with *Grm6* in

only a few ON-CBC subtypes[2] (**Figure 1h',i', S2**). Contrary to the assumption that the

end of the step ERG is mediated entirely by OFF-CBCs, ERG recordings revealed a

complete loss of the d-wave in *Grm6*[-/-];*Grik1*[-/-] mice, suggesting that both GRM6 and

GRIK1 mediate this response *in vivo* (**Figure 5j,k, S4c,d**).

GRIK1 localization to the ribbon synapse is reduced in *Fat3* mutants

To investigate possible cellular origins of the visual deficits, we analyzed the

retina for possible changes in cones and BCs. We found no change in the number or

distribution of cones (27.77 ± 0.91 cones, n=13 sections from 4 eyes vs. 29.00 ± 0.59

cones, n=15 sections from 4 WT eyes) or BCs (detected by staining for VSX2), which

occupied a similar area in WT compared to *Fat3*[ΔTM/ΔTM] mutant retinas (21.09 ± 1.04

arbitrary units, n=15 sections from 4 eyes vs. 21.59 ± 0.97 arbitrary units, n=15 sections

from 4 WT eyes (**Figure S5a-f**). To visualize OFF-CBC morphology, WT and mutant

P2/P3 retinas were injected with AAV8-Grik1-GFP and evaluated at P22. Whereas 100%

(n=12 cells from 5 animals) of OFF-CBC axons terminated and stratified in the OFF

sublaminas of the IPL in WT animals, only 49 ± 17 % of the mutant OFF-CBC axons

(n=7 animals, 22 cells) terminated properly with 56 ± 19 % of axons terminating instead

in the OMPL or in both the OMPL and the IPL (**Figure S5g-i**). Despite this change in

axon position, mutant OFF-CBCs showed typical bipolar morphologies and extended

dendrites that terminated properly in the OPL, as in the WT retina (**Figure S5g-i**). Thus,

unlike its role in ACs, FAT3 is not essential for the position or shape of OFF-CBC

dendritic arbors.

Given the absence of obvious changes in position or morphology of BC dendrites,

we hypothesized that the observed changes in retinal function are due instead to FAT3-

dependent effects on retinal ribbon synapses. Indeed, in addition to binding to a variety

of cytoplasmic effectors important for neuronal migration and neurite retraction, the

FAT3 ICD binds to several proteins associated with synaptic function, such as

HOMER1[6] and the LAR family protein, PTPσ, which is encoded by the *Ptprs* gene, as

shown by GST-pulldown (**Figure 6a**). Further, *Drosophila* Fat-like interacts with the

related RPTP protein dLAR[27], and PTPσ is required for excitatory synapse formation in

hippocampal neurons[28–30]. Since *Ptprs* RNA is detected in both ON- and OFF-BCs[2]

(**Figure S2**), we asked whether cell type-specific synaptic features might be altered in

the absence of FAT3. Immunostaining revealed that PTPσ localized to the post-synaptic

dendrites of CBCs, co-localized with GRIK1 and apposed CtBP2+ ribbons in the cones

(**Figure 6b,c**). However, significantly less PTPσ was detected in the OPL of *Fat3*$^{\Delta TM/\Delta TM}$

mutants (density in the OPL of 4871 $\pm$ 463.5, n=3 mutants vs 8682 $\pm$ 583.0 in n=4 WT)

(**Figure 6d,e,h**). Immunostaining revealed no change in the levels of pre-synaptic

CTBP2+ ribbons (OPL mean fluorescence intensity: 5.00 $\pm$ 0.14 in n= 4 *Fat3*$^{\Delta TM/\Delta TM}$

animals *vs.* 4.80 $\pm$ 0.16 in n=4 WT; 9.70 $\pm$ 0.39 in n= 4 *Fat3*$^{\Delta ICD\text{-}GFP/\Delta ICD\text{-}GFP}$ animals *vs.*

10.03 $\pm$ 0.32 in n=4 WT) (**Figure S6a,b,e,g,h,k**) or post-synaptic HOMER1 in the OPL

of *Fat3* mutants (**Figure 6f,g,i**), indicating that FAT3 is not required for synapse

formation *per se.* Moreover, GRIK1 staining was severely reduced in the OPL of both

Fat3$^{\Delta TM/\Delta TM}$ (**Figure 7a-c**) and *Fat3*$^{\Delta ICD\text{-}GFP/\Delta ICD\text{-}GFP}$ (**Figure 7d-f**) retinas compared to

WT littermates (density in the OPL: 25838 $\pm$ 5028, n=4 *Fat3*$^{\Delta TM/\Delta TM}$ vs 52727 $\pm$ 7204 in

n=4 WT and 11687 $\pm$ 1110 in n= 4 *Fat3*$^{\Delta ICD\text{-}GFP/\Delta ICD\text{-}GFP}$ *vs.* 16119 $\pm$ 1168 in n=4 WT).

BCs also expressed less *Grik1* RNA in *Fat3*$^{\Delta TM/\Delta TM}$ animals compared to WT (**Figure**

S7a-c). Loss of GRIK1 was apparent by 3 weeks of age in *Fat3*$^{\Delta TM/\Delta TM}$ mice, when 30

Hz flicker ERG defects were already present (**Figure S7e-k**). By contrast, we did not

detect a significant change in the expression or localization of the ON-CBC synaptic

protein GRM6, with similar levels of *Grm6* RNA in the INL of mutant and control retinas

(**Figure S7a,b,d**) and of GRM6 in the OPL of mutant and control retinas (**Figure**

S6c,d,f). GRM6 levels were also unchanged in *Fat3*$^{\Delta ICD-GFP/\Delta ICD-GFP}$ mutants (**Figure**

S6i,j,l). Loss of *Ptprs* (**Figure S8a,b**) also resulted in a modest decrease of GRIK1 in

the OPL (**Figure S8c,d,g**) without significantly affecting the ribbons of the

photoreceptors (**Figure S8e,f,h**). However, unlike *Fat3* mutants, *Ptprs*$^{-/-}$ animals did not

show any change in the ERG response to a 30 Hz flickering stimulus, the implicit timing

of the response to a 20 Hz stimulus, or the d-wave amplitude in the step ERG assay

(**Figure S8i-k**). Thus, FAT3 likely impacts not only the presence of PTPσ and GRIK1,

but also additional features of the CBC synapses necessary for responses to high

temporal frequency stimuli.

Discussion

A major goal in neuroscience to understand how the diverse cell types of the

retina, including >80 types of interneurons, influence parallel circuits that transform

information about specific elements of the visual world. A valuable approach is to

characterize retinal physiology and visual behavior in mice carrying mutations in genes

required for proper circuit assembly and/or function. However, the nature of the circuits

that enable responses to stimuli that change with high temporal frequency has been

explored very little, in part because standard assays of retinal function, particularly the

conventional scotopic and photopic ERGs, are not designed to detect signals elicited by

these stimuli. Despite lacking the cellular level resolution that can be achieved by *ex*

vivo patch clamping, the ERG is an elegant and relatively simple measurement that can

be used *in vivo* to reveal behaviorally-relevant changes in overall retinal function from

population of cells. Here, we use flicker and step ERGs to show that retinal activity in

response to high frequency flickering and light-off stimuli is altered in *Fat3* mutant mice,

which also show an impaired behavioral response to flickering stimuli. FAT3 is a multi-

functional transmembrane protein that is expressed by ACs, BCs, and RGCs and is

required for retinal lamination. By analyzing the consequences of *Fat3* deletion from

different retinal cell types, we discovered that the loss of high frequency light response

was not due to changes in AC position or connectivity. Instead, these visual defects

appear to reflect abnormal synaptic responses in BCs. Although the precise origin of the

synaptic defects remains unclear, we found that FAT3 binds to the synaptic protein

PTPσ and that both FAT3 and PTPσ are required for enrichment of GRIK1 at the cone

to OFF-CBC synapse. Further, only the loss of glutamatergic signaling through both ON

and OFF CBCs recapitulates the *Fat3* visual deficits. Together, these data demonstrate

the importance of both ON and OFF CBCs for high temporal frequency light response,

and uncover a role for FAT3 in the formation and/or maintenance of functional cone to

BC synapses.

FAT3 mutant mice show deficits in high frequency light response

Despite their severe defects in neuronal wiring, *Fat3* mutant mice are not blind,

as seen by the preservation of conventional ERG and optomotor responses. However,

when presented with fast flickering stimuli, *Fat3* mice showed severe visual deficits

consistent with loss of signaling from cones to OFF-CBCs. In support of this idea,

abnormal high temporal frequency light response was observed only in mice lacking

Fat3 in BCs and did not correlate with effects on retinal lamination (**Figure 2,3,4**).

Additionally, *Fat3* is expressed by multiple BC types, including GRIK1-positive OFF-

CBCs, and GRIK1 levels were reduced in the OPL of *Fat3* mutant mice (**Figure 1,7**). To

independently assay cone to OFF-CBC signaling, we established a step ERG protocol

and showed that the d-wave was reduced in both *Fat3* and *Grik1*[-/-] mutant mice

compared to WT littermates (**Figure 2,4,5**). Collectively, these data suggest that FAT3

is required for proper OFF-CBC function and hence the ability to detect fast flickering

stimuli. This phenotype is fundamentally different from that which occurs in *Fat3* mutant

ACs, which extend extra neurites that form ectopic synapses.

Both ON and OFF BCs are required for high frequency light response

BCs are the first retinal interneurons that encode, segregate, and relay visual

information into over a dozen pathways for further processing. It had been hypothesized

that the OFF-CBCs mediate high frequency visual signal transmission, based upon

indirect evidence from flicker ERG studies of mice carrying mutations such as *Grm6*[-/- 11].

Here, we found that both ON and OFF CBCs are required for high frequency light

response (**Figure 5**). Inspired by the loss of GRIK1 from *Fat3* mutant OFF-CBC

synapses, we assayed visual responses in mutant mice lacking critical glutamate

receptors specific to either ON (GRM6) or OFF (GRIK1) CBCs. However, in contrast to

the shared effects on the d-wave, *Grik1*$^{-/-}$ mice showed only a modest reduction in the

amplitude of the response to a 30 Hz flicker ERG. This was striking given the drastic

reductions in this measurement in *Fat3* mutants. These findings raised the possibility

that FAT3 is also required for ON-CBCs to transmit high frequency signals. Indeed,

mice lacking both *Grm6* and *Grik1* showed a similar reduction in high frequency light

detection as *Fat3* mutants. Other physiological phenotypes also were seen: *Grik1*$^{-/-}$ mice

had advanced implicit time in response to 20 Hz, while *Grm6*$^{-/-}$ mice had delayed implicit

time, similar to *Fat3* mutants. These data strongly suggest a role for both ON and OFF

classes of CBCs in supporting high frequency visual signal transmission.

OFF-CBC activities are thought to be initiated by two classes of ionotropic GluRs,

namely kainate and AMPA receptors, which were proposed to differentially encode

temporal signals from cones[31]. AMPA receptors also have been reported to mediate

high frequency signaling in cb2, a particular OFF-CBC subtype in ground squirrel retina,

as shown by *ex vivo* patch recording[32]. Here, we found that *Grik1*$^{-/-}$;*Grm6*$^{-/-}$ mice lacked

all responses to 30 Hz flicker ERGs, arguing against a role for AMPA receptors in the

bulk transmission of high temporal frequency stimuli from cone to CBCs in mice, at least

not at the population level that is detectable by *in vivo* ERG. Despite some controversy

on this topic in mice[33,34], AMPA receptors might mediate only slow and/or sustained

OFF-CBCs responses that do not contribute to the transient and/or fast responses

needed for high temporal frequency visual signal transmission, as suggested by a

previous *ex vivo* study using pharmacological blockers with patch recordings[35].

How FAT3 impacts ON-CBC function remains unclear. Although *Fat3* RNA is

weakly expressed in the 5D and 6 ON-CBC subtypes[2] (**Figure S2**), possibly affecting

some ON-CBC functions directly, it is unlikely that an autonomous change in only two

subtypes would alter the ON-CBC population responses that the ERG measures.

Further, *Fat3* mutants showed no detectable change in the level of GRM6, the

glutamate receptor that leads to membrane voltage changes in ON bipolar cells[24]. OFF-

CBCs are connected to ON-BCs via AII amacrine cells in IPL. However, this circuit is

unlikely to affect the flicker ERG, which originates from the cation channels of BCs in or

near the OPL[9,36]. However, it is possible that modulation originating within the IPL

affects OPL signaling through an undescribed backpropagation mechanism, which was

suggested by an *ex vivo* study using pharmacological blockers within the amphibian

retina[37]. We propose three possible explanations for the observations of the role of the

ON-CBC: 1) GRM6 protein levels in ON-CBC are affected in *Fat3* mutants but we are

unable to resolve them, as there is abundant GRM6 in RBCs, the most numerous of all

BCs, 2) FAT3 regulates certain GRM6 functions and/or related signaling pathways (e.g.

G proteins or TRPM1, which would also be masked by their expression in RBCs) that

we did not assay, and/or 3) GRM6 is not directly downstream of FAT3 but can act in

concert with FAT3-dependent signaling in OFF-CBCs and/or ACs to modulate high

frequency visual signal transmission. For example, PTPσ might serve as an

intermediary, as it can act in *trans*, is expressed in both ON and OFF-BCs[2] (**Figure S2**),

and is reduced in *Fat3* mutants (**Figure 6**). Further studies, including detailed patch

clamp analysis, are needed to address these possible explanations.

It is also interesting that the d-wave was completely gone in *Grik1*[-/-];*Grm6*[-/-] eyes,

but only slightly reduced in *Grik1*[-/-] eyes, again suggesting an unexpected role for ON-

BCs in the response to light turning off. One possible explanation concerns the

phenomenon of lateral inhibition. An individual ON-BC can be depolarized instead of

hyperpolarized when there is no light in its peripheral receptive field, i.e. surrounding the

central receptive field[38]. ON-BCs were originally termed "ON-center" BCs to accurately

reflect their response to light in their central and peripheral fields. Since the ERG light

stimulus is full-field, it is possible that the effect of having the light turn off in the

peripheral receptive field of individual GRM6+ ON bipolar cells contributes to the

residual d-wave in *Grik1*[-/-] retina.

Fat3 effects on high temporal frequency light response requires intracellular
448
signaling
449

The discovery of a new synaptic defect in *Fat3* mutant retinas highlights FAT3's

versatility as a signaling molecule. Previous work showed that FAT3 acts through

different motifs in its ICD to control AC migration, neurite retraction, and synapse

localization, likely by recruiting different combinations of cytoplasmic effectors[6]. Here,

we found that the FAT3 ICD is also required for CBC function, possibly acting through a

separate module of synapse-related effectors. The FAT3-ICD interacts with several

known synaptic proteins, including PTPσ, which is one of the four type IIA family of

receptor-type protein tyrosine phosphatase in the LAR-RPTP subfamily[39]. Interactions

between FAT3 and PTPσ appear to regulate the amount of GRIK1 at the synapse,

since PTPσ levels were reduced in the OPL of *Fat3* mutants (**Figure 6**) and GRIK1

levels were reduced in the OPL of both *Fat3* and *Ptprs* mutants (**Figure S8**). Although

PTPσ plays a well-established role in differentiation of the pre-synaptic component of

excitatory synapses in the brain[28,30,39], our findings point to a role on the post-synaptic

side, as suggested previously[29]. Further, the related protein LAR also can be localized

to the post-synaptic compartment and is required for proper surface expression and

clustering of AMPA receptors[40]. Thus, it is possible that PTPσ acts similarly to control

the distribution of GRIK1 in CBCs, either on its own or in collaboration with FAT3. Since

Ptprs mutants do not show the same visual deficits as *Fat3* mutants (**Figure S8**), other

LAR subfamily members may compensate. This might also explain why *Ptprs* mutants

have no obvious changes in retinal lamination[41]. Alternatively, other FAT3-dependent

proteins may enable sufficient GRIK1 activity for OFF-CBC signaling in *Ptprs* mutants.

Although much remains to be learned about the contribution of FAT3-PTPσ interactions

to the synapse, this seems to be a conserved relationship, since *Drosophila* Fat-like and

LAR interact to ensure collective cell migration[27], in this case acting in *trans*[42].

There are several ways that FAT3 might influence synaptic function. One model

is that the FAT3 ICD serves as a scaffold for synaptic proteins that secures them to the

OFF-CBC dendrites in the OPL and thus directly shapes visual signal transmission. This

fits with the fact that FAT3, PTPσ, and GRIK1 are all localized to the OPL, and that

several other synaptic proteins, including the WAVE regulatory complex, also interact

with the FAT3 ICD[6]. Alternatively, FAT3 may ensure directed trafficking of PTPσ and

other proteins to the synapse, echoing its role as a tissue polarity protein and its ability

to promote asymmetric localization of cytoskeletal proteins[8]. These possibilities are not

mutually exclusive, as FAT3 could control synapse assembly during development and

then maintain the synapse in the mature retina. Finally, FAT3 may impact the synapse

through effects on gene expression. In flies, the Fat-like ICD is cleaved and binds to a

transcriptional co-repressor to influence gene expression[43]. Since the LAR ICD can be

internalized and thus inhibit transcription[44], FAT3 and PTPσ could cooperate to control

expression of synaptic genes. This notion is supported by the observation that *Grik1*

RNA is modestly reduced in *Fat3* mutant bipolar cells (**Figure S6**).

Physiological consequences of high frequency light response

Finally, it is interesting to mention some recent studies reporting a benefit of

gamma frequency entrainment. A one-hour daily exposure of 40 Hz light flashes plus

40 Hz sound was reported to alleviate dementia symptoms and neurodegeneration in

mouse models of Alzheimer's disease, and in human patients[45–47]. The data shown here

concerning how high temporal frequency light signals are transmitted at the first

synapse may contribute to our understanding of the mechanisms of gamma frequency

entrainment. These data may also help resolve the different results reported in a recent

study of a mouse model of Alzheimer's disease[48] where 40 Hz light stimulation provided

little benefit.

Limitations of the study

Although our data provide strong evidence that FAT3 impacts cone-CBC

signaling needed for high frequency light response, this study does not show definitively

what is wrong at the level of the synapse. The ERG is a measurement of cellular activity

across a population of cells, i.e. a group of cells from the same class that function

similarly, such as OFF-CBC vs. ON-CBC. Due to the lack of Cre lines specific for all

BCs, or OFF-CBCs, or ON-CBCs, it is not possible to analyze the consequences of

FAT3 loss in these cell types. Additionally, despite being an excellent and relatively

easy tool to assay overall retinal physiology of cell populations *in vivo*, the ERG cannot

distinguish electrical activity limited to a particular subtype, e.g. CBC-5D. ERGs are also

not designed to detect signals carried within distinct microcircuits that perform

transformations of particular features of a visual scene. Patch clamping is necessary to

study the responses of specific BC types at single-cell resolution. Likewise, the large

size of FAT3 and its localization to the cell surface make it difficult to define the precise

nature of its effects on PTPσ, GRIK1 and other uncharacterized players. Our discovery

that CBC activity is compromised in *Fat3* mutants sets the stage for more detailed

analysis in the future.

Acknowledgements

We thank members of the Goodrich and Cepko labs, as well as A.P. Sampath (UCLA), for their insightful comments; Barbara Caldarone, Mouse Behavior Core, Harvard Medical School for assistance on behavioral studies; Paula Montero-Llopis, Microscopy Resources on the North Quad, Harvard Medical School for support on microscopic imaging; C. Wright (Vanderbilt U.), M.E. Greenberg (Harvard Medical School), and Christophe Mulle (University of Bordeaux) for sharing mouse strains; Jeannie Chen (University of Southern California), Frans Vinberg (University of Utah), Shinya Sato (University of California Irvine), Henri Leinonen (University of Eastern Finland), Gabriella Niconchuk, Sylvain Lapan and Emma West (Harvard Medical School), Neal Peachey (Cleveland Clinic), Peter Lukasiewicz and Daniel Kerschensteiner (Washington University in St. Louis) for materials, and technical and advisory support. NIH grants K99EY030951 (to Y.X. before June 30, 2022), EY030912 (V.J.K), Howard Hughes Medical Institute (to C.L.C.); The Edward R. and Anne G. Lefler Center and a Harvard Brain Initiative Bipolar Disorders grant (LVG); Lingang Laboratory startup fund (to Y.X. after July 20, 2022); and a Leonard and Isabelle Goldenson Fellowship and Alice and Joseph Brooks Fund Postdoctoral Fellowship (ECA). The authors also acknowledge support from an RPB unrestricted grant to the Department of Ophthalmology, University of California, Irvine.

Author contributions

Conceptualization, ECA, LVG, CLC, YX; methodology, SKW, SP, SS, LL, VJK; investigation, ECA, YX; writing, ECA, YX, LVG, CLC, VJK; funding acquisition, LVG, CLC, YX; supervision, LVG, CLC, YX.

Declaration of interests

The authors declare no competing interests.

Methods

Animals

The *Fat3*[ΔTM] mouse line lacks exon 23, which contains the coding region for the transmembrane domain. Since no ICD anchored to the membrane has been detected[6,7], this allele is expected to act as a full loss of function. The *Fat3*[floxed] line contains LoxP sites flanking exon 23[7]. The *Fat3*[ΔICD-GFP] mouse line has a deletion of most of the FAT3-ICD, which is replaced by GFP. This line possesses a full extracellular domain anchored to the cell membrane[6]. *Fezf2*[-/-] mice were generated by Hirata and colleagues[49]. Heterozygous mice were used as breeders to obtain wildtype and knockout littermates. *Grik1*[-/-] mice were obtained from Christophe Mulle (University of Bordeaux, France)[50]. *Grm6*[-/-] mice (also known as *Grm6*[nob3]), which was characterized by Maddox and colleagues[51], were purchased from The Jackson Laboratory (ME, Strain #: 016883). *Grik1*[-/-];*Grm6*[-/-] mice were bred in house by crossing the two lines. *Ptprs* KO mice were made by Michel Tremblay's laboratory (McGill University)[52]. Transgenic mice expressing Cre recombinase were obtained from the following sources: *Ptf1a*[CRE] (C. Wright, Vanderbilt U.)[19]; *Islet1*[CRE] (a.k.a. *Isl1*[tm1(cre)Sev]/J)[53] (The Jackson Laboratory, ME. Strain #: 024242); and *Bhlhe22*[CRE] (a.k.a *Bhlhb5*[CRE])[54] (M.E. Greenberg, Harvard Medical School). Mice were maintained on a 12 hour/12 hour light/dark cycle at 18–23 °C and 40–60% humidity. Animals were handled ethically according to protocols approved by the Institutional Animal Care and Use Committee at Harvard Medical School. Genotyping was done using real time PCR (Transnetyx, Cordova, TN).

Electroretinography (ERG)

Mice were dark adapted overnight before *in vivo* ERG recordings. Animals were anesthetized with 100/10 mg/kg ketamine/xylazine cocktail and placed on a heating pad. Their pupils were dilated with a drop of 1% tropicamide solution (Bausch + Lomb). Electrodes were applied to the cornea to pick up the electrical signals from the retina. Eyes were kept moist by a drop of phosphate buffered saline (PBS). With an Espion E3 System (Diagonsys LLC), four types of ERG tests were performed: 1) scotopic test; 2) photopic tests with 1, 10, 100 and 1,000 cd s/m^2 under a 30 cd/m^2 background light to saturate the rod responses; 3) flicker tests at 0.5, 10, 20, 30, 40 and 50 Hz; and 4) step-

light test with a three-second light step of 1,000 cd/m^2. The scotopic and photopic ERGs
were conducted as described previously[55]. The flicker ERG was recorded using 3.162
cd s/m^2 flashes as adapted from a published protocol[11]. The step ERG was created for
this study to probe the d-wave from OFF-bipolar cells.

Optical Coherence Tomography (OCT)

OCT of mouse eyes was conducted using an OCT2 system (Phoenix Research
Labs), as described previously[55]. OCT imaging was performed on the mice *in vivo*
immediately after the ERG tests to confirm the previously observed *ex vivo* histological
changes. Before OCT imaging, a drop of GONAK 2.5% hypromellose solution (Akorn)
was applied to the eye as the immersion medium with the OCT lens.

Flicker-light cued fear conditioning assay

The fear conditioning test for high temporal frequency vision was created by
modifying previously published protocols[16]. The Med Associates (St Albans, VT) system
was used for the tests with four LED lights (two green and two yellow) controlled by a
computer software (Med PC). Mice were videotaped through Media Recorder Software
and the fear response of freezing was analyzed by researchers who were blinded to the
genotypes as a surrogate measurement of memory linked with visual input. On Day 1,
the mice were brought to the electric shock cage individually to get familiar with the
environment and procedure. They were in the cage for 30 minutes under a dim house
light in the background and LEDs turned off (**Figure S2**). On Day 2, mice were brought
back to the electric shock cage, first exposed to static green/yellow LED lights for two
minutes, followed by 30 seconds of 33 Hz LED flicker light (i.e. the cue). Within the last
2 seconds of the cue, a series of 0.7mA shocks were initiated to trigger the fear memory
linked with the cue. This static-flicker-shock cycle was repeated twice more. On Day 3,
contextual memory (i.e. context test) was measured by placing the mouse back into the
conditioning chamber for three minutes (no electric shock was delivered during this
session), and the duration of freezing was recorded. Then, cued memory was measured
by placing the mouse into an altered context, which was composed of different tactile
and olfactory cues. The amount of freezing in the altered context was measured as a

baseline (3 minutes, static light) followed by measurement of freezing during

presentation of the cued stimulus (3 minutes, 33 Hz flicker light). The videos were

analyzed by an examiner blinded to the genotypes to extract the freezing time of all

three conditions (i.e. context, static and 33 Hz flicker).

Optomotor assay

Using an OptoMotry System (CerebralMechanics), the optomotor assay to

measure the visual acuity of mice was conducted as described previously[55]. The testing

grates were set with 100% contrast and were moved at 1.5 Hz temporal frequency. The

visual acuity (i.e. maximal spatial frequency in the unit of cycle/degree) was tested by

an examiner, who was blinded to the genotype. During each testing episode, the

examiner reported either "yes" or "no" to a computer program until the threshold of

acuity was reached. The parameter of each testing episode (i.e. spatial frequency) was

determined by the computer program and blinded to the examiner.

Dissections and immunohistochemistry

Animals of the desired postnatal age (3 or 6 weeks of age, as indicated) were

euthanized by CO_2 inhalation and cervical dislocation. Extraocular tissue, the cornea

and the lens were removed from the eyes and the eyecups were further fixed by

immersion in 4 % paraformaldehyde (PFA, EMS Cat#15710) for 30 minutes (min) at

room temperature or 15 min on ice. After several washes with PBS buffer, the eyes

were submerged in 30% sucrose and kept at 4°C for at least 2 hours (h). After sucrose

cryoprotection, eyes were incubated in NEG-50 (VWR, Cat#84000-154) overnight at

4°C and embedded by freezing in a liquid nitrogen vapor bath. Retinal slices were

obtained by cryosectioning the eyes at 20 µm thickness and mounting on Superfrost®

Plus Micro Slide (VWR, Cat#48311-703). The sections were either stained immediately

or stored at -80°C.

For regular immunohistochemistry, NEG-50 was removed by short incubation in

PBS and then sections were blocked and permeabilized by incubation in 5% Normal

Donkey Serum (NDS, Jackson ImmunoResearch Cat#017-000-121) in Sorenson´s

supplemented with 0.5% Triton-X for 1-2 h at room temperature. Sections were then

incubated in primary antibody diluted in blocking buffer overnight at 4°C. After several

washes with PBS, sections were incubated with fluorescent secondary antibodies

diluted in 5% NDS in Sorenson´s buffer supplemented with 0.02% Triton-X for 1.5-2 h at

room temperature. After final washes, sections were mounted in DAPI-Fluoromount-G

(SouthernBiotech Cat#0100-20). Primary antibodies used for immunohistochemistry

were: rabbit anti-ARR3 (Millipore Sigma, Cat#AB15282), goat anti-Bhlhb5 (1:500; Santa

Cruz Biotechnology, Cat#sc-6045), mouse anti-CtBP2 (1:2,000, BD Biosciences,

Cat#612044), rabbit anti-dsRed (cross-reacts with TdTomato, 1:1,000, Clontech,

Cat#632496), mouse anti-FAT3[6,7] (1:200), chicken anti-GFP (1:500; Aves, Cat#GFP-

1020), mouse anti-GRIK1 (GluR5, 1:200, Santa Cruz Biotechnology, Cat#sc-393420)[25],

sheep anti-GRM6 (1:2,000, a gift from Jeannie Chen, USC and originally developed by

Kirill Martemyanov Lab[56]), goat anti-PTPσ (1:200; R&D Systems, Cat# AF3430), rabbit

anti-VGAT (1:300; SynapticSystems, Cat#131002), and mouse anti-VSX2 (Chx10,

1:100; Santa Cruz Biotechnology, Cat#sc-365519). All secondary antibodies were

diluted 1:1,000 and were: Donkey anti-chicken Alexa Fluor® 488, Donkey anti-goat

Alexa Fluor® 568, Donkey anti-mouse Alexa Fluor® 488, Donkey anti-mouse Alexa

Fluor® 568, Goat anti-mouse Alexa Fluor® 647, Donkey anti-rabbit Alexa Fluor® 488,

Donkey anti-rabbit Alexa Fluor® 568, Donkey anti-rabbit Alexa Fluor® 647, and Donkey

anti-sheep Alexa Fluor® 568.

To detect FAT3 on retinal sections, we performed Hybridization Chain Reaction

Immunohistochemistry (HCR-IHC)[57], according to the manufacturer's instructions

(Molecular Instruments, CA). In brief, on day 1 sections were treated similarly to regular

immunohistochemistry. After overnight incubation of the primary mouse anti-FAT3[7]

(1:200) antibody, sections were rinsed with PBS-0.1% Tween-20 (PBS-T) and

incubated with 1 µg/mL of initiator-labeled anti-mouse secondary antibody (Molecular

Instruments) for 1 h at room temperature. Slides were rinsed with PBS-T and a final

rinse with 5X Saline-Sodium Citrate buffer with 0.1% Tween-20 (SSC-T) and incubated

with amplification buffer (Molecular Instruments) for 30 min at room temperature. H1

and h2 fluorescently-labeled hairpins were separately denatured at 95 °C for 90 s

followed by 30 min incubation at room temperature in the dark. A 60 mM hairpin solution

mix was prepared by adding snap-cooled h1 and h2 hairpins to amplification buffer and

incubated on the slides over night at room temperature in a dark, humidified chamber.

After several washes with SSC-T, sections were mounted with DAPI-Fluoromount-G.

***In situ* hybridization (RNAscope)**

678   Tissue collection was performed similar as to for immunohistochemistry, except

using RNAase-free conditions. For RNAscope *in situ* hybridization, we used

RNAscope® Fluorescent Multiplex Reagent Kit v2 (ACD, Cat#323120) assay following

the manufacturer´s instructions. In brief, retinal sections were post-fixed in 4% PFA for

15 min at room temperature, treated with hydrogen peroxide for 10 min at room

temperature and treated with Protease III for 10 min at 40°C before probe incubation.

Probes were obtained from ACD (see Key Resource table). Immunohistochemistry was

performed after *in situ* hybridization by rinsing the sections in PBS after the final

RNAscope wash and permeabilized and blocked again with 5% NDS/0.5% Triton X-100

Sorenson´s buffer, followed by the regular immunohistochemistry protocol.

***In vivo* viral injection**

690   The AAV-Grik1-GFP plasmid was generated by cloning a previously identified

Grik1 enhancer (CRM4)[15] upstream of a simian virus 40 (SV40) intron, Kozak sequence,

GFP coding sequence, woodchuck hepatitis virus post-transcriptional regulatory

element (WPRE), and polyadenylation sequence. To produce the AAV8-Grik1-GFP

vector, HEK293T cells were triple transfected with a mixture of AAV-Grik1-GFP plasmid,

adenovirus helper plasmid, and rep2/cap8 packaging plasmid. Viral particles were

harvested from the supernatant 72 hours after transfection and purified using an

iodixanol gradient as described previously[58]. The titer of AAV8-Grik1-GFP was

determined by comparing SYPRO Ruby (Molecular Probes) staining for viral capsid

proteins (VP1, VP2, and VP3) to that of a reference vector with known titer.

700   To deliver AAV8-Grik1-GFP into the developing retina, we injected into the

subretinal space as described previously[59,60]. In brief, neonatal P2-3 mouse pups were

anesthetized by chilling on ice. We injected 2.5×10^9 vector genomes (vg) per eye,

which is at a titer not toxic to the eye, diluted in PBS and 0.1% Fast Green (for

visualization) using a pulled borosilicate glass needle with an opening of 0.5-1mm

diameter connected to an Eppendorf FemtoJet injector into the subretinal space. The

pups recovered on a warm pad and upon regaining consciousness they were returned

to their mother. We then let them develop until performing histological procedures at

P22.

Image acquisition

After immunohistochemistry or RNAscope, retinal sections were imaged within

300 μm from the optic nerve head on a Leica SP8 or a Zeiss LSM800 confocal

microscope. The entire sections were imaged in consecutive z-slices separated by 1 μm

using a 40x or 63x oil objective. The z stacks were then projected at maximum

fluorescence intensity using Fiji/ImageJ.

Histochemical quantifications

We assigned random numbers to each image to ensure blinded quantifications.

Only after the quantification was done, the identity of the images was revealed to assign

the values to their corresponding genotype. All the procedures were done under the

same technical parameters, and the comparisons were made between control and

experimental conditions within the same experiment to avoid batch effects. The animal

(N) and sample (i.e. sections, n) numbers, statistical test performed, and *p* values are

indicated in figure legends and/or figures.

We assessed AC migration and the "ectopic synapse score" or "OMPL score' as

described previously[6]. To quantify expression of PTPσ, GRIK1 or GRM6 proteins in the

OPL, the images were thresholded until background signal in the OPL was not

observed. All images from the same experiment were treated the same way using Fiji

(ImageJ). Then, the integrated density was measured to quantify protein expression in

the same total area of each image on the OPL region (44.69 μm x 17.78 μm). To

quantify CtBP2 fluorescence intensity on immunohistochemistry samples we used Fiji

(ImageJ) to measure the Mean Gray Value on areas of the OPL by tracing a rectangle

that took up most of the OPL height. In addition, the Mean Gray Value was measured

on a rectangle traced on the ONL (region where photoreceptors reside and is used as

background signal) to normalize the value of the OPL.

Statistics

To determine significant differences between control and experimental groups,

we used Prism6 software for statistical analysis. After applying a D'Agostino-Pearson

omnibus normality test to determine Gaussian distribution of the samples, we either

used two-tailed *t* test (if the samples followed a Gaussian distribution) or Mann-Whitney

test (if the samples did not follow a Gaussian distribution) to calculate the *p* values. For

ERG data of more than two groups, one-way ANOVA with Dunnett multiple comparison

test was used.

GST Pull down and Western blot

For binding analysis, we performed Western blots of supernatants after pulling

down binding partners from mouse brain protein homogenates with the FAT3-ICD fused

to Glutathione-S-transferase (GST) and GST alone generated previously[6]. Samples

were denatured at 95°C for 10 mins and subjected to SDS-PAGE in a 4–12%

Criterion™ XT Bis-Tris Protein Gel (Bio-Rad) using XT MES Running Buffer (Bio-Rad).

After 2 h at 150 V of electrophoresis, the proteins were transferred to Immobilon-P

PVSF (0.45µm, Sigma-Millipore) in Tris-Glycine buffer supplemented with 20%

methanol for 1 h at 75V. The Immobilon-P membranes were blocked with 5% skim milk

in TBS buffer and then incubated with primary antibodies at 4°C overnight. The primary

antibody used for Western blots was mouse anti-PTPσ (1:1,000, Medimabs, Cat# MM-

0020-P). After several washes with TBS supplemented with 0.5% Tween 20 (Sigma-

Aldrich), the membranes were incubated with a secondary goat anti-mouse HRP

antibody (Biorad, Cat# 170-6516) diluted 1:2,000 for 1-2 h at room temperature. The

signal was developed using Clarity ECL substrate following the manufacturer´s

instructions (Bio-Rad). Western blots were done at least twice with similar results.

External gene expression profile datasets

Gene expression in different types of retinal bipolar cells was analyzed by using

the single-cell RNAseq database and a modified R script that were published

previously[2].

Resource and data availability

The source data for graphs are provided with this paper. For all quantifications,

the raw data are shown along with means and standard errors of the means as well as

the statistical analyses utilized. The original images used to generate these data are

available from the corresponding author upon request. All unique materials generated in

this study are available upon request.

KEY RESOURCES TABLE

Reagent or resource	Source	Identifier
Antibodies		
Rabbit anti-ARR3	Millipore Sigma	Cat#AB15282; RRID:AB_ 1163387
Goat anti Bhlhb5	Santa Cruz	Cat#sc-6045; RRID:AB_2065343
Mouse anti-CTBP2	BD Biosciences	Cat#612044; RRID:AB_399431
Rabbit anti-dsRed	Clontech	Cat#632496; RRID:AB_10013483
Mouse anti FAT3	[6,7]	N/A; RRID:AB_2904260
Chicken anti GFP	Aves	Cat#GFP-1020; RRID:AB_10000240
Mouse anti-GRIK1 (GluR5)	Santa Cruz Biotechnology	Cat#sc-393420; RRID:AB_2716684
Sheep anti-GRM6	[56]	N/A

Goat anti-PTPσ	R&D Systems	Cat#AF3430; RRID:AB_2175157
Mouse anti-PTPσ	Medimabs	Cat# MM-0020-P
Rabbit anti VGAT	Synaptic systems	Cat#131002; RRID:AB_887871
Mouse anti-VSX2 (Chx10)	Santa Cruz Biotechnology	Cat#sc-365519; RRID:AB_10842442
Donkey anti chicken, Alexa Fluor® 488	Jackson ImmunoResearch	Cat#703-545-155; RRID:AB_2340375
Donkey anti goat, Alexa Fluor® 568	Thermo Fisher Scientific	Cat#A11057; RRID:AB_142581
Donkey anti mouse, Alexa Fluor® 488	Abcam	Cat#ab150105; RRID:AB_2732856
Donkey anti mouse, Alexa Fluor® 568	Thermo Fisher Scientific	Cat#A10037; RRID:AB_2534013
Goat anti mouse, Alexa Fluor® 647	Thermo Fisher Scientific	Cat#A-21235; RRID:AB_2535804
Donkey anti rabbit, Alexa Fluor® 488	Thermo Fisher Scientific	Cat#A21206; RRID:AB_2535792
Donkey anti rabbit, Alexa Fluor® 568	Thermo Fisher Scientific	Cat#A10042; RRID:AB_2534017
Donkey anti rabbit, Alexa Fluor® 647	Thermo Fisher Scientific	Cat#A31573; RRID:AB_2536183
Donkey anti sheep, Alexa Fluor® 568	Thermo Fisher Scientific	Cat#A-21099; RRID:AB_2535753
Goat anti mouse – HRP	BioRad	Cat# 170-6516; RRID:AB_11125547
Commercial assays		
RNAscope® Multiplex	ACD	Cat#323100

Fluorescent Reagent Kit v2		
HCR IHC Bundle	Molecular Instruments	N/A
Experimental models: Organisms/Strains		
Mouse: *Ptf1a*[CRE]	C. Wright, Vanderbilt U [19]	MGI:2387812
Mouse: *Isl1*[CRE]	The Jackson Laboratory[53]	Strain #: 024242
Mouse: *Bhlhe22*[CRE]	M.E. Greenberg (Harvard Medical School)[54]	N/A
Mouse: *Fat3*[floxed]	[7]	N/A
Mouse: *Fat3*[ΔTM]	[7]	N/A
Mouse: *Fat3*[ΔICD-GFP]	[6]	N/A
Mouse: *Fezf2*[-/-]	[49]	N/A
Mouse: *Grik1*[-/-]	Christophe Mulle (University of Bordeaux, France)[50]	N/A
Mouse: *Grm6*[-/-]	The Jackson Laboratory[51]	Strain #: 016883
Mouse: *Ptprs*[-/-]	Michel Tremblay[52]	MGI:2158757
***In situ* hybridization Probes**		
RNAscope® Probe- Mm-*Fat3*-O1	ACD	Cat# 509051
RNAscope® Probe- Mm-*Grik1*-C3	ACD	Cat# 438771-C3

Recombinant DNA		
AAV-Grik1-GFP plasmid	This paper	N/A
Viral vectors		
AAV8-Grik1-GFP	This paper	N/A

References

1. Shekhar, K., and Sanes, J.R. (2021). Generating and Using Transcriptomically
Based Retinal Cell Atlases. Annu. Rev. Vis. Sci. *7*, 43–72.

2. Shekhar, K., Lapan, S.W., Whitney, I.E., Tran, N.M., Macosko, E.Z., Kowalczyk,
M., Adiconis, X., Levin, J.Z., Nemesh, J., Goldman, M., et al. (2016).
Comprehensive Classification of Retinal Bipolar Neurons by Single-Cell
Transcriptomics. Cell *166*, 1308-1323.e30.

3. Yan, W., Laboulaye, M.A., Tran, N.M., Whitney, I.E., Benhar, I., and Sanes, J.R.
(2020). Mouse Retinal Cell Atlas: Molecular Identification of over Sixty Amacrine
Cell Types. J. Neurosci. *40*, 5177–5195.

4. Tran, N.M., Shekhar, K., Whitney, I.E., Jacobi, A., Benhar, I., Hong, G., Yan, W.,
Adiconis, X., Arnold, M.E., Lee, J.M., et al. (2019). Single-Cell Profiles of Retinal
Ganglion Cells Differing in Resilience to Injury Reveal Neuroprotective Genes.
Neuron *104*, 1039-1055.e12.

5. Demb, J.B., and Singer, J.H. (2015). Functional Circuitry of the Retina. Annu. Rev.
Vis. Sci. *1*, 263–289.

6. Avilés, E.C., Krol, A., Henle, S.J., Burroughs-Garcia, J., Deans, M.R., and
Goodrich, L. V. (2022). Fat3 acts through independent cytoskeletal effectors to
coordinate asymmetric cell behaviors during polarized circuit assembly. Cell Rep.
38.

7. Deans, M.R., Krol, A., Abraira, V.E., Copley, C.O., Tucker, A.F., and Goodrich, L.
V. (2011). Control of Neuronal Morphology by the Atypical Cadherin Fat3. Neuron
71, 820–832.

8. Krol, A., Henle, S.J., and Goodrich, L. V. (2016). Fat3 and Ena/VASP proteins
influence the emergence of asymmetric cell morphology in the developing retina.
Development *143*, 2172–2182.

9. Puller, C., Ivanova, E., Euler, T., Haverkamp, S., and Schubert, T. (2013). OFF
bipolar cells express distinct types of dendritic glutamate receptors in the mouse
retina. Neuroscience *243*, 136–148.

10. DeVries, S.H., and Schwartz, E.A. (1999). Kainate receptors mediate synaptic
transmission between cones and "Off" bipolar cells in a mammalian retina. Nature

397, 157–160.

11. Tanimoto, N., Sothilingam, V., Kondo, M., Biel, M., Humphries, P., and Seeliger, M.W. (2015). Electroretinographic assessment of rod- and cone-mediated bipolar cell pathways using flicker stimuli in mice. Sci. Rep. *5*, 10731.

12. Stockton, R.A., and Slaughter, M.M. (1989). B-wave of the electroretinogram. A reflection of ON bipolar cell activity. J. Gen. Physiol. *93*, 101–122.

13. Xu, X., and Karwoski, C. (1995). Current source density analysis of the electroretinographic d wave of frog retina. J. Neurophysiol. *73*, 2459–2469.

14. Kolb, H., Fernandez, E., and Nelson, R. (1995). Webvision. Webvision, 1–31.

15. Kishi, J.Y., Lapan, S.W., Beliveau, B.J., West, E.R., Zhu, A., Sasaki, H.M., Saka, S.K., Wang, Y., Cepko, C.L., and Yin, P. (2019). SABER amplifies FISH: enhanced multiplexed imaging of RNA and DNA in cells and tissues. Nat. Methods *16*, 533–544.

16. Gaub, B.M., Berry, M.H., Holt, A.E., Isacoff, E.Y., and Flannery, J.G. (2015). Optogenetic Vision Restoration Using Rhodopsin for Enhanced Sensitivity. Mol. Ther. *23*, 1562–1571.

17. Shi, Q., Colodner, K.J., Matousek, S.B., Merry, K., Hong, S., Kenison, J.E., Frost, J.L., Le, K.X., Li, S., Dodart, J.C., et al. (2015). Complement C3-Deficient Mice Fail to Display Age-Related Hippocampal Decline. J. Neurosci. *35*, 13029–13042.

18. Donner, K. (2021). Temporal vision: measures, mechanisms and meaning. J. Exp. Biol. *224*.

19. Fujitani, Y., Fujitani, S., Luo, H., Qiu, F., Burlison, J., Long, Q., Kawaguchi, Y., Edlund, H., MacDonald, R.J., Furukawa, T., et al. (2006). Ptf1a determines horizontal and amacrine cell fates during mouse retinal development. Development *133*, 4439–4450.

20. Elshatory, Y., Deng, M., Xie, X., and Gan, L. (2007). Expression of the LIM-homeodomain protein Isl1 in the developing and mature mouse retina. J. Comp. Neurol. *503*, 182–197.

21. Suzuki-Kerr, H., Iwagawa, T., Sagara, H., Mizota, A., Suzuki, Y., and Watanabe, S. (2018). Pivotal roles of Fezf2 in differentiation of cone OFF bipolar cells and functional maturation of cone ON bipolar cells in retina. Exp. Eye Res. *171*, 142–

844   154.

845   22.   Amamoto, R., Garcia, M.D., West, E.R., Choi, J., Lapan, S.W., Lane, E.A.,
846   Perrimon, N., and Cepko, C.L. (2019). Probe-Seq enables transcriptional profiling
847   of specific cell types from heterogeneous tissue by RNA-based isolation. Elife *8*.

848   23.   West, E.R., Lapan, S.W., Lee, C.H., Kajderowicz, K.M., Li, X., and Cepko, C.L.
849   (2022). Spatiotemporal patterns of neuronal subtype genesis suggest hierarchical
850   development of retinal diversity. Cell Rep. *38*.

851   24.   Masu, M., Iwakabe, H., Tagawa, Y., Miyoshi, T., Yamashita, M., Fukuda, Y.,
852   Sasaki, H., Hiroi, K., Nakamura, Y., Shigemoto, R., et al. (1995). Specific deficit of
853   the ON response in visual transmission by targeted disruption of the mGluR6
854   gene. Cell *80*, 757–765.

855   25.   Nemitz, L., Dedek, K., and Janssen-Bienhold, U. (2021). Synaptic Remodeling in
856   the Cone Pathway After Early Postnatal Horizontal Cell Ablation. Front. Cell.
857   Neurosci. *15*.

858   26.   Leinonen, H., Pham, N.C., Boyd, T., Santoso, J., Palczewski, K., and Vinberg, F.
859   (2020). Homeostatic plasticity in the retina is associated with maintenance of night
860   vision during retinal degenerative disease. Elife *9*, 1–27.

861   27.   Barlan, K., Cetera, M., and Horne-Badovinac, S. (2017). Fat2 and Lar Define a
862   Basally Localized Planar Signaling System Controlling Collective Cell Migration.
863   Dev. Cell *40*, 467–477.

864   28.   Han, K.A., Ko, J.S., Pramanik, G., Kim, J.Y., Tabuchi, K., Um, J.W., and Ko, J.
865   (2018). PTPσ drives excitatory presynaptic assembly via various extracellular and
866   intracellular mechanisms. J. Neurosci. *38*, 6700–6721.

867   29.   Horn, K.E., Xu, B., Gobert, D., Hamam, B.N., Thompson, K.M., Wu, C.L.,
868   Bouchard, J.F., Uetani, N., Racine, R.J., Tremblay, M.L., et al. (2012). Receptor
869   protein tyrosine phosphatase sigma regulates synapse structure, function and
870   plasticity. J. Neurochem. *122*, 147–161.

871   30.   Roppongi, R.T., Dhume, S.H., Padmanabhan, N., Silwal, P., Zahra, N., Karimi, B.,
872   Bomkamp, C., Patil, C.S., Champagne-Jorgensen, K., Twilley, R.E., et al. (2020).
873   LRRTMs Organize Synapses through Differential Engagement of Neurexin and
874   PTPσ. Neuron.

875   31.   DeVries, S.H. (2000). Bipolar cells use kainate and AMPA receptors to filter visual
876   information into separate channels. Neuron *28*, 847–856.

877   32.   Grabner, C.P., Ratliff, C.P., Light, A.C., and DeVries, S.H. (2016). Mechanism of
878   High-Frequency Signaling at a Depressing Ribbon Synapse. Neuron *91*, 133–145.

879   33.   Puthussery, T., Percival, K.A., Venkataramani, S., Gayet-Primo, J., Grünert, U.,
880   and Rowland Taylor, W. (2014). Kainate receptors mediate synaptic input to
881   transient and sustained OFF visual pathways in primate retina. J. Neurosci. *34*,
882   7611–7621.

883   34.   Borghuis, B.G., Looger, L.L., Tomita, S., and Demb, J.B. (2014). Kainate
884   receptors mediate signaling in both transient and sustained OFF bipolar cell
885   pathways in mouse retina. J. Neurosci. *34*, 6128–6139.

886   35.   Ichinose, T., and Hellmer, C.B. (2016). Differential signalling and glutamate
887   receptor compositions in the OFF bipolar cell types in the mouse retina. J. Physiol.
888   *594*, 883–894.

889   36.   Morgans, C.W., Zhang, J., Jeffrey, B.G., Nelson, S.M., Burke, N.S., Duvoisin,
890   R.M., and Brown, R.L. (2009). TRPM1 is required for the depolarizing light
891   response in retinal ON-bipolar cells. Proc. Natl. Acad. Sci. U. S. A. *106*, 19174–
892   19178.

893   37.   Awatramani, G., Wang, J., and Slaughter, M.M. (2001). Amacrine and ganglion
894   cell contributions to the electroretinogram in amphibian retina. Vis. Neurosci. *18*,
895   147–156.

896   38.   Nelson, R., and Connaughton, V. (1995). Bipolar Cell Pathways in the Vertebrate
897   Retina. Webvision Organ. Retin. Vis. Syst.

898   39.   Cornejo, F., Cortés, B.I., Findlay, G.M., and Cancino, G.I. (2021). LAR Receptor
899   Tyrosine Phosphatase Family in Healthy and Diseased Brain. Front. Cell Dev.
900   Biol. *9*, 3475.

901   40.   Wyszynski, M., Kim, E., Dunah, A.W., Passafaro, M., Valtschanoff, J.G., Serra-
902   Pagès, C., Streuli, M., Weinberg, R.J., and Sheng, M. (2002). Interaction between
903   GRIP and liprin-α/SYD2 is required for AMPA receptor targeting. Neuron *34*, 39–
904   52.

41.   Sapieha, P.S., Duplan, L., Uetani, N., Joly, S., Tremblay, M.L., Kennedy, T.E.,

and Di Polo, A. (2005). Receptor protein tyrosine phosphatase sigma inhibits

axon regrowth in the adult injured CNS. Mol. Cell. Neurosci. *28*, 625–635.

42.   Williams, A.M., and Horne-Badovinac, S. (2023). Fat2 polarizes Lar and Sema5c

to coordinate the motility of collectively migrating epithelial cells. bioRxiv Prepr.

Serv. Biol.

43.   Fanto, M., Clayton, L., Meredith, J., Hardiman, K., Charroux, B., Kerridge, S., and

McNeill, H. (2003). The tumor-suppressor and cell adhesion molecule Fat controls

planar polarity via physical interactions with Atrophin, a transcriptional co-

repressor. Development *130*, 763–74.

44.   Haapasalo, A., Doo, Y.K., Carey, B.W., Turunen, M.K., Pettingell, W.H., and

Kovacs, D.M. (2007). Presenilin/gamma-secretase-mediated cleavage regulates

association of leukocyte-common antigen-related (LAR) receptor tyrosine

phosphatase with beta-catenin. J. Biol. Chem. *282*, 9063–9072.

45.   Adaikkan, C., Middleton, S.J., Marco, A., Pao, P.C., Mathys, H., Kim, D.N.W.,

Gao, F., Young, J.Z., Suk, H.J., Boyden, E.S., et al. (2019). Gamma Entrainment

Binds Higher-Order Brain Regions and Offers Neuroprotection. Neuron *102*, 929-

943.e8.

46.   Martorell, A.J., Paulson, A.L., Suk, H.J., Abdurrob, F., Drummond, G.T., Guan, W.,

Young, J.Z., Kim, D.N.W., Kritskiy, O., Barker, S.J., et al. (2019). Multi-sensory

Gamma Stimulation Ameliorates Alzheimer's-Associated Pathology and Improves

Cognition. Cell *177*, 256-271.e22.

47.   Iaccarino, H.F., Singer, A.C., Martorell, A.J., Rudenko, A., Gao, F., Gillingham,

T.Z., Mathys, H., Seo, J., Kritskiy, O., Abdurrob, F., et al. (2016). Gamma

frequency entrainment attenuates amyloid load and modifies microglia. Nature

540, 230–235.

48.   Soula, M., Martín-Ávila, A., Zhang, Y., Dhingra, A., Nitzan, N., Sadowski, M.J.,

Gan, W.B., and Buzsáki, G. (2023). Forty-hertz light stimulation does not entrain

native gamma oscillations in Alzheimer's disease model mice. Nat. Neurosci.

49.   Hirata, T., Suda, Y., Nakao, K., Narimatsu, M., Hirano, T., and Hibi, M. (2004).

Zinc finger gene fez-like functions in the formation of subplate neurons and

thalamocortical axons. Dev. Dyn. *230*, 546–556.

50.   Mulle, C., Sailer, A., Swanson, G.T., Brana, C., O'Gorman, S., Bettler, B., and
Heinemann, S.F. (2000). Subunit composition of kainate receptors in hippocampal
interneurons. Neuron *28*, 475–484.

51.   Maddox, D.M., Vessey, K.A., Yarbrough, G.L., Invergo, B.M., Cantrell, D.R.,
Inayat, S., Balannik, V., Hicks, W.L., Hawes, N.L., Byers, S., et al. (2008). Allelic
variance between GRM6 mutants, Grm6nob3 and Grm6nob4 results in
differences in retinal ganglion cell visual responses. J. Physiol. *586*, 4409–4424.

52.   Elchebly, M., Wagner, J., Kennedy, T.E., Lanctôt, C., Michaliszyn, E., Itié, A.,
Drouin, J., and Tremblay, M.L. (1999). Neuroendocrine dysplasia in mice lacking
protein tyrosine phosphatase σ. Nat. Genet. *21*, 330–333.

53.   Yang, L., Cai, C.L., Lin, L., Qyang, Y., Chung, C., Monteiro, R.M., Mummery, C.L.,
Fishman, G.I., Cogen, A., and Evans, S. (2006). Isl1Cre reveals a common Bmp
pathway in heart and limb development. Development *133*, 1575–1585.

54.   Ross, S.E., Mardinly, A.R., McCord, A.E., Zurawski, J., Cohen, S., Jung, C., Hu,
L., Mok, S.I., Shah, A., Savner, E.M., et al. (2010). Loss of inhibitory interneurons
in the dorsal spinal cord and elevated itch in Bhlhb5 mutant mice. Neuron *65*,
886–898.

55.   Xiong, W., Wu, D.M., Xue, Y., Wang, S.K., Chung, M.J., Ji, X., Rana, P., Zhao,
S.R., Mai, S., and Cepko, C.L. (2019). AAV cis-regulatory sequences are
correlated with ocular toxicity. Proc. Natl. Acad. Sci. U. S. A. *116*, 5785–5794.

56.   Cao, Y., Masuho, I., Okawa, H., Xie, K., Asami, J., Kammermeier, P.J., Maddox,
D.M., Furukawa, T., Inoue, T., Sampath, A.P., et al. (2009). Retina-specific
GTPase accelerator RGS11/G beta 5S/R9AP is a constitutive heterotrimer
selectively targeted to mGluR6 in ON-bipolar neurons. J. Neurosci. *29*, 9301–
9313.

57.   Schwarzkopf, M., Liu, M.C., Schulte, S.J., Ives, R., Husain, N., Choi, H.M.T., and
Pierce, N.A. (2021). Hybridization chain reaction enables a unified approach to
multiplexed, quantitative, high-resolution immunohistochemistry and in situ
hybridization. Development *148*.

58.   Grieger, J.C., Choi, V.W., and Samulski, R.J. (2006). Production and
characterization of adeno-associated viral vectors. Nat. Protoc. *1*, 1412–1428.

59.   Matsuda, T., and Cepko, C.L. (2004). Electroporation and RNA interference in the

rodent retina in vivo and in vitro. Proc. Natl. Acad. Sci. U. S. A. *101*, 16–22.

60.   Wang, S., Sengel, C., Emerson, M.M., and Cepko, C.L. (2014). A gene regulatory

network controls the binary fate decision of rod and bipolar cells in the vertebrate

retina. Dev. Cell *30*, 513–527.

Figure 1: *Fat3* **RNA is enriched in OFF-cone bipolar cells.**

(a) Schematic representation of retinal layers and their neurons.

(b) *in situ* hybridization for *Fat3* RNA in WT P22 retinas.

(c) *in situ* hybridization for *Fat3* RNA in *Fat3*$^{\Delta TM/\Delta TM}$ P22 retinal tissue. In b and c the RNA puncta are shown in white and the yellow brackets indicate the area of VSX2+ cell bodies. Yellow dashed lines demark the inner nuclear layer (INL) and the outer misplaced plexiform layer (OMPL) in *Fat3*$^{\Delta TM/\Delta TM}$ tissue. The squares demark the insets seen in b' and c' at higher magnification. VSX2 protein is seen in cyan.

(d) Hybridization Chain Reaction-Immunohistochemistry (HCR-IHC) of FAT3 in wild type retinas. Inset demarked in a yellow box in d is shown at higher magnification in d'.

(e) HCR-IHC of FAT3 in *Fat3*$^{\Delta TM/\Delta TM}$ mutant retinas. Inset demarked in a yellow box in e is shown at higher magnification in e'.

(f) Schematic representation of *Grik1* and *Grm6* RNA enrichment in bipolar cells, according to data in Figure S2.

(g) *in situ* hybridization of *Grik1* RNA (magenta) and Grm6 RNA (yellow, g') with immunostaining for VSX2 (cyan). The insets in g and g' are shown at a higher magnification in g'' and g'''.

(h) Triple *in situ* hybridization to *Fat3*, *Grik1* and *Grm6* RNA. Inset in h is seen at higher magnification in i-i'''.

(i) Higher magnification of inset shown in h. Yellow dashed lines in i'' and i''' demark *Grik1* RNA+ cell bodies. *Fat3* RNA (white) is shown together with *Grm6* RNA in i', with *Grik1* RNA in i'' and alone in i'''.

Scale bars: 20µm.

Figure 2: Flicker ERG and vision at high frequency and step ERG of *Fat3*-deficient mice.

(a) Representative flicker ERG raw traces of WT control and $Fat3^{\Delta TM/\Delta TM}$ eyes elicited by 3.162 cd s/m^2 flashes at 20 and 30 Hz frequencies.

(b) Flicker ERG amplitude at 30 Hz for WT control (n=10) and $Fat3^{\Delta TM/\Delta TM}$ (n=10) eyes. Unpaired two-tailed Student's t test.

(c) Flicker ERG implicit time (1st peak) at 20 Hz for WT control (n=10) and $Fat3^{\Delta TM/\Delta TM}$ (n=10) eyes. Unpaired two-tailed Student's t test.

(d) Schematics of fear conditioning and optomotor behavioral experiment. On Day 1, a mouse is brought to the electric-shock cage with a floor of metal bars for habituation of the environment. On Day 2, the mouse is conditioned by electrical shock paired with 33 Hz flashing light. On Day 3 (see Supplementary movies for representative recordings from Fat3 mutant mice), the mouse is first subjected to a contextual check, in which the "Context" measures the freezing time of the mouse after it is brought back to the electric shock cage, which presents a fear-associated context environment, without the shock. "Static" measures the freezing time of the mouse with a static light, after the covering the metal bars and an odor change. Following this measurement, a 33 Hz flickering light is turned on, and the freezing time of the mouse is measured, as the "flicker" time.

(e) Fear conditioning responses as freezing time (sec) from $Fat3^{\Delta TM/+}$ (n=8) and $Fat3^{\Delta TM/\Delta TM}$ (n=9) mice. One-way ANOVA with Dunnett multiple comparison test.

(f) The visual threshold of spatial frequency of WT (n=8) and $Fat3^{\Delta TM/\Delta TM}$ (n=10) mice measured with the optomotor behavioral assay shown in the bottom cartoon in panel d. Unpaired two-tailed Student's t test.

(g) Representative step ERG raw traces of WT (n=10) control and $Fat3^{\Delta TM/\Delta TM}$ (n=10) eyes elicited by a 3-second step light at 1000 cd/m^2 intensity.

(h) Statistics of step ERG amplitudes (b-wave, d-wave and b : d ratio) of WT (n=10) control and $Fat3^{\Delta TM/\Delta TM}$ (n=10) eyes elicited by a 3-second step of light at 1000 cd/m^2 intensity. Unpaired two-tailed Student's t test.

Figure 3: Flicker ERG at high frequency of *Ptf1a*^{CRE} and *Isl1*^{CRE} conditional *Fat3* mice.

(a) Schematic representation of cell classes that express FAT3 in wild type tissue. Cells that express FAT3 are represented in magenta.

(b) Schematic representation of cell classes, i.e. ACs, that lose *Fat3* expression in a *Ptf1a*^{CRE} cKO, shown in black outlines.

(c) Schematic representation of cell classes, i.e. starburst ACs, RGCs and ON-CBCs, that lose *Fat3* expression in an *Isl1*^{CRE} cKO, shown in black outlines.

(d) Flicker ERG amplitude at 30 Hz for the *Ptf1a*^{CRE} *Fat3*cKO condition. Control genotypes are *Ptf1a*^{CRE};*Fat3*$^{fl/+}$ (n=8 eyes) and *Ptf1a*^{cKO} genotypes are *Ptf1a*^{CRE};*Fat3*$^{fl/\Delta TM}$ (n=8 eyes). Unpaired two-tailed Student's t test.

(e) Flicker ERG implicit time at 20 Hz for *Ptf1a*^{CRE};*Fat3*$^{fl/+}$ (control, n=8 eyes) and *Ptf1a*^{CRE};*Fat3*$^{fl/\Delta TM}$ (*Ptf1a*^{cKO}, n=8 eyes). Unpaired two-tailed Student's t test.

(f) Representative flicker ERG raw traces for *Ptf1a*^{CRE};*Fat3*$^{fl/+}$ (control, n=8 eyes) and *Ptf1a*^{CRE};*Fat3*$^{fl/\Delta TM}$ (*Ptf1a*^{cKO}, n=8 eyes).

(g) VGAT immunostaining for control *Isl1*$^{CRE/+}$;*Fat3*$^{fl/+}$ mice.

(h) VGAT immunostaining for *Isl1*$^{CRE/+}$;*Fat3*$^{fl/\Delta TM}$ *Isl1*^{cKO}. TdTomato reporter of Cre expression is seen in g'-h'.

(i) Quantification of the OMPL score for *Isl1*^{CRE} *Fat3*^{cKO}. Controls (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/+}$): 0.095 ± 0.065 (n=14 sections, N=3 animals); *Isl1*^{cKO} (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/\Delta TM}$): 0.825 ± 0.074 (n=19 sections, N=3 animals), Mann-Whitney test.

(j) DAPI and Bhlhb5 immunostaining of control retinas (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/+}$).

(k) DAPI and Bhlhb5 immunostaining of *Isl1*^{CRE} *Fat3*^{cKO} (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/\Delta TM}$) retinas.

(l) Quantification of the number of nuclei per field in the IPL. Controls (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/+}$): 1.25 ± 0.35 (n=12 sections, N=3 animals); *Isl1*^{cKO} (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/\Delta TM}$): 5.11 ± 0.39 (n=18 sections, N=3 animals). *t* test.

(m) Quantification of the number of Bhlhb5+ nuclei per field in the IPL and GCL. Controls (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/+}$): 3.67 ± 0.43 (n=12 sections, N=3 animals); *Isl1*^{cKO} (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/\Delta TM}$): 9.33 ± 0.67 (n=18 sections, N=3 animals). *t* test.

(n) Flicker ERG amplitude at 30 Hz for control (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/+}$, n=6 eyes) and *Isl1*^{cKO} (*Isl1*$^{CRE/+}$;*Fat3*$^{fl/\Delta TM}$, n=6 eyes). Unpaired two-tailed Student's t test.

1064   (o) Flicker ERG implicit time at 20 Hz for control (*Isl1*[CRE/+];*Fat3*[fl/+], n=6 eyes) and

1065   *Isl1*[cKO] (*Isl1*[CRE/+];*Fat3*[fl/ΔTM], n=6 eyes). Unpaired two-tailed Student's t test.

1066   (p) Representative flicker ERG raw traces for control and *Isl1*[CRE] *Fat3*cKO.

1067   Scale bars: 20μm.

1068

1069

1070

Figure 4: High frequency flicker ERG and step ERG of FAT3 intracellular domain

(ICD) deficient mice.

(a) Schematics of molecular structure of FAT3 wild type protein and FAT[ΔICD-GFP].

(b) Immunostaining for GFP in WT retinal sections. The arrow points the OPL.

(c) Immunostaining for GFP in *Fat3*[ΔICD-GFP/ΔICD-GFP] retinal sections. The arrow points

the OPL.

(d) Representative OCT images of *Fat3*[ΔICD-GFP/+] control and *Fat3*[ΔICD-GFP/ΔICD-GFP]

eyes.

(e) Representative flicker ERG raw traces of *Fat3*[ΔICD-GFP/+] control and *Fat3*[ΔICD-

GFP/ΔICD-GFP] eyes elicited by 3.162 cd s/m^2 flashes at 20 and 30 Hz frequencies.

(f) Flicker ERG amplitude at 30 Hz for *Fat3*[ΔICD-GFP/+] control (n=10 eyes) and

Fat3[ΔICD-GFP/ΔICD-GFP] (n=10) eyes. Unpaired two-tailed Student's t test.

(g) Flicker ERG implicit time at 20 Hz for *Fat3*[ΔICD-GFP/+] control (n=10 eyes) and

Fat3[ΔICD-GFP/ΔICD-GFP] (n=10) eyes at 20 Hz. Unpaired two-tailed Student's t test.

(h) Representative step ERG raw traces of *Fat3*[ΔICD/+] control and *Fat3*[ΔICD-GFP/ΔICD-GFP]

eyes elicited by a 3-second step light at 1000 cd/m^2 intensity.

(i) Statistics of step ERG d-wave amplitudes for *Fat3*[ΔICD-GFP/+] control (n=10 eyes)

and *Fat3*[ΔICD-GFP/ΔICD-GFP] (n=10) eyes elicited by a 3-second step of light at 1000

cd/m^2 intensity. Unpaired two-tailed Student's t test.

1090

1091

Figure 5: Immunostaining, high frequency flicker ERG and step ERG of mice

lacking glutamate receptors, GRIK1 and GRM6.

(a) Schematic representation of OFF-BCs and their synapses with cone
photoreceptors. ARR3 (white) labels cones and GRIK1 (magenta) labels
postsynaptic BC dendrites.
(b) Schematic representation of ON-BCs and their synapses with cone
photoreceptors. ARR3 (white) labels cones and GRM6 (magenta) labels
postsynaptic cone and rod BC (RBC) dendrites.
(c) GRIK1 and GRM6 immunostaining of adult WT retina.
(d) GRIK1 and GRM6 immunostaining of $Grik1^{-/-}$ retina.
(e) GRIK1 and GRM6 immunostaining of $Grm6^{-/-}$ retina.
(f) GRIK1 and GRM6 immunostaining of $Grik1^{-/-}$ $Grm6^{-/-}$.
White: ARR3, magenta: GRIK1 and GRM6.
(g) Representative flicker ERG raw traces of WT, $Grik1^{-/-}$, $Grm6^{-/-}$ and $Grik1^{-/-}$ $Grm6^{-/-}$
eyes at 20 and 30 Hz frequencies.
(h) Flicker ERG amplitude at 30 Hz for WT (n=13 eyes), $Grik1^{-/-}$ (n=10 eyes), $Grm6^{-/-}$
(n=12 eyes) and $Grik1^{-/-}$;$Grm6^{-/-}$ (n=10) eyes. One-way ANOVA with Dunnett
multiple comparison test.
(i) Flicker ERG implicit time at 20 Hz for WT (n=13 eyes), $Grik1^{-/-}$ (n=10 eyes),
$Grm6^{-/-}$ (n=12 eyes) and $Grik1^{-/-}$;$Grm6^{-/-}$ (n=10) eyes. One-way ANOVA with
Dunnett multiple comparison test.
(j) Representative step ERG raw traces of WT, $Grik1^{-/-}$, $Grm6^{-/-}$ and $Grik1^{-/-}$ $Grm6^{-/-}$
eyes elicited by a 3-second step light at 1000 cd/m^2 intensity.
(k) Ensemble-averaged flicker d-wave of step ERG amplitudes of WT (n=13 eyes),
$Grik1^{-/-}$ (n=10 eyes), $Grm6^{-/-}$ (n=12 eyes) and $Grik1^{-/-}$;$Grm6^{-/-}$ (n=10) eyes elicited
by a 3-second step of light-OFF at 1000 cd/m^2 intensity. One-way ANOVA with
Dunnett multiple comparison test.
1119
1120

<u>Figure 6</u>: Assay of FAT3 binding to PTPσ and PTPσ localization in WT and FAT3-
ICD mutant.
(a) Binding of FAT3-ICD to PTPσ was assayed using a pull down, with GST fused to
FAT3-ICD.

(b) Immunostaining of PTPσ and (b') CtBP2, a marker of ribbons in photoreceptor
axons.

(c) Immunostaining of PTPσ and (c') GRIK1.

(d) Immunostaining of PTPσ in WT retinas.

(e) Immunostaining of PTPσ in $Fat3^{\Delta TM/\Delta TM}$ retinas.

(f) Immunostaining of HOMER1 and SV2, and (f') HOMER1 alone in WT retinas.

(g) Immunostaining of HOMER1 and SV2, and (f') HOMER1 alone in $Fat3^{\Delta TM/\Delta TM}$
retinas.

(h) Quantification of PTPσ integrated intensity in the OPL. WT Controls: 8682 ± 583
(n=16 sections, N=4 animals); $Fat3^{\Delta TM/\Delta TM}$: 4871 ± 463.5 (n=12 sections, N=3
animals), Mann-Whitney test.

(i) Quantification of HOMER1 integrated intensity in the OPL. WT Controls: $34997 \pm$
2509 (n=8 sections, N=3 animals); $Fat3^{\Delta TM/\Delta TM}$: 32051 ± 2003 (n=8 sections, N=3
animals), t test.
Scale bars: 20µm.

1140

1141

Figure 7: Immunohistochemical assay of GRIK1 in WT and $Fat3^{\Delta TM/\Delta TM}$ retinas.

(a) Immunostaining for GRIK1 in WT retinas.

(b) Immunostaining for GRIK1 in $Fat3^{\Delta TM/\Delta TM}$ retinas. Cone arrestin (ARR3) labels
the cone photoreceptor axonal endings in the OPL in a' and b'.

(c) Quantification of GRIK1 integrated intensity in the OPL. WT Controls: $52727 \pm$
7204 (n=19 sections, N=4 animals); $Fat3^{\Delta TM/\Delta TM}$: 25838 ± 5028 (n=18 sections,
N=4 animals), Mann-Whitney test.

(d) Immunostaining for GRIK1 in WT retinas.

(e) Immunostaining for GRIK1 in $Fat3^{\Delta ICD\text{-}GFP/\Delta ICD\text{-}GFP}$ retinas.
Cone arrestin (ARR3) labels the cone photoreceptor endings in the OPL in d'
and e'.

1153 (f) Quantification of GRIK1 integrated intensity in the OPL. WT Controls: $16119 \pm$

1154   1168 (n=16 sections, N=4 animals); $Fat3^{\Delta ICD /\Delta ICD}$: 11687 ± 1110 (n=15 sections,

1155   N=4 animals), Mann-Whitney test.

1156

Figure 1

Figure 2

Figure 3

a WT

b *Ptf1a*^{CRE} *Fat3*^{cKO} (ACs)

c *Islet1*^{CRE} *Fat3*^{cKO} (pan ON-CBCs, SACs, RGCs)

Figure 4

Figure 5

Figure 6

Figure 7