Title: *In silico* analysis of the altered protein structure and functions caused by existing synaptic

gene mutations in Indian population having neurological disorders.

Pavan Kumar Divi[1,#], Nivedhitha Mukesh[1,#], Saswati Mohapatra[1], Sanjana Kanthimath[1], Nishant

Mishra[1], Karunakar Kar[2], Bibin G Anand[1,*], Anil Annamneedi[1,*]

[1]Department of Biotechnology, School of Bioengineering, SRM Institute of Science and

Technology, Kattankulathur-603203, Tamil Nadu, India

[2]Biophysical and Biomaterials Research Laboratory, School of Life Sciences, Jawaharlal Nehru

University, New Delhi-110067, India

*Correspondence: anila@srmist.edu.in; bibingna@srmist.edu.in

# Equal contribution

ORCiD ID's:

Pavan Kumar Divi: 0009-0000-0559-060X

Nivedhitha Mukesh: 0009-0004-5079-9575

Saswati Mohapatra: 0009-0009-0755-9695

Sanjana Kanthimath: 0009-0007-2217-2443

Nishant Mishra: 0009-0004-4479-417X

Karunakar Kar: 0000-0001-7047-6539

Bibin G Anand: 0000-0001-7339-6657

Anil Annamneedi: 0000-0002-6743-8627

Abstract: Neurological disorders encompass a diverse range of conditions that affect individuals

cognitive, emotional, and social functioning. Though these disorders are multifactorial, genetic

factors play a significant role in the pathogenesis. Synaptic gene mutations or synaptic protein

dysfunction are shown to be closely associated with neuropathology, suggesting the term

synaptopathies. This study aims at understanding the critical role of synaptic compartment by

conducting comprehensive analysis of existing synaptic gene mutations responsible for the

development of three significant disorders in the Indian population: Autism Spectrum Disorder

(ASD), Epilepsy and Schizophrenia. Our *in silico* analysis predicts that mutations in genes RPL10

(rs387906727), GABRA1 (rs121434579) and DRD2 (rs1801028) corresponding to ASD, Epilepsy

and Schizophrenia respectively, are deleterious. Further, molecular docking shows that the mutated

proteins resulting from these SNPs have altered protein-protein interaction with vital interacting

partners, possibly impacting neuronal physiological pathways generally affected in the above-

mentioned neurological disorders. Specifically, the abnormal interaction between D(2) dopamine

receptor and neuronal calcium sensor 1, revealed through molecular docking, may hamper

neurotransmitter regulation. Altogether, our study helps to further our understanding of synaptic

signaling in the context of the Indian population with neurological disorders and potentiates the

usage of a model and dynamic simulation systems to identify novel therapeutic targets aiding in

the development of personalized therapy.

Keywords: Neurological disorders, ASD, Schizophrenia, Epilepsy, Indian Population.

Introduction: The brain is the master regulator of the body, with several signal transduction

pathways critical in controlling all voluntary and involuntary actions, as well as other diverse

biological functions. The human brain is an advanced network that also facilitates cognition,

perception, judgement, memory, and retrieval (Greeno et al., 1996). Synaptic communication is

critical for the transfer of information and these neurological disorders that manifest due to synaptic

dysfunction are broadly termed synaptopathies (Lepeta et al., 2016). Understanding the shared

origins and fundamental distinctions in disorders linked with synaptic dysfunction could lead to

novel therapeutic approaches for neurological disorders. Genetic mutations in genes coding for

proteins ultimately change protein structure and its function and may contribute to a phenotypic

attribute of the disorder. As a result, studying synapses at the molecular level is critical in

determining etiology.

Single nucleotide polymorphism (SNP), one of the most common types of genetic diversity in the

human genome, has an important role in phenotypic variation and individual differentiation within

a species. SNPs can occur in both coding and noncoding areas of the genome, where SNPs in the

coding area are classed as synonymous or non-synonymous (nsSNPs), further classified as

missense or nonsense (Edwards et al., 2007). nsSNPs substitute protein residues, thus impacting

protein stability, dynamics, physiological molecular interactions, formation of stable complexes

with binding partners and even post-translational modifications (Yang et al., 2019). Negative

genetic variation can modify the electrostatic nature of protein surfaces, causing changes in stability

or folding; this influences binding selectivity and affinity, compromising protein function. The

identification of detrimental nsSNPs assists in the establishment of the likely biochemical pathway

causing the disease and the creation of effective inhibitors to target the mutant proteins (Dong et

al., 2021).

Our current study focuses on analyzing the existing gene mutations linked to ASD, Epilepsy and

Schizophrenia with a special focus on the Indian population and their effect on protein structure,

and function in mediating synaptic signaling. ASD and Schizophrenia are two different

neurological conditions, with distinct phenotypic changes, yet display similarities in certain

behaviors such as disturbed social cognition, altered neuroanatomy, disturbed GABAergic

functioning, and co-occurrence of Epilepsy (Jutla et al., 2022; Gao and Penzes., 2015; Canitano

and Pallagrosi., 2017; Keller et al., 2017; Cascella et al., 2009). By studying localized structural

changes and its impact on the protein's orthosteric site and protein binding hot spots, we theorize

on the mode of alteration of protein functions due to the SNPs.

MATERIALS AND METHODS:

Primary list of all mutated genes associated with neurological disorders such as ASD, Epilepsy and

Schizophrenia in the Indian population was gathered from PubMed and Google Scholar

(Supplementary Table 1). Using the SynGo database (Koopmans et al., 2019) the list of genes was

segregated into synaptic and nonsynaptic genes, and only synaptic genes were considered for

further analysis. The SNPs in synaptic genes were further sorted into exonic (synonymous and non-

synonymous) and intronic based on position of occurrence. To sift out highly impactful mutations,

Position-Specific Independent Counts Scores (PSIC) tools such as PolyPhen2 and SIFT were used

(Adzhubei et al., 2013). PolyPhen scores can be interpreted as follows: 0.0 to 0.15 - predicted to

be benign; 0.15 to 1.0 - possibly damaging. 0.85 to 1.0 - confidently predicted to be damaging.

SIFT (Sorting Intolerant From Tolerant) uses sequence homology to predict whether an amino acid

substitution will affect protein function or not. PolyPhen-2 and SIFT scores use the same range,

0.0 to 1.0, but with opposite intents. A variant with a PolyPhen-2 score of 0.0 is predicted to be

benign, whereas a variant with a SIFT score of 1.0 is predicted to be benign. We next compared

different variants corresponding to a single synaptic gene, and further compared all the synaptic

genes to identify the most deleterious gene mutation per neurological condition. The nucleotide

sequence of the respective gene variants that are disease-causing were taken from NCBI the gene

code for *RPL10* is 6314, the gene code for *GABRA1* is 2554 and for *DRD2* the gene code is 1813.

and the mutations were induced by altering the nucleotide sequence (SNP) according to the

respective mutations. Amino acid sequences were identified for the wild type (wild type) and

mutated gene-encoded protein using Mutation Taster (Schwarz et al., 2014). The three-dimensional

protein structures of the wild-type and mutant variants were generated using SWISS-MODEL

(Waterhouse et al., 2018), and sequence identity was generated between wild-type and mutant

proteins. Molecular dynamics (MD) simulations were conducted using GROMACS version 2023.1

(Spoel et al., 2005) with ff19SB (Tian et al., 2020) Amber force field and TIP3P water model. The

proteins were centered in a 10nm simulation box, solvated, and electrically neutralized by 150mM

counter ions (K+/Cl-). Bond constraints were applied using the LINCS (Hess, 2008) algorithm.

First, proteins underwent minimization, followed by 2 staged equilibrations—NVT heating to

298K and NPT equilibration employing V-rescale (Bussi et al., 2007) thermostat and Parrinello-

Rahman (Martonák et al., 2003) pressure coupling algorithms. The final 20ns production was run

at physiological temperature(310K), from which frames at 0ns, 10ns, and 20ns were extracted for

subsequent analysis using GROMACS built-in tools. Furthermore, snapshots and interactions were

visualized and analyzed using Discovery Studio Visualizer (Biovia, 2021).

The list of interacting protein partners to the wild type protein were prepared using the STRING

(Search Tool for the Retrieval of Interacting Genes/Proteins) analysis tool (Szklarczyk et al., 2023),

with highest confidence (0.900) as minimum required interaction score. To check for the binding

differences between the wild type protein and mutated proteins with their interaction partners

ClusPro (Kozakov et al., 2017) was used.

Molecular Simulation Dynamics:

Understanding the mechanisms by which mutations affect protein structure and stability is one of

the most important questions to be answered while studying the correlation between the occurrence

of a mutation and the manifestation of a disease. To validate the fidelity of the putative structure

and to study the structural perturbations in the secondary structure molecular dynamics simulations

followed by post-analysis using Ramachandran Plot analysis was done (Gromiha et al., 2002).

RESULTS:

1. Synaptic gene mutations implicated in the Indian population.

Using the SynGO (cellular component) database, we narrowed down on synaptic genes from the

larger list of genes (Supplementary Table 1) whose mutations are implicated in the Indian

population with neurological disorders (Figure 1 and Supplementary Table 2). Indeed, except for

Epilepsy conditions, the synaptic genes are localised to both pre and post synaptic compartments

in ASD and Schizophrenia.

Intronic variants, despite not contributing to amino acid substitution or changes in the coding

sequence, have been implicated in various disorders (Baskin et al., 2011). Some of these variants

are present in linkage disequilibrium with other exonic variants, whereas other intronic mutations

cause mis-splicing and pseudo-exons (Vaz-Drago et al., 2017). Surprisingly, more than 50% of the

mutations implicated in Schizophrenia, both synaptic and non-synaptic genes are intronic variants,

whereas we did not observe this trend in case of ASD and Epilepsy (Supplementary Table 3 and

4).

2.   Severity of impact of mutations on protein structure

Through the scores of both PolyPhen-2 and SIFT, we have concluded that the three genes *RPL10*,

GABRA1, *DRD2* and their mutations associated with ASD, Epilepsy and Schizophrenia,

respectively, are of highest impact potential (Table 1). All the three genes are predicted to be

damaging or deleterious based on the PolyPhen-2 and SIFT scores. Among the three genes (where

RPL10 encodes for Large ribosomal subunit protein uL16, *GABRA1* encodes Gamma-

aminobutyric acid receptor subunit alpha-1, *DRD2* codes for and D(2) dopamine receptor).

GABRA1 and *DRD2* encoded proteins are vital in neurotransmission related signal transduction

pathways and are highly impaired in the diseases they are associated with, whereas *RPL10* codes

for a ribosomal protein critical for the assembly of the polysomes (Noble, 2000; Maillard et al.,

2022; Klauck et al., 2006).

3.   Molecular Dynamics Simulation:

First, we sought to visualize the structural differences in proteins coded by mutated genes in

comparison with wild type (Supplementary Figure 1). The sequence identity for the

aforementioned proteins are as follows: Large ribosomal subunit protein uL16 - 98.9%**,** Gamma-

aminobutyric acid receptor subunit alpha-1 - 99.07%, and D(2) dopamine receptor - 99.77%.

Further, we analyzed the φ-ψ angles using Discovery Studios to determine changes, if any, in the

secondary structure. Molecular dynamics simulation was done to study the behavior of a protein

molecule in solvent, to mimic its native state. The Ramachandran plot or φ-ψ plot uses torsion

angles to describe polypeptide and protein conformation and elucidates if a conformation is

allowed or not. The plot allows us to validate the fidelity of the wild type and mutated protein of

interest and its associated protein obtained using structural biology techniques (Gromiha et al.,

2002).

Root Mean Square Deviation (RMSD), which measures the stability of a protein by determining

the protein backbone's deviation relative to its initial conformation during the course of the

simulation. Smaller deviation implies more stability of the protein structure (Maruyama et al.,

2023). Room Mean Square Fluctuation (RMSF) is used to determine the regional flexibility of the

protein. The higher the RMSF, the higher is the flexibility.

The radius of gyration (Rg) is defined as the distribution of atoms of a protein around its axis

(Lindahl, 2008). Rg is the length that represents the distance between the point when it is rotating

and at the point where the transfer of energy has the maximum effect (Sneha et al., 2016). A

relatively steady value of the Rg means stable folding of a protein and fluctuation of the Rg may

imply structural destabilization (Liu et al, 2017). Solvent Accessible Surface (SASA) is commonly

used to reflect the solvent-exposed surface area of proteins, and a variation of this parameter can

reveal the dynamic nature of the protein structure. In general, the smaller SASA indicates that the

protein molecule is in a more folded and compact structural state (Durham et al., 2009). Post

analysis of our simulation studies revealed that the *DRD2* wild-type model, in the 0ns frame the

Ser311 residue falls in the disallowed region but falls in the beta turn conformation in the 10ns and

20ns frame. In the mutant model, the Cys311 residue falls in the coil conformation region at 0ns

but shifts to β- turn region in the 10ns and 20ns frame (Supplementary Figure 2a). But despite the

secondary structure both being β-turn, the phi angle is observed to change significantly. This shows

that there are significant structural constraints due to the amino acid substitution. In the root mean

square deviation (RMSD) graph for DRD2 (Supplementary Figure 2b) after the 8ns of simulation,

there is a higher degree of fluctuation in the wild type of protein model than the mutated protein

model. This points in the direction that the mutant model shows more rigidity than the wild-type

model. In the root mean square fluctuation (RMSF) graph of DRD2 (Supplementary Figure 2c),

The Ser311 residue in the wild-type model falls in a higher peak region of the RMSF graph than

Cys311 in the mutant model, indicating relatively higher level fluctuation in its position in the

backbone in comparison with the mutant model, thus supporting the hypothesis that the mutant

model is relatively more stable than the wild type model. Such phenomena may be due to the

residues interacting spatially with free cysteines of non-membrane proteins revealed a predominant

association with hydrophobic residues and a lesser affinity for polar or charged residues

(PMC7102755) or less flexible/dynamic. The radius of gyration (Supplementary Figure 2d) and

solvent-accessible surface (Supplementary Figure 2e) for both wild type and mutant models is

similar within normal ranges of fluctuation. However, we did not find any significant alterations

in protein structure due to mutations in either RPL10 or GABRA1 (Supplementary Figure 3 and

4).

4. Impact of mutation on protein-protein interaction:

It is well understood that point mutations significantly contribute to stabilizing protein structures,

as demonstrated in studies on lysozyme (PMID: 3200317). Therefore, we anticipate that similar

mutational alterations in our protein may also impact its stability. So here in this study we have

identified the interaction partners of each of the impacted proteins in wild-type condition (Figure

2), using the STRING tool. From this list, we have assessed the most affected "interacting partner-

POI (protein of interest) interaction" (Table 2). In the case of ASD, the interaction between

RPL18A and the Large ribosomal subunit protein RPL10 is highly affected (difference in weighted

scores between wild-type and mutated conditions is -48.6), which might result in disruption of

polysome formation. For Epilepsy, the affected interaction is between GABBR2 and GABRA1

(difference in weighted scores is -375.7). We further identified that the strongly affected interaction

is between NCS1 and D(2) dopamine receptors, which is implicated in Schizophrenia (difference

in weighted scores is 890.1) (Table 2).

FREQ encodes Neuronal calcium sensor 1 protein, which is a highly conserved EF hand calcium

binding protein, a critical dopamine receptor interacting protein that has a variety of neuronal

functions like membrane traffic, cell survival, and ion channel and receptor signalling. Neuronal

calcium sensor 1 and D(2) dopamine receptors colocalize at sites of synaptic activity and this

interaction has shown to attenuate ligand-induced D(2) dopamine receptor internalisation and

receptor phosphorylation, enhancing D(2) dopamine receptor signalling (Cravchik et al., 1996;

Kabbani et al., 2002). Altogether our analysis and previous studies predict a strong influence of

genetic mutations on protein-protein interactions thus potentially affecting the neuronal functions.

5. Molecular Docking:

It is vital to analyze the change in the protein-protein interaction between the protein of interest

and its interacting partner to assess the gravity of the mutation. In RPL10, Leu206 participates in

the interaction between RPL10 and RPL18A. But post mutation, none of the initial residues,

including residue 206, interact with RPL18A. In terms of hydrogen bonding, the bonding

responsible for most directional interactions in the protein, there is an increase in number of

hydrogen bonds (20) from the initial number (5). There is also an overall increase in total number

of interacting residues post mutation, from 23 to 36. In wild type RPL10, RPL10Ala186 interacts

with both RPL18AArg161 and RPL18ALeu159 through alkyl bonding. This interaction is

completely absent in mutated RPL10. In addition, RPL10Trp209 interacts with RPL18ALeu154,

RPL18APhe152 and RPL18APro155, an interaction trio also absent in mutant RPL10- RPL18A

interaction (Supplementary Table 5a, 5b).

In GABRA1, some interactions are partially conserved pre and post-mutation i.e., the amino acid

in GABBR2 involved in the interaction with GABRA1 is the same, but the residue number

changes. Nonetheless, the residue interacting post-mutation is less than 5 residues away from the

initial interacting amino acid. But there is an increase in Pi-Alkyl interactions between GABRA1

and GABBR2, from 6 to 20 bonds, highlighting the importance of hydrophobic interactions

between two proteins. There is a reduction in number of interacting residues from 29 to 22. There

is a Phe323-Leu666-Ala322 inteteraction present in wild type GABRA1-GABBR2 interaction,

which is absent in mutant protein where Ala322 is replaced with Asp322. (Supplementary Table

6a, 6b).

In the case of DRD2, Ser311 is not involved in the interaction of the protein with NCS1. The

mutation causes structural changes dire enough that none of the residues in DRD2 wild type model

that interact with NCS1, participate in the interaction pair after mutation (Supplementary Table 7a,

7b). Interestingly, there is a formation of salt bridges observed in mutant DRD2 and NCS1, whereas

salt bridges are not observed in binding between WT DRD2 and NCS1. The introduction of salt

bridges into the binding site may have a destabilising effect on the bonding (Hendsch & Tidor,

1994). Thus, despite not being directly involved in the interaction between DRD2 and NCS1, the

Ser311Cys mutation in DRD2 shows a substantiated effect on the interaction between these

proteins. This is highlighted by the stark difference in the interacting residues of DRD2 in wild

type model and mutant model.

DISCUSSION:

Our current study elucidates the impact of existing single nucleotide polymorphisms on the overall

structure and function of proteins coded by genes whose mutations are implicated in ethnically

Indian populations having neurological disorders. Our *in silico* analysis demonstrates that known

mutations in genes *RPL10*, *GABRA1* and *DRD2*, associated with ASD, Epilepsy and Schizophrenia

respectively, are the most potentially damaging ones with respect to the protein structure and

subsequent protein functions (Figure 2). Interestingly, all the three candidates - Large ribosomal

subunit protein uL16 (coded by *RPL10*), Gamma-Aminobutyric Acid Receptor Subunit alpha-1

(coded by *GABRA1*) and D(2) dopamine receptor (coded by *DRD2*), - are essential for optimum

neurological function in the disorders that their mutations have been correlated in.

Synaptic proteins regulate ion channel localisation, synaptic vesicle cycle, positional priming and

other critical functions (Gundelfinger et al., 2012). Dysfunction in synaptic genes leads to synaptic

protein dysregulation and function. Indeed, except for Epilepsy conditions, the synaptic genes are

localised to both pre and post synaptic compartments in ASD and Schizophrenia. Thus, by studying

the mutations in these genes, we can understand the mechanism by which these mutations may

contribute to the aetiology of synaptopathies mentioned above.

Intronic variants, despite not contributing to amino acid substitution or changes in the coding

sequence, have been implicated in various disorders (Baskin et al., 2011). Some of these variants

are present in linkage disequilibrium with other exonic variants, whereas other intronic mutations

cause mis-splicing and pseudo-exons (Vaz-Drago et al., 2017). Thus, it is vital to study intronic

variations parallel to exonic variants when establishing a roster of susceptibility genes for any

disorder. Surprisingly, more than 50% of the mutations implicated in Schizophrenia, both synaptic

and non-synaptic genes are intronic variants, whereas we did not observe this trend in case of ASD

and Epilepsy (Supplementary table 3 and 4).

ASD is used to describe individuals with a specific pattern of social impairment paired with

repetitive behaviours, often with niche interests and/or sensory behaviours with an early onset.

Subtle anatomical and functional deviations have been observed in the brain through post mortem

studies despite not characterising the disorder by gross brain pathology (Lord et al., 2020). Large

ribosomal subunit protein uL16 is pivotal for joining the 60S and 40S subunits in a late step of

translation initiation, and structural data suggests that it is present in the interface of the 60s subunit

in a cleft between the central protuberance and the L7/L12 stalk in the vicinity of the A and P sites

and the catalytic trans-peptidyl transferase centre (Spahn et al., 2001). Studies showing the

missense mutation *RPL10* Leu206Met in ASD candidates suggest an impairment of a complete

translating ribosome complement, that is, the polysome population rather than basic translation,

per se (Klauck et al., 2006).

Juvenile myoclonic epilepsy, or Janz syndrome, is a prevalent form of epilepsy constituting up to

10% of all epilepsy and 18% of idiopathic generalised epilepsy (IGE) (Camfield et al., 2013). The

GABRA1 gene codes for the α1 subunit of heteropentameric gamma-aminobutyric acid A receptors

(GABAARs), which functions in mediating fast primary inhibition in the brain and is critical for

brain development. Impairment of GABAergic transmission by genetic mutations has been shown

to induce epileptic seizures (Sperk et al., 2004). Ala322Asp mutant GABAA receptors show

reduced amplitude of GABA activated currents *in vitro*, alluding that seizures may be a

consequence of loss of function of GABA gated channels (Cossette et al., 2002). Application of

GABA to HEK cells expressing receptor complexes that included the mutant subunit showed a

reduced chloride flux compared to receptors formed with wild type subunits (Noebels et al., 2002).

Hyperactivity of dopamine in the mesolimbic pathway in the subcortical region augments D(2)

dopamine receptor activation and mediates positive psychotic symptoms of Schizophrenia, where

the manifestation of the illness is divided into positive (delusions, hallucinations, thought

disorganisation, paranoia) or negative (lack of drive and motivation, alogia, social withdrawal)

symptoms, with cognitive impairments. The negative symptoms and cognitive decline are thought

to arise from reduced D(1) dopamine receptor stimulation (Shen et al., 2012; Brisch et al., 2014).

D(2) dopamine receptor regulates the release of dopamine through a negative feedback loop when

localised in the presynaptic compartment. It plays an integral role in maintaining dopamine

concentrations in the synapse, and any dysfunction can lead to hyperdopaminergia, which is

extensively discussed in the Schizophrenia hypothesis (Howes et al., 2009; Kesby et al., 2018).

We further observed that there is a significant percentage of intronic variants in the list of

implicated mutations in the synaptic genes. The intron–exon structure of eukaryotic genes has

played a major role in the creation of new genes through exon shuffling (Berk., 2016), and the

ability to alternatively select different combinations of exons has been critical for generating gene

expression diversity in more complex organisms (Keren et al., 2010). Alterations in pre-mRNA

splicing are increasingly recognized as responsible for monogenic disorders, with about 15% to

50% affecting pre-mRNA splicing (Vaz-Drago et al., 2017). Mutations affecting intronic splicing

enhancers or silencers have been shown to result in mis-regulated splicing and then in most cases

rapidly degrade (Sharp et al., 1987). Some genes are purely snoRNA or microRNA generating,

with no protein-coding capacity (Bortolin and Kiss., 1998; Flynt et al., 2010; Tycowski et al.,

1996). Recent transcriptomic analysis revealed that many introns are actively retained in

polyadenylated transcripts and contribute to down regulating gene expression (Wong et al., 2013;

Yap et al., 2012).

Molecular simulation dynamics predicts the movement of each and every atom present in a protein

in a specified timeframe, using a general model of physics governing interatomic interactions as

the template (Karplus and Mc Cammon, 2002). These simulations can replicate many vital

biomolecular processes including but not limited to conformational change, ligand binding, and

protein folding, determining the positions of the atoms at femtosecond temporal resolution. Most

importantly, MSD simulates how proteins respond at atomic levels to mutations. These simulations

are used hand in hand with experimental structural biology techniques. We also observed how the

mutation affects interacting amino acid residues of RPL10, GABRA1 AND DRD2 with the protein

of interest, there are no similar amino residues interacting in the mutated type and wild type RPL10

with its interacting partner RPL18A. In case of GABRA1 also there no common interacting

residues

In the case of RPL10 and GABRA1, using MSD we can conclude that the mutation does not

drastically impact the structure of the protein, and that the fluctuations are well within normal range

(Supplementary Figure 3a, 5a). This hints at the fact that despite being in the coding region and a

missense SNP, the mutation contributes to the disorder without destabilising the structure. In the

case of DRD2, it can be observed that surprisingly, the mutant variant is more rigid than the wild

type (Supplementary Figure 2d).

Protein-protein interactions are vital for cellular functions, and disruptions in these interactions can

lead to various disorders. For example, in ASD, PTEN-AKAP8L interaction is said to affect

neuronal growth, and both proteins are POI for ASD aetiology (Pintacuda et al., 2023). By

investigating these interactions, we gain insights into disease causes and identify potential drug

targets, offering hope for more effective treatments. Our results reveal, proteins like ion channels

and GPCRs can be majorly affected by mutations, and their interactions with other proteins present

in their vicinity within the cell or matrix are also impacted (Figure 2 and Table 2). Studying the

physiological functions and mechanisms facilitated by the interactions between the unmutated

proteins, and the impact of mutations on these protein interactions aid us in determining the

mechanism of disorder manifestation (Rao et al., 2014; Bugge et al., 2020).Our *in silico* analysis

in the current study demonstrates the effect of the mutations in different genes related to

neurological conditions on their encoded protein structures and predicted the possible alteration in

synaptic signaling, specifically the *DRD2*. In addition to its schizophrenia association, *DRD2* is

one of the strongest candidates linked to ASD, and targeting its activity might be a potential

therapeutic strategy (Annamneedi et al., 2023). Further studies are required to address the altered

interaction between Neuronal calcium sensor 1 and D(2) dopamine receptor, due to the mutation

and its relevance to both schizophrenia and ASD and in developing the common therapeutic

strategies.

Acknowledgements:

Authors acknowledge the support and facilities provided by the Department of Biotechnology,

School of Bioengineering at SRM Institute of Science and Technology, Kattankulathur, India.

Conflict of Interest:

Authors declare no conflict of interest.

Author's contributions:

P.K.D., N.M., S.M., S.K wrote the initial manuscript. P.K.D., N.M., S.M., S.K., N.M contributed

to the data mining and formal analysis. P.K.D, N.M., N.M and B.G.A contributed to bioinformatics,

simulation analysis and image processing. A.A designed and supervised the study. All the authors

contributed to the final version of the manuscript.

References:

1. Adzhubei, Ivan et al. "Predicting functional effect of human missense mutations using PolyPhen-2." Current protocols in human genetics vol. Chapter 7 (2013): Unit 7.20.
2. Annamneedi, Anil et al. "Towards the convergent therapeutic potential of G protein-coupled receptors in autism spectrum disorders." *British journal of pharmacology*, 10.1111/bph.16216. 13 Aug. 2023
3. Baskin, Berivan et al. "Duchenne muscular dystrophy caused by a complex rearrangement between intron 43 of the DMD gene and chromosome 4." *Neuromuscular disorders : NMD* vol. 21,3 (2011): 178-82.
4. Berk, Arnold J. "Discovery of RNA splicing and genes in pieces." Proceedings of the National Academy of Sciences of the United States of America vol. 113,4 (2016): 801-5.
5. Biovia, D.S. (2021) Discovery Studio Visualizer. San Diego.
6. Bortolin, M L, and T Kiss. "Human U19 intron-encoded snoRNA is processed from a long primary transcript that possesses little potential for protein coding." *RNA (New York, N.Y.)* vol. 4,4 (1998): 445-54.
7. Brisch, Ralf, et al. "The role of dopamine in schizophrenia from a neurobiological and evolutionary perspective: old fashioned, but still in vogue." Frontiers in psychiatry 5 (2014): 47.
8. Bugge, Katrine, et al. "Interactions by disorder–a matter of context." Frontiers in Molecular Biosciences 7 (2020): 110.
9. Bussi, Giovanni et al. "Canonical sampling through velocity rescaling." *The Journal of chemical physics* vol. 126,1 (2007): 014101.
10. Camfield, Carol S et al. "Epidemiology of juvenile myoclonic epilepsy." *Epilepsy & behavior : E&B* vol. 28 Suppl 1 (2013): S15-7.
11. Cascella, Nicola G et al. "Schizophrenia and epilepsy: is there a shared susceptibility?." *Neuroscience research* vol. 63,4 (2009): 227-35.
12. Canitano, Roberto, and Mauro Pallagrosi. "Autism Spectrum Disorders and Schizophrenia Spectrum Disorders: Excitation/Inhibition Imbalance and Developmental Trajectories." *Frontiers in psychiatry* vol. 8 69. 1 May. 2017
13. Cossette, Patrick, et al. "Mutation of *GABRA1* in an autosomal dominant form of juvenile myoclonic epilepsy." Nature genetics 31.2 (2002): 184-189.
14. Cravchik, Anibal, David R. Sibley, and Pablo V. Gejman. "Functional analysis of the human D2 dopamine receptor missense variants." Journal of Biological Chemistry 271.42 (1996): 26013-26017.
15. Durham, Elizabeth, et al. "Solvent accessible surface area approximations for rapid and accurate protein structure prediction." Journal of molecular modeling 15 (2009): 1093-1108.
16. Edwards, D., Forster, J.W., Chagné, D., Batley, J. (2007). What Are SNPs?. In: Oraguzie, N.C., Rikkerink, E.H.A., Gardiner, S.E., De Silva, H.N. (eds) Association Mapping in Plants. Springer, New York, NY.

17. Flynt, Alex S., and Eric C. Lai. "Biological principles of microRNA-mediated regulation: shared themes amid diversity." Nature Reviews Genetics 9.11 (2008): 831-842.

18. Gao, R, and P Penzes. "Common mechanisms of excitatory and inhibitory imbalance in schizophrenia and autism spectrum disorders." Current molecular medicine vol. 15,2 (2015): 146-67.

19. Greeno, James G., Allan M. Collins, and Lauren B. Resnick. "Cognition and learning." Handbook of educational psychology 77 (1996): 15-46.

20. Grellscheid, Sushma-Nagaraja, and Christopher WJ Smith. "An apparent pseudo-exon acts both as an alternative exon that leads to nonsense-mediated decay and as a zero-length exon." Molecular and Cellular Biology 26.6 (2006): 2237-2246.

21. Gromiha, M. Michael, et al. "Importance of mutant position in Ramachandran plot for predicting protein stability of surface mutations." Biopolymers: Original Research on Biomolecules 64.4 (2002): 210-220.

22. Gundelfinger, Eckart D., and Anna Fejtova. "Molecular organization and plasticity of the cytomatrix at the active zone." Current opinion in neurobiology 22.3 (2012): 423-430.

23. Hess, Berk. "P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation." *Journal of chemical theory and computation* vol. 4,1 (2008): 116-22.

24. Hendsch, Z. S., & Tidor, B. (1994). Do salt bridges stabilize proteins? A continuum electrostatic analysis. *Protein Science*, *3*(2), 211-226.

25. Howes, Oliver D., and Shitij Kapur. "The dopamine hypothesis of schizophrenia: version III—the final common pathway." Schizophrenia bulletin 35.3 (2009): 549-562.

26. Jutla, Amandeep et al. "Autism spectrum disorder and schizophrenia: An updated conceptual review." *Autism research : official journal of the International Society for Autism Research* vol. 15,3 (2022): 384-412.

27. Kabbani, Nadine, et al. "Interaction with neuronal calcium sensor NCS-1 mediates desensitization of the D2 dopamine receptor." Journal of Neuroscience 22.19 (2002): 8476-8486.

28. Kelemen, Olga, et al. "Function of alternative splicing." Gene 514.1 (2013): 1-30.

29. Keller, Roberto et al. "Autism, epilepsy, and synaptopathies: a not rare association." *Neurological sciences : official journal of the Italian Neurological Society and of the Italian Society of Clinical Neurophysiology* vol. 38,8 (2017): 1353-1361.

30. Keren, Hadas, Galit Lev-Maor, and Gil Ast. "Alternative splicing and evolution: diversification, exon definition and function." Nature Reviews Genetics 11.5 (2010): 345-355.

31. Kesby, J. P., Eyles, D. W., McGrath, J. J., & Scott, J. G. (2018). Dopamine, psychosis and schizophrenia: the widening gap between basic and clinical neuroscience. *Translational psychiatry*, *8*(1), 30.

32. Klauck, S. M., et al. "Mutations in the ribosomal protein gene *RPL10* suggest a novel modulating disease mechanism for autism." Molecular psychiatry 11.12 (2006): 1073-1084.

33. Koopmans, Frank et al. "SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the Synapse." Neuron vol. 103,2 (2019): 217-234.e4.

34. Kozakov, Dima et al. "The ClusPro web server for protein-protein docking." Nature protocols vol. 12,2 (2017): 255-278.

35. Lepeta, Katarzyna, et al. "Synaptopathies: synaptic dysfunction in neurological disorders– A review from students to students." Journal of neurochemistry 138.6 (2016): 785-805.

36. Lindahl, Erik R. "Molecular dynamics simulations." Molecular modeling of proteins (2008): 3-23.

37. Liu, Pinzhi, et al. "Lubricant shear thinning behavior correlated with variation of radius of gyration via molecular dynamics simulations." The Journal of chemical physics 147.8 (2017).

38. Lord, Catherine, et al. "Autism spectrum disorder." Nature reviews Disease primers 6.1 (2020): 1-23.

39. Maillard, Pierre-yves, et al. "Molecular and clinical descriptions of patients with GABAA receptor gene variants (*GABRA1*, GABRB2, GABRB3, GABRG2): A cohort study, review of literature, and genotype–phenotype correlation." Epilepsia 63.10 (2022): 2519-2533.

40. Martonák, R et al. "Predicting crystal structures: the Parrinello-Rahman method revisited." *Physical review letters* vol. 90,7 (2003): 075503.

41. Maruyama, Yutaka, et al. "Analysis of Protein Folding Simulation with Moving Root Mean Square Deviation." Journal of Chemical Information and Modeling 63.5 (2023): 1529-1541.

42. Mirdita, M., Schütze, K., Moriwaki, Y. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679–682 (2022). https://doi.org/10.1038/s41592-022-01488-1

43. Noble, Ernest P. "The *DRD2* gene in psychiatric and neurological disorders and its phenotypes." Pharmacogenomics 1.3 (2000): 309-333.

44. Noebels, Jeffrey L.. "Human Epilepsy Can Be Linked to a Defective Calcium Channel."*Epilepsy currents* vol. 2,3 (2002): 95

45. Pintacuda, Greta, et al. "Protein interaction studies in human induced neurons indicate convergent biology underlying autism spectrum disorders." Cell Genomics 3.3 (2023).

46. Rao, V. Srinivasa, et al. "Protein-protein interaction detection: methods and analysis." International journal of proteomics 2014 (2014).

47. Rong, Yu, et al. "DDRS: Detection of drug response SNPs specifically in patients receiving drug treatment." Computational and Structural Biotechnology Journal 19 (2021): 3650-3657.

48. Schwarz, Jana Marie et al. "MutationTaster2: mutation prediction for the deep-sequencing age." Nature methods vol. 11,4 (2014): 361-2.

49. Sharp, P M, and W H Li. "The codon Adaptation Index--a measure of directional synonymous codon usage bias, and its potential applications." *Nucleic acids research* vol. 15,3 (1987): 1281-95.

50. Shen, Hai-Ying, et al. "Adenosine augmentation ameliorates psychotic and cognitive endophenotypes of schizophrenia." The Journal of clinical investigation 122.7 (2012): 2567-2577.

51. Sneha, P., & Doss, C. G. P. "Molecular dynamics: new frontier in personalized medicine". Advances in protein chemistry and structural biology, 102 (2016), 181-224.

52. Spahn, Christian M T, et al. "Structure of the 80S ribosome from Saccharomyces cerevisiae—tRNA-ribosome and subunit-subunit interactions." Cell 107.3 (2001): 373-386.

53. Sperk, Günther, et al. "GABA and its receptors in epilepsy." Recent advances in epilepsy research (2004): 92-103.

54. Steudle, Friederike, et al. "A novel de novo variant of *GABRA1* causes increased sensitivity for GABA in vitro." Scientific Reports 10.1 (2020): 2379.

55. Szklarczyk, Damian et al. "The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest." Nucleic acids research vol. 51,D1 (2023): D638-D646.

56. Tian, Chuan et al. "ff19SB: Amino-Acid-Specific Protein Backbone Parameters Trained against Quantum Mechanics Energy Surfaces in Solution." *Journal of chemical theory and computation* vol. 16,1 (2020): 528-552.

57. Tycowski, Kazimierz T., Mei-Di Shu, and Joan A. Steitz. "A mammalian gene with introns instead of exons generating stable RNA products." Nature 379.6564 (1996): 464-466.

58. Van Der Spoel, David et al. "GROMACS: fast, flexible, and free." *Journal of computational chemistry* vol. 26,16 (2005): 1701-18.

59. Vaz-Drago, Rita, Noélia Custódio, and Maria Carmo-Fonseca. "Deep intronic mutations and human disease." Human genetics 136 (2017): 1093-1111.

60. Waterhouse, Andrew et al. "SWISS-MODEL: homology modelling of protein structures and complexes." Nucleic acids research vol. 46,W1 (2018): W296-W303.

61. Wong, Justin J-L., et al. "Orchestrated intron retention regulates normal granulocyte differentiation." Cell 154.3 (2013): 583-595.

62. Yang, Yang et al. "AWESOME: a database of SNPs that affect protein post-translational modifications." *Nucleic acids research* vol. 47,D1 (2019): D874-D880.

63. Yap, Karen, et al. "Coordinated regulation of neuronal mRNA steady-state levels through developmentally controlled intron retention." Genes & development 26.11 (2012): 1209-1223.

Figure legends:

Figure 1: Localization of synaptic genes. The list of genes identified from various studies conducted on the Indian population with respect to neurological conditions, are stratified into synaptic genes, both pre and post synapse using SynGO-cellular component.

Figure 2: Protein-protein interactions in wild type condition. Protein-protein interactions form the foundation of complex biological functions. Thus, we compile the interacting partners of our POI using STRING and assessed the change in binding strength between the interacting pairs. The most affected protein-POI is encircled in a box for each potential protein.

Figure 3: Molecular Docking of protein-protein interaction. (a) Protein-protein interaction molecular docking of (i) wild type RPL10 and RPL18A and (ii) mutant RPL10 and RPL18A (Supplementary Table 5). (b) Protein-protein interaction molecular docking of (i) wild type GABRA1 and GABBR2 and (ii) mutant GABRA1 and GABBR2 (Supplementary Table 6). (c) Protein-protein interaction molecular docking of (i) wild type DRD2 and NCS1 and (ii) mutant DRD2 and NCS1 (Supplementary Table 7).

Table descriptions:

Table 1: Position-Specific Independent Counts Scores. To determine if the mutation is potentially damaging, and to assess the degree of impact on the structure and function of the protein, we use PSIC software Polyphen-2 and SIFT.

Table 2: Protein-protein interaction scores (Lowest Energy). To understand the impact of mutation on protein-protein interaction, we use ClusPro and observe the change in lowest energy scores.

Figures:

1. **Synaptic gene mutations implicated in the Indian population**

Figure 1: Localization of synaptic genes. The list of genes identified from various studies conducted on the Indian population with respect to neurological conditions, are stratified into synaptic genes, both pre and post synapse using SynGO-cellular component.

2. Impact of Mutation on Protein-Protein Interaction

Figure 2: Protein-protein interactions in wild type condition. *Protein-protein interactions*
form the foundation of complex biological functions. Thus, we compile the interacting
partners of our POI using STRING and assessed the change in binding strength between
the interacting pairs. The most affected protein-POI is encircled in a box for each potential
protein.

Figure 3: Molecular Docking of protein-protein interaction. (a) Protein-protein
interaction molecular docking of (i) wild type RPL10 and RPL18A and (ii) mutant RPL10
and RPL18A (Supplementary Table 5). (b) Protein-protein interaction molecular docking

of (i) wild type GABRA1 and GABBR2 and (ii) mutant GABRA1 and GABBR2
(Supplementary Table 6). (c) Protein-protein interaction molecular docking of (i) wild type
DRD2 and NCS1 and (ii) mutant DRD2 and NCS1 (Supplementary Table 7).

Severity of impact of mutations on protein structure

DISORDER	GENE	RSID	MUTATION	POLYPHEN-2 SCORE	POLYPHEN-2 PREDICTION	SIFT SCORE	SIFT PREDICTION
ASD	*RPL10*	rs387906727	616C>A L206M	0.999	Probably Damaging	0.008	Deleterious
Epilepsy	*GABRA1*	rs121434579	965C>A Ala322Asp	0.992	Probably Damaging	0.035	Deleterious
Schizophrenia	*DRD2*	rs1801028	932C>G Ser311Cys	0.999	Probably Damaging	0.027	Deleterious

574   ***Table 1: Position-Specific Independent Counts Scores.*** *To determine if the mutation is*
575   *potentially damaging, and to assess the degree of impact on the structure and function of*
576   *the protein, we use PSIC softwares Polyphen-2 and SIFT.*

578 **1. Impact of Mutation on Protein-Protein Interaction**

Disease	Protein of Concern	Interacting Protein (Gene)	Weighted Score- Wild Type Protein	Weighted Score - Mutated Protein	Difference
ASD	Large ribosomal subunit protein uL16 (rs782521991)	RPL13	-887.8	-888.4	0.6
		RPL15	-765.6	-765.9	0.3
		RPL18A	-844.7	-789.3	-55.4
		RPL19	-727.2	-680.1	-47.1
		RPL35	-680.4	-654	-26.4
		RPL8	-904.2	-856.5	-47.7
		RPS11	-741.4	-741	-0.4
		RPS3	-1049.8	-1050.3	0.5
Epilepsy	Gamma-aminobutyric acid receptor subunit alpha-1 (rs121434579)	GABBR2	-1704.2	-1328.5	-375.7
		GABRB2	-1733.3	-1639	-94.3
		GABRB3	-1919.4	-1879.9	-39.5
		GABRG2	-1699.8	-1707.6	7.8
		HAP1	-1967.2	-1630.7	-336.5
		PLCL1	-1481.2	-1476.3	-4.9
Schizophrenia	D(2) dopamine receptor (rs1801028)	ADORA2A	-2139.6	-2140.8	1.2
		ARRB1	-1280.2	-1285.2	5
		ARRB2	-1828	-1828.1	0.1
		GNAI1	-1301.7	-1301.1	-0.6
		GNB1	-1157.1	-1157	-0.1
		NCS1	-1333.4	-2223.5	890.1

579 *Table 2: Protein-protein interaction scores (Lowest Energy).* To understand the impact of
580 *mutation on protein-protein interaction, we use ClusPro and observe the change in lowest*
581 *energy scores.*