Feeding Behavior Modifies the Circadian Variation in RR and QT intervals by Distinct Mechanisms in Mice.

Makoto Ono[1], Don E. Burgess[1], Sidney R. Johnson[1], Claude S. Elayi[2], Karyn A. Esser[3], Tanya S. Seward[1], Carie R. Boychuk[4], Andrés P. Carreño[1], Rebecca A. Stalcup[1], Abhilash Prabhat[1], Elizabeth A. Schroder[1,5*], Brian P. Delisle[1*]

[1]Department of Physiology, University of Kentucky, Lexington, KY, USA
[2]CHI Saint Joseph Hospital, Lexington, KY, USA
[3]University of Florida, Gainesville, FL, USA
[4]Dalton Cardiovascular Research Center, Department of Biomedical Sciences, University of Missouri, Columbia, MO, USA
[5]Department of Internal Medicine, University of Kentucky, Lexington, KY, USA

*Contributed equally to the study

Running Title: Shaping 24-hour Rhythms in RR and QT Intervals

Corresponding authors:

Elizabeth A. Schroder
Department of Internal Medicine
859-323-5571
eschr0@uky.edu
780 Rose Street, Medical Science Building, Rm: MN-512, Lexington, KY 40536-0298

Brian P. Delisle
(859) 323-2797
 brian.delisle@uky.edu
741 S Limestone Street BBSRB B353, Lexington, KY, 40536

Word Count: 7,974

Abstract

Rhythmic feeding behavior is critical for regulating the phase and amplitude in the ≈24-hour variation of the heart rate (RR intervals), ventricular repolarization (QT intervals), and core body temperature in mice. We hypothesized the changes in cardiac electrophysiology associated with feeding behavior were secondary to changes in core body temperature. Telemetry was used to record electrocardiograms and core body temperature in mice during ad libitum-fed conditions and after inverting normal feeding behavior by restricting food access to the light cycle. Light cycle-restricted feeding quickly modified the phase and amplitude of the 24-hour rhythms in RR intervals, QT intervals, and core body temperature to realign with the new feeding time. Heart rate variability analysis and inhibiting β-adrenergic and muscarinic receptors suggested that the changes in the phase and amplitude of the 24-hour rhythms in RR intervals were secondary to changes in autonomic signaling. In contrast, the changes in the QT intervals closely mirrored changes in core body temperature. Studies at thermoneutrality confirmed the daily variation in the QT interval, but not the RR interval, and reflected daily changes in core body temperature (even in ad libitum-fed conditions). Correcting the QT interval for differences in core body temperature helped to unmask QT interval prolongation after starting light cycle-restricted feeding and in a mouse model of long QT syndrome. We conclude feeding behavior alters autonomic signaling and core body temperature to regulate the phase and amplitude in RR and QT intervals, respectively.

Abbreviations:

ECG - Electrocardiogram
pHF - high-frequency power
pLF - low-frequency power
SD - standard deviation
WT - Wild type
$QT_{37°C}$ - QT corrected to $37°C$
ZT - Zeitgeber Time

New and Noteworthy: We used time-restricted feeding and thermoneutrality to demonstrate that different mechanisms regulate the 24-hour rhythms in heart rate and ventricular repolarization. The daily rhythm in heart rate reflects changes in autonomic input, whereas daily rhythms in ventricular repolarization reflect changes in core body temperature. This novel finding has major implications for understanding 24-hour rhythms in mouse cardiac electrophysiology, arrhythmia susceptibility in transgenic mouse models, and interpretability of cardiac electrophysiological data acquired in thermoneutrality.

Introduction

Organisms have evolved circadian rhythms in physiology to anticipate changes during the 24 hours in the external environment.(1) In mammals, circadian rhythms are generated at the cellular level by the circadian clock mechanism, a transcription-translation feedback loop that cycles with a periodicity of ≈24 hours.(2, 3) The circadian clock mechanism in the suprachiasmatic nucleus (SCN) of the hypothalamus entrains to the light cycle, and light-entrained SCN neurons signal to neighboring SCN neurons and other brain nuclei to align neurohumoral signaling, behavioral rhythms, and peripheral circadian clocks throughout the body.(4, 5) Several studies suggest SCN signaling modifies autonomic input to the heart to generate circadian variations in cardiac electrophysiology (i.e., RR and QT intervals).(6, 7) Studies also show that the circadian clock mechanism in cardiomyocytes contributes to 24-hour rhythms in the RR and QT intervals by altering the expression of genes, proteins, and different cardiac ionic currents.(8-14)

Feeding behavior is an important timekeeping cue for aligning circadian clocks in peripheral tissues, including the heart. Mouse studies reveal that restricting feeding to the inverse time of normal feeding behavior (i.e., light cycle-restricted feeding) realigns the phase of circadian mRNA expression in the heart and 24-hour rhythms in cardiac electrophysiology to the new feeding behavior.(15, 16) The effect of light cycle-restricted feeding on the realignment of 24-hour rhythms in cardiac electrophysiology is not secondary to changes in the circadian clocks in the SCN or cardiomyocytes. The reason is the circadian clock in the SCN remains entrained to the light cycle during light cycle-restricted feeding(17), and genetic disruption of the cardiomyocyte circadian clock by inducing deletion of BMAL1 does not prevent realignment of 24-hour rhythms in cardiac electrophysiology with feeding.(18) This study determined how feeding behavior impacts 24-hour rhythms in the phase and amplitude of the RR and QT intervals in mice.

Several studies have demonstrated a strong temperature dependence in the RR and QT interval.(19-29) In addition, previous studies have shown that light cycle-restricted feeding in mice alters the phase and amplitude of the 24-hour rhythm in core body temperature.(17, 30) This raises the intriguing possibility that feeding modifies 24-hour cardiac electrophysiology rhythms by altering core body temperature rhythms.

In this study, we tested whether light-cycle restricted feeding similarly altered the phase and amplitudes of the 24-hour rhythms of the RR intervals, QT intervals, and core body temperature. We found that the changes in the 24-hour rhythm of the RR interval during light cycle-restricted feeding were only partially explained by changes in the core body temperature, and pharmacological studies showed the 24-hour rhythm in the RR interval was strongly influenced by autonomic signaling. In contrast, the changes in the 24-hour rhythm in the QT interval during light cycle-restricted feeding closely mirrored changes in the core body temperature. Correcting the QT interval for core body temperature was important in identifying abnormal changes in the QT interval in transgenic mouse models of long QT syndrome. We conclude that feeding behavior aligns the 24-hour rhythms in the RR and QT intervals by distinct mechanisms in mice.

Methods

Animals. The mice used in this study were male on the 129S6 background strain. The 129S6 mouse strain has been previously characterized in the ΔKPQ mouse model ($Scn5a^{+/\Delta KPQ}$ mice).(31, 32) For studies comparing the genotype-specific differences, age-matched (wild type, WT) and $Scn5a^{+/\Delta KPQ}$ littermates were used. Mice were fed Teklad Global 18% protein rodent diet (2018). All procedures complied with the Association for Assessment and Accreditation of Laboratory Animal Care guidelines and were approved by the Institutional Animal Care and Use Committee at the University of Kentucky. Dr. Peter Carmeliet kindly provided the $Scn5a^{+/\Delta KPQ}$ mouse strain.(33) This study represents new and re-analysis of data collected previously.(15) All studies were performed at room temperature ($23 \pm 2^{\circ}C$) except for data presented in **Figure 7**. These studies were done in mice housed in thermoneutral temperatures ($30 \pm 2^{\circ}C$).(34) After the study, mice were euthanized by anesthetic overdose followed by cervical dislocation.

Telemetry. We used telemetry to measure electrocardiogram (ECG), core body temperature, and locomotor activity. Mice were anesthetized with isoflurane (2-4%) using a nose cone attached to an anesthetic gas vaporizer and implanted with PhysioTel ETA-F10 (Data Sciences International) telemetry transmitter units. ECG telemetry data were recorded at 1000 Hz, and RR, QT, and PR interval analyses were done using MATLAB (MathWorks), similar to those previously described.(8, 15, 18) ECG traces recorded for each hour were aligned to the peak of the R wave to generate an ensemble ECG trace. The beginning of the P wave was defined as the point where the slope of the P wave profile changed by more than 0.001 mV/ms; the Q wave was defined as the point where the slope of the profile changed from negative to positive at the base of the QRS complex; the end of the T wave was defined as the point of intersection of T wave maximum slope with baseline.(35)

Statistical analysis. For some graphs, data are shown as the mean and standard deviation of the mean (SD). For other graphs, a data point for each animal is shown with a horizontal bar at the mean. The statistical JTK_CYCLE package confirmed significant cyclical rhythms at $p<0.05$ before analyzing the data using cosine waveform curve fitting.(36). Data were analyzed in PRISM (GraphPad). Details of statistical analyses for each experiment are reported in the figure legends.

Results

Experiment 1: Determine how long it takes for light cycle-restricted feeding to alter the 24-hour rhythms in cardiac electrophysiology and core body temperature.

Three to four-month-old WT mice housed in 12-hour light/dark cycles at room temperature with ad libitum access to water were used in this study. To initiate light cycle-restricted feeding, food was removed at 9 h after the beginning of the light cycle (Zeitgeber time or ZT = 9). The next day, food presentation was restricted to 7 h during the light cycle, between ZT 2–9 for two weeks. Light cycle-restricted feeding caused mice to lose body mass during the first week (**Figure S1**). However, caloric intake increased after that, and similar to previous reports, light cycle-restricted-fed animals re-gained body mass after two weeks.(37)

We used *in vivo* telemetry to longitudinally record electrocardiogram (ECG) data
from mice before and after switching to light cycle-restricted feeding (**Figure 1A**). Data
recorded shortly after starting light cycle-restricted feeding revealed immediate impacts
on cardiac electrophysiology. Data recorded two weeks after beginning light cycle-
restricted feeding indicated the steady state changes.(17) The top, left panel in **Figure**
1B shows a three-day time series for changes in the hourly mean heart rate (RR
interval), the duration of ventricular repolarization (QT interval), and the duration of atrial
to ventricular conduction (PR interval) in ad libitum-fed conditions. The time series
showed that the amplitude in the daily rhythms in RR, QT, and PR intervals peaked at a
similar time (i.e., had similar acrophases) during the light cycle. However, a new 24-
hour rhythm in the ECG data emerged one day after starting light cycle-restricted
feeding. The new 24-hour rhythms had larger amplitudes and peaked in the dark cycle.
The changed 24-hour rhythms in the ECG data after starting light cycle-restricted
feeding were not transitory and persisted for two weeks.

Previous studies show that light cycle-restricted feeding causes analogous changes
in daily core body temperature rhythms.(17, 38) We found that light cycle-restricted
feeding changed the daily core body temperature rhythm in a way that largely mirrored
the changes in the daily rhythms in RR, QT, and PR intervals (**Figure 1B**).

We quantified the changes in the 24-hour rhythms in cardiac electrophysiology and
core body temperature before and after starting light cycle-restricted feeding. Almost all
of the individual ECG and core body temperature time series recorded from ad libitum-
fed mice, one day after starting light cycle-restricted feeding, and two weeks after
starting light cycle-restricted feeding had significant cyclical rhythms with a periodicity of
≈24 hours (i.e, JTK analysis showed p<0.05) (see **Supplemental Table 1**).

The hourly mean data for RR, QT, and PR intervals and core body temperature for each
mouse were fit with the cosine curve function:

115   Interval = $A*\cos[2\pi*t/T - \emptyset] + m$

to calculate the period (T), the time between the peak amplitudes; acrophase ($\emptyset$), the
phase of the peak amplitude in relation to the start of the light cycle (ZT = 0); the
amplitude (A), one-half of the peak to trough; and the mesor (m), the rhythm adjusted
mean (calculated data are presented in **Supplement Table 1**).(8, 15, 18)

Data showed that light cycle-restricted feeding caused large changes in the
acrophase, amplitude, and mesor of daily rhythms in RR intervals, QT intervals, PR
intervals, and core body temperature (**Figures 2A, 2B**). No or very small additional
changes were seen in the cyclical ECG and core body temperature rhythms two weeks
after starting light cycle-restricted feeding.

We conclude only one day of light cycle-restricted feeding causes large, simultaneous,
and abrupt changes in the phase, amplitude, and mesor in the ≈24-hour rhythms of
cardiac electrophysiology and core body temperature.

Experiment 2: Light cycle-restricted feeding impacts the 24-hour rhythm in the RR
interval by modifying core body temperature and autonomic input to the heart.

We first estimated how much of the 24-hour rhythm in the RR interval is expected to be
driven by daily change in core body temperature (RR$_{\text{core body temperature}}$ interval). We used
a Q$_{10}$ value based on the Q$_{10}$ value reported in isolated WT mouse hearts.(23) For the
RR$_{\text{core body temperature}}$ interval calculation, we used a reference core body temperature of
37°C and an RR interval of 116 ms because these were the mean values in WT mice in
ad libitum-fed conditions.

$$\text{RR}_{\text{core body temperature}} \text{ interval} = 116 \text{ ms} * (0.40)^{((\text{core body temperature}-37.0)/10)}$$

We directly compared the estimated RR$_{\text{core body temperature}}$ interval with the RR interval
measured from WT mice in ad libitum-fed conditions, immediately after starting light
cycle-restricted feeding, and two weeks after starting light cycle-restricted feeding
(**Figure 3**). The data suggest that the RR$_{\text{core body temperature}}$ interval represented a
significant portion of the amplitude in the RR interval during ad libitum-fed conditions.
However, a large difference between the RR$_{\text{core body temperature}}$ and RR interval emerged
during the dark cycle one day after starting light cycle-restricted feeding (**Figure 3A**),
suggesting that additional factors contributed to the amplitude of the 24-hour rhythm in
RR intervals one day after starting light cycle-restricted feeding.

The 24-hour rhythm in the RR interval is classically thought to reflect changes in
autonomic input to the heart and activity.(6, 39, 40) We quantified changes in autonomic
input to the heart by measuring changes in the heart rate variability from mice in ad
libitum-fed conditions and after switching to light cycle-restricted feeding. We followed
the approaches outlined by Fenske et al. (41). Interpolated RR interval time series were
generated for spectral analysis and determination of frequency domain components for
heart rate variability analyses. In mice, the frequency range to classify the high-
frequency (HF) component is 1.5-4.0 Hz, and the frequency range used to classify the
low-frequency (LF) component is 0.4-1.5 Hz (41) The HF component is generally
accepted to correspond to the parasympathetic outflow related to respiratory sinus
arrhythmia, and the LF component is thought to primarily reflect autonomic outflow
associated with the baroreceptor reflex.(42)
158
159 Power Spectral Density $\mathcal{P}(f)$ has units of power per frequency (ms^2/Hz). The HF and
160 LF components represent the power within a frequency band and are related to the
161 integration:
162

$$P_{band} = \int_{f_1}^{f_2} \mathcal{P}(f)df$$

164 The units of the differential cancel the frequency units, so the units for the HF and LF
165 components are in ms^2. Normalization is based on the discrete version of Parseval's
166 theorem. It allows us to consistently approximate the integration for LF and HF
167 components of the variance for the underlying RR time series.(43)
168
In ad libitum-fed conditions, the mean HF component was higher during the light
cycle than the dark cycle. One day after starting light cycle-restricted feeding, the mean
HF component increased during the dark cycle, and this persisted two weeks after

starting light cycle-restricted feeding (**Figure 3A, 3B**). Similar trends were seen for the
LF component but did not reach statistical significance (**Figure 3S**).

In ad libitum-fed conditions, the mean activity was lower during the light cycle than
the dark cycle. This trend persisted initially after starting light cycle-restricted feeding,
but the difference between the light and dark cycle activity levels disappeared after two
weeks.

The individual mouse three-day time series data for the HF and activity components
did not show significant cyclical rhythms for most mice (JTK analysis P>0.05), and the
data were not well described using cosine waveform curve fitting. Therefore, we
analyzed the relative rhythmicity between the RR interval and the HF component or
activity using cross-correlation analyses.(44) As part of the analysis, we included the
cross-correlation between RR interval and core body temperature. We used the R
function "ccf" to calculate the cross-correlation.(45) The formula used in "ccf" to
calculate the cross-correlation between two time series $X(t), Y(t)$ has the form

$$c_{xy}(t) = \frac{1}{n} \sum_{s=\max(1,-t)}^{\min(n-t,n)} [X(s+t) - \bar{X}] [Y(s+t) - \bar{Y}], \; r_{xy}(t) = \frac{c_{xy}(t)}{c_{xy}(0)}$$

Figure 4 shows the cross-correlation for RR-core body temperature, RR-HF, and
RR-activity from mice in ad libitum-fed conditions and after starting light cycle-restricted
feeding. The cross-correlation between the RR-core body temperature and RR-HF
showed strongly correlated rhythmicity that peaked at a lag of 0 in all three conditions,
but the RR-activity cross-correlation only showed correlated rhythmicity in ad libitum-fed
conditions. The data suggested that altered autonomic signaling and not activity may be
important for modifying the daily rhythm in the RR interval during light cycle-restricted
feeding.

We directly tested the involvement of autonomic signaling by doing intraperitoneal
injections of propranolol hydrochloride and atropine sulfate (10 mg/kg each) in six to
eight-month-old telemetered WT mice undergoing light cycle-restricted feeding.(46-48)
Propranolol and atropine inhibit β-adrenergic and muscarinic receptor activation,
respectively. We injected the mice at ZT 23.5 (when the daily rhythm in the RR interval
peaked) and 9.5 (when the RR interval was at its nadir) for two consecutive days. The
feeding onset was delayed one hour to ZT 3 to separate the food anticipatory decrease
in RR interval from the start of the light cycle. Propranolol and atropine injections
caused a loss in the 24-hour rhythms of the RR interval, core body temperature, and the
difference between the RR_core body temperature interval and RR interval (**Figure 5A, 5B, and
5C**).

These data suggest light cycle-restricted feeding modifies the 24-hour rhythm in the
RR interval by altering core body temperature and autonomic input to the heart.

Experiment 3: The 24-hour rhythm in the QT interval during light cycle-restricted feeding
primarily reflects changes in the core body temperature.

The time series data shows that longitudinal changes in the hourly mean QT interval
occur in parallel to the changes in the RR interval and mirror changes in core body
temperature (**Figure 1**). The duration of the QT interval in mice is thought to be
independent of the RR interval.(9, 49-51) Therefore, we quantified the relation between
the QT interval and core body temperature. We used an approach similar to that
described previously to correct the heart rate corrected QT interval to core body
temperature in beagles.(29) The hourly mean QT intervals were plotted against the
corresponding core body temperature for all data collected in ad libitum- and light cycle-
restricted fed mice, and we identified a strong negative linear relationship between the
mean QT interval and core body temperature (**Figure 6A**).(29) The average slope of the
linear QT-core body temperature relation was used to derive a QT-core body
temperature correction formula. We corrected the QT intervals to 37°C (QT$_{37^\circ C}$ interval)
because 37°C was the daily mean in the core body temperature for ad libitum-fed mice.

223   QT$_{37^\circ C}$ = QT interval-(4.3*(37°C-core body temperature))

Correcting the time series of the 24-hour rhythm in the QT interval for core body
temperature caused a loss in the 24-hour rhythm in the QT interval for mice in ad libitum
and light cycle-restricted feeding (**Figure 6B**).

We confirmed the 24-hour rhythm in the QT interval reflected changes in core body
temperature (and not the RR interval) by studying the ECG properties of the mice
housed in thermoneutrality under ad libitum-fed conditions. Studying mice in
thermoneutrality alters autonomic tone to dramatically slow the basal heart rate without
causing large changes in the daily rhythm of the core body temperature.(52) Housing
mice at thermoneutrality slowed the RR interval and generated a large difference in the
RR$_{core\ body\ temperature}$ not seen in ad libitum-fed mice housed at room temperature
(compare **Figure 3, top left panel** to **Figure 7A**). In contrast, both the QT and QT$_{37^\circ C}$
intervals measured in thermoneutrality were qualitatively similar to the QT and QT$_{37^\circ C}$
intervals measured at room temperature (compare **Figure 6B**, **left panel** to **Figure 7A**).
The QT$_{37^\circ C}$ interval in mice housed at thermoneutrality also generated a horizontal line
when plotted as a function of core body temperature (**Figure 7B**). We plotted the hourly
mean QT intervals against the corresponding core body temperature. The negative
linear relationship between the mean QT interval and core body temperature was
similar to mice housed at room temperature (compare **Figure 6A and 7C**).(29)

These data support the concept that the 24-hour rhythms in the RR and QT intervals
are regulated by distinct mechanisms- even in ad libitum fed mice. Specifically, the 24-
hour rhythm of the RR interval is sensitive to changes in core body temperature and
autonomic signaling, whereas the 24-hour rhythm of the QT interval primarily reflects
changes in the core body temperature.

Experiment 4: Determine the impact of core body temperature on the QT interval in a
transgenic mouse model of long QT syndrome.

We previously showed that light cycle-restricted feeding for two weeks prolonged the
RR and QT intervals in *Scn5a*$^{+/\Delta KPQ}$ mice. However, the QT intervals in this study were
not corrected for core body temperature.(15) We quantified the ECG properties and
core body temperature measured in *Scn5a*$^{+/\Delta KPQ}$ mice during ad libitum-fed conditions

and after starting light cycle-restricted feeding. The changes in the 24-hour rhythm of
RR interval, QT interval, PR interval, and core body temperature were similar to those
recorded in WT mice (**Figures 1 and 3S**). Cosine fitting analysis of ECG data from the
$Scn5a^{+/\Delta KPQ}$ mice showed that, one day after starting light cycle-restricted feeding, there
was an immediate change in the phase, increase in amplitude, and change in the mesor
for 24-hour rhythms in the RR intervals, QT intervals, PR intervals, and core body
temperature (**Figure 4S**). We compared the $QT_{37°C}$ interval for WT and $Scn5a^{+/\Delta KPQ}$
mice in ad libitum-fed conditions and after starting light cycle-restricted feeding (**Figure**
8A). The $QT_{37°C}$ interval was not different between WT and $Scn5a^{+/\Delta KPQ}$ mice in ad
libitum-fed conditions (**Figure 8B**). After starting light cycle-restricted feeding, the $QT_{37°C}$
interval became longer in WT and $Scn5a^{+/\Delta KPQ}$ mice. Light cycle-restricted feeding also
unmasked a genotype-specific difference in $QT_{37°C}$ intervals measured from WT and
$Scn5a^{+/\Delta KPQ}$ mice (**Figure 8B**).

Discussion

We determined the mechanisms that underlie the changes in the 24-hour rhythms of
the RR and QT intervals during light cycle-restricted feeding. We found that one day
after starting light cycle-restricted feeding, the 24-hour rhythms in the RR intervals, QT
intervals, PR intervals, and core body temperature changed to a new phase with a
larger amplitude (**Figures 1B, 2A, 2B**). The Changes in the 24-hour rhythm of the RR
interval correlated with changes in heart rate variability (**Figures 3, 4**), and inhibiting β-
adrenergic and muscarinic receptor activation in light cycle restricted-fed mice caused a
loss in the 24-hour rhythm of the RR interval (**Figure 5**). The 24-hour rhythm in the QT
interval showed a strong linear relation with core body temperature (**Figure 6**), and
studies in ad libitum-fed mice housed in thermoneutrality confirmed that the 24-hour
rhythm in the QT interval primarily reflected changes in core body temperature (not the
RR interval) (**Figure 7**). Correcting the QT interval for core body temperature allowed us
to identify abnormal QT interval prolongation in a transgenic mouse model of long QT
syndrome (**Figure 8**).

Feeding behavior impacts the 24-hour rhythm in the RR interval (heart rate) by
modifying autonomic input to the heart.

Studies suggested that the circadian clock in the SCN modified autonomic signaling
to generate the 24-hour rhythm in heart rate.(6, 53-56) These studies were supported
by data showing that ablation of the SCN or transgenic mice lacking the core circadian
clock transcription factor *Bmal1* (e.g., *Bmal1*$^{-/-}$) did not have a 24-hour rhythm in heart
rate.(57-59) Light cycle-restricted feeding is not thought to impact the phase of the
circadian clock mechanism in the SCN;(17) therefore, it was surprising that light cycle-
restricted feeding caused a significant phase shift in the 24-hour rhythm of heart
rate.(15) Light cycle-restricted feeding in mice also shifts the phase of circadian gene
expression in peripheral tissues, including the heart, to align with the timing of feeding
behavior.(17) Several studies implicate the cardiomyocyte circadian clock in regulating
gene expression essential for cardiac excitability, cardiac electrophysiology, and heart
rate.(8-12, 58, 60) One might predict that light cycle-restricted feeding shifts the phase
of cardiomyocyte gene expression to facilitate changes in 24-hour rhythms of heart rate.
However, genetically disrupting the cardiomyocyte-specific circadian clock (by inducing

the deletion of *Bmal1*) did not abolish the light cycle-restricted feeding-mediated phase
shift in the 24-hour rhythm in heart rate.(18) We conclude that autonomic receptor-
mediated signaling largely shapes the 24-hour rhythm in heart rate during light cycle-
restricted feeding.

The 24-hour rhythm in the QT depends on core body temperature.

We now report a robust linear QT-core body temperature relation in mice, and
correcting the QT interval for core body temperature caused a loss in the 24-hour
rhythm of the QT interval. These new data suggest the changes in the 24-hour rhythm
of the QT interval in mice during light cycle-restricted feeding primarily reflect daily
changes in core body temperature (not the RR interval). The idea that the 24-hour
rhythm in the QT interval in mice primarily reflects core body temperature is also
supported by a study showing that the 24-hour rhythm of the QT interval in mice is
independent of the RR interval.(9) core body temperature was not reported in the study,
but the observed change in the amplitude of the daily rhythm in the QT interval (4-5 ms)
is consistent with a core body temperature amplitude typically seen in mice (1°C).(61)

The QT interval is used clinically in humans to assess the risk for certain deadly
ventricular arrhythmias.(62) We extended our light cycle-restricted feeding studies to
quantify changes in the QT interval in a transgenic knock-in mouse model that
expresses a mutation linked to long QT syndrome in people.(33) Correcting the QT
interval for core body temperature allowed direct comparisons of QT intervals at
different core body temperatures across the 24-hour cycle, identified genotype-specific
differences in the QT interval independent of core body temperature, and showed that
light cycle-restricted feeding caused a prolongation of the QT interval in WT and
Scn5a$^{+/\Delta KPQ}$ mice. These studies underscore the importance of monitoring and
considering the influence of core body temperature when measuring QT intervals.

An important question is whether these data may be clinically relevant to humans.
We propose that the daily rhythm in core body temperature also contributes to the 24-
hour rhythm in the heart rate corrected QT (QTc) interval observed in people. Several
studies report a temperature dependence of the QTc interval in people is 10-15 ms for
every 1°C change in core body temperature.(24-29) The peak-to-trough amplitude in the
daily core body temperature rhythm in people is about 1°C, and the peak-to-trough
amplitude in the 24-hour rhythm in the QTc interval is 10-20 ms.(63, 64) Future studies
that measure the QT interval and core body temperature across the 24-hour cycle in
people will help delineate how much of the 24-hour rhythms in the QTc interval reflect
temperature-dependent changes in the properties of ventricular repolarization.
Consideration of the influence of core body temperature on the QTc interval is
warranted when the core body temperature might be different due to physiological,
behavioral, environmental, or pharmacological changes (for example, patients with
hypothermia or sepsis).

Implications

Modern lifestyles disrupt evolutionarily conserved behavioral and environmental
timekeeping cues critical for maintaining healthy circadian homeostatic
mechanisms.(65) Our findings suggest that feeding behavior is a critical timekeeping

cue for daily core body temperature and cardiac electrophysiology rhythms in mice.
However, we did not see feeding as being a critical timekeeping cue for activity rhythms.
This suggests that eating at an inverse time may cause misalignment between 24-hour
rhythms in core body temperature, cardiac electrophysiology, and activity rhythms. If
conserved, modern lifestyles that disrupt feeding rhythms in people may negatively
impact the alignment between core body temperature, cardiac electrophysiological, and
activity rhythms to impair cardiac performance and influence arrhythmogenic risk.

Another implication of this study is that it underscores the importance that daily
rhythms in core body temperature have in shaping 24-hour rhythms in ventricular
repolarization. Continuous core body temperature monitoring is not routinely performed
in people undergoing electrophysiological testing despite its contribution to heart rate
and QTc intervals. This is especially true in studies that monitor changes in heart rate
and QTc intervals across the 24-hour cycle.

Limitations

Observing a temperature dependence on the QT interval is not necessarily new and
has been previously reported.(29) What is new is data suggesting the daily change in
core body temperature is primarily responsible for generating a 24-hour rhythm in the
QT interval. Ad libitum feeding may not be the best control for restricted light-cycle
feeding. A dark cycle-restricted feeding control would rule out any potential effect of
food consumption during the light phase seen in ad libitum-fed mice. Fasting mice
makes them vulnerable to torpor (a transient hibernating-like state that causes a
significant drop in heart rate and core body temperature below 32°C.(22) Our model of
light cycle-restricted feeding did not calorically restrict the mice, and the mice continued
to eat 3-4 grams of food per day during light cycle-restricted feeding (**Figure 1S**) and
gained mass after one week (**Figure 1S**). Another limitation is that this study used only
male mice. Results in female mice may differ. The mice used in the pharmacological
study were older than those used in the initial light-cycle restricted feeding study.
Whether a long $QT_{37^{\circ}C}$ interval is associated with an increased risk for spontaneous
ventricular arrhythmias remains unclear. We found that, during light cycle-restricted
feeding, the daily rhythm in the PR interval simultaneously shifted with the RR and QT
intervals and core body temperature. However, due to the small absolute amplitude in
the PR interval in response to light cycle-restricted feeding, it was difficult to determine if
the PR interval reflected core body temperature, autonomic regulation of the heart, or
both. The analyses in these studies focused on studying hourly mean values and 24-
hour rhythms. We did not quantify the data at higher resolutions or study possible
ultradian rhythms. Many cardiac electrophysiological channels that underlie the
electrical rhythm of the heart, specifically ventricular repolarization, are not only
temperature-sensitive but also time-of-day sensitive. Future studies examining the
expression/activity of key components and circadian clock/autonomic nervous system
markers in light cycle restricted feeding are warranted. Lastly, caution must be used
when extrapolating data from mice to humans.(51, 66)

Summary

Daily rhythms in cardiac electrophysiology align with feeding and activity cycles via
signals originating from the SCN. Restricting food availability to the light cycle disrupts

the association between environmental and behavioral timekeeping cues, resulting in
the alignment of the daily rhythms of cardiac electrophysiology with the new feeding
behavior. The alignment of the 24-hour rhythm in the RR interval primarily reflects
changes in autonomic input to the heart. In contrast, the phase of the QT interval
mirrors changes in core body temperature. Thermoneutrality studies suggest the distinct
regulation of the 24-hour rhythms in RR and QT intervals also occurs in ad libitum-fed
mice. We conclude that feeding-induced changes in daily autonomic signaling and core
body temperature rhythms facilitate the alignment of 24-hour rhythms in heart rate and
ventricular repolarization.

Competing interests
None declared.

Author contributions
All authors contributed to drafting the work and revising it critically for important
intellectual content, approved the final version of the manuscript, and qualified for
authorship.

Funding
This work was supported by National Heart Lung and Blood Institute grants
R01HL153042 and R01HL141343. This publication was supported by the National
Center for Research Resources and the National Center for Advancing Translational
Sciences, National Institutes of Health, through Grant UL1TR001998**.** The content is
solely the responsibility of the authors and does not necessarily represent the official
views of the NIH.

Data Availability
Data will be made available upon reasonable request.

Acknowledgment
We want to thank Drs. Wayne Giles and Chao-Yin Chen (University of California, Davis)
for providing feedback and discussion. We also acknowledge Isabel Stumpf (Centre
College) and David Schneider (Centre College) for technical assistance.

References

1. **Thosar SS, and Shea SA**. Circadian control of human cardiovascular function. *Curr Opin Pharmacol* 57: 89-97, 2021.
2. **Mohawk JA, Green CB, and Takahashi JS**. Central and peripheral circadian clocks in mammals. *Annu Rev Neurosci* 35: 445-462, 2012.
3. **Mohawk JA, and Takahashi JS**. Cell autonomy and synchrony of suprachiasmatic nucleus circadian oscillators. *Trends Neurosci* 34: 349-358, 2011.
4. **Buijs RM, and Kalsbeek A**. Hypothalamic integration of central and peripheral clocks. *Nat Rev Neurosci* 2: 521-526, 2001.
5. **Hastings MH, Maywood ES, and Brancaccio M**. Generation of circadian rhythms in the suprachiasmatic nucleus. *Nat Rev Neurosci* 19: 453-469, 2018.

6.   **Tong M, Watanabe E, Yamamoto N, Nagahata-Ishiguro M, Maemura K,**
Takeda N, Nagai R, and Ozaki Y. Circadian expressions of cardiac ion channel genes
in mouse might be associated with the central clock in the SCN but not the peripheral
clock in the heart. *Biol Rhythm Res* 44: 519-530, 2013.
7.   **Scheer FA, Ter Horst GJ, van Der Vliet J, and Buijs RM**. Physiological and
anatomic evidence for regulation of the heart by suprachiasmatic nucleus in rats. *Am J*
Physiol Heart Circ Physiol 280: H1391-1399, 2001.
8.   **Schroder EA, Burgess DE, Zhang X, Lefta M, Smith JL, Patwardhan A,**
Bartos DC, Elayi CS, Esser KA, and Delisle BP. The cardiomyocyte molecular clock
regulates the circadian expression of Kcnh2 and contributes to ventricular
repolarization. *Heart Rhythm* 12: 1306-1314, 2015.
9.   **Gottlieb LA, Larsen K, Halade GV, Young ME, and Thomsen MB**. Prolonged
QT intervals in mice with cardiomyocyte-specific deficiency of the molecular clock. *Acta*
Physiol (Oxf) 233: e13707, 2021.
10.   **Delisle BP, Stumpf JL, Wayland JL, Johnson SR, Ono M, Hall D, Burgess**
DE, and Schroder EA. Circadian clocks regulate cardiac arrhythmia susceptibility,
repolarization, and ion channels. *Curr Opin Pharmacol* 57: 13-20, 2021.
11.   **D'Souza A, Wang Y, Anderson C, Bucchi A, Baruscotti M, Olieslagers S,**
Mesirca P, Johnsen AB, Mastitskaya S, Ni H, Zhang Y, Black N, Cox C, Wegner S,
Bano-Otalora B, Petit C, Gill E, Logantha S, Dobrzynski H, Ashton N, Hart G,
Zhang R, Zhang H, Cartwright EJ, Wisloff U, Mangoni ME, da Costa Martins PA,
Piggins HD, DiFrancesco D, and Boyett MR. A circadian clock in the sinus node
mediates day-night rhythms in Hcn4 and heart rate. *Heart Rhythm* 18: 801-810, 2021.
12.   **Jeyaraj D, Haldar SM, Wan X, McCauley MD, Ripperger JA, Hu K, Lu Y,**
Eapen BL, Sharma N, Ficker E, Cutler MJ, Gulick J, Sanbe A, Robbins J,
Demolombe S, Kondratov RV, Shea SA, Albrecht U, Wehrens XH, Rosenbaum DS,
and Jain MK. Circadian rhythms govern cardiac repolarization and arrhythmogenesis.
Nature 483: 96-99, 2012.
13.   **Schroder EA, Wayland JL, Samuels KM, Shah SF, Burgess DE, Seward T,**
Elayi CS, Esser KA, and Delisle BP. Cardiomyocyte Deletion of Bmal1 Exacerbates
QT- and RR-Interval Prolongation in Scn5a (+/ΔKPQ) Mice. *Front Physiol* 12: 681011,
2021.
14.   **Schroder EA, Lefta M, Zhang X, Bartos DC, Feng HZ, Zhao Y, Patwardhan**
A, Jin JP, Esser KA, and Delisle BP. The cardiomyocyte molecular clock, regulation of
Scn5a, and arrhythmia susceptibility. *Am J Physiol Cell Physiol* 304: C954-965, 2013.
15.   **Schroder EA, Burgess DE, Manning CL, Zhao Y, Moss AJ, Patwardhan A,**
Elayi CS, Esser KA, and Delisle BP. Light phase-restricted feeding slows basal heart
rate to exaggerate the type-3 long QT syndrome phenotype in mice. *Am J Physiol Heart*
Circ Physiol 307: H1777-1785, 2014.
16.   **Zhang D, Colson JC, Jin C, Becker BK, Rhoads MK, Pati P, Neder TH, King**
MA, Valcin JA, Tao B, Kasztan M, Paul JR, Bailey SM, Pollock JS, Gamble KL, and
Pollock DM. Timing of Food Intake Drives the Circadian Rhythm of Blood Pressure.
Function (Oxf) 2: zqaa034, 2021.
17.   **Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, and Schibler**
U. Restricted feeding uncouples circadian oscillators in peripheral tissues from the
central pacemaker in the suprachiasmatic nucleus. *Genes Dev* 14: 2950-2961, 2000.

18. **Schroder EA, Burgess DE, Johnson SR, Ono M, Seward T, Elayi CS, Esser**
KA, and Delisle BP. Timing of food intake in mice unmasks a role for the
cardiomyocyte circadian clock mechanism in limiting QT-interval prolongation.
Chronobiol Int 1-10, 2021.
19. **Clark AJ**. The effect of alterations of temperature upon the functions of the
isolated heart. *J Physiol* 54: 275-286, 1920.
20. **Tanner JM**. The relationships between the frequency of the heart, oral
temperature and rectal temperature in man at rest. *J Physiol* 115: 391-409, 1951.
21. **Kirschen GW, Singer DD, Thode HC, Jr., and Singer AJ**. Relationship
between body temperature and heart rate in adults and children: A local and national
study. *Am J Emerg Med* 38: 929-933, 2020.
22. **Swoap SJ, and Gutilla MJ**. Cardiovascular changes during daily torpor in the
laboratory mouse. *Am J Physiol Regul Integr Comp Physiol* 297: R769-774, 2009.
23. **Wu Y, Wang Q, Granger J, Gaido OR, Aguilar EN, Ludwig A, Moroni A,**
Bianchet MA, and Anderson ME. HCN channels sense temperature and determine
heart rate responses to heat. *bioRxiv* 2023.
24. **Fleming PR, and Muir FH**. Electrocardiographic changes in induced
hypothermia in man. *Br Heart J* 19: 59-66, 1957.
25. **Hicks CE, McCord MC, and Blount SG, Jr.** Electrocardiographic changes
during hypothermia and circulatory occlusion. *Circulation* 13: 21-28, 1956.
26. **Karjalainen J, and Viitasalo M**. Fever and cardiac rhythm. *Arch Intern Med* 146:
1169-1171, 1986.
27. **Mattu A, Brady WJ, and Perron AD**. Electrocardiographic manifestations of
hypothermia. *Am J Emerg Med* 20: 314-326, 2002.
28. **Yaganti VM, Naik CA, and Dhond AJ**. Electrocardiographic manifestations in a
hypothermic patient: review of Osborn waves. *Indian Heart J* 59: 80-82, 2007.
29. **van der Linde HJ, Van Deuren B, Teisman A, Towart R, and Gallacher DJ**.
The effect of changes in core body temperature on the QT interval in beagle dogs: a
previously ignored phenomenon, with a method for correction. *Br J Pharmacol* 154:
1474-1481, 2008.
30. **Schibler U, Gotic I, Saini C, Gos P, Curie T, Emmenegger Y, Sinturel F,**
Gosselin P, Gerber A, Fleury-Olela F, Rando G, Demarque M, and Franken P.
Clock-Talk: Interactions between Central and Peripheral Circadian Oscillators in
Mammals. *Cold Spring Harb Symp Quant Biol* 80: 223-232, 2015.
31. **Guzadhur L, Pearcey SM, Duehmke RM, Jeevaratnam K, Hohmann AF,**
Zhang Y, Grace AA, Lei M, and Huang CL. Atrial arrhythmogenicity in aged
Scn5a+/DeltaKPQ mice modeling long QT type 3 syndrome and its relationship to Na+
channel expression and cardiac conduction. *Pflugers Arch* 460: 593-601, 2010.
32. **Zhang Y, Wu J, King JH, Huang CL, and Fraser JA**. Measurement and
interpretation of electrocardiographic QT intervals in murine hearts. *Am J Physiol Heart*
Circ Physiol 306: H1553-1557, 2014.
33. **Nuyens D, Stengl M, Dugarmaa S, Rossenbacker T, Compernolle V, Rudy Y,**
Smits JF, Flameng W, Clancy CE, Moons L, Vos MA, Dewerchin M, Benndorf K,
Collen D, Carmeliet E, and Carmeliet P. Abrupt rate accelerations or premature beats
cause life-threatening arrhythmias in mice with long-QT3 syndrome. *Nature Medicine* 7:
1021-1027, 2001.

34.   **Skop V, Guo J, Liu N, Xiao C, Hall KD, Gavrilova O, and Reitman ML**. Mouse
Thermoregulation: Introducing the Concept of the Thermoneutral Point. *Cell Rep* 31:
107501, 2020.
35.   **Lund K, Nygaard H, and Kirstein Pedersen A**. Weighing the QT intervals with
the slope or the amplitude of the T wave. *Ann Noninvasive Electrocardiol* 7: 4-9, 2002.
36.   **Hughes ME, Hogenesch JB, and Kornacker K**. JTK_CYCLE: an efficient
nonparametric algorithm for detecting rhythmic components in genome-scale data sets.
J Biol Rhythms 25: 372-380, 2010.
37.   **Bray MS, Ratcliffe WF, Grenett MH, Brewer RA, Gamble KL, and Young ME**.
Quantitative analysis of light-phase restricted feeding reveals metabolic dyssynchrony in
mice. *Int J Obes (Lond)* 37: 843-852, 2013.
38.   **Satoh Y, Kawai H, Kudo N, Kawashima Y, and Mitsumoto A**. Time-restricted
feeding entrains daily rhythms of energy metabolism in mice. *Am J Physiol Regul Integr*
Comp Physiol 290: R1276-1283, 2006.
39.   **Barazi N, Polidovitch N, Debi R, Yakobov S, Lakin R, and Backx PH**.
Dissecting the Roles of the Autonomic Nervous System and Physical Activity on
Circadian Heart Rate Fluctuations in Mice. *Front Physiol* 12: 692247, 2021.
40.   **Gordan R, Gwathmey JK, and Xie LH**. Autonomic and endocrine control of
cardiovascular function. *World J Cardiol* 7: 204-214, 2015.
41.   **Fenske S, Probstle R, Auer F, Hassan S, Marks V, Pauza DH, Biel M, and
Wahl-Schott C**. Comprehensive multilevel in vivo and in vitro analysis of heart rate
fluctuations in mice by ECG telemetry and electrophysiology. *Nat Protoc* 11: 61-86,
2016.
42.   **Goldstein DS, Bentho O, Park MY, and Sharabi Y**. Low-frequency power of
heart rate variability is not a measure of cardiac sympathetic tone but may be a
measure of modulation of cardiac autonomic outflows by baroreflexes. *Exp Physiol* 96:
1255-1261, 2011.
43.   **Press WH, and Numerical Recipes Software (Firm)**. Numerical recipes in C.
Cambridge, England ; New York, N.Y.: Cambridge University Press,, 1992, p. 1
computer disk 5 1/4 in.
44.   **Levine JD, Funes P, Dowse HB, and Hall JC**. Signal analysis of behavioral and
molecular cycles. *BMC Neurosci* 3: 1, 2002.
45.   **Team Rc**. R: A Language and Environment for Statistical Computing R
Foundation for Statistical Computing. https://www.R-project.org.
46.   **Rose RA, Kabir MG, and Backx PH**. Altered heart rate and sinoatrial node
function in mice lacking the cAMP regulator phosphoinositide 3-kinase-gamma. *Circ*
Res 101: 1274-1282, 2007.
47.   **Krishnaswamy PS, Egom EE, Moghtadaei M, Jansen HJ, Azer J, Bogachev
O, Mackasey M, Robbins C, and Rose RA**. Altered parasympathetic nervous system
regulation of the sinoatrial node in Akita diabetic mice. *J Mol Cell Cardiol* 82: 125-135,
2015.
48.   **Egom EE, Vella K, Hua R, Jansen HJ, Moghtadaei M, Polina I, Bogachev O,
Hurnik R, Mackasey M, Rafferty S, Ray G, and Rose RA**. Impaired sinoatrial node
function and increased susceptibility to atrial fibrillation in mice lacking natriuretic
peptide receptor C. *J Physiol* 593: 1127-1146, 2015.

49.   **Mulla W, Murninkas M, Levi O, and Etzion Y**. Incorrectly corrected? QT interval
analysis in rats and mice. *Front Physiol* 13: 1002203, 2022.
50.   **Mulla W, Gillis R, Murninkas M, Klapper-Goldstein H, Gabay H, Mor M,**
Elyagon S, Liel-Cohen N, Bernus O, and Etzion Y. Unanesthetized Rodents
Demonstrate Insensitivity of QT Interval and Ventricular Refractory Period to Pacing
Cycle Length. *Front Physiol* 9: 897, 2018.
51.   **Ripplinger CM, Glukhov AV, Kay MW, Boukens BJ, Chiamvimonvat N,**
Delisle BP, Fabritz L, Hund TJ, Knollmann BC, Li N, Murray KT, Poelzing S, Quinn
TA, Remme CA, Rentschler SL, Rose RA, and Posnack NG. Guidelines for
assessment of cardiac electrophysiology and arrhythmias in small animals. *Am J*
Physiol Heart Circ Physiol 323: H1137-H1166, 2022.
52.   **Swoap SJ, Li C, Wess J, Parsons AD, Williams TD, and Overton JM**. Vagal
tone dominates autonomic control of mouse heart rate at thermoneutrality. *Am J Physiol*
Heart Circ Physiol 294: H1581-1588, 2008.
53.   **Scheer FA, Hilton MF, Mantzoros CS, and Shea SA**. Adverse metabolic and
cardiovascular consequences of circadian misalignment. *Proc Natl Acad Sci U S A* 106:
4453-4458, 2009.
54.   **Sano H, Hayashi H, Makino M, Takezawa H, Hirai M, Saito H, and Ebihara S**.
Effects of suprachiasmatic lesions on circadian rhythms of blood pressure, heart rate
and locomotor activity in the rat. *Jpn Circ J* 59: 565-573, 1995.
55.   **Schroder EA, and Delisle BP**. Time Restricted Feeding to the Light Cycle
Dissociates Canonical Circadian Clocks and Physiological Rhythms in Heart Rate. *Front*
Pharmacol 13: 910195, 2022.
56.   **Warren WS, Champney TH, and Cassone VM**. The suprachiasmatic nucleus
controls the circadian rhythm of heart rate via the sympathetic nervous system. *Physiol*
Behav 55: 1091-1099, 1994.
57.   **Curtis AM, Cheng Y, Kapoor S, Reilly D, Price TS, and Fitzgerald GA**.
Circadian variation of blood pressure and the vascular response to asynchronous
stress. *Proc Natl Acad Sci U S A* 104: 3450-3455, 2007.
58.   **Schroder EA, Ono M, Johnson SR, Rozmus ER, Burgess DE, Esser KA, and**
Delisle BP. The role of the cardiomyocyte circadian clocks in ion channel regulation
and cardiac electrophysiology. *J Physiol* 600: 2037-2048, 2022.
59.   **Yang G, Chen L, Grant GR, Paschos G, Song WL, Musiek ES, Lee V,**
McLoughlin SC, Grosser T, Cotsarelis G, and FitzGerald GA. Timing of expression
of the core clock gene Bmal1 influences its effects on aging and survival. *Sci Transl*
Med 8: 324ra316, 2016.
60.   **Hayter EA, Wehrens SMT, Van Dongen HPA, Stangherlin A, Gaddameedhi**
S, Crooks E, Barron NJ, Venetucci LA, Neill JSO, Brown TM, Skene DJ, Trafford
AW, and Bechtold DA. Distinct circadian mechanisms govern cardiac rhythms and
susceptibility to arrhythmia. *Nat Commun* 12: 2472, 2021.
61.   **Refinetti R**. Circadian rhythmicity of body temperature and metabolism.
Temperature (Austin) 7: 321-362, 2020.
62.   **Postema PG, and Wilde AA**. The measurement of the QT interval. *Curr Cardiol*
Rev 10: 287-294, 2014.

63.   **Molnar J, Zhang F, Weiss J, Ehlert FA, and Rosenthal JE**. Diurnal pattern of
QTc interval: how long is prolonged? Possible relation to circadian triggers of
cardiovascular events. *J Am Coll Cardiol* 27: 76-83, 1996.
64.   **Browne KF, Prystowsky E, Heger JJ, Chilson DA, and Zipes DP**.
Prolongation of the Q-T interval in man during sleep. *Am J Cardiol* 52: 55-59, 1983.
65.   **Gohari A, Wiebe D, and Ayas N**. Shift working and cardiovascular health.
Chronobiol Int 40: 27-32, 2023.
66.   **Sabir IN, Killeen MJ, Grace AA, and Huang CL**. Ventricular arrhythmogenesis:
insights from murine models. *Prog Biophys Mol Biol* 98: 208-218, 2008.
618
619

Figure Legends

Figure 1. Light cycle-restricted feeding caused large changes in hourly RR intervals, QT intervals, PR intervals, and core body temperature. A. Representative ensemble ECG traces recorded from a mouse in ad libitum-fed conditions (ALF, left column), after initially starting light cycle-restricted feeding (L-RF$_i$, middle column), and two weeks after starting light cycle-restricted feeding (L-RF$_{2wk}$, right column) at different Zeitgeber times (ZT). **B.** Averaged time series data showing three consecutive days of the hourly mean for the RR interval (vermillion), QT interval (sky blue), PR interval (orange), and core body temperature (Tb, reddish purple). The data were plotted as a function of ZT and the shaded regions correspond to the dark cycle. The left column shows the data recorded during ALF, the middle column shows the data recorded during L-RF$_i$ (the boxed arrow indicates 24 hours after the start of light cycle-restricted feeding), and the right column shows the data recorded after L-RF$_{2wk}$. The solid lines are cosine curves fitted to the data (n = 6 mice, data points are averaged data and SD).

Figure 2. Light cycle-restricted feeding modified the properties of the 24-hour rhythms for RR intervals, QT intervals, PR intervals, and core body temperature. The individual mouse time series data for the RR intervals, QT intervals, PR intervals, and core body temperature (Tb) were fit to a cosine wave to calculate the acrophase, amplitude, and mesor for mice in ad libitum-fed conditions (ALF), 24 hours after initially starting light cycle-restricted feeding (L-RF$_i$), and two weeks after starting light cycle-restricted feeding (L-RF$_{2wk}$). **See Supplemental Table 1. A.** The top row shows the acrophase of the RR interval (vermillion), QT interval (sky blue), PR interval (orange), and Tb (reddish purple) for each condition. **B.** The top row shows the amplitude, and the bottom row shows the mesor of the RR interval, QT interval, PR interval, and Tb for each condition. Data were analyzed using a two-way mixed effects model (**see Figure 3S**) with Geisser and Greenhouse correction and Sidǎk correction, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, n = 5-6 mice per group.

Figure 3. Light cycle-restricted feeding altered heart rate variability and activity. A. The top row of graphs shows the averaged time series data for three consecutive days of the hourly mean for the RR interval from **Figure 1** (grey) overlayed with the estimated RR$_{core\ body\ temperature}$ interval (green). The left column shows the data from mice in ad libitum-conditions (ALF), the middle column shows the data initially starting light cycle-restricted feeding (L-RF$_i$), and the third column shows the data two weeks after starting light cycle-restricted feeding (L-RF$_{2wk}$). The middle row of graphs shows the corresponding data for the power of the high-frequency component in heart rate variability (pHF, blue), and the bottom row of graphs shows the corresponding data for cage activity (yellow). Data were plotted as a function of ZT and the shaded regions correspond to the dark cycle (n = 6 mice, data points are averaged data and SD). **B.** The left graph shows the average pHF value for each mouse during the light cycle (open circle) and dark cycle (solid circle) measured during ALF, L-RF$_i$, and L-RF$_{2wk}$. The right graph shows the average activity value for each mouse during the light cycle (open circle) and dark cycle (solid circle) during ALF, L-RF$_i$, and L-RF$_{2wk}$. Data were analyzed

665 using a one-way ANOVA with Geisser and Greenhouse correction and Sidǎk correction,
666 *p<0.05 and **p<0.01, n = 6 mice per group.
667
Figure 4. After starting light cycle-restricted feeding, the RR interval continued to
show correlated rhythmicity with core body temperature and pHF but not activity.
The mean data and SD for the cross-correlation between the time series data for the
RR interval and core body temperature (Tb, top row, reddish purple), power of the high-
frequency component in heart rate variability (pHF, middle row, blue), and activity
(bottom row, yellow) in ad libitum-fed conditions (ALF, left column), after initially starting
light cycle-restricted feeding (L-RF$_i$, middle column), and two weeks after starting light
cycle-restricted feeding (L-RF$_{2wk}$, right column). The peak correlation coefficient at lag 0
is shown (n = 6 mice per group). Differences in the peak correlation coefficient data at
lag 0 were analyzed using a one-way ANOVA with Geisser and Greenhouse correction
and Sidǎk correction *p<0.05 and **p<0.01.
679
Figure 5. Inhibition of autonomic receptor activation caused a loss in the daily
rhythm of the RR interval during light cycle-restricted feeding. A. Time series data
showing the hourly mean in the RR interval (vermillion) and core body temperature (Tb,
reddish purple) recorded 1 week after the start of light cycle-restricted feeding (ZT 3 to
ZT 9). The data were plotted as a function of ZT and the shaded regions represent the
dark cycle. The time series shows the hourly mean data and SD measured during the
light and dark cycle the day before starting the propranolol and atropine injections (LC0,
DC0), and the first and second light and dark cycles after starting the propranolol and
atropine injections at ZT 23.5 and ZT 9.5 (LC1, DC1, LC2, and DC2, respectively,
injection times are denoted by arrows; n = 5 mice). **B.** Shown are the RR interval
(vermillion) and Tb (reddish purple) data recorded during LC0 and DC0, LC1 and DC1,
and LC2 and DC2; the horizontal bar in each data set is the mean data. The data were
analyzed using a two-way ANOVA with Geisser and Greenhouse correction and Tukey
correction, *p<0.05, **p<0.01, ***p<0.001. **C.** Time series of RR interval data from **A.**
overlayed with the averaged time series for the corresponding RR$_{core body temperature}$
interval (blue) before and after injection with propranolol and atropine.
696
Figure 6. Correcting the QT interval for core body temperature caused a loss in
the daily rhythm in the QT interval. A. The left graph shows the hourly mean in the
QT interval plotted as a function of the corresponding core body temperature (Tb) for
each mouse in ad libitum-fed conditions, initially after starting light cycle-restricted
feeding, and two weeks after starting light cycle-restricted feeding (each mouse is
represented by a different color, n = 6). The average slope was used to correct the QT
interval to 37°C (QT$_{37°C}$ interval). The right graph shows the QT$_{37°C}$ interval is plotted as
a function of the corresponding Tb for the same set of mice. Linear regression of the
data for each mouse (red lines) was used to calculate the average slope and R^2 and SD
for both the left and right graphs. **B.** The left column shows the hourly QT$_{37°C}$ interval
and SD calculated during ad libitum-fed conditions (ALF), the middle column shows the
QT$_{37°C}$ interval initially starting light cycle-restricted feeding (L-RF$_i$), and the third column
shows the QT$_{37°C}$ interval two weeks after starting light cycle-restricted feeding (L-
RF$_{2wk}$). Also shown is the corresponding time series data for the hourly mean in the

uncorrected QT interval data from **Figure 1** for direct comparison (grey). Data were
plotted as a function of ZT and the shaded regions represent the dark cycle.
713

Figure 7. Housing mice in thermoneutrality unmasked distinct mechanisms
regulating daily rhythms in RR and QT intervals. A. Representative hourly ensemble
ECG traces recorded from a mouse in thermoneutrality at ZT 4 and 17. Also shown are
the time series data for the hourly mean (and SD) for the RR (sky blue), $RR_{core\ body}$
$_{temperature}$ (blue), QT (purple), and $QT_{37°C}$ (vermillion) intervals measured from mice in ad
libitum-fed conditions housed at thermoneutrality. Data were plotted as a function of ZT
and the shaded regions correspond to the dark cycle (n = 6 mice). **B.** The top graph
plots the $QT_{37°C}$ interval as a function of the corresponding core body temperature (Tb)
for the same set of mice in **A**. Each mouse is represented by a different color. The
bottom graph shows the hourly mean in the uncorrected QT interval plotted as a
function of the corresponding Tb. Linear regression (red lines) analysis for each mouse
was used to calculate the mean slope and R^2 (including SD) for both data sets.
726

Figure 8. QT interval correction for core body temperature unmasked QT interval
prolongation in a mouse model of long QT syndrome. A. The left column shows the
hourly $QT_{37°C}$ interval and SD calculated from $Scn5a^{+/\Delta KPQ}$ mice (blue) during ad libitum-
fed conditions (ALF), the middle column shows the $QT_{37°C}$ interval measured initially
after starting light cycle-restricted feeding (L-RF_i), and the third column shows the
$QT_{37°C}$ interval measured two weeks after starting light cycle-restricted feeding (L-
RF_{2wk}). The graph also shows the mean hourly $QT_{37°C}$ interval measured from WT mice
from **Figure 6** for comparison (grey) **C.** The graph shows $QT_{37°C}$ intervals measured
from WT (grey) or $Scn5a^{+/\Delta KPQ}$ mice (blue) in ALF, L-RF_i, and L-RF_{2wk}. The horizontal
bar in each data set is the mean (Data were analyzed using a two-way ANOVA with
Geisser and Greenhouse correction and Sidǎk correction, n = 6, *p<0.05,).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

A.

Slope = -4.3 ± 0.6
R^2 = 0.68 ± 0.07

Slope = 0.03 ± 0.04
R^2 = 0.04 ± 0.03

B.

☐ WT QT interval
■ WT $QT_{37°C}$ interval

Figure 6.

WT housed at 30°C

Figure 7.

WT and *Scn5a*$^{+/\Delta KPQ}$ housed at 22°C

A.

□ WT QT$_{37°C}$ interval
■ *Scn5a*$^{+/\Delta KPQ}$ QT$_{37°C}$ interval

Figure 8.