Feeding Behavior Modifies Daily Rhythms in Autonomic Signaling and Core Body Temperature to Shape 24-hour Rhythms in Cardiac Electrophysiology

Makoto Ono[1], Don E. Burgess[1], Sidney R. Johnson[1], Claude S. Elayi[2], Karyn A. Esser[3], Tanya S. Seward[1], Carie R. Boychuk[4], Rebecca A. Stalcup[1], Abhilash Prabhat[1], Elizabeth A. Schroder[1,5]*, Brian P. Delisle[1]*

[1]Department of Physiology, University of Kentucky, Lexington, KY, USA
[2]CHI Saint Joseph Hospital, Lexington, KY, USA
[3]University of Florida, Gainesville, FL, USA
[4]Dalton Cardiovascular Research Center, Department of Biomedical Sciences, University of Missouri, Columbia, MO, USA
[5]Department of Internal Medicine, University of Kentucky, Lexington, KY, USA

*Contributed equally to the study

Running Title: Feeding Shapes 24-hour Rhythms in Cardiac Ephys

Corresponding authors:

Elizabeth A. Schroder
Department of Internal Medicine
859-323-5571
eschr0@uky.edu
780 Rose Street, Medical Science Building, Rm: MN-512, Lexington, KY 40536-0298

Brian P. Delisle
(859) 323-2797
 brian.delisle@uky.edu
741 S Limestone Street BBSRB B353, Lexington, KY, 40536

Word Count: 7,974

Key Words: Cardiac Electrophysiology**,** Time of Feeding**,** Core Temperature**,** Circadian Rhythms**,** Autonomic Nervous System

Abstract

Background: Circadian rhythms in physiology optimize organismal behavior to anticipate predictable environmental changes. Modern lifestyles can cause misalignment between ≈24-hour rhythms in feeding behavior from daily environmental cycles. Feeding behavior is a potent behavioral zeitgeber (timekeeping cue). We determined how shifting the timing of feeding behavior impacts the alignment of ≈24-hour rhythms in HR and cardiac electrophysiology in wild-type (WT) and $Scn5a^{+/\Delta KPQ}$ mice.

Methods: Mice eat most of their daily calories during the dark cycle. We shifted limited their food access to the light cycle (light-cycle restricted feeding or L-RF). Telemetry was used to continuously record changes in electrocardiogram (ECG), core body temperature (Tb), and activity. Cosinor and cross correlation analyses were used to analyze rhythmicity and 24-hour alignment between variables.

Results: L-RF for one day caused the 24-hour rhythms of the RR, QT, and PR intervals to shift 7-9 hours. The changes in the 24-hour ECG rhythms before and after the start of L-RF mirrored changes in daily body temperature (Tb) rhythm and correlated with changes in the high frequency (pHF) component of heart rate variability. Injecting mice undergoing L-RF with propranolol and atropine reduced the pHF component and caused a loss in the 24-hour rhythm in the RR interval and Tb. We identified a strong linear relation between the the QT interval and Tb. Normalizing the QT interval for Tb eliminated the 24-hour rhythm in the QT interval and identified abnormal changes in the QT interval in WT and $Scn5a^{+/\Delta KPQ}$ mice during L-RF.

Conclusion: Altering the timing of daily feeding behavior reshapes 24-hour rhythms in cardiac electrophysiology in mice by modifying daily rhythms in autonomic signaling and Tb. There is a strong linear relation between the QT interval and Tb. Normalizing the QT for Tb is important for identifying QT interval prolongation independent of Tb.

Abbreviations:

ALF - ad libitum fed
Electrocardiogram - ECG
L-RF - Light cycle restricted feeding
HR - Heart rate – (HR)
HRV – heart rate variability
MAP - Mean arterial pressure
pHF – high-frequency power
pLF – low-frequency power
Tb - core body temperature
SD- standard deviation
WT - Wild type
QT_{37^oC} – QT corrected to 37^{o}C
ZT – Zeitgeber Time

Introduction

Modern lifestyles, including shift work and nighttime eating, disrupt evolutionarily conserved behavioral and environmental timekeeping cues critical for maintaining healthy circadian homeostatic mechanisms.[1] Disruption of these timekeeping cues and subsequent homeostasis is linked to an increased incidence of cardiovascular disease.[2,3] Understanding mechanistically how misaligned timekeeping cues ultimately impact cardiac function will provide new insight into how modern lifestyles can be improved to positively influence cardiac health and decrease arrhythmia susceptibility.

Daily rhythms of cardiac function usually align with daily feeding and activity cycles.[4] In mice, mean heart rate (HR) and mean arterial pressure (MAP) are highest during the dark cycle when mice are active and consume most of their daily calories.[5] Restricting food availability to the light cycle (light cycle-restricted feeding or L-RF) disrupts the alignment between daily feeding behavior and activity cycles. Importantly, L-RF in mice shifts the phase of the daily rhythms in HR and MAP (i.e., increases in HR and MAP) to align to the time of feeding and not activity, resulting in significantly higher HR and MAP during the light cycle compared to the dark cycle.[6,7] These data suggest that feeding behavior largely shapes the 24-hour rhythms in cardiovascular physiology. The purpose of this study is to determine the mechanism by which feeding behavior impacts the 24-hour rhythm of HR and cardiac electrophysiology.

Previous studies show that L-RF in mice shifts the phase and increases the amplitude of the daily core body temperature (Tb) rhythm.[8,9] Several studies show that HR increases and decreases with Tb.[10-14] We tested the idea that the L-RF induced change in the daily Tb rhythm facilitates drives the realignment of the daily rhythms in HR and cardiac electrophysiology to feeding behavior.

Methods

Animals. All the mice used in this study were male SV129 mice. Wild type (WT) and *Scn5a$^{+/\Delta KPQ}$* mice used in longitudinal L-RF studies were 3-4 months old, and mice used for pharmacological studies were 6-8 months old. All procedures complied with the Association for Assessment and Accreditation of Laboratory Animal Care guidelines and were approved by the Institutional Animal Care and Use Committee at the University of Kentucky. Dr. Peter Carmeliet kindly provided the *Scn5a$^{+/\Delta KPQ}$* mouse strain.[15] This study represents new and reanalysis of data collected previously.[6]

Light Cycle Restricted Feeding. Mice were housed in 12-hour light/dark cycles at room temperature with ad libitum access to water. To initiate L-RF, food was removed at 9 h after the beginning of the light cycle (Zeitgeber time or ZT = 9). The next day, food presentation was restricted for 7 h during the light cycle, between ZT 2–9 for up to two weeks unless otherwise noted. L-RF caused mice to lose body mass during the first week (**Figure S1**). However, caloric intake increased after that, and L-RF animals gained body mass after two weeks.[16] Telemetry recordings were performed in ad libitum fed conditions (ALF) before L-RF, shortly after starting L-RF, and two weeks after the start of L-RF.

ECG, Tb, and Locomotor Activity Measurements by Telemetry. We used telemetry to measure electrocardiogram (ECG), Tb, and locomotor activity. Mice were anesthetized

with isoflurane and implanted with PhysioTel ETA-F10 (Data Sciences International)
telemetry transmitter units. ECG telemetry data were recorded at 1000 Hz, and RR and
QT interval analyses were done using MATLAB (MathWorks), similar to previously
described.[6,17,18] ECG traces recorded for each hour were aligned to the peak of the R
wave to generate an averaged ECG trace. The start of the P wave was defined as the
point where the slope of the P wave profile changed by more than 0.001 mV/ms; the Q
wave was defined as the point where the slope of the profile changed from negative to
positive at the base of the QRS complex; the end of the T wave was defined as the
point of intersection of T wave maximum slope with baseline.[19]

The hourly mean data for RR, QT, and PR intervals and Tb were fit with the cosine
curve function:

$$\text{Interval} = A*\cos[2\pi*t/T - \emptyset] + m$$

to calculate the period (T), the time between the peak amplitudes; acrophase ($\emptyset$), the
phase of the peak amplitude in relation to the start of the light cycle (ZT = 0); the
amplitude (A), one-half of the peak to trough amplitude; and the mesor (m), the rhythm
adjusted mean.[6,17,18]

Heart rate variability (HRV) was calculated as outlined previously.[20] RR interval time
series were generated for R wave peak detection. The frequency range used to classify
low frequency was 0.4-1.5 Hz, and the frequency range to classify high frequency was
1.5-4.0 Hz.[20]

We used cross-correlation between time series to compare the rhythmicity and phase
alignment of the RR interval and Tb, the strength of the high frequency (pHF) HRV
component, or activity.[21] The RR interval time series was lagged against the time series
for the Tb, pHF component, or activity. If the time series are in phase, the peak cross-
correlation will occur at lag 0. A regular rise and fall in the cross-correlation between
variables with a lag of 0 indicates correlated rhythmicity between the different time
series. We used the R function "ccf" to calculate the cross-correlation between two data
sets.[22] The formula used in "ccf" to calculate the cross-correlation between two time
series $X(t), Y(t)$ has the form

$$c_{xy}(t) = \frac{1}{n} \sum_{s=\max(1,-t)}^{\min(n-t,n)} [X(s+t) - \bar{X}] [Y(s+t) - \bar{Y}], \, r_{xy}(t) = \frac{c_{xy}(t)}{c_{xy}(0)}$$

This study also corrected the hourly mean in the QT interval for Tb. We used an
approach similar to that described previously to generate the Tb correction formula.[23]
The hourly mean QT intervals were plotted against the corresponding Tb for all data
collected in ALF, after the start of L-RF, and two weeks after the start of L-RF. The data
were described using linear regression, and the R^2 for a linear relationship was
significant for each data set. The average slope of the linear QT-Tb relation was used to
derive the correction formula. We corrected the QT intervals to 37°C ($QT_{37°C}$ interval)
because 37°C was the daily mean in the Tb for mice in ALF.

$$QT_{37°C} = QT \text{ interval} - (4.3*(37°C-Tb))$$

We demonstrated that the $QT_{37^\circ C}$ interval generated a horizontal line in the corrected
$QT_{37^\circ C}$ -Tb plots to validate the formula.

Propranolol and atropine injections. We assessed the effect of L-RF on RR interval
following intraperitoneal (IP) injection of propranolol hydrochloride and atropine sulfate
(propranolol and atropine; 10 mg/kg each) to block β-adrenergic and muscarinic
receptors, respectively, similar to that previously reported.[24-26] We did longitudinal
studies and injected telemetered mice undergoing L-RF at ZT 23.5 (when the daily
rhythm in the RR interval peaked) and 9.5 (when the RR interval was at its nadir) for two
consecutive days. The L-RF protocol for these mice was to provide food access
between ZT 3 and ZT 9. Feeding onset was delayed one hour to separate the food
anticipatory decrease in RR interval before feeding from the start of the light cycle.

Statistical analysis. For some graphs, data are shown as the mean and standard
deviation of the mean (SD). For other graphs, a data point for each animal is shown with
a horizontal bar at the mean. The statistical JTK_CYCLE package was used to confirm
statistically significant cyclical rhythms before analyzing the data using cosine waveform
curve fitting.[27] Data were analyzed in PRISM (GraphPad). Data between two groups
were analyzed using a T-test, and data sets with more than two groups were analyzed
using a 1-way or 2-way ANOVA. Post-hoc testing was done using Šidák. $p<0.05$ was
considered significant.

Results

L-RF simultaneously modifies 24-hour rhythms in HR, cardiac electrophysiology, and
Tb.

This study aimed to identify the mechanism(s) for the impact of L-RF in mice on daily
rhythms in HR (as measured by the RR interval) and cardiac electrophysiology. We
used *in vivo* telemetry to longitudinally record ECG data from mice in ALF, after the
initial start of L-RF, and two weeks after the start of L-RF (**Figure 1A**). Data recorded
shortly after starting L-RF revealed the acute impact L-RF had on cardiac
electrophysiology, and data recorded two weeks after starting L-RF indicated steady-
state (i.e., chronic) changes L-RF had on cardiac electrophysiology.[9] The top panel in
Figure 1B shows a three-day time series for changes in the hourly mean in the RR
interval, the duration of ventricular repolarization (the QT interval), and the duration of
atrial to ventricular conduction (the PR interval). All three of the time series for the
hourly mean of the RR, QT, and PR intervals displayed daily rhythms that aligned with
each other and peaked at a similar time during the light cycle (unshaded areas).
However, one day after starting L-RF, the daily rhythms in ECG data exhibited large
simultaneous shifts in phase, resulting in peaks (i.e., longer wave durations and slower
HRs) during the dark cycle. These 'dark cycle' peaks were also notably larger in
amplitude than before L-RF started. These changes were not transitory and persisted
throughout L-RF (i.e., two weeks after the start of L-RF).

L-RF causes similar causes similar shifts in daily Tb rhythms.[9,28] We tested whether
L-RF-induced changes in Tb rhythm (i.e., robust drops in Tb during the dark cycle)
occurred simultaneously with the changes in the ECG. We compared the hourly mean
in Tb for the same days as ECG data from mice in ALF, after starting L-RF, and two

weeks after starting L-RF (**Figure 1B**). The daily Tb rhythm mirrored daily rhythms in RR, QT, and PR intervals before and after the start of L-RF.

The individual ECG and Tb time series recorded from mice in ALF, one day after starting L-RF, and two weeks after starting L-RF were significant cyclical rhythms well described using cosine curve fitting. We used cosine curve fitting to the individual mouse data to calculate the phase shifts caused by L-RF (**Figures 1B and 2A**). Data showed that the acrophase of the daily rhythms in the RR intervals, QT intervals, PR intervals, and Tb shifted 7-9 hours shortly after starting L-RF. Only small additional changes were seen after two weeks in L-RF. Although L-RF caused abrupt changes in the phase of the ECG and Tb data, the periodicity of the daily ECG and Tb rhythms remained ≈24 hours (**Figure 2A**).

Cosine curve fitting allowed us to calculate the impact that L-RF had on the mesor and amplitude of the 24-hour rhythms in RR intervals, QT intervals, PR intervals, and Tb (**Figure 2B**). Compared to data recorded in ALF, the mesor and amplitudes of 24-hour rhythms in RR and QT intervals increased one day after starting L-RF and persisted as long as recorded (two weeks after starting L-RF). Analogous changes were seen in the mesor and amplitude of the Tb rhythm. In contrast, the L-RF-induced changes in the 24-hour rhythm of the PR interval did not occur immediately. Alterations in mesor and amplitude of the PR interval were only detected two weeks after starting L-RF.

L-RF-induced changes in the RR interval correlate with changes in HRV but not activity

The 24-hour rhythm in the RR interval is classically thought to reflect changes in efferent signaling from the autonomic nervous system (ANS), endocrine system, and daily activity cycles.[29-31] We quantified changes in ANS cardiac regulation by measuring HRV in the frequency domain.[20] We focused on the pHF component as an index for parasympathetic motor efferent regulation of cardiac function since L-RF caused an overall slowing of HR. For mice in ALF, the amplitude of the pHF component was higher during the light cycle compared to the dark cycle. One day after starting L-RF, the pHF component increased during the dark cycle (**Figure 1B**), and the increase persisted two weeks after starting L-RF.

We also measured the time series in activity from the mice in ALF, after starting L-RF, and two weeks after starting L-RF (**Figure 1B**). In ALF mice, mouse activity is highest during the dark cycle. However, the activity peaks during the dark cycle were less apparent after starting L-RF because activity profiles showed multiple peaks during the light and dark cycles.

The individual mouse three-day time series data for the pHF component and activity rhythms did not show robust significant cyclical rhythms, and they were not well described by cosine waveform curve fitting. Therefore, we analyzed the relative rhythmicity between the RR interval and pHF or activity using cross-correlation analyses.[21] As part of the analysis, we included the cross-correlation between RR-Tb. **Figure 3** shows the cross-correlation results for the RR-Tb, RR-pHF, and RR-activity from mice in ALF, one day after starting L-RF, and two weeks after starting L-RF. The RR-Tb and RR-pHF showed correlated rhythmicity that peaked with a lag of 0. A qualitatively similar correlated rhythmicity was seen in the cross-correlation for RR-

activity during ALF, but the correlated rhythmicity was lost after starting L-RF. The data
suggest Tb and autonomic signaling may be important for shaping the daily rhythm in
the RR interval recorded from mice in L-RF.

Propranolol and atropine cause a loss in daily RR interval and Tb rhythms during L-RF

We determined how propranolol and atropine injections impacted the 24-hour
rhythm in mean RR interval and Tb from mice in L-RF (fed from ZT 3 to 9). Mice were
subjected to L-RF for one week and then injected with propranolol and atropine at ZT
23.5 (**Figure 4A**). We confirmed that propranolol and atropine injection reduced the
amplitude of the pHF and pLF components (**Figure 4B**). We found that the injection
only caused minor changes in the mean RR interval and Tb when compared to the
same time on the previous day (**Figure 2S, Figure 4A,** and **Figure 4B**). Repeating the
propranolol and atropine injection at ZT 9.5 also caused only small changes in the mean
RR interval and Tb compared to the same time on the previous day (**Figure 2S and
4A**). However, several hours after the injection at ZT 9.5, the RR interval did not
increase and Tb did not decrease during the dark cycle (**Figure 4C**). Repeating the
injections of propranolol and atropine at ZT 23.5 and 9.5 for a second day also
prevented the increase in the RR interval and the decrease in Tb during the subsequent
dark cycle. The data demonstrate that injecting mice in L-RF with propranolol and
atropine resulted in a loss in the daily rhythm of the RR interval and Tb in L-RF mice
(**Figure 4C**). We conclude that, during L-RF, the daily rhythm in the RR interval is
strongly linked to changes in the autonomic signaling and the daily Tb rhythm.

L-RF reveals a robust linear QT-Tb relation in WT mice

The time series data show that longitudinal changes in the mean QT interval are
similar to the RR interval and mirror the Tb rhythm (**Figure 1**). Previous studies
demonstrate that QT interval duration in mice is independent of RR interval.[32-35]
Therefore, we quantified the linear relation between the hourly QT interval and Tb using
data recorded from mice in ALF, initially after starting L-RF, and two weeks after starting
L-RF similar to that described for beagles (**Figure 5A**).[23] We found a strong negative
linear relationship, and we used the average slope of linear regression analyses to
normalize the QT interval data to a mean Tb of ALF (37°C) (**Figure 5A**). Normalizing the
time series of the QT interval for a Tb of 37°C caused a loss in the 24-hour rhythm in
the QT interval for mice in ALF and L-RF (**Figure 5B**).

*Correcting the QT interval unmasks differences in the QT interval between WT and
Scn5a$^{+/\Delta KPQ}$ mice*

To determine the physiological significance for the QT interval dependence on Tb
we studied a transgenic mouse model of long QT type 3 syndrome (*Scn5a$^{+/\Delta KPQ}$*) before
and after L-RF similar to that described for WT mice in **Figure 1**. We measured ECG
properties and Tb in ALF, immediately after starting L-RF, and two weeks after starting
L-RF. The changes in the daily rhythms of the ECG and Tb data after starting L-RF
were qualitatively and quantitatively similar to those recorded in WT mice (**Figure 6A,
6B, Figure 3S**). Cosine fitting analysis of ECG data from the *Scn5a$^{+/\Delta KPQ}$* mice showed
that one day after starting L-RF, there was an immediate shift in the phase for RR, QT,
and PR intervals and Tb, an increase in the amplitudes of the rhythms, an increase in

the mesor of the RR, QT, and PR intervals, and a decrease in the mesor for Tb (**Figure
3S**). There were some genotype-specific differences in the mesor and amplitude for RR
intervals, QT intervals, and Tb between the WT and $Scn5a^{+/\Delta KPQ}$ mice. We calculated
the QT_{37^oC} interval for $Scn5a^{+/\Delta KPQ}$ mice in ALF, after starting L-RF, and two weeks after
starting L-RF, and we compared the the QT_{37^oC} interval data form the $Scn5a^{+/\Delta KPQ}$ mice
with WT mice (**Figure 6B**). The QT_{37^oC} interval was not different between WT and
$Scn5a^{+/\Delta KPQ}$ mice in ALF. However, after starting L-RF, the QT_{37^oC} interval measured in
$Scn5a^{+/\Delta KPQ}$ mice became longer than the QT_{37^oC} interval measured from WT mice
(**Figure 3S**). We also identified a genotype-specific difference in how L-RF impacted the
QT_{37^oC} interval in WT and $Scn5a^{+/\Delta KPQ}$ mice. In WT mice, L-RF increased the QT_{37^oC}
interval only after two weeks, but in $Scn5a^{+/\Delta KPQ}$ mice, the QT_{37^oC} interval increased only
after one day (**Figure 3S**).

Discussion

In this study, we demonstrate that one day after starting L-RF, 24-hour rhythms in
mean RR, QT, and PR intervals and Tb shifted simultaneously by 7-9 hours to a new
phase and remained at this new phase for the duration of L-RF. The 24-hour rhythms in
L-RF were shaped by increased ECG interval durations and lower Tb measured during
the dark cycles. The increase of the RR interval correlated with the changes in the
amplitude of the pHF component in HRV analysis and did not depend on the activity.
Repeated inhibition of β-adrenergic and muscarinic receptors in mice undergoing L-RF
prevented the increase in RR interval and decrease in Tb during the dark cycle-
resulting in the loss of the daily RR interval and Tb rhythms in mice during L-RF. We
also identified a strong linear relation between the hourly mean in the QT interval and
Tb. Correcting the QT interval for Tb allowed us to identify abnormal changes in the QT
interval measured from WT and $Scn5a^{+/\Delta KPQ}$ mice independent of Tb.

Studies suggested that the circadian clock in the suprachiasmatic nucleus (SCN)
modified ANS signaling to generate the 24-hour rhythm in HR.[4,30,36-38] These studies
were supported by data showing that ablation of SCN or transgenic mice lacking the
core circadian clock transcription factor *Bmal1* (e.g., $Bmal1^{-/-}$) did not have a 24-hour
rhythm in HR.[39-41] L-RF is not thought to impact the circadian clock mechanism in the
SCN;[9] therefore, it was surprising that our previous work demonstrated that L-RF
caused a large phase shift in the 24-hour rhythm of the HR.[6] L-RF in mice also shifts the
phase of circadian gene expression in peripheral tissues, including the heart, to align
with the timing of feeding behavior.[9] Several studies implicate the cardiomyocyte
circadian clock in regulating gene expression essential for cardiac excitability, cardiac
electrophysiology, and HR.[18,35,40,42-45] One might predict then that L-RF shifts the phase
of cardiomyocyte gene expression to facilitate the phase shift in HR and cardiac
electrophysiology. However, genetically disrupting the cardiomyocyte-specific circadian
clock (by inducing the deletion of *Bmal1*) does not abolish the L-RF mediated phase
shift in the daily HR and cardiac electrophysiological rhythms.[17] Additionally, we tested
for a difference in the HR measured from mice in ALF vs. L-RF after injecting
propranolol and atropine (**Figure 5S**). We did not detect a difference in HR between
these two groups after injection. Together, the data demonstrate that the observed
changes in HR with L-RF are not secondary to changes in cardiomyocyte circadian

clock signaling or intrinsic to the heart. The changes in HR with L-RF appear to be
secondary to daily changes in autonomic signaling and Tb.

The QT interval is used clinically to assess people's risk for deadly ventricular
arrhythmias.[46] In humans, the QT interval changes as a function of the RR interval, and
several formulas can correct the QT interval for HR. Although HR correction formulas
for the QT interval in mice exist,[47] the most commonly used correction formula was
derived by measuring the average HR and corresponding QT interval across the 24-
hour cycle. The temperature dependence of the QT interval and daily rhythm in Tb were
not considered. Importantly, recent studies show mice have a weak HR adaptation of
the QT interval[32,33,48] and applying the QT interval correction leads to overcorrection and
errors. We suspected that the daily rhythm in the QT interval used to derive the QT
interval correction in mice may have reflected daily rhythms in Tb and not HR.
Consistent with this idea, we now report a robust linear QT-Tb relation in mice, and
correcting the QT interval for Tb caused a loss in the daily rhythm of the QT interval in
mice during ALF and L-RF. These new data raise the possibility that the daily rhythm in
the QT interval in mice primarily reflects Tb. This possibility is also supported by a study
showing the 24-hours rhythm of the QT interval in mice is independent of the daily RR
rhythm.[35] Tb was not measured in this study, but the observed change in the amplitude
of the daily rhythm in the QT interval (4-5 ms) is consistent with an expected amplitude
change of the daily Tb rhythm in mice (1°C).[49]

We extended our L-RF studies to quantify the changes in the QT interval in a
transgenic knock-in mouse model that expresses a mutation linked to long QT
syndrome in people.[15] Correcting the QT interval for Tb allowed direct comparisons of
QT intervals at different Tb across the 24-hour cycle, identified genotype-specific
differences in the QT interval independent of Tb, and showed that L-RF caused a
prolongation of the QT interval in $Scn5a^{+/\Delta KPQ}$ mice (but not WT mice) after only one
day. These results are qualitatively different if the QT interval is not corrected for Tb and
underscore the importance of monitoring and considering the impact of Tb when
measuring ECGs in mice.

Implications

Modern lifestyles disrupt evolutionarily conserved behavioral and environmental
timekeeping cues critical for maintaining healthy circadian homeostatic mechanisms.[1]
We found that a misalignment of the timing of feeding behavior relative to the light/dark
cycle causes dramatic changes in the daily rhythms of HR and cardiac
electrophysiology in mice. These changes appeared to be driven by changes in ANS
signaling and Tb. Our findings suggest feeding behavior is a critical timekeeping cue for
the daily Tb and cardiac electrophysiology rhythms in mice. If conserved, modern
lifestyles that disrupt evolutionary conserved feeding rhythms in people may negatively
impact the alignment between daily circadian and cardiac electrophysiological to impair
cardiac performance and influence arrhythmogenic risk.

Another implication of this study is that it underscores the importance that daily
rhythms in Tb have in shaping 24-hour rhythms in cardiac electrophysiology.
Continuous monitoring of Tb is not routinely performed in people undergoing
electrophysiological testing despite representing an important contributor to the HR and

QTc intervals. This is especially true in studies that monitor changes in HR and QTc
interval across the 24-hour cycle. Several studies estimate that the Q10 in the HR of
mice is between 2-2.5.[13,14] The Q10 in HR reflect a conserved structural and functional
property of HCN channels that directly couples temperature to cell membrane
excitability.[14] A HR with a Q10 of 2-2.5 predicts that 50-75% of the observed amplitude
in the daily rhythm of the RR interval is secondary to the daily rhythm Tb. Our data
suggest Tb is also important for shaping the 24-hour rhythm in the QT interval in mice.
Based on previously published data, we propose that the daily rhythm in Tb also
contributes to the 24-hour rhythm in the QTc interval observed in people. Several
studies report a temperature dependence of the QTc interval in people as 10-15 ms for
every 1°C change.[23,50-54] The peak to trough amplitude in the daily Tb rhythm in people
is about 1°C, and the peak to trough amplitude in the daily rhythm in the QTc is 10-20
ms.[55,56] Future studies that measure both ECG parameters and Tb across the 24-hour
cycle in people will help to delineate how much of the 24-hour rhythms in cardiac
electrophysiology are due to daily rhythms in Tb. Importantly, measuring the Tb and the
QTc interval together may help to delineate abnormal changes in the QT interval
secondary to repolarization abnormalities vs. abnormal changes in Tb.

Limitations

These studies were performed in mice at room temperature, and mice housed at
room temperature are vulnerable to torpor during fasting. Torpor is a transient
hibernating-like state that causes a large drop in Tb (below 32°C) and slow HR.[13] Our
model of L-RF did not calorically restrict the mice. The mice continued to eat 3-4 grams
of food per day during L-RF (**Figure 1S**), gained mass after one week in L-RF (**Figure**
1S), and Tb analysis showed only one mouse briefly had a Tb < 33°C after starting L-
RF (**Figure 4S**). Another limitation this study used only male mice. We chose male mice
because female mice are more prone to bouts of torpor.[13] It also remains unclear
whether a longer $QT_{37°C}$ interval is associated with an increased risk for spontaneous
ventricular arrhythmias. The lack of spontaneous arrhythmias in $Scn5a^{+/\Delta KPQ}$ mice is not
surprising because the absolute incidence of spontaneous arrhythmias in these mice
and people with LQT3 is extremely low. We found that, during L-RF, the daily rhythm in
the PR interval simultaneously shifted with the RR and QT intervals and Tb. However,
due to the small amplitude and relatively slow change in the PR interval in response to
L-RF, it was difficult to determine if Tb contributed to changes in the PR interval. Lastly,
caution must be used when extrapolating ECG data obtained from mice to people.[34,57]

Summary

Daily rhythms in HR and cardiac electrophysiology align with feeding and activity
cycles via signals originating from the SCN. However, when food availability becomes
restricted to the light cycle, this disrupts the association between environmental and
behavioral timekeeping cues, and daily rhythms in HR and cardiac electrophysiology
align with the new feeding behavior and Tb. We conclude that L-RF-induced changes in
the daily autonomic signaling and Tb rhythms are important for facilitating the
realignment 24-hour rhythms in HR and cardiac electrophysiology, and the daily
variation in the Tb rhythm helps to shape the daily rhythms in the RR and QT intervals.

Competing interests
None declared.
347
348 **Author contributions**
349 All authors contributed to drafting the work and revising it critically for important
350 intellectual content, approved the final version of the manuscript, and qualified for
351 authorship.
352

353 **Funding**
354 This work was supported by National Heart Lung and Blood Institute grants
355 R01HL153042 and R01HL141343.
356
357 **Acknowledgement**
358 We would like to thank Drs. Wayne Giles and Chao-Yin Chen (University of California,
359 Davis) for providing feedback and discussion. We would also like to acknowledge Isabel
360 Stumpf (Centre College) and David Schneider (Centre College) for technical assistance.
361

References

1. Gohari A, Wiebe D, Ayas N. Shift working and cardiovascular health. *Chronobiol Int*. 2023;40:27-32. doi: 10.1080/07420528.2021.1933000

2. Yang L, Luo Y, He L, Yin J, Li T, Liu S, Li D, Cheng X, Bai Y. Shift Work and the Risk of Cardiometabolic Multimorbidity Among Patients With Hypertension: A Prospective Cohort Study of UK Biobank. *J Am Heart Assoc*. 2022;11:e025936. doi: 10.1161/JAHA.122.025936

3. Xu C, Weng Z, Liang J, Liu Q, Zhang X, Xu J, Li Q, Zhou Y, Gu A. Shift Work, Genetic Factors, and the Risk of Heart Failure: A Prospective Study of the UK Biobank. *Mayo Clin Proc*. 2022;97:1134-1144. doi: 10.1016/j.mayocp.2021.12.003

4. Schroder EA, Delisle BP. Time Restricted Feeding to the Light Cycle Dissociates Canonical Circadian Clocks and Physiological Rhythms in Heart Rate. *Front Pharmacol*. 2022;13:910195. doi: 10.3389/fphar.2022.910195

5. Li P, Sur SH, Mistlberger RE, Morris M. Circadian blood pressure and heart rate rhythms in mice. *Am J Physiol*. 1999;276:R500-504. doi: 10.1152/ajpregu.1999.276.2.R500

6. Schroder EA, Burgess DE, Manning CL, Zhao Y, Moss AJ, Patwardhan A, Elayi CS, Esser KA, Delisle BP. Light phase-restricted feeding slows basal heart rate to exaggerate the type-3 long QT syndrome phenotype in mice. *Am J Physiol Heart Circ Physiol*. 2014;307:H1777-1785. doi: 10.1152/ajpheart.00341.2014

7. Zhang D, Colson JC, Jin C, Becker BK, Rhoads MK, Pati P, Neder TH, King MA, Valcin JA, Tao B, et al. Timing of Food Intake Drives the Circadian Rhythm of Blood Pressure. *Function (Oxf)*. 2021;2:zqaa034. doi: 10.1093/function/zqaa034

8. Schibler U, Gotic I, Saini C, Gos P, Curie T, Emmenegger Y, Sinturel F, Gosselin P, Gerber A, Fleury-Olela F, et al. Clock-Talk: Interactions between Central and Peripheral Circadian Oscillators in Mammals. *Cold Spring Harb Symp Quant Biol*. 2015;80:223-232. doi: 10.1101/sqb.2015.80.027490

9. Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U. Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev*. 2000;14:2950-2961. doi: 10.1101/gad.183500

10. Clark AJ. The effect of alterations of temperature upon the functions of the isolated heart. *J Physiol*. 1920;54:275-286. doi: 10.1113/jphysiol.1920.sp001929

11. Tanner JM. The relationships between the frequency of the heart, oral temperature and rectal temperature in man at rest. *J Physiol*. 1951;115:391-409. doi: 10.1113/jphysiol.1951.sp004677

12. Kirschen GW, Singer DD, Thode HC, Jr., Singer AJ. Relationship between body temperature and heart rate in adults and children: A local and national study. *Am J Emerg Med*. 2020;38:929-933. doi: 10.1016/j.ajem.2019.158355

13. Swoap SJ, Gutilla MJ. Cardiovascular changes during daily torpor in the laboratory mouse. *Am J Physiol Regul Integr Comp Physiol*. 2009;297:R769-774. doi: 10.1152/ajpregu.00131.2009

14. Wu Y, Wang Q, Granger J, Gaido OR, Aguilar EN, Ludwig A, Moroni A, Bianchet MA, Anderson ME. HCN channels sense temperature and determine heart rate responses to heat. *bioRxiv*. 2023. doi: 10.1101/2023.09.02.556046

15. Nuyens D, Stengl M, Dugarmaa S, Rossenbacker T, Compernolle V, Rudy Y, Smits JF, Flameng W, Clancy CE, Moons L, et al. Abrupt rate accelerations or premature beats cause life-threatening arrhythmias in mice with long-QT3 syndrome. *Nature Medicine*. 2001;7:1021-1027. doi: 10.1038/nm0901-1021

16. Bray MS, Ratcliffe WF, Grenett MH, Brewer RA, Gamble KL, Young ME. Quantitative analysis of light-phase restricted feeding reveals metabolic dyssynchrony in mice. *Int J Obes (Lond)*. 2013;37:843-852. doi: 10.1038/ijo.2012.137

17. Schroder EA, Burgess DE, Johnson SR, Ono M, Seward T, Elayi CS, Esser KA, Delisle BP. Timing of food intake in mice unmasks a role for the cardiomyocyte circadian clock mechanism in limiting QT-interval prolongation. *Chronobiol Int*. 2021:1-10. doi: 10.1080/07420528.2021.2011307

18. Schroder EA, Burgess DE, Zhang X, Lefta M, Smith JL, Patwardhan A, Bartos DC, Elayi CS, Esser KA, Delisle BP. The cardiomyocyte molecular clock regulates the circadian expression of Kcnh2 and contributes to ventricular repolarization. *Heart Rhythm*. 2015;12:1306-1314. doi: 10.1016/j.hrthm.2015.02.019

19. Lund K, Nygaard H, Kirstein Pedersen A. Weighing the QT intervals with the slope or the amplitude of the T wave. *Ann Noninvasive Electrocardiol*. 2002;7:4-9. doi: 10.1111/j.1542-474x.2001.tb00132.x

20. Fenske S, Probstle R, Auer F, Hassan S, Marks V, Pauza DH, Biel M, Wahl-Schott C. Comprehensive multilevel in vivo and in vitro analysis of heart rate fluctuations in mice by ECG telemetry and electrophysiology. *Nat Protoc*. 2016;11:61-86. doi: 10.1038/nprot.2015.139

21. Levine JD, Funes P, Dowse HB, Hall JC. Signal analysis of behavioral and molecular cycles. *BMC Neurosci*. 2002;3:1. doi: 10.1186/1471-2202-3-1

22. Team Rc. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. https://www.R-project.org. 2020.

23. van der Linde HJ, Van Deuren B, Teisman A, Towart R, Gallacher DJ. The effect of changes in core body temperature on the QT interval in beagle dogs: a previously ignored phenomenon, with a method for correction. *Br J Pharmacol*. 2008;154:1474-1481. doi: 10.1038/bjp.2008.265

24. Rose RA, Kabir MG, Backx PH. Altered heart rate and sinoatrial node function in mice lacking the cAMP regulator phosphoinositide 3-kinase-gamma. *Circ Res*. 2007;101:1274-1282. doi: 10.1161/CIRCRESAHA.107.158428

25. Krishnaswamy PS, Egom EE, Moghtadaei M, Jansen HJ, Azer J, Bogachev O, Mackasey M, Robbins C, Rose RA. Altered parasympathetic nervous system regulation of the sinoatrial node in Akita diabetic mice. *J Mol Cell Cardiol*. 2015;82:125-135. doi: 10.1016/j.yjmcc.2015.02.024

26. Egom EE, Vella K, Hua R, Jansen HJ, Moghtadaei M, Polina I, Bogachev O, Hurnik R, Mackasey M, Rafferty S, et al. Impaired sinoatrial node function and increased susceptibility to atrial fibrillation in mice lacking natriuretic peptide receptor C. *J Physiol*. 2015;593:1127-1146. doi: 10.1113/jphysiol.2014.283135

27.   Hughes ME, Hogenesch JB, Kornacker K. JTK_CYCLE: an efficient
nonparametric algorithm for detecting rhythmic components in genome-scale
data sets. *J Biol Rhythms*. 2010;25:372-380. doi: 10.1177/0748730410379711

28.   Satoh Y, Kawai H, Kudo N, Kawashima Y, Mitsumoto A. Time-restricted feeding
entrains daily rhythms of energy metabolism in mice. *Am J Physiol Regul Integr*
Comp Physiol. 2006;290:R1276-1283. doi: 10.1152/ajpregu.00775.2005

29.   Barazi N, Polidovitch N, Debi R, Yakobov S, Lakin R, Backx PH. Dissecting the
Roles of the Autonomic Nervous System and Physical Activity on Circadian Heart
Rate Fluctuations in Mice. *Front Physiol*. 2021;12:692247. doi:
10.3389/fphys.2021.692247

30.   Tong M, Watanabe E, Yamamoto N, Nagahata-Ishiguro M, Maemura K, Takeda
N, Nagai R, Ozaki Y. Circadian expressions of cardiac ion channel genes in
mouse might be associated with the central clock in the SCN but not the
peripheral clock in the heart. *Biol Rhythm Res*. 2013;44:519-530. doi:
10.1080/09291016.2012.704801

31.   Gordan R, Gwathmey JK, Xie LH. Autonomic and endocrine control of
cardiovascular function. *World J Cardiol*. 2015;7:204-214. doi:
10.4330/wjc.v7.i4.204

32.   Mulla W, Murninkas M, Levi O, Etzion Y. Incorrectly corrected? QT interval
analysis in rats and mice. *Front Physiol*. 2022;13:1002203. doi:
10.3389/fphys.2022.1002203

33.   Mulla W, Gillis R, Murninkas M, Klapper-Goldstein H, Gabay H, Mor M, Elyagon
S, Liel-Cohen N, Bernus O, Etzion Y. Unanesthetized Rodents Demonstrate
Insensitivity of QT Interval and Ventricular Refractory Period to Pacing Cycle
Length. *Front Physiol*. 2018;9:897. doi: 10.3389/fphys.2018.00897

34.   Ripplinger CM, Glukhov AV, Kay MW, Boukens BJ, Chiamvimonvat N, Delisle
BP, Fabritz L, Hund TJ, Knollmann BC, Li N, et al. Guidelines for assessment of
cardiac electrophysiology and arrhythmias in small animals. *Am J Physiol Heart*
Circ Physiol. 2022;323:H1137-H1166. doi: 10.1152/ajpheart.00439.2022

35.   Gottlieb LA, Larsen K, Halade GV, Young ME, Thomsen MB. Prolonged QT
intervals in mice with cardiomyocyte-specific deficiency of the molecular clock.
Acta Physiol (Oxf). 2021;233:e13707. doi: 10.1111/apha.13707

36.   Scheer FA, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and
cardiovascular consequences of circadian misalignment. *Proc Natl Acad Sci U S*
A. 2009;106:4453-4458. doi: 10.1073/pnas.0808180106

37.   Sano H, Hayashi H, Makino M, Takezawa H, Hirai M, Saito H, Ebihara S. Effects
of suprachiasmatic lesions on circadian rhythms of blood pressure, heart rate
and locomotor activity in the rat. *Jpn Circ J*. 1995;59:565-573. doi:
10.1253/jcj.59.565

38.   Warren WS, Champney TH, Cassone VM. The suprachiasmatic nucleus controls
the circadian rhythm of heart rate via the sympathetic nervous system. *Physiol*
Behav. 1994;55:1091-1099. doi: 10.1016/0031-9384(94)90392-1

39.   Curtis AM, Cheng Y, Kapoor S, Reilly D, Price TS, Fitzgerald GA. Circadian
variation of blood pressure and the vascular response to asynchronous stress.
Proc Natl Acad Sci U S A. 2007;104:3450-3455. doi: 10.1073/pnas.0611680104

40. Schroder EA, Ono M, Johnson SR, Rozmus ER, Burgess DE, Esser KA, Delisle BP. The role of the cardiomyocyte circadian clocks in ion channel regulation and cardiac electrophysiology. *J Physiol*. 2022;600:2037-2048. doi: 10.1113/JP282402

41. Yang G, Chen L, Grant GR, Paschos G, Song WL, Musiek ES, Lee V, McLoughlin SC, Grosser T, Cotsarelis G, et al. Timing of expression of the core clock gene Bmal1 influences its effects on aging and survival. *Sci Transl Med*. 2016;8:324ra316. doi: 10.1126/scitranslmed.aad3305

42. Delisle BP, Stumpf JL, Wayland JL, Johnson SR, Ono M, Hall D, Burgess DE, Schroder EA. Circadian clocks regulate cardiac arrhythmia susceptibility, repolarization, and ion channels. *Curr Opin Pharmacol*. 2021;57:13-20. doi: 10.1016/j.coph.2020.09.015

43. Jeyaraj D, Haldar SM, Wan X, McCauley MD, Ripperger JA, Hu K, Lu Y, Eapen BL, Sharma N, Ficker E, et al. Circadian rhythms govern cardiac repolarization and arrhythmogenesis. *Nature*. 2012;483:96-99. doi: 10.1038/nature10852

44. D'Souza A, Wang Y, Anderson C, Bucchi A, Baruscotti M, Olieslagers S, Mesirca P, Johnsen AB, Mastitskaya S, Ni H, et al. A circadian clock in the sinus node mediates day-night rhythms in Hcn4 and heart rate. *Heart Rhythm*. 2021;18:801-810. doi: 10.1016/j.hrthm.2020.11.026

45. Hayter EA, Wehrens SMT, Van Dongen HPA, Stangherlin A, Gaddameedhi S, Crooks E, Barron NJ, Venetucci LA, Neill JSO, Brown TM, et al. Distinct circadian mechanisms govern cardiac rhythms and susceptibility to arrhythmia. *Nat Commun*. 2021;12:2472. doi: 10.1038/s41467-021-22788-8

46. Postema PG, Wilde AA. The measurement of the QT interval. *Curr Cardiol Rev*. 2014;10:287-294. doi: 10.2174/1573403x10666140514103612

47. Mitchell GF, Jeron A, Koren G. Measurement of heart rate and Q-T interval in the conscious mouse. *Am J Physiol*. 1998;274:H747-751. doi: 10.1152/ajpheart.1998.274.3.H747

48. Roussel J, Champeroux P, Roy J, Richard S, Fauconnier J, Le Guennec JY, Thireau J. The Complex QT/RR Relationship in Mice. *Sci Rep*. 2016;6:25388. doi: 10.1038/srep25388

49. Refinetti R. Circadian rhythmicity of body temperature and metabolism. *Temperature (Austin)*. 2020;7:321-362. doi: 10.1080/23328940.2020.1743605

50. Fleming PR, Muir FH. Electrocardiographic changes in induced hypothermia in man. *Br Heart J*. 1957;19:59-66. doi: 10.1136/hrt.19.1.59

51. Hicks CE, McCord MC, Blount SG, Jr. Electrocardiographic changes during hypothermia and circulatory occlusion. *Circulation*. 1956;13:21-28. doi: 10.1161/01.cir.13.1.21

52. Karjalainen J, Viitasalo M. Fever and cardiac rhythm. *Arch Intern Med*. 1986;146:1169-1171.

53. Mattu A, Brady WJ, Perron AD. Electrocardiographic manifestations of hypothermia. *Am J Emerg Med*. 2002;20:314-326. doi: 10.1053/ajem.2002.32633

54. Yaganti VM, Naik CA, Dhond AJ. Electrocardiographic manifestations in a hypothermic patient: review of Osborn waves. *Indian Heart J*. 2007;59:80-82.

55. Molnar J, Zhang F, Weiss J, Ehlert FA, Rosenthal JE. Diurnal pattern of QTc interval: how long is prolonged? Possible relation to circadian triggers of

cardiovascular events. *J Am Coll Cardiol*. 1996;27:76-83. doi: 10.1016/0735-
1097(95)00426-2
56.   Browne KF, Prystowsky E, Heger JJ, Chilson DA, Zipes DP. Prolongation of the
Q-T interval in man during sleep. *Am J Cardiol*. 1983;52:55-59. doi:
10.1016/0002-9149(83)90068-1
57.   Sabir IN, Killeen MJ, Grace AA, Huang CL. Ventricular arrhythmogenesis:
insights from murine models. *Prog Biophys Mol Biol*. 2008;98:208-218. doi:
10.1016/j.pbiomolbio.2008.10.011
550
551

Figure Legends

Figure 1. L-RF causes a simultaneous shift in the daily rhythms of the RR, QT, and PR intervals, Tb, and strength of the pHF component. A. Representative hourly averaged ECG traces recorded from a mouse in ALF (left), after initially starting L-RF (middle), or 2 weeks after starting L-RF (right). **B.** Averaged time series (n = 6 mice, data points are averaged data and SD) data showing 3 consecutive days of the hourly mean for the RR interval (vermillion), QT interval (sky blue), PR interval (orange), Tb (reddish purple), pHF (blue), and activity (yellow). The data are plotted as a function of ZT and the shaded regions correspond to the dark cycle. Left column shows the data recorded during ALF, middle column shows the data recorded initially after starting L-RF (boxed arrow indicates 24 hours after the start of L-RF), and right column shows the data recorded 2 weeks after starting L-RF. The solid lines shown for the time series ECG and Tb data in ALF, 24 hours after the start of L-RF, and 2 weeks after the start of L-RF are cosine curve fit to the averaged data.

Figure 2. L-RF causes a simultaneous shift and increased amplitude in the 24-hour rhythms of the mean RR, QT, and PR intervals and Tb. The individual mouse time series data for the RR, QT, and PR intervals, and Tb were fit to a cosine wave to calculate the acrophase, period, mesor, and amplitude for mice in ALF, 24 hours after initially starting L-RF (L-RF$_i$), and 2 weeks after starting L-RF (L-RF$_{2wk}$). **A.** The top row shows the acrophase of the RR interval (vermillion), QT interval (sky blue), PR interval (orange), and Tb (reddish purple) for each condition. The second row, left graph, shows the difference in the acrophase of the RR interval, QT interval, PR interval, and Tb measured from mice in ALF compared to those from mice 24 hours after initially starting L-RF or 2 weeks after starting L-RF. The second row, right graph, shows the period of the RR interval, QT interval, PR interval, and Tb from mice in ALF, 24 hours after initially starting L-RF, and 2 weeks after starting L-RF. **B.** The top row shows the mesor and second row shows the amplitude of the RR interval, QT interval, PR interval, and Tb for each condition (n = 6 mice per group, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. The horizontal bar in each data set is the mean).

Figure 3. Cross-correlation analysis between the RR interval and Tb or strength of the pHF component before and after L-RF shows correlated and aligned rhythmicity. The average of the cross-correlation between the time series data for the RR interval and Tb (top row, reddish purple), pHF component (middle row, blue), and activity (bottom row, yellow) in ALF (left column), 24 hours after initially starting L-RF (middle column), and 2 weeks after starting L-RF (right column). The peak correlation coefficient at lag 0 is shown (n = 6 mice per group, data points are mean data and SD, *p<0.05 and **p<0.01).

Figure 4. Propranolol and atropine prevent the daily rhythms in RR interval and Tb in mice during L-RF. A. Time series data showing the hourly mean in the RR interval (vermillion) and Tb (reddish purple) recorded 1 week after the start of L-RF (ZT 3 to ZT 9). The data are plotted as a function of ZT and the shaded regions represent the dark cycle. The time series shows the average of the hourly mean data and SD

measured during the light and dark cycle the day before starting the propranolol and
atropine injections (LC0, DC0), and the first and second light and dark cycles after
injecting mice at ZT 23.5 and ZT 9.5 (LC1, DC1, LC2, and DC2, respectively, injection
times are denoted by arrows; n = 5 mice). **B.** Shown are the 24-hour means in the pHF
(sky blue) and pLF (blue) components measured from the mice during the four before
and after the first propranolol and atropine injections (horizontal bar in each data set is
the mean, *p<0.05). **C.** Shown are the mean RR interval (vermillion) and Tb (reddish
purple) data recorded during the light and dark cycle on day before (LC0 and DC0) or
the days after starting the injections with propranolol and atropine (LC1, DC1, LC2 and
DC2; horizontal bar in each data set is the mean, *p<0.05; ***p<0.001).

Figure 5. Correcting the QT-interval for Tb causes a loss in the daily rhythm in the
QT interval. A. Left graph shows the hourly mean in the QT interval plotted as a
function of the corresponding Tb for each mouse in ALF, initially after starting L-RF, or 2
weeks after starting L-RF (each mouse is represented by a different color, n = 6). The
right graph shows the $QT_{37^{\circ}C}$ interval is plotted as a function of the corresponding Tb for
the same set of mice. Linear regression of the data for each mouse (red lines) were
used to calculate the average slope and R^2 and SD for both the left and right graphs. **B.**
Averaged time series for the hourly mean $QT_{37^{\circ}C}$ interval and SD (sky blue) in ALF,
initially after starting L-RF, and 2 weeks after the start of L-RF. Also shown is the
corresponding averaged time series data for the hourly mean in the uncorrected QT
interval data from **Figure 1** for direct comparison (grey). Data are plotted as a function
of ZT and the shaded regions represent the dark cycle.

Figure 6. L-RF impacts the daily rhythms of the RR, QT, and PR intervals and Tb
in *Scn5a*$^{+/\Delta KPQ}$ mice. A. Representative of hourly averaged ECG traces recorded from
WT or *Scn5a*$^{+/\Delta KPQ}$ mice in ALF (left), after initially starting L-RF (middle), or 2 weeks
after starting L-RF (right) at different ZT. The top row shows the times of day that the
QT intervals were longest, and the bottom row corresponds to times of day when the
QT intervals were shortest. **B.** Averaged time series (n = 6 mice, data points are mean
data and SD) data showing 3 consecutive days of the hourly mean for the RR interval
(vermillion), QT interval (sky blue), PR interval (orange), Tb (reddish purple), and the
$QT_{37^{\circ}C}$ interval (blue). The data are plotted as a function of ZT and the shaded regions
correspond to the dark cycle. The time series for the $QT_{37^{\circ}C}$ interval also includes the
$QT_{37^{\circ}C}$ interval measured from WT mice (grey) for direct comparison. Left column
shows the data recorded during ALF, middle column shows the data recorded initially
after starting L-RF (boxed arrow indicates 24 hours after the start of L-RF), and right
column shows the data recorded 2 weeks after starting L-RF. The solid lines shown for
the time series ECG and Tb data in ALF, 24 hours after the start of L-RF, and 2 weeks
after the start of L-RF are cosine curve fit to the averaged data.

Supplemental Material
642
643   **Supplemental Figure Legends**
644
Figure 1S. L-RF initially causes a decrease in mass, but the mice regain and
increase mass after 1 week in L-RF. The left graphs shows the daily body mass of
mice of mice in ALF (yellow) or after initially starting L-RF (blueish green, n = 5 mice per
group, **p<0.01 ***p<0.001); the middle graph shows the average weekly mass in mice
during the first 2 weeks of L-RF (n = 6, *p<0.05 and **p<0.01); and the right graph
shows the average food consumption per mouse/day during each day after the start of
L-RF for 2 weeks (n=6).
652
Figure 2S. Injecting L-RF mice with propranolol and atropine has little impact on
mean RR or Tb interval 2 hours after the injection. The graphs show the 15 min
mean in the **A.** RR interval or **B.** Tb. The left panel of each graph shows the data on the
day before the injection (blue), and the right panel of each graph shows the data
immediately before and after injection (vermillion). The injection time is denoted by an
arrow labeled P+A. The left graphs in **A.** and **B.** show the data corresponding to the
transition of the dark (shaded) to light cycle on the day before the injection (LC0) or the
light cycle after the injection (LC1). The injection time was ZT 23.5. The right graphs in
A. and **B.** show the data corresponding to light cycle on the day before the injection
(LC0) or the light cycle after the injection (LC1). The injection time was ZT 9.5. (n = 5,
*p<0.05).
664
Figure 3S. L-RF causes genotype specific differences in the 24-hour rhythms of
the mean RR, QT, and PR intervals and Tb measured from WT and $Scn5a^{+/\Delta KPQ}$
mice. A. The individual mouse time series data for the RR, QT, and PR intervals, and
Tb were fit to a cosine wave to calculate the mesor and amplitude for $Scn5a^{+/\Delta KPQ}$ mice
in ALF, 24 hours after initially starting L-RF (L-RF$_i$), and 2 weeks after starting L-RF (L-
RF$_{2wk}$). The top row shows the mesor and second row shows the amplitude of the RR
interval (vermillion), QT interval (sky blue), PR interval (orange), and Tb (reddish purple)
measured from $Scn5a^{+/\Delta KPQ}$ mice for each condition. The horizontal bar in each data set
is the mean. Also included are the corresponding WT data presented in Figure 2 (grey)
for comparison (n = 6 mice per group, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001). **B.**
The graph shows the average of the mean QT_{37^oC} interval measured from WT (grey) or
$Scn5a^{+/\Delta KPQ}$ mice (blue) in ALF, one day after starting L-RF, and two weeks after
starting L-RF. The horizontal bar in each data set is the mean (n = 6, *p<0.05).
678
679   **Figure 4S. L-RF does not cause torpor.** Shown are the Tb profiles for each individual
680   WT mice in ALF, 24 hours after initially starting L-RF (L-RF$_i$), and 2 weeks after starting
681   L-RF (L-RF$_{2wk}$). Only one mouse showed a transient decrease in the Tb < 33^oC
682   (asterisk).
683
Figure 5S. L-RF does not alter HR after injection with propranolol and atropine.
We injected mice in ALF (yellow) or after two weeks of L-RF (bluish green) and
recorded the mean RR interval using ECGenie (Mouse Specifics, Inc., Boston, MA,

USA). The data were collected between ZT 0.5-3. The ECGenie system comprises a
platform with electrodes connected to an amplifier (e-MOUSE) that measures ECG non-
invasively. The mice were injected with propranolol and atropine (P+A) 15 minutes prior
to recording the ECGs (n = 4 mice per group, the horizontal bar in each data set is the
mean).

Figure 1.

Figure 2.

Figure 3.

A.

P+A injection

Food availability

LC0 LC1 LC2

RR interval (ms)

Tb (°C)

DC0 DC1 DC2

ZT (hrs)

B.

pHF (ms²)

Con   P+A

pLF (ms²)

Con   P+A

C.

RR interval (ms)

Con   P+A   P+A

LC0 DC0 LC1 DC1 LC2 DC2

Tb (°C)

Con   P+A   P+A

LC0 DC0 LC1 DC1 LC2 DC2

Figure 4.

Figure 5.

Figure 6.

Supplement Figure 1.

A.

B.

Supplement Figure 2.

Supplement Figure 3.

Supplement Figure 4.

Supplement Figure 5.