ATF6 activation alters colonic lipid metabolism causing tumor-associated microbial adaptation

Authors

Olivia I. Coleman[1] *, Adam Sorbie[1,2] *, Alessandra Riva[1], Miriam von Stern[1], Stephanie Kuhls[1], Denise M. Selegato[3], Nikolai Köhler[4], Jakob Wirbel[5], Tim Kacprowski[6,7,8], Andreas Dunkel[9], Josh K. Pauling[4], Johannes Plagge[10], Diego Mediel-Cuadra[11], Sophia Wagner[11], Ines Chadly[1], Sandra Bierwith[1], Tingying Peng[11], Thomas Metzler[12], Clemens Schafmayer[13], Sebastian Hinz[13], Christian Röder[14], Christoph Röcken[15], Michael Zimmermann[3], Philip Rosenstiel[16], Katja Steiger[12], Moritz Jesinghaus[17], Gerhard Liebisch[18], Josef Ecker[10,19], Christina Schmidt[20], Georg Zeller[3], Klaus-Peter Janssen[21], Dirk Haller[1,22] #

Authors' affiliation

[1] Chair of Nutrition and Immunology, School of Life Sciences, Technische Universität München, Freising-Weihenstephan, Germany.

[2] Institute for Stroke and Dementia Research (ISD), University Hospital, LMU Munich, Munich, Germany.

[3] Molecular Systems Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany

[4] LipiTUM, Chair of Experimental Bioinformatics, School of Life Sciences Weihenstephan, Technische Universität München, Freising-Weihenstephan, Germany.

[5] Structural and Computational Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany.

[6] Division Data Science in Biomedicine, Peter L. Reichertz Institute for Medical Informatics of TU Braunschweig and Hannover Medical School, Braunschweig, Germany.

[7] Braunschweig Integrated Centre of Systems Biology (BRICS), TU Braunschweig, Braunschweig Germany.

[8] Department of Computational Biology of Infection Research, Helmholtz Centre for Infection Research (HZI), Braunschweig, Germany.

[9] Chair of Food Chemistry and Molecular Sensory Science, School of Life Sciences, Technische Universität München, Germany.

[10] ZIEL Institute of Food & Health, Research Group Lipid Metabolism, Technische Universität München, Freising, Germany.

[11] Helmholtz AI, Helmholtz Munich, 85764 Neuherberg, Germany.

[12] Comparative Experimental Pathology (CEP), School of Medicine and Health, Technische Universität München, Munich, Germany.

[13] Department of General Surgery, University Hospital Rostock, Rostock, Germany.

[14] Institute for Experimental Cancer Research, University Medical Center Schleswig-Holstein, Kiel, Germany.

[15] Department of Pathology, Christian-Albrechts University, Kiel, Germany.

[16] Institute of Clinical Molecular Biology, University hospital Schleswig-Holstein, Campus Kiel
and Kiel University, Kiel, Germany.

[17] Institute of Pathology, Universitätsklinikum Marburg, Baldingerstraße, 35043, Marburg,
Germany.

[18] Institute of Clinical Chemistry and Laboratory Medicine, University Hospital Regensburg,
Regensburg, Germany.

[19] Institute of Clinical Chemistry and Laboratory Medicine, University Hospital Regensburg,
Regensburg, Germany

[20] Faculty of Medicine and Heidelberg University Hospital, Institute for Computational
Biomedicine, Heidelberg University, Heidelberg, Germany.

[21] Department of Surgery, Klinikum rechts der Isar, Technische Universität München,
München, Germany.

[22] ZIEL Institute for Food and Health, Technische Universität München, Freising, Germany

*These authors contributed equally to this project.

Corresponding author:

Prof. Dr. Dirk Haller
Technical University of Munich
Chair of Nutrition and Immunology
Gregor-Mendel-Str. 2
85354 Freising, Germany
Email: dirk.haller@tum.de
Phone: + 49 8161-71 2026

Abstract

Endoplasmic reticulum unfolded protein responses (UPR^{ER}) contribute to cancer development and the activating transcription factor 6 (ATF6) is involved in microbiota-dependent tumorigenesis. Here, we substantiate the clinical relevance of ATF6 in early-onset and late colorectal cancer patient cohorts. Transcriptional analysis in intestinal epithelial cells (IEC) of ATF6 transgenic mice (nATF6^{IEC}) identified bacteria-specific changes in cellular metabolism enriched for fatty acid biosynthesis. Untargeted metabolomics and isotype-labeling confirmed ATF6-related enrichment of long chain fatty acids in colonic tissue of patients, mice and organoid cultures. FASN inhibition and microbiota transfer in germ-free nATF6^{IEC} mice confirmed the causal involvement of ATF6-induced lipid alterations in tumorigenesis. The selective expansion of tumor-relevant microbial taxa was mechanistically linked to long chain fatty acid exposure, using bioorthogonal non-canonical amino acid tagging (BONCAT) and growth analysis of Desulfovibrio isolates. We postulate chronic ATF6 signaling in the epithelium to select for tumor-promoting microbiota by altering lipid metabolism.

Keywords: Endoplasmic reticulum (ER), Activating transcription factor 6 (ATF6), intestinal epithelial cells (IEC), colorectal cancer, metabolomics, lipid metabolism, microbiota

Graphical Abstract: Chronic ATF6 signaling in the colonic epithelium alters lipid metabolism to select a tumor-promoting microbiota

- Biallelic expression of activated ATF6 (p50 nuclear fragment) in intestinal epithelial cells (nATF6[IEC]) induces spontaneous colon tumors in SPF but not GF mice

- Mechanistically, biallelic SPF nATF6[IEC] mice alter colonic lipid metabolism, including the upregulation of LCFAs and Fasn

- Inhibition of FASN prevents colon tumor formation in mice, and reduces the tumor-promoting potential of the intestinal microbiota (FMT)

- Exposure of fl/fl control mouse microbiota to LCFAs *ex vivo* translationally activates tumor-associated bacteria, including *Desulfovibrio fairfieldensis*

- Human CRC patients show ATF6 upregulation, FASN co-occurrence and increased LCFAs in T tissue

- ATF6 activity links with CRC-associated microbiota in patients, including *Desulfovibrio*

Created with BioRender.com

nATF6: activated activating transcription factor 6; LCFA: long-chain fatty acids; SAFA: saturated fatty acids; Fasn: fatty acid synthase; C75 i.p.: intraperitoneal injection of the Fasn inhibitor C75.

Introduction

The endoplasmic reticulum (ER) in mammalian cells forms an extensive tubular-reticular network that acts as a gatekeeper controlling the synthesis and folding of proteins, and the synthesis of cellular lipids [1,2]. The accumulation of un- and misfolded ER proteins activates the unfolded protein response (UPR[ER]), a highly conserved group of intricately regulated signaling pathways that aim to restore ER homeostasis. PKR-like ER kinase (PERK), inositol-requiring enzyme 1 (IRE1) and activating transcription factor 6 (ATF6) constitute the three main arms of the UPR[ER] [3]. ER-stress results in the dissociation of membrane-bound ATF6 complexes from the ER chaperone glucose-regulating protein 78 (GRP78), subsequent cleavage by the site-1 and site-2 proteases at the Golgi apparatus, and translocation of the p50 active cytosolic N-terminal portion of the transcription factor (nATF6) into the nucleus to bind the ER-stress response element (ERSE) [4,5].

Numerous studies recognize the UPR[ER] as a fundamental mediator in cellular physiology and the pathogenesis of inflammatory disorders, metabolic diseases and cancer, including colorectal cancer (CRC) [6-12]. Enhanced expression of GRP78, a downstream target of nATF6, correlates with growth, invasion and metastasis of tumors [13]. Furthermore, ATF6 mRNA expression and polymorphisms are associated with CRC metastasis and relapse [14,15], and hepatocellular carcinoma [16], respectively. We could show that chronic transgenic expression of nATF6 specifically in intestinal epithelial cells (IECs) induces spontaneous and microbiota-dependent colon adenoma formation in a murine model of early-onset CRC (nATF6[IEC]) [11]. Accumulating evidence unarguably renders the intestinal microbiota as an important regulator of host health and a causal player in the development and progression of CRC. Our findings that dysbiosis precedes tumorigenesis in nATF6[IEC] mice and that nATF6-expressing germ-free mice remain tumor-free [11], clearly indicate a mechanistic link between ATF6 signaling and the intestinal microbiota in the context of colon tumorigenesis. What remains to be elucidated, is how ATF6 signaling induces dysbiosis at the pre-tumor stage.

The multifaceted possibilities of physiological outcomes in response to the UPR[ER] underlines the necessity to understand the mechanistic contribution of UPR[ER] signaling to disease pathology. Here we firstly set out to identify the relevance of ATF6 in CRC patient populations, and secondly to understand the consequence of ATF6 signaling on the murine host. We characterize ATF6 expression in multiple independent German CRC cohorts and causally link chronic ATF6 activity to altered lipid metabolism and tumor-relevant microbiota changes.

Results

High ATF6 expression defines a subset of CRC patients

We previously identified increased *ATF6* expression to be associated with reduced disease-free survival in colorectal cancer (CRC) patients [11]. To substantiate these findings, we quantified ATF6 protein expression via immunohistochemistry (IHC) staining in a German CRC patient cohort (cohort 1; *n=959*) using QuPath [17,18]. ATF6 antibody specificity was confirmed through negative staining in human heart muscle, and the appropriate isotype control staining in CRC tissue (Extended Data Fig. 1a). ATF6 expression was compared between the tumor-center (ATF6-TC) and the tumor invasive front (ATF6-IF), showing similar expression levels between these two localizations (data not shown), and subsequently averaged to determine the ATF6 H-score (capturing both the intensity and the percentage of stained cells) (Fig. 1a). Findings show that ATF6 upregulation occurs in 38% of CRC patients that can be classed as ATF6-high in terms of their nuclear (NUC) and cytoplasmic (CYT) expression (Fig. 1a; Extended Data Fig. 1b,c). Stratification into all possible combinations of NUC and CYT ATF6 staining into low (L) and high (H) shows that all NUC H patients are also CYT H (38%), with 0% of NUC H patients being CYT L (Extended Data Fig. 1d). The other combinations of NUC L/CYTL and NUC L/CYT H make up 21% and 41% of the CRC patients, respectively (Extended Data Fig. 1d). In the same cohort 1, we assessed the global levels of endoplasmic reticulum (ER) stress through IHC staining of glucose regulating protein 78 (GRP78), the master chaperone of the ER. Interestingly, quantification of GRP78 using QuPath showed that, despite the wide range of GRP78 H-scores amongst the CRC cohort, there is no difference in GRP78 H-scores between ATF6-high and ATF6-low CRC patient groups (Extended Data Fig. 1e,f). To validate the above findings, we trained a deep learning model to predict ATF6 expression levels of CRC patient biopsy images. The model was trained on an independent CRC patient cohort (cohort 2; *n=50*) with classification and regression objectives using 5-fold cross validation. In line with our results, external testing results identified 39% (classification task) and 35.9% (regression task) of CRC patients in cohort 1 as ATF6-high (Fig. 1b-e, Extended Data Fig. 1g,h), confirming the robustness and applicability of our AI methodology. To further support our findings, we quantified ATF6 expression using QuPath in a third German CRC patient cohort (cohort 3; *n=256*). Here we identified a subpopulation of 19% of CRC patients to be ATF6-high (Fig. 1f). To investigate a potentially dissimilar effect of ATF6 expression in patients with early-onset CRC (age<50; EOCRC) versus late-onset CRC (age>50; LOCRC), we quantified ATF6 expression in additional patients from cohort 3 with EOCRC (cohort 3; *n=55* EOCRC). Findings show similar expression levels in EOCRC, with 11% of patients classified as ATF6-high (Fig. 1g). Together, these results clearly demonstrate that high ATF6 expression defines a subpopulation of CRC patients (19-38% LOCRC; 11% EOCRC).

ATF6 alters colonic fatty acid metabolism in the presence of bacteria

In our ATF6 transgenic mouse model (nATF6^{IEC}), we previously established a causal role of the microbiota in adenoma formation [11]. In agreement with these findings, biallelic nATF6^{IEC} mice (tg/tg) show a reduced survival (Fig. 2a), and develop colonic tumors with an incidence of 100% under specific pathogen-free (SPF) conditions, while controls (fl/fl) and monoallelic (tg/wt) nATF6 expression, as well as germ-free (GF) tg/tg mice remain tumor-free (Fig. 2b). To characterize the transcriptional response induced by chronic ATF6 activity and to further understand the contribution of bacteria, we performed bulk mRNA sequencing of colonic epithelial cells in fl/fl and tg/tg mice at the pre-tumor time point (5wk, highlighted in orange) under SPF and GF conditions (Fig. 2c). Transgenic chronic activation of nATF6 in SPF tg/tg mice results in a total of 1941 differentially expressed genes (DEGs; threshold: -0.5 > log2FC > 0.5, p adj. < 0.05) compared to fl/fl controls, of which 1236 DEGs are upregulated and 705 DEGs are downregulated (Fig. 2d; Supplementary Table 1, sheet 1). Functional analysis using the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways showed that almost half (16/36; blue labels) of the top and bottom ranked pathways, according to the normalized enrichment score (NES), are related to metabolic pathways in SPF tg/tg mice (Fig. 2e, Supplementary Table 2, sheet 1). Under GF conditions, tg/tg mice showed a total of 706 DEGs, of which 491 DEGS are upregulated and 215 DEGs are downregulated (Extended Data Fig. 2a; Supplementary Table 1, sheet 2). Again, half of the top and bottom ranked KEGG pathways (14/28; blue labels) in GF tg/tg mice are linked to metabolic pathways (Extended Data Fig. 2b, Supplementary Table 2, sheet 2). Metabolic KEGG pathways associated with SPF tg/tg mice (tumor development) but not with GF tg/tg mice (no tumor development) are beta alanine metabolism, butanoate metabolism, fatty-acid (FA) metabolism, glycolysis gluconeogenesis, metabolism of xenobiotics by cytochrome P450 and nicotinate and nicotinamide metabolism (Fig. 2e). Of these, FA metabolism represents the highest ranked KEGG pathway upregulated in SPF tg/tg mice. To characterize the specificity of ATF6-related activation in the context of all UPR arms, we specifically looked at UPR-related genes, including the chronic SPF model, GF model, as well as our acute (tamoxifen-inducible transgenic model at 4 days post ATF6-activation; nATF6 Vil-Cre^{ERT2Tg}) SPF model (Extended Data Fig. 2c). We see that eight out of the nine top regulated genes are also described to be ATF6 targets (marked in red), with considerable overlap in the other UPR^{ER} arms. The panel reveals that the expression level of target genes is amplified by chronicity of ATF6-activation and bacterial presence.

To disentangle the transcriptional regulation between chronic tg/tg signaling under SPF and GF conditions, differentiating between the tumor-developing and non-tumor developing phenotype, we applied Signature Regulatory Clustering (SiRCle) [19]. We defined three states of genes in tg/tg mice compared to fl/fl mice for each colonization status group, namely No Change (not regulated, -0.5 < log2FC < 0.5, p adj. > 0.05), UP (upregulated, log2FC > 0.5 and

p. adj. <0.05) and DOWN (downregulated, log2FC <-0.5 and p adj. <0.05) (Fig. 2f,
Supplementary Table 3, sheet 1). Following the flow of genes between SPF and GF conditions
identified eight regulatory clusters named after the biological phenotype they include, for
instance Core-ATF6 UP (Fig. 2f; Supplementary Table 3, sheet 2). Depicted next to the Alluvial
plot are the number and percentage of genes found in each of the clusters. Additionally, we
performed functional analysis using KEGG pathways (threshold: >5% of pathway genes
detected, p adj. < 0.2, supplementary Table 4), and identified transcription factors that drive
individual clusters (supplementary Table 5). After the regulatory cluster "none" (84.70%), the
highest percentages of DEGs were found in the clusters "SPF-UP" (5.93 %), "SPF-DOWN"
(4.27%) and "Core-ATF6 UP" (2.65%) (Fig. 2f). As anticipated, more than half (*P4hb*, *Atf6*,
Calr, *Hspa5*, *Pdia4*, *Anxa6*, *Manf*, *Sdf2l1*, *Hsp90b1*, *Creld2*, *Nucb2*, *C1qtnf1*, *Herpud1* and
Dnajb11) of the highly DEGs in the Core-ATF6 UP cluster (24 DEGs; black labels) are
classically associated with ER protein folding and calcium homeostasis, and localized within
the ER (Extended Data Fig. 2d). Transcription factors *Atf4* and *Atf6* were identified as the
drivers of the CORE-ATF6 UP cluster (Extended Data Fig. 2d, Supplementary Table 5, sheet
1). Highly DEGs in the Core-ATF6 DOWN cluster (black labels) include *Cyba*, *Ces1g* and
Cyp4b1, which are associated with intestinal ROS production and defense against bacteria at
the colonic epithelium [20], hydrolysis of lipids and xenobiotics [21], and the hydroxylation of fatty
acids and alcohols [22], respectively (Extended Data Fig. 2e). Functionally, the CORE-ATF6
DOWN cluster associates with five KEGG pathways, including the drug metabolism
cytochrome p450 pathway, associated with cellular metabolism, homeostasis and the
detoxification of drugs [23] (Extended Data Fig. 2e, Supplementary Table 4, sheet 1). *Arntl* is the
identified transcription factor driving the CORE-ATF6 DOWN cluster, responsible for
maintenance of the circadian rhythm and has been shown to be downregulated in cancers with
an association to anti-cancer roles [24] (Extended Figure 2e, Supplementary Table 5, sheet 2).
Highly regulated DEGs in the SPF DOWN cluster (black labels) include genes that have been
associated with tumor suppression and CRC [25-27] (*Meg3*, *Cwh43, Naprt)*, lipid metabolism and
detoxification [28] (*Ces1f*), and bactericidal/antimicrobial activity [29] (*Iapp*) (Extended Data Fig.
2f). The majority of KEGG pathways associated with the SPF DOWN cluster are metabolic
pathways, including the drug metabolism cytochrome p450 pathway also identified in the
CORE-ATF6 DOWN cluster (Extended Data Fig. 2f, Supplementary Table 4, sheet 2).
Most relevant to our murine model and the role of ATF6 signaling in colon adenoma
development in tg/tg mice is the SPF UP cluster. The highest DEG in this cluster (black label)
is ER localized *Serp2*, classically involved in the UPR and protein glycosylation (Extended
Data Fig. 2g). Among the other highly DEGs (black labels) in this cluster, *Nup88*, *Dmxl1* and
Igf2bp3 are associated with dysplasia and cancer, including CRC [30-32]. Importantly, and in line
with our findings above (Fig. 2e), the SPF UP cluster contains 23 differentially expressed FA

metabolism-related genes (blue labels) (Fig. 2g, Extended Data Fig. 2g). Of note, additional

analysis of the pre-tumor time point using our inducible nATF6 Vil-Cre[ERT2Tg] murine model that

addresses the transcriptional response induced by acute (4d) activation of nATF6,

demonstrates that the upregulation of the majority (13/23) of these FA metabolism-related

genes (blue labels) occurs already early on after nATF6 activation (Extended Data Fig. 2h,

Supplementary Table 1, sheet 3). In accordance with our tumor phenotype, one of the two

identified KEGG pathways associated with the SPF UP cluster is cell cycle (Extended Data

Fig. 2g, Supplementary Table 4, sheet 3). Three transcription factors were identified as drivers

of the SPF UP cluster (Extended Data Fig. 2g, Supplementary Table 5, sheet 3), namely *E2f4*

(control of cell cycle, increased expression in cancers[33]), *Foxm1* (oncogenic transcription

factor, control of cell cycle, regulation of CRC metastasis and progression [34]) and *NFyb*

(promotes invasion, metastasis and differentiation, associated with multiple cancers, activator

or repressor in a context dependent manner [35]).

ATF6 positively correlates with FASN in CRC patients

Alterations in the lipidomic profile have been observed in several cancer types [36-38], including

CRC [39], which marks lipid metabolic rewiring as a phenotypic hallmark of cancer. Upregulation

of fatty acid synthase (FASN), a key multi-enzyme complex of *de novo* lipid biosynthesis [40], is

associated with aggressive disease and poor prognosis in CRC [41,42]. We recently described a

robust CRC-specific lipid signature based on quantitative lipidomics in a multi-cohort and

cross-center study, identifying triacylglycerol species and FASN gene expression as

differential markers of cancerous tissue and bad prognosis, respectively [43]. *Fasn* was shown

to be upregulated in response to both chronic and acute activation of nATF6 in our analyses

(Fig 2.g, Extended Data Fig. 2g,h). Here, we stained a random subset (n=181) of tissue cores

from our human CRC patient cohort 1 for FASN and correlated the QuPath quantified FASN

H-score with our ATF6 classification (from Fig. 1a) in the same patients. Exemplary images

show that patients clearly differed in their FASN expression levels (Fig. 2h). Strikingly, FASN

H-scores were significantly higher in CRC patients classified as ATF6-high compared to those

classified as ATF6-low (Fig. 2i), identifying a positive correlation between ATF6 and FASN

expression in CRC patient tissue. In a further approach, the publicly available Pan-Cancer

Atlas (*n=10.967* cases including approx. 600 CRC cases) and TCGA (*n=640* CRC cases)

databases were screened for genomic amplifications/gains (amp/gain) in either *ATF6* alone

(*ATF6*-only), *FASN* alone (*FASN*-only) or in both genes simultaneously (*ATF6+FASN*). The

relationship between *ATF6*-high and *FASN*-high patients is evident through their association

classified as co-occurrence, despite not reaching significance in the TCGA database

(*p=0.297*), likely due to lower case numbers compared to the Pan-Cancer Atlas database

(*p<0.001*) (Extended Data Fig. 3a). While the IHC-quantified ATF6 expression did not

associate with CRC patient disease–free survival in our human cohorts (Extended Data Fig.
3b-d), Kaplan-Meier analysis in both databases showed a significantly reduced disease-free
survival in cases that were classed as *ATF6*-only amp/gain (p<0.001 pan-cancer, p=0.0122
TCGA) and those classed as combined *ATF6* and *FASN* amp/gain (p<0.0001 pan-cancer,
p<0.0001 TCGA), compared to those cases unaltered in either of the two genes (Extended
Data Fig. 3e,f). Importantly, disease-free survival was significantly reduced in cases of
combined *ATF6* and *FASN* amp/gain compared to those showing *ATF6*-only amp/gain in the
Pan-Cancer Atlas (*p=0.0394*), with a trend observed in the TCGA database (*p=0.4046*)
(Extended Data Fig. 3e,f). Amp/gain modifications were not associated with overall survival in
the two databases (Extended Data Fig. 3g,h).

ATF6 drives FA elongation and LCFAs accumulate in mice and CRC patients

ATF6 signaling clearly alters epithelial metabolism. To understand its effects on the
metabolome, we performed untargeted metabolomics of nATF6^{IEC} cecal content at pre-tumor
(5wk), tumor (12wk) and late-tumor (20wk) time points using LC-TOF-MS (left side schematic,
Fig 3a). In parallel, we performed untargeted metabolomics of CRC patient tumor tissue (T)
and patient-matched tumor adjacent (T adjacent) tissue in a cohort of 259 EOCRC and LOCRC
patients (cohort 4; right side schematic, Fig. 3a). In our nATF6^{IEC} murine model, principal
component analysis revealed a drastic difference in the metabolome between tumor-bearing
mice (red) and non-tumor mice (grey), while metabolite profiles clustered in closer proximity
for all three genotypes at the pre-tumor time point (Extended Data Fig. 4a). Differential
enrichment analysis at the tumor time point demonstrated a depletion of peptide metabolites
and sphingolipids, while long-chain fatty acids (LCFA) and other complex lipid species were
enriched in T mice (Extended Data Fig. 4b). Principal component analysis showed that lipid
profiles of non-tumor (NT) mice separate from tumor (T) mice already at the pre-tumor time
point, with increasing diversity over time (Extended Data Fig. 4c). Differential metabolite
analysis of the significantly regulated metabolites, shown between all genotypes and time
points, demonstrated that tg/tg associated metabolites are mostly comprised of
lysophospholipids and LCFAs (Extended Data Fig. 4d). In the CRC patient cohort 4, we
identified 56 FAs that were significantly regulated in T tissue compared to T adjacent tissue
(Table 1). We next examined the overlap between LCFAs (≥C20) which were significantly
regulated in CRC patient T tissue and in T mice. Multiple fatty acids were significantly regulated
in both, including Docosatetranoic acid (FA 22:4), Docosapentanoic acid (FA 22:5),
Docosahexanoic acid (FA 22:6), and putative FAs: FA 20:3, C20 hydroxy FAs, FA 24:5 and
FA 24:6 (Fig. 3b,c). Stratifying by timepoint, an unknown C20 hydroxy fatty acid was the only
LCFA enriched in tg/tg mice at the pre-tumor time point, however the number of significant
shared LCFAs increased at the tumor and late-tumor time points, indicating that tumor onset

increases the LCFA pool (Fig 3b). When further stratifying significantly regulated FAs in CRC patient cohort 4 into EOCRC and LOCRC, we identified two LCFAs (C20:2 hydroxy FA and C20:3 FA) to also be significantly regulated in EOCRC T tissue compared to T adjacent tissue (Extended Data Fig. 4e). FAs 22:4, 22:5 and 22:6 have been validated by comparison with pure standards (Extended Data Fig. 4f).

To further characterize the altered lipid environment in tg/tg mice, we quantified total FAs in colonic tissue using GC-MS. Analyses revealed an increase in saturated fatty acids (SAFAs) in T compared to NT mice, which also correlated with tumor number (Extended Data Fig. 4g,h). Furthermore, T mice show a clear elongation of SAFAs including FA 20:0 and FA 22:0 compared to NT mice (Fig. 3d). To determine whether elongation is a direct consequence of ATF6 signaling, we assessed ATF6-mediated elongation of FAs in *ex vivo* colon organoid cultures. We first phenotypically characterized colon organoids derived from fl/fl controls and tg/tg mice at the late-tumor time point (20wk). Both fl/fl and tg/tg crypts formed viable organoids in culture, including cyst formation (d1) and budding (d7) of cysts (Extended Data Fig. 4i). Strikingly, tg/tg organoids in culture did not show any significant increase in organoid size (Extended Data Fig. 4j) or the number of budding crypts per cyst (Extended Data Fig. 4k) compared to fl/fl controls, indicating that the hyper-proliferative phenotype observed *in vivo* in the presence of the microbiota is lost in this *ex vivo* culture system. In a separate experiment, intestinal organoids from nATF6 Vil-Cre^{ERT2} (fl/fl^{ERT2} and tg/tg^{ERT2}) mice were cultured, and *ATF6* expression induced by adding 500nM (Z)-4-hydroxytamoxifen (4-OHT) to the culture medium for 24h. This was followed by a 24h exposure of organoids to deuterated eicoseinoic acid (D$_3$-FA 20:0) (Fig. 3e, Supplementary Fig. 1). Importantly, tg/tg^{ERT2} intestinal organoids showed a significantly increased potential of FA elongation, quantified as the percentage of metabolized D$_3$-FA20:0 to D$_3$-FA:22:0, compared to fl/fl^{ERT2} controls (Fig. 3f). The above findings demonstrate that ATF6 activation is the driver of FA elongation in the absence of microbial and immune signaling.

ATF6-related lipid profiles are causally linked to tumor-associated microbiota

We next sought to investigate whether chronic *Atf6* activation would alter the luminal and mucosa-associated microbiota. To isolate the impact of tumor formation, we first examined changes in microbiota composition at 5wks, before tumor onset. We observed augmented modulation of the mucosa-associated microbiota with increasing *Atf6* gene dose as shown by mean GUniFrac dissimilarity relative to control, between genotypes, an effect comparable to changes observed in the luminal environment (Fig. 4a). Global microbiota shifts were observed between all timepoints and genotypes, in both luminal and mucosal datasets, and appeared more distinct at later timepoints, particularly in tg/tg mice (Extended Data Fig. 5a,b; luminal PERMANOVA, p=0.001, mucosal PERMANOVA p=0.001). We applied machine learning

models (Random forest (RF), Ridge regression (RR), LASSO (L), Elastic-net (E) and L1-penalized regression (LL)) to both sample types, demonstrating that the mucosal microbiota can better classify the tumorigenic phenotype compared to the luminal microbiota. Supporting increased utility of mucosal over luminal microbiota, four out of five models trained on mucosal microbiota data showed higher AUC values across all leave-one-out cross validation repetitions (Extended Data Fig. 5c). Moreover, we observed a significant correlation between tumor number and Shannon Effective diversity in mucosal data which was absent in luminal (Extended data Fig. 5d,e), further supporting this finding and suggesting a link between tumor burden and microbial diversity.

To examine whether these changes in microbiota community structure were associated with shifts in metabolite profiles, we performed a multi-omic data integration using sPLS-DA analysis of luminal untargeted metabolomic profiles and luminal and mucosal 16S data [44]. To delineate between genotype-driven and tumor-driven alterations, we performed two integrations that model genotype and phenotype separately. By sorting tg/tg discriminating features according to importance, we identified Lactobacillus/Limosilactobacillus and Desulfovibrio zOTUs as well as multiple LCFAs and lysophospholipids as microbial and metabolite contributions, respectively (Fig. 4b). Metabolite contributions to phenotype discrimination were similar and comprised mostly LCFAs and Lysophospholipids. However microbial contributions differed greatly, with mostly zOTUs classified as Parabacteroides and Romboutsia, indicating the presence of tumors differentially modulates microbiota, compared to *Atf6* activation alone (Extended data Fig. 6a). Next, to identify microbiota-correlated metabolites, we generated relevance networks using the network function of mixOmics DIABLO R package [44]. Importantly, several fatty acids, including an unidentified C20 hydroxy fatty acid, nervonic acid (FA 24:1), Docosapentanoic acid (FA 22:5) and Docosatetranoic acid (FA22:4), correlated positively with tg/tg discriminative zOTUs, particularly those classified as Lactobacillus/Limosilactobacillus, and additionally with Desulfovibrio in the mucosal dataset (Extended data Fig. 6b,c, Supplementary Table 6). LCFA compounds are known to be toxic to bacteria, requiring adaptation mechanisms to allow growth in FA-rich milieus. Using PICRUSt2, we generated predictions of genomic content [45,46], following a spatially-resolved analysis of 16S rRNA amplicon sequencing-based bacterial profiles using individually defined 0.5 x 0.5 cm sites along the longitudinal colonic axis in fl/fl controls and tg/tg mice at the tumor (12wk) time point. We identified genes involved in bacterial responses to FAs in the proximal tumor-susceptible region of the colon, which are known to be involved in FA detoxification, catabolism and efflux (Extended Data Fig. 7a). Of this FA-response signature, oleate hydratase (*ohyA*) and the FA efflux pump (*farE*) were observed to be upregulated in tumor mice compared to controls (Extended Data Fig. 7a). *OhyA*-mediated fatty acid hydroxylation enables growth of certain bacteria in lipid-rich environments[47]. Spatial mapping of $\log_{10}$

transformed relative abundance showed clear enrichment of *ohyA* across all sites in tg/tg mice

compared to fl/fl controls (Fig. 4c). The relative abundance of *ohyA* was increased in tumor-

bearing tg/tg mice compared to fl/fl controls (Fig. 4d). Furthermore, the ratio of *ohyA*+ to *ohyA*-

taxa was increased in all three mucosal phenotypes (non-tumor (NT), tumor adjacent (TA) and

tumor (T)) in tg/tg mice, compared to fl/fl controls (Extended Data Fig. 7b). *FarE* constitutes a

lipid efflux pump associated with antimicrobial FA resistance to limit the cellular accumulation

of toxic FAs [48]. Similar to *ohyA*, we observed an increased relative abundance across all tg/tg

sites (Fig. 4e), and increased relative abundance of *farE* in tumor-bearing tg/tg mice (Fig. 4f).

Furthermore, the ratio of *farE*+ to *farE*- taxa was increased in all tg/tg mucosal phenotypes,

compared to fl/fl controls (Extended Data Fig. 7c). Taken together, our findings clearly indicate

an early pre-tumor microbial adaptation to the ATF6-altered FA-rich intestinal environment, to

shape a tumor-associated bacterial lipid-response signature.

To understand the causal role played by FASN in colonic tumorigenesis in our nATF6^{IEC} murine

model, we exposed fl/fl and tg/tg SPF mice to bi-weekly intraperitoneal injections (i.p.) of the

FASN inhibitor C75 from the age of 3 weeks (pre-tumor time point) until 12 weeks (tumor time

point) (Fig. 4g). Interestingly, FASN inhibition reduced tumor incidence in tg/tg mice from 80%

to 0% with no tumors observed in the tg/tg colon (Extended Data Fig. 7d,e). In line with this,

FASN inhibition was able to reduce tumor number and tumor volume in the colon of tg/tg mice

(Extended Data Fig. 7f,g). To validate the link between FA synthesis and microbiota-dependent

tumorigenesis in nATF6^{IEC} mice, we performed microbiota transfer experiments from C75-

treated mice and RPMI-treated controls into GF tg/tg mice and fl/fl controls (Fig. 4g). Compared

to fl/fl recipient mice, all tg/tg recipient mice show reduced survival (Extended Data Fig. 7h).

Strikingly, the tumor-promoting capacity of the C75-treatment-derived microbiota was reduced

from a tumor incidence of 73% to 25% (Fig. 4h). In line with this, the microbiota from C75-

treated mice reduced both tumor number and volume to levels comparable to tumor-free fl/fl

controls (Fig. 4i, Extended Data Fig. 7i). Taken together, these findings clearly demonstrate

that microbiota changes associated with ATF6-induced lipid alterations is causally involved in

inducing a more aggressive tumorigenic phenotype in nATF6^{IEC} mice.

LCFAs selectively activate the growth of tumor-related bacteria

To assess whether LCFAs directly impact bacteria, we exposed the complex microbiota of 5

week-old fl/fl mice to previously identified (Fig. 3b, Table 1) ATF6-associated LCFAs (24:1,

22:0, 20:0, 22:6, 22:5, 20:3 and 22:4). Our approach applied biorthogonal non-canonical amino

acid tagging (BONCAT) combined with fluorescence-activated cell sorting (FACS) to identify

translationally active bacteria in response to LCFA exposure (Fig. 5a). Confocal microscopy

images displayed a significantly higher proportion of translationally active cells after LCFA

supplementation compared to the controls (Fig. 5b, Supplementary Fig. 2). Microscopy images

were confirmed with quantitative measure of translationally active bacterial cells exhibiting a
higher percentage of translational activity compared to the controls (ANOVA, p<0.0001 for all
comparisons) (Fig. 5c). A distinct separation between sorted and unsorted communities along
with a significant decrease of zOTUs in the sorted fraction, demonstrates a specific microbial
response to LCFA exposure (PERMANOVA, p<0.0001; Wilconox test: p<0.0001) (Extended
Data Fig. 8 a,b). In total, we identified 50 zOTUs that were significantly enriched in response
to one or multiple LCFAs, while only one zOTUs was depleted (Wald test, p<0.05) (Fig. 5d,
Extended Data Fig. 8c). Strikingly, LCFA selectively activated tumor-associated bacteria
including those of the Desulfovibrionaceae family (e.g., *Desulfovibrio fairfieldensis* and
Bilophila spp.) (Fig 5 d), while the overall microbiota profiles were similar in response to the
different LCFAs (Extended Data Fig. 8d; PERMANOVA, p=0.8045). Supporting the *in vivo*
relevance of these findings, we examined the overlap between LCFA-enriched zOTUs and
differentially abundant zOTUs at the pre-tumor timepoint, identifying *Desulfovibrio fairfieldensis*
as one of three significantly enriched zOTUs in tg/tg mice at the pre-tumor timepoint in both
the mucosal and luminal environments (Fig. 5e, Extended Data Fig. 8e). The higher
translational activity in Desulfovibrionaceae was accompanied by a significant enhancement
in bacterial growth when exposed to individual LCFAs, as indicated by the increased expansion
of *Desulfovibrio fairfieldensis* (two-way ANOVA, p<0.0001 for all comparisons) (Fig. 5f,
Extended Data Fig. 8f). A similar growth-promoting effect was observed for another
Desulfovibrio isolate, *D. piger*, further confirming the ability of Desulfovibrio spp. to selectively
grow in the presence of LCFAs (p<0.0001 for all comparisons; Extended Data 8g). Additional
zOTUs responding to the LCFA-rich environment included Sutterella spp., Lachnospiraceae,
Muribaculaceae, *Turicimonas muris*, *Ruminococcus gnavus*, Oscillibacter spp. (Fig. 5d). The
above demonstrates that LCFAs translationally activate specific bacteria, including tumor-
associated taxa, providing a direct link between tumor-associated fatty acid profiles and the
selective expansion of bacterial taxa in mice.
Recent developments in identifying microbial signatures in the publicly available The Cancer
Genome Atlas (TCGA) dataset [49], allowed us to compare computationally inferred transcription
factor activity using previously validated transcription factor-target gene relationships (RNA-
seq data) and microbial profiles (identified in whole-exome sequencing data from the same
samples) in CRC patients. In a hypothesis-free approach, ATF6 was identified among the most
significant transcription factors, as calculated using a linear mixed effects model, whose
activities correlated positively with the presence of CRC-enriched genera, including
Campylobacter, Porphyromonas, Fusobacterium, Selonomonas and Desulfovibrio (Fig. 5 g,h).
Both *ATF6* activity and bacterial profiles are strongly associated with microsatellite instability
(MSI) status of the cancer, which could be a potential confounding factor, but we observed a
strong correlation of the CRC-enriched genera with ATF6 activity even after correcting our

association models for MSI status (Extended Data Fig. 8h). Of note is the observed overlap in
specific tumor- and ATF6-associated taxa between mouse and human microbiota (Fig.
5d,e,g,h and Extended Data Fig. 8h), namely *D. fairfieldensis*, Oscillibacter spp.,
Desulfovibrionaceae spp. Bilophila spp., supporting a causal link between ATF6-mediated
LCFA synthesis and tumor-relevant changes in the microbiota.
473

Discussion

As the second leading cause of deaths worldwide [50], early and accurate diagnosis as well as novel treatment strategies are imperative in the management of CRC. As a fundamental mediator in the pathogenesis of CRC, the UPR^{ER} represents a promising host cellular mechanism for preventive and therapeutic intervention. While our mechanistic understanding of the role of ATF6 signaling in CRC remains limited, its expression has been associated with poor prognosis in CRC patients [14,15]. In the present study, we demonstrate the relevance of ATF6 expression in CRC patients across multiple independent German cohorts by identifying a clear sub-population (19-38% LOCRC; 11% EOCRC) of patients that present with high ATF6 expression, underlining the importance of ATF6 in CRC. Most importantly, we identified a mechanism through which ATF6-mediated changes in cellular lipid metabolism of the intestinal epithelium contribute to the development of tumor-promoting microbiota changes at an early disease stage.

We previously showed that chronic ATF6 activity specifically in murine intestinal epithelial cells (nATF6^{IEC}) results in microbiota-dependent early-onset colonic adenomas [11]. The necessity of microbial triggers in tumor induction was shown via GF mice, which remained tumor-free [11]. Furthermore, the transfer of pre-tumor murine cecal content into GF mice revealed the dysbiotic microbiota from tumor-susceptible mice to have an increased capacity to induce tumors [11]. Importantly, the development of dysbiosis in nATF6^{IEC} mice precedes tumor formation [11], suggesting that tumor-independent mechanisms at early stages of the ATF6-driven pathogenesis initiate microbial changes. To ascertain these mechanisms, we set out to characterize the transcriptional response induced by chronic ATF6 activation in the presence (SPF) and absence (GF) of bacteria at the pre-tumor stage. We identify the FA metabolism KEGG pathway as one of the top ranked metabolic pathways altered in tumor-susceptible SPF mice when compared to GF mice, and detect 23 upregulated FA metabolism-related genes. The majority are upregulated already in response to acute ATF6 activation, confirming FA alterations to be an early event occurring in direct response to ATF6 signaling in the presence of the microbiota. While the specific contribution of ATF6 to lipid metabolism is less well understood, the UPR^{ER} is known to play an important role in lipid metabolism [51-54]. Lipid metabolic rewiring forms a phenotypic hallmark of cancer, with altered lipidome profiles identified in tumors of numerous organs [37,55]. Ecker *et al.* recently identified a robust CRC tumor-specific triglyceride species signature that correlates with postoperative disease-free survival in a multicohort study spanning two clinical centres [43]. Moreover, gene expression of *FASN*, which forms a multi-enzyme complex responsible for catalyzing FA synthesis, was prognostically detrimental [43]. Strikingly, the *in vivo* inhibition of FASN in nATF6^{IEC} mice results in the prevention of colon adenoma development. We show that our tumor-associated cecal

metabolite signature is mostly comprised of lysophospholipids and LCFAs, with an overlap of
several significantly altered LCFAs between tg/tg mice and CRC patient tumor tissue,
confirming the human relevance of the observations in our transgenic murine tumor model.
Analyses of murine colonic tissue demonstrated increases in SAFAs and LCFAs. These
findings are in line with a study profiling FAs in tissue and plasma of CRC patients and healthy
controls, which identified a CRC-associated FA panel and found that LCFAs were the most
increased FAs [56]. Our experimental approach in *ex vivo* organoids demonstrates that FA
elongation is a direct consequence of ATF6 signaling, occurring in the absence of microbial
signals.

A handful of studies have identified pro-oncogenic bacteria enriched in adenoma samples,
however, questions remain over whether these taxa initiate disease, since adenoma formation
induces metabolic alterations that likely impact the local microenvironment and alter microbial
composition [57-59]. Our knowledge on which triggers may enable taxa to expand early on, and
potentially become tumor-promoting, is sparse. We hypothesize that ATF6-driven lipid
alterations impact the intestinal microbiota as an early event of tumorigenesis in nATF6[IEC]
mice. Our multi-omic data integration of 16S rRNA microbiota (luminal and mucosal) profiling
and   cecal   metabolites   identified   several   zOTUs,   including   those   classified   as
Lactobacillus/Limosilactobacillus and Desulfovibrio, to be correlated with several LCFAs
associated with our tumor-susceptible mice. Importantly, and in support of our hypothesis,
PICRUSt2 predictions of genomic content confirm that the ATF6-driven FA-rich intestinal
environment results in microbial adaptation in the form of a lipid-response signature that is
associated with genes involved in FA detoxification, catabolism and efflux. Our data suggest a
clear link between ATF6-induced lipid alterations and changes in the intestinal microbiota
preceding tumor formation. Strikingly, FMT in GF mice using cecal content of FASN inhibitor-
treated mice clearly demonstrated that the microbiota associated with ATF6-induced lipid
alterations is causally involved in inducing a more aggressive tumorigenic phenotype in
nATF6[IEC] mice. To identify selective taxa of the intestinal microbiota responding to LCFAs, we
here applied an approach to directly expose the microbiota of fl/fl control mice *ex vivo* to our
identified LCFAs, showing that tumor-associated taxa are translationally activated in response
to the LCFAs. Among these was *Desulfovibrio fairfieldensis*, which was significantly enriched
in tg/tg mice at the pre-tumor time point in both the mucosal and luminal environments. In
addition to CRC-associated lipid signatures, CRC-associated microbial signatures have also
been identified [60,61]. In a hypothesis-free approach analyzing transcription factor activity and
microbial profiles in CRC patients, we show *ATF6* activity to positively associate with the
presence of such a risk profile associated bacterial genera, identifying a clear link between
ATF6 signaling and the microbiota in CRC pathogenesis. Interestingly, and in confirmation of
our murine data, we identified Desulfovibrio among the ATF6-associated CRC genera.

In conclusion, we identify a clear mechanistic sequence of events in the initiation of colon tumorigenesis whereby ATF6 signaling drives a clinically relevant pathologic response by altering lipid metabolism to induce microbial adaptation, ultimately leading to a tumor-relevant bacterial dysbiosis. Although beyond the scope of this study, future work should focus on single-cell transcriptomics to characterize the ATF6 transcriptional response at the cellular level, as well as extending gnotobiotic mouse experiments with individual bacterial candidates, to identify causal tumor drivers and elucidate mechanisms of tumor initiation. Together, the above highlights novel mechanistic findings in an exciting new avenue of research bringing together cellular stress responses, lipid metabolism and the intestinal microbiota in the context of CRC. This evident triangular interplay presents new targetable vulnerabilities in CRC that warrant future research efforts to identify potential future prevention and treatment strategies.

Funding

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Collaborative Research Center CRC1335 (Aberrant immune signals in cancer; project no. 360372040; P11) and CRC 1371 (Microbiome signatures: functional relevance in the digestive tract; project no. 395357507; P13 and P18), as well as the Bundesministerium für Bildung und Forschung (BMBF, Microbiome-based Prevention of Early-Onset Colon and Rectal Cancer (Mi-EOCRC); project no. 100509410).

Competing interests: The authors declare no competing interests.

Author contributions

OC and DH wrote the manuscript. AS, AR, MS, DS and CS supported manuscript writing. OC, AS, AR, MS, SB, SK, IC, MS, DS and DH designed the experiments and performed data analysis. AS, MS and SK performed mouse experiments and respective mouse tissue and microbiota processing and analysis. KS performed IHC staining on human cohort TMAs. OC performed the quantification and data analysis of human IHC staining. TM supported quantification of human IHC staining. MJ provided CRC cohort 1 tissue and supported data analysis. KPJ provided CRC cohort 2 tissue and generated TCGA survival curves. CS, SH, CR (Christoph Röcken) and CR (Christian Röder) provided CRC cohort 3 tissue and supported data analysis. MZ provided untargeted metabolomics analyses of CRC cohort 4. DMC, SW and TP performed ATF6 quantification validation classification and regression models. JW and GZ performed microbial and transcription factor TCGA analyses. SB prepared mouse tissue for mRNA sequencing samples and supported mRNA sequencing analyses. PR performed mRNA sequencing. CS and TK performed mRNA sequencing data analyses. AD, DS and MZ performed untargeted metabolomics. JP (Josh Pauling) and NK performed metabolomics analysis and integration. JP (Johannes Plagge), JE and GL performed fatty acid analyses and data analysis.

Data Availability

Raw and processed RNASeq data are available at GEO under accession GSE247122. Human CRC metabolomics data is available in the MetaboLights Database [62] (https://www.ebi.ac.uk/metabolights/) under accession number MTBLS7387. 16S

590 rRNA gene amplicon sequence data have been deposited in the NCBI Short Read Archive
591 under PRJNA1227744. Source data are provided with this paper.

Acknowledgements

ORCID IDs

Olivia Coleman https://orcid.org/0000-0002-4075-7820

Adam Sorbie https://orcid.org/0000-0002-3330-4612

Alessandra Riva https://orcid.org/0000-0002-5724-7616

Denise Selegato https://orcid.org/0000-0001-7642-6524

Nikolai Köhler https://orcid.org/0000-0002-5103-883X

Jakob Wirbel https://orcid.org/0000-0002-4073-3562

Christina Schmidt https://orcid.org/0000-0002-3867-0881

Tim Kacprowski https://orcid.org/0000-0002-5393-2413

Andreas Dunkel https://orcid.org/0000-0002-4445-6144

Johannes Plagge https://orcid.org/0000-0002-7180-8221

Tingying Peng https://orcid.org/0000-0002-7881-1749

Thomas Metzler https://orcid.org/0000-0002-1620-5559

Michael Zimmermann https://orcid.org/0000-0002-5797-3589

Philipp Rosenstiel https://orcid.org/0000-0002-9692-8828

Katja Steiger https://orcid.org/0000-0002-7269-5433

Moritz Jesinghaus https://orcid.org/0000-0002-0018-5661

Gerhard Liebisch https://orcid.org/0000-0003-4886-0811

Georg Zeller https://orcid.org/0000-0003-1429-7485

Klaus-Peter Janssen https://orcid.org/0000-0002-4707-7887

Dirk Haller http://orcid.org/0000-0002-6977-4085

Online Methods

Ethics statement regarding animal experiments. All animal experiments, as well as maintenance and breeding of mouse lines, were approved by the Committee on Animal Health Care and Use of the state of Upper Bavaria (Regierung von Oberbayern; AZ ROB-55.2-1-54-2532-217-2014, AZ ROB-55.2-2532.Vet_02-20-58, AZ TVA 55.2-2532.Vet_02-18-149, AZ TVA 55.2-2532.Vet_02-18-121, AZ TVA 55.2-2532.Vet_02-19-006) and performed in strict compliance with the EEC recommendations for the care and use of laboratory animals (Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes).

Ethics statement regarding patient samples. Immunohistochemistry staining and quantification of ATF6 were performed in three independent human cohorts: a German large single-center cohort comprising 1004 colorectal cancer patients that underwent surgical resection spanning two decades, for which 959 patient tissue samples could be quantified (cohort 1) [17], a German cohort consisting of 104 CRC patients followed up over 20 years, for which 50 patient samples could be quantified (cohort 2), and a German single center cohort comprising more than 1,500 resected CRC patient samples spanning approximately 25 years, for which 55 EOCRC (age <50y) and 256 LOCRC (age >50y) were obtained and could be quantified (cohort 3). The use of surgically resected human tissue samples was approved by the local Ethics Committee of the Technical University of Munich (TUM) (ref.-no. 252/16 s, cohort 1), by the Ethics Committee of the Medical Faculty of TUM (ref.-no. 1926/7, 375/16S and 2022-169-S-KH, cohort 2), and by the local Ethics Committee of the University Hospital Schleswig-Holstein (ref.-no. A 110/99, cohort 3). CRC patient cohort 3 tissue specimens were supplied by the biobank TRIBanK (Translational Interdisciplinary Biobank Kiel) together with the Institute of Pathology, University Hospital Schleswig-Holstein, Kiel, Germany. LC-MS analyses were performed in a fourth German human cohort consisting of 259 patients diagnosed with EOCRC and LOCRC and included one tumor tissue (T) and one adjacent tissue (T adjacent) for each patient (cohort 4). The use of surgically resected human tissue samples was approved by the local Ethics Committee of the Medical School – The Christian-Albrechts-University of Kiel (ref.-no. A 156/03), the Ethics Committee of the Medical School of the University of Rostock (ref.-no. A 2019-0187) and the Ethics Committee of the EMBL (ref.-no. 2024/HE000062 - MiEOCRC). For all human cohorts in this study, both sexes were deliberately included, with slightly more male cases, corresponding to the incidence in the overall population. Sex-dependent differences were not observed in our data. Patient samples from all four cohorts were obtained after prior informed written consent, and information on sex was acquired by self-reporting.

Immunohistochemical staining and quantification of ATF6. TMA cores were immunohistochemically stained with an anti-ATF6 antibody (Sigma-Aldrich, HPA-005935) diluted 1:100, following antigen retrieval in an EDTA buffer (30 min, pH 9), or an anti-FASN antibody (LS Bio, LS-B3636) or anti-GRP78 (Cell Signaling, 3177s) diluted 1:200 following antigen retrieval in a Citrate buffer (20 min, pH 6). Nuclei were counterstained with hematoxylin (Waldeck). Stainings were performed using the Leica Bond RXm (Leica) and scanned using the Aperio AT2 (Leica Biosystems). ATF6- and FASN-stained TMA cores were quantified using the QuPath Software [18]. Cells were identified through positive cell detection of the hematoxylin-positive nucleus. ATF6 staining intensities for the nuclear and the cytoplasmic regions were classified according to defined thresholds of the DAB OD max and DAB OD mean, respectively. FASN staining was quantified based on cytoplasmic regions only. Furthermore, the object classifier was trained and applied to distinguish between epithelial cells and stromal cells. Exported epithelial histochemical-scores (H-scores, capturing both the intensity and the percentage of stained cells) for the nucleus and the cytoplasm were used for further patient cohort analyses

ATF6 expression analysis using AI. Whole slide images were segmented using QuPath into individual cores and scaled down to 312x312 pixels. This pre-processing step yielded a dataset of 176 TMA cores from a cohort of 50 patients (cohort 2). For our deep learning model, a ResNet18 was trained (arXiv:1512.03385v1 [cs.CV]), freezing all layers except the final convolutional and fully connected layers for the classification task and freezing the first layer for the regression task. This allowed leveraging pre-trained features while fine-tuning the model's output layers for our specific task. To improve model generalization, data augmentation techniques were applied, including random horizontal and vertical flips and random rotations. In the final layer, dropout regularization was incorporated to prevent overfitting. A 5-fold cross-validation was conducted to evaluate our model's performance. Each fold used data from 40 patients for training and 10 patients for validation. Throughout cross-validation, the best models were recorded based on their accuracy (classification task) or loss (regression task) in the validation dataset. Their weights were preserved for subsequent use. In our final prediction task of classifying samples as ATF6-high or ATF6-low, a weighted average ensemble strategy was employed by combining the predictions of the five best models. In a final evaluation step, our ensemble's predictions were tested on an entirely independent cohort comprising 918 patients (cohort 1).

Transgenic nATF6^{IEC} mice (SPF and GF). nATF6^{IEC} mice were generated as previously described [11] and housed under SPF and GF conditions (12 h light/dark cycles and 24-26°C) at the Technical University of Munich (Weihenstephan, Germany). All mice were fed a standard chow diet (autoclaved V1124-300; Ssniff, Soest, Germany) *ad libitum.* Mice were cage-

separated according to sex but not according to genotype. Both sexes were used for all experiments in this study. For the determination of the tumor phenotype, mice were sacrificed and the colon excised. The presence and number of tumors was recorded for each mouse as macroscopically observed.

Isolation of murine primary colonic IECs. Colonic tissue was longitudinally opened, and transferred to 20 mL Dulbecco's modified Eagle medium (Gibco) containing 10% fetal calf serum, 1% L-glutamine, and 0.8% antibiotics/antimycotics (IEC isolation medium) supplemented with 1 mmol/L dithiothreitol (Roth). The tissue was vigorously vortexed for 1 minute, incubated under continuous shaking (37°C, 15 minutes) and vortexed again for 1 minute. The cell suspension was centrifuged (7 minutes, 300xge) and the cell pellet was resuspended in 5 mL IEC isolation medium. After vortexing for 1 minute, the remaining tissue pieces were incubated in 20 mL PBS (10 minutes, 37°C) containing 1.5 mM EDTA under continuous shaking. The cell suspension was vortexed again followed by another centrifugation to pellet the cells. Both cell suspension fractions were combined and purified by centrifugation through a 20%/40% (in medium/PBS) discontinuous Percoll gradient (GE Healthcare Life Sciences) at 600xg for 30 minutes. The IEC cells at the interface between the two Percoll phases were collected and washed twice (with medium/PBS). The isolated IECs were lysed in RA1 RNA lysis buffer (Macherey-Nagel).

RNA Isolation and library construction. RNA of colonic IECs was isolated according to the manufacturer's instructions (NucleoSpin RNAII kit, Macherey-Nagel) and measured by NanoDrop ND-1000 spectrophotometer (Thermo Fisher Scientific). TruSeq stranded Total RNA kit (Illumina) was used for the construction of strand-specific sequencing libraries from the isolated colonic IEC RNA. Sequencing was performed on an Illumina HiSeq2500 (75-nucleotide (nt) paired-end reads for each run, 1/10 lanes per sample.

RNASeq preprocessing. After the initial trimming of reads using trim_galore (version 0.6.4) to satisfy a Phred score of at least 30, RNASeq data was processed with the 'nf-core/rnaseq' pipeline (version 1.4.2) and aligned to the GRCm38 genome using the 'STAR' aligner (version 2.6.1d). Quantification of transcripts was performed using Salmon (version 0.14.1). All sequence properties were inspected in MultiQC and passed the quality thresholds.

RNAseq functional analysis. Gene Set enrichment analysis (GSEA) was performed using the differential expression results ranked by t-value [63]. KEGG gene set was downloaded from the Molecular Signatures Database [63-65]. Gene names were translated into mouse gene names before the analysis using scibiomart (v.), a wrapper around the API from BioMart. GSEA was performed using the packages fgsea [66] (v1.24.0) and GSEABase [67] (v 1.60.0). Next we performed regulatory clustering based on regulatory rules (Supplementary Table 3, sheet 2) using sircleR package function sircleRCM_2Cond [19]. In brief, each gene was assigned to "UP"

(log2FC > 0.5, *p*.adj <0.05), "DOWN" (log2FC < -0.5, *p*.adj <0.05) or "No Change"
(0.5 > log2FC > -0.5 and/or *p*.adj > 0.05) based on its expression change between fl/fl versus
tg/tg. The background method chosen was "C1&C2", which means a gene had to be detected
in both conditions in order to be included in the clustering. Plots were generated using the
EnhancedVolcano package [68] (v1.16.0) and ggplot2 [69] (v 3.4.2).

Untargeted metabolomics of murine cecal content or tissue. Sample preparation was
performed as previously described [70]. Briefly, mouse cecal content (~20mg) or tissue (~25mg)
was mixed with 1ml methanol-based extraction solvent in a 2ml bead beater tube (CKMix 2ml,
Bertin Technologies) filled with ceramic beads (1.4 mm and 2.8 mm ceramic beads). Samples
were then homogenized using a bead beater (Precellys Evolution, Bertin Technologies)
supplied with a Cryolys cooling module (Bertin Technologies, cooled with liquid nitrogen) three
times for a duration of 20 seconds, at a speed of 8000rpm. Subsequently, the resulting
suspension was centrifuged for 10 minutes, at 6000rpm. Finally, 100µl of supernatant was
mixed with 20µl internal standard solution (7 µmol/l) and injected into the Liquid
chromatography time-of-flight mass spectrometer (LC-TOF-MS) system for untargeted
analysis. Untargeted LC-TOF-MS analysis was carried out as described in Metwaly *et al.* [70].
Briefly, untargeted analysis was performed on an ExionLC Ultra-high performance liquid
chromatography (UHPLC) system (Sciex), connected to a 6600 TripleTOF instrument (Sciex)
operating in positive and negative electrospray mode and calibrated using ESI positive and
negative calibration solutions (Sciex). UHPLC phase separation was performed in reverse as
well as hydrophilic interaction stationary phase (HILIC), on a Kinetex C18 column
(Phenomenex) and ACQUITY BEH Amide column (Water) respectively. Mass spectrometry
was performed using SWATH mode, with fragment spectra recorded in high-sensitivity mode.
Metabolomics data processing was performed as described previously [70]. Briefly, Raw data
files were converted into Reifycs Abf (Analysis Base File) files and subsequent untargeted
peak picking was performed using MS-DIAL software (version 3.5274) [71]. Alignment was
performed across all samples and the peak area of individual features were exported for further
analysis using the R statistical software environment. Peak normalisation was based on QC
samples, employing the method described in Wehrens *et al.* [72]. All features were then
combined into a single table and further normalised according to sample weight, before
analysis.

Fatty acid quantification. For FA quantification in mouse tissues and investigation of
eicosanoic acid elongation in organoids, FA methyl esters (FAMEs) were generated by acetyl
chloride and methanol treatment, and extracted with hexan [73]. Total FA analysis was carried
out using a Shimadzu 2010 GC-MS system (Duisburg, Germany). FAMEs were separated on
a BPX70 column (10 m length, 0.10 mm diameter, 0.20 µm film thickness) from SGE using

helium as the carrier gas. The initial oven temperature was 50°C and was programed to increase at 40°C/min to 155°C, 6°C/min to 210°C, and finally 15°C/min to 250°C. D3-eicosanoic acid and its elongation products in organoids were quantified by single-ion monitoring of specific fragment ions (D3-eicosanoic acid, 329 m/z; D3-docosanoic acid, 357 m/z). The internal standard was C21:0-iso.

Untargeted metabolomics of CRC patient tissue sample preparation. Solid tissues were prepared for LC-MS analysis by organic solvent extraction. In brief, 200 µL of 0.1 mm zirconia/silica beads (BioSpec Products) and 500 µL of organic solvent (acetonitrile:methanol, 1:1) supplemented with an internal standard mixture were added to 70-100 mg of pre-weighed solid material. Internal standard mixture consisted of phenylalanine-*d5*, tryptophan-*d5*, ibuprofen-*d4*, tolfenamic acid-*d4*, estriol-*d3*, diclofenac-*d4*, warfarin-*d5*, oxfendazole-*d3*, chloramphenicol-*d5*, nafcillin-*d5* and caffeine-*d9*, each to a final concentration of 80 nM. The material was homogenized by mechanical disruption with a bead beater (BioSpec Products) set for 2 minutes and 30 seconds on the highest setting at room temperature. After incubation for at least 1 h at -20°C, samples were centrifuged (3220 rcf, 4°C) for 15 min. For C18 LC-MS acquisition, 15 µL of supernatant was diluted with 15 µL H2O.

Untargeted metabolomics of CRC patient tissue LC-MS data acquisition. Chromatographic separation was performed by reversed-phase chromatography using an Agilent 1200 Infinity UHPLC system. The qTOF instrument (Agilent 6550) was operated in negative scanning mode (50–1,700 *m/z*). Online mass calibration was performed using a second ionization source and a constant flow of reference solution (*m/z* 112.9857 and 1033.9881). InfinityLab Poroshell HPH-C18 column (Agilent, 2.1 x 100 mm, 1.9 Micron particle size) was used for chromatography. LC solvent A was water with 0.1% formic acid (v/v) and solvent B was methanol with 0.1% formic acid (v/v). Column operated at 45 °C. 5 µL of sample were injected at the following gradients: 0 min 5% B; 0.1 min 5% B; 5.50 min 95% B; 6.49min 95% B; 6.5 min 5% B. The time between injections was 0.5 min. The qTOF was operated in negative scanning mode (50 - 1700 m/z) and the following source parameters: VCap: 3500 V, nozzle voltage: 2000 V, gas temp: 225 °C; drying gas 13 L/min; nebulizer: 20 psig; sheath gas temp 225 °C; sheath gas flow 12 L/min. Tandem mass spectrometry analysis (LC-MS/MS) was performed for fatty acids using the chromatographic separation and source parameters described above and the targeted-MS/MS mode of the instrument with a preferred inclusion list for parent ion with 20 ppm tolerance, Iso width set to 'narrow width' and collision energy to either 10, 20 or 40 eV.

Semi-targeted metabolomics of CRC patient tissue LC-MS data processing. An *in-silico* library of 134 fatty acids was mined again C18-MS negative data for the extraction of area values of FA-putative features (Supplementary Table 7). The area under the curve from the

targeted analysis was integrated using the MassHunter Quantitative Analysis Software
(Agilent, version 7.0) based the accurate high-resolution mass of these reference analytes and
the following parameters: signal threshold of 50,000; mass tolerances of 0.002 amu or 20 ppm;
retention time tolerance of 0.2 min. The resulting table was further normalized in R 4.4.1 using
RStudio/2023.06.1 according to established protocols [74,75].

Validation of significant fatty acids of CRC patient tissue. To ensure the annotation of the
significant fatty acids, we have performed level-1 metabolite annotation, which is the direct
comparison of an authentic chemical standard analyzed under identical analytical conditions.
For this, we have used standards of docotetranoic acid, docopentanoic acid and docohexanoic
acid, as well as samples of CRC-patients that have shown high abundance of these
metabolites. Correct annotation occurs when all three orthogonal properties of the metabolite
of interest (retention time, high-resolution mass, and fragmentation mass spectrum) are similar
between the analyte and the CRC-sample.

Murine primary crypt isolation and intestinal organoid culture. Murine intestinal crypts
were isolated following digestion with EDTA (30 min; 4°C) and mechanical dissociation. The
supernatant was filtered through a 70µm cell strainer, pelleted by centrifugation, and
resuspended in Matrigel (BD Biosciences). 25µl Matrigel®-organoid domes were plated in 48-
well plates (Eppendorf) together with 250µl IntestiCult™ Organoid Growth Medium
(STEMCELL Technologies) and maintained in a humidified 5% CO_2 atmosphere at 37°C.
Organoids were passaged every 7 days by mechanical disruption using a 1ml syringe with a
20G needle, pelleted by centrifugation, and embedded in fresh Matrigel.

Induction of *ex vivo* recombination and fatty acid stimulation of intestinal organoids.
Intestinal organoids from nATF6 Vil-Cre[ERT2] (fl/fl[ERT2] and tg/tg[ERT2]) mice [11] were passaged three
days before the start of the experiment and cultivated in self-made crypt culture medium (CCM;
advanced DMEM/F12 medium (Gibbco), 2mM GlutaMax (Gibbco), 10mM HEPES, penicillin,
streptomycin and amphotericin (all Sigma-Aldrich) supplemented with N2, B27 (both Gibbco),
1mM N-acetylcysteine (Sigma-Aldrich), 50ng/ml EGF (ImmunoTools), 100ng/ml noggin and
0.5µg/ml R-spondin 1 (both PeproTec)). *ATF6* overexpression in intestinal organoids was
induced by adding 500nM (Z)-4-hydroxytamoxifen (4-OHT; LKT) to the culture medium for 24h.
After induction, organoids were exposed to Eicosanoic 20,20,20-D3 acid (30µM; Larodan) in
CCM for 24h. Organoids were subsequently harvested in 0.2% SDS in ddH_2O.

Sampling of murine cecal content and colonic mucosa-associated microbiota. Cecal
content was harvested immediately after euthanization and stored at -80°C before DNA
extraction. The murine colon was excised and opened longitudinally before clearing of colonic
content using a sterile needle and sterile PBS washes. Tissue sections were then excised
using a sterile scalpel pre-treated with DNA Away™ (Fisher Scientific) to destroy

contaminating DNA fragments. For spatial microbiota analyses, the entire colon was laid on a
grid and dissected into individual 0.5 cm x 0.5 cm sections and subsequently frozen at -80°C
before DNA extraction. All steps were carried out in a laminar flow cabinet.

Microbial DNA extraction from murine cecal content and colonic tissue. DNA extraction
from frozen cecal content was carried out using a modified version of the protocol described
by Godon et al [76]. Briefly, cecal samples were homogenized by vortexing and transferred to
autoclaved screw-cap Eppendorf tubes containing 500mg silica beads (0.1mm Carl Roth) and
kept on ice. To this, 600µl of DNA stabilizer (Macherey Nagel), 250µl 4M Guanidine
thiocyanate and 500µl 5% N-laurolyl-sarcosine was added. Samples were incubated under
moderate shaking (700rpm) for 1 hour at 70°C. Lysis was achieved via mechanical disruption
with a FastPrep ®-24 bead beater (MP Biomedicals) using three 40-second cycles at a speed
of 6.5 m/s. 15mg Polyvinylpolypyrrolidone (PVPP; Sigma-Aldrich) was added to homogenate
to remove phenol contamination, before centrifugation at 15,000×g for 3 minutes at 4°C. The
resulting supernatant was recovered, and centrifuged again under the same conditions,
resulting in a clear supernatant containing lysed bacterial cells. To this mixture, 10mg/ml
RNase was added to degrade bacterial RNA and incubated at 37°C for 30 minutes under
constant shaking (700rpm). DNA extraction from colonic tissue was performed using
enzymatic digestion. Where possible samples were extracted using the same kit batch, to limit
differences in inherent kit contamination between different batches [77]. Tissues were placed in
180µl sterile lysis buffer (20Mm Tris/HCL, 2Mm EDTA, 1% Triton-X100; pH 8 supplemented
with 20mg/ml lysozyme) and incubated for 1 hour in a shaking incubator (950rpm) at 37°C.
10mg/ml Proteinase K (Macherey Nagel) was then added, and samples were incubated for 1-
3 hours (until complete lysis of the tissue was obtained) at 56°C under moderate shaking
(950rpm). The resulting genomic DNA was then purified using the NucleoSpin® gDNA clean-
up kit (Macherey Nagel), according to the manufacturer's instructions. The concentration and
purity of extracted genomic DNA was determined using a Nanodrop ND-1000
Spectrophotometer. Samples were stored at -20°C before sequencing.

16S rRNA gene amplicon sequencing. High throughput amplicon sequencing of the 16S
rRNA gene was carried out as previously described [70]. In brief, the V3 and V4 hypervariable
regions were amplified via a 2-step protocol, using the 341f and 785r primer pair (341f 5'-
CCTACGGGNGGCWGCAG-3', 785r 5'-GACTACHVGGGTATCTAATCC-3'). With 10x15
cycles for cecal content samples and 15x15 for mucosal samples. Samples were barcoded
with a double index, according to Kozich *et al.* [78]. The resulting amplicons were purified using
AGENCOURT AMPure XP Beads (Beckman Coulter, Krefeld, Germany), pooled in equimolar
ratios and then sequenced on an Illumina MiSeq system (Illumina), in paired-end mode (2 x
275bp). Raw reads were processed using IMNGS, which wraps the UPARSE/USEARCH

software pipeline [79,80]. Sequences were demultiplexed, trimmed to first base with a quality

score <3 and merged. To remove spurious sequences, a length filter was applied based on

the expected amplicon size of 444bp, removing those smaller than 300 and larger than 600bp.

The resulting paired sequences were then dereplicated and denoised using UNOISE3, to

generate zero-radius OTUs or zOTUs, reflecting true biological sequences [81]. Taxonomy was

assigned using the RDP classifier version 2.11 [82]. Samples with less than 5000 total reads

were excluded from further analysis. Since low biomass samples are susceptible to

contamination which can lead to spurious conclusions, all mucosal samples were further

processed with the R package decontam to remove putative contaminants [83].

Supervised classification of 16S rRNA sequencing. Supervised classification of mucosal

and luminal microbial profiles in non-tumor (fl/fl and tg/wt) and T (tg/tg) mice was performed

using the SIAMCAT R package, utilising Random forest (RF), Ridge regression (RR), LASSO

(L), Elastic-net (E) and L1-penalized regression (LL), which were chosen based on the ease

of interpretation of model output [84,85]. To account for differences in sample size between

mucosal and luminal data, mucosal data was randomly subsampled to match luminal data.

Prior to model training, low prevalent features - defined as those present in less than one-third

of samples - were removed, and the data transformed using a centred log-ratio transform. All

models were trained using five-fold CV, or in case of smaller sample size (<50 total samples),

leave-one-out CV (LOOCV), each with five rounds of resampling.

Multi-omics integration. Integration of metabolomic and microbiota data was performed on

log-transformed, scaled, and centred log-ratio (clr) transformed metabolite and zOTU matrices,

respectively. General associations between the microbiota (luminal and mucosal) and

metabolites were identified using a multi-block sparse partial least squares discriminant

analysis (sPLS-DA) model with seven components and 10-fold CV, as implemented in the

DIABLO framework of the mixOmics R package [44,86]..

Functional potential prediction. To calculate predicted functional profiles based on 16S

rRNA sequencing data, we utilised PICRUSt2 version 2.3 [45]. FASTA files of representative

sequences and minimum sum scaled zOTU tables were used as input for the command:

picrust2_pipeline.py, which runs the full PICRUSt2 pipeline, aligning and placing the

sequences into a reference tree, calculation of 16S copy number, Enzyme Commission (EC)

and KEGG orthologs abundances, adjustment of these by 16S abundance and finally infers

MetaCyc pathways by collapsing EC numbers according to their associated metabolic

pathway. All PICRUSt2 data were generated in a high-performance computing environment,

utilising the Linux-cluster system at the Leibniz Rechenzentrum, Garching.

Intraperitoneal injections of mice with FASN inhibitor C75. The FASN inhibitor C75

(Biomol) was dissolved in RPMI 1640 (Sigma Aldrich) to a final concentration of 1.5 mg/mL,

aliquoted and stored at -20°C until used. C75 or RPMI were injected intraperitoneally (i.p.; 10 ml/kg body weight) biweekly, starting at the age of three weeks until twelve weeks of age (tumor time point). When necessary, mice were given a subcutaneous depot (1:1 mixture (v/v) of 5% glucose and ringer's lactate (B. Braun)) to alleviate potential side-effects of C75. Mice were euthanized with CO_2 at the age of twelve weeks or when abortion criteria were met.

Fecal microbiota transfer into GF mice. Cecal content from C75-treated mice and RPMI-treated controls was instantly suspended at 1:10 weight/volume in filter-sterilized phosphate-buffered saline (PBS)/40% glycerol and stored at –80°C. For gavage, cecal content solutions were centrifuged (3 minutes, 300g, 4°C) to pellet debris, followed by centrifugation (10 minutes, 8000g, 4°C) to collect microbes. This fraction was resuspended in an equal volume of sterile PBS. Each recipient mouse (tg/tg or fl/fl) was gavaged with 100 μL of the bacterial suspension at the age of 4 weeks. Recipient mice were housed in microbiota-specific isolators and euthanized after 12 weeks, or when abortion criteria (as defined in ethical proposals) were fulfilled.

Ex vivo **anaerobic incubations with LCFAs in cecal content of fl/fl mice.** Cecal content was collected from 10 wild-type mice at 5 weeks of age and immediately transferred to an anaerobic chamber (10% H2, 90% N2). To ensure sufficient material, cecal contents of two mice of the same genotype and litter were combined. Cecal content (200-300 mg each) was homogenized in 1X PBS and filtered through a 40 μm filter (Corning, Germany) to remove larger particles. The samples were then transferred to sterile falcon tubes containing a final concentration of 50 μM of the cellular activity marker L-azidohomoalanine (AHA) (Baseclick GmbH, Germany) and between 400 pM and 50 nM of seven long-chain fatty acids (LCFAs), including FAs 20:0, 20:3, 22:0, 22:4, 22:5, 22:6, and 24:1, dissolved in either ethanol or DMF at a final concentration of 1%. Specifically, ethanol and DMF were included as solvent controls for FAs 20:3, 22:4, 22:5, 22:6, 24:1 and 20:0, 22:0, respectively, while 2 mg/mL glucose was the positive control for each experiment. Samples were incubated under anaerobic conditions for 6 hours. After incubation, a fraction of the samples was frozen for DNA extraction, and other aliquots were washed twice with PBS and fixed in 1:1 ethanol for further analysis by fluorescence-activated cell sorting (FACS) [87].

Bioorthogonal non-canonical amino acid tagging (BONCAT) and bacterial cell quantification. To assess the ability of gut microbiota to utilize LCFAs, we applied BONCAT, a fluorescence-based single-cell labeling technique. BONCAT is based on the incorporation of the non-canonical amino acid AHA followed by fluorescent labeling of AHA-containing proteins via azide-alkyne click chemistry [88]. The click chemistry reaction was carried out on microscopy slides, as previously described (Hatzenpichler et al. 2014, doi: 10.1111/1462-2920.12436). Samples were counterstained with 4′,6-diamidino-2-phenylindole (DAPI). Microscopy pictures

were captured using a confocal microscope (Olympus, fluoview, FV10i, Germany) and processed with Fiji software [89] (**Supplementary Figure 1**). Translationally active cells were quantified with flow cytometry using absolute counting beads (CountBrightTM, Invitrogen, ThermoFisher Scientific, Germany) according to the manufacturer's instructions [87].

BONCAT-FACS and DNA extraction. BONCAT combined with FACS was employed to identify bacteria capable of utilizing LCFAs. Samples were processed according to [88]. Briefly, ethanol-fixed samples were first washed with 1X sterile PBS, resuspended in 96% ethanol, and centrifuged. The BONCAT dye solution was then added, and the samples were incubated for 30 minutes in the dark at room temperature [88]. Following incubation, the samples were washed three times with 1X sterile PBS and filtered through a 35 mm nylon mesh using BD tubes (12 x 75 mm, Corning, Germany) prior to sorting. Cy5-positive bacterial cells were sorted using a FACS Melody instrument (BD, Germany), collected in sterile tubes, and stored at -80°C until DNA extraction. The gating strategy is depicted in Extended Data Fig. 8 for fl/fl mice. DNA from both sorted and unsorted bacterial cells was extracted using the QIAamp DNA Mini Kit (Qiagen, Germany), following the manufacturer's instructions and according to [87].

***Desulfovibrio fairfieldensis* and *Desulfovibrio piger* cultivation in LCFA supplemented media.** *Desulfovibrio fairfieldensis* (CSUR P7329, from Collection de Souches de l'Unité des Rickettsies) and *Desulfovibrio piger* (CLA-AA-H201, from Human intestinal bacterial collection) were grown in postgate medium (PG; DSMZ 63). Strains were initially grown in PG medium broth at 37 °C in an anaerobic chamber (5 % CO_2, 5 % H_2, 90% N_2). Once the cultures reached their maximum OD_{600}, they were diluted to 0.05 OD_{600} into fresh PG medium broth supplemented with 10µM of each of the seven long-chain fatty acids (LCFAs), including arachidic acid (20:0), homo-γ-linolenic acid (20:3), docosanoic acid (22:0), docosatetraenoic acid (22:4), docosapentaenoic acid (22:5), docosahexaenoic acid (22:6), and nervonic acid (24:1). PG medium with *D. fairfieldensis* or *D. piger* was used as control. Bacteria suspensions were aliquoted into a sterile 96-well flat-bottom plate (BRANDplates, BRAND, Germany) and incubated for 24 h at 37 °C in a microplate reader (Cerillo, Germany) under anaerobic conditions. Growth data were measured every 30 minutes. Every experiment was conducted five times for *D. fairfieldensis* and tree times for *D. piger*, with three technical replicates. LCFA added to postgate medium in the same amount as to bacteria-inoculated tubes, served as blanks for each condition. The average OD-values of the blanks were subtracted from each technical replicate of the corresponding sample incubated with each LCFA. Of these, the average and standard deviation were calculated and plotted.

TCGA Dataset. For both CRC-related TCGA projects (TCGA-COAD and TCGA-READ), gene expression profiles were downloaded from the firebrowse.org website (RSEM normalized gene expression). Using those tables, transcription factor activities were inferred using the

DoRothEA R package (version 1.10.0) [90] with default parameters (using transcription factor-target interactions of medium- or higher confidence (A-C) and using the viper algorithm for statistical inference). Bacterial profiles derived from whole exome sequencing of the same samples were extracted from the supplemental material of Dohlman *et al.* [49]. For all samples present in both data types (610 samples overlap: 451 COAD, 159 READ), the inferred activity of ATF6 was associated with bacterial presence, using the project information as random effect in a random effect model implemented in the lmerTest R package [91]. Since we observed microsatellite instability (MSI) status to be very strongly associated with both bacterial profiles and ATF6 activity, we additionally included MSI-status as another random effect in a second association model. Bacterial genera were tested for CRC-enrichment with a random effects model for all patients with paired tumor and normal tissue samples and significantly CRC-enriched genera are highlighted in Figure 1 and Extended Data 1. All *p*-values were corrected for multiple hypothesis testing using the Benjamini-Hochberg procedure [92].

Statistical analysis. For data analysis, unless otherwise stated below, statistical analyses were carried out using R (R software foundation, Vienna, Austria) or GraphPad Prism (version 9.00; GraphPad Software, San Diego, CA, USA). For flow cytometry analyses, data processing and analysis were performed with FlowJo™ v10.10.0 software (BD Life Sciences). (Reference: FlowJo™ Software (for Windows) Version v10.10.0. Ashland, OR: Becton, Dickinson and Company; 2023). For ATF6 quantification of IHC staining, the H-score cut-off was determined using the online *Cutoff Finder* Web Application for Biomarker cutoff optimization [93]. For comparing the mean of two groups, unpaired student's T-test or Wilcoxon test was used where appropriate. Differences between multiple groups were tested using ANOVA or Kruskal-Wallis tests. For testing differences in frequency, Fisher's exact test was used. To test for differences in beta diversity, multivariate statistical testing was conducted using Permutational multivariate analysis of variance (PERMANOVA) [94].For hierarchical clustering of metabolite intensities, Ward's method was used [95]. Multiple comparisons were controlled using the method of Benjamini & Hochberg [96].

For semi-targeted metabolomics of CRC patient tissue, statistical analysis and plotting were performed in R 4.4.1 using RStudio/2023.06.1. The statistical significance of molecular feature in T versus T adjacent was assessed with paired t-test (t.test function in R), and p-values were FDR-corrected for multiple hypotheses testing using the Benjamini-Hochberg procedure (p.adjust function in R with 'fdr' parameter). Significant metabolites were selected when their intensity had a corrected p-value < 0.05.

For 16S rRNA gene amplicon sequencing, the decontam package utilises the relationship between pooled DNA concentration prior to sequencing and the prevalence and abundance of taxa in negative controls to identify putative contaminants. Identification of contaminants was

performed with the isContaminant function using default parameters. Downstream analyses were performed using Rhea and phyloseq [97,98]. Briefly, zOTU tables were normalized using minimum sum scaling or relative abundance. Alpha diversity was measured using Shannon Effective and diversity. To assess differences between groups, beta diversity was calculated based on generalized UniFrac distance (GUniFrac) [99]. Mean GUniFrac dissimilarity for a given sample was calculated as the dissimilarity to the respective control mean. Differentially abundant taxa were identified using the Linear Discriminant analysis Effect size (LEfSe) algorithm, with a Linear discriminant analysis threshold (LDA) threshold of 3.0, to limit false positives [100,101].

For BONCAT-FACS experiments, statistical analysis was performed using R statistical software (https:// www.r-project.org/). Raw sequencing data were pre-processed using the web platform IMNGS2 (https://www.imngs2.org/). Contaminants originating from PCR and kit reagents were detected and removed using the R package decontam [102]. zOTU counts were subsampled to a number of reads smaller than the smallest library (1000 reads). Beta diversity was calculated using generalized UniFrac distances [103] and visualized using multidimensional scaling (MDS) [97]. Permutational multivariate analysis of variance (perPERMANOVA) was performed to determine the significance of samples grouping. Alpha-diversity was evaluated based on taxa richness [97] and group comparison was calculated using Wilcoxon test. Significant enriched and depleted zOTUs between sorted and unsorted fractions were performed using DEseq2 [104].

Significant zOTUs sequences were identified using the 16S-based ID tool of EzBioCloud (https://www.microbiologyresearch.org/content/journal/ijsem/10.1099/ijsem.0.001755).

The variation in microbial communities across the seven LCFAs was further analyzed using the UpSetR package in R, which employs a matrix-based layout to represent set intersections and their sizes [105].

For quantifications of the percentage of translationally active cells, ANOVA and Tukey's test for multiple comparisons were applied.

For growth curves, two-way ANOVA was used to compare *D. fairfieldensis* or *D. piger* stimulated with each LCFA with the control without LCFAs.Variables were expressed as mean ± standard deviation (SD), a probability value (p-value) less than 0.05 was considered statistically significant and Bonferroni method was used for multiple comparison. For single time point comparison one-way ANOVA was used and Dunnet's test for multiple comparisons.

Unless otherwise specified all data are presented as the mean ± standard deviation, *p*-values < 0.05 are considered statistically significant ($p<0.05=*$, $p<0.01=**$, $p<0.01=***$, $p<0.001=****$).

 Figure legends

 Fig. 1: High ATF6 expression defines a subset of CRC patients. a, QuPath quantified ATF6
H-score of nuclear (NUC) and cytoplasmic (CYT) ATF6 expression in immunohistochemically
stained CRC patient tissue samples (cohort 1, n=959). Significance was calculated using the
Student's two-tailed unpaired T-Test ($p<0.0001$), indicated is the mean. Representative
images of ATF6-high and ATF6-low staining in the tumor centre (TC) and the invasive front
(IF). Percentage shows subpopulation of CRC patients in each ATF6-stained category (ATF6-
low, light red; ATF6-high, dark red). Scale bars: 300µm overview, 50µm zoom-in. **b,** Confusion
matrix depicting the classification model of ATF6 H-scores quantified using QuPath (true
labels) versus ATF6 H-scores predicted by the classification algorithm (predicted labels) for
ATF6-high and ATF6-low CRC patients. **c,** Pie chart showing percentage of ATF6-high and
ATF6-low CRC patients identified in CRC cohort 2 (n=50) using the classification model. **d,**
Regression model of ATF6 H-scores quantified using QuPath (true values) versus ATF6 H-
scores predicted by the regression algorithm (predicted values) for nuclear (NUC) and
cytoplasmic (CYT) ATF6 expression. **e,** Pie chart showing percentage of ATF6-high and ATF6-
low CRC patients identified in CRC cohort 2 (n=50) using the regression model. **f,** QuPath
quantified ATF6 H-score of nuclear (NUC) and cytoplasmic (CYT) ATF6 expression in
immunohistochemically stained CRC patient tissue samples (cohort 3, LOCRC, n=256).
Significance was calculated using the Student's two-tailed unpaired T-Test ($p<0.0001$),
indicated is the mean. Representative images of ATF6-high and ATF6-low staining.
Percentage shows subpopulation of CRC patients in each ATF6-stained category. Scale bars:
300µm overview, 50 µm zoom-in. **g,** QuPath quantified ATF6 H-score of nuclear (NUC) and
cytoplasmic (CYT) ATF6 expression in immunohistochemically stained CRC patient tissue
samples (cohort 3, EOCRC, n=55). Significance was calculated using the Student's two-tailed
unpaired T-Test, indicated is the mean. Representative images of ATF6-high and ATF6-low
staining. Percentage shows subpopulation of CRC patients in each ATF6-stained category.
Scale bars: 300µm overview, 50µm zoom-in. A significant p-value <0.05 is represented as an
asterisk: $*p < 0.05$, $**p < 0.01$. $***p < 0.001$, $****p < 0.0001$.

Fig. 2: ATF6 alters colonic fatty acid metabolism in the presence of bacteria a, Survival
curve showing percent survival of fl/fl, tg/wt, tg/tg specific pathogen-free (SPF) and tg/tg germ-
free (GF) mice between 0-25 weeks of age (n per genotype = 6-12). **b,** Tumor incidence
(percentage) of SPF fl/fl, tg/wt and tg/tg mice and GF tg/tg mice at the pre-tumor (5wk), tumor
(12wk) and late-tumor (15+wk) time points. Grey represents no tumor, red represents tumor.
Number of mice is stated in each column. **c,** Schematic of non-tumor (NT) and tumor (T)
phenotypes associated with GF and SPF fl/fl and tg/tg mice, highlighting the genotypes and

colonization status used for the RNA sequencing analyses (orange box and arrow) (n=6). **d,**

Volcano plot of detected genes in fl/fl versus tg/tg nATF6[IEC] mice showing the number and

percentage of genes that are unchanged (No Change), upregulated in tg/tg mice (UP) or

downregulated in tg/tg mice (DOWN). Threshold: -0.5 > log2FC >0.5, *p* adj. <0.05. **e,** Lollipop

graph of the top and bottom regulated KEGG pathways in tg/tg versus fl/fl nATF6[IEC] SPF mice

(*p* adj. <= 0.25, −1 =>NES>= 1, with a minimum of 50% of genes detected in the pathway).

Metabolic pathways are highlighted in blue. NES, normalized enrichment score. **f,** Alluvial plot

depicting the flows used to define SiRCle clusters in a comparison between SPF and GF mice.

Each data type (SPF and GF) has been labelled as a column, with one of three states (UP, No

Change, DOWN) defined for each column based on the results for differential analysis between

fl/fl versus tg/tg in that data type. Shown are the flows between each data type and state, which

define the regulatory clusters (third column). The number and percentage of genes in the flow

are depicted next to each regulatory cluster. **g,** Lollipop graph of the FA metabolism related

genes from the SPF UP cluster (blue) shown for GF (grey) and SPF (red) mice. **h,**

Representative images of low H-score and high H-score FASN immunohistochemically stained

in CRC patient tissue samples (cohort 1, n=181). Scale bars: 300µm overview, 50µm zoom-

in. **i,** QuPath quantified FASN H-score with patients grouped into ATF6-low and ATF6-high

(according to QuPath quantified ATF6 H-score). Significance was calculated using the

Student's two-tailed unpaired T-Test (*p*=0.0105), indicated is the mean. A significant *p*-value

<0.05 is represented as an asterisk: **p* < 0.05, ***p* < 0.01. ****p* < 0.001, *****p* < 0.0001.

1105

Fig. 3: ATF6 drives FA elongation and LCFAs accumulate in mice and CRC

patients. a, Schematic depicting metabolomic analysis strategy in nATF6[IEC] mice and human

CRC patients (cohort 4). Created with BioRender.com. **b,** Boxplots comparison of log-

transformed fatty acid intensities in caecal content, comparing fl/fl to tg/tg mice, stratified by

timepoint. Whiskers extend to the minimum and maximum values. Only fatty acid metabolites

which were significant in mouse data and overlapped with CRC patient data are shown.

Statistical significance was calculated using pairwise T-tests and adjusted for multiple

comparisons using the Benjamini-Hochberg procedure. **c,** Boxplot comparison of fatty acid

intensities in tumor tissue and tumor-adjacent tissue of samples from cohort 4. The plot

displays the intensity differences for nine fatty acids that shown significant abundance in tumor

samples (left to right): hydroxyeicosadenoic acid (FA 20:2), hydroxy-dihomo-linolenic acid (FA

20:3), hydroxy-arachidonic acid (FA 20:4), dihomo-linoleic acid (FA 20:3), docosatetranoic

acid (FA 22:4), docosapentanoic acid (FA 22:5), docosahexanoic acid (FA 22:6),

tetracosapentanoic acid (FA 24:5) and herring acid (FA 24:6). Whiskers extend to the minimum

and maximum values, excluding outliers, which are shown as individual points. Paired T-Test

was used to calculate *p*-values (and FDR corrected). Only fatty acids with p-values (FDR

corrected) <0.05 are shown. **d,** Percentage of saturated fatty acids (SAFA) in NT and T tissue after quantification of total FA using GC-MS. Statistical analysis was perfomed using using an unpaired Wilcoxon test (p=0.0043). **e,** Scheme showing the treatment of intestinal organoids from nATF6 Vil-CreERT2 (fl/fl^{ERT2} and tg/tg^{ERT2}) mice cultured *ex vivo* (n=11-12). Organoids were passaged at D0, ATF6 expression in all wells was induced at D3 with 500nM (Z)-4-hydroxytamoxifen (4-OHT) and treated with D3-FA 20:0 at D4, before harvesting at D5 (see also Supplementary Figure 1). Created with BioRender.com. **f,** Percentage of metabolized D$_3$-FA20:0 to D$_3$-FA:22:0 in organoids of tg/tg^{ERT2}, compared to fl/fl^{ERT2} controls using GC-MS. Significance was calculated using the Student's two-tailed unpaired T-Test (p=0.0184). A significant p-value <0.05 is represented as an asterisk: *$p < 0.05$, **$p < 0.01$. ***$p < 0.001$, ****$p < 0.0001$.

Fig. 4: ATF6-related lipid profiles are causally linked to tumor-associated microbiota. a, Mean generalized UniFrac distance relative to control across fl/fl, tg/wt and tg/tg samples, at pre-tumor (5wk) and tumor-onset (12wk) in luminal (left) and mucosal communities. *P* values were calculated using a pairwise t-test and adjusted for multiple comparisons using the Benjamini-Hochberg procedure. Whiskers extend to the minimum and maximum values (n=6-7 mice per group). **b,** Loadings plots of omic features selected by the sPLS-DA model, discriminating tg/tg mice from other genotypes. Luminal zOTUs are shown in yellow, mucosal zOTUs in red and metabolites in blue. Features are sorted by importance (n = 17-18 per genotype). **c,** Spatial maps of Log10 transformed predicted relative abundance of oleate hydratase (ohyA) positive ASVs along the entire colon in fl/fl and tg/tg mice (n=6 mice per group). **i,** Predicted relative abundance of ohyA in fl/fl and tg/tg mice. **d,** Relative abundance of oleate hydratase (ohyA) in tg/tg and fl/fl mice p=4.7e-10). **e,** Spatial maps of Log10 transformed predicted relative abundance of FA efflux pump (farE) positive ASVs along the entire colon in fl/fl and tg/tg mice (n=6 mice per group). **f,** Predicted relative abundance of fatty acid efflux pump, farE in tg/tg and fl/fl mice (p=<2.26e-16). **g,** Schematic showing the experimental design for in vivo C75 FASN inhibitor intervention and transfer experiment. SPF mice were intraperitoneally (i.p.) injected either with the RPMI control or the C57 FASN inhibitor bi-weekly from the age of 3 weeks. After an intervention period of 9 weeks, mice were sacrificed at 12 weeks of age. Arrows indicate i.p. injections. **h,** Tumor incidence (percentage) of GF fl/fl and tg/tg mice after gavage with donor material from either RPMI-treated (controls) or C75 FASN inhibitor-treated mice. **i,** Tumor number of GF fl/fl and tg/tg mice after gavage with donor material from either RPMI-treated (controls) or C75 FASN inhibitor-treated mice. Points are colored by donor. Statistical significance was calculated using an ordinary one-way ANOVA with Tukey`s multiple comparison test (p=0.0205). A significant p-value <0.05 is represented as an asterisk: *$p < 0.05$, **$p < 0.01$. ***$p < 0.001$, ****$p < 0.0001$.

Fig. 5: LCFAs selectively activate the growth of tumor-related bacteria. a, Schematic overview of bioorthogonal non-canonical amino acid tagging (BONCAT). Mouse cecal content was incubated in anaerobic conditions and stimulated with seven long-chain fatty acids (LCFAs), including arachidic acid (20:0), homo-γ-linolenic acid (20:3), docosanoic acid (22:0), docosatetraenoic acid (22:4), docosapentaenoic acid (22:5), docosahexaenoic acid (22:6), and nervonic acid (24:1) and along with the cellular activity marker L-azidohomoalanine (AHA). Translationally active bacterial cells were labeled by azide-alkyne click chemistry, sorted with FACS and sequenced by 16S rRNA gene amplicon sequencing. Created with BioRender.com. **b,** Representative confocal microscopic images of a mouse cecal content stimulated with nervonic acid (24:1). Pink: active cells (BONCAT-Cy5); blue: all cells (DAPI); merge image. Scale bar is 10 µm. **c,** Percentage of translationally active bacteria after LCFA amendment. LCFA are compared to their respective solvent control: 20:0 and 22:0 were compared to DMF (DMF), 20:3, 22:4, 22:5, 22:6 and 24:1 were compared to ethanol (Etoh). P values were calculated by ANOVA and Tukey's test for multiple comparisons (mean ± sd: 5.8 ± 1.6% for LCFAs in total; 0.9 ± 0.51 for the controls; ANOVA, $p < 0.0001$ for all comparisons, n=80). Error bars represent standard deviation of the mean. **d,** Heatmap displayed log2-fold changes of significantly enriched or depleted zOTUs in the translationally active fraction compared to the unsorted group for each LCFA as calculated with the Wald test ($p < 0.05$, n=160). All the zOTUs (51) listed in the heatmap are significantly enriched or depleted. zOTUs were identified using the 16S-based ID tool of EzBioCloud. Numbers in parenthesis correspond to the barplots of Extended Data 7d where each barplot illustrate the shared and unique zOTUS between all LCFAs. **e,** Relative abundance of a *Desulfovibrio fairfieldensis* zOTU, in luminal (left) and mucosal (right) communities comparing fl/fl, tg/wt and tg/tg mice at the 5-week timepoint. Whiskers extend to the minimum and maximum values. Statistical significance was calculated using pairwise t-tests and adjusted for multiple comparisons using the Benjamini-Hochberg procedure . **f,** Growth curve of *Desulfovibrio fairfieldensis* in postgate medium, supplemented with different LCFAs. Each curve represents averaged values of n = 5 biological replicates in three technical replicates. Long-chain fatty acids (20:0, 22:0, 20:3, 22:4, 22:5, 22:6 and 24:1) were added to postgate medium to a concentration of 10µM each. The control (gray) contains only postgate medium with bacteria. All LCFA supplementations show significant increase of growth compared to the control (two-way ANOVA, Bonferroni test for multiple comparisons, $p < 0,0001$ for all comparisons, n=735). Error bars shown as dotted line represent standard deviation of the mean. **g,** Plot of ATF6 activity in CRC patients (TCGA Database) when bacterial genus is present (x-axis) and the significance (y-axis). Marked in orange/red are those genera significantly associated with ATF6 activity, with red dots representing classical CRC-associated genera. **h,** Plot of ATF6 activity in CRC patients (TCGA

1196   Database) when bacteria are absent (grey) or present (red) for the CRC-associated genera

1197   marked red in h (*p* values*:* Campylobacter = 0.00123, Desulfovibrio = 0.00507, Porphyromonas

1198   = 0.00287, Fusobacterium = 0.00507, Selenomonas = 0.00507). A significant *p*-value <0.05

1199   is represented as an asterisk: *p < 0.05, **p < 0.01. ***p < 0.001, ****p < 0.0001.

1200

Extended Data Fig. 1: ATF6 and GRP78 analyses in human CRC patients. a,

Representative images of ATF6 antibody staining in heart tissue (top) and the isotype control

in CRC patient tissue (bottom). Scale bars: 300µm overview, 50µm zoom-in. **b,** Nuclear (NUC)

cutoff for the ATF6 H-score as determined using the online *Cutoff Finder* Web Application for

Biomarker cutoff optimization, dividing patients into ATF6-high and ATF6-low. **c,** Cytoplasmic

(CYT) cutoff for the ATF6 H-score as determined using the online *Cutoff Finder* Web

Application for Biomarker cutoff optimization, dividing patients into ATF6-high and ATF6-low.

d, Donut chart depicting the percentage of patients with all observed combinations of nuclear

and cytoplasmic staining (NUC L/CYT L = low nuclear and low cytoplasmic staining, NUC

L/CYT H = low nuclear and high cytoplasmic staining, NUC H/CYT L = high nuclear and low

cytoplasmic staining, NUC H/CYT H = high nuclear and high cytoplasmic staining). **e,** QuPath

quantified GRP78 H-score of ATF6-low and ATF6 high patients in immunohistochemically

stained CRC patient tissue samples (cohort 1, n=959). Significance was calculated using the

Student's two-tailed unpaired T-Test. **f,** Representative images of GRP78 antibody staining in

CRC patient tissue samples, stratified according to H-score and isotype control. Scale bars:

300µm overview, 50µm zoom-in. **g,** Plot showing the classification model training accuracy,

displaying the mean training and validation accuracy and their standard deviation. **h,** Plot

showing the classification model training loss, displaying the mean training and validation loss

and their standard deviation. A significant *p*-value <0.05 is represented as an asterisk: *p <

0.05, **p < 0.01. ***p < 0.001, ****p < 0.0001.

1221

Extended Data Fig. 2: Gene expression and pathway analysis in response to ATF6

activation under GF and SPF conditions. a, Volcano plot of detected genes in fl/fl versus

tg/tg nATF6[IEC] mice showing the number and percentage of genes that are unchanged (No

Change), upregulated in tg/tg mice (UP) or downregulated in tg/tg mice (DOWN). Threshold: -

0.5 > log2FC >0.5, *p* adj. <0.05. **b,** Lollipop graph of the top regulated KEGG pathways in tg/tg

versus fl/fl nATF6[IEC] GF mice (*p* adj. <= 0.25, −1 =>NES>= 1, with a minimum of 50% of genes

detected in the pathway). Metabolic pathways are highlighted in blue. NES, normalized

enrichment score. **c,** Lollipop graph of log2 FC of UPR-related genes. ATF6-pathway related

genes are highlighted in red. Colour represents enrichment (yellow = Acute SPF, blue =

Chronic SPF, green = Chronic GF) and size, -log10 adjusted p-value. **d,** Volcano Plot of genes

in the regulatory cluster CORE-ATF6 UP. Upset plot representing the number of genes detected for the respective transcription factors driving the cluster. **e,** Volcano Plot of genes in the regulatory cluster CORE-ATF6 DOWN. Upset plot representing the number of genes detected in the cluster-associated KEGG pathways (threshold: >5% of pathway genes detected, p adj. < 0.2). Upset plot representing the number of genes detected and the respective transcription factors driving the cluster. **f,** Volcano Plot of genes in the regulatory cluster SPF DOWN. Upset plot representing the number of genes detected in the cluster-associated KEGG pathways (threshold: >5% of pathway genes detected, p adj. < 0.2). **g,** Volcano Plot of genes in the regulatory cluster SPF UP. Genes related to fatty acid metabolism are highlighted in blue. Upset plot representing the number of genes detected in the cluster-associated KEGG pathways (threshold: >5% of pathway genes detected, p adj. < 0.2). Upset plot representing the number of genes detected and the respective transcription factors driving the cluster. **h,** Volcano plot of detected genes in fl/fl versus tg/tg following acute activation of ATF6 in nATF6 Vil-Cre[ERT2] SPF mice showing the number and percentage of genes that are unchanged (No Change), upregulated in tg/tg mice (UP) or downregulated in tg/tg mice (DOWN). Threshold: -0.5 > log2FC >0.5, *p* adj. <0.05. Genes related to fatty acid metabolism are highlighted in blue. A significant *p*-value <0.05 is represented as an asterisk: *p < 0.05, **p < 0.01. ***p < 0.001, ****p < 0.0001.

Extended Data Fig. 3: ATF6 and FASN co-occurrence in CRC patient cohorts. a, Table showing the analysis of *ATF6* amplifications/gain and *FASN* amplifications/gain in the pan-cancer and TCGA Datasets, indicating the log2 odds ratio (OR), *p* value and observed association/tendency. **b,** Kaplan-Meier analysis of disease-free survival shown for ATF6-low (blue) and ATF6-high (red) at different months after surgery in CRC cohort 1. **c,** Kaplan-Meier analysis of overall survival shown for ATF6-low (green) and ATF6-high (blue) at different months after surgery in CRC cohort 3 EOCRC patients. **d,** Kaplan-Meier analysis of overall survival shown for ATF6-low (green) and ATF6-high (blue) at different months after surgery in CRC cohort 3 LOCRC patients. **e,** Kaplan-Meier analysis of disease-free survival of patients in the Pan-Cancer Atlas database (n=4.129/10.967 cases including approx. 600 CRC cases) with patients grouped into FASN+ATF6 unaltered, ATF6-only amp/gain or ATF6+FASN amp/gain. **f,** Kaplan-Meier analysis of disease-free survival of patients in The Cancer Genome Atlas database (TCGA; n=494/640 CRC cases) with patients grouped into FASN+ATF6 unaltered, ATF6-only amp/gain or ATF6+FASN amp/gain. **g,** Kaplan-Meier analysis of overall survival of patients in the Pan-Cancer Atlas database (n=8.345/10.967 cases including approx. 600 CRC cases) with patients grouped into FASN+ATF6 unaltered, ATF6-only amp/gain or ATF6+FASN amp/gain. **f,** Kaplan-Meier analysis of overall survival of patients in The Cancer

1268   Genome Atlas database (TCGA; n=543/640 CRC cases) with patients grouped into
1269   FASN+ATF6 unaltered, ATF6-only amp/gain or ATF6+FASN amp/gain.

1270

Extended Data Fig. 4: Metabolite profiles in nATF6^{IEC} mice and CRC patients. a, PCA of

metabolite profiles comparing fl/fl (circles), tg/wt (squares) and tg/tg (triangles) mice at 5wk,

12wk and 20wk. Grey indicates non-tumor (NT) and red tumor (T). **b,** Volcano plot of

differentially enriched metabolites between fl/fl and tg/tg mice across all time points (Log2 fold

change threshold = 1.0, FDR threshold = 0.1)**. c,** PCA of luminal lipid profiles between tumor-

susceptible/tumor (T) and non-tumor (NT) mice at 5wk, 12wk and 20wk. **d,** Heatmap of scaled

differentially enriched metabolites, sorted by genotype and age. Rows are clustered using

Ward's method. **e,** Boxplot comparison of fatty acid intensities in tumor and tumor-adjacent

tissue stratified for EOCRC and LOCRC. Displayed are intensity differences for two fatty acids

that show significant abundance in tumor tissue for EOCRC in addition to EOCRC. Only fatty

acids with p-values (FDR corrected) <0.05 are shown. Statistical significance was determined

using a paired T-test. **f,** Mirror plots of MS/MS spectra for the annotation of fatty acids at 10,

20 and 40eV. The plots compare the experimental MS/MS spectra (top, blue) with the standard

compound spectra (bottom, red) for selected fatty acids. Correct annotation occurs when

retention time, high-resolution mass, precursor and diagnostic fragment ions are similar

between the analyte and the CRC-sample. **g,** Percentage of saturated fatty acids (SAFA) in

NT (n=6) and T (n=5) tissue after quantification of total FA using GC-MS. **h,** Percentage of

SAFA plotted against tumor number after quantification of total FA using GC-MS. **i,**

Representative images of intestinal organoids on day 1 (d1) and day 7 (d7) after culture, for

the fl/fl and tg/tg genotype. Scale bars: 50μm. **j,** Quantification of organoid size as the sum of

two dimensions in μm at d1 and d7 for the fl/fl and tg/tg genotype. **k**, Quantification of the

number of crypts per cyst at d7 for the fl/fl and the tg/tg genotype. A significant p-value <0.05

is represented as an asterisk: *p < 0.05, **p < 0.01. ***p < 0.001, ****p < 0.0001.

1294

Extended Data Fig. 5: Luminal and mucosal microbiota profiles in nATF6^{IEC} mice. a, Non-

metric multidimensional scaling (NMDS) plot of Beta diversity profiles based on generalized

UniFrac distance comparing genotypes (fl/fl, tg/wt and tg/tg) at each timepoint (5, 12 and 20-

weeks) in luminal (n=54) and **b,** mucosal data (n=62). **c,** ROC curves comparing classification

accuracy of Random Forest (RF), Ridge regression (RR)), LASSO (L), Elastic-net (E) and L1-

penalized regression (LL) models built on luminal and mucosal data, respectively. Mucosal

data was randomly subsampled to match luminal sample size (total n=62, random subsample

n=54). Models were trained using repeated 5-fold cross-validation (CV). **d,** Spearman's rank

correlation between Shannon Effective and tumor number in mucosal and **e,** luminal data**.**

1304

Extended Data Fig. 6: Multi-omics data integration of metabolite and microbial profiles. **a,** Loadings plots of omic features selected by the sPLS-DA model, discriminating T from NT mice. Luminal zOTUs are shown in yellow, mucosal zOTUs in red and metabolites in blue. Features are sorted by importance (n=52). **b,** Relevance network of highly correlated (Absolute correlation >0.65) metabolites and mucosal zOTUs for the genotype sPLS-DA model (n=52). **c,** Relevance network of highly correlated (Absolute correlation >0.65) metabolites and luminal zOTUs for the genotype sPLS-DA model (n=52).

Extended Data Fig. 7: Bacterial lipid stress response and FASN inhibition. a, Heatmap of bacterial FA-response genes predicted by PICRUSt2 for fl/fl controls and non-tumor (NT), tumor-adjacent (TA) and tumor (T) mucosal phenotypes of tg/tg mice. Spatial sites along the colon and biological replicates (mouse) are numerically labelled underneath. Only tumor-susceptible sites (sites 1-10) are shown (n=6 mice per group and 15-17 colonic sites per mouse). **b,** Ratio of ohyA positive to ohyA negative taxa across various mucosal phenotypes in fl/fl and tg/tg mice . **c,** Ratio of farE positive to farE negative taxa across various mucosal phenotypes in fl/fl and tg/tg mice. **d,** Tumor incidence (percentage) of SPF fl/fl and tg/tg mice after termination of the intervention with either RPMI (controls) or C75 FASN inhibitor. Grey represents no tumor, red represents tumor. Each bar shows the number of mice for each genotype. **e,** Representative macroscopic images of the colon of the tg/tg genotype on RPMI or C75. Marked with circles and asterisks are tumors identified in the control group. **f,** Tumor number in SPF fl/fl and tg/tg mice after invervention with RPMI or C75. **g,** Tumor volume in SPF fl/fl and tg/tg mice after intervention with RPMI or C75. **h,** Survival curve showing percent survival of fl/fl and tg/tg GF mice colonised with either RPMI-treated or C75-treated microbiota between 0-12 weeks of age. **i,** Tumor volume (mm^3) in GF fl/fl and tg/tg mice after gavage with donor material from either RPMI or C75 treated mice. Points are colored by donor. Statistical significance was calculated using pairwise t-tests and adjusted for multiple comparisons using the Benjamini-Hochberg procedure (p=0.0317). A significant *p*-value <0.05 is represented as an asterisk: *$p < 0.05$, **$p < 0.01$. ***$p < 0.001$, ****$p < 0.0001$.

Extended Data Fig. 8. Bacterial responses to LCFAs. a, Alpha diversity richness showing significant difference between sorted and unsorted fraction (Wilcoxon test: p<0.0001, n=160). Boxplot: boxplot medians (center lines), interquartile ranges (box ranges), whisker ranges. **b,** Beta-diversity MDS plot shows clustering of samples based on sorted and unsorted fractions (perMANOVA, p = 0.0015, n=160). Unsorted fraction= blue, sorted fraction= green. **c,** Venn diagram intersections shows 10 zOTUS shared between all LCFAs. 15 are shared between the saturated LCFAs (20:0 and 22:0), 9 are unique for 22:6 and 5 are shared between 22:6

and 20:3. The total size of each set is represented on the right barplot. The bottom plot represents all possible intersections, while their occurrence are shown in the top barplot. Each row corresponds to a set, and each column corresponds to one segment in a Venn diagram. A set is part of the intersection only when its cell is filled. Numbers in parenthesis on the top of each barplot represents which zOTUs belongs to each intersection. The corresponding zOTUS groups are highlighted in green in Figure 5d. **d,** Beta-diversity MDS plot displayed the active fraction across the different LCFAs. No significant difference was found between LCFAs (perMANOVA, $p>0.05$, n=70). **e,** Relative abundance of zOTUs classified as Parasutterella and Lachnospiraceae, in mucosal (top), mucosal (middle) and luminal (bottom) communities respectively, comparing fl/fl, tg/wt and tg/tg mice at the 5-week timepoint. Whiskers extend to the minimum and maximum values. **f,** *Desulfovibrio fairfieldensis* growth in postgate medium, supplemented with different LCFAs, at 18 hours. Barplots represent averaged values of n = 5 biological replicates in three technical replicates. *P*-values were calculated by one-way ANOVA and Dunnet's test for multiple comparisons, n=120. Error bars represent standard deviation of the mean. **g,** Growth curve of *Desulfovibrio piger* in postgate medium, supplemented with different LCFAs. Each curve represents averaged values of n = 3 biological replicates in three technical replicates. Long-chain fatty acids (20:0, 22:0, 20:3, 22:4, 22:5, 22:6 and 24:1) were added to postgate medium to a concentration of 10µM each. The control (gray) contains only postgate medium with bacteria. All LCFA supplementations show significant increase of growth compared to the control (two-way ANOVA, Bonferroni test for multiple comparisons, $p < 0,0001$ for all comparisons, n=441). Error bars shown as dotted line represent standard deviation of the mean. **h,** Association of ATF6 activity with CRC-enriched genera after correcting for MSI status, and correlation of the CRC-enriched genera (highlighted in red, 1-5) with ATF6 activity for MSS and MSI in the absence and presence of the respective bacteria. **i,** Plot of ATF6 activity in CRC patients (TCGA Database) when bacterial genus is present (x-axis) and the significance (y-axis). Marked in orange/red are those genera significantly associated with ATF6 activity, with red dots representing classical CRC-associated genera. A significant *p*-value <0.05 is represented as an asterisk: *$p < 0.05$, **$p < 0.01$. ***$p < 0.001$, ****$p < 0.0001$.

References

1   Braakman, I. & Bulleid, N. J. Protein folding and modification in the mammalian endoplasmic reticulum. *Annu Rev Biochem* **80**, 71-99, doi:10.1146/annurev-biochem-062209-093836 (2011).

2   Jacquemyn, J., Cascalho, A. & Goodchild, R. E. The ins and outs of endoplasmic reticulum-controlled lipid biosynthesis. *EMBO Rep* **18**, 1905-1921, doi:10.15252/embr.201643426 (2017).

3   Rutkowski, D. T. & Kaufman, R. J. A trip to the ER: coping with stress. *Trends Cell Biol* **14**, 20-28 (2004).

1379	4	Shen, J., Chen, X., Hendershot, L. & Prywes, R. ER stress regulation of ATF6 localization by
1380		dissociation of BiP/GRP78 binding and unmasking of Golgi localization signals. *Dev Cell* **3**, 99-
1381		111 (2002).
1382	5	Yoshida, H., Matsui, T., Yamamoto, A., Okada, T. & Mori, K. XBP1 mRNA is induced by ATF6 and
1383		spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* **107**,
1384		881-891 (2001).
1385	6	Kaser, A. *et al.* XBP1 links ER stress to intestinal inflammation and confers genetic risk for
1386		human inflammatory bowel disease. *Cell* **134**, 743-756, doi:10.1016/j.cell.2008.07.021 (2008).
1387	7	Bertolotti, A. *et al.* Increased sensitivity to dextran sodium sulfate colitis in IRE1beta-deficient
1388		mice. *J Clin Invest* **107**, 585-593 (2001).
1389	8	Shkoda, A. *et al.* Interleukin-10 blocked endoplasmic reticulum stress in intestinal epithelial
1390		cells: impact on chronic inflammation. *Gastroenterology* **132**, 190-207,
1391		doi:10.1053/j.gastro.2006.10.030 (2007).
1392	9	Stengel, S. T. *et al.* Activating Transcription Factor 6 Mediates Inflammatory Signals in Intestinal
1393		Epithelial Cells Upon Endoplasmic Reticulum Stress. *Gastroenterology* **159**, 1357-1374.e1310,
1394		doi:10.1053/j.gastro.2020.06.088 (2020).
1395	10	Hanaoka, M. *et al.* Expression of ATF6 as a marker of pre-cancerous atypical change in
1396		ulcerative colitis-associated colorectal cancer: a potential role in the management of dysplasia.
1397		*J Gastroenterol*, doi:10.1007/s00535-017-1387-1 (2017).
1398	11	Coleman, O. I. *et al.* Activated ATF6 Induces Intestinal Dysbiosis and Innate Immune Response
1399		to Promote Colorectal Tumorigenesis. *Gastroenterology* **155**, 1539-1552 e1512,
1400		doi:10.1053/j.gastro.2018.07.028 (2018).
1401	12	Waldschmitt, N. *et al.* C/EBP homologous protein inhibits tissue repair in response to gut injury
1402		and is inversely regulated with chronic inflammation. *Mucosal Immunol* **7**, 1452-1466,
1403		doi:10.1038/mi.2014.34 (2014).
1404	13	Lee, A. S. Glucose-regulated proteins in cancer: molecular mechanisms and therapeutic
1405		potential. *Nat Rev Cancer* **14**, 263-276, doi:10.1038/nrc3701 (2014).
1406	14	Lin, Y. H. *et al.* Multiple gene expression classifiers from different array platforms predict poor
1407		prognosis of colorectal cancer. *Clin Cancer Res* **13**, 498-507, doi:10.1158/1078-0432.Ccr-05-
1408		2734 (2007).
1409	15	Schewe, D. M. & Aguirre-Ghiso, J. A. ATF6alpha-Rheb-mTOR signaling promotes survival of
1410		dormant tumor cells in vivo. *Proc Natl Acad Sci U S A* **105**, 10519-10524,
1411		doi:10.1073/pnas.0800939105 (2008).
1412	16	Wu, X. *et al.* A missense polymorphism in ATF6 gene is associated with susceptibility to
1413		hepatocellular carcinoma probably by altering ATF6 level. *Int J Cancer* **135**, 61-68,
1414		doi:10.1002/ijc.28649 (2014).
1415	17	Jesinghaus, M. *et al.* Morphology Matters: A Critical Reappraisal of the Clinical Relevance of
1416		Morphologic Criteria From the 2019 WHO Classification in a Large Colorectal Cancer Cohort
1417		Comprising 1004 Cases. *Am J Surg Pathol* **45**, 969-978, doi:10.1097/PAS.0000000000001692
1418		(2021).
1419	18	Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci Rep*
1420		**7**, 16878, doi:10.1038/s41598-017-17204-5 (2017).
1421	19	Mora, A., Schmidt, C., Balderson, B., Frezza, C. & Bodén, M. SiRCle (Signature Regulatory
1422		Clustering) model integration reveals mechanisms of phenotype regulation in renal cancer.
1423		*bioRxiv*, 2022.2007.2002.498058, doi:10.1101/2022.07.02.498058 (2022).
1424	20	Zhu, L. *et al.* Germline Variants of CYBA and TRPM4 Predispose to Familial Colorectal Cancer.
1425		*Cancers (Basel)* **14**, doi:10.3390/cancers14030670 (2022).
1426	21	Lian, J., Nelson, R. & Lehner, R. Carboxylesterases in lipid metabolism: from mouse to human.
1427		*Protein Cell* **9**, 178-195, doi:10.1007/s13238-017-0437-z (2018).
1428	22	Thesseling, F. A. *et al.* Novel insights into oxidation of fatty acids and fatty alcohols by
1429		cytochrome P450 monooxygenase CYP4B1. *Arch Biochem Biophys* **679**, 108216,
1430		doi:10.1016/j.abb.2019.108216 (2020).

23 Zhao, M. *et al.* Cytochrome P450 Enzymes and Drug Metabolism in Humans. *Int J Mol Sci* **22**,
doi:10.3390/ijms222312808 (2021).
24 Kiessling, S. *et al.* Enhancing circadian clock function in cancer cells inhibits tumor growth. *BMC*
biology **15**, 13, doi:10.1186/s12915-017-0349-7 (2017).
25 Zuo, S., Wu, L., Wang, Y. & Yuan, X. Long Non-coding RNA MEG3 Activated by Vitamin D
Suppresses Glycolysis in Colorectal Cancer via Promoting c-Myc Degradation. *Front Oncol* **10**,
274, doi:10.3389/fonc.2020.00274 (2020).
26 Chu, C. M. *et al.* Gene expression profiling of colorectal tumors and normal mucosa by
microarrays meta-analysis using prediction analysis of microarray, artificial neural network,
classification, and regression trees. *Dis Markers* **2014**, 634123, doi:10.1155/2014/634123
(2014).
27 Duarte-Pereira, S. *et al.* Extensive regulation of nicotinate phosphoribosyltransferase (NAPRT)
expression in human tissues and tumors. *Oncotarget* **7**, 1973-1983,
doi:10.18632/oncotarget.6538 (2016).
28 Gan, C. *et al.* Carboxylesterase 1 family knockout alters drug disposition and lipid metabolism.
Acta Pharm Sin B **13**, 618-631, doi:10.1016/j.apsb.2022.10.017 (2023).
29 Wang, L. *et al.* Antimicrobial activity of human islet amyloid polypeptides: an insight into
amyloid peptides' connection with antimicrobial peptides. *Biological Chemistry* **393**, 641-646,
doi:doi:10.1515/hsz-2012-0107 (2012).
30 Emterling, A. *et al.* Clinicopathological significance of Nup88 expression in patients with
colorectal cancer. *Oncology* **64**, 361-369, doi:10.1159/000070294 (2003).
31 Merkulova, M. *et al.* Mapping the H(+) (V)-ATPase interactome: identification of proteins
involved in trafficking, folding, assembly and phosphorylation. *Sci Rep* **5**, 14827,
doi:10.1038/srep14827 (2015).
32 Chen, L. J. *et al.* IGF2BP3 promotes the progression of colorectal cancer and mediates
cetuximab resistance by stabilizing EGFR mRNA in an m(6)A-dependent manner. *Cell Death Dis*
14, 581, doi:10.1038/s41419-023-06099-y (2023).
33 Garneau, H., Paquin, M. C., Carrier, J. C. & Rivard, N. E2F4 expression is required for cell cycle
progression of normal intestinal crypt cells and colorectal cancer cells. *J Cell Physiol* **221**, 350-
358, doi:10.1002/jcp.21859 (2009).
34 Chu, X. Y. *et al.* FOXM1 expression correlates with tumor invasion and a poor prognosis of
colorectal cancer. *Acta Histochem* **114**, 755-762, doi:10.1016/j.acthis.2012.01.002 (2012).
35 Fang, Z. *et al.* NFYB-induced high expression of E2F1 contributes to oxaliplatin resistance in
colorectal cancer via the enhancement of CHK1 signaling. *Cancer Lett* **415**, 58-72,
doi:10.1016/j.canlet.2017.11.040 (2018).
36 Cifkova, E. *et al.* Determination of lipidomic differences between human breast cancer and
surrounding normal tissues using HILIC-HPLC/ESI-MS and multivariate data analysis. *Anal*
Bioanal Chem **407**, 991-1002, doi:10.1007/s00216-014-8272-z (2015).
37 Eggers, L. F. *et al.* Lipidomes of lung cancer and tumour-free lung tissues reveal distinct
molecular signatures for cancer differentiation, age, inflammation, and pulmonary
emphysema. *Sci Rep* **7**, 11087, doi:10.1038/s41598-017-11339-1 (2017).
38 Rohrig, F. & Schulze, A. The multifaceted roles of fatty acid synthesis in cancer. *Nat Rev Cancer*
16, 732-749, doi:10.1038/nrc.2016.89 (2016).
39 Coleman, O., Ecker, M. & Haller, D. Dysregulated lipid metabolism in colorectal cancer. *Curr*
Opin Gastroenterol **38**, 162-167, doi:10.1097/MOG.0000000000000811 (2022).
40 Porstmann, T. *et al.* SREBP activity is regulated by mTORC1 and contributes to Akt-dependent
cell growth. *Cell Metab* **8**, 224-236, doi:10.1016/j.cmet.2008.07.007 (2008).
41 Zaytseva, Y. Y. *et al.* Cancer cell-associated fatty acid synthase activates endothelial cells and
promotes angiogenesis in colorectal cancer. *Carcinogenesis* **35**, 1341-1351,
doi:10.1093/carcin/bgu042 (2014).

42   Zaytseva, Y. Y. *et al.* Inhibition of fatty acid synthase attenuates CD44-associated signaling and
reduces metastasis in colorectal cancer. *Cancer Res* **72**, 1504-1517, doi:10.1158/0008-
5472.CAN-11-4057 (2012).
43   Ecker, J. *et al.* The Colorectal Cancer Lipidome: Identification of a Robust Tumor-Specific Lipid
Species Signature. *Gastroenterology* **161**, 910-923 e919, doi:10.1053/j.gastro.2021.05.009
(2021).
44   Singh, A. *et al.* DIABLO: an integrative approach for identifying key molecular drivers from
multi-omics assays. *Bioinformatics* **35**, 3055-3062, doi:10.1093/bioinformatics/bty1054
(2019).
45   Douglas, G. M. *et al.* PICRUSt2 for prediction of metagenome functions. *Nat Biotechnol* **38**,
685-688, doi:10.1038/s41587-020-0548-6 (2020).
46   Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes and the BioCyc
collection of Pathway/Genome Databases. *Nucleic Acids Res* **42**, D459-471,
doi:10.1093/nar/gkt1103 (2014).
47   Subramanian, C., Frank, M. W., Batte, J. L., Whaley, S. G. & Rock, C. O. Oleate hydratase from
Staphylococcus aureus protects against palmitoleic acid, the major antimicrobial fatty acid
produced by mammalian skin. *J Biol Chem* **294**, 9285-9294, doi:10.1074/jbc.RA119.008439
(2019).
48   Alnaseri, H. *et al.* Inducible Expression of a Resistance-Nodulation-Division-Type Efflux Pump
in Staphylococcus aureus Provides Resistance to Linoleic and Arachidonic Acids. *J Bacteriol*
197, 1893-1905, doi:10.1128/JB.02607-14 (2015).
49   Dohlman, A. B. *et al.* The cancer microbiome atlas: a pan-cancer comparative analysis to
distinguish tissue-resident microbiota from contaminants. *Cell Host Microbe* **29**, 281-298 e285,
doi:10.1016/j.chom.2020.12.001 (2021).
50   Bray, F. *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality
worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* **68**, 394-424,
doi:10.3322/caac.21492 (2018).
51   So, J. S. *et al.* Silencing of lipid metabolism genes through IRE1alpha-mediated mRNA decay
lowers plasma lipids in mice. *Cell Metab* **16**, 487-499, doi:10.1016/j.cmet.2012.09.004 (2012).
52   Sriburi, R. *et al.* Coordinate regulation of phospholipid biosynthesis and secretory pathway
gene expression in XBP-1(S)-induced endoplasmic reticulum biogenesis. *J Biol Chem* **282**, 7024-
7034, doi:10.1074/jbc.M609490200 (2007).
53   Lee, A. H., Scapa, E. F., Cohen, D. E. & Glimcher, L. H. Regulation of hepatic lipogenesis by the
transcription factor XBP1. *Science* **320**, 1492-1496, doi:10.1126/science.1158042 (2008).
54   Bommiasamy, H. *et al.* ATF6alpha induces XBP1-independent expansion of the endoplasmic
reticulum. *J Cell Sci* **122**, 1626-1636, doi:10.1242/jcs.045625 (2009).
55   Vriens, K. *et al.* Evidence for an alternative fatty acid desaturation pathway increasing cancer
plasticity. *Nature* **566**, 403-406, doi:10.1038/s41586-019-0904-1 (2019).
56   Mika, A. *et al.* Rearrangements of Blood and Tissue Fatty Acid Profile in Colorectal Cancer -
Molecular Mechanism and Diagnostic Potential. *Front Oncol* **11**, 689701,
doi:10.3389/fonc.2021.689701 (2021).
57   Dejea, C. M. *et al.* Patients with familial adenomatous polyposis harbor colonic biofilms
containing tumorigenic bacteria. *Science* **359**, 592-597, doi:10.1126/science.aah3648 (2018).
58   Kim, M. *et al.* Fecal Metabolomic Signatures in Colorectal Adenoma Patients Are Associated
with Gut Microbiota and Early Events of Colorectal Cancer Pathogenesis. *mBio* **11**,
doi:10.1128/mBio.03186-19 (2020).
59   Yachida, S. *et al.* Metagenomic and metabolomic analyses reveal distinct stage-specific
phenotypes of the gut microbiota in colorectal cancer. *Nat Med* **25**, 968-976,
doi:10.1038/s41591-019-0458-7 (2019).
60   Thomas, A. M. *et al.* Metagenomic analysis of colorectal cancer datasets identifies cross-cohort
microbial diagnostic signatures and a link with choline degradation. *Nat Med* **25**, 667-678,
doi:10.1038/s41591-019-0405-7 (2019).

61   Wirbel, J. *et al.* Meta-analysis of fecal metagenomes reveals global microbial signatures that
are specific for colorectal cancer. *Nat Med* **25**, 679-689, doi:10.1038/s41591-019-0406-6
(2019).

62   Haug, K. *et al.* MetaboLights-an open-access general-purpose repository for metabolomics
studies and associated meta-data. *Nucleic acids research* **41**, D781-D786,
doi:10.1093/nar/gks1004 (2013).

63   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550,
doi:10.1073/pnas.0506580102 (2005).

64   Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739-1740,
doi:10.1093/bioinformatics/btr260 (2011).

65   Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res*
28, 27-30, doi:10.1093/nar/28.1.27 (2000).

66   Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv*, 060012, doi:10.1101/060012
(2021).

67   Morgan M, F. S., Gentleman R. GSEABase: Gene set enrichment data structures and methods.
R package version 1.62.0 (2023).

68   Blighe K, R. S., Lewis M. EnhancedVolcano: Publication-ready volcano plots with enhanced
colouring and labeling. *R package version 1.18.0* (2023).

69   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).

70   Metwaly, A. *et al.* Integrated microbiota and metabolite profiles link Crohn's disease to sulfur
metabolism. *Nat Commun* **11**, 4322, doi:10.1038/s41467-020-17956-1 (2020).

71   Tsugawa, H. *et al.* MS-DIAL: data-independent MS/MS deconvolution for comprehensive
metabolome analysis. *Nat Methods* **12**, 523-526, doi:10.1038/nmeth.3393 (2015).

72   Wehrens, R. *et al.* Improved batch correction in untargeted MS-based metabolomics.
Metabolomics **12**, 88, doi:10.1007/s11306-016-1015-8 (2016).

73   Ecker, J., Scherer, M., Schmitz, G. & Liebisch, G. A rapid GC-MS method for quantification of
positional and geometric isomers of fatty acid methyl esters. *J Chromatogr B Analyt Technol*
Biomed Life Sci **897**, 98-104, doi:10.1016/j.jchromb.2012.04.015 (2012).

74   Shah, A. K. P. *et al.* Statistical analysis of feature-based molecular networking results from non-
targeted metabolomics data. *Nature Protocols* **20**, doi:10.1038/s41596-024-01046-3 (2025).

75   Pakkir Shah, A. K. *et al.* The Hitchhiker's Guide to Statistical Analysis of Feature-based
Molecular Networks from Non-Targeted Metabolomics Data. doi:10.26434/chemrxiv-2023-
wwbt0 (2023).

76   Godon, J. J., Zumstein, E., Dabert, P., Habouzit, F. & Moletta, R. Molecular microbial diversity
of an anaerobic digestor as determined by small-subunit rDNA sequence analysis. *Appl Environ*
Microbiol **63**, 2802-2813 (1997).

77   Salter, S. J. *et al.* Reagent and laboratory contamination can critically impact sequence-based
microbiome analyses. *BMC Biol* **12**, 87, doi:10.1186/s12915-014-0087-z (2014).

78   Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D. Development of a
dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on
the MiSeq Illumina sequencing platform. *Applied and environmental microbiology* **79**, 5112-
5120, doi:10.1128/AEM.01043-13 (2013).

79   Lagkouvardos, I. *et al.* IMNGS: A comprehensive open resource of processed 16S rRNA
microbial profiles for ecology and diversity studies. *Sci Rep* **6**, 33721, doi:10.1038/srep33721
(2016).

80   Edgar, R. C. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat*
Methods **10**, 996-998, doi:10.1038/nmeth.2604 (2013).

81   Edgar, R. C. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon sequencing.
bioRxiv, 081257, doi:10.1101/081257 (2016).

82   Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for rapid assignment
of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* **73**, 5261-5267,
doi:10.1128/AEM.00062-07 (2007).
83   Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A. & Callahan, B. J. Simple statistical
identification and removal of contaminant sequences in marker-gene and metagenomics data.
Microbiome **6**, 226, doi:10.1186/s40168-018-0605-2 (2018).
84   Wirbel, J. *et al.* Microbiome meta-analysis and cross-disease comparison enabled by the
SIAMCAT machine learning toolbox. *Genome Biol* **22**, 93, doi:10.1186/s13059-021-02306-1
(2021).
85   Sidak, D., Schwarzerova, J., Weckwerth, W. & Waldherr, S. Interpretable machine learning
methods for predictions in systems biology from omics data. *Front Mol Biosci* **9**, 926623,
doi:10.3389/fmolb.2022.926623 (2022).
86   Rohart, F., Gautier, B., Singh, A. & Le Cao, K. A. mixOmics: An R package for 'omics feature
selection and multiple data integration. *PLoS Comput Biol* **13**, e1005752,
doi:10.1371/journal.pcbi.1005752 (2017).
87   Riva, A. *et al.* Identification of inulin-responsive bacteria in the gut microbiota via multi-modal
activity-based sorting. *Nature communications* **14**, 8210, doi:10.1038/s41467-023-43448-z
(2023).
88   Hatzenpichler, R. *et al.* In situ visualization of newly synthesized proteins in environmental
microbes using amino acid tagging and click chemistry. *Environ Microbiol* **16**, 2568-2590,
doi:10.1111/1462-2920.12436 (2014).
89   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**,
676-682, doi:10.1038/nmeth.2019 (2012).
90   Garcia-Alonso, L., Holland, C. H., Ibrahim, M. M., Turei, D. & Saez-Rodriguez, J. Benchmark and
integration of resources for the estimation of human transcription factor activities. *Genome*
Res **29**, 1363-1375, doi:10.1101/gr.240663.118 (2019).
91   Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear Mixed
Effects Models. *Journal of Statistical Software* **82**, 1 - 26, doi:10.18637/jss.v082.i13 (2017).
92   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate - a Practical and Powerful
Approach to Multiple Testing. *J R Stat Soc B* **57**, 289-300, doi:DOI 10.1111/j.2517-
6161.1995.tb02031.x (1995).
93   Budczies, J. *et al.* Cutoff Finder: a comprehensive and straightforward Web application
enabling rapid biomarker cutoff optimization. *PLoS One* **7**, e51862,
doi:10.1371/journal.pone.0051862 (2012).
94   Anderson, M. J. A new method for non-parametric multivariate analysis of variance. *Austral*
Ecol **26**, 32-46, doi:DOI 10.1046/j.1442-9993.2001.01070.x (2001).
95   Ward, J. H. Hierarchical Grouping to Optimize an Objective Function. *Journal of the American*
Statistical Association **58**, 236-244, doi:10.1080/01621459.1963.10500845 (1963).
96   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*
57, 289-300, doi:10.1111/j.2517-6161.1995.tb02031.x (1995).
97   Lagkouvardos, I., Fischer, S., Kumar, N. & Clavel, T. Rhea: a transparent and modular R pipeline
for microbial profiling based on 16S rRNA gene amplicons. *PeerJ* **5**, e2836,
doi:10.7717/peerj.2836 (2017).
98   McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. *PLoS One* **8**, e61217, doi:10.1371/journal.pone.0061217
(2013).
99   Chen, J. *et al.* Associating microbiome composition with environmental covariates using
generalized UniFrac distances. *Bioinformatics* **28**, 2106-2113,
doi:10.1093/bioinformatics/bts342 (2012).
100 Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome Biology* **12**,
doi:ARTN R60

10.1186/gb-2011-12-6-r60 (2011).

101   Nearing, J. T. *et al.* Microbiome differential abundance methods produce different results
1637   across 38 datasets. *Nature Communications* **13**, doi:ARTN 342

1638   10.1038/s41467-022-28034-z (2022).

1639   102   Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A. & Callahan, B. J. Simple statistical
1640   identification and removal of contaminant sequences in marker-gene and metagenomics data.
1641   *Microbiome* **6**, doi:ARTN 226

1642   10.1186/s40168-018-0605-2 (2018).

1643   103   Chen, J. *et al.* Associating microbiome composition with environmental covariates using
1644   generalized   UniFrac   distances.   *Bioinformatics*   **28**,   2106-2113,
1645   doi:10.1093/bioinformatics/bts342 (2012).

1646   104   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
1647   RNA-seq data with DESeq2. *Genome biology* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).

1648   105   Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of
1649   intersecting   sets   and   their   properties.   *Bioinformatics*   **33**,   2938-2940,
1650   doi:10.1093/bioinformatics/btx364 (2017).

1651

Fig.1

a ATF6 H-score — NUC, CYT (****)

ATF6- TC
ATF6-High — H-score NUC 261.48, CYT 293.28 — 38 %
ATF6-Low — H-score NUC 55.48, CYT 121.23 — 62 %

ATF6- IF
ATF6-High — H-score NUC 113.06, CYT 238.3
ATF6-Low — H-score NUC 17.34, CYT 70.09

b True Labels vs Predicted Labels
ATF6-High / ATF6-High: 311, ATF6-High / ATF6-Low: 44
ATF6-Low / ATF6-High: 47, ATF6-Low / ATF6-Low: 516
% of row total

c ATF6-High 39 %, ATF6-Low 61 %

d Predicted values vs True values (NUC, CYT)

e ATF6-High 35.9 %, ATF6-Low 64.1 %

f ATF6-LOCRC
ATF6 H-score — NUC, CYT (****)
ATF6-High 19 %
ATF6-Low 81 %

g ATF6-EOCRC
ATF6 H-score — NUC, CYT
ATF6-High 11 %
ATF6-Low 89 %

Fig.2

Fig. 3

Fig. 4

Fig.5