Time-dependent Interaction Modification Generated from Plant-soil Feedback

Heng-Xing Zou[*,a], Xinyi Yan[b], Volker H.W. Rudolf[a]

a. Program in Ecology and Evolutionary Biology, Department of BioSciences, Rice University, Houston, TX 77005

b. Department of Integrative Biology, The University of Texas at Austin, Austin, TX 78712

* Corresponding author. Email address: hxzou.ecology@gmail.com.

ORCIDs: Zou: 0000-0002-1965-6910; Yan: 0000-0003-0480-5044; Rudolf: 0000-0002-9214-2000

Author Contributions: H.-X.Z. conceived the idea, and designed the research with assistance from X.Y.; H.-X.Z performed the research with assistance from V.H.W.R; H.-X.Z. wrote the first version of the manuscript, and all authors contributed to revisions.

Keywords: community assembly, higher-order interactions, plant-soil feedback, plant-microbe interactions, phenology

Running Title: Time-dependent Interaction Modifications

Article Type: Letter

Word Count (Main Text): 4991

Word Count (Abstract): 149

Number of References: 92

List of Elements: Figures 1-4, Table 1

Data Availability Statement: Code and data used to generate all results have been deposited at Dryad: https://doi.org/10.5061/dryad.b8gtht7kf

Abstract

Pairwise interactions between species can be modified by other community members, leading to emergent dynamics contingent on community composition. Despite the prevalence of such higher-order interactions, little is known about how they are linked to the timing and order of species' arrival. We generate population dynamics from a mechanistic plant-soil feedback model, then apply a general theoretical framework to show that the modification of a pairwise interaction by a third plant depends on its germination phenology. These time-dependent interaction modifications emerge from concurrent changes in plant and microbe populations and are strengthened by higher overlap between plants' associated microbiomes. The interaction between this overlap and the specificity of microbiomes further determines plant coexistence. Our framework is widely applicable to mechanisms in other systems from which similar time-dependent interaction modifications can emerge, highlighting the need to integrate temporal shifts of species interactions to predict the emergent dynamics of natural communities.

Introduction

Complex communities in nature often reassemble every growing season. The relative timing of species life history events in each season, i.e., phenology, can affect species interactions and community structure (Fukami 2015; Rudolf 2019; Zou & Rudolf 2023a). In pairwise systems, the shift in species interactions and community states with changes in species arrival times is well described by the framework of priority effects. However, theories studying the effects of variation in arrival sequence on pairwise systems (Song *et al.* 2020; Zou & Rudolf 2023b) have fallen behind the numerous experiments conducted in multispecies communities (Dickie *et al.* 2012; Drake 1991; Pu & Jiang 2015; Song *et al.* 2021b; Uricchio *et al.* 2019). The focus on pairwise interactions limits our perspective on widespread higher-order interactions that arise when the presence of a third species affects the interactions between two species (Billick & Case 1994; Werner & Peacor 2003; Wootton 1994) and are important to maintaining biodiversity (Levine *et al.* 2017). As species-rich communities in nature face widespread reshuffling in phenology and timing of interactions (Kharouba *et al.* 2018; Parmesan 2006), we need a general framework for higher-order interactions that includes this temporal dimension of community structure.

Documented in systems from microbes and plants to crustaceans and amphibians (Blackford *et al.* 2020; Fragata *et al.* 2022; Grainger *et al.* 2019; Rudolf 2018; Shorrocks & Bingley 1994), priority effects are generated by either positive frequency dependence or temporal changes in species traits due to different arrival times. Here, "arrival" can refer to both colonizations over many generations such as succession, and recurring arrival time such as the annual reassembly of communities (Zou & Rudolf 2023a). For example, early-colonizing bacteria may grow to higher abundance and exclude late arrivers (Grainger *et al.* 2019), and plants can be affected by the

chemical composition or microbiomes in soil previously occupied by other species (Kardol *et al.* 2007). Because these changes in relative abundances or traits are not instantaneous but develop over time, the relative arrival time between two species directly modifies their interaction strengths. Similarly, in three-species communities, how the third species affects a pairwise interaction (interaction modification) can depend on its arrival time, which can change its density, traits, or both, and that of other community members. For instance, increasing the coverage of barnacles decreases bird predation of limpets in intertidal communities (Wootton 1993). If the coverage (*density*) of barnacles depends on the length they colonized the patch before limpets arrive, then this change in coverage over time could lead to a temporal change in the strength of interaction modification. In an aquatic predator-prey system, morphological changes of snails induced by the presence of a first predator increased its risk towards a second predator (Hoverman & Relyea 2008). Because changes in morphological *traits* are not instantaneous, the length of the acclimation period with the first predator could affect the amount of increased predation risk. These temporal changes in interaction modification are the higher-order equivalent of priority effects, and we term them "time-dependent interaction modification." This term includes all forms of higher-order interactions that can change with species' arrival times.

    In plant communities, such time-dependent interaction modification could arise from plant-soil feedback. Early-arriving plants modify the soil environment, e.g., via allelopathy, changes in the microbiome, and litter addition from previous residents, which reduces the fitness of the later arriving plant (Inderjit & van der Putten 2010; van der Putten *et al.* 2016; Suding & Goldberg 1999). However, the strength of the plant-soil feedback often depends on how long the plant conditions the soil (Ke *et al.* 2021; Lepinay *et al.* 2018). Furthermore, plant-plant interactions

can also be mediated by soil microbiome cultivated by a third species arriving before them

(Zhang *et al.* 2020). These studies indicate the potential of time-dependent interaction

modification in plant communities, but this relationship is still unknown.

To fill this conceptual gap, we explore how time-dependent interaction modifications can

arise in nature. We generate population dynamics from a mechanistic plant-soil feedback model

that focuses on the interaction between plants and their microbiomes. Then, we apply a

framework of time-dependent interaction modification to the simulated data to reveal the

relationship between the temporal structure of communities and their higher-order interactions.

Methods

Plant-soil Feedback Model

We simulate seasonal population dynamics of plants using a mechanistic, discrete-time

model of plant-soil feedback to explore how time-dependent interaction modifications can arise

in natural communities (Figure 1). Each year has T time steps. We use τ to represent each

specific time step within the year ($1 \leq \tau \leq T$), and each plant species germinates from dormancy

at a specific time step $\tau = p_x$, where p_x is the germination phenology of the plant x. Note that a

"year" defines the suitable growing season for plants and microbiomes, which may be a few

months per calendar year. After germination, each plant cultivates microbiomes which affects the

mortality rates of all plants (Bever *et al.* 1997). At the end of their life cycles, plants reproduce

and senesce. All plants finish their life cycles within a year, regardless of their germination

phenology. We assume that life cycles of all plants have identical lengths of l time steps

($p_x + l \leq T$), such that the plant germinating earliest will also reproduce and senesce earliest in

the year.

Within the life cycles, the mortality of plant x depends on its density ($N_x(\tau)$) and the

accumulated effects from all microbiomes ($\sum_y m_{xy} S_y$; Suding *et al.* 2013), where m_{xy} denotes

the effect of microbiome associated with plant y on plant x, S_y the density of the microbiome

associated with plant y, and d_x the density-dependent mortality rate:

$$N_x(\tau + 1) = N_x(\tau)\left(1 - d_x N_x(\tau) + \sum_y m_{xy} S_y(\tau)\right) \text{ when } p_x < \tau < p_x + l \text{ (Eqn. 1)}.$$

At the end of their life cycle, plants produce seeds (D_x) with fecundity λ_x:

$$D_x(\tau + 1) = \lambda_x N_x(\tau) \text{ when } \tau = p_x + l \text{ (Eqn. 2)}.$$

Seeds germinate at the assigned phenology (p_x) at the beginning of the next year without

additional mortality, i.e., $N_x(\tau + 1) = D_x(\tau)$ when $\tau = p_x$.

To highlight the effect of plant-soil feedback, we focus on interspecific interactions arising

from changes in microbiomes (Figure 1) rather than resource competition. This represents a

natural scenario where each species is limited by different resources (Tilman 1982).

Within the year, the population of microbiomes ($S_x(\tau)$) increases proportionally with plant

density at the rate $v_{xy} N_y$ once the plants emerge, where v_{xy} is the rate of cultivation by plant y,

and is limited by a density-dependent mortality ($\mu_x S_x(\tau)$):

$$S_x(\tau + 1) = S_x(\tau)\left(1 + \sum_y v_{xy} N_y(\tau) - \mu_x S_x(\tau)\right) \text{ when } \tau < T \text{ (Eqn. 3)}.$$

Eqn. 3 specifies that when plants are not present (e.g., before seed germination or after

senescence), their associated microbiome immediately starts to decay (Esch & Kobe 2021). We

assume a constant proportion (μ_{year}) of each microbiome does not carry over to the next year:

$$S_x(\tau + 1) = \left(1 - \mu_{year}\right)S_x(\tau) \text{ when } \tau = T \text{ (Eqn. 4)}.$$

Time-dependent Interaction Modification

Higher-order interactions are often measured from fitting population data to statistical

models (Kleinhesselink *et al.* 2022; Letten & Stouffer 2019). Here, we extend the previous

framework of higher-order interactions to define and quantify time-dependent interaction

modification in the plant-soil feedback model (Eqn. 1-4). Consider a three-species community

that assembles over time (Figure 1), where the change in density (N) of a species between time

steps can be characterized by a pairwise Ricker model:

$$\frac{N_i(t+1)}{N_i(t)} = \lambda_i \exp\left(-\alpha_{ii}N_i(t) - \alpha_{ij}N_j(t) - \alpha_{ik}N_k(t)\right) \text{ (Eqn. 5)}.$$

λ_i is the intrinsic growth rate of species i, and α_{ij} is the per-capita effect from species j to i.

In a pairwise system with priority effects (Figure 1), the per-capita interaction from species j to i

can depend on the difference between their germination phenology within a year, $p_j - p_i$, i.e.,

their phenological difference Δp_{ij} (Rudolf 2019; Zou & Rudolf 2023a) (Figure 1). In this case,

we can expand Eqn. 5 into:

$$\frac{N_i(t+1)}{N_i(t)} = \lambda_i \exp\left(-\alpha_{ii}N_i(t) - f_{ij}(\Delta p_{ij})N_j(t) - \alpha_{ik}N_k(t)\right) \text{ (Eqn. 6)}.$$

On the other hand, higher-order interactions in this three-species system can be modeled by

the following variant of the Ricker model (Kleinhesselink *et al.* 2022):

$$\frac{N_i(t+1)}{N_i(t)} = \lambda_i \exp\left(-\alpha_{ii}N_i(t) - \alpha_{ij}N_j(t) - \alpha_{ik}N_k(t) - \beta_{i(ij)}N_i(t)N_j(t) - \beta_{i(ik)}N_i(t)N_k(t) - \right.$$

$$\left. \beta_{i(jk)}N_j(t)N_k(t)\right) \text{ (Eqn. 7)},$$

where the β coefficients quantify the strength of the higher-order interaction (HOI), e.g., $\beta_{i(jk)}$

quantifies how densities of species j and k modify each other's effect on species i (Letten &

Stouffer 2019).

If phenological differences between species can affect the strengths of both pairwise and

higher-order interactions by changing the corresponding per-capita coefficients (Figure 1), in the

most general form, Eqn. 7 can be written as:

$\frac{N_i(t+1)}{N_i(t)} = \lambda_i\, exp\left(-\alpha_{ii}N_i(t) - f_{ij}\left(\Delta p_{ij}\right)N_j(t) - f_{ik}(\Delta p_{ik})N_k(t) - f_{i(ij)}{}'\left(p_i, p_j\right)N_i(t)N_j(t) - \right.$

$\left. f_{i(ik)}{}'(p_i, p_k)N_i(t)N_k(t) - f_{i(jk)}{}'\left(p_i, p_j, p_k\right)N_j(t)N_k(t)\right)$

130   (Eqn. 8),

where $f_{i(jk)}{}'\left(p_i, p_j, p_k\right)$ defines how the HOI coefficient, $\beta_{i(jk)}$, depends on the phenology of all

three species. If we consider species i and j as the "focal pair" and fix their phenology while

shifting species k's phenology, then $f_{ij}\left(\Delta p_{ij}\right)$ and p_j become constants, and $\beta_{i(jk)}$ only depends

on p_k. We maintain this assumption here to simplify our analyses of three-species communities.

To detect time-dependent interaction modification, we compare $\beta_{i(jk)}$ values fitted from

population dynamics with different phenology of species k (p_k). If interaction modifications are

absent, $\beta_{i(jk)}$ should be close to 0; if $\beta_{i(jk)}$ is nonzero but constant for all p_k, then the interaction

modification does not depend on time. This definition does not consider time-dependent

interaction modification via temporal changes in species density, which is beyond our scope.

The accurate quantification of time-dependent interaction modifications requires density

gradients of all three species and a temporal gradient of at least one species, a scale rarely

feasible for empirical studies. A simplified quantification of time-dependent interaction

modifications fits pairwise models between focal species i and j (e.g., Eqn. 5) to experimental

populations with and without species k, then examines how the difference of the two pairwise

interaction coefficients change with the phenology of species k (p_k). Under this definition, the

time-dependent interaction modification combines the effects of both the density and phenology

of species k, and its detection requires much less density gradients of all three species (Appendix

II). In practice, it simplifies the experimental design but carries stronger assumptions on the

community with and without the third species k. We discuss the mathematical links between the

methods and their pros and cons in Appendix II, but focus here on the first approach (Eqn. 8) to

facilitate direct comparison to previous studies (Kleinhesselink *et al.* 2022; Letten & Stouffer

2019). In both approaches, the interpretation of fitted higher-order interactions depends on the

specific system and its biological processes, and the statistical power of the phenomenological

model structure.

Simulation

In nature, plants can share pathogenic or mutualistic microbes (Crawford *et al.* 2019; Gilbert

& Webb 2007; Sedio & Ostling 2013; Spear & Broders 2021), which can be cultivated by

multiple plants in the same community. We simulate this overlap in microbiomes by allowing for

the plant x to cultivate not only its associated microbiome but also those associated with other

plants ($v_{yx} \neq 0$ for all $x \neq y$). Each plant's total capacity of cultivation ($V_x = v_{xx} + \sum_y v_{yx}$ for

all $x \neq y$) is constant, reflecting a finite pool of resources plants can provide to microbiomes. To

simulate different levels of shared microbiomes among plants, we partition each plant's

cultivation rate to simulate three scenarios: no overlap ($\sum_y v_{yx} = 0V_x$), low overlap ($\sum_y v_{yx} =$

$0.25V_x$), and high overlap in microbiomes ($\sum_y v_{yx} = 0.5V_x$). The strength of interspecific

cultivation is equally split between microbiomes associated with other plants and therefore

depends on the number of plant species. These values ensure that the microbiome always

receives the largest (or equal) cultivation effect from its host compared to others.

We focus on the role of soil pathogens only and assume that microbiomes increase the

mortality within the year (i.e., $m_{xy} < 0$; see Table 1), although our methods can be readily

extended to mutualistic and mixed microbiomes. We explored the effect of specificity of the

pathogens by changing the microbiome's effect on its host vs. other plants: if $m_{xx} = m_{yx}$, the

microbiome is a generalist pathogen to all plants; if $|m_{xx}| > |m_{yx}|$ or $|m_{xx}| < |m_{yx}|$, the

microbiome is a specialist on its host, or plants other than its host. To simulate the specificity of

microbiomes found in nature, we calculated the distribution of m_{xx}/m_{yx} from an empirical

metaanalysis (Yan *et al.* 2022), then randomly drew this proportion to partition the total capacity

of feedback. See Appendix I for detailed methods.

Each year has 12 time steps ($T = 12$) and the length of plant life cycles (l) is 6 time steps

($l = 6$); therefore, the earliest germination phenology is 1 (the beginning of the year), and the

latest 6. Plants' germination phenology does not change over years or with other plants. This

range ensures enough time for species interaction while allowing for a proper gradient of

phenological differences between plants. We choose identical fecundity λ and effects of

microbiomes m_{xy} (Table 1) for all plants, such that any differences in population dynamics

would arise from phenological differences between plant species, different overlaps of

microbiomes, and different host specificity of microbiomes.

To accurately obtain interaction coefficients (Eqn. 8), we used a response surface design

(commonly used in empirical studies; Inouye 2001; Kleinhesselink *et al.* 2022) with 6 densities

of each plant species, yielding 36 and 216 combinations of initial densities for two-plant and

three-plant communities. We ran all response surface simulations for one year, representing the

typical duration of plant experiments. We simulate two-plant communities with a gradient of 11

phenological differences from $\Delta p_{ij} = p_j - p_i = -5$ (species *j* early by 5 time steps) to $\Delta p_{ij} = 5$

(species i early by 5 time steps). In three-plant communities, when a third plant is added to the

focal pair, it could germinate either before (scenario I), between (scenario II), or after the focal

species pair (scenario III; Figure 3). For simplicity, we examined scenarios I and III by fixing the

phenology of the focal pair (i and j) both at the latest (time 6; scenario I) or at the earliest (time

1; scenario III) and changed the phenology of the third plant p_k ranging from 1 to 6. This allows

all plants to germinate simultaneously in some cases. To explore the case where plant k

germinates between the focal pair (scenario II), we let plant i germinate at the earliest (time 1)

and j at the latest (time 6) and again let $1 \leq p_k \leq 6$.

We fit a discrete-time Ricker model containing all higher-order interaction terms to plant

population at the end of the year (Eqn. 8). The Ricker model structure fits our simulated data best

among several other discrete-time population models (Hart *et al.* 2018) based on Akaike

Information Criterion (AIC; Appendix I, Figure S1). To evaluate whether a model with higher-

order coefficients indeed significantly improves model performance, we fitted three-plant

population dynamics to pairwise Ricker models (Eqn. 1) with either plants i or j as the focal

species, then compared both the root-mean-squared error (RMSE) and the AIC values of the

pairwise (Eqn. 1) and higher-order Ricker model (Eqn. 8) at different p_k. Finally, to test the

sensitivity to model parameters, we randomly drew 100 sets of microbial effects (m) and

cultivation rate (V) from a uniform distribution (Table 1), and fitted Eqn. 8 to obtain a

distribution of higher-order coefficients.

We run the model for 50 years (600 time steps) with equal initial densities of plants and

microbiomes to evaluate coexistence; the duration ensures that all populations reach equilibrium.

We conducted all simulations in R 4.2.1 (R Core Team 2022) and fitted models using the

function *nlsLM()* in the R package *minpack.lm* (Elzhov *et al.* 2016). The code is available at

https://doi.org/10.5061/dryad.b8gtht7kf.

Results

Two-plant Communities

We first examine two-plant communities to analyze the causes and consequences of priority

effects. The late plant always has a lower population, and this disadvantage is smaller with

smaller differences between the germination phenology of the two plants (Figure 2A). The effect

of the early plant on the late plant increases when it germinates earlier. Changes of α_{ij} and α_{ji}

with differences in germination phenology Δp_{ij} for both plants are symmetrical (Figure 2B) and

nonlinear. Note that when one plant germinates very late ($| \Delta p_{ij} | \geq 4$), its effect on the early

plant is negligible because the temporal overlap between the two plants is too short, and the

effect it experiences from the early plant slightly decreases because the early plant's microbiome

declines without its host.

The overlap between plants' microbiomes interacts with the specificity of microbiomes to

affect the final populations and interaction coefficients. When the microbiome affects other

plants more negatively than its host ($|\mathrm{m_{xx}}| < |m_{yx}|$, for $x \neq y$), each plant receives less

feedback from its microbiome, and the early plant tends to exclude the late plant. However, with

increasing overlap the early plant cultivates the late plant's microbiome, which subsequently

exerts negative feedback to the early plant, promoting coexistence (Figure S5). When the

microbiome affects its host the most ($|\mathrm{m_{xx}}| > |m_{yx}|$), increasing overlap leads to a stronger

negative interaction between plants and promotes monodominance (Figure S6).

Three-plant Communities: Population Dynamics

We examine patterns of equilibrium population density following three scenarios of plant phenology (plant k germinates before, between, or after the focal pair; Figure 3). In general, the earlier a plant germinates relative to its competitors, the higher its equilibrium population (Figure 3). However, when all three plants germinate at different times (scenario II), both the earliest and the last plants benefit from the later germination of the intermediate species, while only the intermediated species itself is negatively affected. This alternating pattern indicates a time-dependent interaction chain: When plant k germinates later than plant i (scenario II), plant i experiences lower total microbiome densities and reaches a higher equilibrium population. Because plant k germinates later it has also less time to cultivate its microbiome. This combination of reduced density and cultivation time of the intermediate plant (k) results in a decline in the total density of the microbiome, which benefits the last plant j (Figure S3).

Like in two-plant communities, levels of overlap in microbiome cultivation among plants interact with the specificity of microbiomes. When all microbial effects are equal, larger overlaps (higher inter- vs. intraspecific cultivation) lead to higher equilibrium populations because plants with earlier phenology devote less to cultivating their microbiomes, leading to a slower increase of microbiomes before the late plants germinate and subsequently a lower overall density of microbiomes (Figure S3). Larger overlap promotes coexistence if the microbial effect on its host is smaller than to others ($|m_{xx}| < |m_{yx}|$) but promotes monodominance if $|m_{xx}| > |m_{yx}|$ (Figure S7-S8).

Time-dependent Interaction Modifications

We examine patterns of time-dependent interaction modifications using the higher-order

coefficient associated with densities of all three plants ($\beta_{i(jk)}$ for plant i, $\beta_{j(ik)}$ for plant j, and

$\beta_{k(ij)}$ for plant k) following the same scenarios of plant phenology (Figure 4). Models that

include higher-order interaction terms almost always fit best, except when fitted higher-order

coefficients are close to 0, despite additional parameters (Figure S16-S17). These results support

the observed time-dependent interaction modifications quantified by our methods. The fitted

higher-order coefficients are generally smaller in value than pairwise coefficients, but the

comparison of the strength of higher-order versus pairwise interaction is contingent on species

densities (Eqn. 7).

In general, all higher-order interaction coefficients depend on the relative phenology of

plants and the level of overlap in associated microbiomes (Figure 4). When plant k germinates

before both plants i and j (the focal pair; scenario I, the strengths of higher-order interactions

experienced by late plants i and j ($\beta_{i(jk)}$ and $\beta_{j(ik)}$) decreases with earlier germination

phenology of plant k without overlap, increases then decreases with low overlap, and decreases

nonlinearly with high overlap of microbiomes. Without overlap in microbiomes, plants i and j's

microbiomes are not cultivated by early-germinating plant k, and the focal pair's exposure to

plant k's microbiome also decreases as plant k germinates earlier. The combined effects lead to

negative $\beta_{i(jk)}$ and $\beta_{j(ik)}$ values, weakening the *overall* interactions experienced by plants i and

j.

When plant k germinates between plants i and j (scenario II), the first plant i always

experiences the weakest higher-order interaction. Similar to scenario I, without overlap in

microbiomes, the latest-germinating plant j is less affected by other plants' microbiomes, while

the cascading effect between the three plants observed in population dynamics further weakens

the interaction it experiences. By germinating later, plant k decreases its temporal overlap with

the earliest plant i and its microbiome. These effects are reflected by the negative $\beta_{j(ik)}$ and

$\beta_{k(ij)}$ values. With higher overlap in microbiomes, the early-germinating plants cultivate

microbiomes of late-germinating plants, which experience stronger higher-order interactions.

When plant k germinates after both plants i and j (scenario III), higher-order interactions

experienced by plants i and j ($\beta_{i(jk)}$ and $\beta_{j(ik)}$) generally decrease with later phenology of plant

k because its effect on the microbiomes associated with the focal pair diminishes, while the

higher-order effect of the early plants on the late germinating plant k ($\beta_{k(ij)}$) generally increases

because the effects from the focal pair get stronger when they germinate relatively earlier.

Although microbiomes' specificity (i.e., the strength of m_{xx} vs. m_{yx}) interacts with the

overlap of microbiomes to determine the long-term coexistence between three plants, qualitative

patterns of time-dependent interaction modifications are unchanged (Figure S9-S10). Drawing

microbial effects and cultivation rates from uniform distributions, or from the distribution

informed by the range found in experiments (where $m_{xx}/m_{yx} \sim lognormal(0.109, 1.168)$; see

Appendix I) changes the strengths but not patterns of time-dependent interaction modifications

(Figure S12-S15).

We focus on terms that involve all three plants ($\beta_{i(jk)}$, $\beta_{j(ik)}$, and $\beta_{k(ij)}$) because they are

unique to three-plant communities. All other higher-order terms in Eqn. 8 change with the

phenology of plant k, and both the magnitudes and the temporal patterns shift along a continuum

with increasing overlap in microbiomes (Figure S11).

Discussion

To understand the present and future of ecological communities, we need to first understand

their history (Fukami 2015; Gause 1934). Here, we bring this temporal perspective to

multispecies communities with higher-order interactions. Using a mechanistic model of plant-

soil feedback, we show that the germination phenology of a third plant determines how it

modifies interaction between other plants. The strength of this interaction modification depends

on the overlap in associated microbiomes between all plants which control indirect interaction

effects in this system. Together, these results highlight the time-dependence of higher-order

interactions in complex communities and provide a road map to further exploring the temporal

dimension of species interactions in diverse communities.

Our framework puts higher-order interactions in a temporal context. In our model, the

accumulation of microbiomes over time affects the strength of plant-plant interactions. The

germination phenology of a third plant marks the starting point of its microbiome cultivation,

subsequently the total microbiome densities of the community and their effects on plants.

Importantly, the interaction modification changes with phenological differences because the

underlying biological process is not instantaneous, despite models of community dynamics that

often assume otherwise. Non-instantaneous processes in real ecosystems can lead to time lags

between different populations, especially when their life cycles are of similar time scales (e.g.

the hare-lynx cycle; Nisbet 1997; Stenseth *et al.* 1998; Adler *et al.* 2008). The ubiquity of these

temporal processes implies that all higher-order interactions have the potential to be time-

dependent.

Higher-order interactions can arise from various ecological mechanisms. When one species

alters the environment, the modification can often affect how other species interact, leading to

higher-order interactions (Wootton 1993; Zhang *et al.* 2020). When these environmental

modifications progress with time, they can also lead to priority effects (Chappell *et al.* 2022).

The temporal process of environmental modification is therefore a potential source of time-

dependent interaction modification in complex communities. In our model, the cultivation of

microbiomes by plants modifies the fitness of other plants, and the different germination

phenology affects the temporal extent of these environmental modifications, leading to time-

dependent interaction modifications. Similarly, one species can alter the traits of other species,

which can also progress with time, leading to time-dependent interaction modifications

(Hoverman & Relyea 2008; O'Keeffe *et al.* 2021; Werner & Peacor 2003). Theoretical studies

often mechanistically model individual species' demographic parameters or traits that give rise to

higher-order interactions, such as growth rates (Kleinhesselink *et al.* 2022; Lai *et al.* 2022),

fecundity (Mayfield & Stouffer 2017), and size (Levine *et al.* 2017). Since these traits often

change with species' arrival times (Blackford *et al.* 2020; Rasmussen *et al.* 2014) the strengths of

resulting higher-order interactions should also be time-dependent. Incorporating time-

dependence in these traits is essential to understand changes in higher-order interactions over

time in complex communities.

In our model, the interaction between two generalist-specialist gradients not only determines

the magnitude of higher-order interactions but also the coexistence of plants. A plant can

specialize in cultivating its microbiome (no overlap), or cultivate all other plants' microbiomes

(total overlap), whereas a microbiome (in our model, pathogenic) can specialize in its host or

other plants (higher microbial effects), or equally affect all plants (uniform microbial effects). If

a microbiome specializes in plants other than its host, its host would effectively have lower self-

limitation than limitation to other plants, leading to monodominance. However, higher overlap

allows it to limit itself by cultivating other plants' microbiomes promoting coexistence. In this

case, higher overlap promotes both time-dependent interaction modifications and coexistence.

On the other hand, when a microbiome specializes on its host, higher overlap promotes

monodominance by increasing the microbiomes of other plants, raising the limitation to others

above that of the host plant. These dynamics arising from the two pairs of generalist vs. specialist

gradients provide a mechanistic explanation of the strengths of higher-order interactions and

coexistence between plants, but not necessarily a direct relationship between the two. Our results

provide a detailed perspective on how the composition of generalist vs. specialist can shape

dynamics across communities (Garzon-Lopez *et al.* 2015; Holt & Lawton 1993; Jiao & Cortez

2022; Snyder & Ives 2001). Although the temporal aspect of these indirect interactions is studied

in the context of pairwise priority effects (Clay *et al.* 2019; Jiao & Cortez 2022), future studies

need to consider the role of time-dependent interaction modifications in these intrinsically

complex communities (e.g., O'Keeffe *et al.* 2021).

In nature, plants often share mutualists or pathogens, especially among those more

phylogenetically related (Davison *et al.* 2015; Gilbert & Webb 2007; Põlme *et al.* 2018), and the

sharing is often assumed in studies that explicitly model mutualists and pathogens (Abbott *et al.*

2021; Jiang *et al.* 2020). Although the net antagonistic effect of soil microbes can be common

(and assumed in our model; Lekberg *et al.* 2018), mutualistic effects from microbiomes can

change niche or fitness differences between plants (Kandlikar *et al.* 2019; Ke & Wan 2020; Yan

et al. 2022), affecting population dynamics and fitted interaction coefficients. The pathogenic vs.

mutualistic roles of microbiomes can further interact with the generalist-specialist gradients in

our model (Semchenko *et al.* 2022). For instance, specialist mutualists can amplify fitness

differences between plants and promote monodominance or positive frequency dependence (Xi

et al. 2021); more abundant plants can be more exposed to their specialist pathogens, leading to

conspecific negative density dependence that favors biodiversity (Hülsmann *et al.* 2021; Liang *et*

al. 2016; Sedio & Ostling 2013). Studying the specific roles of microbiomes will greatly improve

our understanding of pairwise and higher-order interactions between plants and mechanisms of

coexistence in highly diverse communities.

Increasing evidence suggests that the plant-soil feedback is a highly dynamic process,

challenging the classic assumption that microbial dynamics are much faster than plant dynamics

such that feedback between plant and microbiome is instantaneous (the "separation of time

scales"; Eppinga *et al.* 2018; Kandlikar *et al.* 2019; Mack *et al.* 2019; Ke & Wan 2020). We

model the change of microbial community compositions over the same time scale as plants and

the decay of plant-associated microbiomes after the senescence of the host. In nature, these

common processes can lead to rich dynamics of microbiomes and strongly affect the strengths of

plant-soil feedback (Chung 2023; Dombrowski *et al.* 2017; Esch & Kobe 2021; Hannula *et al.*

2019, 2021; Ke *et al.* 2021; Ke & Levine 2021; Lepinay *et al.* 2018; Tan *et al.* 2021), and in our

model, they lead to temporal changes in pairwise and higher-order interactions between plants.

Life histories of plants and soil microbes vary greatly across ecosystems. Several

biologically realistic assumptions in our model may therefore only apply to certain communities.

We model large differences in germination phenology, which represents systems where native

and invasive species respond differently to environmental cues (Cleland *et al.* 2015; Wainwright

& Cleland 2013). In some annual plant communities where germination is highly synchronized

(e.g., California annual grassland; Chiariello 1989; Bart *et al.* 2017), the potential for priority

effects and time-dependent interaction modifications may be reduced. We assume fixed lengths

of life cycles among plants. However, the plasticity of life history could lead to novel tradeoffs

between species, affecting the presence and strengths of higher-order interactions

(Kleinhesselink *et al.* 2022; Levine *et al.* 2022). We also assume no mortality or dormancy of

seeds. Including a seed bank can promote coexistence by buffering plants against competition or

environmental stress (Lennon *et al.* 2021), subsequently affecting the strength of time-dependent

interaction modifications. Finally, we assume that all microbiomes are present at the same

density before plants germinate. This fixed baseline does not consider the potential differences in

the life cycles and phenology of soil microbiomes (Rudgers *et al.* 2020) or the potential for

early-germinating plants to associate preferentially with early microbes. Nevertheless, our model

provides an important baseline for future empirical and theoretical works on the temporal

dynamics of complex plant communities.

Our framework of time-dependent interaction modification lays the groundwork for

quantifying temporal changes pairwise and higher-order interactions in multispecies

communities, advancing the understanding of their assembly over time (Song *et al.* 2020, 2021b,

a). However, our results do not directly address the link between higher-order interactions and

coexistence (e.g., Bairey *et al.* 2016; Buche *et al.* 2023; Gibbs *et al.* 2022; Grilli *et al.* 2016).

Resolving higher-order interactions along the temporal axis reveals new processes that could

stabilize or destabilize communities, such as fluctuations of arrival times (phenological

variations) over years (Carter *et al.* 2018; Rudolf 2019; Theobald *et al.* 2017; Zou & Rudolf

2023b), further contributing to the context-dependency of the complexity-stability relationship.

The concept of time-dependent interaction modification combines the two burgeoning fields,

higher-order interactions, and time-explicit ecology. This temporal layer of realism is essential to

understanding the processes shaping complex communities in nature, and to predicting how they

will change as climate change continues to reshuffle species phenology worldwide.

Author Contributions

H.-X.Z. conceived the idea, developed the methods, performed the simulation, and analyzed the results with assistance from X.Y. and V.H.W.R. H.-X.Z. and X.Y. designed the plant-soil feedback model. H.-X.Z. wrote the first draft and all authors contributed to revisions.

Acknowledgments

We thank Po-Ju Ke, Gaurav Kandlikar, and Tom E.X. Miller for comments on the manuscript and members of the Levine Lab at Princeton University for thoughtful discussions. Comments from the editor and three anonymous reviewers greatly improved the manuscript. Funding was provided by NSF DEB-1655626. X.Y. is supported by the Stengl-Wyer Fellowship.

429 **Figures and Tables**

430 Table 1. Parameters used for simulations. All values except for phenology are not plant-specific

431 and follow the general notations in Eqn. 1-4. Note that subscripts x and y can refer to any

432 species, but i, j, and k refer to specific species.

Definition	Symbol	Values
Initial density of plants (as seeds)	$D(0)$	1
Initial and baseline density of microbiomes	$S(0)$	0.1
Length of the year	T	12
Length of plants' life cycle	l	6
Plant mortality rate within the life cycle	d	0.1
Microbiome mortality rate within the year	μ	0.1
Microbiome mortality at the end of the year	$μ_{year}$	0.7
Plant fecundity	λ	3
Total capacity of cultivation (plants to microbiomes)	V	0.3; or drawn from uniform distribution $(0.1, 0.3)$
Cultivation rate from plant x to its associated microbiome	v_{xx}	V_x if no overlap, $0.75V_x$ if small overlap, $0.5V_x$ if high overlap
Cultivation rate from plant x to other plants' associated microbiome	v_{yx} for all plants $x \neq y$	0 if no overlap, $0.25V_x/n$ if small overlap, $0.5V_x/n$ if high overlap; n is the number of plant species other than itself
Effects of microbiomes on plants	m_{xy} for all plants x and y	-0.1 or -0.3; or drawn from uniform distribution $(-0.5, -0.1)$
Phenology of plants i, j and k	p_i, p_j, p_k	1 (earliest) to 6 (latest)
Phenological difference between the focal pair i and j	Δp_{ij}	-5 to 5

Figure 1. A roadmap of the methods. (1) We first construct a plant-soil feedback model (two-

plant model shown for simplicity). With no overlap of associated microbiomes between plants,

each plant can cultivate only its associated microbiome (v_{ii} and v_{jj}), which in turn affects both

plant species (m_{ii}, m_{ij}, m_{ji}, and m_{jj}). With overlap in microbiomes, each plant species can also

cultivate the microbiome associated with the other plant species (v_{ij} and v_{ji}, black arrows);

higher overlap leads to higher interspecific cultivation rates. (2) We then apply the above model

to communities with two and three plants and their microbiomes to simulate plant population

dynamics. Each plant germinates at a certain time, leading to three distinctive relative arrival

times, Δp_{ij}, Δp_{jk}, and Δp_{ik}. (3) Finally, we fit phenomenological models to population dynamics

generated in (2) to evaluate priority effects and time-dependent interaction modifications. In two-

plant communities (i and j; left), trait-dependent priority effects occur when the per-capita

interaction from j to i (α_{ij}) is determined by the relative arrival time between j and i ($f_{ij}(p_i, p_j)$.

In three-plant communities, the presence and arrival time of plant k can determine the interaction

between the other two species ($\beta_{i(jk)} = f_{i(jk)}{}'(p_i, p_j, p_k)$), leading to time-dependent interaction

modification.

Figure 2. Results of two-plant communities with equal microbial effects ($|m_{xx}| = |m_{yx}|$). A.

Population dynamics of the plant-soil feedback model with two plants showing a clear early-

arriver advantage ($\Delta p_{ij} = 4$). Densities of plants and microbiomes are in solid and dashed lines.

Equilibrium densities (after 50 years) of plants and microbiomes are shown in solid and empty

dots. Dynamics of seeds are not shown. B. Fitted interspecific interaction coefficients between

plants i and j along a temporal gradient of relative arrival times Δp_{ij},. A $\Delta p_{ij} < 0$ means that j

arrives early, $\Delta p_{ij} > 0$ means that i arrives early.

Figure 3. Equilibrium population (after 50 years) of three plant species in three-plant

communities with equal microbial effects ($|m_{xx}| = |m_{yx}|$). Vertical red and blue dashed lines

indicate the germination phenology of plants i and j. Scenario I shows plant k germinating

before the focal pair i and j, Scenario II shows k germinating between the focal pair, and

Scenario III shows k germinating after the focal pair. When species arrive at the same time, their

equilibrium population is identical; points and lines of each plant are jittered for visibility.

Figure 4. Time-dependent interaction modification in three-plant communities with equal

microbial effects ($|m_{xx}| = |m_{yx}|$). The higher-order interaction coefficients $\beta_{i(jk)}$ and $\beta_{j(ik)}$ are

fitted according to Eqn. 8. Vertical red and blue dashed lines indicate the germination phenology

of plants i and j. Scenario I shows plant k germinating before the focal pair i and j, Scenario II

shows k germinating between the focal pair, and Scenario III shows k germinating after the focal

pair. When the focal pair germinates at the same time (Scenarios I and III), the two plants are

equivalent, and higher-order interaction coefficients $\beta_{i(jk)}$ and $\beta_{j(ik)}$ are identical; points and

lines of each coefficient are jittered for visibility.

Reference

Abbott, K.C., Eppinga, M.B., Umbanhowar, J., Baudena, M. & Bever, J.D. (2021). Microbiome influence on host community dynamics: Conceptual integration of microbiome feedback with classical host–microbe theory. *Ecology Letters*, 24, 2796–2811.

Adler, F.R., Pearce-Duvet, J.M.C. & Dearing, M.D. (2008). How Host Population Dynamics Translate into Time-Lagged Prevalence: An Investigation of Sin Nombre Virus in Deer Mice. *Bull. Math. Biol.*, 70, 236–252.

Bairey, E., Kelsic, E.D. & Kishony, R. (2016). High-order species interactions shape ecosystem diversity. *Nat Commun*, 7, 12285.

Bart, R.R., Tague, C.L. & Dennison, P.E. (2017). Modeling annual grassland phenology along the central coast of California. *Ecosphere*, 8, e01875.

Bever, J.D., Westover, K.M. & Antonovics, J. (1997). Incorporating the Soil Community into Plant Population Dynamics: The Utility of the Feedback Approach. *The Journal of Ecology*.

Billick, I. & Case, T.J. (1994). Higher Order Interactions in Ecological Communities: What Are They and How Can They be Detected? *Ecology*, 75, 1529–1543.

Blackford, C., Germain, R.M. & Gilbert, B. (2020). Species Differences in Phenology Shape Coexistence. *The American Naturalist*, 195, E168–E180.

Buche, L., Bartomeus, I. & Godoy, O. (2023). Multitrophic higher-order interactions modulate species persistence. *The American Naturalist*.

Carter, S.K., Saenz, D. & Rudolf, V.H.W. (2018). Shifts in phenological distributions reshape interaction potential in natural communities. *Ecology Letters*, 21, 1143–1151.

Chappell, C.R., Dhami, M.K., Bitter, M.C., Czech, L., Herrera Paredes, S., Barrie, F.B., *et al.* (2022). Wide-ranging consequences of priority effects governed by an overarching factor. *eLife*, 11, e79647.

Chiariello, N.R. (1989). Phenology of California Grasslands. In: *Grassland structure and function: California annual grassland*, Tasks for vegetation science (eds. Huenneke, L.F. & Mooney, H.A.). Springer Netherlands, Dordrecht, pp. 47–58.

Chung, Y.A. (2023). The temporal and spatial dimensions of plant–soil feedbacks. *New Phytologist*, 237, 2012–2019.

Clay, P.A., Dhir, K., Rudolf, V.H.W. & Duffy, M.A. (2019). Within-host priority effects systematically alter pathogen coexistence. *American Naturalist*.

Cleland, E.E., Esch, E. & Mckinney, J. (2015). Priority effects vary with species identity and origin in an experiment varying the timing of seed arrival. *Oikos*.

Crawford, K.M., Bauer, J.T., Comita, L.S., Eppinga, M.B., Johnson, D.J., Mangan, S.A., *et al.* (2019). When and where plant-soil feedback may promote plant coexistence: a meta-analysis. *Ecology Letters*, 22, 1274–1284.

Davison, J., Moora, M., Öpik, M., Adholeya, A., Ainsaar, L., Bâ, A., *et al.* (2015). Global assessment of arbuscular
mycorrhizal fungus diversity reveals very low endemism. *Science*, 349, 970–973.

Dickie, I.A., Fukami, T., Wilkie, J.P., Allen, R.B. & Buchanan, P.K. (2012). Do assembly history effects attenuate
from species to ecosystem properties? A field test with wood-inhabiting fungi. *Ecology Letters*, 15,
133–141.

Dombrowski, N., Schlaeppi, K., Agler, M.T., Hacquard, S., Kemen, E., Garrido-Oter, R., *et al.* (2017). Root
microbiota dynamics of perennial Arabis alpina are dependent on soil residence time but
independent of flowering time. *ISME J*, 11, 43–55.

Drake, J.A. (1991). Community-assembly mechanics and the structure of an experimental species ensemble.
American Naturalist, 137, 1–26.

Elzhov, T.V., Mullen, K.M., Spiess, A.-N., Bolker, B., Mullen, M.K.M. & Suggests, M. (2016). Package 'minpack.
lm.' *Title R Interface Levenberg-Marquardt Nonlinear Least-Sq. Algorithm Found MINPACK Plus
Support Bounds.*

Eppinga, M.B., Baudena, M., Johnson, D.J., Jiang, J., Mack, K.M.L., Strand, A.E., *et al.* (2018). Frequency-
dependent feedback constrains plant community coexistence. *Nat Ecol Evol*, 2, 1403–1407.

Esch, C.M. & Kobe, R.K. (2021). Short-lived legacies of Prunus serotina plant–soil feedbacks. *Oecologia*, 196,
529–538.

Fragata, I., Costa-Pereira, R., Kozak, M., Majer, A., Godoy, O. & Magalhães, S. (2022). Specific sequence of
arrival promotes coexistence via spatial niche pre-emption by the weak competitor. *Ecology Letters*,
25, 1629–1639.

Fukami, T. (2015). Historical contingency in community assembly: integrating niches, species pools, and
priority effects. *Annual Review of Ecology, Evolution, and Systematics*.

Garzon-Lopez, C.X., Ballesteros-Mejia, L., Ordoñez, A., Bohlman, S.A., Olff, H. & Jansen, P.A. (2015). Indirect
interactions among tropical tree species through shared rodent seed predators: a novel mechanism
of tree species coexistence. *Ecology Letters*, 18, 752–760.

Gause, G.F. (1934). *The Struggle for Existence: A Classic of Mathematical Biology and Ecology*. Courier Dover
Publications.

Gibbs, T., Levin, S.A. & Levine, J.M. (2022). Coexistence in diverse communities with higher-order interactions.
Proceedings of the National Academy of Sciences, 119, e2205063119.

Gilbert, G.S. & Webb, C.O. (2007). Phylogenetic signal in plant pathogen–host range. *Proceedings of the
National Academy of Sciences*, 104, 4979–4983.

Grainger, T.N., Letten, A.D., Gilbert, B. & Fukami, T. (2019). Applying modern coexistence theory to priority
effects. *Proceedings of the National Academy of Sciences of the United States of America*, 116, 6205–
6210.

Grilli, J., Rogers, T. & Allesina, S. (2016). Modularity and stability in ecological communities. *Nature
Communications*.

Hannula, S.E., Heinen, R., Huberty, M., Steinauer, K., De Long, J.R., Jongen, R., *et al.* (2021). Persistence of plant-
mediated microbial soil legacy effects in soil and inside roots. *Nat Commun*, 12, 5686.

Hannula, S.E., Kielak, A.M., Steinauer, K., Huberty, M., Jongen, R., De Long, J.R., *et al.* (2019). Time after Time:
Temporal Variation in the Effects of Grass and Forb Species on Soil Bacterial and Fungal
Communities. *mBio*, 10, 10.1128/mbio.02635-19.

Hart, S.P., Freckleton, R.P. & Levine, J.M. (2018). How to quantify competitive ability. *Journal of Ecology*, 106,
1902–1909.

Holt, R.D. & Lawton, J.H. (1993). Apparent Competition and Enemy-Free Space in Insect Host-Parasitoid
Communities. *The American Naturalist*, 142, 623–645.

Hoverman, J.T. & Relyea, R.A. (2008). Temporal environmental variation and phenotypic plasticity: A
mechanism underlying priority effects. *Oikos*, 117, 23–32.

Hülsmann, L., Chisholm, R.A. & Hartig, F. (2021). Is Variation in Conspecific Negative Density Dependence
Driving Tree Diversity Patterns at Large Scales? *Trends in Ecology & Evolution*, 36, 151–163.

Inderjit & van der Putten, W.H. (2010). Impacts of soil microbial communities on exotic plant invasions.
Trends in Ecology & Evolution, 25, 512–519.

Inouye, B.D. (2001). Response surface experimental designs for investigating interspecific competition.
Ecology, 82, 2696–2706.

Jiang, J., Abbott, K.C., Baudena, M., Eppinga, M.B., Umbanhowar, J.A. & Bever, J.D. (2020). Pathogens and
mutualists as joint drivers of host species coexistence and turnover: Implications for plant
competition and succession. *American Naturalist*, 195, 591–602.

Jiao, J. & Cortez, M.H. (2022). Exploring How a Generalist Pathogen and Within-Host Priority Effects Alter the
Risk of Being Infected by a Specialist Pathogen. *The American Naturalist*, 200, 815–833.

Kandlikar, G.S., Johnson, C.A., Yan, X., Kraft, N.J.B. & Levine, J.M. (2019). Winning and losing with microbes:
how microbially mediated fitness differences influence plant diversity. *Ecology Letters*, 22, 1178–
1191.

Kardol, P., Cornips, N.J., van Kempen, M.M.L., Bakx-Schotman, J.M.T. & van der Putten, W.H. (2007). Microbe-
mediated plant--soil feedback causes historical contingency effects in plant community assembly.
Ecological monographs.

Ke, P.-J. & Levine, J.M. (2021). The Temporal Dimension of Plant-Soil Microbe Interactions: Mechanisms
Promoting Feedback between Generations. *The American Naturalist*, 198, E80–E94.

Ke, P.-J. & Wan, J. (2020). Effects of soil microbes on plant competition: a perspective from modern
coexistence theory. *Ecological Monographs*, 90, e01391.

Ke, P.-J., Zee, P.C. & Fukami, T. (2021). Dynamic plant–soil microbe interactions: the neglected effect of soil
conditioning time. *New Phytologist*, 231, 1546–1558.

Kharouba, H.M., Ehrlén, J., Gelman, A., Bolmgren, K., Allen, J.M., Travers, S.E., *et al.* (2018). Global shifts in the
phenological synchrony of species interactions over recent decades. *Proceedings of the National*
Academy of Sciences of the United States of America, 115, 5211–5216.

Kleinhesselink, A.R., Kraft, N.J.B., Pacala, S.W. & Levine, J.M. (2022). Detecting and interpreting higher-order
interactions in ecological communities. *Ecology Letters*, 25, 1604–1617.

Lai, H.R., Chong, K.Y., Yee, A.T.K., Mayfield, M.M. & Stouffer, D.B. (2022). Non-additive biotic interactions
improve predictions of tropical tree growth and impact community size structure. *Ecology*, 103,
e03588.

Lekberg, Y., Bever, J.D., Bunn, R.A., Callaway, R.M., Hart, M.M., Kivlin, S.N., *et al.* (2018). Relative importance of
competition and plant–soil feedback, their synergy, context dependency and implications for
coexistence. *Ecology Letters*, 21, 1268–1281.

Lennon, J.T., den Hollander, F., Wilke-Berenguer, M. & Blath, J. (2021). Principles of seed banks and the
emergence of complexity from dormancy. *Nat Commun*, 12, 4807.

Lepinay, C., Vondráková, Z., Dostálek, T. & Münzbergová, Z. (2018). Duration of the conditioning phase affects
the results of plant-soil feedback experiments via soil chemical properties. *Oecologia*, 186, 459–470.

Letten, A.D. & Stouffer, D.B. (2019). The mechanistic basis for higher-order interactions and non-additivity in
competitive communities. *Ecol Lett*, 22, 423–436.

Levine, J.I., Levine, J.M., Gibbs, T. & Pacala, S.W. (2022). Competition for water and species coexistence in
phenologically structured annual plant communities. *Ecology Letters*, 25, 1110–1125.

Levine, J.M., Bascompte, J., Adler, P.B. & Allesina, S. (2017). Beyond pairwise mechanisms of species
coexistence in complex communities. *Nature*, 546, 56–64.

Liang, M., Liu, X., Gilbert, G.S., Zheng, Y., Luo, S., Huang, F., *et al.* (2016). Adult trees cause density-dependent
mortality in conspecific seedlings by regulating the frequency of pathogenic soil fungi. *Ecology*
Letters, 19, 1448–1456.

Mack, K.M.L., Eppinga, M.B. & Bever, J.D. (2019). Plant-soil feedbacks promote coexistence and resilience in
multi-species communities. *PLOS ONE*, 14, e0211572.

Mayfield, M.M. & Stouffer, D.B. (2017). Higher-order interactions capture unexplained complexity in diverse
communities. *Nat Ecol Evol*, 1, 0062.

Nisbet, R.M. (1997). Delay-Differential Equations for Structured Populations. In: *Structured-Population Models*
in Marine, Terrestrial, and Freshwater Systems, Population and Community Biology Series (eds.
Tuljapurkar, S. & Caswell, H.). Springer US, Boston, MA, pp. 89–118.

O'Keeffe, K.R., Simha, A. & Mitchell, C.E. (2021). Indirect interactions among co-infecting parasites and a
microbial mutualist impact disease progression. *Proceedings of the Royal Society B: Biological*
Sciences, 288, 20211313.

Parmesan, C. (2006). Ecological and Evolutionary Responses to Recent Climate Change. *Annual Review of*
Ecology, Evolution, and Systematics.

Põlme, S., Bahram, M., Jacquemyn, H., Kennedy, P., Kohout, P., Moora, M., *et al.* (2018). Host preference and
network properties in biotrophic plant–fungal associations. *New Phytologist*, 217, 1230–1239.

Pu, Z. & Jiang, L. (2015). Dispersal among local communities does not reduce historical contingencies during
metacommunity assembly. *Oikos*, 124, 1327–1336.

van der Putten, W.H., Bradford, M.A., Pernilla Brinkman, E., van de Voorde, T.F.J. & Veen, G.F. (2016). Where,
when and how plant–soil feedback matters in a changing world. *Functional Ecology*, 30, 1109–1121.

R Core Team. (2022). R: A language and environment for statistical computing.

Rasmussen, N.L., Van Allen, B.G. & Rudolf, V.H.W. (2014). Linking phenological shifts to species interactions
through size-mediated priority effects. *Journal of Animal Ecology*.

Rudgers, J.A., Afkhami, M.E., Bell-Dereske, L., Chung, Y.A., Crawford, K.M., Kivlin, S.N., *et al.* (2020). Climate
Disruption of Plant-Microbe Interactions. *Annu. Rev. Ecol. Evol. Syst.*, 51, 561–586.

Rudolf, V.H.W. (2018). Nonlinear effects of phenological shifts link interannual variation to species
interactions. *Journal of Animal Ecology*.

Rudolf, V.H.W. (2019). The role of seasonal timing and phenological shifts for species coexistence. *Ecology*
Letters.

Sedio, B.E. & Ostling, A.M. (2013). How specialised must natural enemies be to facilitate coexistence among
plants? *Ecology Letters*, 16, 995–1003.

Semchenko, M., Barry, K.E., de Vries, F.T., Mommer, L., Moora, M. & Maciá-Vicente, J.G. (2022). Deciphering the
role of specialist and generalist plant–microbial interactions as drivers of plant–soil feedback. *New*
Phytologist, 234, 1929–1944.

Shorrocks, B. & Bingley, M. (1994). Priority effects and species coexistence: Experiments with fungal-breeding
Drosophila. *Journal of Animal Ecology*, 63, 799–806.

Snyder, W.E. & Ives, A.R. (2001). Generalist Predators Disrupt Biological Control by a Specialist Parasitoid.
Ecology, 82, 705–716.

Song, C., Fukami, T. & Saavedra, S. (2021a). Untangling the complexity of priority effects in multispecies
communities. *Ecology Letters*, 24, 2301–2313.

Song, C., Rohr, R.P., Vasseur, D. & Saavedra, S. (2020). Disentangling the effects of external perturbations on
coexistence and priority effects. *Journal of Ecology*, 108, 1677–1689.

Song, C., Uricchio, L.H., Mordecai, E.A. & Saavedra, S. (2021b). Understanding the emergence of contingent and
deterministic exclusion in multispecies communities. *Ecol Lett*, 24, 2155–2168.

Spear, E.R. & Broders, K.D. (2021). Host-generalist fungal pathogens of seedlings may maintain forest
diversity via host-specific impacts and differential susceptibility among tree species. *New Phytologist*,
231, 460–474.

Stenseth, N.C., Falck, W., Chan, K.-S., Bjørnstad, O.N., O'Donoghue, M., Tong, H., *et al.* (1998). From patterns to
648   processes: Phase and density dependencies in the Canadian lynx cycle. *Proceedings of the National*
649   *Academy of Sciences*, 95, 15430–15435.

650   Suding, K.N., Stanley Harpole, W., Fukami, T., Kulmatiski, A., MacDougall, A.S., Stein, C., *et al.* (2013).
651   Consequences of plant–soil feedbacks in invasion. *Journal of Ecology*, 101, 298–308.

652   Suding, K.NasH. & Goldberg, D.E. (1999). Variation in the effects of vegetation and litter on recruitment across
653   productivity gradients. *Journal of Ecology*, 87, 436–449.

654   Tan, J., Kerstetter, J.E. & Turcotte, M.M. (2021). Eco-evolutionary interaction between microbiome presence
655   and rapid biofilm evolution determines plant host fitness. *Nat Ecol Evol*, 5, 670–676.

656   Theobald, E.J., Breckheimer, I. & HilleRisLambers, J. (2017). Climate drives phenological reassembly of a
657   mountain wildflower meadow community. *Ecology*, 98, 2799–2812.

658   Tilman, D. (1982). Resource competition and community structure. *Monographs in population biology*.

659   Uricchio, L.H., Daws, S.C., Spear, E.R. & Mordecai, E.A. (2019). Priority effects and nonhierarchical competition
660   shape species composition in a complex grassland community. *American Naturalist*, 193, 213–226.

661   Wainwright, C.E. & Cleland, E.E. (2013). Exotic species display greater germination plasticity and higher
662   germination rates than native species across multiple cues. *Biological Invasions*.

663   Werner, E.E. & Peacor, S.D. (2003). A Review of Trait-Mediated Indirect Interactions in Ecological
664   Communities. *Ecology*, 84, 1083–1100.

665   Wootton, J.T. (1993). Indirect Effects and Habitat Use in an Intertidal Community: Interaction Chains and
666   Interaction Modifications. *The American Naturalist*, 141, 71–89.

667   Wootton, J.T. (1994). The nature and consequences of indirect effects in ecological communities. *Annual*
668   *Review of Ecology and Systematics*.

669   Xi, N., Adler, P.B., Chen, D., Wu, H., Catford, J.A., van Bodegom, P.M., *et al.* (2021). Relationships between plant–
670   soil feedbacks and functional traits. *Journal of Ecology*, 109, 3411–3423.

671   Yan, X., Levine, J.M. & Kandlikar, G.S. (2022). A quantitative synthesis of soil microbial effects on plant species
672   coexistence. *Proceedings of the National Academy of Sciences*, 119, e2122088119.

673   Zhang, Z., Liu, Y., Brunel, C. & van Kleunen, M. (2020). Soil-microorganism-mediated invasional meltdown in
674   plants. *Nature Ecology & Evolution*, 1–10.

675   Zou, H.-X. & Rudolf, V.H.W. (2023a). Bridging theory and experiments of priority effects. *Trends in Ecology &*
676   *Evolution*.

677   Zou, H.-X. & Rudolf, V.H.W. (2023b). Priority Effects Determine How Dispersal Affects Biodiversity in Seasonal
678   Metacommunities. *The American Naturalist*, 202, 140–151.
679

1. Construct the mechanistic plant-soil feedback model

2. Change the phenology of plants, simulate population dynamics

3. Fit phenomenological models to population dynamics

Two-plant community,
pairwise priority effects

Three-plant community,
pairwise priority effects and
time-dependent interaction modifications