PAH DEFICIENT PATHOLOGY IN HUMANIZED c.1066-11G>A PHENYLKETONURIA MICE

Ainhoa Martínez-Pizarro[a,c,d], Sara Picó[a,e,1], Arístides López-Márquez[a,c,2], Claudia Rodriguez-López[a,e], Elena Montalvo[a,c,d], Mar Alvarez[a,c,d], Margarita Castro[b,c,d], Santiago Ramón-Maiques[c,f], Belén Pérez[a,b,c,d], José J Lucas[a,e], Eva Richard[a,b,c,d], Lourdes R Desviat[a,b,c,d]

[a]Centro de Biología Molecular Severo Ochoa UAM-CSIC, Universidad Autónoma de Madrid, Madrid, Spain

[b] Centro de Diagnóstico de Enfermedades Moleculares (CEDEM)

[c]Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), ISCIII

[d]Instituto de Investigación Sanitaria Hospital La Paz (IdiPaz)

[e]Centro de Investigación Biomédica en Red de Enfermedades Neurodegenerativas (CIBERNED), ISCIII

[f]Instituto de Biomedicina de Valencia (IBV-CSIC), Valencia, Spain

[1]*present address:* Department of Biomedical Sciences, Faculty of Biology and Medicine, University of Lausanne, Lausanne, Vaud, Switzerland.

[2] *present address:* Institut de Recerca de Sant Joan de Déu, Barcelona, Spain

Correspondence should be addressed to:

Lourdes R Desviat

Centro de Biología Molecular Severo Ochoa, Universidad Autónoma de Madrid

Nicolás Cabrera 1, 28049 Madrid, Spain

Phone: 34-911964566

Email: lruiz@cbm.csic.es; lourdes.ruiz@uam.es

Abstract

We have generated using CRISPR/Cas9 technology a partially humanized mouse model of the neurometabolic disease phenylketonuria (PKU) carrying the prevalent splice intronic variant *PAH* c.1066-11G>A. This variant creates a strong alternative splice acceptor site, leading to the inclusion of 9 nucleotides coding for a 3 amino acid insertion between Q355 and Y356 of the phenylalanine hydroxylase (PAH) protein. Homozygous *Pah* c.1066-11A mice, with a partially humanized intron 10 sequence with the variant, accurately recapitulate the splicing defect and present almost undetectable hepatic PAH protein or activity. They exhibit fur hypopigmentation, lower brain and body weight and reduced survival. Blood and brain phenylalanine (Phe) levels are elevated, along with decreased tyrosine (Tyr), tryptophan (Trp) and monoamine neurotransmitters levels. *Pah* c.1066-11A mice present behavioral deficits, mainly hypoactivity and diminished social interaction, as well as locomotor deficiencies. Abnormal hind-limb clasping reflex and microophtalmia are observed. Immunohistochemistry analyses of GFAP and Iba1 markers revealed changes in the morphology of glial cells and a significant increase in GFAP and Iba1 staining signals. Western blot analyses in mouse liver lysates and in gene edited HepG2 cells with the variant showed reduced levels of chaperones DNAJC12 and HSC70/HSP70 and an increase in autophagy markers LAMP1 and LC3BII, suggesting that the mutant pQ355_Y356insGLQ PAH protein could aggregate complexed with DNAJC12 and HSP70 proteins, with the aggregates being processed by autophagy. This humanized model corresponding to a severe, classical PKU phenotype represents a useful tool for pathophysiology research in PKU and for novel therapies development.

Keywords

phenylketonuria; PKU; phenylalanine hydroxylase; PAH; humanized mouse model; splice mutation; CRIPSR/Cas

Introduction

Phenylketonuria (PKU; OMIM#261600) is probably the best studied and the most frequent inborn error of amino acid metabolism, caused by autosomal recessive deficiency of the hepatic enzyme phenylalanine hydroxylase (PAH; EC 1.14.16.1), catalysing the (6R)-5,6,7,8-tetrahydrobiopterin (BH_4) dependent hydroxylation of L-phenylalanine (Phe) to L-tyrosine (Tyr). The global PKU prevalence was recently estimated to be approximately 1:24,000 newborns (Hillert et al., 2020), with a documented large frequency variability depending on the population and geographical location, being around 1:10,000 in USA and parts of Europe (Hillert et al., 2020). In PAH deficiency, high Phe levels cause brain dysfunction in untreated patients, exhibiting intellectual disability, seizures and behavioural problems, although a broad clinical spectrum exists, from severe, classical PKU to the mildest form, mild hyperphenylalaninemia (HPA) with normal brain development (van Spronsen et al., 2021). Early diagnosis is based on newborn screening and implementation of Phe-restricted dietary treatment, which has been the gold standard for many decades (Adams et al., 2023). This prevents major neurological alterations, although significant challenges remain, associated to poor adherence and suboptimal outcomes. In the last years, new treatment options have become available, a synthetic form of BH_4 (sapropterin, Kuvan®) which allows a less restrictive or absence of diet for a group of patients with mild PKU forms, and a pegylated form of phenylalanine ammonia lyase, indicated for adult patients (van Spronsen et al., 2021). Other treatments are in preclinical or clinical development, such as mRNA, gene and gene editing therapies, or use of a probiotic *Escherichia coli* Nissle strain, engineered to overexpress enzymes metabolizing gut Phe (Chen et al., 2022).

In order to investigate PKU pathophysiology and as tools for preclinical novel treatment testing, several mouse models have been generated. First models were obtained by N-ethyl-N-nitrosourea germline mutagenesis, with *Enu1* mice representing a mild HPA phenotype and *Enu2* and *Enu3* as models for severe, classical PKU (Haefele et al., 2001; McDonald and Charlton, 1997). However, the murine mutations present in these models do not correspond to human variants, and, with the advent of gene editing technologies, mouse "avatar" models carrying variants identified in human patients (p.R261Q, p.R408W) have now been generated (Aubi et al., 2021; Li et al., 2021). However, there are no mouse models with *PAH* splicing variants, despite their frequency in the PKU mutational spectrum. The second most frequent variant worldwide is c.1066-11G>A (global allele frequency 6.4%) (Hillert et al., 2020), which is often called the "Mediterranean mutation," as it accounts for the majority of the mutant alleles in Southern Europe (11%), and is also common in Latin America (9,4%). The

homozygous genotype c.[1066-11G>A];[1066-11G>A] is the second most prevalent genotype worldwide (2,6% globally) and the first in Southern Europe (5,2%) (Hillert et al., 2020), associated to a classical, non BH$_4$-responsive severe phenotype (www.biopku.org).

The *PAH* c.1066-11G>A variant, located in intron 10, creates a strong splice acceptor site used instead of the natural one, leading to the aberrant in-frame insertion of 9 nucleotides in the mature mRNA, coding for three extra amino acids between residues 355 and 356 of the protein (pQ355_Y356insGLQ). The initial study reported the presence of immunoreactive protein in a liver biopsy from a homozygous patient but total absence of catalytic activity (Dworniczak et al., 1991). Expression analysis in a prokaryotic system of the mutant protein (with the 3 extra amino acids) fused to maltose binding protein showed that it was recovered exclusively as high molecular weight aggregates, with no catalytic activity (Pey et al., 2003). The exact liver pathomechanism associated to the mutant protein is yet unknown, stressing the need of finding an adequate *in vivo* model in which to test potential therapies.

The fact that intronic sequences are not conserved between mouse and human, and that, even with an equivalent nucleotide change, the splicing outcome may be different due to the distinct genomic context of the mutation (Rivera-Barahona et al., 2015), hinders the precise modelling of splicing defects in mice. To that end, based on the interest in the prevalent c.1066-11G>A variant, in this work we have used CRISPR/Cas9 technology to generate a novel PKU mouse model with a partially humanized *Pah* intron 11 carrying the variant. After confirming the altered *Pah* splicing profile in liver of these so called *Pah* c.1066-11A *knock in* (*KI*) mice, accurately recapitulating the human molecular defect, we have characterised their biochemical, histological and behavioural phenotypes. We have also used molecular approaches to investigate pathological mechanisms underlying mutant protein expression in liver. In parallel, we have generated a stable isogenic cellular model of the splicing variant through CRISPR/Cas gene editing. To his end, since PAH is expressed mainly in liver, we decided to use HepG2 cells as the most suitable cell line to study the characteristics of the mutant protein in a human genetic background. The novel mouse avatar model and the edited HepG2 cell line, both carrying the c.1066-11A variant corresponding to a frequent classical PKU phenotype, are ideally suited for preclinical research and development of RNA therapies targeting the splicing defect or pharmacological compounds targeting the mutant protein.

Methods

Cell culture and treatments. Human hepatoma cell lines were grown in Minimum Essential Medium (MEM, Sigma Aldrich) supplemented with 5% fetal bovine serum (FBS), 1% glutamine and 0.1% antibiotic mix (penicillin/streptomycin) under standard cell culture conditions (37ºC, 95% relative humidity, 5% CO_2).

Minigenes construction. For evaluation of *in vitro* splicing, a minigene construction including *PAH* exon 10, full intron 10, exon 11, 1958 bp of intron 11, and exon 12 cloned in pcDNA3.1+ (Heintz et al., 2012) was used. A mutant minigene with the c.1066-11G>A variant was generated by site-directed mutagenesis with QuikChange Lightning Kit (Agilent Technologies, Santa Clara, CA, USA). Sanger sequencing confirmed the identity of the constructs. To study the effect of humanizing 90 bp of the 3' end of intron 10 of the mouse *Pah* gene containing the c.1066-11G>A mutation for the subsequent generation of the murine model, a hybrid minigene was constructed with this design and including from *Pah* exon 10 to exon 12, with a shortened intron 11. The sequence of interest, with a total length of 2532 base pairs, was synthesized and cloned into pcDNA3.1+ between the restriction enzymes KpnI and XhoI, using the GeneArt service (ThermoFisher Scientific, USA).

Transient transfection and splicing analysis. For minigene assays, Hep3B cells were seeded in six-well plates at a density of $4x10^5$ in 2 ml 10% MEM and grown overnight. Cells were transfected with a total DNA amount of 2 µg per well using lipofectamine 2000 (Invitrogen) and harvested by trypsinization after 48 h. Total RNA was extracted using Trizol Reagent (ThermoFisher) and phenol-chloroform. cDNA synthesis was performed using NZY First-Strand cDNA Synthesis Kit (NZYtech). Splicing analysis was carried out by PCR amplification with FastStart Taq Polymerase (Roche) using specific primers (Table 1). End-point PCR amplification products were analyzed by 3% agarose gel electrophoresis.

Generation of the *KI* cell line HepG2 *PAH* c.1066-11A. The *KI* cell model was generated via CRISPR/Cas9 using Breaking Cas software (https://bioinfogp.cnb.csic.es/tools/breakingcas/) (Oliveros et al., 2016) to design guides (crRNA) with high specificity to introduce the c.1066-11G>A variant in the PAH gene and to identify potential off-targets (Table 2). To generate the sgRNA complexes, crRNA and tracrRNA (Integrated DNA Technologies, IDT) were mixed in equimolar concentrations. The ribonucleoprotein complex (RNP) was formed by the combination

of 24 pmoles of Cas9 (Streptococcus pyogenes, IDT) and 24 pmoles of sgRNA. Approximately $6x10^5$ HepG2 cells per well were seeded in a 6-well plate with the RNP complex and 2.4 pmoles of the DNA template for homology driven repair (HDR) (Ultramer oligonucleotide, IDT) (Table 2) using Lipofectamine CRISPRMAX™ (ThermoFisher Scientific Waltham). After 48h, the culture media was changed to fresh media. Three days post-transfection, cells were dispersed at low density into 150 mm dishes in MEM medium. About 15-20 days later, large colonies were picked and expanded. Genomic DNA extraction of different clones was carried out using QIAamp DNA Mini Kit (Qiagen). DNA concentration was measured using Nanodrop One spectrophotometer (ThermoFisher Scientific). The region surrounding the desired change was amplified by PCR and subsequently digested with the restriction enzyme DdeI (New Englands Biolabs) to detect positive clones, which were confirmed by Sanger sequencing. The three predicted highest risk off-target sites were also amplified using specific primers (Table 2) and sequenced.

Generation of humanized *KI* mouse *Pah* c.1066-11A. PKU humanized mice were generated in a 50% C57BL/6J and 50% CBA/CA background via CRISPR/Cas9-mediated gene editing, by the institutional core facility (CNB-CBM mouse transgenesis). The CRISPR sgRNA (300 ng/mL) was co-injected into fertilized oocytes with HiFi Cas9 protein (600 ng/mL) and DNA Template (400 ng/mL) (Table 3), all of them synthesized and purchased from IDT. The DNA template harbored 90 base pairs of the 3' end of human *PAH* intron 10 containing the c.1066-11G>A pathogenic variant flanked by mouse *Pah* intron 10 sequence. One animal was found to be heterozygous for the c.1066-11G>A variant and the correct wild-type sequence from a total of 14 pups born. Confirmation of the on-target and the verification of the absence of the four highest risk off-target sites were performed from genomic DNA tail biopsies by PCR amplification with specific primers (Table 3) and subsequent Sanger analysis.

Mouse husbandry and colony expansion. All animal studies were approved by the Institutional Ethical Committee for Animal Experimentation (Universidad Autónoma de Madrid, reference CEI 963-A026) and by the Regional Environment Department (Comunidad de Madrid, reference PROEX 194/19). Animal housing and maintenance protocols followed the local authority guidelines. Mice were maintained on standard chow. All the experiments were carried out in a pathogen-free environment at the animal facility of the Centro de Biología Molecular Severo Ochoa, in accordance with the Spanish Law of Animal Protection. Food and water were available ad libitum, and

mice were maintained in a temperature-controlled environment on a 12/12 h light-dark cycle.

The colony was backcrossed up to F10 with wild-type mice (C57BL/6J). Heterozygous mice were used for breeding and the resultant homozygous mutant *Pah* c.1066-11A and wild-type (WT) mice have been used in the experiments presented in this work. All mice used were adults (3–5 months-old) and males and females were included in all experiments. Body weight was monitored every week. At the time/age stated for each experiment, mice were sacrificed in a carbon dioxide chamber and perfused with 50 ml of NaCl. Immediately after, brain and liver were surgically excised, snap-frozen in liquid nitrogen and stored at –80 °C.

BH$_4$-loading test. A treatment solution of BH$_4$ (Sigma Aldrich, T4425) (2mg/ml BH$_4$, 2% ascorbic acid, and 10% DMSO; in a dosage of 20 mg BH4/kg body weight) or placebo solution (2% ascorbic acid and 10% DMSO), was injected daily intraperitoneally into PKU mice (n = 5 mice per group) in 2 doses for 4 days. Blood samples were collected onto filter paper cards after 3 h fasting, before the first injection and during the 4 days, and Phe was quantified by fluorimetry using Neonatal Phenylalanine kit (Labsystems Diagnostics).

Amino acid and neurotransmitters determination in brain. For analysis of amino acid concentrations in brain, half-brains were homogenized using Ultra-Turrax device (IKA) in 5 volumes/weight 10% trichloroacetic acid containing norvaline (150 µM). The homogenates were clarified by centrifugation at 500 g, 5 min at 4$\square$C and supernatants stored at –80°C until analysis by ion exchange chromatography with ninhydrin. For neurotransmitters analysis half-brains were processed by adding 3 volumes/weight of a lysis buffer solution (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.1% Triton X-100, 1 mM EDTA, and 10% glycerol) with protease inhibitor. Analysis of biogenic amine neurotransmitters metabolites, 5-hydroxyindolacetic acid (5-HIAA), homovanillic acid (HVA), 3-metoxy-hydroxy-phenylglycol (MHGP), 3-O-methyldopa (3-OMD), 5-hydroxytryptophan (5-OH-TRP), dihydroxyphenylacetic acid (DOPAC), dopamine and serotonin in brain lysates was performed by ion pair high-performance liquid chromatography (HPLC) with electrochemical detection (ESA, Coulocem III) as reported (Ormazabal et al., 2005). Briefly, samples were deproteinized with VWR Centrifugal Filters, diluted 1:2 in the chromatography mobile phase and 50µl were injected into HPLC. Measured brain homogenate monoamine neurotransmitter concentrations were corrected for the protein content of the homogenate and expressed as pmol/mg protein.

Amino acid determination in serum. In order to obtain serum specimens, whole blood was collected by cardiac puncture and transferred to an Eppendorf tube. Subsequently, blood samples were left for 1 h at 37ºC and at 4ºC overnight to facilitate coagulation, followed by two consecutive centrifugation steps (1200 g, 5 min, and 4 °C) after which the respective supernatants were pipetted out to a clean tube. The isolated serum fractions were stored at –80 °C and amino acids measured by ion exchange chromatography (visualized with ninhydrin).

PAH enzymatic activity assay. For PAH enzyme activity determination, cells were lysed by adding Na-Hepes buffer (50 mM HEPES pH 7.0, 500 mM NaCl) followed by three freeze-thawing cycles. Proteins were collected in the supernatant fraction after centrifugation 10 min at 16,000 g at 4ºC. Liver lysates were prepared with Na-Hepes buffer, shredding the tissue with ultra-turrax and centrifuged at 16000 g for 30 min. PAH activity in the homogenates was measured at room temperature (RT≈25°C) using 10 μg of total protein in each assay, with 1 mM L-Phe in Na–Hepes pH 7.0, containing catalase (0.1 mg/ml). After 5 min preincubation at 25°C, ferrous ammonium sulfate (100 μM) was added, and the reaction was triggered after 1 min by adding 75 μM BH_4 and 5 mM DTT (final concentrations in the assay). The reactions were allowed to run for 30 min at RT and stopped with 3 volumes of 12% perchloric acid, followed by incubation at 4°C for 1 h and at -20°C overnight. The mix was centrifuged at 16000 g, 15 min at 4°C and supernatants stored at –20°C until analysis. Under these conditions, PAH activity was linear to the amount of protein in the extracts. L-Tyr formed was quantified by HPLC with fluorometric detection as described previously (Allen et al., 1999).

Cells, liver and brain protein analysis. HepG2 cells were harvested by trypsinization and treated with RIPA lysis buffer (10 mM Tris-HCl 7.5, 150 mM NaCl, 0.1% Triton, and 10% glycerol), protease and phosphatase inhibitors. Samples were subjected to three freeze-thawing cycles and proteins were collected in the supernatant fraction after centrifugation at 16,000 g for 10 min at 4°C. Mouse liver and brain proteins were isolated by disrupting the tissue in lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.1% Triton X-100, 1 mM EDTA, and 10% glycerol) using ultra-turrax followed by centrifugation at 16,000 *g* for 30 min at 4°C. Protein concentration in the supernatant was measured using the Bradford method (Bio-Rad Laboratories).For western blot analysis, equal amounts of protein (50 μg) were loaded into 4-12% SDS-polyacrylamide gels. Proteins were transferred into a nitrocellulose membrane using the iBlot Dry Blotting System (Thermo Fisher Scientific). For the analysis of total

proteins, membranes were blocked for 1 hour with 5% nonfat milk, in 0.1% TBS-tween and incubated overnight with the corresponding primary antibody: PAH (1:1000, Santa Cruz Biotechnology, sc-271258), Ubiquitin (1:1000, Santa Cruz Biotechnology, sc-8017), Hsp70 (1:1000, Novus Biological, NB110-61582), DNAJC12 (1/1000, Abcam, ab167425), LC3B (1:1000, Cell Signaling Technology, #2775), LAMP1 (1:1000, Cell Signaling Technology, #3243) and p62/SQSTM1 (1:5000, Novus biologicals, H00008878-M01). Secondary antibodies were horse anti-mouse (1:2000, Cell Signaling Technology, 7076S) and goat anti rabbit (1:5000, Cell Signaling Technology,7074S). Antibody against GAPDH (1:5000, Abcam, ab8245) and β-actin (1:2000, Origene, TA811000S) was used as loading control. Enhanced chemiluminescence reagent (ECL, Cytiva) or SuperSignal West Femto (ThermoFisher) were used for protein detection. Band intensity for each protein was quantified with BioRad GS-900 Densitometer (Bio-Rad) and ImageLab program.

mRNA quantification. For gene expression analysis, cDNA was obtained by retrotranscription of 500 ng of total RNA from mouse liver or HepG2 cells using NZY First-Strand cDNA synthesis kit (NZYTech). *Pah, Hsp70, Dnajc12*, genes were amplified with specific primers (Table 1), using Perfecta SYBR Green FastMix kit (Quanta Biosciences) in a CFX Opus 384 (Bio-Rad). *Gapdh* was used as endogenous control and quantification was done using the $2^{-\Delta\Delta Ct}$ method.

Immunohistochemistry. Brain hemispheres were extracted and fixed overnight in 4% paraformaldehyde, cryopreserved for 72h in 30% sucrose in PBS and included in optimum cutting temperature (OCT) compound (TissueTek, Sakura Finetek Europe, 4583), immediately frozen and stored at –80 °C until use. Using a cryostat (Thermo-Fisher Scientific) 30 μm sagittal sections were cut, collected and stored free floating at –20 °C in glycol containing buffer (30% glycerol, 30% ethylene glycol in 0.02 M phosphate buffer). Mouse sections were washed in PBS, immersed in 0.3% H_2O_2 in PBS for 45 min and blocked for 1h in blocking solution (PBS containing 0.5% fetal bovine serum, 0.3% Triton X-100 and 1% BSA) and then incubated overnight at 4°C with the primary antibody diluted in blocking solution. After washing, sections were incubated with biotinylated anti-mouse or anti-rabbit secondary antibody and then with avidin-biotin complex using the Elite Vectastain kit (Vector Laboratories, PK-6101 and PK-6102). Chromogen reactions were performed using diaminobenzidine (SIGMAFAST DAB, Sigma-Aldrich, D4293) for 10 min. Sections were mounted on glass slides and coverslipped with Mowiol (Calbiochem, 475904). An Axioskop2 plus (Zeiss) microscope with an DMC6200 camera (Leica) was used to capture the images.

Transmission electron microscopy. Mice were completely anaesthetized with an intraperitoneal tiopental injection (100 µl/10 g), perfused transcardially with 50 ml of NaCl and then with a solution of 4% formaldehyde and 2 % glutaraldehyde in 0.1M phosphate buffer pH 7.4 Spinal cords were postfixed in the same solution for 2 h at room temperature and overnight at 4ºC. Spinal cord lumbar fragments were postfixed in 1% osmium tetroxide in 0.1 M cacodylate buffer, dehydrated in ethanol and embedded in Epon-Araldite. Serial ultrathin sections of the lumbar transverse section of the spinal cord were collected on pioloform-coated, single-hole grids, and stained with uranyl acetate and lead citrate. The sections were analysed with a JEM-1010 transmission electron microscope (Jeol, Japan) equipped with a side-mounted CCD camera Mega View III from Olympus Soft Imaging System GmBH (Muenster, Germany). Myelinated fibers were sampled randomly in the white matter tracts of the spinal cord at a magnification of 3000X.

Behavioral testing. Open field test: Locomotor activity was measured in clear Plexiglas® boxes measuring 27.5cm x 27.5cm, outfitted with photo-beam detectors for monitoring horizontal and vertical activity. Activity was recorded with a MED Associates' Activity Monitor (MED Associates) and were analyzed with the MED Associates' Activity Monitor Data Analysis v.5.93.773 software. Mice were placed in the center of the open-field apparatus and left to move freely. Data were individually recorded for each animal during 15 min. Ambulatory walked distance and velocity were measured. Rotarod test: Motor coordination was measured in an accelerating rotarod apparatus (Ugo Basile). Mice were pre-trained during two days at a constant speed, the first day: 4 trials of 1 min at 4 rpm and the second day: 4 trials of 1 min at 8 rpm. On the third day, rotarod was set to accelerate from 4 to 40 rpm over 5 min and mice were tested four times. During accelerating trials, the latency to fall from the rod was measured. Inverted grid test: Muscle strength was examined by placing the mice in the center of a wire grid that was rotated to an inverted position over 10 s and held steadily 50 cm above a padded surface. Latency to fall from the grid was measured (maximum: 300 s, two trials). Social approaching test: Test was conducted in a Plexiglas three-chambered box with openings between the chambers. The first day (habituation), mice were allowed to explore for 10 min the empty box. The next day (test), mice were placed in the center of the box, containing two wire cages placed in opposite corners of the lateral chambers, one empty ("Empty cage") and the other with an unknown (gender paired) mouse on it ("Occupied cage"). Mice were recorded for 10 min and the time spent in each chamber was measured. Preference index was measured as [time in Occupied cage – time in Empty cage]/[time in Empty cage + time in Occupied

cage]*100. <u>Vertical pole</u>: Mice were placed in the top of a 50 cm high vertical 1 cm diameter pole. The pole is mounted on a triangular base stand and placed in the home cage so that mice might prefer to descend to the floor of cage. Mice were trained once to turn around and descend the pole. Then, the animal is placed again in the top of the pole and the recording starts when the animal begins the turning movement. The time to turn completely downward and total time to descend to the floor are measured.

In silico structural analysis

The inserted sequence GLQ was first modeled by replacing residues [356]YCL[358] in the crystal structure of human PAH (PDB entry 6HYC) using COOT (Emsley et al., 2010) and selecting rotamers for side chains that diminished steric clashes with nearby elements. Next, we used AlphaFold (Jumper et al., 2021) and ColabFold (Mirdita et al., 2022) to predict the structure of the human protein with the wild-type sequence (UniProt P00439) or bearing the 3 amino acid insertion. The analysis of the predicted model was done with COOT. Figures were prepared with PyMOL (Schrödinger, LLC).

Data analysis

All results are expressed as mean ± SEM, unless stated otherwise. The unpaired Student´s *t* test was used to evaluate the significance of differences between groups. Critical values for significance of $P<0.05$, $P<0.01$ or $P<0.001$ were used throughout the study.

Results

The c.1066-11G>A PAH variant minigene mimics disease-associated mis-splicing

Bioinformatic analysis of *PAH* intron 10 sequence confirmed the creation of a 3' splice acceptor site with variant c.1066-11G>A (MaxEnt score 5.5), which is predicted to be used instead of the natural splice site (MaxEnt score 3.2). Functional analysis in the pcDNA3.1 minigene (Martinez-Pizarro et al., 2018) confirmed the splicing defect (Figure 1A). As previously reported, residual exon 11 skipping is observed in the WT minigene, due to a naturally weak 3' splice site and a suboptimal 5' splice site (Martinez-Pizarro et al., 2018) (Figure 1A).

Lack of PAH in c.1066-11G>A variant-engineered HepG2 cells

We next studied the expression of the mutant protein in a human hepatic context. To that end, we used CRISPR/Cas9 to introduce the c.1066-11G>A variant in

HepG2 cells. After confirming the presence of the variant in homozygous fashion in the selected clone and the resulting splicing defect by RT-PCR (data not shown), western blot analysis showed almost no detectable immunoreactive protein (Figure 1B), which correlated with total absence of catalytic PAH activity (non-detectable).

The humanized *Pah* c.1066-11g>a mouse model exhibits a classical PKU phenotype

The *KI Pah* c.1066-11A mouse model was generated using CRISPR/Cas9 and HDR gene editing. Examination of intron 10 sequence revealed no similarity between murine and human sequences, as expected. In fact, the GGGG sequence where the variant is located is not present in the murine sequence. Therefore, our strategy was to replace murine intron 10 sequence, adjacent to exon 11, with the human sequence carrying the variant. However, humanization of splicing sequences in a murine genomic context can have undesirable effects on the splicing outcome, as previously reported (Barman-Aksözen et al., 2017; Vázquez-Domínguez and Garanto, 2022). In order to confirm the correct recognition of the human sequence in the murine context and the recapitulation of the splicing defect, we first validated the strategy using minigenes, reproducing the construct designed for the mouse gene editing experiment. Murine exon 10, a partially humanized intron 10, murine exon 11, a shortened intron 11 and exon 12 were cloned in the pcDNA3.1 vector. This minigene in its WT version resulted in a correctly spliced exon 10-11-12 transcript and a transcript corresponding to exon 11 skipping. The mutant (c.1066-11A) hybrid minigene accurately recapitulated the splicing defect, with a sole transcript in which we confirmed the insertion of 9 extra nucleotides due to the use of the newly created splice site (Supplementary Figure 1A, B).

The humanized mouse model homozygous for the c.1066-11A variant was generated by CRISPR/Cas9 gene editing using the same design and the heterozygous founder mouse was identified by PCR genotyping and subsequent sequencing analysis (Supplementary Figure 1C). The founder mouse was backcrossed to C57BL6/J background up to 10 generations. Mating of heterozygous mice yielded the expected mendelian ratios of wild-type (WT), heterozygous and homozygous *Pah* c.1066-11A offspring. In all experiments with the newly generated *Pah* c.1066-11A mice, age and sex-matched WT littermates were used as the control group. RT-PCR from liver RNA samples of homozygous mice confirmed the splicing defect resulting in the aberrant insertion of extra 9 nucleotides in the mature mRNA (Supplementary Figure 1D). Normal *Pah* mRNA levels were confirmed by RT-qPCR analysis (data not shown).

Pah c.1066-11A mice exhibited high mortality at weaning and lethal seizures at 6-7 months of age, thus resulting in reduced survival (Figure 2A). They display reduced body weight and the fur hypopigmentation characteristic of mouse PKU models (Figure 2 B, C). Besides, most *Pah* c.1066-11A mice exhibit microphthalmia (Figure 2C) which has previously been reported in other PKU models with a severe phenotype and correlates with the ocular abnormalities observed in untreated PKU patients (Hopf et al., 2020; Singh et al., 2021).

Western blot analysis in liver samples showed nearly undetectable hepatic PAH protein in *Pah* c.1066-11A PKU model (Figure 2D). Accordingly, lack of hepatic PAH activity was confirmed (Figure 2E). Heterozygous mice exhibited slightly reduced levels of PAH protein (76%) and intermediate-low PAH activity (23%).

Biochemical analyses revealed high blood serum Phe (1885.7±71.2 µM *versus* 59.3±2.5 µM for WT) and low Tyr levels (28.8±1.6 µM *versus* 68.2±4.4 µM for WT), corresponding to severe, classical PKU. Female mice exhibited higher blood Phe levels as compared to males (2126.8±35 vs 1135 ± 53 µM). Sex differences in blood Phe levels have been previously reported in PKU mouse models (Winn et al., 2018), although they have not been described in PKU patients. In brain, Phe levels were also high in both sexes of the *Pah* c.1066-11A PKU model (153.4±3.9 µM *versus* 13.6±0.7 µM for WT) and Tyr levels were low (6.4±0.3 µM *versus* 14.6±1.2 µM for WT). Among all the remaining analysed amino acids, cystine was found significantly elevated in the *Pah* c.1066-11A mice serum, and lower levels were found for tryptophan, asparagine, glutamine, glycine, proline, alanine, methionine, ornithine, lysine and arginine. In brain, serine, glycine, lysine, histidine, and cystathionine were found elevated while glutamine, leucine and isoleucine were found in reduced levels in *Pah* c.1066-11A mice (Supplementary Tables 1 and 2). No significant differences were observed between males and females.

Monoamine neurotransmitters and related metabolites (dopamine, serotonin, DOPAC, 5-OHTrp, HVA and 5-HIAA) were significantly lower in brain of homozygous *Pah* c.1066-11A mice of both sexes as compared to WT mice (Figure 3).

We also examined the response to BH_4 treatment in the novel PKU mouse model. BH_4 was administered at 10 mg/kg body weight twice a day for four consecutive days and Phe levels measured in dried blood spots sampled daily after 3 hours fasting. As seen in Supplementary Figure 2, no significant changes in blood Phe content were detected in BH_4 treated PKU mice compared to those receiving placebo (the same solution without BH_4).

Humanized *Pah c.1066-11A* mice exhibit neurological alterations

Different behavioural tests were carried out in 12–14-week-old mice (Figure 4). First, mice were analysed in the open field test to evaluate their general locomotor activity. *Pah* c.1066-11A mice showed decreased total ambulatory distance covered, as well as decreased average velocity, indicative of a general hypoactivity phenotype in these mice (Figure 4A). Curiously, the decreased ambulatory activity is more pronounced in the centre of the arena, what might be suggestive of anxiety-like behaviour (Figure 4A). Motor coordination and neuromuscular fitness were also assessed in the rotarod and inverted grid tests, respectively. In the rotarod, *Pah* c.1066-11A mice showed a significant decrease in their latency to fall from the accelerating rod, compared to WT mice, thus evidencing a deficit in motor coordination (Figure 4B). The latency to fall was also markedly decreased in the inverted grid test (Figure 4C). Interestingly, *Pah* c.1066-11A mice also showed hindlimb clasping, a common phenotype in a number of mouse models of neurodegeneration, including certain cerebellar ataxias and striatal disorders (Figure 4D). Furthermore, *Pah* c.1066-11A mice were unable to perform the vertical pole test; they immediately fell down, as they were not able to hold on to it, thus further revealing a deficit in motor coordination and neuromuscular fitness. Finally, *Pah* c.1066-11A mice were also subjected to the social approach test. In general, *Pah* c.1066-11A mice behaved similar to WT mice in this test. However, when exploring the occupied cage, *Pah* c.1066-11A mice showed a decreased number of physical contacts with the other mouse, what might be indicative of diminished sociability (Figure 4E).

We next examined selected neuronal and glial markers by immunohistochemistry analysis in different brain regions of 20 weeks-old *Pah* c.1066-11A mice, as compared to age matched WT mice. No differences were found when staining for the neuronal marker NeuN (data not shown). On the contrary, both GFAP and Iba1 staining signals were significantly increased in *Pah* c.1066-11A mice, revealing changes in the morphology of both astrocytes and microglial cells, as shown for the dentate gyrus of the hippocampal formation (Figure 5). In contrast, Olig2 staining was significantly reduced indicating a deficit in oligodendrocytes, as shown for cortex, striatum and corpus callosum (Figure 6A), which is in line with the reported myelinization deficits in PKU. Indeed, we could confirm decreased myelinization in brains of the *Pah* c.1066-11A mice by electron microscopy, with decreased thickness of the myelin sheaths and loss of myelinated axons in the lateral tract of the spinal cord (Figure 6B).

The pQ355_Y356insGLQ mutant PAH protein is highly unstable

Structural predictions for the pQ355_Y356insGLQ mutant protein resulting from the c.1066-11G>A variant indicate loss of H-bonds relevant for the correct position of α-helix 11 and steric clashing of the inserted Leu with L333 (Supplementary Figure 3). AlphaFold predicts an additional turn of the helix in the mutant protein but introduces a torsion in order to maintain the interactions of Y356 as in the WT. Therefore, it is probable that the insertion of 3 amino acids hampers the correct arrangement of residues in this region, thereby affecting protein folding. This correlates with the low levels of detected protein *in vitro* and *in vivo*.

Mutant forms of the PAH protein undergoing misfolding and/or aggregation have been reported to be targets for ubiquitin-mediated proteasomal degradation or selective autophagy (Jung-Kc et al., 2019). Accordingly, we examined the impact of the expression of the mutant p.Q355_Y356insGLQ protein in those processes, as well as on the levels of the co-chaperone DNAJC12 and of Hsp70 that participate in the correct folding and stabilisation of the PAH protein (Anikster et al., 2017; Gallego et al., 2020). Western blot analysis in mouse liver lysates showed no significant changes in the ubiquitination profile between WT and *Pah* c.1066-11A mice samples. However, both DNAJC12 and Hsp70 protein levels were decreased (Figure 7A). No significant differences in gene expression levels were detected for the corresponding *Dnajc12* and *Hsp70* genes by RT-qPCR analysis (data not shown). Overall, the results could be indicative of some degree of intracellular co-aggregation of the molecular chaperones with the mutant protein.

PAH protein aggregates may be processed by selective autophagy rather than by the ubiquitin-dependent proteasome system (Aubi et al., 2021; Jung-Kc et al., 2019). We detected altered levels of autophagic markers in *Pah* c.1066-11A mice livers, with higher levels of glycosylated LAMP1 protein and an increase in the conversion of LC3-I to the lower migrating form LC3-II (Figure 7B).

To confirm these results in a human genetic context, we analyzed these markers in the gene edited HepG2 cells carrying the c.1066-11A variant. DNAJC12 protein levels were also decreased, and we could detect decreased levels of LC3BI/II and an increase in glycosylated LAMP1 (Figure 7C, D).

We recently showed that coexpression of DNAJC12 with different PAH mutant constructs in COS7 cells results in a mutation-dependent effect, with an increase in protein levels and activity for some unstable mutants (Gallego et al., 2020). To examine the effect of DNAJC12 overexpression on the mutant p.Q355_Y356insGLQ protein, we transfected the edited HepG2 cells with the pReceiver/DNAJC12 plasmid (Gallego et

al., 2020) and analysed PAH protein levels after 72h. The results showed no significant change in the low levels of mutant PAH protein present in the edited cells (Figure 7E).

DISCUSSION

The development of personalized medicine for genetic diseases requires preclinical testing in appropriate models. PAH variant c.1066-11G>A, a splicing defect resulting in an in-frame insertion of 3 amino acids, is the second most common pathogenic variant causing PKU worldwide, prevalent in Southern Europe and Latin America. Innovative RNA-based therapeutics such as splice switching antisense oligonucleotides (Havens and Hastings, 2016) or HULC lncRNA mimics (Li et al., 2021), gene editing approaches (Böck et al., 2022; Brooks et al., 2023; Villiger et al., 2018) or use of specific pharmacological chaperones (Pey et al., 2008; Santos-Sierra et al., 2012), are theoretically applicable to target either the mutant pre-mRNA or the resulting protein, with the aim of correcting the pathological phenotype of this frequent variant. Due to the limitation in access to patients' tissues where the *PAH* gene is expressed (liver and to a lesser extent, kidney) the splicing defect was studied using minigenes and in a gene edited HepG2 cells homozygous for the c.1066-11G>A variant, in which almost no detectable PAH protein and activity were detected. While *in vitro* studies can be performed in cellular models, for a successful clinical translation of potential therapies, a specific *in vivo* model is needed, in which the gene is expressed in its natural context. This allows to evaluate efficacy in target gene engagement, functional protein expression, downstream functional readouts, therapeutic compounds' uptake and biodistribution, and to perform potential toxicology studies.

As mouse and human intronic sequences are non-homologous, in the present work we have employed CRISPR/Cas9 technology to replace the murine sequence of *Pah* intron 10, proximal to exon 11, with the human sequence carrying the c.1066-11G>A variant. We verified that no modifications of the predicted off-target sites were detected in the humanized mouse and that the genome-editing approach did not interfere with *Pah* gene expression in liver, where precise recapitulation of the splicing defect was confirmed. As expected, no functional PAH protein was detectable in liver resulting in a phenotype resembling the classical form of PKU, with Phe levels above 1200 µmol/L and corresponding low Tyr and Trp levels. Elevated Phe and low Tyr levels were also detected in brain samples of homozygous *Pah* c.1066-11A mice, along with significant dopamine and serotonin deficiencies, which have been implicated in the pathogenesis of the neurological, behavioural and cognitive deficits observed in

untreated patients and in PKU mouse models (de Groot et al., 2010; Winn et al., 2018). Correspondingly, *Pah* c.1066-11A mice exhibit locomotor and behavioural abnormalities, as well as altered hind-limb clasping which is a common feature of different neurological and neurodegenerative diseases (Lieu et al., 2013). This is associated with impaired brain connectivity, possibly mediated by changes in monoaminergic transmission to different brain regions.

Neuropathological evaluation of different PKU mouse models have shown white matter alterations (Bregalda et al., 2023; Ferreira et al., 2021), decreased synaptic connections but normal neuron numbers (Horling et al., 2015), in agreement with brain histology reports from untreated PKU patients (Bauman and Kemper, 1982; Rovelli and Longo, 2023). Classical PKU patients, left untreated, exhibit intellectual disability, developmental delay, epilepsy, autistic features and movement disorders. Our results obtained with the novel PKU mouse model created are in agreement with these observations. Immunohistochemistry analysis in the *Pah* c.1066-11>A mice confirmed an altered brain physiology, with changes in the morphology of astrocytes and microglia, decreased levels of oligodendrocytes and normal neuronal content. Hypomyelination is probably the most important factor in the observed motor impairment with neurological alterations in PKU (Rovelli and Longo, 2023). Elevated Phe levels inhibit 3-hydroxy-3-methylglutaryl coenzyme A reductase (HMGR), thus reducing cholesterol biosynthesis in the PKU brain (Shefer et al., 2000). Cortex and corpus callosum appear to be specifically vulnerable to high Phe levels (Rovelli and Longo, 2023), consistent with our Olig2 immunohistochemistry observations. Other factors contributing to the observed hypomyelination in PKU have been put forward: epigenetic modifications, oxidative stress, defects in calcium homeostasis or altered protein synthesis at the myelin sheath (Bregalda et al., 2023).

Our results demonstrate that the *Pah* c.1066-11>A mice recapitulate PKU disease biology, with biochemical and clinical alterations observed in the most severe classical PKU phenotype, consistent with almost no detectable PAH protein and activity in liver. It thus represents a useful "avatar" mouse model for a frequent PKU variant, for physiopathology studies and for assessing efficacy of various innovative therapeutic modalities. In this sense, for splicing variants, our group has efficiently used splice-switching AONs to correct the resulting splice defects (Gallego-Villar et al., 2014; Martínez-Pizarro et al., 2022; Perez et al., 2010). The c.1066-11G>A variant is in close proximity to the natural 3' splice acceptor site, hindering the design of an AON that could effectively block the novel splice site created by the variant, without causing additional mis-splicing defects such as exon skipping. Indeed, the insertion of 9 extra

nucleotides due to the similar variant c.3191-11T>A in the *ABCA4* gene causing Stargardt disease, could not be corrected by the single AON that could be designed, taking into account the limitations imposed by the target sequence region (Tomkiewicz et al., 2021). Other strategies targeting binding sites of splice factors involved in the negative regulation of exon 11 splicing can be envisaged. However, the newly created 3' splice acceptor site has a higher splice score compared to that of the natural one (MaxEnt score 5.5 versus 3.2). In addition, several reports have shown that the spliceosome preferably uses the first AG-dinucleotide downstream of the branch point (BP) sequence (Chua and Reed, 2001), thus the region between the BP and the 3' splice acceptor site is devoid of AGs (AG-exclusion zone) (Gooding et al., 2006). Future studies are needed to test possible AON-mediated splice correction for the c.1066-11G>A variant.

Alternatively, base editing approaches, which have shown to be effective in correcting PKU variants in different mouse models (Böck et al., 2022; Brooks et al., 2023; Villiger et al., 2018), could provide a novel therapeutic approach for PKU patients with this specific variant. In this sense, the created avatar PKU mouse model with the "humanized" mutant (c.1066-11>A) intron 10 fragment of the endogenous *Pah* gene is an alternative preclinical model suitable for assessing the efficacy of adenine base editing as a therapeutic intervention which could be applicable to a high number of PKU patients worldwide.

In this study we confirmed that the mutant p.Q355_Y356insGLQ PAH protein resulting from the c.1066-11G>A variant was found to be highly unstable, in both *in vitro* (HepG2) and *in vivo* (mouse) models generated in this work. Previous studies using a prokaryotic expression system showed that the mutant protein fused with maltose binding protein was recovered exclusively as high molecular weight aggregates (Pey et al., 2003). In this work, the reduction of immunodetected levels of DNAJC12 and, to a lesser extent, of Hsp70, indicated a possible coaggregation of the mutant PAH protein with these chaperones. This has been described for several unstable human *PAH* variants and for the highly unstable *Pah* variant p.V106A present in the Enu1/1 mouse model (Jung-Kc et al., 2019). Our results support a role of DNAJC12 in the processing of misfolded aggregation-prone p.Q355_Y356insGLQ mutant protein by the autophagy system. Overexpression of DNAJC12 in the edited HepG2 cells did not counteract the stability defect. Our group recently described a mutation-dependent effect of the overexpression of DNAJC12 on missense PAH mutant proteins transiently expressed in COS7 cells; for some there was an increase in protein and activity levels while for others the effects on stability were aggravated

(Gallego et al., 2020). The results suggested that the outcome may depend on the exact pathogenic mechanism of each variant, with DNAJC12 guiding the Hsp70 machinery towards the specific folding or degradation of the client protein. Further studies are necessary to reveal the interactions between the mutant PAH protein and the quality control system and the exact mechanism implicated in its degradation.

In summary, our work has provided a translationally relevant murine model of classical PKU resulting from a frequent PAH variant, as well as the corresponding human hepatoma cellular model, to be used as useful experimental tools to advance research in PKU pathophysiology and personalised therapies.

ACKNOWLEDGEMENTS

This work was funded by Spanish Ministry of Science and Innovation and European Regional Development Fund (grants PID2019-105344RB-I00/AEI/10.13039/501100011033 and PID PID2022-137238OB-I00) and by Fundación Ramón Areces (XX National Call 2020). Centro de Biología Molecular Severo Ochoa receives an institutional grant from Fundación Ramón Areces.

AUTHORS CONTRIBUTIONS

L.R.D. conceptualized this study. A.M-P., S.P., A.L-M., C L-R. performed experiments, analysed data and participated in drafting and preparing of figures. E.M., M.A. and M.C. provided technical support for the experiments. S.R-M. performed in silico structural predictions. B.P., J.J.L., E.R. and L.R.D. acquired funding and provided supervision. J.J.L. and L.R.D. drafted and revised the manuscript. All authors reviewed the final draft.

REFERENCES

Adams, A.D., Fiesco-Roa, M.Ó., Wong, L., Jenkins, G.P., Malinowski, J., Demarest, O.M., Rothberg, P.G., Hobert, J.A., ACMG Therapeutics Committee. Electronic address: documents@acmg.net, 2023. Phenylalanine hydroxylase deficiency treatment and management: A systematic evidence review of the American College of Medical Genetics and Genomics (ACMG). Genet Med 25, 100358. https://doi.org/10.1016/j.gim.2022.12.005

Allen, K.R., Degg, T.J., Rushworth, P.A., Smith, M., Henderson, M.J., 1999. Measurement of phenylalanine and tyrosine in plasma by high-performance liquid chromatography using the inherent fluorescence of aromatic amino acids. Ann Clin Biochem 36 (Pt 2), 207–11.

Anikster, Y., Haack, T.B., Vilboux, T., Pode-Shakked, B., Thony, B., Shen, N., Guarani, V., Meissner, T., Mayatepek, E., Trefz, F.K., Marek-Yagel, D., Martinez, A., Huttlin, E.L., Paulo, J.A., Berutti, R., Benoist, J.F., Imbard, A., Dorboz, I., Heimer, G., Landau, Y., Ziv-Strasser, L., Malicdan, M.C.V., Gemperle-Britschgi, C., Cremer, K., Engels, H., Meili, D., Keller, I., Bruggmann, R., Strom, T.M., Meitinger, T., Mullikin, J.C., Schwartz, G., Ben-Zeev, B., Gahl, W.A., Harper, J.W., Blau, N., Hoffmann, G.F., Prokisch, H., Opladen, T., Schiff, M., 2017. Biallelic Mutations in DNAJC12 Cause Hyperphenylalaninemia, Dystonia, and Intellectual Disability. Am J Hum Genet 100, 257–266. https://doi.org/S0002-9297(17)30002-2 [pii] 10.1016/j.ajhg.2017.01.002

Aubi, O., Prestegård, K.S., Jung-Kc, K., Shi, T.-J.S., Ying, M., Grindheim, A.K., Scherer, T., Ulvik, A., McCann, A., Spriet, E., Thöny, B., Martinez, A., 2021. The Pah-R261Q mouse reveals oxidative stress associated with amyloid-like hepatic aggregation of mutant phenylalanine hydroxylase. Nat Commun 12, 2073. https://doi.org/10.1038/s41467-021-22107-1

Barman-Aksözen, J., C´wiek, P., Bansode, V.B., Koentgen, F., Trüb, J., Pelczar, P., Cinelli, P., Schneider-Yin, X., Schümperli, D., Minder, E.I., 2017. Modeling the ferrochelatase c.315-48C modifier mutation for erythropoietic protoporphyria (EPP) in mice. Disease Models & Mechanisms 10, 225. https://doi.org/10.1242/dmm.027755

Bauman, M.L., Kemper, Th.L., 1982. Morphologic and histoanatomic observations of the brain in untreated human phenylketonuria. Acta Neuropathologica 58, 55–63. https://doi.org/10.1007/BF00692698

Böck, D., Rothgangl, T., Villiger, L., Schmidheini, L., Matsushita, M., Mathis, N., Ioannidi, E., Rimann, N., Grisch-Chan, H.M., Kreutzer, S., Kontarakis, Z., Kopf, M., Thöny, B., Schwank, G., 2022. In vivo prime editing of a metabolic liver disease in mice. Sci Transl Med 14, eabl9238. https://doi.org/10.1126/scitranslmed.abl9238

Bregalda, A., Carducci, C., Viscomi, M.T., Pierigè, F., Biagiotti, S., Menotta, M., Biancucci, F., Pascucci, T., Leuzzi, V., Magnani, M., Rossi, L., 2023. Myelin basic protein recovery during PKU mice lifespan and the potential role of microRNAs on its regulation. Neurobiol Dis 180, 106093. https://doi.org/10.1016/j.nbd.2023.106093

Brooks, D.L., Carrasco, M.J., Qu, P., Peranteau, W.H., Ahrens-Nicklas, R.C., Musunuru, K., Alameh, M.-G., Wang, X., 2023. Rapid and definitive treatment of phenylketonuria in variant-humanized mice with corrective editing. Nat Commun 14, 3451. https://doi.org/10.1038/s41467-023-39246-2

Chen, A., Pan, Y., Chen, J., 2022. Clinical, genetic, and experimental research of hyperphenylalaninemia. Front Genet 13, 1051153. https://doi.org/10.3389/fgene.2022.1051153

Chua, K., Reed, R., 2001. An Upstream AG Determines Whether a Downstream AG Is Selected during Catalytic Step II of Splicing. Molecular and Cellular Biology 21, 1509–1514. https://doi.org/10.1128/MCB.21.5.1509-1514.2001

de Groot, M.J., Hoeksma, M., Blau, N., Reijngoud, D.J., van Spronsen, F.J., 2010. Pathogenesis of cognitive dysfunction in phenylketonuria: review of hypotheses. Mol Genet Metab 99 Suppl 1, S86-9. https://doi.org/10.1016/j.ymgme.2009.10.016 S1096-7192(09)00300-X [pii]

Dworniczak, B., Aulehla-Scholz, C., Kalaydjieva, L., Bartholome, K., Grudda, K., Horst, J., 1991. Aberrant splicing of phenylalanine hydroxylase mRNA: the major cause for Phenylketonuria in parts of Southern Europe. Genomics 11, 242–946.

Emsley, P., Lohkamp, B., Scott, W.G., Cowtan, K., 2010. Features and development of Coot. Acta Crystallogr D Biol Crystallogr 66, 486–501. https://doi.org/10.1107/S0907444910007493

Ferreira, B.K., Rodrigues, M.T., Streck, E.L., Ferreira, G.C., Schuck, P.F., 2021. White matter disturbances in phenylketonuria: Possible underlying mechanisms. Journal of Neuroscience Research 99, 349–360. https://doi.org/10.1002/jnr.24598

Gallego, D., Leal, F., Gamez, A., Castro, M., Navarrete, R., Sanchez-Lijarcio, O., Vitoria, I., Bueno-Delgado, M., Belanger-Quintana, A., Morais, A., Pedron-Giner, C., Garcia, I., Campistol, J., Artuch, R., Alcaide, C., Cornejo, V., Gil, D., Yahyaoui, R., Desviat, L.R., Ugarte, M., Martinez, A., Perez, B., 2020. Pathogenic variants of DNAJC12 and evaluation of the encoded co-chaperone as a genetic modifier of hyperphenylalaninemia. Hum Mutat. https://doi.org/10.1002/humu.24026

Gallego-Villar, L., Viecelli, H.M., Perez, B., Harding, C.O., Ugarte, M., Thony, B., Desviat, L.R., 2014. A sensitive assay system to test antisense oligonucleotides for splice suppression therapy in the mouse liver. Mol Ther Nucleic Acids 3, e193. https://doi.org/10.1038/mtna.2014.44 mtna201444 [pii]

Gooding, C., Clark, F., Wollerton, M.C., Grellscheid, S.-N., Groom, H., Smith, C.W.J., 2006. A class of human exons with predicted distant branch points revealed by analysis of AG dinucleotide exclusion zones. Genome Biol 7, R1. https://doi.org/10.1186/gb-2006-7-1-r1

Haefele, M.J., White, G., McDonald, J.D., 2001. Characterization of the mouse phenylalanine hydroxylase mutation Pah(enu3). Mol Genet Metab 72, 27–30. https://doi.org/10.1006/mgme.2000.3104

Havens, M.A., Hastings, M.L., 2016. Splice-switching antisense oligonucleotides as therapeutic drugs. Nucleic Acids Res 44, 6549–63. https://doi.org/10.1093/nar/gkw533

Heintz, C., Dobrowolski, S.F., Andersen, H.S., Demirkol, M., Blau, N., Andresen, B.S., 2012. Splicing of phenylalanine hydroxylase (PAH) exon 11 is vulnerable: molecular pathology of mutations in PAH exon 11. Mol Genet Metab 106, 403–11. https://doi.org/S1096-7192(12)00202-8 [pii] 10.1016/j.ymgme.2012.05.013

Hillert, A., Anikster, Y., Belanger-Quintana, A., Burlina, A., Burton, B.K., Carducci, C., Chiesa, A.E., Christodoulou, J., Đorđević, M., Desviat, L.R., Eliyahu, A., Evers, R.A.F., Fajkusova, L., Feillet, F., Bonfim-Freitas, P.E., Giżewska, M., Gundorova, P., Karall, D., Kneller, K., Kutsev, S.I., Leuzzi, V., Levy, H.L., Lichter-Konecki, U., Muntau, A.C., Namour, F., Oltarzewski, M., Paras, A., Perez, B., Polak, E., Polyakov, A.V., Porta, F., Rohrbach, M., Scholl-Bürgi, S., Spécola, N., Stojiljković, M., Shen, N., Santana-da Silva, L.C., Skouma, A., van Spronsen, F., Stoppioni, V., Thöny, B., Trefz, F.K., Vockley, J., Yu, Y., Zschocke, J., Hoffmann, G.F., Garbade, S.F., Blau, N., 2020. The Genetic Landscape and Epidemiology of Phenylketonuria. Am J Hum Genet 107, 234–250. https://doi.org/10.1016/j.ajhg.2020.06.006

Hopf, S., Nowak, C., Hennermann, J.B., Schmidtmann, I., Pfeiffer, N., Pitz, S., 2020. Saccadic reaction time and ocular findings in phenylketonuria. Orphanet J Rare Dis 15, 124. https://doi.org/10.1186/s13023-020-01407-7

Horling, K., Schlegel, G., Schulz, S., Vierk, R., Ullrich, K., Santer, R., Rune, G.M., 2015. Hippocampal synaptic connectivity in phenylketonuria. Human Molecular Genetics 24, 1007–1018. https://doi.org/10.1093/hmg/ddu515

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W.,

Kavukcuoglu, K., Kohli, P., Hassabis, D., 2021. Highly accurate protein structure prediction with AlphaFold. Nature 596, 583–589. https://doi.org/10.1038/s41586-021-03819-2Jung-Kc, K., Himmelreich, N., Prestegård, K.S., Shi, T.-J.S., Scherer, T., Ying, M., Jorge-Finnigan, A., Thöny, B., Blau, N., Martinez, A., 2019. Phenylalanine hydroxylase variants interact with the co-chaperone DNAJC12. Hum Mutat 40, 483–494. https://doi.org/10.1002/humu.23712

Li, Y., Tan, Z., Zhang, Y., Zhang, Z., Hu, Q., Liang, K., Yao, J., Ye, Y., Li, Y.-C., Li, C., Liao, L., Xu, J., Xing, Z., Pan, Y., Chatterjee, S.S., Nguyen, T.K., Hsiao, H., Egranov, S.D., Putluri, N., Coarfa, C., Hawke, D.H., Gunaratne, P.H., Tsai, K.-L., Han, L., Hung, M.-C., Calin, G.A., Namour, F., Guéant, J.-L., Muntau, A.C., Blau, N., Sutton, V.R., Schiff, M., Feillet, F., Zhang, S., Lin, C., Yang, L., 2021. A noncoding RNA modulator potentiates phenylalanine metabolism in mice. Science 373, 662–673. https://doi.org/10.1126/science.aba4991

Lieu, C.A., Chinta, S.J., Rane, A., Andersen, J.K., 2013. Age-Related Behavioral Phenotype of an Astrocytic Monoamine Oxidase-B Transgenic Mouse Model of Parkinson's Disease. PLOS ONE 8, e54200. https://doi.org/10.1371/journal.pone.0054200

Martinez-Pizarro, A., Dembic, M., Perez, B., Andresen, B.S., Desviat, L.R., 2018. Intronic PAH gene mutations cause a splicing defect by a novel mechanism involving U1snRNP binding downstream of the 5' splice site. PLoS Genet 14, e1007360. https://doi.org/10.1371/journal.pgen.1007360

Martínez-Pizarro, A., Leal, F., Holm, L.L., Doktor, T.K., Petersen, U.S.S., Bueno, M., Thöny, B., Pérez, B., Andresen, B.S., Desviat, L.R., 2022. Antisense Oligonucleotide Rescue of Deep-Intronic Variants Activating Pseudoexons in the 6-Pyruvoyl-Tetrahydropterin Synthase Gene. Nucleic Acid Ther 32, 378–390. https://doi.org/10.1089/nat.2021.0066

McDonald, J.D., Charlton, C.K., 1997. Characterization of mutations at the mouse phenylalanine hydroxylase locus. Genomics 39, 402–405. https://doi.org/10.1006/geno.1996.4508

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., Steinegger, M., 2022. ColabFold: making protein folding accessible to all. Nat Methods 19, 679–682. https://doi.org/10.1038/s41592-022-01488-1

Oliveros, J.C., Franch, M., Tabas-Madrid, D., San-Leon, D., Montoliu, L., Cubas, P., Pazos, F., 2016. Breaking-Cas-interactive design of guide RNAs for CRISPR-Cas

experiments for ENSEMBL genomes. Nucleic Acids Res 44, W267-71. https://doi.org/10.1093/nar/gkw407

Ormazabal, A., Garcia-Cazorla, A., Fernandez, Y., Fernandez-Alvarez, E., Campistol, J., Artuch, R., 2005. HPLC with electrochemical and fluorescence detection procedures for the diagnosis of inborn errors of biogenic amines and pterins. J Neurosci Methods 142, 153–8.

Perez, B., Rodriguez-Pascau, L., Vilageliu, L., Grinberg, D., Ugarte, M., Desviat, L.R., 2010. Present and future of antisense therapy for splicing modulation in inherited metabolic disease. J Inherit Metab Dis 33, 397–403. https://doi.org/10.1007/s10545-010-9135-1

Pey, A.L., Desviat, L.R., Gámez, A., Ugarte, M., Pérez, B., 2003. Phenylketonuria: genotype-phenotype correlations based on expression analysis of structural and functional mutations in PAH. Hum Mutat 21, 370–378. https://doi.org/10.1002/humu.10198

Pey, A.L., Ying, M., Cremades, N., Velazquez-Campoy, A., Scherer, T., Thöny, B., Sancho, J., Martinez, A., 2008. Identification of pharmacological chaperones as potential therapeutic agents to treat phenylketonuria. J Clin Invest 118, 2858–2867. https://doi.org/10.1172/JCI34355

Rivera-Barahona, A., Sánchez-Alcudia, R., Viecelli, H.M., Rüfenacht, V., Pérez, B., Ugarte, M., Häberle, J., Thöny, B., Desviat, L.R., 2015. Functional characterization of the spf/ash splicing variation in OTC deficiency of mice and man. PLoS One 10, e0122966. https://doi.org/10.1371/journal.pone.0122966

Rovelli, V., Longo, N., 2023. Phenylketonuria and the brain. Molecular Genetics and Metabolism 139. https://doi.org/10.1016/j.ymgme.2023.107583

Santos-Sierra, S., Kirchmair, J., Perna, A.M., Reiss, D., Kemter, K., Roschinger, W., Glossmann, H., Gersting, S.W., Muntau, A.C., Wolber, G., Lagler, F.B., 2012. Novel pharmacological chaperones that correct phenylketonuria in mice. Hum Mol Genet 21, 1877–87. https://doi.org/dds001 [pii] 10.1093/hmg/dds001

Shefer, S., Tint, G.S., Jean-Guillaume, D., Daikhin, E., Kendler, A., Nguyen, L.B., Yudkoff, M., Dyer, C.A., 2000. Is there a relationship between 3-hydroxy-3-methylglutaryl coenzyme a reductase activity and forebrain pathology in the PKU mouse? J Neurosci Res 61, 549–563. https://doi.org/10.1002/1097-4547(20000901)61:5<549::AID-JNR10>3.0.CO;2-0

Singh, K., Cornell, C.S., Jackson, R., Kabiri, M., Phipps, M., Desai, M., Fogle, R., Ying, X., Anarat-Cappillino, G., Geller, S., Johnson, J., Roberts, E., Malley, K., Devlin, T., DeRiso, M., Berthelette, P., Zhang, Y.V., Ryan, S., Rao, S., Thurberg, B.L., Bangari, D.S., Kyostio-Moore, S., 2021. CRISPR/Cas9 generated knockout mice lacking phenylalanine hydroxylase protein as a novel preclinical model for human phenylketonuria. Sci Rep 11, 7254. https://doi.org/10.1038/s41598-021-86663-8

Tomkiewicz, T.Z., Suárez-Herrera, N., Cremers, F.P.M., Collin, R.W.J., Garanto, A., 2021. Antisense Oligonucleotide-Based Rescue of Aberrant Splicing Defects Caused by 15 Pathogenic Variants in ABCA4. Int J Mol Sci 22, 4621. https://doi.org/10.3390/ijms22094621

van Spronsen, F.J., Blau, N., Harding, C., Burlina, A., Longo, N., Bosch, A.M., 2021. Phenylketonuria. Nat Rev Dis Primers 7, 36. https://doi.org/10.1038/s41572-021-00267-0

Vázquez-Domínguez, I., Garanto, A., 2022. Considerations for Generating Humanized Mouse Models to Test Efficacy of Antisense Oligonucleotides. Methods Mol Biol 2434, 267–279. https://doi.org/10.1007/978-1-0716-2010-6_18

Villiger, L., Grisch-Chan, H.M., Lindsay, H., Ringnalda, F., Pogliano, C.B., Allegri, G., Fingerhut, R., Häberle, J., Matos, J., Robinson, M.D., Thöny, B., Schwank, G., 2018. Treatment of a metabolic liver disease by in vivo genome base editing in adult mice. Nat Med 24, 1519–1525. https://doi.org/10.1038/s41591-018-0209-1

Winn, S.R., Scherer, T., Thöny, B., Ying, M., Martinez, A., Weber, S., Raber, J., Harding, C.O., 2018. Blood phenylalanine reduction corrects CNS dopamine and serotonin deficiencies and partially improves behavioral performance in adult phenylketonuric mice. Mol Genet Metab 123, 6–20. https://doi.org/10.1016/j.ymgme.2017.10.009

Table 1. Primers used for gDNA and cDNA amplification.

Name	Sequence 5'-3	Method
PAH 10-11-12 S	5'-GGTAACGGAGCCAACATGGTTTACTG-3'	RT-PCR human minigene
PAH 10-11-12 AS	5'- AGACTCGAGGGTAGTCTATTATCTGTT-3'	
pcDNA3.1 F	5'-ACCCACTGCTTACTGGCTTA-3'	RT-PCR mouse minigene
pcDNA3.1 R	5'-GCAACTAGAAGGCACAGTCG-3'	
hPAH exon 10 F	5'-ACTGTGGAGTTTGGGCTCTG-3'	RT-PCR HepG2 cells
hPAH exon 12 R	5'-ACTGAGAAGGGCCGAGGTAT-3'	
mPAH exon 10 F	5'-GGCTCTTGTCATCCTTTGGA-3'	RT-PCR mice liver
mPAH exon 11 R	5'-TACTCCTGGCAGGCTGTCTT-3'	
hPAH intron 10 F	5'-ACCTATGCATGCAGCCTTGT-3'	PCR HepG2 edited cells (mutation detection)
hPAH intron 11 R	5'- AGGGTGGAGAACATGGGAGA-3'	
mPAH exon 10 F	5'-GGCTCTTGTCATCCTTTGGA-3'	PCR wt allele genotyping
mPAH WT R	5'-GCTTCTTCCATCTCCCCATT-3'	
mPAH exon 10 F	5'-GGCTCTTGTCATCCTTTGGA-3'	PCR mutant allele genotyping
-11G>A human R	5'-CAGTATTCCCTGCTGCATCC-3'	
hPAH ex1-2 F	5'-GACTTTGGACAGGAAACAAGC-3'	qPCR human PAH
hPAH ex2-3 R	5'-CATTCTCCTCAAATAAGCGCA-3'	
hDNAJC12 F	5'-CAGTGAAGACGTCAATGCACT-3'	qPCR human Dnajc12
hDNAJC12 R	5'-GGTTGAAGCCAGCTCCTCT-3'	
hHSP70 F	5'-GATCAACGACGGAGACAAGC-3'	qPCR human Hsp70
hHSP70 R	5'-GCTGCGAGTCGTTGAAGTAG-3'	
mPah ex6-7 F	5'-TTCTGCAGACTTGTACTGGT-3'	qPCR mouse PAH
mPah ex7-8 R	5'-ACAGATATCAGGTTCAGGTG-3'	
mDnajc12 F	5'-CTCTCCTCGGTTGAGCAAAT-3'	qPCR mouse Dnajc12
mDnajc12 R	5'-ATGCATCGAAGTTTTCACTG-3'	
mHsp70 F	5'-GCGCTCGAATCCTATGCCTT-3'	qPCR mouse Hsp70
mHsp70 R	5'-GTGCACGAACTCCTCCTTGT-3'	

Table 2. DNA template, gRNA and top predicted off-targets for gene editing in HepG2 cells.

DNA TEMPLATE	5'-CCTATGGGATGCAGCAGGGAATACTGATCCTGATTTAACAGTGATAATAACTTTTCACTTAGGGCCTACAGTACTGCTTATCAGAGAAGCCAAAGCTTCTCCCCCTGGAGCTGGAGAAGA-3'				
GUIDE (gRNA)	5'-UCUCUGAUAAGCAGUACUGU -3'				

OFF-TARGET SEQUENCE*	MM*	GENOMIC LOCALIZATION	REGION	GENE	PRIMERS (5'-3')
T**G**TA**GG**A**C**AAGCAGTACTGTTGG	4	Chr8: 118873153-118873175	INTRON	SAMD12-AS1	5'-TGGTGCACTAAAGCAGCCAT-3' 5'-TGCCAAGGCTACGGAAAGTT-3'
TCTCT**CACAG**GCAGTACTGTTGG	3	Chr8: 118873153-118873175	UPSTREAM	SCFD2	5'-TGGGTTCTGTTTGTTCCACC-3' 5'-AGTGGTTTGATAGGTGGGGG-3'
TC**AA**TGA**GG**AGCAGTACTGTGGG	4	Chr21: 39462538-39462560	EXON	GET1-SH3BGR (IKZF3)	5'-AGCAGGCACATAAGGGTGTT-3' 5'-GCCCACTGTCCTTAGTGGAT-3'

*mismatches (MM) shown in bold

Table 3. DNA template, gRNA and top predicted off-targets for gene editing in mice.

DNA TEMPLATE	5'-AGTCCATGCTAAATGTCTCAGTTTTAGGGAATTTGCAAGAGAAGGGGCACAAATGGCCTATGGGATGCAGCAGGGAATACTGATCCTGATTTAACAGTGATAATAACTTTTCACTTGGGGCCTACAGTACTGTTTATCAGACAAGCCAAAGCTCCTGCCCC-3'				
GUIDE (gRNA)	5'-GUUUUAGGGAAUUUGCACUAA-3'				

OFF-TARGET SEQUENCE*	MM*	GENOMIC LOCALIZATION	REGION	GENE	PRIMERS (5'-3')
TTT**A**TAGGGAATTTGCACT**G**TGG	3	Chr15: 32795435-32795457	EXON	Gm32618	5'-GGGAACACCAGTTGCTACTAGC-3' 5'-GCACCAGAGACTCTCCAGCA-3'
TATC**AA**GGGAATTTGCACTATGG	4	Chr7: 134570396-134570418	INTRON	Dock1	5'-GGTAAAGTCCTCAGGTGGTCA-3' 5'-CCATGAGCCAACATTACACC-3'
GTT**A**TATGGA**G**TTTGCACTAAGG	3	Chr7: 89552521-89552543	EXON	Ccdc81	5'-CAGTGCTCTGGCTTCGTGTT-3' 5'-CACACACCTGCATCAAGTCA-3'
TAT**A**TAGGTAATTTGCACTAAGG	4	Chr10: 120001151-120001173	INTRON	Irak3	5'-GGCAAGAGCAAGAGGAGAAGA-3' 5'-GAACCCAGAGCTCTCAGGTCAT-3'

*mismatches (MM) shown in bold

LEGENDS TO FIGURES

Figure 1. Functional analysis of the c.1066-11G>A PAH variant. A) Schematic representation of minigene construction in the pcDNA3.1 vector, RT-PCR after transfection in Hep3B cells with wild-type (WT) and mutant minigenes and Sanger sequencing. **B)** Western Blot for immunodetection of PAH from HepG2 WT and mutant cell lines. COS cells were used as negative control. Tubulin was used as loading control.

Figure 2. Characterization of *Pah* c.1066-11A (PKU) mice. A) PKU mice survival. **B)** Body weight distribution by genotype during 12 weeks. Data are presented as mean of body weight. **C)** Hypopigmentation (left photography, 12 weeks-old mice) and microphthalmia (right photography, 20 weeks-old mice) in homozygous PKU mice compared with wild-type (WT). **D)** Western Blot for immunodetection of PAH from WT, heterozygous (Het) and PKU mice livers. GAPDH was used as loading control. **E)** Overview of PAH specific activity in liver lysates. Data are presented as mean ± SEM. **$P<0.01$, ***$P<0.001$.

Figure 3. Brain neurotransmitters and related metabolites levels in *Pah* c.1066-11A (PKU) mice. Measures of neurotransmitters in brain lysates by HPLC. Data are presented as mean ± SEM. **$P<0.01$, ***$P<0.001$.

Figure 4. Behavioural performance of *Pah* c.1066-11A (PKU) mice. A) Distance travelled and velocity in open field test. **B)** Hind-limb clasping. **C)** Latency to fall off the rotarod. **D)** Latency to fall off the inverted grid test. **E)** Time spent interacting with empty cage and unfamiliar mouse in social approach test. Data are presented as mean ± SEM. *$P<0.05$, **$P<0.01$, ***$P<0.001$.

Figure 5. Analysis of different types of neuronal markers. Immunohistochemistry brain image of Iba1 **(A)** and GFAP **(B)** in PKU mice. Data are presented as mean ± SEM. *$P<0.05$, **$P<0.01$, ***$P<0.001$.

Figure 6. Myelinization defects. A) Immunohistochemistry brain images of Olig2 in PKU mice. Data are presented as mean ± SEM (n = 6 WT and 7 PKU). *$P<0.05$. **B)** Transmission electron microscopy (TEM) images of lumbar transverse sections of the spinal cord of WT and *Pah* c.1066-11A (PKU) mice.

Figure 7. Evaluation of the effect of the c.1066-11G>A variant on PAH folding and stability. (A, B) Western blot analysis of PAH, ubiquitin and DNAJC12 and HSP70 co-chaperones and autophagy markers LAMP1, LC3BII and p62 in liver samples from 12-week-old mice. **(C, D)** Western blot analysis in HepG2 cells. **E)** PAH levels detected by

western blot in HepG2 cells 72 h after transfection with 2 µg of the plasmid coding for DNAJC12. GAPDH or β-actin were used as loading controls.

Figure 1

A)

5'ss (9.5)   3'ss (3.2)   5'ss (9.2)   3'ss (8.9)

| Exon 10 | Intron 10 | Exon 11 | Intron 11 (shortened) | Exon 12 |

pcDNA3.1   556 bp   1958 bp   pcDNA3.1

100 bp   134 bp   118 bp

B)

HepG2

KDa	WT	-11G>A	COS	
51				PAH
55				Tubulin

bp

	WT	-11 G>A

388
379
245

10 — 11 12

10 11 12

10 12

Exon 10   Exon 11

C C T T T G G T G A A T T A C A G T A C T G C T T A T C A G A G

WT

Exon 10   Intron 10   Exon 11

A A T T A C A G G G C C T A C A G T A C T G C T T A T C A G A G

-11 G>A

Figure 2

Figure 3

Figure 4

A) Open field test

WT
(n=22)
PKU
(n=20)

B) Rotarod

WT
(n=22)
PKU
(n=20)

C) Inverted grid

WT
(n=13)
PKU
(n=15)

D) Hind clasping reflex

E) Social approach

WT
(n=19)
PKU
(n=17)

Figure 5

A) Iba1

B) GFAP

Figure 6

A) Olig2

B) TEM

Figure 7