1   **Plasma proteome signature of canine acute haemorrhagic diarrhoea syndrome (AHDS)**

2   Lukas Huber[1], Benno Kuropka[2], Pavlos G. Doulidis[1], Elisabeth Baszler[1], Lukas Martin[1], Anda Rosu[1],
3   Lisa Kulmer[1], Carolina Frizzo Ramos[3], Alexandro Rodríguez-Rojas[1*], Iwan A. Burgener[1*].
4
[1]Department for Companion Animals, Small Animal Internal Medicine, University of Veterinary
Medicine, Vienna, Austria.
[2] Institute of Chemistry and Biochemistry, Freie Universität Berlin, Berlin, Germany
[3] The Interuniversity Messerli Research Institute, Medical University Vienna and University of
Veterinary Medicine Vienna, Vienna, Austria.
*Correspondence: Alexandro.Rojas@vetmeduni.ac.at, Iwan.Burgener@vetmeduni.ac.at

Keywords: AHDS, haemorrhagic, canine diarrhoea, plasma proteome, proteomics, biomarker

Abstract

Acute haemorrhagic diarrhoea is a common complaint in dogs. In addition to causes like intestinal
parasites, dietary indiscretion, intestinal foreign bodies, canine parvovirus infection, or
hypoadrenocorticism, acute haemorrhagic diarrhoea syndrome (AHDS) is an important and
sometimes life-threatening differential diagnosis. There is some evidence supporting the link
between *Clostridium perfringens* toxins and AHDS. These toxins may be partially responsible for the
epithelial cell injury, but the pathogenesis of AHDS is still not fully understood. Recent studies have
suggested that severe damage to the intestinal mucosa and associated barrier dysfunction can
trigger chronic gastrointestinal illnesses. Besides bloodwork and classical markers for AHDS such as
protein loss and intestinal bacterial dysbiosis, we focused mainly on the plasma-proteome to
identify systemic pathological alterations during this disease and searched for potential biomarkers
to improve the diagnosis. To accomplish this, we used liquid chromatography-mass spectrometry.
We compared the proteomic profiles of 20 dogs with AHDS to 20 age-, breed-, and sex-matched
control dogs. We identified and quantified 207 plasmatic proteins, from which dozens showed
significantly altered levels in AHDS. Serpina3, Lipopolysaccharide-binding protein, several Ig-like
domain-containing proteins, Glyceraldehyde-3-phosphate dehydrogenase and Serum amyloid A
were more abundant in plasma from AHDS affected dogs. In contrast, other proteins such as
Paraoxonase, Selenoprotein, Amine oxidases, and Apolipoprotein C-IV were significantly less
abundant. Many of those proteins we are known to be associated with inflammation. Other proteins
like Serpina3 and RPLP1 play an important role in oncogenesis. Some proteins and their role have
not yet been described in dogs with diarrhoea. Our study opens new avenues that could contribute
to the understanding of the aetiology and pathophysiology of AHDS.

Introduction

Acute Haemorrhagic Diarrhoea Syndrome (AHDS) in dogs is a potentially life-threatening gastrointestinal disorder with a high prevalence [1–3]. Characterised by a sudden onset of bloody diarrhoea, vomiting, and profound dehydration resulting in haemoconcentration, AHDS treatment poses significant challenges for veterinary practitioners [4]. The condition can progress rapidly, leading to severe complications and, in some small number of cases, the outcome can be fatal if not adequately addressed [1]. Persistent vomiting and profuse diarrhoea lead to rapid bodily fluid loss, severe dehydration, and an electrolyte imbalance [7]. Dehydration affects the dog's overall health, leading to weakness, lethargy, and potential organ dysfunction. Blood loss can lead to anaemia and contribute to morbidity [8]. In severe cases, AHDS can trigger systemic inflammatory response syndrome (SIRS), an aggressive immune reaction that causes widespread inflammation. This reaction can progress to sepsis or even septic shock, leading to a life-threatening drop in blood pressure and organ dysfunction [9]. Septic shock requires immediate and intensive medical intervention to decrease the risk of death but is, in any case, associated with a high mortality [5]. Despite its prevalence and numerous publications investigating this disease, the aetiology of AHDS is unknown. Several pathogens are associated with AHDS in dogs, including some viruses, bacteria, and parasites. In AHDS, significant damage to the lining of the stomach and intestines occurs. It is suspected that the involved pathogens could directly attack and destroy the intestinal cell lining, leading to haemorrhage and ulceration [5]. Some strains of *Clostridium perfringens*, a bacterium that is an ordinary member of the canine microbiome, produce the pore-forming toxins NetE and NetF, which are commonly discussed in the literature as a major cause of gastrointestinal epithelial necrosis leading to AHDS [6]. As the incidence of AHDS is next to parasitic diseases and alimentary indiscretion probably the highest among acute gastrointestinal diseases in dogs, understanding the underlying aetiology and pathophysiology, identifying predisposing factors and developing effective management strategies are critical in mitigating the impact of this syndrome on canine health [13]. While MS-based proteomic analysis in veterinary medicine is less extensive than in human clinical research, notable studies have focused on samples from dogs afflicted with diverse diseases [14]. The application of mass spectrometry for quantitative proteome analysis in veterinary medicine holds significant promise for enhancing our understanding of the underlying mechanisms of various diseases like AHDS in dogs. These techniques enable researchers to identify and quantify proteins with a higher degree of precision, aiding in discovering of potential biomarkers and therapeutic

targets. Although the current body of research is comparatively limited, the growing attention to

this cutting-edge approach in veterinary science is encouraging [15].

This article reports a quantitative proteomic study of canine plasma from patients affected by AHDS

and its comparison with plasma from healthy controls. The main goal of this study is to describe the

general perturbation of the blood induced by AHDS and the exploration of new potential biomarkers

that could help to understand the pathogenesis of AHDS better, improve future diagnostics and aid

in the treatment and follow-up of the disease.

Material and Methods

Animals and sample collection.

All the samples used for this work were surplus materials from previous studies. The previous studies

were approved by the Ethics Committee of the University of Veterinary Medicine Vienna and the

Austrian Federal Ministry of Science and Research (BMBWF-68.205/0216-V/3b/2019, 2021-

0.518.574), and methods for sample collection were carried out in accordance with relevant Austrian

guidelines and regulations. In this work, we analysed the samples of 20 client-owned dogs presented

to the Clinic for Small Animal Internal Medicine of the University of Veterinary Medicine Vienna,

Austria, over a period of one year and diagnosed with AHDS according to current diagnostic criteria

from a still ongoing prospective study. We compared them to a healthy control group matched

according to age, sex and a phylogenetic tree. Data collected from these animals included age, sex,

breed, body weight, physical examination findings and clinicopathological values. Further diet and a

history of previous gastrointestinal diseases were especially investigated during the anamnesis. Only

dogs with acute bloody diarrhoea for less than three days, with haemoconcentration with a

haematocrit of ≥50% and without a known history of chronic gastrointestinal disorders were

included in the study.

As a control group, we used clinically healthy client-owned dogs (N=20) with no signs of disease that

were presented to the Clinic for Small Animal Internal Medicine over a six-month period for clinical

examination and blood sampling to be enrolled in a behavioural study from the Interuniversity

Messerli Institute of Research (Vienna, Austria). Inclusion followed a thorough history, physical

examination, and blood sampling. The samples were analysed including complete blood count,

plasma biochemical profile and electrolytes measurements. All samples showed no significant

pathological alterations.

The samples from both groups, diagnosed with AHDS and the healthy control group, were grouped

into five independent pool sets, each created using the "tidyverse" R package to generate

randomisation and group selection as previously described [15]. The procedure involved assigning a

number to each sample. Upon execution, the computer script generated five random lists, each

containing four individuals. These lists were used to prepare the pools. Samples were pooled to limit

the effect of individual samples.

Quantitative proteome analysis by label-free liquid chromatography–mass spectrometry (LC–MS).

Blood samples were collected in 2 ml Vacuette tubes with lithium heparin (clinical chemistry) 13x75

green cap-white ring PREMIUM (Greiner Bio-One GmbH, Bad Haller Str. 32, 4550, Austria) and the

plasma was centrifugated at 2000 x g for 5 minutes and stored in -20ºC. Five microliters of plasma

from each of the 40 individuals was collected from the supernatant after a centrifugation step at 20

000 x g at 4 °C. For each experimental replicate, five samples containing the pooled plasma of four

randomly selected individuals were prepared. From each plasma pool, 3 µl were transferred to a

tube containing 120 µl urea denaturing buffer (6 M urea, 2 M thiourea, and 10 mM HEPES; pH 8.0).

Proteins were reduced by adding 5 µl dithiothreitol (10 mM) and incubated for 30 min at room

temperature. Then, the samples were alkylated by the addition of 5 µl iodoacetamide solution (55

mM) and incubated at room temperature for another 30 min in the dark. All samples were diluted

by adding four volumes of ammonium bicarbonate buffer (50 mM) to reduce the urea concentration.

Thereafter, samples were digested overnight at 37 °C after adding 1 µg of trypsin protease (Thermo

Scientific, USA). Next day, the samples were acidified by adding a final concertation of 5%

acetonitrile and 0.3% of TFA and subsequently desalted via C18 StageTips with Empore™ C18

Extraction Disks [16]. Peptides eluted from the StageTips were dried by vacuum centrifugation.

Peptides were reconstituted in 80 µl of 0.05% TFA, 4% acetonitrile and 1 µl of each sample was

applied to an Ultimate 3000 reversed-phase capillary nano liquid chromatography system connected

to a Q Exactive HF mass spectrometer (Thermo Fisher Scientific). Samples were injected and

concentrated on a PepMap100 C18 trap column (3 µm, 100 Å, 75 µm inner diameter [i.d.] × 20mm,

nanoViper; Thermo Scientific) equilibrated with 0.05% TFA in water. After switching the trap column

inline, LC separations were performed on an Acclaim PepMap100 C18 capillary column (2 µm, 100

Å, 75 µm i.d. × 500mm, nanoViper; Thermo Scientific) at an eluent flow rate of 300 nl/min. Mobile

phase A contained 0.1% (v/v) formic acid in water, and mobile phase B contained 0.1% (v/v) formic

acid and 80% (v/v) acetonitrile in water. The column was pre-equilibrated with 5% mobile phase B

followed by an increase to 44% mobile phase B over 100 min. Mass spectra were acquired in a data-

dependent mode, utilizing a single MS survey scan (m/z 300–1650) with a resolution of 60,000, and

MS/MS scans of the 15 most intense precursor ions with a resolution of 15,000. The dynamic

exclusion time was set to 20 s, and the automatic gain control was set to 3×10^6 and 1×10^5 for MS

and MS/MS scans, respectively.

MS and MS/MS spectra were processed using the MaxQuant software package (version 2.0.3.0) with

the implemented Andromeda peptide search engine [17]. Data were searched against the *Canis*

lupus familiaris reference proteome (ID: UP000002254; downloaded from Uniprot.org on

29.03.2023; 59102 sequences) using the default parameters and enabling the options label-free

quantification (LFQ) and match between runs. Data filtering and statistical analysis were conducted

with the Perseus 1.6.14 software [18]. Only proteins that were identified and quantified with LFQ

intensity values in at least three (out of five) replicates (within at least one of the two experimental

groups) were used for downstream analysis. Missing values were replaced from normal distribution

(imputation) using the default settings.

Statistical analysis.

Statistical differences in protein expression levels (LC-MS) were determined using Student's t-tests

with a permutation-based False Discovery Rate (FDR) of 0.05 implemented in the Perseus software

[18]. Proteins with a minimum 2-fold intensity change compared to control (log2 fold-change ≥ 2 or

log2-fold change ≤ -2) and a *p*-value ≤ 0.05 (FDR adjusted *p*-value) were considered significantly

affected.

To compare the blood markers, we employed Student's t-tests to assess the significance of

differences between two groups (AHDS and control) when variances were found to be equal through

the Levene's test. If variances were unequal, Welch's test was utilised instead. All *p*-values less than

or equal to 0.05 were considered significant. These test were performed using R computational

language platform in R-studio version 2023.09.0+463 [19].

Results

Breeds represented in this study in the AHDS group included Crossbreed (4), Chihuahua (3), Maltese

(2), Pomeranian (2), Yorkshire Terrier (2), Chinese Crested (1), English Bulldog (1), Golden Retriever

(1), Havanese (1), Labrador Retriever (1), Poodle (1) and Schnauzer (1). The control group included

Crossbreed (8), Poodle (3), German Shepherd Dog (2), English Toy Terrier (1), French Bulldog (1),

Golden Retriever (1), Husky (1), Labrador (1), Pomeranian (1) and Västgötaspets (1).

The median age of the dogs in the study group was 3.5 years (1–11 years), and the age of the dogs

in the control group was 4.5 years (1-10 years). The mean body weight was 5.3kg (1.5-27.5kg), while

the control group weighed 15.9 kg (3.5-33kg). We included 12 female (6 intact and 6 neutered) and

8 male (5 neutered and 3 intact) dogs in the AHDS group. In the control group 14 female (9 intact

and 5 neutered) and 6 male (1 intact and 5 neutered) dogs were examined.

All the dogs in the control group were regarded as clinically healthy and showed no abnormalities at

physical examination. Of the AHDS dogs the most common reported presenting complains besides

diarrhoea were vomiting (75%, 15/20), apathy (45%, 9/20) and anorexia (45%, 9/20) and the most

frequent clinical signs were tachycardia (>100beats/min) (85%, 17/20) and abdominal discomfort on

palpation (85%, 17/20). Complete blood count and a plasma biochemistry panel containing glucose

level, urea, creatinine, total proteins (Tp), albumin, ALT, and ALKP, were carried out for all dogs except

one from the AHDS group.

A comparative analysis of blood cell counts between the AHDS and control groups revealed

significant differences in various markers (Figure 1). The AHDS group exhibited a significantly lower

mean corpuscular volume (MCV) with $p=0.0019$ along with reduced lymphocyte counts ($p = 0.0013$)

and eosinophil counts ($p=0.013$). In contrast, sickened dogs demonstrated a higher abundance of

leukocytes ($p=0.00024$), monocytes ($p=0.0061$), and neutrophils ($p=0.0009$) than the control group.

The rest of the blood cell markers were non-significant (see Figure S1).

Similar to the changes in the complete blood cell count, also the biochemistry panel shared

significant differences between the AHDS cohort and healthy controls. AHDS patients displayed a

notable decrease in the abundance of total proteins (TP, $p=<0.001$), also corroborated by lower levels

of albumin ($p=0.00022$). Conversely, AHDS dogs exhibited elevated levels of glucose ($p= 0.0019$) and

alkaline phosphatase ($p= 0.033$). These findings are summarised in Figure 2, while the results for

other plasma biomarkers can be found in Figure S2.

Figure 1. Boxplot analysis of significant blood cell markers in the AHDS and control groups. This figure presents boxplots showcasing the significant variations in blood cell markers observed between the AHDS and control groups. Statistical differences were determined using either the student's t-test for samples with equal variance or the Welch test for samples exhibiting differing variances. Significance levels were established as *p* < 0.05, significant *; *p* < 0.01**, highly significant; and *p* < 0.001**, very significant.

Figure 2. Boxplot analysis of significant biochemistry parameters in the AHDS and control groups. This figure provides a boxplot analysis depicting significant variations in select plasma biochemistry markers between the AHDS and control groups. Statistical differences were assessed using either the student's t-test for samples with equal variance or the Welch test for samples displaying varying variances. Significance levels were established as $p < 0.05$, significant *; $p < 0.01$**, highly significant; and $p < 0.001$***, extremely significant.

Next, we used label-free quantification mass spectrometry to analyse canine plasma from patients affected AHDS in comparison with healthy controls. This study aims to uncover average traits or typical patterns in AHDS with the potential for discovering new biomarkers. The experiment was conducted using pooled samples, comprising ten independent plasma pools (five from AHDS samples and five from control samples), each collected from four randomly selected individuals.

A comprehensive examination of the data revealed that the pools from AHDS patients exhibited a distinct proteome signature compared to those from healthy dogs. Principal component analysis of all quantified proteins from both groups demonstrated the segregation of samples into two separate clusters, indicating the sensitivity of the proteomic analysis in capturing these differences (see Figure 3A).

Among the 207 quantified proteins, we identified 95 proteins exhibited elevated levels, and 67 showed diminished abundance in the plasma of AHDS-affected individuals compared to healthy

dogs, highlighting the discernible impact of the disease (Figure 3B, Table S1). Considering these

findings, our focus is directed toward proteins that demonstrated the most significant increments

or reductions associated with AHDS.

Notably, within the group of proteins increased in AHDS, the following top 8 proteins stand out:

SERPINA3, Lipopolysaccharide-binding protein (encoded by the locus LOC485869), Ribosomal

protein lateral stalk subunit P1 (RPLP1), Actin alpha 1 of skeletal muscle (ACTA1), Glyceraldehyde-

3-phosphate dehydrogenase (GAPDH), Serum amyloid A (SAA1), Inter-alpha-trypsin inhibitor heavy

chain 4 (ITIH4) and Lactotransferrin (LTF).

271 **Figure 3.** (A) Principal Component Analysis (PCA) of quantitative proteome analysis reveals the distinct separation
272 between healthy dogs and those affected by AHDS. (B) Volcano plots were generated to compare plasma pools from
273 healthy dogs with those from five AHDS-affected individuals (n=4 each). Gene names were employed instead of protein
274 names. Upregulated proteins are denoted by green dots, signifying higher abundance in AHDS dogs compared to healthy
275 ones (p-value < 0.05 and a log2 fold-change greater than 2). Conversely, downregulated proteins are indicated by red

dots, indicating lower abundance in AHDS group compared to the control (*p*-value < 0.05 and a log2 fold-change smaller than -2). Non-significantly differentially expressed proteins are depicted as grey dots. A comprehensive list of identified and quantified proteins is available in Table S1.

Notably, a distinct case exists within our dataset - the variable region of lambda light chains, a pivotal component within all immunoglobulins, exhibited altered levels in AHDS samples compared to the control group. These proteins exhibit notable diversity, and in our study, we identified 14 distinct protein variants showing higher abundance in AHDS (see Table S1). In a converse trend, two variants experienced downregulation. These specific protein groupings have been denoted as IGLV@, as referenced in specific databases [20].

Other proteins that exhibited significantly increased abundance in AHDS include Serum amyloid A (SAA1), Inter-alpha-trypsin inhibitor heavy chain 4 (ITIH4), and Lactotransferrin (LTF).

Proteins with a significantly increased abundance in the control group, or conversely, a lower abundance in the AHDS group, comprised Pregnancy zone protein-like (encoded by the gene LOC611458), Paraoxonase (PON1), Selenoprotein P (SELENOP), Insulin-like growth factor binding protein acid labile (IGFALS), and two Amine oxidases (AOC3 and LOC112655648). Other proteins that were depleted in the AHDS group involved Apolipoprotein M (APOM), complement component 4 binding protein alpha (C4BPA), and coagulation factor XII (F12), Carboxypeptidase N (CPN2), Sex hormone binding globulin (SHBG), and the HGF activator (HGFAC).

Using the dataset of quantified proteins, we conducted gene ontology enrichment analysis [21] to investigate the functional significance of differentially expressed proteins (DEPs) in AHDS patients compared to the healthy control group (Figure 4). The analysis highlighted some homologous proteins related to Alzheimer's disease in humans, as well as Parkinson's and Huntington's diseases. Additionally, notable protein classes in both upregulated and downregulated groups included coagulation-related proteins such as those involved in the plasminogen activation cascade and inflammation-related proteins.

Figure 4. The figure illustrates the Gene Ontology (GO) analysis results conducted on the plasma proteome of dogs affected by AHDS. The analysis categorised the proteins according to their pathways using Panther software [21]. The pie charts provide an overview of the distribution of proteins based on their assigned Gene Ontology terms. Each slice represents a specific functional category, and the numbers represent the number of hits or proteins implicated in a particular category.

We have utilised a network analysis rooted in protein-protein interactions, drawing upon functional insights sourced from the STRING database, to visually depict the overarching protein alterations in the plasma of ADHS patients. This thorough examination encompasses both direct (physical) and indirect (functional) associations among proteins [22]. To elucidate the impact of AHDS on plasmatic protein levels, we superimposed our proteomic datasets onto the known protein-protein interaction database for dogs. Notably, the proteins exhibiting up-regulation and down-regulation in AHDS displayed a remarkable degree of interconnectivity (Figure 5). This striking observation hints at extensive proteome-wide adjustments in response to the pathological condition, offering insight into the modifications within various protein clusters.

Figure 5. Network visualization of proteins overrepresented (green) and depleted (red) in plasma of dogs with AHDS
compared to the plasma of healthy dogs using STRING database [23]. The network diagram depicts the interactions
between various canine plasma proteins using as inclusion a significant FDR-adjusted *p*-value regardless of the level of
expression. Each node represents a unique protein, and the edges connecting nodes indicate known or predicted
interactions. Node colour indicates the type of interaction evidence, ranging from experimental data to curated
databases and text mining.

Discussion

Collecting intestinal tissue samples from animals can be challenging in most cases for ethical or clinical issues. However, when it comes to investigating the development of diarrhoea in animals, plasma emerges as an optimal choice for sampling. This preference arises from its comparatively stable chemical and physical properties, rich protein content, and relatively reliable collection techniques [24]. We can add to these facts that since haemorrhagic diarrhoea affects the overall physiological functioning of the body, plasma protein profiling offers valuable insights into these perturbations, making it a very suitable material for diagnosis and biomarker discovery.

It is important to note that in cases of intestinal blood loss, patients experience a depletion of electrolytes and water, causing haemoconcentration, accompanied by the loss of specific proteins that impact the whole body's homeostasis [25]. This explains why the total protein and albumin levels in AHDS dogs are lower compared to the control group although they are haemoconcentrated. These haemoconcentration and selective protein losses trigger a reactive response that in turn affects the levels of other proteins. As a result, the findings presented here may reflect a delicate trade-off among these factors and the overall homeostatic disruption observed in AHDS patients.

The classical clinical blood analysis revealed significant imbalances in blood cell counts and plasma biochemistry. The discrepancies can be partially attributed to factors such as intestinal blood loss and electrolyte imbalances, which result in haemoconcentration [4]. Neutrophilia and monocytosis as well as a rise in blood glucose fit to the picture of a stress leukogram, triggered by the effects of adrenalin and cortisol during acute illness, but can also be seen in systemic inflammation in general. In some instances, these factors may lead to alterations that either overestimate or underestimate the quantification of specific biomarkers. This phenomenon arises from the complex interplay of various mechanisms within the organism. For instance, the body may attempt to replenish specific blood components when possible [26]. At the same time, alterations in other parameters can relate to immune responses or the expenditure of proteins, such as coagulation factors.

 Some of the altered proteins found in the ADHS samples are already known to be involved in other gastrointestinal disorders. For instance, an increase of the amine oxidases (AOC3 and AOX2) is considered an indicator of intestinal injury caused by some anti-cancer drug treatments [27–29]. Another good example is the protein SERPINA3, the protein which showed the strongest increase in the ADHS plasma samples compared to healthy controls. SERPINA3, also known as Alpha-1-antichymotrypsin (ACT), belongs to the serine protease inhibitor (serpin) family. Its primary function is to regulate and inhibit the activity of certain proteases, specifically chymotrypsin-like enzymes. SERPINA3 acts as an inhibitor of chymotrypsin-like proteases. By binding to these proteases and

forming a stable complex, SERPINA3 helps to prevent excessive proteolysis, which is essential for maintaining proper protein balance and preventing tissue damage [30]. In some inflammatory processes, SERPINA3 is also implicated in the modulation of certain components of the immune system and cytokines, affecting immune responses and inflammation. Changes in SERPINA3 levels have been associated with various inflammatory conditions and diseases. SERPINA3 may also play a role in regulating fibrinolysis, the breaking down of blood clots. It interacts with plasmin, an enzyme involved in fibrinolysis, and can influence the balance between clot formation and clot dissolution [31]. SERPINA3 is also involved in tissue repair and remodelling, modulating tissue responses to injury and promoting healing. Some studies have pointed out that SERPINA3 levels may be altered in specific cancer types and neurodegenerative diseases [30].

Another highly elevated protein is the RPLP1, which belongs to the ribosomal P complex, which consists of the acidic ribosomal P proteins. The presence of this intracellular protein in the bloodstream may serve as an indicator of significant tissue damage and the release of cellular contents into the circulatory system in AHDS affected dogs. RPLP1 overexpression not only immortalises primary cells but also plays a significant role in their transformation. Furthermore, there is evidence of RPLP1 protein being upregulated in certain human cancers [32]. In the same direction, it is known that Glyceraldehyde-3-phosphate dehydrogenase, another upregulated protein in AHDS, is overexpressed in colorectal cancer onset [33]. These findings could be another link between gastrointestinal affection like AHDS and the development of future tumours, which is another reason to enhance better understanding and management of the disease.

It is also intriguing to speculate about the relationship between some of the protein classes that exhibited altered levels in this study and their roles in conditions associated with the brain, including Alzheimer's disease, Parkinson's disease, and Huntington's disease as shown here with the gene ontology analysis. For instance, older dogs often develop canine cognitive dysfunction, a condition that shares many similarities with Alzheimer's disease. Dogs exhibit cognitive deficits that can be compared to human symptoms, such as disorientation, memory loss, changes in behaviour, and the presence of beta-amyloid plaques, which are commonly detected in both the extracellular space as senile plaques and around blood vessels in their brains. [34]. Various clinical, epidemiological, and immunological findings provide strong indications that the gut-brain axis is significantly and deeply impacted by alterations of gut microbiome [35].

Managing diarrhoea in dogs, whether related to conditions like AHDS or similar ailments, holds significance for the well-being of animal companions and human health, as coined by the concept of One Health [36]. Certain microorganisms found in these faecal deposits could serve as sources of

zoonotic transmission for parasites such as *Giardia* or potential bacterial pathogens like *Campylobacter*, *E. coli*, and *Salmonella*. Hence, understanding and controlling diarrhoea in dogs is crucial for safeguarding both animal and human health.

The study's reliance on a relatively small cohort of just 40 participants raises some limitations that must be considered when interpreting the results. The sample size may not provide a comprehensive representation of the broader population, limiting the generalizability of the findings. Furthermore, an additional challenge arises from the noticeable differences in body weight between the groups within the study, and the breed are not exactly matching. These variations can introduce confounding variables that may obscure some effects of the factors under investigation. Differences in breed and body weight can influence the outcomes in ways that are unrelated to our primary research question, potentially leading to skewed or biased results. Therefore, while the study provides valuable insights, these limitations should be considered when drawing conclusions or extrapolating the findings to a larger and more diverse population.

Our study opens several promising avenues for future research by uncovering the alteration of several proteins in plasma that were not previously reported in the literature for AHDS. First, we need to investigate further the role of the identified biomarker candidates in other diarrhoeic diseases to understand better if altered proteins are specific for AHDS or if changes in their expression levels also occur in other confounding conditions. A comparison with the proteomic profile in diseases like Parvovirosis or "inflammatory bowel disease" (IBD) would be a possible next step. Second, we should explore the connection between severe intestinal mucosal damage, barrier dysfunction, and the development of other chronic gastrointestinal diseases.

To improve the AHDS diagnostic accuracy and treatment strategies, future studies should focus on refining and expanding the pool of potential biomarkers we have identified here. Considering a broader range of canine populations and disease stages, additional studies are required to extend the validity of our results. Collaborative efforts to validate and translate these candidates into clinical practice could ultimately improve the prognosis and management of AHDS in dogs. Taken together, our research provides a foundation upon which future investigations can continue to advance our understanding and enhance the care of dogs affected by AHDS. To the best of our knowledge, this is the first study that employs a quantitative plasma proteomic approach to gain a deeper understanding of AHDS.

Conclusions

In conclusion, acute haemorrhagic diarrhoea syndrome in dogs presents a significant clinical
challenge, often leading to severe health complications. While several factors, including *Clostridium*
perfringens toxins, have been implicated in its development, the exact pathogenesis of AHDS
remains elusive. Our study took a comprehensive approach, exploring the proteomic profiles of dogs
afflicted with AHDS and comparing them to reasonably well-matched control dogs. To the best of
our knowledge, this is the first study that employs a quantitative plasma proteomic approach to gain
a deeper understanding of AHDS. This in-depth analysis uncovered substantial alterations in the
levels of numerous proteins present in plasma, shedding light on potential biomarkers for diagnostic
and therapeutic advancements and possibly therapeutic interventions. SerpinA3,
Lipopolysaccharide-binding protein, Ig-like domain-containing proteins, Glyceraldehyde-3-
phosphate dehydrogenase, and Serum amyloid A emerged as candidates with increased abundance
in AHDS-affected dogs. In contrast, others exhibited decreased levels, such as Paraoxonase,
Selenoprotein, Amine oxidases, and Apolipoprotein C-IV.

Acknowledgements. For mass spectrometry, we would like to acknowledge the assistance of the
Core Facility BioSupraMol supported by the Deutsche Forschungsgemeinschaft (DFG).

Ethical Statement. This investigation was carried out with surplus samples from previous studies.
These previous studies underwent rigorous scrutiny and received approval from the institutional
ethical committee at the Veterinary University of Vienna Research (BMBWF-68.205/0216-
V/3b/2019, 2021-0.518.574).

Data Availability. We have included all data generated or analysed during this comprehensive study
within this published article and its supplementary information files.

References

1. Pugh, C.A.; Bronsvoort, B.M. de C.; Handel, I.G.; Querry, D.; Rose, E.; Summers, K.M.; Clements, D.N. Incidence Rates and Risk Factor Analyses for Owner Reported Vomiting and Diarrhoea in Labrador Retrievers - Findings from the Dogslife Cohort. *Prev. Vet. Med.* **2017**, *140*, 19–29, doi:10.1016/J.PREVETMED.2017.02.014.

2. Burrows, C.F. Canine Hemorrhagic Gastroenteritis [Diseases of Unknown Etiology, Dogs]. *Journal-American Anim. Hosp. Assoc.* **1977**, *13*, 451–458.

3. Mortier, F.; Strohmeyer, K.; Hartmann, K.; Unterer, S. Acute Haemorrhagic Diarrhoea Syndrome in Dogs: 108 Cases. *Vet. Rec.* **2015**, *176*, 627, doi:10.1136/VR.103090.

4. Busch, K.; Unterer, S. Update on Acute Hemorrhagic Diarrhea Syndrome in Dogs. *Adv. Small Anim. Care* **2022**, *3*, 133–143, doi:10.1016/j.yasa.2022.06.003.

5. Unterer, S.; Lechner, E.; Mueller, R.S.; Wolf, G.; Straubinger, R.K.; Schulz, B.S.; Hartmann, K. Prospective Study of Bacteraemia in Acute Haemorrhagic Diarrhoea Syndrome in Dogs. *Vet. Rec.* **2015**, *176*, 309, doi:10.1136/VR.102521.

6. Sindern, N.; Suchodolski, J.S.; Leutenegger, C.M.; Mehdizadeh Gohari, I.; Prescott, J.F.; Proksch, A.L.; Mueller, R.S.; Busch, K.; Unterer, S. Prevalence of Clostridium Perfringens NetE and NetF Toxin Genes in the Feces of Dogs with Acute Hemorrhagic Diarrhea Syndrome. *J. Vet. Intern. Med.* **2019**, *33*, 100–105, doi:10.1111/JVIM.15361.

7. Unterer, S.; Strohmeyer, K.; Kruse, B.D.; Sauter-Louis, C.; Hartmann, K. Treatment of Aseptic Dogs with Hemorrhagic Gastroenteritis with Amoxicillin/Clavulanic Acid: A Prospective Blinded Study. *J. Vet. Intern. Med.* **2011**, *25*, 973–979, doi:10.1111/J.1939-1676.2011.00765.X.

8. Unterer, S.; Busch, K. Acute Hemorrhagic Diarrhea Syndrome in Dogs. *Vet. Clin. North Am. Small Anim. Pract.* **2021**, *51*, 79–92, doi:10.1016/J.CVSM.2020.09.007.

9. Sänger, F.; Unterer, S.; Werner, M.; Dörfelt, R. C-Reactive Protein as a Tool for Monitoring Response to Treatment in Dogs with Acute Hemorrhagic Diarrhea Syndrome. *Front. Vet. Sci.* **2023**, *9*, 1019700, doi:10.3389/FVETS.2022.1019700/BIBTEX.

10. Skotnitzki, E.; Suchodolski, J.S.; Busch, K.; Werner, M.; Zablotski, Y.; Ballhausen, B.D.; Felix Neuerer, |; Unterer, S. Frequency of Signs of Chronic Gastrointestinal Disease in Dogs after an Episode of Acute Hemorrhagic Diarrhea. **2021**, doi:10.1111/jvim.16312.

11. Hubbard, K.; Skelly, B.J.; Mckelvie, J.; Wood, J.L.N. Risk of Vomiting and Diarrhoea in Dogs. *Vet. Rec.* **2007**, *161*, 755–757, doi:10.1136/VR.161.22.755.

12. Stavisky, J.; Pinchbeck, G.L.; German, A.J.; Dawson, S.; Gaskell, R.M.; Ryvar, R.; Radford, A.D. Prevalence of Canine Enteric Coronavirus in a Cross-Sectional Survey of Dogs Presenting at Veterinary Practices. *Vet. Microbiol.* **2010**, *140*, 18–24, doi:10.1016/J.VETMIC.2009.07.012.

13. Jugan, M.C.; KuKanich, K.; Freilich, L. Clinical Response in Dogs with Acute Hemorrhagic Diarrhea Syndrome Following Randomized Probiotic Treatment or Fecal Microbiota Transplant. *Front. Vet. Sci.* **2023**, *10*, 1050538, doi:10.3389/FVETS.2023.1050538/BIBTEX.

14. Ceciliani, F.; Eckersall, D.; Burchmore, R.; Lecchi, C. Proteomics in Veterinary Medicine: Applications and Trends in Disease Pathogenesis and Diagnostics. *Vet. Pathol.* **2014**, *51*, 351–362, doi:10.1177/0300985813502819.

15. Doulidis, P.G.; Kuropka, B.; Ramos, C.F.; Rodríguez-Rojas, A.; Burgener, I.A. Label-Free Quantification LC-Mass Spectrometry Proteomic Analysis of Blood Plasma in Healthy Dogs. *bioRxiv* **2023**, 2023.06.13.544743, doi:10.1101/2023.06.13.544743.

16. Rappsilber, J.; Mann, M.; Ishihama, Y. Protocol for Micro-Purification, Enrichment, Pre-Fractionation and Storage of Peptides for Proteomics Using StageTips. *Nat. Protoc.* **2007**, *2*, 1896–1906, doi:10.1038/nprot.2007.261.

17. Tyanova, S.; Temu, T.; Carlson, A.; Sinitcyn, P.; Mann, M.; Cox, J. Visualization of LC-MS/MS

Proteomics Data in MaxQuant. *Proteomics* **15**, 1453–1456, doi:10.1002/pmic.201400449.

18. Tyanova, S.; Temu, T.; Sinitcyn, P.; Carlson, A.; Hein, M.Y.; Geiger, T.; Mann, M.; Cox, J. The
Perseus Computational Platform for Comprehensive Analysis of (Prote)Omics Data. *Nat.*
Methods **2016**, *13*, 731–740, doi:10.1038/nmeth.3901.

19. R Core Team *R: A Language and Environment for Statistical Computing.*; R Foundation for
Statistical Computing: Vienna, Austria, 2017;

20. IGLV@ Immunoglobulin Lambda Variable Cluster [Homo Sapiens (Human)] - Gene - NCBI
Available   online:
https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=DetailsSearch&Term=3546   (accessed
on 8 August 2023).

21. Mi, H.; Muruganujan, A.; Ebert, D.; Huang, X.; Thomas, P.D. PANTHER Version 14: More
Genomes, a New PANTHER GO-Slim and Improvements in Enrichment Analysis Tools. *Nucleic*
Acids Res. **2019**, *47*, D419–D426, doi:10.1093/nar/gky1038.

22. Szklarczyk, D.; Gable, A.L.; Lyon, D.; Junge, A.; Wyder, S.; Huerta-Cepas, J.; Simonovic, M.;
Doncheva, N.T.; Morris, J.H.; Bork, P.; et al. STRING V11: Protein-Protein Association Networks
with Increased Coverage, Supporting Functional Discovery in Genome-Wide Experimental
Datasets. *Nucleic Acids Res.* **2019**, *47*, D607–D613, doi:10.1093/nar/gky1131.

23. Szklarczyk, D.; Franceschini, A.; Wyder, S.; Forslund, K.; Heller, D.; Huerta-Cepas, J.; Simonovic,
M.; Roth, A.; Santos, A.; Tsafou, K.P.; et al. STRING V10: Protein–Protein Interaction Networks,
Integrated   over   the   Tree   of   Life.   *Nucleic   Acids   Res.*   **2015**,   *43*,   D447–D452,
doi:10.1093/nar/gku1003.

24. Yan, Z.; Zhang, K.; Wang, G.; Wang, L.; Zhang, J.; Qiu, Z.; Guo, Z.; Zhang, K.; Li, J. Differential
Proteomic of Plasma Provides a New Perspective on Scientific Diagnosis and Drug Screening
for   Dampness   Heat   Diarrhea   in   Calves.   *Front.   Vet.   Sci.*   **2022**,   *9*,
doi:10.3389/FVETS.2022.986329/FULL.

25. Camilleri, M.; Sellin, J.H.; Barrett, K.E. Pathophysiology, Evaluation, and Management of
Chronic   Watery   Diarrhea.   *Gastroenterology*   **2017**,   *152*,   515,
doi:10.1053/J.GASTRO.2016.10.014.

26. Field, M. Intestinal Ion Transport and the Pathophysiology of Diarrhea. *J. Clin. Invest.* **2003**,
111, 931, doi:10.1172/JCI18326.

27. Sato, Y.; Tanaka, Y.; Imai, T.; Okumura, N.; Matsuhashi, N.; Takahashi, T.; Shimokawa, T.;
Yoshida, K. Serum Diamine Oxidase Activity Derived from Response to Chemotherapy Affects
Adverse Events and Serum Amino Acid Levels. *Support. Care Cancer* **2022**, *30*, 9369–9377,
doi:10.1007/s00520-022-07362-2.

28. Moriyama, K.; Kouchi, Y.; Morinaga, H.; Irimura, K.; Hayashi, T.; Ohuchida, A.; Goto, T.;
Yoshizawa, Y. Diamine Oxidase, a Plasma Biomarker in Rats to GI Tract Toxicity of Oral
Fluorouracil   Anti-Cancer   Drugs.   *Toxicology*   **2006**,   *217*,   233–239,
doi:10.1016/j.tox.2005.09.017.

29. Goto, T.; Matsubara, T.; Yoshizawa, Y.; Sasaya, S.; Nemoto, H.; Sanada, Y.; Moriyama, K.; Kouchi,
Y. Diamine Oxidase as Blood Biomarker in Rats and Humans to G1 Tract Toxicity of Fluorouracil
Anti-Cancer Drugs. *Japanese J. Cancer Chemother.* **2011**, *38*, 765–769.

30. de Mezer, M.; Rogaliński, J.; Przewoźny, S.; Chojnicki, M.; Niepolski, L.; Sobieska, M.;
Przystańska, A. SERPINA3: Stimulator or Inhibitor of Pathological Changes. *Biomed. 2023, Vol.*
11, Page 156 **2023**, *11*, 156, doi:10.3390/BIOMEDICINES11010156.

31. Chapin, J.C.; Hajjar, K.A. Fibrinolysis and the Control of Blood Coagulation. *Blood Rev.* **2015**,
29, 17, doi:10.1016/J.BLRE.2014.09.003.

32. Artero-Castro, A.; Perez-Alea, M.; Feliciano, A.; Leal, J.A.; Genestar, M.; Castellvi, J.; Peg, V.;
Ramón y Cajal, S.; Lleonart, M.E. Disruption of the Ribosomal P Complex Leads to Stress-
Induced Autophagy. *Autophagy* **2015**, *11*, 1499, doi:10.1080/15548627.2015.1063764.

33.   Tarrado-Castellarnau, M.; Diaz-Moralli, S.; Polat, I.H.; Sanz-Pamplona, R.; Alenda, C.; Moreno,
586   V.; Castells, A.; Cascante, M. Glyceraldehyde-3-Phosphate Dehydrogenase Is Overexpressed in
587   Colorectal Cancer Onset. *Transl. Med. Commun. 2017 21* **2017**, *2*, 1–9, doi:10.1186/S41231-
588   017-0015-7.
589   34.   Mihevc, S.P.; Majdic, G. Canine Cognitive Dysfunction and Alzheimer's Disease – Two Facets
590   of the Same Disease? *Front. Neurosci.* **2019**, *13*, doi:10.3389/FNINS.2019.00604.
591   35.   Appleton, J. The Gut-Brain Axis: Influence of Microbiota on Mood and Mental Health. *Integr.*
592   *Med. A Clin. J.* **2018**, *17*, 28.
593   36.   WHO One Health GLOBAL Available online: https://www.who.int/health-topics/one-
594   health#tab=tab_1 (accessed on 7 September 2022).
595   37.   Overgaauw, P.A.M.; Vinke, C.M.; van Hagen, M.A.E.; Lipman, L.J.A. A One Health Perspective
596   on the Human-Companion Animal Relationship with Emphasis on Zoonotic Aspects. *Int. J.*
597   *Environ. Res. Public Health* **2020**, *17*, doi:10.3390/IJERPH17113789.