**Habitat remediation followed by managed connectivity reduces unwanted changes in evolutionary trajectory of high extirpation risk populations**

Gina F. Lamka[1] and Janna R. Willoughby[1]

1. College of Forestry, Wildlife, and Environment, 602 Duncan Dr., Auburn University, Auburn, AL 36849

**ABSTRACT**

As we continue to convert green spaces into roadways and buildings, connectivity between

populations and biodiversity will continue to decline. In threatened and endangered species, this

trend is particularly concerning because the cessation of immigration can cause increased

inbreeding and loss of genetic diversity, leading to lower adaptability and higher extirpation

probabilities in these populations. Unfortunately, monitoring changes in genetic diversity from

management actions such as assisted migration and predicting the extent of introduced genetic

variation that is needed to prevent extirpation is difficult and costly in situ. Therefore, we

designed an agent-based model to link population-wide genetic variability and the influx of

unique alleles via immigration to population stability and extirpation outcomes. These models

showed that management of connectivity can be critical in restoring at-risk populations and

reducing the effects of inbreeding depression; increased connectivity prevented extirpation for

the majority of scenarios we considered (71.5% of critically endangered populations and 100%

of endangered and vulnerable populations). However, the rescued populations were more similar

to the migrant source population (average $F_{ST}$ range 0.05 – 0.10) compared to the historical

recipient population (average $F_{ST}$ range 0.23 – 0.37). This means that these management actions

not only recovered the populations from the effects of inbreeding depression, but they did so in a

way that changed the evolutionary trajectory that was predicted and expected for these

populations prior to the population crash. This change was most extreme in populations with the

smallest population sizes, which are representative of critically endangered species that could

reasonably be considered candidates for restored connectivity or translocation strategies.

Understanding how these at-risk populations change in response to varying management

interventions has broad implications for the long-term adaptability of these populations and can

improve future efforts for protecting locally adapted allele complexes when connectivity is

restored.

**INTRODUCTION**

Habitat fragmentation is increasing globally, a concern for sustainability because habitat

fragmentation impairs ecosystem functions and reduces biodiversity up to 75% (Haddad et al.

2015). As habitat fragmentation continues to encroach on wild environments, connectivity

between populations will remain a top priority for population management, particularly for

threatened and endangered species due to the often-existing concerns of decreased genetic

diversity and increased inbreeding and the potential for these concerns to be exacerbated by

increased fragmentation. However, the minimum connectivity required to support population

recovery is difficult to predict because the number of migrants needed varies with population

characteristics such as extinction risk, migration rate, and genetic makeup of the populations.

Here, we quantify these interactions to understand how to maintain genetic variation in

recovering populations when repairing corridors or otherwise encouraging movement of genetic

variants is desirable.

When genetic diversity is diminished and inbreeding increases, populations often face increased

inbreeding depression through the expression of recessive traits and accumulation of maladapted

alleles (Coltman and Slate 2003; Del Castillo et al. 2011). Small populations, in particular, are

often at high extinction risk due to these effects because they are typically isolated and lose

genetic variants quickly through drift (Caughley 1994). Furthermore, populations with high

levels of inbreeding depression can enter an extinction vortex when inbreeding depression and

genetic drift reduce genetic diversity and fitness in a positive feedback loop (Caughley 1994;

Drake et al. 2006; Drake 2008; Braumann 2010; Messer 2016). However, increasing or restoring

connectivity can alleviate these stressors by introducing new genetic variants that mask

deleterious recessive alleles, thereby decreasing inbreeding depression.

Theoretically, connectivity of subpopulations via one migrant per population per generation is

sufficient to retain genetic variation while allowing divergence in allele frequencies; empirically,

however, the minimum of one individual per population per generation is often insufficient to

achieve neutral or positive population growth for struggling populations (Mills and Allendorf

1996). As such, the genetic connectivity of passing alleles via immigration must also

complement demographic connectivity so that the benefits of migrants are reflected in population

growth and stability, a goal especially important for management of fragmented and threatened

species (Lowe and Allendorf 2010). The benefits of migration – either naturally with corridors or

via artificial translocations of individuals from one area to another – depend on the carrying

capacity of the source and recipient populations, the rate of migration, the growth rate of the

recipient population, the reproductive success of migrants, the competition for resources against

other species, and the frequency of repeated migration such that the outcome of immigration

events are dependent on examination of these factors before using migration as a conservation

tool (Thrimawithana et al. 2013; Backus and Baskett 2021).

Supplementing populations can help sustain populations long term, but at the expense of a

decreased effective population size – and therefore decreased diversity in the gene pool – for the

first few generations (Ryman-Laikre effect; Ryman and Laikre 1991). This drop in the number of

effective breeders in the first generations post-migration arises because the reproductive rate of

new individuals tends to be favored, leading to variance in mean fitness between receiving and

migrant source lineages (Hagen et al. 2019). This effect is especially pronounced when a greater

proportion of the migrating population contains new genetic variants than the receiving

population (Christie et al. 2012; Hagen et al. 2020), so identifying the number of new individuals

to introduce can be a balancing act between increasing the number of adults while attempting to

maintain genetic diversity and fitness. Therefore, establishing if an increase in population size or

population fitness is the conservation goal and post-release monitoring of population-wide

reproductive success, maintaining an equal sex ratio, and evaluating kinship among offspring is

especially important in the first years following assisted migrations.

Because funding is nearly always a limiting factor, most agencies must consider management

strategies and the time and resources that population managers must devote to sustaining the

species (Waldron et al. 2013; White et al. 2022). Translocation actions for large carnivores can

cost at least a few thousand dollars per individual (Weise et al. 2014), with this cost increasing

significantly when accounting for unsuccessful reintroductions (Weise et al. 2014). Further,

tracking technology, post-translocation monitoring, and other mitigation strategies constitute a

higher proportion of project funds than the translocations themselves (Weise et al. 2014; Julien et

al 2022) and as a result, some recovery programs choose to focus on data collection rather than

action (Buxton et al. 2020). Delaying or shortening the frequency of assisted migration efforts

can decrease financial costs, but at a risk of decreased genetic diversity and long-term viability

(Martinez-Abrain et al. 2011; Hilbers et al. 2019). Although many have proposed frameworks for

species management via translocations (McLachlan et al. 2007; Schwartz and Martin 2013;

Williams and Dumroese 2013; Sansilvestri et al. 2015; Barbosa and Tella 2019; Berger-Tal et al.

2019) and for monitoring genetic diversity (Weeks et al. 2011; Furlan et al. 2020; Chen et al.

2021; Hogg et al. 2021; DeWoody et al. 2023), implementation across taxa is still lacking. Thus,

identifying how we can most efficiently use funding to support at-risk populations by restoring

migration patterns is important for effective management.

Oftentimes, management practices have the goal of both increasing population size and genetic

diversity by restoring gene flow via outbreeding with genetically distinct individuals (Weeks et

al. 2011; Stepkovitch et al. 2022). Consistently monitoring the genetic diversity of wild

populations – both vulnerable and otherwise – before and after management intervention is both

costly and difficult *in situ*. Here, we establish a model that can simulate genetic diversity across

hundreds of years to evaluate the risks and benefits of increasing migration into high extinction

risk populations. Using simulations, we address two questions relevant to understanding

evolutionary trends in populations of threatened and endangered species. First we answer the

question: How do genetic characteristics of the migrant source population, specifically the

population-specific unique genetic variants that determine individual fitness, influence recipient

population stability and evolution? We hypothesized that populations receiving migrants with

lower heterozygosity (and were accordingly less fit) would be less stable over time compared to

populations receiving more fit migrants that have more genetic variants, since low fitness

individuals may have reduced effects on the recipient populations' gene pool. The second

question we address is: How does recipient population size (e.g., carrying capacity) and trend

(e.g., increasing, decreasing) alter the magnitude of the population's evolutionary response to

immigration? We hypothesized that small recipient populations would represent the migrant

gene pools more than moderately sized recipient populations because periods of rapid population

growth following an influx of migrants would favor new migrant-brought alleles. These aims

will help guide future decisions when considering migration (via corridors or translocations) for

managing species on the brink of extinction.

**METHODS**

*Demographic model steps*

We developed a forward-time agent-based model in R that tracks individuals and their multi-

locus genotypes for 350 years (Figure 1). Parameters were selected and evaluated separately to

examine their independent and interaction effects (Table 1). We incorporated the effects of

migration by creating two populations: a migrant-receiving population and a migrant source

population, from which migrants were randomly selected to disperse into the recipient

population. In all iterations, the recipient populations were allowed to stabilize for 100 years with

a carrying capacity of 1000 individuals (K = 1000) and we then subjected some recipient

populations to a 50-year population size reduction to simulate the effects of environmental

change; this reduction was comprised of a population size decline (i.e., decreasing trend) over 10

years and a subsequent 40-year period of maintenance at this population size minima. We varied

the degree of population size reduction, with three possible reduction proportions corresponding

to IUCN (International Union for Conservation of Nature) classifications: vulnerable species had

a 30% decline in 10 years (vulnerable carrying capacity = 700 individuals); endangered had a

70% decline in 10 years (endangered carrying capacity = 300 individuals); and critically

endangered populations had a 90% decline in 10 years (critically endangered carrying capacity =

100 individuals; Table 1). After the 50-year reduced population period, we simulated habitat

restoration by allowing the populations to grow again (i.e., increasing trend), up to the original

carrying capacity (1000 individuals), and tracked the populations' demographic and evolutionary

responses for 200 years. Additionally, we included populations that did not face environmental

degradation or population decline to serve as controls. Below, we describe the basic model

structure and function, and provide further details about each model function in the Supporting

Information.

Simulations began by initializing the populations so that the recipient and migrant source

populations were at carrying capacity (total recipient population size started at 1000 individuals;

total migrant source population size was 5000 individuals) and all individuals were assigned

1100 SNP genotypes. SNPs were split into two types: neutral and migrant-associated. The 1000

neutral SNPs were randomized as either homozygous or heterozygous according to one of two

minimum allele frequency ranges for the same allele in both populations and were used for

downstream genetic analyses (i.e., $0.05 \leq p \leq 0.15$ or $0.40 \leq p \leq 0.50$; Table 1). The 100 migrant-

associated SNPs were population specific such that all individuals were homozygous for the

source or the alternate recipient population allele and were used to characterize the proportion of

migrant ancestry in the population but do not influence fitness. In subsequent generations, each

allele had a 0.000001% chance of mutating ($\mu = 1.0 \times 10^{-8}$), which is similar to empirical

estimates of mutation (Nachman and Crowell 2000). Starting individuals were randomly

assigned sex and age along a Poisson distribution centered around the age of maturity (1 year). In

subsequent years, each surviving individual was aged by 1 year and newly created individuals

(see reproduction description below) were assigned genotypes following patterns of Mendelian

genetics.

We enacted migration events where random migrants from the source population entered the

receiving population and quantified the demographic and evolutionary response of the recipient

populations. Within each set of parameter values, migrants entered the recipient populations at a

set frequency: 1 migrant per generation, as well as at rates common for assisted migration

management (Anderson et al 2005; Kim et al. 2011; Sasmal et al. 2013; Nash et al. 2020; Morris

et al. 2021) including 100 individuals at a single time period (i.e., 100 individuals in year 151),

which we refer to as burst migration, and 25 individuals at four time periods (i.e., 25 individuals

in years 151, 165, 181, 195), which we refer to as pulsed migration. As a basis for comparison,

we also included scenarios where migration was completely absent (Table 1). For each assisted

migration scenario, we considered the effects of these events both during the population minima

when habitat was still being restored (burst immigration year 125; pulse immigration years 125,

140, 155, 170) and after habitat restoration when the population was permitted to grow (burst

immigration year 151; pulse immigration years 151, 165, 181, 195; Table 1) to determine if

bringing in new individuals before habitat quality is restored would have the same effect on the

genetic diversity of the population as bringing new individuals after habitat management and

population monitoring occurs.

After the migrants entered the population, individuals were randomly paired for mating and

reproduction, and parental pairs that included at least one migrant were preferentially chosen to

produce offspring so that the number of migrants closely approximated the number of effective

migrants; when migration rates were high and the number of breeding individuals was less than

the number of migrants, not all migrants were able to reproduce and, occasionally at small

population sizes, migrants did not breed when opposite sex mates were not available. We

subsequently considered the effects of this migrant-biased procedure and found that there were

similar trends between preferentially choosing migrant parental pairs and completely random

mating without a migrant bias (see Supplemental Materials and Supplementary Figure 1). We

included population demographic effects in our model to reflect the decreased chance of

potential mates interacting with each other at small population sizes (Allee effect; Drake and

Kramer 2011) by reducing the chance of finding mates as population sizes decreased; the

probability of finding a mate was the complement of the reciprocal of the total number of adult

pairs. All mated pairs produced 1-2 offspring, and each year the total number of offspring

produced followed the logistic growth equation:

$$N_{t+1} = \left(1 + r\left(1 - \frac{N_t}{K}\right)\right) * N_t$$

where the population size ($N_{t+1}$) was determined by the per capita growth rate ($r = 1$), carrying

capacity (K = 1000 individuals), and the population size prior to reproduction ($N_t$). Density

independent variation in population size was introduced where the value of $N_{t+1}$ was randomly

generated along a normal distribution with a mean of $N_{t+1}$ and standard deviation of 1.

Individuals were removed from the population in two ways, one relating to fitness effects and

one relating to random environmental or competition effects. To reflect fitness effects, we used

the neutral SNPs (i.e., not including source or migrant-receiving population-specific SNPs) to

force a fitness cost to reduced genetic diversity; when an individual reached maturity, we

removed individuals with a probability equal to the reciprocal of its genomic heterozygosity,

such that there was a higher chance of mortality with decreasing heterozygosity. To reflect other

causes of mortality in our population, each year we assumed the cumulative probability of death

of an individual was equal to the quotient of an individual's age and the maximum lifespan.

After age nine, individuals were removed from the population through this method. Unless the

recipient population size was reduced to 10 or fewer individuals – which we considered

functionally extirpated – these events were repeated for 350 years.

*Genetic model steps*

We devised several monitoring measures for our recipient populations that persisted across all

350 years. Using the 1000 neutral SNPs, we calculated yearly observed and expected

heterozygosity as measures of individual and population-wide genetic diversity by dividing the

number of heterozygous neutral alleles over the total number of neutral alleles. We used the

migrant-associated SNPs to quantify the proportion of migrant loci over the total number of

population-source-associated loci to calculate migrant ancestry in the recipient populations and

quantified the number of mating individuals to monitor population fitness. We quantified

inbreeding ($F_{IS}$; Supplemental Figure 3) using the *hierfstat* package to examine how immigration

of migrants into the recipient populations and random genetic drift changed the progression of

the populations. We analyzed divergence ($F_{ST}$) in two ways: the divergence between the

historical recipient populations in year 0 compared to each consecutive year of the contemporary

recipient populations as well as the historical migrant source populations compared to the

contemporary recipient populations in each year using the *hierfstat* package in R (Goudet 2005;

Weir and Goudet 2017). Along with the number of effective migrants and the number of

effective parents, we also recorded the total number of individuals, sex ratio (Supplemental

Figure 4), and number of adults present in each year. The number of migrants in the population

and the proportion of migrant-specific alleles in the population served as measures of the

demographic and genetic effects of migration on the recipient populations. Finally, the lifetime

reproductive success of all individuals was calculated by determining the number of mates,

number of offspring, and the number of offspring that survived to maturity for all individuals

(Supplemental Figure 5). Significant differences among variables within each year were

determined by comparing confidence intervals (84% quantiles approximating α = 0.05; Payton et

al. 2003). All scripts and figures were created using RStudio (R version 4.2.1) and all scripts are

available at https://github.com/ginalamka/ComplexModel_ABM/.

**RESULTS**

Across simulation runs, we found a number of trends that suggest our model was operating

within expectations of population genetic theory. For example, during the habitat destruction

period, census population size declined to at or below the level as described in IUCN extinction

risk designations due to the combination of reduced carrying capacity, Allee effects, and age-

related removals (i.e., when $K = 1000$, $N_C \approx 750$; Figure 2A). Further, when there was no

immigration in our models, increasing the severity of population decline corresponded to

proportional decreases in genetic diversity and increases in genetic divergence between historic

and contemporary populations (Figure 2). For example, populations with the largest declines in

population size (90% decline, critically endangered) resulted in a 52.0% reduction in

heterozygosity across neutral SNPs whereas populations with the smallest tested decline (30%

decline, vulnerable) had a 14.8% decrease in heterozygosity between years 100 to 350 (Figure

2B). Despite these losses in genetic diversity and the resulting reduced fitness for some

individuals, populations returned to pre-bottleneck sizes following habitat restoration (Figure

2A). However, after recovery, populations had diverged substantially from the historical

recipient populations and at rates correlated with population crash intensity when the population

was small (Figure 1C). These populations also diverged from the historical migrant source

population at rates correlating to population crash intensity when the recipient population size

was small (Figure 2D). Because there was no migration in this set of analyses, these outputs

depict the rate of genetic drift between these two populations. Combined, these outputs illustrate

the development of unique evolutionary trajectories that can be altered by population size

reductions that ultimately influence individual fitness.

*Migrant genetic variance and population stability*

We were interested in how migrant-specific genetic characteristics influenced evolution and

stability in the migrant-recipient populations. When comparing genetic diversity in populations

with a single migrant per generation, we found that starting minor allele frequency (and therefore

lower heterozygosity compared to higher initial minor allele frequencies) in the migrant source

and recipient populations were not associated with any noticeable effect on average migration

ancestry represented in the recipient populations over time (Figure 3A). However, heterozygosity

changes in the neutral SNPs differed when starting minor allele frequencies differed; when the

source and recipient populations had high heterozygosity initially and no migration,

heterozygosity declined by 32.8% over the full 350-year period, although one migrant per

generation was effective in slowing this process to a 11.3% loss over the same time period

(Figure 3B). Similarly, when the starting minor allele frequencies in both migrant source and

recipient populations were small, heterozygosity decreased by 33.6% without migration and was

slowed to a 1% loss with migration across the 350 years. When the starting minor allele

frequencies in the recipient populations were low but high in the migrant source populations, one

migrant per generation increased heterozygosity, particularly through the population decline

(years 50-150 resulted in a 70.7% increase due to the influx of new alleles), and ended year 350

with heterozygosities nearest to the populations that started with high minor allele frequencies in

both populations with one migrant per generation ($H_{high > low} = 0.42$ with a 134.7% increase years

1-350, $H_{high > high} = 0.44$ with a 11.3% decrease years 1-350; Figure 3B). Finally, when the

migrant source populations had a low starting allele frequency that migrated into the recipient

populations with high initial heterozygosity, the resulting heterozygosity was less than if there

were no migrants; migrants with low fitness reduced the recipient population's heterozygosity by

46.9% (Figure 3B).

We measured divergence ($F_{ST}$) in two ways: the contemporary recipient populations at each year

compared to the historical recipient populations and each year between the contemporary

recipient and historical source populations. We found that the contemporary recipient

populations diverged from the historical recipient populations most slowly when both recipient

and source populations had high starting minor allele frequencies and when there was no

migration (Figure 3C). When there was a slow influx of migrants and the recipient populations

had limited genetic diversity compared to the source populations, the genetic variation of the

migrants influenced recipient population divergence, such that the populations diverged

significantly faster than all other migrant-recipient population pairs before, during, and after the

population size reduction period (depicted as non-overlapping CIs; Figure 3C). This reflects the

influx of many new alleles from the migrants into the recipient populations that substantially

increased migrant offspring fitness in the contemporary recipient populations. However, all

populations with migrants resulted in lower divergence from the historical source populations

than in populations without migration; when migrants had lower starting allele frequencies than

the recipient populations, the rate of divergence from the historical source populations decreased

the fastest, but $F_{ST}$ was still higher than if both populations contained similar starting allele

frequencies or when migrants were more fit than the recipient populations (Figure 3D).

*Recipient population size and migration effects*

We compared the genetic diversity effects on populations resulting from a single migrant per

generation to no migration when the migrant recipient populations were reduced to one of three

extinction risk levels. Despite a brief period of decline in genetic diversity when the population

size dropped (years 100 – 150), we found that heterozygosity was maintained in all combinations

of migration and population crash sizes (years 150 - 350; Figure 4B). Comparing within each

extinction risk category, we found that connected populations tended to be more genetically

diverse than the unconnected populations. However, even when connected by one migrant per

generation, populations reduced to the critically endangered size had similar heterozygosity as

unconnected vulnerable and unconnected non-bottlenecked populations (evidenced by

overlapping CIs; Figure 4B). These trends were driven directly by the increasing influence of the

migrants; over time, the proportion of migrant ancestry in the recipient populations increased in

all populations regardless of the magnitude of the population crash (i.e., the relative extinction

risk categories), but populations with the largest declines ended with the largest proportion of

migrant ancestry (nearly 100% in the critically endangered populations; Figure 4A). Because the

single-migrant populations with lower extinction risk lost fewer alleles during the population

crash, these populations were less penetrable by new migrant-brought alleles, meaning that

migrant ancestry accumulated less in these populations.

These patterns of differences among extinction risk categories related to migrant population

ancestry was also reflected in population divergence ($F_{ST}$) trends, where unconnected

populations diverged less from the historical recipient populations than their connected recipient

population counterparts (Figure 4C). In addition, the severity of the population crash was

inversely proportional to the strength of divergence observed; at the end of the 350-year

observation period, mean $F_{ST}$ between the historical recipient populations and the contemporary

recipient populations at year 350 ranged from 0.23 in the vulnerable populations to 0.37 in the

critically endangered populations when connected by one migrant per generation and from 0.11

in the vulnerable populations to 0.27 in the critically endangered populations in the absence of

migration (Figure 4C). When comparing divergence of the recipient populations to the historical

migrant source populations, we found that connected and unconnected populations had differing

trends; unconnected populations drifted from the historical migrant source populations whereas

the connected recipient populations tended to become more similar, with the size of the

population crash relating to the magnitude of this divergence. At the end of the year 350, mean

$F_{ST}$ between the historical migrant source populations and the contemporary connected recipient

populations ranged from 0.05 to 0.10 and the unconnected populations ranged from 0.41 to 0.48

in the critically endangered and vulnerable populations, respectively (Figure 4D). Although the

range between the minimum and maximum $F_{ST}$ within the connected and unconnected

populations was similar, the connected populations moved much further from the initial starting

value ($F_{ST} = 0.36$) compared to the unconnected populations, indicating accelerated divergence

in the populations connected by migration compared to the populations without migration.

We evaluated changes in genetic diversity, accumulation of migrant ancestry, and population

divergence among the four migration scenarios (one migrant per generation, burst of 100

migrants at a single time point, pulse of 25 migrants each at four time points, no migrants) within

each extinction risk category and in the absence of a population crash (Figure 5). Trends among

migration scenarios within each extinction risk category were similar, with the largest within-

group differences in the critically endangered category (90% decline); heterozygosity remained

consistent over each time step except during the population crash (years 100 to 150; Figure 5D-

5F). Within each extinction risk category, burst and pulsed migration scenarios resulted in

similar heterozygosities, surpassing heterozygosity in the one migrant per generation scenario

only in critically endangered populations as the population size increased, with final

heterozygosity similar in all populations with migrants in all extinction risk groups by year 350

(i.e., overlapping confidence intervals). In contrast to populations that contained migrants, the

final mean heterozygosity was significantly lower in the no migration scenarios in all extinction

risk groups (Figure 5D-5F). These trends were driven by migrant contributions to these

populations; as the proportion of migrant SNPs represented in the recipient populations

increased, neutral alleles originating in the migrant source populations also increased in

frequency. Populations connected by a single migrant per generation had a larger proportion of

alleles with migrant ancestry in most generations compared among all other migration schemes

and within each population crash size category (Figure 5A-5C). As individuals migrated and the

contemporary recipient populations increased to its historic size, alleles with migrant ancestry

were maintained due to the increased fitness of these lineages relative to some of the existing

families in each population (i.e., increased heterozygosity due to increased fitness in migrant-

resident pair offspring compared to offspring with both resident parents). Furthermore, the burst

migration scheme resulted in increased migrant ancestry compared to the pulsed scheme; in the

endangered population crash size category, for example, the burst introduction of 100 individuals

resulted in 10% more migrant ancestry than pulsed migration (38% vs 28% migrant ancestry

years 200 – 350). Thus, even though heterozygosity was equivalent in burst and pulse

populations from years 200 – 350, the genetic variants had different sources suggesting the

potential for different evolutionary and ecological outcomes.

Overall, divergence increased between the contemporary recipient populations and the historical

recipient populations over time (Figure 5G-5I) and decreased between the contemporary

recipient populations and historical migrant source populations each year (except in the absence

of migration; Figure 5J-5L). As populations returned to their historic sizes, pulsed migrations

diverged from the initial population the least, followed by no migration, burst, and single-

individual migration diverged the most, with magnitudes for each related to the size of the

population crash (i.e., larger percent population reduction resulted in more divergence). These

divergence patterns contrast with divergence of the recipient populations to the historical migrant

source populations; for all population crash categories, the populations not connected by

migration drifted away from the historical migrant source populations. All three sets of migrant

connected populations ended more similar to the historical migrant source populations than they

started, with a single migrant per generation and burst migrations being the most similar to the

migrant source populations immediately after habitat recovery (year 150) and pulsed migration

being significantly less similar that same year. After assisted migrations concluded, the burst and

pulsed migration populations diverged away from the historical migrant source populations

along their own evolutionary trajectories. Comparing across extinction risk categories and within

migration schemes, the smaller the population minima, the more diverged these populations were

from the historical recipient populations (Figure 5G-I) and the more similar to the historical

migrant source populations (Figure 5J-5L).

Consistent with trends comparing unconnected populations to populations connected by a single

migrant per generation, burst and pulsed migrations resulted in population genetic and

divergence outcomes that scaled with the severity of the population crash. For example, in both

burst and pulsed migrations, critically endangered populations resulted in a higher proportion of

migrant SNPS and were more diverged from the historical recipient populations than other

extinction risk categories (Figure 5). Similarly, these populations were less diverged from the

historical migrant source populations than the other population categories under the burst and

pulse migration model. Combined, these trends represent the effects of migrant ancestry that

build up at rates proportional to the recipient population size as long as migration continues.

Finally, we considered how the timing of assisted migration events affected genetic diversity of

the recipient population. Specifically, we compared population outcomes when migration

occurred during population minima (and therefore during habitat restoration) against when

migration occurred during recovery phases (after habitat restoration; Figure 6). Burst migrations

resulted in similar outcomes between these two timing scenarios, although heterozygosity was

slightly higher and divergence from the historical recipient populations slightly lower when

migration was initiated during the recovery phase (Figure 6A and 6E). In contrast to this, pulse

migration that occurred prior to the population recovery period resulted in almost double the

frequency of migrant alleles compared to if migration had occurred after population recovery

(67% vs. 38%; Figure 6B). As a result, pulsed migration that happens during population recovery

resulted in populations that more closely resembled the historical recipient populations and were

more diverged from the historical migrant source populations in year 350 compared to when

migration occurred before the population started growing (Figure 6F and 6H).

**DISCUSSION**

Although one migrant per population per generation is regarded often as the ideal population

connectivity scenario because higher genetic diversity is generally predictive of less inbreeding

and more stable populations (DeWoody et al. 2021; Kardos et al. 2021), connectivity can have

unintended effects if local alleles that are swamped by migrant variants ultimately reduce fitness

(Lenormand 2002). Here, we consider the contrasting effects of inbreeding depression and

connectivity in the context of conservation by modeling the effects of migration on recovery of

at-risk populations. Specifically, we consider populations from critically endangered,

endangered, and vulnerable species, vary migration rates and frequencies as well as migrant

source population characteristics, and consider the long-term effects that various management

decisions can have on these conservation targets. Overall, we find that the timing and rate of

migration is a critical predictor of the genetic make-up of recovered populations and that some

combinations of these parameters can send newly connected populations on entirely new

evolutionary trajectories. Through this work, we add to the body of evidence that identifies

habitat suitability, management actions, and population connectivity as the most limiting factors

for a species' long-term viability (Bouzat et al. 2009; Baling et al. 2016; Bubac et al. 2019; Hilty

et al. 2020; Goicolea et al. 2022; Gonzalez et al. 2023; Li et al. 2023; Smith et al. 2023) and

provide a model that can evaluate the risks and benefits of common management strategies.

Similar to other theoretical and empirical research relating the severity of population crashes to

population viability and the ability of migration and selection to effectively swamp local alleles

(Fagan and Holmes 2005; Lenormand 2002), here we show mechanistically how extinction risk

(i.e., population size) influences the relationship between these forces; our model shows that

reduced heterozygosity, high divergence from local alleles, and high relatedness to migrant

source populations occurs in connected populations with high intensity crashes. Ultimately,

predicting the populations' response to a single migrant per generation or assisted burst or pulsed

migrations depends on the system and goal of management interventions. For example, gene

flow can constrain adaptation to local conditions by passing on migrant derived traits, potentially

lowering fitness via outbreeding depression when fecundity is low or turnover is high (Frankham

et al. 2011; Weeks et al. 2011). In our model, accumulation of migrant ancestry alleles occurred

at a faster rate with low frequency but constant immigration as compared to burst or pulsed

assisted migration strategies, sometimes resulting in lower genetic diversity in the long term

(Figure 5). However, migrations also act as direct support in preventing extirpation that, when it

occurs, removes many of the locally adapted allele complexes. Thus, balancing the risks of

increased connectivity with the risks of complete population loss is tied to population-specific

pressures and management goals.

We investigated the effects of the average minor allele frequency in the migrant source and

recipient populations as a potential avenue for lessening the migration risks to populations on the

edge of extirpation (Figure 3). This is based on the idea that migration and selection can be

balanced across populations and gene variants will be maintained in the recipient populations

even with differing population-wide fitness (Lenormand 2002). In lab-based mesocosms,

Trinidadian guppy populations with lower starting genetic variation and smaller effective

population sizes benefitted from gene flow more than populations with higher diversity and

larger sizes, suggesting that evolutionary history can be predictive of individual and population

level fitness (Miller et al. 2019). Although the accumulation of migrant-derived alleles was

similar in the high and low minor allele frequency comparisons, the divergence from the

historical recipient populations was greatest when one or both migrant-connected populations

had a low starting allele frequency and divergence from the historical migrant source populations

was smallest when migrants have high starting allele frequencies. Because populations with high

conservation need are often small and have reduced genetic diversity, this supports the idea that

identifying populations with similar allelic variants, especially putatively adaptive variants, may

provide an avenue for demographic support with less risk to the existing population (Weeks et al.

2011; La Haye et al. 2017; Bertola et al. 2022). Here, we underscore the importance of this

consideration as our model suggests that the neutral portions of recipient populations are unlikely

to return to their historical genetic composition once migration ends (Figure 3C and 3D).

Maintaining behavioral, morphological, and metabolic adaptations expressed via locally adapted

alleles is often an important consideration in introducing gene flow among populations.

Although one migrant per generation can result in increased heterozygosity over time, the

tradeoff is that approximately 50-100% migrant derived alleles will drive this increase in

diversity, resulting in a population that more closely resembles the migrant source population –

and therefore retains migrant-favored traits – rather than traits that have accumulated in the

historical recipient population (Figure 4). Increased migration and selection in favor of migrant

traits further drives these demographic responses, which occurs at faster time scales than genetic

drift (see Supplementary Figure 1), despite the effect of this relationship being proportional to

population crash sizes. In the case of rare species, maintaining these derived traits is imperative

to the sustainability of the species.

The inverse relationship between genetic diversity and the presence of admixed alleles is

especially evident in populations that exhibit introgression and hybridization (Allendorf et al.

2001); backcrossing of previously isolated populations homogenizes historically unique alleles.

While homogenization can contribute to the loss of rare traits, introgression can support

populations in rapidly changing environments that require more than the current standing genetic

diversity (Benjamin-Fink and Reilly 2017; Ranke et al. 2020). Unfortunately, until admixture

occurs in wild populations, it is hard to predict if hybridization of distinct populations will result

in more adaptive phenotypes (potentially due to increased heterozygosity) or disruption of co-

adapted gene complexes.

Assisted migration, such as the burst and pulse migrations simulated here, develop alternative

evolutionary trajectories, evidenced by a similar increase in heterozygosity but different rates of

decreased divergence from the migrant source populations. Burst and pulse migrations result in

overall similar heterozygosities across all extinction risk categories, but a slower retention of

migrant derived alleles in pulsed migrations, depicting a lessened Ryman-Laikre effect in this

management plan (Figure 5). Therefore, pulse migrations seem to be more effective than burst

migrations for increasing diversity and population size while maintaining local alleles. For

example, in an endangered *Cricetus cricetus* population, genetic diversity initially decreased

after the first translocation when reintroduced individuals were genetically different from the

receiving population, but additional supplementation increased genetic diversity regardless of the

initial diversity of the source and recipient populations (La Haye et al. 2017). This disparate

adaptive potential may be due to alternative responses to selection; gene-frequency estimates

(e.g., heterozygosity) responds to selection in the short-term while allelic-diversity estimates

(e.g., migrant ancestry) reflect long-term adaptation and total selection responses because

segregating sites within a subdivided population are more strongly constrained by natural

selection than neutral alleles (Cabellero and García-Dorado 2013). In addition to the genetic

benefits of pulse migrations, when management utilizes multiple translocations, recruitment of

migrants into the population may increase via increased learning of newly translocated

individuals from those surviving from a previous reintroduction, higher availability of density-

dependent resources, relaxed Allee effects, a buffer against high mortality from stochastic natural

disturbance, and feasibility of more frequent but smaller management actions (IUCN/SCC 2013;

La Haye et al. 2017; Martin et al. 2017; Lewis et al. 2022).

Just as the frequency of introductions alters the demographic and genetic effects in the receiving

population, we examined how the timing of translocations influences the accumulation of traits

with migrant ancestry. When migrations occur before the habitat is restored and while the

population remains at low population sizes, accumulation of migrant traits and increased

divergence from the historical population occurs long-term, altering the demographic and

evolutionary trajectories compared to if the habitat quality increased and the population was

permitted to grow post restoration (Figure 6). Waiting to begin translocations until habitat quality

increases is an effective management strategy (see Figure 2 in Martin et al. 2017), as long as the

population is not extirpated during the waiting period. In our system, critically endangered

populations without gene flow were eradicated within 18-65 generations of population decline

commencement (Supplementary Figure 2), which could be before assisted migrations are

implemented if the habitat is unsuitable for supplementation. Though gene flow can temporarily

increase population viability, without addressing the cause of the decline — often habitat

destruction — the long-term sustainability is still at risk (Grant et al. 2019). While translocations

were and still often are regarded as the last-ditch effort due to long-lasting effects on the

population and environment, we recommend active and proactive restoration efforts over natural

regeneration (Watson and Watson 2015), especially for populations with high risk of extirpation

and when proper techniques and genetic considerations are implemented (Chipman et al. 2008;

Weeks et al. 2011; Berger-Tal et al. 2019; Morris et al. 2021).

Of course, instead of active population management and monitoring, many of these difficulties in

assisted migrations could be alleviated with increased habitat quality and decreased

fragmentation; corridors that allow for gene flow among populations are an effective strategy

that can boost viability (Sharma et al. 2013; Christie and Knowles 2015; Hevroy et al. 2017; this

study). Although unassisted gene flow via habitat restoration and connectivity could be more

time and cost effective long-term than repeated assisted gene flow, the desired genetic diversity,

relatedness of migrant source and recipient populations, and the desired rate of population

growth must be considered. Indeed, our simulations suggest that populations with connectivity

via one migrant per generation are slower to grow in size and retain migrant alleles faster than

populations with assisted migrations but can still be more useful for conservation planning than

continued population fragmentation. Additional investigations into the adaptability of

populations, perhaps through epigenomic modifications or selection based on changing

environmental conditions could be additionally useful in population management. Further,

examination of other genetic problems that are reflected in small populations like mutational

load, inbreeding depression, and fixation of maladaptive alleles are promising avenues for future

viability analyses. Ultimately, protecting and maintaining species at the brink of extinction will

benefit biodiversity and the community by building engagement and collaboration (Kremen and

Merenlender 2018; Zellmer and Goto 2022), in addition to the genetic and demographic benefits

of connected populations to species sustainability.

Data Archiving Statement: All scripts and data for this study are available at

https://github.com/ginalamka/ComplexModel_ABM/.

**Literature Cited**

Allendorf, F. W., R. F. Leary, P. Spruell, and J. K. Wenburg. 2001. The problems with hybrids: setting conservation guidelines. Trends in Ecology and Evolution 16:613-622.

Anderson, D. P., M. G. Turner, J. D. Forester, J. Zhu, M. S. Boyce, H. Beyer, and L. Stowell. 2005. Scale-dependent summer resource selection by reintroduced elk in Wisconsin, USA. Journal of Wildlife Management 69:298-310.

Backus, G. A. and M. L. Baskett. 2021. Identifying robust strategies for assisted migration in a competitive stochastic community. Conservation Biology 35:1809-1820.

Baling, M., D. Stuart-Fox, D. H. Brunton, and J. Dale. 2016. Habitat suitability for conservation translocation: The importance of considering camouflage in cryptic species. Biological Conservation 203:298-305.

Barbosa, A. E. A. and J. L. Tella. 2019. How much does it cost to save a species from extinction? Costs and rewards of conserving the Lear's macaw. Royal Society Open Science 6:190190.

Benjamin-Fink, N. and B. K. Reilly. 2017. Conservation implications of wildlife translocations; The state's ability to act as conservation units for wildebeest populations in South Africa. Global Ecology and Conservation 12:46-58.

Berger-Tal, O., D. T. Blumstein, and R. R. Swaisgood. 2019. Conservation translocations: A review of common difficulties and promising directions. Animal Conservation 23:121-131.

Bertola, L. D., S. M. Miller, V. L. Williams, V. N. Naude, P. Coals, S. G. Dures, P. Henschel, M. Chege, E. A. Sogbohossou, A. Ndiaye, M. Kiki, A. Gaylard, D. K. Ikanda, M. S. Becker, and P. Lindsey. 2022. Genetic guidelines for translocations: Maintaining intraspecific diversity in the lion (*Panthera leo*). Evolutionary Applications 15:22-39.

Bouzat, J. L., J. A. Johnson, J. E. Toepfer, S. A. Simpson, T. L. Esker, and R. L. Westemeier. 2009. Beyond the beneficial effects of translocations as an effective tool for the genetic restoration of isolated populations. Conservation Genetics 10:191-201.

Braumann, C. A. 2010. Environmental versus demographic stochasticity in population growth. In: González Velasco, M., Puerto, I., Martínez, R., Molina, M., Mota, M., Ramos, A. (eds) Workshop on Branching Processes and Their Applications. Lecture Notes in Statistics(), vol 197. Springer, Berlin, Heidelberg.

Bubac, C. M., A. C. Johnson, J. A. Fox, and C. I. Cullingham. 2019. Conservation translocations and post-release monitoring: Identifying trends in failures, biases, and challenges from around the world. Biological Conservation 238:108239.

Buxton, R. T., S. Avery-Gomm, H-Y. Lin, P. A. Smith, S. J. Cooke, and J. R. Bennett. 2020. Half of resources in threatened species conservation plans are allocated to research and monitoring. Nature Communications 11:4668.

Cabellero, A. and A. García-Dorado. 2013. Allelic diversity and its implications for the rate of
adaptation. Genetics 195:1373-1384.

Caughley, G. 1994. Directions in conservation biology. Journal of Animal Ecology 63:215-244.

Chen, Z., L. Grossfurthner, J. L. Loxterman, J. Masingale, B. A. Richardson, T. Seaborn, B.
Smith, L. P. Waits, and S. R. Narum. 2021. Applying genomics in assisted migration under
climate change: Framework, empirical applications, and case studies. Evolutionary Applications
15:3-21.

Chipman, R., D. Slate, C. Rupprecht, and M. Mendoza. 2008. Downside risk of wildlife
translocation. Developments in Biologicals 131:223-232.

Christie, M. R. and L. L. Knowles. 2015. Habitat corridors facilitate genetic resistance
irrespective of species dispersal abilities or population sizes. Evolutionary Applications 8:454-
463.

Christie, M. R., M. L. Marine, R. A. French, R. S. Waples, and M. S. Blouin. 2012. Effective
size of a wild salmonid population is greatly reduced by hatchery supplementation. Heredity
109:254-260.

Coltman, D. W. and J. Slate. 2003. Microsatellite measures of inbreeding: a meta-analysis.
Evolution 57:971-983.

Del Castillo, R. F., S. Trujillo-Argueta, N. Sanchez-Vargas, and A. C. Newton. 2011. Genetic
factors associated with population size may increase extinction risks and decrease colonization
potential in a keystone tropical pine. Evolutionary Applications 4:574-588.

DeWoody, J. A., A. M. Harder, S. Mathur, and J. R. Willoughby. 2021. The long-standing
significance of genetic diversity in conservation. Molecular Ecology 30:4147-4154.

DeWoody, J. A., J. Y. Jeon, J. W. Bickham, E. J. Heenkenda, S. Janjua, G. F. Lamka, A. J.
Mularo, A. Black, A. Bruniche-Olsen, and J. R. Willoughby. 2022. The Threatened Species
Imperative: Conservation assessments would benefit from population genomic insights. PNAS
119:e2210685119.

Drake, J. M. 2008. Population viability analysis. Encyclopedia of Ecology 2901-2907.

Drake, J. M. and A. M. Kramer. 2011. Allee effects. Nature Education Knowledge 3:2.

Drake, J. M., K. L. S. Drury, D. M. Lodge, A. Blukacz, N. D. Yan, and G. Dwyer. 2006.
Demographic stochasticity, environmental variability, and window of invasion risk for
*Bythotrephes longimanus* in North America. Biological Invasions 8:843-861.

Fagan, W. F. and E. E. Holmes. 2005. Quantifying the extinction vortex. Ecology Letters 9:51-
60.

Frankham, R., J. D. Ballou, M. D. B. Eldridge, R. C. Lacy, K. Ralls, M. R. Dudash, and C. B.
Fenster. 2011. Predicting the probability of outbreeding depression. Conservation Biology
25:465-475.

Furlan, E. M., B. Gruber, C. R. M. Attard, R. N. E. Wager, A. Kerezsy, L. K. Faulks, L. B.
Beheregaray, and P. J. Unmack. 2020. Assessing the benefits and risks of translocations in
depauperate species: A theoretical framework with an empirical validation. Journal of Applied
Ecology 57:831-841.

Goicolea, T., R. G. Mateo, M. J. Aroca-Fernandez, A. Gaston, J. I. Garcia-Viñas, and M. C.
Mateo-Sanchez. 2022. Considering plant functional connectivity in landscape conservation and
restoration management. Biodiversity and Conservation 31:1591-1608.

Gonzalez, A. M., N. Espejo, D. Armenteras, K. A. Hobson, K. J. Kardynal, G. W. Mitchell, N.
Mahony, C. A. Bishop, P. J. Negret, and S. Wilson. 2023. Habitat protection and restoration:
Win-win opportunities for migratory birds in the Northern Andes. Perspectives in Ecology and
Conservation 21:33-40.

Goudet, J. 2005. HIERFSTAT, a package for R to compute and test hierarchical $F$-statistics.
Molecular Ecology Notes 5:184-186.

Grant, L., C. Johnson, and C. Thiessen. 2019. Evaluating the efficacy of translocation:
maintaining habitat key to long-term success for an imperiled population of an at-risk species.
Biodiversity and Conservation 28:2727-2743.

Haddad, N. M., L. A. Brudvig, J. Clobert, K. F. Davies, A. Gonzales, R. D. Holt, T. E. Lovejoy,
J. O. Sexton, M. P. Austin, C. D. Collins, W. M. Cook, E. I. Damschen, R. M. Ewers, B. L.
Foster, C. N. Jenkins, A. J. King, W. F. Laurance, D. J. Levey, C. R. Margules, B. A. Melbourne,
A. O. Nicholls, J. L. Orrock, D.-X. Song, and J. R. Townshend. 2015. Habitat fragmentation and
its lasting impact on Earth's ecosystems. Science Advances 1:e1500052.

Hagen, I. J., A. J. Jensen, G. H. Bolstad, O. H. Diserud, K. Hindar, H. Lo, and S. Karlsson. 2019.
Supplementary stocking selects for domesticated genotypes. Nature Communications 10:199.

Hagen, I. J., O. Ugedal, A. J. Jensen, H. Lo, E. Holthe, B. Bjøru, B. Florø-Larsen, H. Sægrov, H.
Skoglund, and S. Karlsson. 2020. Evaluation of genetic effects on wild salmon populations from
stock enhancement. ICES Journal of Marine Science 78:900-909.

Hevroy, T. H., M. L. Moody, and S. L. Krauss. 2018. Population genetic analysis reveals barriers
and corridors for gene flow within and among riparian populations of a rare plant. AoB Plants
10:plx065.

Hilbers, J. P., M. A. J. Huijbregts, and A. M. Schipper. 2019. Predicting reintroduction costs for
wildlife population under anthropogenic stress. Journal of Applied Ecology 57:192-201.

Hilty, J., G. L. Worboys, A. Keeley, S. Woodley, B. J. Lausche, H. Locke, M. Carr, I. Pulsford,
J. Pittock, J. W. White, D. M. Theobald, J. Levine, M. Reuling, J. E. M. Watson, R. Ament and

G. M. Tabor. 2020. Guidelines for conserving connectivity through ecological networks and corridors. Best Practice Protected Area Guidelines Series No. 30. Gland, Switzerland: IUCN.

Hogg, C. J., K. Ottewell, P. Latch, M. Rossetto, J. Biggs, A. Gilbert, S. Richmond, and K. Belov. 2022. Threatened Species Initiative: Empowering conservation action using genomic resources. PNAS 119:e2115643118.

IUCN/SSC. 2013. Guidelines for reintroductions and other conservation translocations. Version 1.0. Gland, Switzerland: IUCN Species Survival Commission, viiii.

Julien, M., B. Colas, S. Muller, and B. Schatz. 2022. Dataset of costs of the mitigation hierarchy and plant translocations in France. Data in Brief 40:107722.

Kardos, M., E. E. Armstrong, S. W. Fitzpatrick, S. Hauser, P. W. Hendrick, J. M. Miller, D. A. Tallmon, and W. C. Funk. 2021. The crucial role of genome-wide genetic variation in conservation. PNAS 118:e2104642118.

Kim, Y-K., Y-L. Hong, M-S. Min, K. S. Kim, Y-J. Kim, I. Voloshina, A. Myslenkov, G. J. D. Smith, N. D. Cuong, H. H. Tho, S-H. Han, D-H. Yang, C-B. Kim, and H. Lee. 2011. Genetic status of Asiatic black bear (*Ursus thibetanus*) reintroduced into South Korea based on mitochondrial DNA and microsatellite loci analysis. Journal of Heredity 102:165-174.

Kremen, C. and A. M. Merenlender. 2018. Landscapes that work for biodiversity and people. Science 362:eaau6020.

La Haye, M. J. J., T. E. Reiners, R. Raedts, V. Verbist, and H. P. Koelewijn. 2017. Genetic monitoring to evaluate reintroduction attempts of a highly endangered rodent. Conservation Genetics 18:877-892.

Lenormand, T. 2002. Gene flow and the limits to natural selection. Trends in Ecology & Evolution 17:183-189.

Lewis, J. C., K. J. Jenkins, P. J. Happe, D. J. Manson, and P. C. Griffin. 2022. Post-release survival of translocated fishers: implications for translocation success. Journal of Wildlife Management 86:e22192.

Li, W., P. Liu, N. Yang, S. Chen, X. Guo, B. Wang, and L. Zhang. 2023. Improving landscape connectivity through habitat restoration: application for Asian elephant conservation in Xishuangbanna Prefecture, China. Integrative Zoology 0:1-17. Doi:10.1111/1749-4877.12713

Lowe, W. H. and F. W. Allendorf. 2010. What can genetics tell us about population connectivity? Molecular Ecology 19:3038-3051.

Martin, J. A., R. D. Applegate, T. V. Dailey, M. Downey, B. Emmerich, F. Hernández, M. M. McConnell, K. S. Reyna, D. Rollins, R. E. Ruzicka, and T. M. Terhune II. 2017. Translocation and population restoration technique for northern bobwhites: A review and synthesis. National Quail Symposuim Proceedings 8:11.

Martinez-Abrain, A., H. M. Regan, C. Viedma, E. Villuendas, M. A. Bartolome, J. A. Gomez,
and D. Oro. 2011. Cost-effectiveness of translocation options for a threatened waterbird.
Conservation Biology 25:726-735.

McLachlan, J. S., J. J. Hellman, and M. W. Schwartz. 2007. A framework for debate of assisted
migration in an era of climate change. Conservation Biology 21:297-302.

Messer, P. W. 2016. Neutral models of genetic drift and mutation. Encyclopedia of Evolutionary
Biology 119-123.

Miller, M. L., J. A. Kronenberger, and S. W. Fitzpatrick. 2019. Recent evolutionary history
predicts population but not ecosystem-level patterns. Ecology and Evolution 9:14442-14452.

Mills, L. S. and F. W. Allendorf. 1996. The one-migrant-per-generation rule in conservation and
management. Conservation Biology 10:1509-1518.

Morris, S. D., B. W. Brook, K. E. Moseby, C. N. Johnson. 2021. Factors affecting success of
conservation translocations of terrestrial vertebrates: A global systematic review. Global Ecology
and Conservation 28:e01630.

Nachman, M. W. and S. L. Crowell. 2000. Estimate of the mutation rate per nucleotide in
humans. Genetics 156:297-304.

Nash, D. J., N. Humphries, and R. A. Griffiths. 2020. Effectiveness of translocation in mitigating
reptile-development conflict in the UK. Conservation Evidence 17:7-11.

Payton, M. E., M. H. Greenstone, and N. Schenker. 2003. Overlapping confidence intervals or
standard error intervals: What do they mean in terms of statistical significance? Journal of Insect
Science 1:34.

Ranke, P. S., S. Skjelseth, I. J. Hagen, A. M. Billing, A. A. B. Pedersen, H. Parn, T. H. Ringsby,
B.-E. Saether, and H. Jensen. 2020. Multi-generational genetic consequences of reinforcement in
a bird metapopulation. Conservation Genetics 21:603-612.

Ryman, N. and L. Laikre. 1991. Effects of supportive breeding on the genetically effective
population size. Conservation Biology 5:325-329.

Sansilvestri, R., N. Frascaria-Lacoste, and J. F. Fernandez-Manjarres. 2015. Reconstructing a
deconstructed concept: Policy tools for implementing assisted migration for species and
ecosystem management. Environmental Science and Policy 51:192-201.

Sasmal, I., J. A. Jenks, L. P. Waits, M. G. Gonda, G. M. Schroeder, and S. Datta. 2013. Genetic
diversity in a reintroduced swift fox population. Conservation Genetics 14:93-102.

Schwartz, M. W. and T. G. Martin. 2013. Translocation of imperiled species under changing
climates. Annals of the New York Academy of Sciences 1286:15-28.

Sharma, S., T. Dutta, J. E. Maldonado, T. C. Wood, H. S. Panwar, and J. Seidensticker. 2013.
Forest corridors maintain historical gene flow in a tiger metapopulation in the highlands of
central India. Proceedings of the Royal Society B 280:20131506.

Smith, J. B., D. A. Keiter, S. J. Sweeney, R. S. Miller, P. E. Schlichting, and J. C. Beasley. 2023.
Habitat quality influences trade-offs in animal movement along the exploration-exploitation
continuum. Scientific Reports 13:4814.

Stepkovitch, B., R. T. Kingsford, and K. E. Moseby. 2022. A comprehensive review of
mammalian carnivore translocations. Mammal Review 52:554-572.

Thrimawithana, A. H., L. Ortiz-Catedral, A. Rodrigo, and M. E. Hauber. 2013. Reduced total
genetic diversity following translocations? A metapopulation approach. Conservation Genetics
14:1043-1055.

Waldron, A., A. O. Mooers, D. C. Miller, N. Nibbelink, D. Redding, T. S. Kuhn, J. T. Roberts,
and J. L. Gittleman. 2013. Targeting global conservation funding to limit immediate biodiversity
declines. PNAS 110:12144-12148.

Watson, D. M. and M. J. Watson. 2015. Wildlife restoration: Mainstreaming translocations to
keep common species common. Biological Conservation 191:830-838.

Weeks, A. R., C. M. Sgro, A. G. Young, R. Frankham, N. J. Mitchell, K. A. Miller, M. Byrne, D.
J. Coates, M. D. B. Eldridge, P. Sunnucks, M. F. Breed, E. A. James, and A. A. Hoffmann. 2011.
Assessing the benefits and risks of translocations in changing environments: a genetic
perspective. Evolutionary Applications 4:709-804.

Weise, F. J., K. J. Stratford, and R. J. van Vuuren. 2014. Financial costs of large carnivore
translocations – accounting for conservation. PLOS ONE 9:e105042.

White, T. B., S. O. Petrovan, A. P. Christie, P. A. Martin, and W. J. Sutherland. 2022. What is
the price of conservation? A review of the status quo and recommendations for improving cost
reporting. BioScience 72:461-471.

Weir, B. S. and J. Goudet. 2017. A unified characterization of population structure and
relatedness. Genetics 206:2085-2103.

Williams, M. I. and R. K. Dumroese. 2013. Preparing for climate change: Forestry and assisted
migration. Journal of Forestry 111:287-297.

Zellmer, A. J. and B. S. Goto. 2022. Urban wildlife corridors: Building bridges for wildlife and
people. Frontiers in Sustainable Cities 4: 954089.

**Figure 1.** Schematic representation of migrant source and recipient population models. (A) The recipient population was allowed to reach equilibrium and stabilize for 100 years with a carrying capacity of 1000 individuals and was then subjected to a 50-year population decline to simulate environmental change; this population size decline happened over 10 years, and the population was held at the population minima for 40 years at three levels representing IUCN (International Union for Conservation of Nature) and state agency classifications: vulnerable for species that have a > 10% decline in 10 years, $K_B = 700$; endangered with a 50-70% decline in 10 years, $K_B = 300$; and critically endangered with a 80-90% decline in 10 years, $K_B = 100$. After this reduced population period, we simulated habitat restoration by allowing the population to grow again, up to the historical carrying capacity. Additional simulations for populations that did not face population decline were used as a control. (B) Migrants entered the recipient population at a set frequency: 1 migrant per generation, 100 individuals at a single time period (e.g., 100 individuals in year 151) and 25 individuals at four time periods (e.g., 25 individuals in years 151, 165, 181, 195). As a basis for comparison, we also included scenarios where migration was completely absent. (C) We tracked the demographic and evolutionary response of the recipient populations across 350 years.

**Figure 2.** Population genetic and demographic responses in populations compared among extinction risk categories in the absence of migration. The census population size (A) is illustrated for 100 replicates of the simulation runs, where each line represents a single iteration. Here, the census size is depicted as less than carrying capacity to match the point in the model when population response parameters were quantified (i.e., after Allee effects, reproduction, and subsequent mortality). Population response is summarized across these replicates as observed heterozygosity (B), divergence from the historical recipient populations over time (C), and

divergence between the recipient and historical migrant source populations at each year (D). In

the absence of migration, a drop in census size can push populations on a new evolutionary

trajectory with different allele frequencies as a result of the loss of genetic variants that occurred

due to drift. Lines in B-D represent mean values across the 100 replicate runs and polygons

represent confidence intervals scaled to compare evolutionary outcomes between each parameter

set, assuming alpha = 0.05 (i.e., 95% confidence intervals).

**Figure 3.** Genetic and demographic responses to migration in populations with different starting

minor allele frequencies and a population crash to 70% of the historical population size. Solid

lines indicate a single migrant per generation and dotted lines depict scenarios without migration.

Starting minor allele frequencies were either low (0.05 – 0.15) in both the migrant source and

recipient populations (cyan), high (0.4 – 0.5) in both populations (purple), high in the migrant

source populations and low in the receiving populations (pink), or low in the migrant source

populations and high in the receiving populations (blue). Lines represent mean values across 100

replicate runs and polygons represent confidence intervals scaled to compare evolutionary

outcomes between each parameter set, assuming alpha = 0.05 (i.e., 95% confidence intervals).

The proportion of migrant ancestry (A), observed heterozygosity (B), divergence from the

historical recipient populations over time (C), and divergence each year from the historical

migrant source populations (D) are depicted in each panel. These outcomes suggest that the

minor allele frequencies influence evolutionary trajectory of populations connected by migration

when migrations occur during a population crash, such that these recipient populations with

migrants from populations with higher minor allele frequencies are less diverged from the

migrant source populations compared to recipient populations with migrants with lower minor

allele frequencies.

**Figure 4.** Population genetic and demographic outcomes of incorporating migration into

population crash recovery. Populations were shrunk (years 100-150) to sizes reflecting common

extinction risk classifications (critically endangered, 90% reduction; endangered, 70% reduction;

vulnerable, 30% reduction; no population reduction), and movement of a single individual per

year (solid line) was used to support and bolster these populations through remediation (year

151) and population recovery. These trends were compared to the same demographic patterns but

without migration (dotted lines). The proportion of migrant ancestry present in the recipient

populations (A), observed heterozygosity (B), divergence of the recipient populations from the

historical populations over time (C), and divergence of the recipient populations from the

migrant source populations each year (D) illustrate the new evolutionary trends resulting from

these migration decisions. Lines represent mean values across 100 replicates and shaded areas

represent the confidence intervals needed to compare among parameter sets assuming alpha =

0.05 (i.e., 95% confidence intervals).

**Figure 5.** Population genetic and demographic outcomes in populations compared among

extinction risk categories while incorporating migration into population crash recovery.

Populations were shrunk (years 100-150) to sizes reflecting common extinction risk

classifications (critically endangered, 90% reduction; endangered, 70% reduction; vulnerable

30% reduction; no population reduction), and movement of a single individual per year (solid

line), burst migration of 100 individuals once (year 151; dashed line), and four pulse migrations

of 25 individuals (years 151, 165, 181, 195; dashed and dotted line) was used to support and

bolster these populations through remediation (year 151) and population recovery. These trends

were compared to the same demographic patterns but without migration (dotted line) and in the

absence of a population crash (grey lines). The proportion of migrant ancestry present in the

recipient populations (A-C), observed heterozygosity (D-F), divergence of the recipient

populations from the historical population over time (G-I), and divergence of the recipient

populations from the historical migrant source populations each year (J-L) illustrate the new

evolutionary trends resulting from these management decisions every 50 years. Lines represent

mean values across 100 replicates and error bars represent the confidence intervals needed to

compare among parameter sets assuming alpha = 0.05 (i.e., 95% confidence intervals).

**Figure 6.** Population genetic and demographic responses with a population crash to 70% of the

starting population with different timings of management intervention. Assisted migrations were

either burst translocations (100 individuals once) or pulsed (25 individuals 4 times). Light grey

lines depict assisted migrations implemented concurrently with habitat restoration (year 125 for

burst; years 125, 140, 155, and 170 for pulse) and black lines indicate migrations that occurred

after habitat restoration was completed (year 151 for burst; years 151, 165, 181, and 195 for

pulse). Lines represent mean values across 100 replicate runs and polygons represent confidence

intervals scaled to compare evolutionary outcomes between each parameter set, assuming alpha

= 0.05 (i.e., 95% confidence intervals). The proportion of migrant ancestry (A), observed

heterozygosity (B), divergence from the historical recipient populations over time (C), and

divergence each year from the historical migrant source populations (D) are depicted in each

panel. These outcomes suggest that implementing a pulse migration after restoration and with

population growth will supplement populations with similar levels of increased genetic diversity

and with less influence of alleles with migrant ancestry.

**Table 1.** Description of parameters and values tested. These parameters describe the starting evolutionary point for the recipient

populations, the extent to which population size was reduced to reflect IUCN (International Union for Conservation of Nature)

extinction risk designations, and the rate and timing of migration events.

| Parameter | Description | Values tested |
|---|---|---|
| Minor allele frequency | Initial minor allele frequency (p) for migrant source and recipient populations | Low minor allele frequency (0.05-0.15); high minor allele frequency (0.40-0.50) |
| Population crash size | Modified carrying capacity forced on recipient populations from year 100 to 150 to simulate habitat destruction (or degradation); note that census size could be below carrying capacity due to Allee effects and mortality | Carrying capacity (K) reflecting IUCN designations: Vulnerable K = 700, Endangered K = 300, Critically Endangered K = 100 |
| Migration rate | Rate at which individuals moved from migrant source populations into recipient populations | No migration; 1 migrant per generation; single burst of 100 individuals; four pulses of 25 individuals |
| Migrant timing | For burst and pulsed migration, at what point during the population crash and recovery were assisted migrations applied | Migrants were moved to the recipient population during habitat restoration; Migrants were moved to the recipient population only after habitat restoration was completed |

## A. Focal recipient population

Populations of 1000 individuals (K) were monitored during a habitat destruction-induced crash and subsequent environmental remediation and recovery.

## B. Migrant source population

Receiving populations were supported with different amounts of immigration during the crash and recovery periods.

Migrant source population

One migrant per generation
Burst of 100 individuals in a single year
Pulses of 25 individuals in four years

## C. Evolutionary outcomes

We quantified population evolutionary trajectories that resulted from the differing immigration patterns in the recovered receiving populations.

Habitat destruction and subsequent population crash. Small population sizes ($K_B$) were set following IUCN extinction risk category cutoffs for population decline.

Habitat remediation leading to population recovery to historic carrying capacity (K).