Working Memory Gating in Obesity is Moderated by Striatal Dopaminergic Gene Variants

Nadine Herzog[1,2], Hendrik Hartmann[1,3,4], Lieneke Katharina Janssen[1,5], Arsene Kanyamibwa[4], Maria Waltmann[1,6], Peter Kovacs[7], Lorenz Deserno[6,8], Sean James Fallon[9], Arno Villringer[1], Annette Horstmann[1,3,4]

1. Department of Neurology, Max Planck Institute for Human Cognitive & Brain Sciences, Leipzig, Germany
2. International Max Planck Research School NeuroCom, Leipzig, Germany
3. Collaborative Research Centre 1052, University of Leipzig, Leipzig, Germany
4. Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Helsinki, Finland
5. Institute of Psychology, Otto von Guericke University Magdeburg, Magdeburg, Germany
6. Department of Child and Adolescent Psychiatry, University of Würzburg, Würzburg, Germany
7. Medical Department III - Endocrinology, Nephrology, Rheumatology, University of Leipzig Medical Center, Leipzig, Germany
8. Department of Psychiatry and Psychotherapy, Technische Universität Dresden, Dresden, Germany
9. School of Psychology, University of Plymouth, Plymouth, UK

Key words

obesity, dopamine, working memory gating, updating, maintenance, Taq1A, COMT, DARPP, C957T, amino acid ratio

Abstract

Everyday life requires an adaptive balance between distraction-resistant maintenance of information and the flexibility to update this information when needed. These opposing mechanisms are proposed to be balanced through a working memory gating mechanism. Prior research indicates that obesity may elevate the risk of working memory deficits, yet the underlying mechanisms remain elusive. Dopaminergic alterations have emerged as a potential mediator. However, current models suggest these alterations should only shift the balance in working memory tasks, not produce overall deficits. The empirical support for this notion is currently lacking, however. To address this gap, we pooled data from three studies (N = 320) where participants performed a working memory gating task. Higher BMI was associated with overall poorer working memory, irrespective of whether there was a need to maintain or update information. However, when participants, in addition to BMI level, were categorized based on certain putative dopamine-signaling characteristics (Single Nucleotide Polymorphisms; specifically, Taq1A and DARPP-32), distinct working memory gating effects emerged. These SNPs, primarily associated with striatal dopamine transmission, appear to be linked with differences in updating, specifically, among high-BMI individuals. Moreover, blood amino acid ratio, which indicates central dopamine synthesis capacity, combined with BMI, shifted the balance between distractor-resistant maintenance and updating. These findings suggest that both dopamine-dependent and dopamine-independent cognitive effects exist in obesity. Understanding these effects is crucial if we aim to modify maladaptive cognitive profiles in individuals with obesity.

INTRODUCTION

In order to function efficiently in a dynamic environment, we must be able to resist distractions while simultaneously being open to update information in response to evolving goals and task requirements. This tension demands a delicate balance, which is thought to be governed by one of our core cognitive control systems - our working memory (WM). Computational and neurophysiological theories propose a metaphorical "gate" that regulates the access to WM (Badre, 2012; O'Reilly and Frank, 2006). When the gate is closed, WM representations are isolated from perceptual input and interference is prevented. When the gate is opened, rapid updating is allowed. Evidence strongly implicates the prefrontal cortex (PFC) in distractor-resistant maintenance, while updating is thought to be supported by the striatum (Miller and Cohen, 2001; D'Ardenne et al., 2012; Braver and Cohen, 2000). Importantly, the neurotransmitter dopamine plays a crucial role in balancing these complementary processes. Within the PFC, tonic dopamine levels mediate maintenance in an inverted-U-shaped manner: very high and very low levels promote gate opening, while medium levels promote gate closing (Cools and D'Esposito, 2011). Within the striatum, phasic increases in dopamine are needed to signal WM updating (Hazy, Frank, and O'Reilly, 2006; D'Ardenne et al., 2012). Importantly, the effectiveness of phasic rises in dopamine to override PFC tonic signals depends on initial baseline dopamine levels in the striatum (Cools and D'Esposito, 2011; Schouwenburg, Aarts, & Cools, 2010; Ranganath & Jacob, 2016). Decreases in tonic dopamine levels in the striatum seem to raise the threshold for updating signals, thus potentially hindering updating (O'Reilly and Frank, 2006). Supporting this, worse updating of WM contents can be observed in unmedicated patients with Parkinson's disease (Fallon et al. 2017), older individuals (Podell et al., 2012), or more generally, in individuals with lower dopamine synthesis capacity (Colzato et al., 2013). Notably, dominance in one WM process typically comes at the cost of the other (Dreisbach & Fröber, 2019; Dreisbach et al., 2005; Fallon and Cools, 2017, 2019a, 2019b). Consequently, an individual's capacity to ignore or update (ir)relevant information may vary according to their baseline dopamine levels (e.g. Cools & D'Esposito, 2011; Furman et al., 2021, Jongkees, 2020).

Interestingly, the intricate relationship between dopamine levels and working memory gating might be key in further understanding discrepancies in the literature regarding working memory functioning in obesity. While many studies show reduced (general) working memory in obese individuals (e.g., Yang et al., 2020, 2019, 2018; Gonzales et al., 2010; Coppin et al., 2014), there are others who do not find such associations (e.g., Calvo et al., 2014; Schiff et al., 2016; Alarcon et al., 2016). Based on the above considerations these inconsistencies may be due to prior studies not clearly differentiating between distractor-resistant maintenance and updating in the context of working memory. This distinction may be crucial, however, as indirect evidence hints at potential specific alteration in these two sub-processes in obesity. For instance, obesity has been associated with aberrant dopamine transmission, with there being an abundance of literature linking obesity to changes in D2 receptor availability in the striatum (see e.g. Horstmann et al., 2015). However, results are not consensual, with studies reporting decreased, increased, or unchanged D2 receptor availability in obesity (Ribeiro et al., 2023; Janssen & Horstmann, 2022; see Darcey et al. (2023) for a potential explanation). Additionally, there are reports of differences in dopamine transporter (DAT) availability in both obese humans (Chen et al., 2008; but also see Pak et al., 2023) and rodents (Narayanaswami et al., 2013; Jones et al., 2021; Hamamah et al., 2023). The observed changes in dopamine are often interpreted as being due to chronic

dopaminergic overstimulation resulting from overeating (Volkow & Wise, 2005; Volkow et al., 2008) and altered reward sensitivity as a consequence thereof (Blum et al., 1996). Considering that working memory gating is highly dependent on dopamine signaling, such changes could theoretically alter the balance between maintenance and updating processes in obesity. Next to this, obesity has frequently been associated with functional and structural changes in WM gating-related brain areas, implying another pathway through which working memory gating might get affected. At the level of the prefrontal cortex (PFC), studies have reported reduced gray matter volume and compromised white matter microstructure in individuals with obesity (Debette et al., 2014; Kullmann et al., 2016; Morys et al., 2024; Lv et al., 2024), and functional changes become evident with frequent reports of decreased activity in the dorsolateral PFC during tasks requiring cognitive control (e.g., Morys et al., 2018; Xu et al., 2017). Notably, Han et al. (2022) observed significantly lower spontaneous dlPFC activity during rest, potentially indicating reduced baseline dlPFC activity in obesity. On the level of the striatum, gray matter volume seems to correlate positively with measures of obesity (Horstmann et al., 2011), and individuals with obesity show greater activation of the dorsal striatum in response to high-calorie food stimuli compared to normal-weight individuals, indicating a stronger dopamine-dependent reward response to food cues (Stice et al., 2008; Small et al., 2003). Additionally, changes in connectivity between and within the striatum and PFC in obesity, both structurally (Li et al., 2023) and functionally (Verdejo-Román et al., 2017a, 2017b; Contreras-Rodríguez et al., 2017) have been reported. Although these studies mostly investigate brain function in relation to food and reward processing, changes in these areas may also impair the ability to adequately engage in working memory gating processes, as activity in affective (reward) and cognitive fronto-striatal loops immensely overlap (Janssen et al., 2019). On the behavioral level, individuals with obesity consistently demonstrate impairments in food-specific (Janssen et al., 2017) but also non-food specific goal-directed behavioral control (Janssen et al., 2020) and reinforcement learning (Weydmann et al., 2023). It seems that difficulties with integrating negative feedback may be central to these alterations (Mathar et al., 2017; Kastner et al., 2017), which could explain a potential insensitivity to the negative consequences associated with (over) eating. Crucially, in humans, a substantial contribution to (reward) learning is mediated by working memory processes (Moustafa et al., 2008; Collins & Frank, 2012, 2018; Collins et al., 2014, 2017; Westbrook et al., 2024). The observed difficulties in reward learning in obesity may hence partly be rooted in a failure to update working memory with new reward information, suggesting cognitive issues that extend beyond mere difficulties in valuation processes. However, empirical support for this interpretation is currently lacking. A more nuanced understanding of the effects of obesity on working memory is crucial, however, as it could lead to more targeted intervention options.

In the present study we therefore aim to examine potential obesity-dependent alterations in working memory gating. To this end, we pooled together data on Body-Mass-Index (BMI; kg/m²) and a working memory gating task from three different studies conducted in our lab. In light of the behavioral and neuropharmacological findings discussed above, we hypothesized that individuals with a high BMI would display worse updating, potentially offset by enhanced distractor-resistant maintenance.

Given dopamine's central role in WM gating, such behavioral patterns might be driven by the altered dopamine signaling observed in obesity (as discussed above). However, in addition to this, inherent predispositions with respect to dopamine signaling have also contribute. In this

context, several single-nucleotide polymorphisms (SNPs) related to dopamine transmission have garnered significant attention in recent years. For instance, Catechol-O-methyltransferase (COMT) Val158Met activity primarily influences dopamine breakdown in the PFC (Tunbridge et al., 2004; Sesack, Hawrylak, Matus, Guido, & Levey, 1998), and carrying the Met allele of this SNP is associated with reduced COMT activity, leading to higher synaptic dopamine levels (Bilder et al., 2004). Consistent with this, individuals with the Met allele tend to perform better on tasks that require stable maintenance of working memory representations compared to those with the Val allele (Berryhill et al., 2013; Farrell et al., 2012; Savitz, Solms, & Ramesar, 2006). Furthermore, the Taq1A polymorphism has been associated with D2 receptor density in the striatum. A-allele carriers of this polymorphism exhibit lower receptor density and show distinct performance patterns on tasks involving working memory updating (Pohjalainen et al., 1998; Jönsson et al., 1999; Eisenstein et al., 2016; Stelzel et al., 2010; Persson et al., 2015; Li et al., 2019). Interestingly, Taq1A and COMT have been demonstrated to interactively affect working memory functioning (Berryhill et al., 2013; Xu et al., 2007, Reuter et al., 2006, Gracia-Gracia et al., 2011; Stelzel et al., 2009, Wishart et al., 2011, Persson & Stenfors, 2018). Consequently, we aim to examine this interactive effect and assess if it varies with BMI. In addition, PPP1R1B and C957T gene polymorphisms have also been linked to working memory (Hotte et al., 2006; Ma et al., 2022; Smith, Swift-Scanlan & Boettiger, 2014; Xu et al., 2007; Klaus et al., 2019; Jacobsen et al., 2006). The PPP1R1B polymorphism (rs907094) codes for Dopamine and cAMP-regulated neuronal phosphoprotein (DARPP-32) - a protein that potently modulates dopamine D1-dependent synaptic plasticity in the striatum (Ouimet et al., 1984; Calabresi et al., 2000; Lindskog et al., 2006; Girault & Nairn, 2021). The C957T (rs6277) polymorphism, on the other hand, is known to impact dopamine D2 mRNA translation (Duan et al., 2003) and postsynaptic D2 receptor availability in the striatum (Hirvonen et al., 2005). Both polymorphisms have also been associated with (diet-induced) weight gain (Sharma & Fulton, 2013; Hu et al., 2006, Müller et al., 2012). Their effect on BMI-dependent WM gating, however, remains unknown. In order to test the impact of these four candidate polymorphisms we also added participants' genetic information to our analyses. We hypothesized that BMI-dependent distractor-resistant maintenance and/or updating of working memory representations would be modulated by (1) an interaction of COMT and Taq1A, a main effect of (2) DARPP-32, and/or a main effect of (3) C957T.

In addition to our primary investigations, we further conducted exploratory analyses on a subsample of our data. Two of the three studies had data available on the ratio of phenylalanine and tyrosine to other large neutral amino acids. This ratio represents the peripheral dopamine precursor availability and can be considered a potential proxy for central dopamine synthesis capacity (Leyton et al., 2004; Montgomery et al., 2003). Existing evidence suggests that this measure may be linked to WM performance in a diet-dependent manner (Hartmann et al., 2020). By looking at amino acid ratio and its connection to BMI-dependent WM gating, we sought to assess the possible influence of dopamine at the system level.

RESULTS

Sample descriptives

Three participants were excluded from the analyses, as they performed below chance (<50% correct in all four conditions). One subject was excluded as they reported that they didn't perform the task properly during the post-task strategy assessment. The final sample thus consisted of 320 participants. The average age of the sample was 26.93 years (SD = 6.82, min = 12.17, max = 49.75). There were 166 males. Mean BMI was 26.38 kg/m² (SD = 6.35, min = 17.51, max = 45.54). Mean IQ was 105.41 (SD = 10.61, min = 71, max = 122). Data for DFSQ was missing for 5 subjects. Mean DFSQ score was 54.89 (SD = 11.61, min = 33, max = 97). Please refer to Table 1 for a list of full sample characteristics (per study).

Taq1A Genotype Moderates the Association between BMI and Working Memory Updating, Independent of COMT

As expected, results for model 1 showed a significant main effect of BMI on overall task performance (X^2 = 16.80, df = 1, $p_{corrected}$ < 0.001), such that BMI was negatively associated with accuracy (OR = 0.84, CI = 0.78 – 0.91, see Figure 2). Against our main hypothesis, however, there was no difference in this effect between the working memory conditions. The two-way interaction between BMI and condition was insignificant (X^2 = 2.66, df = 3, $p_{corrected}$ > 1), indicating no evidence for BMI to have different effects across our working memory conditions. We found significant main effects of IQ, gender, tiredness, and concentration (all corrected p-values < 0.008). As expected, IQ and concentration were positively associated with task performance (OR_{IQ} = 1.24, CI_{IQ} = 1.14 – 1.35; $OR_{concentration}$ = 1.30, $CI_{concentration}$ = 1.20 – 1.41), while tiredness predicted task performance in a negative manner (OR = 0.87, CI = 0.80 – 0.95). Males performed worse than females on the task (OR = 0.86, CI =0.82 – 0.97). Please refer to Table 2 for the full model output displaying the original, uncorrected p-values.

When investigating the interactive effects of COMT and Taq1A on BMI-dependent WM gating (model 2), results reveal that the four-way interaction of BMI x condition x COMT x Taq1A was non-significant (X^2 = 4.09, df = 6, $p_{corrected}$ > 1). This indicates that the two SNPs did not have the expected differential effects on WM gating. There were no main effects of COMT (X^2 = 0.159, df = 2, $p_{corrected}$ > 1) or Taq1A (X^2 = 1.13, df = 1, $p_{corrected}$ > 1), and all other two- or three-way interactions involving COMT were insignificant (all corrected p-values > 0.34). Interestingly, however, we could observe a significant three-way interaction between Taq1A genotype, BMI, and condition (X^2 = 12.40, df = 3, $p_{corrected}$ = 0.024), indicating that Taq1A genotype might moderate BMI-dependent effects on working memory gating. Please refer to Table 3 for the full model output with original, uncorrected p-values.

To further investigate the significant Taq1A x condition x BMI interaction, we ran simple effects analyses, testing the Taq1A-BMI interaction separately for each condition. These analyses showed that the BMI-genotype interaction was significant in the update condition (p = 0.002), but not in the other three conditions (all p-values > 0.079), suggesting that the effect was specific to updating and hence might drive the observed overall three-way interaction (Fig. 3).

Further *post hoc* examination of the effects on updating revealed that, the association between BMI and performance was significant for A1-carriers (95%CIs: -0.488 to -0.190), with 33.9% lower probability to score correctly per unit change in BMI, but non-significant for non-A1-carriers (95%CIs: -0.153 to 0.129; 1.22% lower probability). Interestingly, compared to all other conditions, in the update condition, the negative association between BMI and task performance was weakest for non-A1-carriers (estimate = -0.012, SE = 0.072, but strongest for A1-carriers (estimate = -0.339, SE = 0.076; see Figure 3 and Table S6), emphasizing that genotype impacts this condition the most. To further check if this difference in slope was statistically significant across conditions, we stratified the sample into Taq1A subgroups (A1+ vs. A1-) and assessed whether BMI affected task performance differently across conditions separately for each subgroup. This analysis revealed no significant difference in the relationship between BMI and task performance across conditions among A1+ individuals ($p_{BMI*condition}$ = 0.219). However, within the A1- subgroup, a significant interaction effect between BMI and condition emerged ($p_{BMI*condition}$ = 0.049). Collectively, these findings suggest that the absence of the A1-allele is linked to improved task performance, particularly in the context of updating, where it seems to mitigate the otherwise negative effects of BMI.

In order to determine whether our results stemmed from mere match/non-match response biases or from proper ignoring/updating, we conducted a follow-up analysis, investigating the effects of the probe type presented at the end of each trial. The probe could either be the target item, a completely novel item, or a distractor item, meaning that the probe was one of the items that had to be encoded initially, but then be overwritten. Thus, a distractor probe measures the cognitive challenge of updating in its most exact form, while a target or novel item primarily assay match/non-match responses. For this analysis, we thus subset our data, including updating trials only, and re-ran our model augmented with the factor probe type. Results showed a significant main effect of probe type (X^2 = 94.11, df = 2, p < 0.001). trials where the probe was a distractor were the hardest (mean accuracy = 86.44%), followed by target probe trials (mean accuracy = 91.58%), and novel probe trials (94.96%). The three-way interaction between probe type, BMI, and Taq1A genotype was not significant (X^2 = 1.645, df = 2, p = 0.439), indicating that the probe type did not affect the BMI-Taq1A interaction in updating trials. High-BMI A-allele carriers were worse than non-carriers in all three probe trial types similarly. However, this pattern was most pronounced in the distractor condition (see Figure 4).

DARPP-32 Genotype Moderates the Association between BMI and Working Memory updating

Investigation of the effects of DARPP-32 on BMI-dependent working memory gating, revealed a similar picture to Taq1A. We found a significant 3-way interaction of DARPP-32, BMI, and condition (X^2 = 20.21, df = 3, $p_{corrected}$ < 0.001), such that DARPP-32 interacted with BMI in the update condition only ($p_{posthoc}$ = 0.006). For the full model output with original, uncorrected p-values, please refer to Table 4. Once more, further examination of the observed DARPP-32, BMI, and condition interaction showed that, in the update condition, the negative association between BMI and task performance was weakest and non-significant for A/A (estimate = -0.044, SE = 0.066; 95%CIs: -0.174 to 0.086), but strongest and significant for G-carrying individuals (estimate = -0.324, SE = 0.079; 95%CIs: -0.478 to – 0.170). See Table S7 and Figure 5. Splitting the sample in to DARPP subgroups (A/A vs. G-carrier) revealed that in both subgroups, there was

significant interaction effect of BMI and condition on task performance ($p_{A/A}$ = 0.034, $p_{G\text{-carrier}}$ = 0.003). In the case of DARPP, it hence appears that carrying the disadvantageous G-allele could exacerbate the negative effects of BMI, while the more advantageous allele (A/A) might mitigate them - once again particularly in the context of updating.

To further explore this association, we ran a follow-up analysis, investigating whether the type of the probe item had an effect. We subset our data, including only updating trials, and re-ran our model augmented with the factor probe type. Results revealed a significant main effect of probe type (p < 0.001), and a significant three-way interaction between probe type, BMI, and DARPP genotype (X^2 = 10.792, df = 2, p = 0.005). Post hoc analyses indicated that this interaction was driven by a significant DARPP-BMI interaction in distractor (p = 0.046) and target (p = 0.008) trials, while there was no such interaction in novel trials (p = 0.242; see Figure 6).

No Association of C957T with BMI-dependent Working Memory Gating
Our analysis revealed no significant main effect of the C957T polymorphism (X^2 = 0.03, df= 1, $p_{corrected}$ > 1). All other main effects stayed significant (all corrected p < 0.012), except for the effect of BMI (X^2 = 3.49, df= 1, $p_{corrected}$ = 0.247). Furthermore, we found no substantial evidence for two- or three-way interactions involving the C957T polymorphism (all corrected p > 0.186), suggesting that C957T does not significantly interact with BMI or one of our working memory conditions. See Table 5 for the full model output. Because the main effect of BMI dissipated when including C957T in the model, we ran an additional exploratory analysis to check whether this polymorphism directly related to BMI. Linear regression, predicting BMI by genotype, showed no association between the two (p = 0.2432), indicating that BMI effect is probably not masked by the presence of the C957T polymorphism. See Table S8.

BMI-dependent Alterations in Working Memory Gating are Associated with Peripheral Dopamine Synthesis Capacity
When investigating potential influences of dopamine changes on the system level (model 5), we found a significant three-way interaction between amino acid ratio, BMI, and condition (X^2=10.88, df = 3, $p_{corrected}$ < 0.049). Post hoc simple effects analyses suggested that this interaction seems to be driven by differential performance specifically in update vs. ignore ($X2$ = 5.57, df = 1, p = 0.018). As BMI increases, higher ratios of amino acids promote better performance in updating, but worse performance in ignoring (see Figure 7, upper panel). All other comparisons (update vs. control short; ignore vs. control long; control long vs. control short, update vs. control long, ignore vs. control short) did not yield significant differential relationships between amino acid ratio and BMI (all p values > 0.168).

The main effects of BMI, condition, and amino acid ratio were insignificant (all $p_{corrected}$ > 1). The main effect of z-IQ (X^2= 11.64, df = 1, $p_{corrected}$ = 0.002) and z-concentration (X^2= 18.60, df = 1, $p_{corrected}$ < 0.001) were significant, both relating positively to performance (OR_{IQ} = 1.28, $OR_{concentration}$ = 1.33). The interactions between BMI and condition (X^2= 9.80, df = 3, $p_{corrected}$ = 0.081), and between amino acid ratio and condition (X^2= 8.69, df = 3, $p_{corrected}$ = 0.135) were not significant. BMI and amino acid ratio showed no significant two-way interaction (X^2 = 0.322, df = 1, $p_{corrected}$ > 1). Because there was an extreme BMI data point, we re-ran the model excluding this data point to check whether the results still hold. The three-way interaction between Amino Acid Ratio, BMI, and condition became trend-significant ($p_{corrected}$ = 0.063, see Table S1 & Figure S1).

DISCUSSION

The present investigation sought to evaluate whether obesity might be associated with impairments in working memory gating. Consistent with previous literature (Yang et al., 2020, 2019, 2018; Gonzales et al., 2010; Coppin et al., 2014, Hartmann et al., 2023) we found evidence for impairments in overall working memory in individuals with a high BMI. Yet, we could not observe the expected interaction of BMI and condition, indicating no specific associations between BMI and WM gating. Interestingly, however, distinct effects of BMI on gating became apparent when taking into account potential changes in inherent dopamine signaling. Specifically, Taq1A and DARPP-32 particularly affected performance in the updating condition. Against our expectation, however, we did not find evidence for an interaction effect of COMT and Taq1A on BMI-dependent working memory gating, nor did we find any effects of C957T.

Selective BMI-Genotype Effects on Working Memory Updating

Our findings are partially in line with our hypothesis. While we did observe the expected worsening of updating WM contents in individuals with a high BMI, this effect was not exclusive to updating. Only when participants - along with BMI - were categorized based on certain putative dopamine-signaling characteristics, distinct effects on updating became apparent. This finding is compelling as it demonstrates a rarely observed selective effect.

Notably, it were the Taq1A and DARPP-32 SNPs that selectively modulated WM updating in a BMI-dependent manner. Intriguingly, both of these SNPs are associated with, predominantly, striatal dopamine signaling (Hemmings & Greengard, 1986, Meyer-Lindenberg et al., 2007, Gluskin & Mickey, 2016), implying a targeted modulation of processes occurring within the striatum. Both, the A-allele (Taq1A) as well as the G-allele (DARPP-32), have previously been considered risk alleles for various conditions and behaviors involving maladaptive cognitive flexibility, such as addiction (Smith et al., 2008; Munafo, Matheson, & Flint, 2007; Deng et al., 2015), Schizophrenia (Meyer-Lindenberg et al., 2007; González-Castro, 2016; Albert et al., 2002), or impaired reinforcement learning (Frank et al., 2007; Doll et al., 2011). Noteworthy, our data revealed that differences in updating appeared to be driven by the non-risk allele groups. Despite increasing BMI, performance remained stable. This pattern suggests that possessing the more advantageous genotype could potentially mitigate the generally negative effects of a high BMI on working memory updating. Moreover, in the normal-weight BMI range, carriers of a risk allele (in both, Taq1A and/or DARPP) slightly outperformed their non-risk allele carrying counterparts. This is especially intriguing as it emphasizes that carrying a "risk-allele" can in fact be advantageous under certain cognitive demands – a claim that has also been put forward by e.g. Stelzel et al. (2010).

Finally, the effects of genotype were particularly pronounced in trials where the probe was a distractor, suggesting that the effect is primarily due to "real" updating, i.e. when initially encoded items need overwriting, as opposed to simple match/non-match responses (as in novel vs. target probe items).

Potential Mechanistic Accounts

Mechanistically, our findings are potentially due to differential go/no-go path activation in the basal ganglia – pathways that are crucially involved in governing working memory gating. In essence, the D1 pathway modulates the 'go' signaling responsible for updating, while the D2 pathway facilitates 'no-go' signaling crucial for distractor-resistant maintenance (Frank & O`Reilly, 2006).

Considering Taq1A, evidence points at increased striatal dopamine synthesis and corresponding increases in striatal BOLD signals in A-carriers compared to non-carriers (Laakso et al. 2005; Stelzel et al., 2010). These findings suggest that the phasic dopamine signal needed to trigger "go" (i.e. updating), might be enhanced in A-carriers. This aligns with the idea that A-carriers, who possess fewer D2 receptors (Thompson et al., 1997; Pohjalainen et al., 1998; Jönsson et al., 1999), fall more within the ambit of the D1/go-dominant regime (Klein et al., 2007). Our data support this speculation by revealing slightly better updating performance in A-carriers in the normal-weight BMI range. However, as BMI increases, the possession of a greater D2 receptor density seems to become advantageous, as evidenced by the lack of a negative correlation between BMI and updating performance in non-A carriers. We speculate that this phenomenon could potentially be attributed to the compensating effects of this genotype. While individuals with fewer D2 receptors (A1+) may have quicker saturation of receptors regardless of dopamine levels, in those with more D2 receptors (A1-) saturation may be slower. This could contribute to a more finely tuned balance between "go" and "no-go" signaling, despite potential alterations in dopamine tone in obesity (Horstmann et al., 2015; but also see Darcey et al., 2023 or Janssen & Horstmann, 2022). Clearly, the current data cannot provide empirical evidence for these speculations, and further discrete research is needed to establish firm conclusions.

Regarding DARPP, we found that carrying the G-allele significantly exacerbated the negative effects of BMI, while the more advantageous allele (A/A) mitigated them, once again particularly in the context of updating. Interestingly, the G-allele has been associated with reduced striatal D1 efficacy (Meyer-Lindenberg et al., 2007). Moreover, Frank et al. (2007) and Doll et al. (2011) showed that carrying a G-allele was associated with worse go-learning – a process requiring activation of the same go-pathway that is likely to be activated during updating of working memory contents. Similarly, Frank and colleagues (2009) demonstrate that the G-carrier group compared to the A/A allele group displayed worse approach learning – again a process relying on go-path activation. However, there were no effects of DARPP on no-go learning (Frank et al., 2007), which requires D2-mediated no-go path activation. Our results hence broadly align with the literature and suggest that, particularly, markers of striatal go-signaling modulate BMI-dependent effects on working memory updating.

Collectively, our observations hint at the potential of advantageous genotypes to moderate the adverse impacts of high BMI on cognitive functions.

Possible Accounts for the Absence of COMT and C957T Effects

Considering that according to the prevailing models, PFC and striatum interact to foster effective WM gating (Cools & D'Esposito, 2011), the question arises as to why we could not observe the expected COMT-Taq1A interaction on BMI-dependent working memory gating. We posit several explanations for the absence of the anticipated interaction.

Firstly, these polymorphisms may indeed exert a limited interactive effect on working memory gating. In line with this notion, prior findings concerning the interactive effects of COMT and Taq1A on working memory have yielded contradictory results. For instance, Garcia-Garcia et al. (2011) and Stelzel et al. (2009) reported patterns of COMT-Taq1A interactions in the context of working memory updating that were consistent in terms of direction of effect. In contrast, Wishart et al. (2011) observed an opposing interaction pattern, while Persson & Stenfors (2018) did not identify any COMT-Taq1A interaction at all. All these studies explored genotype

interactions using paradigms that either assessed the two memory processes separately (Gracia-Garcia et al., 2011; Wishart et al., 2011; Persson & Stenfors, 2018) or in a manner that they were not distinctly discernible (Berryhill et al., 2013). None of them examined the comprehensive interaction of COMT, Taq1A, and working memory updating vs. ignoring within a single paradigm, as we did here. Secondly, it has recently been debated whether COMT has a noteworthy effect on cognition. Some meta-analyses find (small) effects (Barnett et al., 2007), while others don't (Geller et al., 2017, Barnett et al., 2008; also see Goldman et al., 2009; Wacker, 2011; and Barnett et al., 2011 for a discussion on the meta-analysis from Barnett et al., 2008).

Beyond this, the absence of significant effects related to COMT could further be interpreted as underscoring the selectiveness of our observed effects. COMT effects are predominantly observed in the PFC (Mier, Kirsch & Meyer-Lindenberg, 2010; Egan et al., 2011; Käenmäki et al., 2010) and rather tied to maintenance of working memory contents (Nolan et al., 2004; Rosa et al., 2010). This lends weight to the interpretation that distractor-resistant maintenance, or prefrontal processes, remain unaffected by BMI. In a similar vein, also our findings concerning the C957T polymorphism bolster the selective nature of our findings. Much like COMT, this polymorphism is presumably more involved in working memory maintenance, that is, prefrontal-related functioning. Supporting this notion, Xu and colleagues (2007) found an association between the C957T polymorphism and specifically maintenance of (phonological and serial) information, but not with other tasks requiring updating. Furthermore, this polymorphism has also been associated with D2 binding potential in extrastriatal regions (Hirvonen et al., 2009b) and greater WM-related activity in PFC (Li, Bäckmann & Persson, 2019). Lastly, the results with respect to C957T involvement in striatal-dependent cognition are mixed (see e.g. discussion part in Baker, Stockwell, & Holroyd, 2013), indicating that C957T may not be a good candidate for influencing striatal-dependent processes.

Selective Modulation of Working Memory Gating: System-Level Dopamine versus Genetic Profiles
Our BMI-gene findings show a selective modulation of working memory updating, as opposed to the previously observed trade-off between ignoring and updating (e.g. Fallon and Cools, 2017; see Cools (2019) for an extensive review). To the best of our knowledge, none of the previous studies investigating working memory gating in relation to dopamine signaling have found such a selective modulation. We speculate that this is because previous studies looked at broader changes in the dopamine system, i.e. by using drug manipulations or comparing Parkinson's vs. healthy controls, rather than particular genetic profiles. Such broad "system-level" dopamine changes may impact both PFC-facilitated distracter resistance and striatal-dependent updating. This, in turn, might foster the commonly observed inverted-U-shaped relationship between dopamine and cognition: In cases where baseline dopamine levels are low, a dopamine increase (for instance, through agonists) would enhance ignoring, albeit at the expense of updating. Conversely, at medium baseline dopamine levels an increase would lead to impaired ignoring, potentially benefiting updating (for a more detailed discussion, see Cools and D'Esposito, 2011). Indeed, we also see this pattern when looking at system-level dopaminergic changes: depending on BMI, low (or high) peripheral dopamine synthesis capacity (as indicated by blood amino acid ratios) was associated with worsening of distractor-resistant maintenance, while improving updating (or vice versa). It should be noted that the sample for our amino acid analyses was much

smaller (N = 160) than the one used in our SNP analysis (N = 320), and the BMI-range for this sub-sample was narrower (mean = 23.63, SD = 2.78, min = 18.63, max = 36.42). This was because only two of the three studies had the data on amino acids available. Interestingly, the system-level effect of amino acid ratio becomes visible in a healthy- to overweight BMI range, indicating that already small changes in BMI can promote different dopamine-dependent cognitive profiles.

Strengths and Limitations
A major strength of this study is that it was the first to probe gene-gene interactions on a direct working memory maintenance and updating comparison. This is a notable advantage, as previous studies have usually examined these aspects in separate paradigms, which might have contributed to the heterogeneous results regarding SNP interaction effects on working memory (see above). Another main strength of our study is the sizable sample. However, despite this relatively large size, our sample was still not big enough for systematic and reliable analysis of multiple gene-gene interactions. This would be of interest however, as possible interactions of all SNPs presented in this paper have been shown (Zmigrod & Robbins, 2021; Frank & Hutchison, 2009; Smith, Swift-Scanlan & Boettiger, 2014, Xu et al., 2007). Such analyses would require even larger cohorts, however, as the effect sizes of single SNPs are usually small. We nevertheless report the outcome of such highly explorative models in our supplements for the purpose of transparency and to guide future studies. Yet, those results should be interpreted with caution. Furthermore, an additional limitation is that our data is slightly skewed towards participants within the normal BMI range. The effective sample size to detect meaningful genotype effects (e.g. for COMT or C957T) might thus have been too small, particularly at higher BMI levels. Future studies may address this limitation by recruiting a more balanced sample, including more individuals with higher BMI. Additionally, the correlational nature of our findings highlights the need for more direct experimental manipulations of dopaminergic processes in obesity. Previous studies have established a causal link between dopamine and WM gating through drug manipulations (Fallon et al., 2017, 2019). Applying a similar approach to an obese sample could help establish a clearer causal link between dopamine activity and WM gating in the context of obesity. Lastly, the sample used for this study was very heterogeneous, as it was pooled from three separate studies. The BMI distribution, for instance, was significantly different depending on gender. BMI was higher in females. This was because females were overrepresented in the BEDOB study, which had the largest BMI range (refer to Table 1). Although we ran control analyses to account for this heterogeneity, we cannot exclude the possibility that certain properties of the data distribution could have influenced our results.

Overall Conclusions
Overall, our data aligns with previous evidence for working memory impairments in obesity. However, selective effects of BMI on working memory gating - specifically updating - become visible only when accounting for genetic markers of striatal dopamine transmission. This level of specificity adds a new nuance to existing research as it demonstrates condition-specific effects of BMI on working memory. Previous research, that generally utilized drug manipulations, consistently demonstrated system-level modulations, leading to a trade-off between ignoring and updating information. While we also observe this trade-off when examining more comprehensive

system-level relationships (i.e. blood amino acid ratio), the specificity of our SNP-related findings to updating sets them apart from previous studies. Our results hence pave the way for new individualized treatments for obesity, as they highlight the potential of advantageous genotypes to mitigate the adverse effects of high BMI on cognitive functions that require updating of information, and suggest that previously documented deficits in reward learning, which partially rely on information updating, could potentially be targeted more specifically when taking genotypes into account.

MATERIALS AND METHODS
Participants
The data used in this study were collected in the scope of three separate pre-registered cross-sectional studies, which are all part of a larger line of research in the O'Brain Lab: GREADT (see https://osf.io/w9e5y), BEDOB (see https://osf.io/fyn6q), and WORMCRI (see https://osf.io/zdmkx). Prior to participation, participants were screened for a history of clinical drug or alcohol abuse, neurological or psychiatric disorders, and first-degree relative history of neurological or psychiatric disorders. Symptoms of depression were assessed via a screening interview using the Structured Clinical Interview for DSM-IV (SCID, Wittchen 1997; in BEDOB & WORMCRI) or Beck Depression Inventory (BDI, Beck et al., 1996; in GREADT).

Study Design
All measures relevant to the present study were collected in a comparable manner. In all studies, participants were first asked to come to the lab for a screening session where in- and exclusion criteria were checked. Weight and height were measured to calculate BMI. After inclusion, blood samples were taken from the participants to assess COMT Val[158]Met, Taq1A, C957T, and DARPP-32 genotypes. Analysis of these SNPs was performed in the laboratory for 'Adiposity and diabetes genetics' at the Medical Research Center, University Leipzig, Leipzig, Germany. In WORMCRI and GREADT, we also took serum blood samples in order to extract information on the amino acid profiles. Participants, therefore, came overnight-fasted for these two studies. Serum blood samples were analyzed at the "Institut für Laboratoriumsmedizin, Klinische Chemie und Molekulare Diagnostik (ILM)" Universitätsklinikum Leipzig, Germany. After the blood draw, participants did a number of neuropsychological tests among which were the digit span task (Wechsler, 2008; assessing baseline working memory), and a proxy for IQ (in BEDOB: "Wortschatztest" (Schmidt & Metzler, 1992; assessing verbal IQ); in GREADT and WORMCRI: "Wiener Matrizen Test" (Formann, Waldherr & Piswanger, 2011). After that, participants filled in several questionnaires, of which the Dietary Fat and free Sugar Questionnaire (DFSQ; Francis & Stevenson, 2013; Fromm & Horstmann, 2019; assessing eating behavior), was subject to all three studies. On the second test day, participants completed the working memory task (described below), either during fMRI (GREADT & BEDOB) or during EEG (WORMCRI). After completion of the task, all participants were asked to indicate the level of tiredness and concentration they felt during the task on a 10-point Likert scale. For a more detailed description of each study's design, please refer to the respective pre-registration mentioned above.

Working Memory Task
Participants completed a modified version of a delayed match-to-sample task originally designed by Fallon and Cools (2014). This modified version has already been described in Hartmann et al. (2023) and Herzog et al. (2023). The task comprises 4 conditions (Fig. 1). In the ignore condition, testing distractor-resistant maintenance, participants first have to memorize two target stimuli, signaled by the letter "T" centered in between the two. Next, they are presented with two new stimuli, this time marked by a centered "N", indicating non-targets which have to be ignored. After that, participants are presented with a probe stimulus and have to determine whether one of the first two target stimuli matches the presented probe. In the update condition, participants are first shown two target stimuli (centered "T"). After that, they see a new set of target stimuli (again indicated by a centered "T"). These two new stimuli replace the previously presented stimuli as the target and thus have to be evaluated for a match when the probe is presented subsequently. The two control conditions do not have any interference and are matched to the temporal delay between encoding the to-be-matched targets and probe. The probe is presented for 2000 msec. The task is separated into four blocks, with each block entailing 8 trails of each condition, interleaved among all blocks. Each block thus consists of 32 trials. The total number of trials in the task amounts to 128. Feedback is presented after each of those blocks. Each trial is separated

by a jittered inter-trial interval ranging from 2000 to 6000 msec. The stimuli are randomly computer-generated, monochromatic RGB "spirographs". The primary outcome measure is accuracy. The total duration of the task is approximately 30 minutes.

Statistical Analyses of Behavioral Data

All behavioral analyses were performed in R in RStudio v4.2.2 (R Core Team, 2015; RStudio Team, 2016). Given the within-subject design of our study, we used generalized linear mixed models (GLM) of the '*lme4*' package to analyze the primary outcome measure of the working memory task: accuracy. We ran a logistic regression using *glmer()* with a binomial link function. We used trial-by-trial information for each subject with binary coded response (0 = incorrect; 1 = correct). Trials with a reaction time < 200 ms and > 2000 ms were excluded, as those trials can be considered false alarms and misses, respectively. Trials with a reaction time > 2000 ms were excluded, as they reflect misses. To first test our main hypothesis that working memory gating is altered, depending on BMI, we built a trial-based regression model including the interaction of the within-subject factor condition (ignore vs. update vs. control long vs. control short) and the continuous between-subject factor BMI. We further probed the influence of several potential covariates: study (GREADT vs. BEDOB vs. WORMCRI), IQ, Age, DFSQ, binge-eating phenotype, tiredness, concentration, and gender. Using the *anova()* function from the '*stats*' package, we compared AIC and BIC (Akaike, 1979; Stone, 1979) of the full model against a simpler version of the model. We found the best-fitting model (lowest AIC and BIC) to include IQ, tiredness, concentration, and gender (see supplementary materials). Due to model convergence problems, the continuous predictors BMI, IQ, tiredness, and concentration were z-scored. Furthermore, the model did not converge with a maximal random structure (including the within-subject factor 'condition'). The random structure of the model was thus reduced to include the factor 'subject' only, thereby accounting for the repeated measures taken from each subject. The final model was:

$$(1) \quad accuracy \sim condition * BMI + IQ + tiredness + concentration + gender + (1 \,|subject)$$

To test how BMI-dependent working memory gating is moderated by the respective dopamine proxy (SNP or amino acid ratio), we ran 4 additional models, each including the respective between-subject factor as an additional factor of interest. Model fit was again assessed using AIC and BIC for each model (see supplementary materials). Again, due to converges, the random structure of the models included the factor 'subject' only. The final models were:

$$(2) \quad accuracy \sim COMT * Taq1A * condition * BMI + IQ + tiredness + concentration + gender + (1|subject)$$

$$(3) \quad accuracy \sim DARPP * condition * BMI + IQ + tiredness + concentration + gender + (1|subject)$$

$$(4) \quad accuracy \sim C957T * condition * BMI + IQ + tiredness + concentration + gender + (1|subject)$$

$$(5) \quad accuracy \sim amino\ acid\ ratio * condition * BMI + IQ + concentration + gender + (1|subject)$$

As we ran 4 additional models testing similar hypotheses, all main results for these models were corrected for multiple comparisons using bonferroni correction, i.e. p-values were multiplied by 4. Model outputs were called using the *Anova()* function, from the '*car*' package. Reported odds ratios (OR) are retrieved from exponentiating the log-odds coefficients called with the *summary()* function.

Acknowledgements
The authors thank Sylvia Stasch, Miriam Huml, Eva Burmeister, Elena Cesnaite and Lisa Okhof for helping with recruitment of participants and data collection. Furthermore, we thank Ines Müller from Peter Kovacs lab at the University of Leipzig for the analyses of our blood samples. Last but not least, we thank Susan Prejawa for her assistance in study organization and financial management.

Author contribution statement
NH, LJ and AH designed the studies. NH, LJ, HH, and MW collected the data. NH analyzed the data. NH drafted the manuscript. Code review was done by HH. HH, AK, LJ, PK, AV, MW, LD, SF, and AH critically revised and approved the final manuscript.

Conflict of interest statement
No potential conflicts of interest relevant to this article were reported

Data availability statement
Data and scripts used for the analysis will be available at https://github.com/O-BRAIN/WM_SNP upon publication

Declaration of Generative AI and AI-assisted Technologies in the writing process
During the preparation of this work, the author used ChatGPT to improve readability and language. After using this tool, the author reviewed and edited the content as needed and takes full responsibility for the content of the publication

REFERENCES

Akaike, H. (1979). A Bayesian extension of the minimum AIC procedure of autoregressive model fitting. Biometrika, 66(2), 237-242

Albert KA, Hemmings HC Jr, Adamo AI, Potkin SG, et al. (2002). Evidence for decreased DARPP-32 in the prefrontal cortex of patients with schizophrenia. Arch. Gen. Psychiatry 59: 705-712

Anticevic, A., Repovs, G., Krystal, J. H., & Barch, D. M. (2012). A broken filter: prefrontal functional connectivity abnormalities in schizophrenia during working memory interference. Schizophrenia research, 141(1), 8-14.

Badre, D. (2012). Opening the gate to working memory. Proceedings of the national academy of ofciences, 109(49), 19878-19879. https://www.pnas.org/doi/abs/10.1073/pnas.1216902109

Baker, T. E., Stockwell, T., & Holroyd, C. B. (2013). Constraints on decision making: implications from genetics, personality, and addiction. *Cognitive, Affective, & Behavioral Neuroscience*, *13*, 417-436

Barnett, J. H., Jones, P. B., Robbins, T. W., & Müller, U. (2007). Effects of the catechol- O - methyltransferase Val 158 Met polymorphism on executive function: A meta-analysis of the Wisconsin Card Sort Test in schizophrenia and healthy controls. Molecular Psychiatry, 12(5), 502–509. https://doi.org/10.1038/sj.mp.4001973

Barnett, J. H., Scoriels, L., & Munafò, M. R. (2008). Meta-Analysis of the Cognitive Effects of the Catechol-O-Methyltransferase Gene Val158/108Met Polymorphism. Biological Psychiatry, 64(2), 137–144. https://doi.org/10.1016/j.biopsych.2008.01.005

Barnett, J. H., Scoriels, L., & Munafò, M. R. (2011). Reply to: With Great Power Comes Great Responsibility: Meta-Analytic Proof for a Timeless Insight. Biological Psychiatry, 69(10), e39. https://doi.org/10.1016/j.biopsych.2010.11.028

Beck, A. T., Steer, R. A., & Brown, G. K. (1996). Manual for the beck depression inventory-II. San Antonio, TX: Psychological Corporation, 1, 82.

Berryhill, M. E., Wiener, M., Stephens, J. A., Lohoff, F. W., & Coslett, H. B. (2013). COMT and ANKK1-Taq-Ia genetic polymorphisms influence visual working memory. PLoS One, 8(1), Article e55862. https://doi.org/10.1371/journal.pone.0055862

Bilder, R. M., Volavka, J., Lachman, H. M., & Grace, A. A. (2004). The catechol-O-methyltransferase polymorphism: Relations to the tonic–phasic dopamine hypothesis and neuropsychiatric phenotypes. *Neuropsychopharmacology*, *29*(11), 1943-1961

Blum, K., Sheridan, P. J., Wood, R. C., Braverman, E. R., Chen, T. J. H., phd, J. C., & Comings, D. E. (1996). The D2 dopamine receptor gene as a determinant of reward deficiency syndrome. *Journal of the Royal Society of Medicine*, *89*(7), 396-400.

Braver, T. S., & Cohen, J. D. (2000). On the control of control: The role of dopamine in regulating prefrontal function and working memory. Control of cognitive processes: Attention and performance XVIII, (2000)

Calabresi, P., Gubellini, P., Centonze, D., Picconi, B., Bernardi, G., Chergui, K., & Greengard, P. (2000). Dopamine and cAMP-regulated phosphoprotein 32 kDa controls both striatal long-term depression and long-term potentiation, opposing forms of synaptic plasticity. *Journal of Neuroscience, 20*(22), 8443-8451

Chatham, C. H., & Badre, D. (2015). Multiple gates on working memory. *Current opinion in behavioral sciences*, *1*, 23-31

Chen, P. S., Yang, Y. K., Yeh, T. L., Lee, I. H., Yao, W. J., Chiu, N. T., & Lu, R. B. (2008). Correlation between body mass index and striatal dopamine transporter availability in healthy volunteers—a SPECT study. *Neuroimage*, *40*(1), 275-279.

Collins, A. G., & Frank, M. J. (2012). How much of reinforcement learning is working memory, not reinforcement learning? A behavioral, computational, and neurogenetic analysis. *European Journal of Neuroscience*, *35*(7), 1024-1035.

Collins, A. G., Brown, J. K., Gold, J. M., Waltz, J. A., & Frank, M. J. (2014). Working memory contributions to reinforcement learning impairments in schizophrenia. *Journal of Neuroscience*, *34*(41), 13747-13756.

Collins, A. G., Ciullo, B., Frank, M. J., & Badre, D. (2017). Working memory load strengthens reward prediction errors. *Journal of Neuroscience*, *37*(16), 4332-4342.

Collins, A. G., & Frank, M. J. (2018). Within-and across-trial dynamics of human EEG reveal cooperative interplay between reinforcement learning and working memory. *Proceedings of the National Academy of Sciences*, *115*(10), 2502-2507.

Colzato, L. S., Jongkees, B. J., Sellaro, R., & Hommel, B. (2013). Working memory reloaded: tyrosine repletes updating in the N-back task. Frontiers in behavioral neuroscience, 7, 200.

Contreras-Rodríguez, O., Martín-Pérez, C., Vilar-López, R., & Verdejo-Garcia, A. (2017). Ventral and dorsal striatum networks in obesity: link to food craving and weight gain. *Biological psychiatry*, *81*(9), 789-796.

Cools, R., & D'Esposito, M. (2011). Inverted-U–shaped dopamine actions on human working memory and cognitive control. Biological psychiatry, 69(12), e113-e125

Cools, R. (2019). Chemistry of the adaptive mind: lessons from dopamine. Neuron, 104(1), 113-131.

Coppin, G., Nolan-Poupart, S., Jones-Gotman, M., & Small, D. M. (2014). Working memory and reward association learning impairments in obesity. *Neuropsychologia*, *65*, 146-155.

Darcey, V. L., Guo, J., Chi, M., Chung, S. T., Courville, A. B., Gallagher, I., & Hall, K. D. (2023). Striatal dopamine tone is positively associated with body mass index in humans as determined by PET using dual dopamine type-2 receptor antagonist tracers. *medRxiv*.

D'Ardenne, K., Eshel, N., Luka, J., Lenartowicz, A., Nystrom, L. E., & Cohen, J. D. (2012). Role of prefrontal cortex and midbrain dopamine system in working memory updating. Proceedings of the National Academy of Sciences, 109(49), 19900-19909.

Deng, X. D., Jiang, H., Ma, Y., Gao, Q., Zhang, B., Mu, B., & Liu, Y. (2015). Association between DRD2/ANKK1 TaqIA polymorphism and common illicit drug dependence: Evidence from a meta-analysis. Human Immunology, 76(1), 42-51

Doll, B. B., Hutchison, K. E., & Frank, M. J. (2011). Dopaminergic genes predict individual differences in susceptibility to confirmation bias. *Journal of Neuroscience*, *31*(16), 6188-6198

Duan, J., Wainwright, M. S., Comeron, J. M., Saitou, N., Sanders, A. R., Gelernter, J., & Gejman, P. V. (2003). Synonymous mutations in the human dopamine receptor D2 (DRD2) affect mRNA stability and synthesis of the receptor. Human molecular genetics, 12(3), 205-216.

Durstewitz, D., & Seamans, J. K. (2008). The dual-state theory of prefrontal cortex dopamine function with relevance to catechol-o-methyltransferase genotypes and schizophrenia. Biological psychiatry, 64(9), 739-749.

Egan, M. F., Goldberg, T. E., Kolachana, B. S., Callicott, J. H., Mazzanti, C. M., Straub, R. E., & Weinberger, D. R. (2001). Effect of COMT Val108/158 Met genotype on frontal lobe function and risk for schizophrenia. *Proceedings of the National Academy of Sciences*, *98*(12), 6917-6922.

Eisenstein, S. A., Bogdan, R., Love-Gregory, L., Corral-Frías, N. S., Koller, J. M., Black, K. J., et al. (2016). Prediction of striatal D2 receptor binding by DRD2/ANKK1 TaqIA allele status. Synapse (New York, N.Y.), 70(10), 418. https://doi.org/10.1002/ SYN.21916

Fallon, S. J., Mattiesing, R. M., Muhammed, K., Manohar, S., & Husain, M. (2017). Fractionating the neurocognitive mechanisms underlying working memory: Independent effects of dopamine and Parkinson's disease. Cerebral Cortex, 27(12), 5727-5738.

Formann, A. K., Waldherr, K., & Piswanger, K. (2011). Wiener Matrizen-Test 2 (WMT-2): Ein Rasch-Skalierter Sprachfreier Kurztest zur Erfassung der Intelligenz [Viennese Matrices Test 2 (WMT-2): A Rapid-Scaled, Language-Free Short-Circuit Test for the Assesment of Intelligence]. Göttingen: Hogrefe.

Francis, H., & Stevenson, R. (2013). Validity and test-retest reliability of a short dietary questionnaire to assess intake of saturated fat and free sugars: a preliminary study. Journal of Human Nutrition and Dietetics, 26(3), 234–242. https://doi.org/10.1111/jhn.12008

Frank, M. J., & O'Reilly, R. C. (2006). A mechanistic account of striatal dopamine function in human cognition: psychopharmacological studies with cabergoline and haloperidol. *Behavioral neuroscience*, *120*(3), 497.

Frank, M. J., Moustafa, A. A., Haughey, H. M., Curran, T., & Hutchison, K. E. (2007). Genetic triple dissociation reveals multiple roles for dopamine in reinforcement learning. Proceedings of the National Academy of Sciences, 104(41), 16311-16316.

Frank, M. J., & Hutchison, K. (2009). Genetic contributions to avoidance-based decisions: striatal D2 receptor polymorphisms. Neuroscience, 164(1), 131-140

Fromm, S. P., & Horstmann, A. (2019). Psychometric Evaluation of the German Version of the Dietary Fat and Free Sugar-Short Questionnaire. Obesity Facts, 1–11. https://doi.org/10.1159/000501969

Furman, D.J., Zhang, Z., Chatham, C.H., Good, M., Badre, D., Hsu, M., Kayser, A.S. (2021) Augmenting Frontal Dopamine Tone Enhances Maintenance over Gating Processes in Working Memory. J Cogn Neurosci;33(9):1753-1765. doi: 10.1162/jocn_a_01641. PMID: 33054556; PMCID: PMC8647935.

Garcia-Garcia, M., Barceló, F., Clemente, I. C., & Escera, C. (2011). COMT and ANKK1 gene–gene interaction modulates contextual updating of mental representations. Neuroimage, 56(3), 1641-1647.

Geller, S., Wilhelm, O., Wacker, J., Hamm, A., & Hildebrandt, A. (2017). Associations of the COMT Val158Met polymorphism with working memory and intelligence–A review and meta-analysis. Intelligence, 65, 75-92.

Girault, J. A., & Nairn, A. C. (2021). DARPP-32 40 years later. *Advances in Pharmacology*, *90*, 67-87.

Gluskin, B. S., & Mickey, B. J. (2016). Genetic variation and dopamine D2 receptor availability: a systematic review and meta-analysis of human in vivo molecular imaging studies. *Translational psychiatry*, *6*(3), e747-e747

Goldman, D., Weinberger, D. R., Malhotra, A. K., & Goldberg, T. E. (2009). The Role of COMT Val158Met in Cognition. Biological Psychiatry, 65(1), e1–e2. https://doi.org/10.1016/j.biopsych.2008.07.032

Gonzáles, M. M., Tarumi, T., Miles, S. C., Tanaka, H., Shah, F., & Haley, A. P. (2010). Insulin sensitivity as a mediator of the relationship between BMI and working memory-related brain activation. *Obesity*, *18*(11), 2131-2137.

González-Castro, T. B., Hernandez-Diaz, Y., Juárez-Rojop, I. E., López-Narváez, M. L., Tovilla-Zárate, C. A., Genis-Mendoza, A., & Alpuin-Reyes, M. (2016). The role of C957T, TaqI and

Ser311Cys polymorphisms of the DRD2 gene in schizophrenia: systematic review and meta-analysis. Behavioral and Brain Functions, 12(1), 1-14

Hamamah, S., Hajnal, A., & Covasa, M. (2023). Reduced Striatal Dopamine Transporter Availability and Heightened Response to Natural and Pharmacological Stimulation in CCK-1R-Deficient Obese Rats. *International journal of molecular sciences*, *24*(11), 9773.

Han, X. D., Zhang, H. W., Xu, T., Liu, L., Cai, H. T., Liu, Z. Q., & Yuan, T. F. (2022). How impulsiveness influences obesity: the mediating effect of resting-state brain activity in the dlPFC. *Frontiers in Psychiatry*, *13*, 873953.

Hartmann, H., Pauli, L. K., Janssen, L. K., Huhn, S., Ceglarek, U., & Horstmann, A. (2020). Preliminary evidence for an association between intake of high-fat high-sugar diet, variations in peripheral dopamine precursor availability and dopamine-dependent cognition in humans. *Journal of Neuroendocrinology*, *32*(12), e12917.

Hartmann, H., Janssen, L. K., Herzog, N., Morys, F., Fängström, D., Fallon, S. J., & Horstmann, A. (2023). Self-reported intake of high-fat and high-sugar diet is not associated with cognitive stability and flexibility in healthy men. *Appetite*, *183*, 106477.

Hazy, T. E., Frank, M. J., & O'Reilly, R. C. (2006). Banishing the homunculus: making working memory work. Neuroscience, 139(1), 105-118

Hemmings, H. C. Jr., and Greengard, P. (1986). DARPP-32, a dopamine- and adenosine 3′:5′-monophosphate- regulated phosphoprotein: regional, tissue, and phylogenetic distribution. J. Neurosci. 6, 1469–1481.

Herzog, N. D., Hartmann, H., Janssen, L., Waltmann, M., Deserno, L., Fallon, S. J., & Horstmann, A. (2023). Working Memory Gating in Obesity: Insights from a Case-Control fMRI Study. *medRxiv*, 2023-09.

Hilbert, S., Nakagawa, T. T., Puci, P., Zech, A., & Bühner, M. (2014). The Digit Span Backwards Task. Http://Dx.Doi.Org/10.1027/1015-5759/A000223. https://doi.org/10.1027/1015-5759/A000223

Hirvonen, M., Lumme, V., Hirvonen, J., Pesonen, U., Nagren, K., Vahlberg, T. (2009). C957T polymorphism of the human dopamine D2 receptor gene predicts extrastriatal dopamine receptor availability in vivo, Prog. Neuropsychopharmacol. Biol. Psychiatry 33 (4) 630–636, https://doi.org/10.1016/j.pnpbp.2009.02.021

Hirvonen, M., Laakso, A., Någren, K. et al. Erratum: C957T polymorphism of the dopamine D2 receptor (DRD2) gene affects striatal DRD2 availability in vivo. Mol Psychiatry 10, 889 (2005). https://doi.org/10.1038/sj.mp.4001707

Horstmann, A., Fenske, W. K., & Hankir, M. K. (2015). Argument for a non-linear relationship between severity of human obesity and dopaminergic tone. Obesity Reviews, 16(10), 821-830

Hotte, M., Thuault, S., Lachaise, F., Dineley, K. T., Hemmings, H. C., Nairn, A. C., & Jay, T. M. (2006). D1 receptor modulation of memory retrieval performance is associated with changes in pCREB and pDARPP-32 in rat prefrontal cortex. *Behavioural brain research*, *171*(1), 127-133.

Hu, J., Redden, D. T., Berrettini, W. H., Shields, P. G., Restine, S. L., Pinto, A., & Allison, D. B. (2006). No evidence for a major role of polymorphisms during bupropion treatment. Obesity, 14(11), 1863-1867

Jacobsen, L. K., Pugh, K. R., Mencl, W. E., & Gelernter, J. (2006). C957T polymorphism of the dopamine D2 receptor gene modulates the effect of nicotine on working memory performance and cortical processing efficiency. *Psychopharmacology*, *188*, 530-540.

Janssen, L. K., Duif, I., van Loon, I., Wegman, J., de Vries, J. H., Cools, R., & Aarts, E. (2017). Loss of lateral prefrontal cortex control in food-directed attention and goal-directed food choice in obesity. *Neuroimage*, *146*, 148-156.

Janssen, L. K., Mahner, F. P., Schlagenhauf, F., Deserno, L., & Horstmann, A. (2020). Reliance on model-based and model-free control in obesity. *Scientific reports*, *10*(1), 22433.

Janssen, L. K., & Horstmann, A. (2022). Molecular imaging of central dopamine in obesity: A qualitative review across substrates and radiotracers. *Brain sciences*, *12*(4), 486.

Jones, S. R., & Fordahl, S. C. (2021). Bingeing on High-Fat Food Enhances Evoked Dopamine Release and Reduces Dopamine Uptake in the Nucleus Accumbens. Obesity, 29(4), 721-730.

Jongkees, B. J. (2020). Baseline-dependent effect of dopamine's precursor L-tyrosine on working memory gating but not updating. *Cognitive, Affective, & Behavioral Neuroscience*, *20*(3), 521-535

Jönsson, E. G., Nothen, M. M., Grünhage, F., Farde, L., Nakashima, Y., Propping, P., et al. (1999). Polymorphisms in the dopamine D2 receptor gene and their relationships to striatal dopamine receptor density of healthy volunteers. Molecular Psychiatry, 4(3), 290–296. http://www.ncbi.nlm.nih.gov/pubmed/10395223.

Käenmäki, M., Tammimäki, A., Myöhänen, T., Pakarinen, K., Amberg, C., Karayiorgou, M., & Männistö, P. T. (2010). Quantitative role of COMT in dopamine clearance in the prefrontal cortex of freely moving mice. *Journal of neurochemistry*, *114*(6), 1745-1755.

Kastner L, Kube J, Villringer A, Neumann J. Cardiac concomitants of feedback and prediction error processing in reinforcement learning. Front Neurosci. 2017; 11:598.

Klaus, K., Butler, K., Curtis, F., Bridle, C., & Pennington, K. (2019). The effect of ANKK1 Taq1A and DRD2 C957T polymorphisms on executive function: a systematic review and meta-analysis. Neuroscience & Biobehavioral Reviews, 100, 224-236.

Klein, T. A., Neumann, J., Reuter, M., Hennig, J., von Cramon, D. Y., & Ullsperger, M. (2007). Genetically determined differences in learning from errors. *Science*, *318*(5856), 1642-1645

Laakso, A., Pohjalainen, T., Bergman, J., Kajander, J., Haaparanta, M., Solin, O., & Hietala, J. (2005). The A1 allele of the human D2 dopamine receptor gene is associated with increased activity of striatal L-amino acid decarboxylase in healthy subjects. *Pharmacogenetics and genomics*, *15*(6), 387-391.

Li, X., Bäckman, L., & Persson, J. (2019). The relationship of age and DRD2 polymorphisms to frontostriatal brain activity and working memory performance. Neurobiology of Aging, 84, 189-199.

Li, G., Hu, Y., Zhang, W., Wang, J., Ji, W., Manza, P., & Wang, G. J. (2023). Brain functional and structural magnetic resonance imaging of obesity and weight loss interventions. *Molecular psychiatry*, *28*(4), 1466-1479.

Lindskog, M., Kim, M., Wikström, M. A., Blackwell, K. T., & Kotaleski, J. H. (2006). Transient calcium and dopamine increase PKA activity and DARPP-32 phosphorylation. *PLoS computational biology*, *2*(9), e119.

Lv, H., Zeng, N., Li, M., Sun, J., Wu, N., Xu, M., & Wang, Z. (2024). Association between body mass index and brain health in adults: a 16-year population-based cohort and mendelian randomization study. *Health Data Science*, *4*, 0087.

Ma, H., Qiu, R., Zhang, W., Chen, X., Zhang, L., & Wang, M. (2022). Association of PPP1R1B polymorphisms with working memory in healthy Han Chinese adults. *Frontiers in Neuroscience*, *16*, 989046.

Mathar, D., Neumann, J., Villringer, A., & Horstmann, A. (2017). Failing to learn from negative prediction errors: Obesity is associated with alterations in a fundamental neural learning mechanism. *Cortex*, *95*, 222-237

Meyer-Lindenberg, A., Straub, R. E., Lipska, B. K., Verchinski, B. A., Goldberg, T., Callicott, J. H., & Weinberger, D. R. (2007). Genetic evidence implicating DARPP-32 in human frontostriatal structure, function, and cognition. *The Journal of clinical investigation*, *117*(3), 672-682

Mier, D., Kirsch, P., & Meyer-Lindenberg, A. (2010). Neural substrates of pleiotropic action of genetic variation in COMT: a meta-analysis. *Molecular psychiatry*, *15*(9), 918-927.

Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. Annual review of neuroscience, 24(1), 167-202.

Morys, F., Tremblay, C., Rahayel, S., Hansen, J. Y., Dai, A., Misic, B., & Dagher, A. (2024). Neural correlates of obesity across the lifespan. *Communications Biology*, *7*(1), 656.

Moustafa, A. A., Sherman, S. J., & Frank, M. J. (2008). A dopaminergic basis for working memory, learning and attentional shifting in Parkinsonism. *Neuropsychologia*, *46*(13), 3144-3156.

Munafo, M. R., Matheson, I. J., & Flint, J. (2007). Association of the DRD2 gene Taq1A polymorphism and alcoholism: a meta-analysis of case–control studies and evidence of publication bias. Molecular psychiatry, 12(5), 454-461

Müller, D. J., Zai, C. C., Sicard, M., Remington, E., Souza, R. P., Tiwari, A. K., & Kennedy, J. L. (2012). Systematic analysis of dopamine receptor genes (DRD1–DRD5) in antipsychotic-induced weight gain. *The pharmacogenomics journal*, *12*(2), 156-164.

Narayanaswami, V., Thompson, A. C., Cassis, L. A., Bardo, M. T., & Dwoskin, L. P. (2013). Diet-induced obesity: dopamine transporter function, impulsivity and motivation. *International journal of obesity*, *37*(8), 1095-1103.

Nolan, K. A., Bilder, R. M., Lachman, H. M., & Volavka, J. (2004). Catechol O-methyltransferase Val158Met polymorphism in schizophrenia: differential effects of Val and Met alleles on cognitive stability and flexibility. *American Journal of Psychiatry*, *161*(2), 359-361

O'Reilly, R. C., & Frank, M. J. (2006). Making working memory work: a computational model of learning in the prefrontal cortex and basal ganglia. Neural computation, 18(2), 283-328

Pak, K., Seok, J. W., Lee, M. J., Kim, K., & Kim, I. J. (2023). Dopamine receptor and dopamine transporter in obesity: A meta-analysis. *Synapse*, *77*(1), e22254.

Persson, J., & Stenfors, C. (2018). Superior cognitive goal maintenance in carriers of genetic markers linked to reduced striatal D2 receptor density (C957T and DRD2/ANKK1-TaqIA). PLoS One, 13(8), e0201837.

Podell, J. E., Sambataro, F., Murty, V. P., Emery, M. R., Tong, Y., Das, S., & Mattay, V. S. (2012). Neurophysiological correlates of age-related changes in working memory updating. Neuroimage, 62(3), 2151-2160.

Pohjalainen, T., Rinne, J. O., Någren, K., Syv¨ alahti, E., & Hietala, J. (1998). Sex differences in the striatal dopamine D2receptor binding characteristics in vivo. American Journal of Psychiatry, 155(6), 768–773. https://doi.org/10.1176/ ajp.155.6.768

Ouimet, C. C., Miller, P. E., Hemmings, H. C., Walaas, S. I., & Greengard, P. (1984). DARPP-32, a dopamine-and adenosine 3': 5'-monophosphate-regulated phosphoprotein enriched in dopamine-innervated brain regions. III. Immunocytochemical localization. *Journal of Neuroscience*, *4*(1), 111-124

Rac-Lubashevsky, R., & Frank, M. J. (2021). Analogous computations in working memory input, output and motor gating: Electrophysiological and computational modeling evidence. PLoS computational biology, 17(6), e1008971.

Ranganath, A., & Jacob, S. N. (2016). Doping the mind: dopaminergic modulation of prefrontal cortical cognition. The Neuroscientist, 22(6), 593-603

Ribeiro, G., Maia, A., Cotovio, G., Oliveira, F. P., Costa, D. C., & Oliveira-Maia, A. J. (2023). Striatal dopamine D2-like receptors availability in obesity and its modulation by bariatric surgery: A systematic review and meta-analysis. *Scientific Reports*, *13*(1), 4959.

Rosa, E. C., Dickinson, D., Apud, J., Weinberger, D. R., & Elvevåg, B. (2010). COMT Val158Met polymorphism, cognitive stability and cognitive flexibility: an experimental examination. *Behavioral and brain functions*, *6*(1), 1-6.

Schmidt, K.-H., & Metzler, P. (1992). Wortschatztest: WST. Beltz.

Sesack, S. R., Hawrylak, V. A., Matus, C., Guido, M. A., & Levey, A. I. (1998). Dopamine axon varicosities in the prelimbic division of the rat prefrontal cortex exhibit sparse immunoreactivity for the dopamine transporter. *Journal of Neuroscience*, *18*(7), 2697-2708.

Sharma, S., & Fulton, S. (2013). Diet-induced obesity promotes depressive-like behaviour that is associated with neural adaptations in brain reward circuitry. *International journal of obesity*, *37*(3), 382-389.

Small, D. M., Jones-Gotman, M., & Dagher, A. (2003). Feeding-induced dopamine release in dorsal striatum correlates with meal pleasantness ratings in healthy human volunteers. *Neuroimage*, *19*(4), 1709-1715.

Smith, L., Watson, M., Gates, S., Ball, D., & Foxcroft, D. (2008). Meta-analysis of the association of the Taq1A polymorphism with the risk of alcohol dependency: a HuGE gene-disease association review. American journal of epidemiology, 167(2), 125-138

Smith, C. T., Swift-Scanlan, T., & Boettiger, C. A. (2014). Genetic polymorphisms regulating dopamine signaling in the frontal cortex interact to affect target detection under high working memory load. *Journal of cognitive neuroscience*, *26*(2), 395-407.

Stelzel, C., Basten, U., Montag, C., Reuter, M., & Fiebach, C. J. (2009). Effects of dopamine-related gene–gene interactions on working memory component processes. *European Journal of Neuroscience, 29*(5), 1056-1063

Stelzel, C., Basten, U., Montag, C., Reuter, M., & Fiebach, C. J. (2010). Frontostriatal involvement in task switching depends on genetic differences in D2 receptor density. Journal of Neuroscience, 30(42), 14205-14212.

Stice, E., Spoor, S., Bohon, C., Veldhuizen, M. G., & Small, D. M. (2008). Relation of reward from food intake and anticipated food intake to obesity: a functional magnetic resonance imaging study. *Journal of abnormal psychology*, *117*(4), 924.

Stone, M. (1979). Comments on model selection criteria of Akaike and Schwarz. Journal of the Royal Statistical Society. Series B (Methodological), 276-278

Thompson, J., Thomas, N., Singleton, A., Piggot, M., Lloyd, S., Perry, E. K., & Ferrier, I. N. (1997). D2 dopamine receptor gene (DRD2) TaqI A polymorphism: reduced dopamine D2 receptor binding in the human striatum associated with the A1 allele. *Pharmacogenetics and Genomics*, *7*(6), 479-484

Tunbridge, E. M., Bannerman, D. M., Sharp, T., & Harrison, P. (2004). Catechol-o-methyltransferase inhibition improves set-shifting performance and elevates stimulated dopamine release in the rat prefrontal cortex. *Journal of Neuroscience*, *24*(23), 5331-5335

van Schouwenburg, M., Aarts, E., & Cools, R. (2010). Dopaminergic modulation of cognitive control: distinct roles for the prefrontal cortex and the basal ganglia. Current pharmaceutical design, 16(18), 2026-2032.

Verdejo-Román, J., Fornito, A., Soriano-Mas, C., Vilar-López, R., & Verdejo-García, A. (2017a). Independent functional connectivity networks underpin food and monetary reward sensitivity in excess weight. *Neuroimage*, *146*, 293-300.

Verdejo-Román, J., Vilar-López, R., Navas, J. F., Soriano-Mas, C., & Verdejo-García, A. (2017b). Brain reward system's alterations in response to food and monetary stimuli in overweight and obese individuals. *Human Brain Mapping*, *38*(2), 666-677.

Volkow, N. D., & Wise, R. A. (2005). How can drug addiction help us understand obesity? *Nature neuroscience*, *8*(5), 555-560.

Volkow, N. D., Wang, G. J., Fowler, J. S., & Telang, F. (2008). Overlapping neuronal circuits in addiction and obesity: evidence of systems pathology. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *363*(1507), 3191-3200.

Volkow, N. D., Wang, G. J., Fowler, J. S., Tomasi, D., & Baler, R. (2012). Food and drug reward: overlapping circuits in human obesity and addiction. Brain imaging in behavioral neuroscience, 1-24.

Wacker, J. (2011). With great power comes great responsibility: Meta-analytic proof for a timeless insight. *Biological Psychiatry*, *69*(10), e37

Wang, G. J., Volkow, N. D., Logan, J., Pappas, N. R., Wong, C. T., Zhu, W., & Fowler, J. S. (2001). Brain dopamine and obesity. The Lancet, 357(9253), 354-357.

Wechsler, D. (2008). Wechsler adult intelligence scale–Fourth Edition (WAIS–IV). *San Antonio, TX: NCS Pearson*, *22*(498), 1.

Westbrook, A., van den Bosch, R., Hofmans, L., Papadopetraki, D., Maatta, J. I., Collins, A. G. E., & Cools, R. (2024). Striatal Dopamine Can Enhance Learning, Both Fast and Slow, and Also Make it Cheaper. *bioRxiv*, 2024-02.

Weydmann, G., Miguel, P. M., Hakim, N., Dube, L., Silveira, P. P., & Bizarro, L. (2023). How are overweight and obesity associated with reinforcement learning deficits? A systematic review. *Appetite*, 107123.

Wittchen, H.-U. Strukturiertes klinisches Interview für DSM-IV: SKID. Achse I: Psychische Störungen: Interviewheft und Beurteilungsheft; eine deutschsprachige, erweiterte Bearbeitung der amerikanischen Originalversion des SCID-I. (Hogrefe, Verlag für Psychologie, 1997).

Xu, H., Kellendonk, C. B., Simpson, E. H., Keilp, J. G., Bruder, G. E., Polan, H. J., & Gilliam, T. C. (2007). DRD2 C957T polymorphism interacts with the COMT Val158Met polymorphism in human working memory ability. Schizophrenia research, 90(1-3), 104-107.

Xu, X., Deng, Z. Y., Huang, Q., Zhang, W. X., Qi, C. Z., & Huang, J. A. (2017). Prefrontal cortex-mediated executive function as assessed by Stroop task performance associates with weight loss among overweight and obese adolescents and young adults. *Behavioural brain research*, *321*, 240-248.

Yang, Y., Shields, G. S., Guo, C., & Liu, Y. (2018). Executive function performance in obesity and overweight individuals: A meta-analysis and review. Neuroscience & Biobehavioral Reviews, 84, 225-244

Yang, Y., Shields, G. S., Wu, Q., Liu, Y., & Guo, C. (2019). Obesity is associated with poor working memory in women, not men: Findings from a nationally representative dataset of US adults. *Eating behaviors*, *35*, 101338.

Yang, Y., Shields, G. S., Wu, Q., Liu, Y., Chen, H., & Guo, C. (2020). The association between obesity and lower working memory is mediated by inflammation: Findings from a nationally representative dataset of US adults. *Brain, Behavior, and Immunity*, *84*, 173-179.

Zmigrod, L., & Robbins, T. W. (2021). Dopamine, Cognitive Flexibility, and IQ: Epistatic Catechol-O-MethylTransferase: DRD2 Gene–Gene Interactions Modulate Mental Rigidity. Journal of Cognitive Neuroscience, 34(1), 153-179.

FIGURES AND TABLES

Figure 1. Schematic illustration of the task structure and experimental conditions. The task consists of three task phases. In the encoding phase, participants have to remember two target stimuli (signaled by the letter "T"), or are presented with a centered cross (short control trials). In the interference phase, participants either have to ignore two non-target stimuli (ignore trials; signaled by the letter "N") or allow two new stimuli (again marked by a "T") to replace the previously remembered target stimuli (update trials). No-interference trials (short and long control) do not require any manipulations in the interference phase. At the end of each trial, participants evaluate whether a presented figure was a target figure or not. Figure reused from Hartmann et al. (2023) with permission.

Table 1. Sample Characteristics

project	all			BEDOB			GREADT			WORMCRI		
N (male)	320 (166)			156 (43)			86 (86)			78 (37)		
	mean (sd)	min	max	mean (sd)	min	max	mean (sd)	min	max	mean (sd)	min	max
BMI	26.38 (6.35)	17.51	45.54	29.172 (7.695)	17.51	45.54	24.025 (2.799)	18.632	36.419	23.217 (2.735)	18.929	29.888
IQ	105.41 (10.61)	71	122	101.575 (11.979)	71	122	109.151 (7.249)	91	118	107.731 (10.416)	74	118
Age	26.93 (6.82)	12.17	49.75	26.879 (8.907)	12.167	49.75	26.756 (4.474)	18	40	26.799 (3.859)	20.106	36.290
DFS	54.89 (11.61)	33	97	55.839 (10.163)	35	91	57.046 (15.107)	33	97	50.584 (8.546)	34	71

Figure 2. Main effect of BMI on working memory performance (model 1). Increasing BMI was associated with worse performance ($p_{corrected} < 0.001$, OR = 0.84). This trend was similar for all four conditions, as there was no interaction between BMI and condition ($p_{corrected} > 1$). Shaded areas represent the 95% confidence intervals.

Table 2. Full output model 1 with uncorrected p-values.

	Chisq	Df	Pr(>Chisq)
(Intercept)	3623.78	1	<0.001
condition	282.00	3	< .001
zBMI	16.80	1	< .001
zIQ	25.10	1	< .001
Gender	10.50	1	0.001
zWM_tired	36.00	1	< .001
zWM_conc	9.19	1	0.002
condition:zBMI	2.66	3	0.447

N = 320

Marginal R^2 / Conditional R^2 = 0.069 / 0.172

Figure 3. Interaction of Taq1A genotype, BMI, and condition on working memory performance (model 2). The two-way interaction of Taq1A and BMI was significant in the update condition only (p = 0.002). In this condition, carrying the A-allele led to a decrease in performance with each increasing unit of BMI (33.9%; SE = 7.58), while there was only -1.22% (SE = 7.21%) change in non-carriers. Shaded areas represent the 95% confidence intervals.

Table 3. Full output model 2 with uncorrected p-values.

	Chisq	Df	Pr(>Chisq)
(Intercept)	3228.77	1	<0.001
condition	212.81	3	< .001
COMT	0.16	2	0.923
Taq1A	1.13	1	0.288
zBMI	22.15	1	< .001
zIQ	24.09	1	< .001
Gender	7.53	1	0.006
zWM_tired	12.39	1	< .001
zWM_conc	30.80	1	< .001
condition:COMT	10.30	6	0.113
condition:Taq1A	4.69	3	0.196
COMT:Taq1A	2.47	2	0.291
condition:zBMI	3.49	3	0.322
COMT:zBMI	0.86	2	0.650
Taq1A:zBMI	2.98	1	0.085
condition:COMT:Taq1A	2.09	6	0.911
condition:COMT:zBMI	6.29	6	0.391
condition:Taq1A:zBMI	12.40	3	0.006
COMT:Taq1A:zBMI	3.68	2	0.159
condition:COMT:Taq1A:zBMI	4.09	6	0.665

N = 318

Marginal R^2 / Conditional R^2 = 0.076 / 0.173

Figure 4. Interaction of Taq1A genotype, BMI, and probe type in updating trials only. There was no significant three-way interaction between probe type, BMI and Taq1A (p = 0.439). The BMI - Taq1A interaction was in a similar direction in all trials. There was a significant main effect of probe type (p < 0.001). Trials where the probe was a distractor had lowest probability to be correct. Shaded areas represent the 95% confidence intervals.

Figure 5. Interaction of DARPP-32, BMI, and condition on working memory performance (model 3). The two-way interaction of DARPP-32 and BMI was significant in the update condition only (p = 0.006). In this condition, carrying the G-allele led to a decrease in performance with each increasing unit of BMI (32.4%; SE = 7.86), while there was only -4.39% (SE = 6.61) change in A/A homozygots. Shaded areas represent the 95% confidence intervals.

Table 4. Full output final model 3 with uncorrected p-values.

	Chisq	Df	Pr(>Chisq)
(Intercept)	3511.81	1	< .001
DARPP	0.03	1	0.853
zBMI	17.18	1	< .001
condition	274.62	3	< .001
zIQ	25.10	1	< .001
zWM_conc	35.27	1	< .001
zWM_tired	10.52	1	0.001
Gender	9.17	1	0.002
DARPP:zBMI	0.18	1	0.668
DARPP:condition	1.00	3	0.801
BMI:condition	3.61	3	0.307
DARPP:BMI:condition	20.21	3	< .001

N = 320

Marginal R^2 / Conditional R^2 = 0.071 / 0.173

Figure 6. Interaction of DARPP-32 genotype, BMI, and probe type for update trials only. There was a significant three-way interaction between probe type, BMI and DARPP-32 (p = 0.005). Post hoc analyses showed that the BMI - DARPP interaction was significant in distractor (p = 0.046) and target (p = 0.008) trials, but not in trials where the probe was a novel item (p = 0.242). There was a significant main effect of probe type (p < 0.001). Trials where the probe was a distractor had lowest probability to be correct. Shaded areas represent the 95% confidence intervals.

Table 5. Full output final model 4 with uncorrected p-values.

	Chisq	Df	Pr(>Chisq)
(Intercept)	328.55	1	< .001
C957T	0.03	1	0.859
zBMI	3.49	1	0.062
condition	48.06	3	< .001
zIQ	25.30	1	< .001
zWM_conc	33.66	1	< .001
zWM_tired	10.54	1	0.001
Gender	8.85	1	0.003
C957T:zBMI	0.36	1	0.548
C957T:condition	7.97	3	0.047
BMI:condition	0.31	3	0.958
C957T:BMI:condition	0.07	3	0.995

N = 318

Marginal R^2 / Conditional R^2 = 0.070 / 0.171

Figure 7. Interaction of Amino Acid Ratio, BMI and condition (model 5). For illustration purposes, amino acid ratio was artificially grouped into high, middle, and low. The difference in condition ($p_{ignore\ vs.\ update}$ < 0.001) becomes especially apparent when looking at individuals with high amino acid ratios: With each increasing unit of BMI, performance gets worse in ignore, but better in update. There were no significant differences in the relationship of amino acid ratio and BMI comparing all other conditions against each other (all p > 0.168). Shaded areas represent the 95% confidence intervals.

Table 6. Full output final model 5 with uncorrected p-values.

	Chisq	Df	Pr(>Chisq)
(Intercept)	13.31	1	< .001
AAratio	0.58	1	0.444
zBMI	0.98	1	0.321
condition	3.43	3	0.330
zIQ	11.64	1	< .001
zWM_conc	18.60	1	< .001
Gender	5.08	1	0.024
AAratio:zBMI	0.32	1	0.570
AAratio:condition	8.69	3	0.034
BMI:condition	9.80	3	0.020
AAratio:BMI:condition	10.88	3	0.012

N = 160

Marginal R^2 / Conditional R^2 = 0.068 / 0.170