Tonically active GABAergic neurons in the dorsal periaqueductal gray control the initiation and execution of instinctive escape

A. Vanessa Stempel[* 1,2 ✉]**, Dominic A. Evans**[* 1,2]**, Oriol Pavón Arocas**[1]**, Federico Claudi**[1]**, Stephen C. Lenzi**[1]**, Elena Kutsarova**[2]**, Troy W. Margrie**[1]**, and Tiago Branco**[1 ✉]

[1]UCL Sainsbury Wellcome Centre for Neural Circuits and Behaviour, London, United Kingdom
[2]Max Planck Institute for Brain Research, Frankfurt am Main, Germany
[*]equal contribution
✉correspondence to vanessa.stempel@brain.mpg.de, t.branco@ucl.ac.uk

To avoid predation, animals perform defensive actions that are both instinctive and adaptable to the environment. In mice, the decision to escape from imminent threats is implemented by a feed-forward circuit in the midbrain, where excitatory VGluT2[+] neurons in the dorsal periaqueductal gray (dPAG) compute escape initiation and escape vigour from threat evidence. Here we show that GABAergic VGAT[+] neurons in the dPAG dynamically control this process by modulating the excitability of excitatory escape neurons. Using *in vitro* patch clamp and *in vivo* neural activity recordings in freely behaving mice we found that VGAT[+] dPAG neurons fire action potentials tonically in the absence of synaptic inputs and are a major source of synaptic inhibition to VGluT2[+] dPAG neurons. Activity in these spontaneously firing VGAT[+] cells transiently decreases at escape onset and increases during escape, peaking at escape termination. Optogenetically increasing or decreasing VGAT[+] dPAG activity bidirectionally changes the probability of escape when the stimulation is delivered at the time of threat onset, and the duration of escape when delivered after escape initiation. We conclude that the activity of tonically firing VGAT[+] dPAG neurons sets a threshold for escape initiation and controls the execution of the flight locomotor action.

Introduction

Escape behaviour comprises of a set of locomotor actions that move an animal away from threat. While these actions can be reflexive and stereotyped to ensure short reaction times and fast movements, it is equally important for survival that they are flexible (1–3). For example, the probability of initiating escape depends on predation risk and threat history (4–11), as well as on the internal state (12–16). In environments where predators abound, it is more likely that ambiguous sensory stimuli might signal the presence of predators and thus the probability of initiating escape should increase (17). Likewise, when faced with a competing motivation, such as hunger, the probability of defensive behaviour decreases (12, 18).

Beyond flexibility in escape initiation, it is equally important that the execution of escape is flexible. The flight to safety should not be ballistic (i.e. along a pre-determined trajectory with fixed kinetics) but instead should allow for continuous adjustments of vigour and trajectory to deal with variability in the terrain, adjust escape to unexpected events and terminate escape at the appropriate place and time (19–22). Accordingly, recent work has shown that mice on a fast shelter-directed escape trajectory can slow down and navigate around an obstacle that suddenly appears, while maintaining the goal of reaching safety (20). This degree of flexibility in escape initiation and execution suggests the presence of modulatory components acting on the neural circuits controlling instinctive escape. In particular, the ability of decreasing the sensitivity to threat and slowing down locomotion during escape suggests the presence of a regulatory inhibitory network.

In mammals, the key circuit for commanding escape behaviour is the periaqueductal gray (PAG), located in the midbrain (23–25). The PAG is an interface between the forebrain and central pattern generators in the hindbrain and spinal cord that acts as a critical neural circuit node for integrating sensory and state-dependent information, as well as for executing instinctive behaviours (25–39). In the dorsal PAG (dPAG), excitatory VGluT2[+] neurons receive information about imminent visual and auditory threats from the superior colliculus through a feed-forward excitatory connection, and their activation determines the onset of escape and controls escape vigour (31). While modulation of threat processing and escape behaviour can be accomplished through a variety of different circuits (40–46), the network of excitatory neurons in the dPAG stands out as a key control node (24, 29, 31, 47–50). Previous work has demonstrated the presence of an inhibitory tone in the PAG that could in principle control the excitability of the excitatory network to modulate escape. For example, local application of a GABA$_A$ receptor antagonist in the PAG elicits responses to a previously sub-threshold stimulus (23, 24). Disinhibition of the PAG has been suggested as a mechanism for initiating other instinctive behaviours such as freezing (51), and single-unit recordings from GABAergic neurons in the lateral and ventral PAG have shown that their baseline firing rates are higher than the excitatory neurons (35, 51). Here we have tested the hypothesis that local GABAergic neurons within the dPAG control escape behaviour by setting the excitability of the dPAG escape network.

Results

GABAergic neurons in dPAG fire tonically in the absence of synaptic input. To investigate how inhibitory neurons in the dPAG might control the excitability of the local excitatory neurons, we performed patch clamp recordings from identified VGAT[+] and VGluT2[+] neurons in acute midbrain

Fig. 1. Properties of GABAergic inhibition in the dorsal PAG. A. Schematic of a patch clamp recording from visually identified VGAT⁺ neurons in the dPAG. **B.** Loose-seal cell-attached recordings of VGAT⁺ dPAG neurons. Example current traces of spontaneously spiking VGAT⁺ neurons without (control, top) and with synaptic blockers (2 mM kynurenic acid and 50 μM picrotoxin, bottom) added to the bath. **C.** Summary of VGAT⁺ dPAG neuron firing rate for control (n = 39 cells, N = 20 mice) and in synaptic blockers (n = 45 cells, N = 13 mice. **D** and **E.** Same as A, for VGluT2⁺ dPAG neurons in control conditions. **F.** Summary of VGluT2⁺ dPAG neuron firing rate for control (n = 42 cells, N = 13 mice) and in 50 μM picrotoxin (n = 30 cells, N = 9). Comparison between VGAT⁺ and VGluT2⁺ controls: p<0.0001, Mann-Whitney test. **G.** Example whole-cell patch clamp recording of spontaneous IPSCs onto a VGluT2⁺ dPAG neuron in the presence of 2 mM kynurenic acid at holding potential of -70mV. **H.** Left panel: summary graph of the mean spontaneous IPSC frequency recorded in VGluT2⁺ dPAG neurons from n = 22, N = 8 mice. Right panel: summary graph of the normalised mean spontaneous IPSC frequency after bath application of SR-95331 (n = 5, N = 4 mice) and time-matched controls (n = 9, N = 7 mice). **I.** Average spontaneous IPSC from the same neuron as in G. **J.** Summary graphs of the mean peak phasic conductance (left panel), unitary IPSC charge (middle panel) and average IPSC charge (right panel) from n = 11 cells, N = 2 mice. **K.** Left: Schematic of the experiment for panels K–N. AAV-DIO-ChR2 and AAV-DIO-hM4D were co-expressed in the dPAG of VGAT-Cre animals and patch clamp recordings were performed from presumptive, non-spiking excitatory neurons. Right: Connectivity rate between ChR2-expressing VGAT⁺ neurons and presumptive excitatory dPAG neurons (n = 17 / 20 cells; N = 2 mice) with an example trace of inhibitory currents recorded in voltage-clamp configuration upon blue light stimulation (5 pulses, 1 ms duration, 20 Hz inter-stimulus interval). **L.** Summary graphs of the evoked IPSC amplitude (left panel) and synaptic delay of this connection (right panel) (n = 16 cells, N = 2 mice). **M.** Left: Same experiment as shown in K, with bath application of CNO. Right: Example traces of spontaneous IPSCs recorded in 2 mM kynurenic acid before (blue trace) and after (orange trace) bath application of 10 μM CNO. **N.** Summary graph of the normalised change of spontaneous IPSC frequency in control conditions vs. CNO (n = 6 cells each, N = 2 mice). Box-and-whisker plots show median, IQR and range, as well as mean (+) and individual data points.

slices *in vitro*. While whole-cell recordings of VGAT⁺ dPAG neurons showed biophysical and synaptic input properties similar to those of VGluT2⁺ dPAG cells (Figure S1, for comparison to VGluT2⁺ dPAG neurons, see (31)), recordings in loose-seal cell-attached configuration to monitor spiking non-invasively showed a marked difference between these two cell populations. VGAT⁺ neurons fired action potentials spontaneously at a mean firing rate of 4.7 ± 0.70 Hz, which was not significantly altered after pharmacological block of glutamatergic and inhibitory synaptic transmission (mean firing rate = 6.2 ± 0.84 Hz; P = 0.21, Mann-Whitney test between control and blockers; Figure 1A, B; see Figure S2 for anatomical locations). Using inter-spike interval (ISI) analysis and and a measure of variability based on the coefficient of variation (CV_2) (52), we found the firing of VGAT⁺ dPAG neurons to be highly regular, as expected from a dominant cell intrinsic mechanism (CV_2 = 0.33 ± 0.03; Poisson firing statistics have a CV_2 of 1 and a perfectly regular neuron has a CV_2 of 0; Figure S3A, B). In contrast, the majority of VGluT2⁺ dPAG neurons were not spontaneously active both in control conditions (92.8 %, 39 out of 42 cells, firing rate < 0.04 Hz; mean firing rate across all neurons = 0.11 ± 0.068 Hz) or after block of inhibitory synaptic transmission (mean firing rate in picrotoxin = 0.039

± 0.033 Hz; P = 0.85, Mann-Whitney tests between control and picrotoxin; Figure 1D–F). These data show that VGAT⁺ dPAG neurons are well positioned to provide sustained inhibition to the local network through input-independent action potential firing.

VGluT2⁺ neurons in dPAG are phasically inhibited by local GABAergic neurons. To assess the impact of local tonically active GABAergic neurons on dPAG excitability, we performed whole-cell recordings from VGluT2⁺ dPAG neurons *in vitro* and first isolated spontaneous inhibitory post-synaptic currents (IPSCs) by blocking glutamatergic synaptic transmission. We found frequent phasic inhibition (mean IPSC frequency = 6.5 ± 0.83 Hz) that was selectively blocked by a GABAₐ receptor antagonist (SR-95531: mean fraction = 0.13 ± 0.05; control: mean fraction = 1.17 ± 0.3, P = 0.002, Mann-Whitney test, Figure 1G, H). Phasic inhibitory currents had a peak conductance of 0.94 ± 0.16 nS and a unitary IPSC charge of 0.28 ± 0.04 pC, generating a continuous average charge over time of 2.12 ± 0.68 pC (Figure 1I, J). These data show that VGluT2⁺ dPAG neurons receive sustained phasic GABAₐ receptor-mediated inhibition with an inhibitory charge transfer that should in principle be large enough to regulate their input-output function *in vivo* (as e.g.

shown for cerebellar granule cells that have a similarly high input resistance (53).

Next, we tested the contribution of local VGAT⁺ dPAG neurons to inhibition of VGluT2⁺ dPAG cells. We used a combined opto- and chemogenetic strategy to co-express an inhibitory DREADD (designer receptor exclusively activated by designer drugs, hM4Di-mCherry) and channelrhodopsin-2 (ChR2-YFP) in the dPAG of VGAT-Cre mice to selectively and bidirectionally change the activity of VGAT ⁺ dPAG neurons while recording inhibitory currents in excitatory dPAG cells (see Methods). We first optogenetically activated VGAT⁺ dPAG neurons and found monosynaptic inhibitory inputs in 85 % of glutamatergic dPAG neurons (mean latency = 2.0 ± 0.15 ms), with a mean evoked IPSC amplitude of -62.58 ± 18.31 pA and a facilitating paired pulse ratio (1.17 ± 0.16; Figure 1K, L). We then chemogenetically blocked synaptic transmission from VGAT⁺ dPAG neurons with Clozapine-N-Oxide (CNO, 10 μM), which reduced the frequency of spontaneous IPSCs onto glutamatergic neurons by 70 % in comparison to control recordings (normalised change in spontaneous IPSC frequency for CNO: mean = 0.59 ± 0.14; control: mean = 1.29 ± 0.17; P = 0.01, unpaired t-test with Welch's correction; Figures 1M, N and S3A). Application of CNO also significantly reduced the amplitude of ChR2-evoked IPSCs (normalised peak amplitude CNO = 0.39 ± 0.16; control = 1.35 ± 0.33; CNO vs. control: P = 0.042, unpaired t-test; Figure S3C, D) further confirming the efficacy and specificity of our chemogenetic inhibition approach. It is important to note that since these experiments rely on small targeted injections and co-expression of two viral constructs within the same cells, our quantification of the contribution of local VGAT⁺ dPAG to spontaneous inhibition is likely to be an underestimate. Together these data show that VGAT⁺ neurons in the dPAG provide phasic and sustained monosynaptic GABA$_A$ receptor-mediated inhibition onto VGluT2⁺ dPAG cells.

VGAT⁺ dPAG neurons are spontaneously active in vivo and modulated during escape behaviour. We next measured the activity of VGAT⁺ dPAG neurons *in vivo* to determine whether they are a source of sustained inhibition as suggested by our *in vitro* experiments, and how their activity relates to escape behaviour. Given the characteristic firing profile of VGAT⁺ dPAG neurons *in vitro*, we first performed chronic silicon probe recordings from the dPAG of freely moving mice to determine whether there are single units with similar activity. Analysis of firing rates for periods when mice were spontaneously exploring the behavioural arena (see Methods) revealed the existence of neurons with high baseline firing rates across the recorded population (9/19 neurons with mean firing rates above 8 Hz; mean frequency = 32.2 ± 7.0 Hz; N = 3 mice; Figure S3E–G). This is consistent with previous reports of high baseline firing rates of VGAT⁺ neurons *in vivo* in other subdivisions of the PAG (35, 51), and with synaptic input driving neurons to higher firing rates than *in vitro*.

To directly record neural activity from molecularly-identified VGAT⁺ neurons, we performed *in vivo* calcium

Fig. 2. Calcium activity profiles of VGAT⁺ dPAG neurons during exploration and instinctive escape. A. Video frame during the behavioural assay of a mouse escaping to shelter (coloured in blue) from the designated threat zone opposite the shelter after stimulation with an innately threatening visual or auditory stimulus (here: a dark expanding disk). **B.** Top left: Example FOV in the dPAG of a VGAT-Cre mouse injected with AAV-flex-GCaMP6s. Bottom left: Overlay of the same FOV with the extracted and numbered cell masks. Right: Example traces of spontaneous calcium activity (Z-scored) of the indicated cell masks during baseline exploration. The speed of the animal is indicated in yellow. **C.** Summary of the mean spontaneous event frequency per animal. **D.** Fraction of cells active during escape. **E.** Example escape trials of one animal during one session. Coordinates of the mouse's location are shown over time and normalised to the distance from shelter (shelter entrance = 0, arena end = 1, shelter location indicated in blue). Time points of behavioural tags (dots) after stimulus onset (orange line) are shown for: startle (light orange), escape turn (green), escape run (blue) and escape stop (grey). Note that the trial with the long-latency escape onset corresponds to trial 7 in panel F. **F.** Heatmap of the Z-scored calcium signal of one example neuron during the escape trials shown in E, with the peak calcium signal aligned to stimulus-onset, escape turn and escape stop respectively. **G.** Heatmaps of the Z-scored calcium activity of all recorded neurons (mean calcium signal across trials), divided by cluster (cluster 1, top; cluster 2; bottom), time warped between behavioural tags and aligned by their peak calcium signal, N = 8 mice. **H.** Average signal of cluster 1 (orange line) and cluster 2 (purple line), together with the average of all neurons for trials where animals failed to escape after stimulus detection (grey dotted line). **I.** The distribution of the peak calcium signal of neurons in cluster one relative to escape termination (mean peak signal = +0.17 ± 0.12 s; white marker, not different from 0 s; P = 0.16, one-sample t-test). Box-and-whisker plots show median, IQR and range, as well as mean (+).

imaging of VGAT⁺ dPAG neurons expressing GCaMP6s in freely moving mice using a head-mounted miniature micro-

scope (Figure S4). In line with the *in vitro* cell-attached and *in vivo* single-unit recordings during exploration, we found that VGAT⁺ dPAG neurons were active while naïve animals freely explored the arena (median calcium event rate = 0.1 Hz, IQR = [0.090, 0.11 Hz], 209 neurons, N = 8 mice; Figures 2A, B and S5A). To test whether this activity varied with the locomotor state of the animal, we calculated a locomotion modulation index for each neuron (see Methods)(54), and found that only a small fraction of cells was significantly modulated by movement (4.5 % of cells positively, 4.5 % of cells negatively modulated; Figure S5B). Together with the previous results, these data consolidate the view that VGAT⁺ dPAG neurons provide a sustained inhibitory tone on the dPAG network.

To understand how local VGAT⁺ dPAG activity might be involved in threat-evoked escape, we presented threatening overhead looming or ultrasound stimuli in the presence of a shelter (55, 56). During stimulus-evoked escape to shelter, more than half of all active cells showed escape-related activity (median across mice = 54%, IQR = [46%, 76%], 123 neurons, N = 8 mice, Figures 2D–F, S5C–F). To compare the activity profiles of neurons across trials, where the escape sequence is stereotyped but has a variable duration, we linearly time warped each neuron's activity to the median time-series of discrete events in the escape sequence to generate an average over trials (see Methods). Across the recorded population, the neuronal activity profiles were best described by two clusters identified by K-means clustering using scores from principal component analysis on the pooled mean time-warped responses (Figures 2G, H and S5G; clustering on top three principal components explaining 86% of the variance), with neurons belonging to both clusters present in all animals (Figure S5H, I): the first cluster showed a gradual increase in activity from threat stimulus onset, peaking at escape termination (mean z-score over escape sequence = +0.95 ± 0.04 Z, n = 62 neurons; Figure 2I); whereas the second cluster exhibited a transient decrease in activity with the trough coinciding with escape onset (mean z-score over escape sequence -0.33 ± 0.05 Z, n = 61 neurons; 100 trials, N = 8 mice). The mean population activity was not modulated in trials where the threat stimulus failed to elicit escape (Figure 2H, grey dotted line; mean post-stimulation signal per neuron = 0.009 ± 0.06 Z, one-sample t-test against 0, P = 0.88, 66 trials, N = 7 mice), which is similar to glutamatergic dPAG neurons (31). To further test whether the activity of VGAT⁺ dPAG neurons was specifically modulated during escape, we compared activity profiles between fast spontaneous runs outside the threat stimulus context with slow escapes, where the average locomotion speed was lower than the spontaneous runs. We found that VGAT⁺ dPAG activity did not change during exploratory runs (*cluster 1, runs during exploration*, mean signal -0.06 ± 0.14 Z and -0.11 ± 0.15 Z in the 2 s before and after run onset respectively, two-tailed paired t-test, P = 0.80; n = 18 neurons, 66 trials, N = 4 mice) but changed significantly during slow escapes (*cluster 1, slow escapes*, mean signal -0.42 ± 0.22 Z before and 0.22 ± 0.21

Z after, two-tailed paired t-test, P = 0.047, n = 12 neurons, 11 trials, N = 4 mice; Figure S5J, K). In addition, cluster population activity was not correlated to the peak speed, mean acceleration or mean deceleration of escape movement (Figure S6). Together this indicates that in the behavioural context tested, changes in the activity of VGAT⁺ dPAG neurons are specific for the escape action rather than sensory stimuli or locomotion. These results suggest that VGAT⁺ dPAG neurons contribute to escape initiation by decreasing their activity in response to threat, thereby releasing the excitatory dPAG neurons from inhibition and facilitating their recruitment to initiate escape. In addition, the gradual increase in activity during escape indicates that VGAT⁺ dPAG neurons might also participate in shutting down the excitatory drive in the dPAG network to terminate the escape action.

Activity manipulations of VGAT⁺ neurons bidirectionally modulate escape initiation and termination. The neural activity profiles of VGAT⁺ dPAG neurons recorded *in vivo* predict that: (1) manipulating their activity at the time of threat presentation should bidirectionally affect escape initiation, and (2) modulating activity during the escape action should perturb the duration of the escape action. We tested these predictions using bilateral, optogenetic manipulations of VGAT⁺ dPAG neurons during instinctive escape (activation with ChR2 and inactivation with iChloC; Figure S7A–C). First, we tested the effect of activity manipulations on escape initiation. Optogenetic activation of VGAT⁺ dPAG neurons at the time of threat presentation strongly reduced escape probability (mean escape probability = 0.88 ± 0.04 for threat only vs. 0.13 ± 0.05 for threat with laser on, P = 0.0005, paired two-tailed t-test; Figure 3A, C), but had no effect when it preceded the threat stimulus (end of the laser pulse train coincident with threat stimulus onset; mean escape probability = 0.92 ± 0.08 for threat with laser preceding stimulus onset vs. threat only, P = 0.44, Mann Whitney test; Figure S7F). Conversely, optogenetic inhibition of VGAT⁺ neurons during presentation of a weakly-threatening, low-contrast stimulus increased escape probability significantly (mean escape probability = 0.28 ± 0.06 for threat only, 0.78 ± 0.1 for threat with laser on; P = 0.0014, paired two-tailed t-test; Figure 3B, D). These results confirm that activity of VGAT⁺ dPAG neurons plays an important role in setting the escape threshold.

We next tested the role of VGAT⁺ neurons in escape execution. Optogenetic stimulation of VGAT⁺ dPAG neurons with ChR2 after escape initiation often led to premature termination of escape, with the animal stopping before reaching the shelter (normalised mean escape termination distance from shelter, laser off = -0.10 ± 0.01, laser on = +0.09 ± 0.01; P < 0.001, paired two-tailed t-test; Figure 3E). Inactivation of VGAT⁺ neurons during escape had the opposite effect, with animals frequently overshooting their preferred target location of the arena shelter, resulting in a significant increase in total escape distance (laser off = 49.9 ± 4.6 cm, laser on = 69.4 ± 4.8 cm, P = 0.0097, unpaired t-test; Figure S8).

Fig. 3. VGAT⁺ dPAG neurons bidirectionally modulate escape. A. Left: example raster plots of mouse speed for the same animal during a high-threat stimulus (a sequence of three high-contrast overhead looming spots) with 8 escapes in 9 trials (control, top) and 0 escapes in 11 trials during simultaneous optogenetic activation of the dPAG VGAT⁺ neurons (ChR2, 50 Hz, 5 s; bottom). Coloured bar indicates escape (red) and no escape (grey) trials. Laser OFF and Laser ON trials were interleaved during the experiment, and trials are sorted by reaction time. Right: Single trial mouse tracking of the same trials showing successful escapes to shelter (red dots = escape termination) for control (top) and during ChR2 stimulation (blue, bottom). **B.** Same as A, for optogenetic inactivation with iChloC (3–5 s, continuous). Left: example raster plots for the same animal during a low-threat stimulus (a sequence of three low-contrast looming spots) with 8 escapes in 25 control trials (top) and 18 escapes out of 24 trials during optogenetic inhibition of VGAT⁺ neurons (bottom). Right: Single trial mouse tracking showing failed and successful escapes to shelter (red dots = escape termination) for control (top) and during iChloC stimulation (orange, bottom). **C.** Summary plot of escape probability in response to threat presentation with and without simultaneous optogenetic activation of VGAT⁺ dPAG neurons (n = 100 trials, N = 5 mice). **D.** Summary plot of escape probability in response to optogenetic inactivation and threat presentation (mean escape probability = 0.12 ± 0.09 Laser ON only, n = 108 trials, N = 4 mice). **E.** Left: Single mouse tracking showing control escape trials (Laser OFF) and trials with ChR2 stimulation *after* escape onset (red dot: escape termination; blue line: Laser ON). Right: Summary plots of the normalised mean escape termination distance from shelter during Laser OFF and Laser ON trials (N = 3 mice). Box-and-whisker plots show median, IQR and range, as well as mean (+).

Importantly, neither manipulation affected the mean speed during baseline exploratory locomotion (ChR2: P = 0.41, iChloc: P = 0.88; two-tailed paired t-tests; Figure S5D–H) suggesting a dedicated function of VGAT⁺ dPAG neurons in the execution of escape behaviour.

Discussion

We have described a locally connected population of GABAergic neurons in the dPAG that tonically fire action potentials and provide monosynaptic inhibition to dPAG excitatory neurons that is mediated by GABA_A receptors. In contrast with the sparse connectivity between VGluT2⁺ dPAG neurons (31), we find that these glutamatergic neurons receive input from local GABAergic dPAG neurons with a high connectivity rate, resulting in frequent inhibitory synaptic events that generate a sustained barrage of inhibition. The presence and role of a tonic inhibitory mechanism in the PAG was postulated decades ago (24) and recent work has described both long-range and local disinhibitory motifs to be a main element of instinctive behaviour initiation (28, 42, 57–

59). Our results add to the emerging view that sustained inhibition is a key property inherent to the functional architecture of the PAG and demonstrate its role in instinctive escape behaviour.

Our neural activity recordings from freely behaving mice show that VGAT⁺ dPAG are active as animals explore the environment, and that approximately half of the recorded population shows a transient decrease in activity before escape onset. We think that this could serve to disinhibit the VGluT2⁺ dPAG neurons that drive escape (31, 60) and facilitate escape initiation. This view is strongly supported by the optogenetic silencing experiments, in which inhibition of VGAT⁺ dPAG neurons increased escape probability. Together with the finding that optogenetic activation of the same neuron population can decrease escape probability, these results show that VGAT⁺ dPAG neurons are a key node for determining the escape initiation threshold, in addition to the synaptic properties between the threat input circuits and the dPAG (31). The existence of a sustained source of inhibition in the dPAG might be important for dampening the excitability of the ex-

citatory neurons and prevent initiation of escape behaviour in the absence of threat input, while also providing a means for fast, bidirectional modulation of escape. It will be interesting in the future to determine whether changes to the baseline firing of VGAT$^+$ dPAG neurons are a mechanism to support plastic changes to the escape threshold, such as during learned suppression of escape (4). It will also be important to establish whether the function of VGAT$^+$ dPAG neurons in escape is exclusively mediated by intra-PAG inhibition or whether they also directly modulate other targets, such as the cuneiform nucleus of the mesencephalic locomotor region which has been previously reported to play a role in defensive behaviours (61–63).

In addition to setting a threshold for initiating escape behaviour, our results suggest that VGAT$^+$ dPAG neurons are also important for the ongoing execution of escape, in particular for terminating the flight action. A large fraction of this population shows a gradual increase in activity between escape onset and escape termination at the shelter, and optogenetically increasing or decreasing their activity can cause animals to undershoot or overshoot their target, respectively. Given the known role of glutamatergic dPAG neurons in locomotor initiation and speed control during escape (31, 34, 48, 64), these results can be interpreted as VGAT$^+$ dPAG neurons acting to shut down firing in the excitatory neurons that drive escape. These results are in line with the previous finding that VGAT$^+$ neurons in the lateral PAG have a role in both the initiation and execution phases of hunting behaviour (35). An important unanswered question is whether there are distinct subsets of VGAT$^+$ dPAG neurons, one for controlling initiation and the other for execution. Our analysis identified two broad clusters based on the activity profile during escape, which may arise if there are groups of VGAT$^+$ neurons integrating different input pathways (35). It will be interesting in the future to determine if such input segregation exists. The observation that we were able to change both escape initiation and execution by manipulating activity of the entire population is consistent with the main output of the two clusters being glutamatergic dPAG neurons, with the timing of VGAT$^+$ neuron activation being the main determinant of the effect on escape behaviour.

While the activity of glutamatergic dPAG neurons is positively correlated to escape speed (31), we found that the population activity of GABAergic neurons is not significantly modulated by speed but instead peaks at escape termination. This observation supports a general framework for locomotor-correlated activity in brainstem regions, where glutamatergic neurons are the key population for regulating speed and are under the control of local inhibitory networks (65). The finding that GABAergic cells in the dPAG mediate the termination of escape behaviour further adds to the view that the dPAG is an integral part of the supraspinal circuits for the coordination of goal-directed locomotion (65–68).

Studies of escape behaviour in the wild have highlighted how flexible the flight action must be since animals often engage in a complex series of stop-and-go events while computing and executing escape trajectories directed to safe loca-

tions (1, 10, 19, 69–71). Our results raise the hypothesis that inhibitory neurons in the dPAG might integrate spatial and sensory information to implement flexible speed adjustments during escape and determine when safety has been reached through modulation of their firing rates (41, 72, 73).

Acknowledgements
This work was funded by a Wellcome Senior Research Fellowship (214352/Z/18/Z), by the Sainsbury Wellcome Centre Core Grant from the Gatsby Charitable Foundation and Wellcome (GAT3755 and 219627/Z/19/Z) and by a European Research Council grant (Consolidator no. 864912) (T.B.), German Research Foundation postdoctoral fellowships (E.K., project no. 515465001; A.V.S., project no. STE 2605/1), the UCL Wellcome 4-year PhD Programme in Neuroscience (O.P.A.), the SWC PhD Programme (F.C., S.C.L.) and the Max Planck Society (E.K., A.V.S.). We thank P. Iordanidou, T. Okbinoglu and M. Strom for technical assistance; the SWC Neurobiological Research Facility and MPI for Brain Research animal facility for animal care, FabLabs for technical support, and K. Betsios for programming the data acquisition software. We thank F. Rau, P. Zatka-Haas, A. Fratzl and all members of the Branco lab for discussions; D. Campagner, J. Kohl, S. Keshavarzi for comments on the manuscript. This manuscript was typset using a modified version of the HenriquesLab bioRxiv template.

Author Contributions
A.V.S. and T.B. conceived the project with critical input from D.A.E. A.V.S, O.P.A. and E.K. performed *in vitro* electrophysiology. A.V.S. and D.A.E. performed *in vivo* calcium imaging and optogenetic manipulation experiments, F.C. contributed to preprocessing and analysis of the calcium imaging dataset. S.C.L. performed *in vivo* single-unit recordings, supervised by T.W.M. A.V.S and T.B. supervised the project. A.V.S., T.B. and D.A.E. wrote the manuscript and all authors read and commented on the manuscript.

Competing Financial Interests
The authors declare there are no competing interests.

Bibliography

1. D. Evans, A. Stempel, R. Vale, and T. Branco. Cognitive Control of Escape Behaviour. *Trends in Cognitive Sciences*, 23(4):334–348, 2019.
2. D. H. Edwards, W. J. Heitler, and F. B. Krasne. Fifty years of a command neuron: The neurobiology of escape behavior in the crayfish. *Trends in Neurosciences*, 22(4):153–161, 1999.
3. M. Y. Peek and G. M. Card. Comparative approaches to escape. *Current Opinion in Neurobiology*, 41:167–173, 2016.
4. S. C. Lenzi, L. Cossell, B. Grainger, S. F. Olesen, T. Branco, and T. W. Margie. Threat history controls flexible escape behavior in mice. *Current Biology*, 32(13):2972–2979.e3, 2022.
5. S. L. Lima and L. M. Dill. Behavioral decisions made under the risk of predation: a review and prospectus. *Canadian Journal of Zoology*, 68(4):619–640, 1990.
6. M. C. Ferrari, J. J. Trowell, G. E. Brown, and D. P. Chivers. The role of learning in the development of threat-sensitive predator avoidance by fathead minnows. *Animal Behaviour*, 70(4):777–784, 2005.
7. A. Collier and J. Y. Hodgson. A Shift in Escape Strategy by Grasshopper Prey in Response to Repeated Pursuit. *Southeastern Naturalist*, 16(4):503–515, 2017.
8. W. E. Cooper and W. G. Frederick. Optimal flight initiation distance. *Journal of Theoretical Biology*, 244(1):59–67, 2007.
9. L. M. Dill. Distance-to-cover and the escape decisions of an African cichlid fish, Melanochromis chipokae. *Environmental Biology of Fishes*, 27(2):147–152, 1990.
10. S. Edut and D. Eilam. Rodents in open space adjust their behavioral response to the different risk levels during barn-owl attack. *BMC ecology*, 3:10, 2003.
11. W. Cresswell, S. J. Butler, M. J. Whittingham, and J. L. Quinn. Very short delays prior to escape from potential predators may function efficiently as adaptive risk-assessment periods. *Behaviour*, 146(6):795–813, 2009.
12. A. Filosa, A. J. Barker, M. Dal Maschio, and H. Baier. Feeding State Modulates Behavioral Choice and Processing of Prey Stimuli in the Zebrafish Tectum. *Neuron*, 90(3):596–608, 2016.
13. M. Thaker, A. Vanak, S. Lima, and D. Hews. Stress and Aversive Learning in a Wild Vertebrate: The Role of Corticosterone in Mediating Escape from a Novel Stressor. *The American Naturalist*, 175(1):50–60, 2010.
14. D. Hawlena, H. Kress, E. R. Dufresne, and O. J. Schmitz. Grasshoppers alter jumping biomechanics to enhance escape performance under chronic risk of spider predation. *Functional Ecology*, 25(1):279–288, 2011.
15. E. P. Azevedo, B. Tan, L. E. Pomeranz, V. Ivan, R. Fetcho, M. Schneeberger, K. R. Doerig, C. Liston, J. M. Friedman, and S. A. Stern. A limbic circuit selectively links active escape to food suppression. *eLife*, 9:1–23, 2020.
16. I. de Araujo Salgado, C. Li, C. J. Burnett, S. Rodriguez Gonzalez, J. J. Becker, A. Horvath, T. Earnest, A. V. Kravitz, and M. J. Krashes. Toggling between food-seeking and self-preservation behaviors via hypothalamic response networks. *Neuron*, 111(18):2899–2917.e6, 2023.
17. W. E. Cooper Jr and D. T. Blumstein. *Escaping From Predators : An Integrative View of Escape Decisions / Edited by William E. Cooper, Jr, Daniel T. Blumstein.* Cambridge University Press, Cambridge, 2015.

18. A. C. Schadegg and J. Herberholz. Satiation level affects anti-predatory decisions in foraging juvenile crayfish. *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology*, 203(3):223–232, 2017.

19. D. A. Blank. Escaping behavior in goitered gazelle. *Behavioural Processes*, 147(October 2017):38–47, 2018.

20. P. Shamash, S. F. Olesen, P. Iordanidou, D. Campagner, N. Banerjee, and T. Branco. Mice learn multi-step routes by memorizing subgoal locations. *Nature Neuroscience*, 2021.

21. P. Domenici, J. M. Blagburn, and J. P. Bacon. Animal escapology I: theoretical issues and emerging trends in escape trajectories. *Journal of Experimental Biology*, 214(15):2463–2473, 2011.

22. D. Eilam. Die hard: A blend of freezing and fleeing as a dynamic defense - Implications for the control of defensive behavior. *Neuroscience and Biobehavioral Reviews*, 29(8):1181–1191, 2005.

23. R. Bandler, A. Depaulis, and M. Vergnes. Identification of midbrain neurones mediating defensive behaviour in the rat by microinjections of excitatory amino acids. *Behavioural Brain Research*, 15(2):107–119, 1985.

24. A. Depaulis and R. Bandler, editors. *The Midbrain Periaqueductal Gray Matter: Functional, Anatomical, and Neurochemical Organization*. Springer New York, New York, 1st editio edition, 1991.

25. M. M. Behbehani. Functional characteristics of the midbrain periaqueductal gray. *Progress in Neurobiology*, 46(6):575–605, 1995.

26. M. B. Shaikh and A. Siegel. GABA-mediated regulation of feline aggression elicited from midbrain periaqueductal gray. *Brain Research*, 507(1):51–56, 1990.

27. J. S. Lonstein and J. M. Stern. Role of the midbrain periaqueductal gray in maternal nurturance and aggression: C-fos and electrolytic lesion studies in lactating rats. *Journal of Neuroscience*, 17(9):3364–3378, 1997.

28. J. Kohl, B. M. Babayan, N. D. Rubinstein, A. E. Autry, B. Marin-Rodriguez, V. Kapoor, K. Miyamishi, L. S. Zweifel, L. Luo, N. Uchida, and C. Dulac. Functional circuit architecture underlying parental behaviour. *Nature*, 556(7701):326–331, 2018.

29. Y. Li, J. Zeng, J. Zhang, C. Yue, W. Zhong, Z. Liu, Q. Feng, and M. Luo. Hypothalamic Circuits for Predation and Evasion. *Neuron*, 97(4):911–924.e5, 2018.

30. S.-G. Park, Y.-C. Jeong, D.-G. Kim, M.-H. Lee, A. Shin, G. Park, J. Ryoo, J. Hong, S. Bae, C.-H. Kim, P.-S. Lee, and D. Kim. Medial preoptic circuit induces hunting-like actions to target objects and prey. *Nature Neuroscience*, 21(March):364–372, 2018.

31. D. A. Evans, A. V. Stempel, R. Vale, S. Ruehle, Y. Lefler, and T. Branco. A synaptic threshold mechanism for computing escape decisions. *Nature*, 558(7711):590–594, 2018.

32. K. Tschida, V. Michael, J. Takatoh, B.-X. Han, S. Zhao, K. Sakurai, R. Mooney, and F. Wang. A Specialized Neural Circuit Gates Social Vocalizations in the Mouse. *Neuron*, pages S0896–6273(19)30478–7, 2019.

33. A. L. Falkner, D. Wei, A. Song, L. W. Watsek, I. Chen, P. Chen, J. E. Feng, and D. Lin. Hierarchical Representations of Aggression in a Hypothalamic-Midbrain Circuit. *Neuron*, 106(4):637–648.e6, 2020.

34. H. Deng, X. Xiao, and Z. Wang. Periaqueductal Gray Neuronal Activities Underlie Different Aspects of Defensive Behaviors. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 36(29):7580–8, 2016.

35. H. Yu, X. Xiang, Z. Chen, X. Wang, J. Dai, X. Wang, P. Huang, Z. dong Zhao, W. L. Shen, and H. Li. Periaqueductal gray neurons encode the sequential motor program in hunting behavior of mice. *Nature Communications*, 12(1), 2021.

36. J. Signoret-Genest, N. Schukraft, S. L. Reis, D. Segebarth, K. Deisseroth, and P. Tovote. Integrated cardio-behavioral responses to threat define defensive states. *Nature Neuroscience*, 26(3):447–457, 2023.

37. N. S. Canteras and M. Goto. Fos-like immunoreactivity in the periaqueductal gray of rats exposed to a natural predator. *NeuroReport*, 10(2), 1999.

38. S. R. Mota-Ortiz, M. H. Sukikara, J. Bittencourt, M. V. Baldo, C. Elias, L. F. Felicio, and N. S. Canteras. The periaqueductal gray as a critical site to mediate reward seeking during predatory hunting. *Behavioural Brain Research*, 226(1):32–40, 2012.

39. E. Vaughn, S. Eichhorn, W. Jung, X. Zhuang, and C. Dulac. Three-dimensional Interrogation of Cell Types and Instinctive Behavior in the Periaqueductal Gray. *bioRxiv*, page 2022.06.27.497769, 2022.

40. A. Fratzl, A. M. Koltchev, N. Vissers, Y. L. Tan, A. Marques-Smith, A. V. Stempel, T. Branco, and S. B. Hofer. Flexible inhibitory control of visually evoked defensive behavior by the ventral lateral geniculate nucleus. *Neuron*, 109(23):3810–3822.e9, 2021.

41. W. Wang, P. J. Schuette, J. Nagai, B. C. Tobias, F. M. Fernando, S. Ji, M. A. de Lima, M. Q. La-Vu, S. Maesta-Pereira, M. Chakerian, S. J. Leonard, L. Lin, A. L. Severino, C. M. Cahill, N. S. Canteras, B. S. Khakh, J. C. Kao, and A. Adhikari. Coordination of escape and spatial navigation circuits orchestrates versatile flight from threats. *Neuron*, 109(11):1848–1860.e8, 2021.

42. J. P. Fadok, S. Krabbe, M. Markovic, J. Courtin, C. Xu, L. Massi, P. Botta, K. Bylund, C. Müller, A. Kovacevic, P. Tovote, and A. Lüthi. A competitive inhibitory circuit for selection of active and passive fear responses. *Nature*, 542(7639):96–100, 2017.

43. B. A. Silva, C. Mattucci, P. Krzywkowski, E. Murana, A. Illarionova, V. Grinevich, N. S. Canteras, D. Ragozzino, and C. T. Gross. Independent hypothalamic circuits for social and predator fear. *Nature neuroscience*, 16(12):1731–3, 2013.

44. P. Wei, N. Liu, Z. Zhang, X. Liu, Y. Tang, X. He, B. Wu, Z. Zhou, Y. Liu, J. Li, Y. Zhang, X. Zhou, L. Xu, L. Chen, G. Bi, X. Hu, F. Xu, and L. Wang. Processing of visually evoked innate fear by a non-canonical thalamic pathway. *Nature communications*, 6:6756, 2015.

45. L. Wang, I. Chen, and D. Lin. Collateral Pathways from the Ventromedial Hypothalamus Mediate Defensive Behaviors. *Neuron*, 85(6):1344–1358, 2015.

46. B. A. Silva, C. T. Gross, and J. Gräff. The neural circuits of innate fear: detection, integration, action, and memorization. *Learning & Memory*, 23(10):544–555, 2016.

47. M. L. Brandão, V. Z. Anseloni, J. E. Pandóssio, J. E. De Araújo, and V. M. Castilho. Neurochemical mechanisms of the defensive behavior in the dorsal midbrain. *Neuroscience and Biobehavioral Reviews*, 23(6):863–875, 1999.

48. Y. Lefler, D. Campagner, and T. Branco. The role of the periaqueductal gray in escape behavior. *Current Opinion in Neurobiology*, 60:115–121, 2020.

49. F. M. Reis, D. Mobbs, N. S. Canteras, and A. Adhikari. Orchestration of innate and conditioned defensive actions by the periaqueductal gray. *Neuropharmacology*, 228:109458, 2023.

50. M. E. Masferrer, B. A. Silva, K. Nomoto, S. Q. Lima, and C. T. Gross. Differential encoding of predator fear in the ventromedial hypothalamus and periaqueductal grey. *Journal of Neuroscience*, 40(48):9283–9292, 2020.

51. P. Tovote, M. S. Esposito, P. Botta, F. Chaudun, J. P. Fadok, M. Markovic, S. B. E. Wolff, C. Ramakrishnan, L. Fenno, K. Deisseroth, C. Herry, S. Arber, and A. Luthi. Midbrain circuits for defensive behaviour. *Nature*, 534(7606):206–212, 2016.

52. G. R. Holt, W. R. Softky, C. Koch, and R. J. Douglas. Comparison of discharge variability in vitro and in vivo in cat visual cortex neurons. *Journal of neurophysiology*, 75(5):1806–1814, 1996.

53. I. Duguid, T. Branco, M. London, P. Chadderton, and M. Häusser. Tonic inhibition enhances fidelity of sensory information transmission in the cerebellar cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 32(32):11132–43, 2012.

54. J. U. Henschke, A. T. Price, and J. M. Pakan. Enhanced modulation of cell-type specific neuronal responses in mouse dorsal auditory field during locomotion. *Cell Calcium*, 96: 102390, 2021.

55. M. Yilmaz and M. Meister. Rapid innate defensive responses of mice to looming visual stimuli. *Current Biology*, 23(20):2011–2015, 2013.

56. R. Mongeau, G. A. Miller, E. Chiang, and D. J. Anderson. Neural correlates of competing fear behaviors evoked by an innately aversive stimulus. *J Neurosci*, 23(9):3855–3868, 2003.

57. P. Tovote, J. P. Fadok, and A. Lüthi. Neuronal circuits for fear and anxiety. *Nature Reviews Neuroscience*, 16(6):317–331, 2015.

58. Z. Xie, H. Gu, M. Huang, X. Cheng, C. Shang, T. Tao, D. Li, Y. Xie, J. Zhao, W. Lu, Z. Zhang, C. Zhan, Z. Tang, F. Zhang, and P. Cao. Mechanically evoked defensive attack is controlled by GABAergic neurons in the anterior hypothalamic nucleus. *Nature Neuroscience*, 25(1): 72–85, 2022.

59. W. Han, L. A. Tellez, M. J. Rangel, S. C. Motta, X. Zhang, I. O. Perez, N. S. Canteras, S. J. Shammah-Lagnado, A. N. van den Pol, and I. E. de Araujo. Integrated Control of Predatory Hunting by the Central Nucleus of the Amygdala. *Cell*, 168(1):311–324.e18, 2017.

60. H. Deng, X. Xiao, and Z. Wang. Periaqueductal Gray Neuronal Activities Underlie Different Aspects of Defensive Behaviors. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 36(29):7580–8, 2016.

61. G. Gatto and M. Goulding. Locomotion Control: Brainstem Circuits Satisfy the Need for Speed. *Current Biology*, 28(6):R256–R259, 2018.

62. E. Tsang, C. Orlandini, R. Sureka, A. H. Crevenna, E. Perlas, I. Prankerd, M. E. Masferrer, and C. T. Gross. Induction of flight via midbrain projections to the cuneiform nucleus. *PLoS ONE*, 18(2):e0281464, 2023.

63. R. P. Bindi, C. C. Guimarães, A. R. de Oliveira, F. F. Melleu, M. A. de Lima, M. V. Baldo, S. C. Motta, and N. S. Canteras. Anatomical and functional study of the cuneiform nucleus: A critical site to organize innate defensive behaviors. *Annals of the New York Academy of Sciences*, 1521(1):79–95, 2023.

64. A. Bittencourt, E. Nakamura-Palacios, H. Mauad, S. Tufik, and L. Schenberg. Organization of electrically and chemically evoked defensive behaviors within the deeper collicular layers as compared to the periaqueductal gray matter of the rat. *Neuroscience*, 133(4):873–892, 2005.

65. M. J. Ferreira-Pinto, L. Ruder, P. Capelli, and S. Arber. Connecting Circuits for Supraspinal Control of Locomotion. *Neuron*, 100(2):361–374, 2018.

66. N. Josset, M. Roussel, M. Lemieux, D. Lafrance-Zoubga, A. Rastqar, and F. Bretzner. Distinct Contributions of Mesencephalic Locomotor Region Nuclei to Locomotor Control in the Freely Behaving Mouse. *Current Biology*, 28(6):884–901.e3, 2018.

67. S. Arber and R. M. Costa. Networking brainstem and basal ganglia circuits for movement. *Nature Reviews Neuroscience*, 23(6):342–360, 2022.

68. R. Leiras, J. M. Cregg, and O. Kiehn. Brainstem Circuits for Locomotion. *Annual Review of Neuroscience*, 45:63–85, 2022.

69. S. P. Ventura, D. C. Passos, L. L. Machado, G. Horta, and C. A. Galdino. Escape tactics by a neotropical montane lizard: a comparison of flight responses against natural and nonnatural predators. *Acta Ethologica*, 20(1):9–15, 2017.

70. P. Domenici and G. D. Ruxton. Escape trajectories and strategies during pursuit. In W. E. C. Jr and D. T. Blumstein, editors, *Escaping from Predators: An Integrative View of Escape Decisions*. Cambridge University Press, Cambridge, 2015.

71. W. E. Cooper. Directional escape strategy by the striped plateau lizard (Sceloporus virgatus): Turning to direct escape away from predators at variable escape angles. *Behaviour*, 153(4):401–419, 2016.

72. P. Shamash, S. Lee, A. M. Saxe, and T. Branco. Mice identify subgoal locations through an action-driven mapping process. *Neuron*, 111(12):1966–1978.e8, 2023.

73. D. Campagner, R. Vale, Y. L. Tan, P. Iordanidou, O. Pavón Arocas, F. Claudi, A. V. Stempel, S. Keshavarzi, R. S. Petersen, T. W. Margrie, and T. Branco. A cortico-collicular circuit for orienting to shelter during escape. *Nature*, 613(7942):111–119, 2023.

74. O. Pavón Arocas and T. Branco. Preparation of acute midbrain slices containing the superior colliculus and periaqueductal Gray for patch-clamp recordings. *PLOS ONE*, 17(8): e0271832, 2022.

75. A. Mathis, P. Mamidanna, K. M. Cury, T. Abe, V. N. Murthy, M. W. Mathis, and M. Bethge. DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nature Neuroscience*, 21(9):1281–1289, 2018.

76. A. Giovannucci, J. Friedrich, P. Gunn, J. Kalfon, B. L. Brown, S. A. Koay, J. Taxidis, F. Najafi, J. L. Gauthier, P. Zhou, B. S. Khakh, D. W. Tank, D. B. Chklovskii, and E. A. Pnevmatikakis. CaImAn an open source tool for scalable calcium imaging data analysis. *eLife*, 8, 2019.

77. G. Paxinos and K. Franklin. *The Mouse Brain in Stereotaxic Coordinates*. Academic Press, 2nd editio edition, 2001.

78. N. A. Steinmetz, C. Aydin, A. Lebedeva, M. Okun, M. Pachitariu, M. Bauza, M. Beau, J. Bhagat, C. Böhm, M. Broux, S. Chen, J. Colonell, R. J. Gardner, B. Karsh, F. Kloosterman, D. Kostadinov, C. Mora-Lopez, J. O'Callaghan, J. Park, J. Putzeys, B. Sauerbrei, R. J. J. van Daal, A. Z. Vollan, S. Wang, M. Welkenhuysen, Z. Ye, J. T. Dudman, B. Dutta, A. W. Hantman, K. D. Harris, A. K. Lee, E. I. Moser, J. O'Keefe, A. Renart, K. Svoboda, M. Häusser, S. Haesler, M. Carandini, and T. D. Harris. Neuropixels 2.0: A miniaturized high-density

probe for stable, long-term brain recordings. *Science*, 372(6539):eabf4588, 2021.

79. A. P. Buccino, C. L. Hurwitz, S. Garcia, J. Magland, J. H. Siegle, R. Hurwitz, and M. H. Hennig. SpikeInterface, a unified framework for spike sorting. *eLife*, 9:e61834, 2020.

80. M. Pachitariu, S. Sridhar, and C. Stringer. Solving the spike sorting problem with Kilosort. *bioRxiv*, page 2023.01.07.523036, 2023.

81. D. Mayerich, L. Abbott, and B. McCormick. Knife-edge scanning microscopy for imaging and reconstruction of three-dimensional anatomical structures of the mouse brain. *Journal of Microscopy*, 231(1):134–143, 2008.

82. T. Ragan, L. R. Kadiri, K. U. Venkataraju, K. Bahlmann, J. Sutin, J. Taranda, I. Arganda-Carreras, Y. Kim, H. S. Seung, and P. Osten. Serial two-photon tomography for automated ex vivo mouse brain imaging. *Nature Methods*, 9(3):255–258, 2012.

83. J. Kim and Y. Li. Chronic activation of CB2 cannabinoid receptors in the hippocampus increases excitatory synaptic transmission. *The Journal of physiology*, 593(4):871–86, 2015.

84. F. Claudi, L. Petrucco, A. Tyson, T. Branco, T. W. Margrie, and R. Portugues. BrainGlobe Atlas API: a common interface for neuroanatomical atlases. *The Journal of Open Source Software*, 5(54):2668, 2020.

85. A. L. Tyson, M. Vélez-Fort, C. V. Rousseau, L. Cossell, C. Tsitoura, S. C. Lenzi, H. A. Obenhaus, F. Claudi, T. Branco, and T. W. Margrie. Accurate determination of marker location within whole-brain microscopy images. *Scientific Reports*, 12(1):867, 2022.

Methods

Animals. Adult (6–14 weeks old), male and female VGAT-ires-Cre (Jackson Laboratory, stock #028862), VGluT2-ires-Cre (Jackson Laboratory, stock #016963), crossed VGAT::tdTomato (Jackson Laboratory, stock # 007908) and VGAT::eYFP or VGluT2::eYFP (Jackson Laboratory, stock #006148) mice were housed with ad libitum access to water and chow on a 12 h reversed light cycle and tested during the light phase. A subset of *in vitro* patch clamp experiments was made from Gad2-T2a-NLS-mCherry animals (Jackson Laboratory, stock #023140) (see below). Silicon probe recordings were performed in 8–10 weeks old, male C57BL/6J mice, obtained from commercial suppliers (Charles River). Animals were used for scientific purposes in accordance with the the UK Animals (Scientific Procedures) Act of 1986 and the Animal Welfare and Ethical Review Body (AWERB), guidelines stated in Directive 2010/63/EU of the European Parliament on the protection of animals used for scientific purposes, institutional guidelines and following approval by the relevant authorities.

General surgical procedures. Drugs were prepared on the day of surgery, diluted in sterile 0.9 % NaCl saline and warmed before use. Animals were anaesthetised with an intraperitoneal (i.p.) injection of ketamine (100 mg/kg) and xylazine (12 mg/kg), and carprofen (5 mg/kg) was administered subcutaneously for perioperative analgesic care. The hair on the forehead was shaved using a trimmer (Contura, Wella), and the skin disinfected. Isoflurane (0.5–1.5 % in oxygen, 1 L/min) was used to maintain anaesthesia. The skin was then cut to expose the skull, and a large craniotomy was made using a hand-held dental drill using a 0.5 or 0.7 mm stainless steel burr (Fine Science Tools) to expose an area between the confluence of transverse and sagittal sinus, and the anatomical border between the superior and inferior colliculi (AP from lambda: +0.1 to -1.0 mm; ML: ± 0.6 mm). A durotomy was performed using a 30G hypodermic needle. The connective tissue covering the confluence of sagittal and transverse sinus was gently pulled forward anteriorly using fine forceps to reveal the underlying superior colliculi. Viral vectors were delivered using a pulled cut glass pipette (10μl Wiretrol II with a Sutter P-1000) that was mounted on a stereotaxic injector (Janelia Research Campus, Ronal Tool Company) coupled to an oil hydraulic micromanipulator (MO-10, Narishige) on a stereotaxic frame (Model 963, Kopf Instruments), at ~20 nl/min. After injection, the needle was kept at the site of injection for 10–20 min, before being retracted. Fibre optic and GRIN lens cannula implants were first slowly lowered to 50 μm beyond their final location, and then retracted. For lens cannula implants, a single 400 μm diameter fibre optic with a 60º conical tip was pre-inserted at the desired location and withdrawn, to allow for easier successive insertion of the 500 μm lens implant. The craniotomy was covered with a silicone sealant (Kwik-Cast, WPI) before implants were secured using light-cured dental cement (RelyX Unicem 2, 3M). The skin wound was glued (Vetbond, 3M) or sutured (6-0, Vicryl Rapide). To ensure maximal stability during calcium imaging, the fissures between the interparietal and occipital plates were glued after the implantation of GRIN lens cannulae and before wound closure. Coordinate measurements are given from lambda and skull surface. For precise rostrocaudal and mediolateral targeting of fibre placements and viral injections, coordinates were confirmed and, if necessary, re-aligned after the craniotomy and durotomy were performed based on the border between the superior and inferior colliculi (~0.65–0.8 mm behind lambda) and the small midline blood vessel that runs sagittal between the two superior colliculi.

Viruses. The following viruses were used in this study and are referred to by abbreviations in the text. For chemogenetic inhibition experiments, AAV2/2-hsyn-DIO-hM4D(Gi)-mCherry was obtained from Addgene (plasmid #44362, 4.6×10^{12} GC/ml). For optogenetic activation, AAV2/2-EF1a-DIO-hChR2(H134R)-EYFP-WPRE (3.9×10^{12} GC/ml) or AAV2/2-EF1a-DIO-hChR2(H134R)-mCherry-WPRE (6.6×10^{12} GC/ml; Deisseroth, UNC Gene Therapy Vectore Core) were used. Optogenetic inhibition experiments were performed with AAV2/9-EF1a-DIO-iChloC-2A-tDimer (3.75×10^{12} GC/ml; a gift from S. Wiegert and T. Oertner) or AAV2/1-EF1a-DIO-iChloC-2A-dsRed (5×10^{13} GC/ml; Addgene plasmid #70762, a gift from T. Margrie). For calcium imaging experiments, AAV2/1-hsyn-DIO-GCaMP6s-WPRE (6.25×10^{12} GC/ml) was acquired from Penn Vector Core.

Electrophysiology recordings *in vitro*. Patch clamp recordings were performed from identified glutamatergic and GABAergic neurons in acute, coronal slices of the midbrain containing the PAG and as previously described (31, 74).

Slice preparation. Slices were prepared by one of two alternative slicing methods (see lab protocol by Pavon Arócas and Branco (74) for details). In brief, animals were either anaesthetised using isoflurane and decapitated directly (path A) or after intracardial perfusion with ice-cold artificial cerebrospinal fluid (aCSF) (path B). Path A: The brain was extracted and submerged in carbogenated ice-cold sucrose-based aCSF containing (in mM): 87 NaCl, 26 NaHCO$_3$, 50 sucrose, 10 glucose, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 3 MgCl$_2$, 0.5 CaCl$_2$. Tissue blocks containing the midbrain PAG were mounted on a vibratome (VT1200S, Leica) and coronal slices

(-4.8 to -4.3 mm from bregma) of 250–300 μm thickness were cut at 0.05–0.06 mm/s. After cutting, slices were kept in sucrose-based aCSF at near-physiological temperature (35º C) for 30 min before being transferred to a beaker with 35º C-warm aCSF (containing in mM: 125 NaCl, 26 NaHCO$_3$, 10 glucose, 2.5 KCl, 2 CaCl$_2$, 1 MgCl$_2$, 1 NaH$_2$PO$_4$) that was then stored at room temperature for at least half an hour before recordings began. Path B: For a subset of experiments, the slicing and recovery aCSF contained (in mM): 96 NMDG, 96 HCl, 25 NaHCO$_3$, 25 D-Glucose, 20 HEPES, 5 Sodium L-ascorbate, 12 N-actyl-L-cystein, 10 MgSO$_4$, 3 Sodium pyrate, 3 Myo-inositol, 2 Thiourea, 1.25 NaH$_2$PO$_4$, 0.5 Ca$_2$Cl, 0.01 Taurine. Slices were kept at 32–34º C to recover for a period of <15 min and then held in HEPES-based aCSF until recordings began. For both paths, slices were stored at RT for ~4–5 hours after preparation before being discarded and all aCSF was equilibrated with 95% O$_2$, 5% CO$_2$.

General setup. Individual midbrain slices containing the PAG were transferred to a submerged chamber, perfused with ACSF heated to 32–33º C at a rate of 2–3 ml/min and visualised on an upright microscope with oblique contrast (Slicescope, Scientifica) using a 60x or 40x objective (Olympus). VGluT2- and VGAT-positive dPAG neurons were identified based on location, fluorescence protein expression and cell-attached firing properties. In a subset of recordings, putative glutamatergic neurons were identified by being VGAT- or GAD2-negative (not expressing a fluorophore; mCherry or eYFP) and non-spiking in cell-attached configuration before break-in. Given that virtually all GABAergic neurons are tonically spiking and excitatory neurons are not, this is a reliable method to identify presumptive excitatory dPAG neurons. Patch clamp recordings were performed with an EPC 800 (HEKA) amplifier and monitored using an HMO1002 oscilloscope (Rohde & Schwarz). Signals were filtered at 5 kHz, digitised with 16-bit resolution at 25 kHz using a PCI 6035E A/D board and recorded in LabVIEW (both National Instruments) using custom software. A subset of recordings was performed and digitised using a Multiclamp 700B amplifier with an Axon Digidata 1550B Digitiser, and recorded using Clampex (Molecular Devices).

Whole-cell and loose-seal patch clamp recordings. Pipettes were pulled from borosilicate filamented glass capillaries (Harvard Apparatus, 1.5 mm OD, 0.85 mm ID) with a P-10 or PC-100 Narishige puller to a final resistance of ~4–7 MΩ. Unless stated otherwise, for whole-cell patch clamp recordings, pipettes were backfilled with filtered internal solution containing (in mM): 130 KMeSO$_3$, 10 KCl, 10 HEPES, 5 Na-phosphocreatine, 4 Mg-ATP, 1 EGTA, 0.5 Na$_2$-GTP, 285–290mOsm, pH was adjusted to 7.35 with KOH. In some recordings, 1 mg/ml biocytin or neurobiotin was added on the day. To analyse inhibitory postsynaptic currents (IP-SCs) onto VGluT2 neurons, a 'high-chloride' internal solution was used containing in mM: 155 KCl, 10 HEPES, 2 EGTA, 2 Na-ATP, 2 Mg-ATP, 0.3 Na-GTP or 140 KCl, 10 HEPES, 5 Na-phosphocreatine, 2 Mg-ATP, 0.5 Na-GTP, 0.2 EGTA, 285–290mOsm, pH adjusted to 7.35 with KOH. For

loose-seal cell-attached recordings, pipettes were backfilled with aCSF and a seal with a resistance of 10–20 MΩ was established. Action currents in voltage-clamp mode were recorded continuously for 5–6 min. A holding potential of 0 mV was maintained through current injections (within a range of ± 20 pA) and the access resistance was monitored throughout the recording. For whole-cell recordings, the resting membrane potential was determined in current clamp directly after establishing the whole-cell configuration and experiments were only continued if cells had a resting membrane potential more hyperpolarised than -45 mV (not corrected for liquid junction potential). In a subset of recordings, cells were further characterised by recording their membrane responses to current injections of increasing strengths (-40 to + 140 pA, increment: 20 pA, 500 ms duration). The access resistance was monitored continuously, and was compensated in current-clamp recordings. Only cells with a stable access resistance < 30 MΩ were analysed. Voltage-clamp recordings were performed at a holding potential of -60 mV with the exception of IPSC recordings using the high-chloride internal solution, which were performed at -70 mV.

Stimulation. Electrical stimulation was performed using a stimulus isolator (DS2A, Digitimer) with a pipette placed in the deeper layers of the superior colliculus. Pulse trains were delivered every 15 s with a stimulation intensity of 0.02–4.3 mA, a pulse duration of 100 μs and an interpulse interval of 50 ms. For optogenetic stimulation experiments, recordings were made > 30 days after injection of AAV2-DIO-ChR2-mCherry into the dPAG of VGAT-Cre mice. Wide-field 490 nm LED illumination (pe-100, CoolLED) was used to excite ChR2 expressed in VGAT dPAG neurons. Pulse trains (5 pulses, 50 ms interpulse interval, 1 ms pulse duration, maximum light intensity = 2.7 mW) were delivered every 30 s. The stimulation intensity for both electrical and optogenetic stimulation was adjusted for each experiment.

Pharmacology. To block glutamatergic synaptic transmission, kynurenic acid (Sigma) was added to the aCSF as powder to a final concentration of 2 mM and sonicated until dissolved. GABA$_A$ and glycinergic receptor-mediated synaptic transmission was blocked using 50 μM picrotoxin (Tocris Bioscience), which was prepared from a 100 mM stock solution containing DMSO (final DMSO concentration: 1/2000). GABA$_A$ receptor-mediated synaptic transmission was blocked using 10 μM SR-95531 hydrobromide (Gabazine, Hellobio). Water-soluble clozapine N-oxide dihydrochloride (CNO, Hellobio) was bath applied at a final concentration of 10 μM from a 100 mM stock solution. Control recordings (matched in duration to experiments with bath application of drugs) were performed interleaved and, if possible, in the same animals. Drugs were added to the aCSF as stated in the text or figure legends.

Data analysis. Membrane potentials were not corrected for liquid junction potential. The spontaneous firing frequency of neurons recorded in loose-seal cell-attached configuration was analysed in a period of 2–4 min after seal formation. For whole-cell recordings, the resting membrane potential was calculated during current steps in a 100 ms

baseline window in current clamp mode. The input resistance (R_{in}) was calculated from the steady-state voltage measured in response to a hyperpolarising test pulse (-40 pA, 500 ms duration) at holding potential of -60 mV. The membrane time constant was calculated by fitting a single exponential to the decay of the test pulse ($y = y0 + Ae^{(-(x-x0)/\tau)}$). The mean action potential firing frequency of VGAT neurons in response to current injection of increasing strengths (20–140 pA, 20 pA increments) was calculated from two consecutive repetitions per cell. Spontaneous inhibitory postsynaptic currents (IPSCs) onto glutamatergic dPAG neurons were detected using a threshold algorithm, and their frequency and peak amplitude were analysed from a continuous 30 s time window every 1 min or continuously and binned by minute. A subset of spontaneous IPSC recordings was analysed using threshold-based IPSC detection in Igor Pro (Version 9, Wavemetrics) with TaroTools (written by Taro Ishikawa, Jikei University School of Medicine, Tokyo, Japan). The peak phasic conductance was calculated from the cell's mean peak current amplitude divided by the holding potential ($G = I / V_{hold}$). The mean unitary charge was calculated from the average spontaneous IPSC for each cell, integrating the area over a 50 ms window from IPSC onset. The average charge was estimated by multiplying the unitary charge by the frequency of events for that cell. For estimating the change in IPSC frequency in SR-95531 and CNO respectively, drugs were bath applied after a stable baseline recording of 2–5 min and recordings were continued for > 5 min. The mean baseline IPSC frequency was calculated from 2 min before bath application of the drug, and the mean IPSC frequency after drug wash-in was calculated for min 7–8 for CNO and min 5–7 for SR-95531 and their respective controls. Pharmacological effects of each drug were statistically assessed in comparison to time-matched controls (see above). Mean peak amplitudes of evoked postsynaptic currents were calculated from averages of a minimum of 5 sweeps both for baseline and drug conditions (where applicable). The paired pulse ratio of a synaptic input was calculated by dividing the mean peak amplitude of the second pulse by the first pulse. Statistical tests were performed on all cells pooled across animals. The coefficient of variation was calculated for each cell as the standard deviation of the inter-spike intervals (ISI) over the mean ISI. CV_2 was calculated after Holt et al. (52) as:

$$CV_2 = \frac{2 \mid \Delta t_{i+1} - \Delta t_i \mid}{\Delta t_{i+1} + \Delta t_i}$$

for each spike i, and the average CV_2 was taken for each cell.

Behavioural recordings.

Setup and acquisition. Behavioural experiments were performed in a rectangular arena made from Perspex (L: 60–80 cm × W: 20 cm × H: 40cm) with an infrared light-transmitting red shelter (L: 10 cm × W: 19 cm × H; 13.5 cm) in one corner of the arena, and as previously described (31). Briefly, a screen made of 100 μm thick drafting film (90 cm × 70 cm, Elmstock) was suspended 64 cm above the arena floor, and an ultra-short throw projector (PF1000U, LG) was used to project a grey background onto the screen, providing a uniform luminance of 3–4 lux (measured on the floor of the arena). Behaviour was recorded at 50 frames per second with a near-infrared camera (acA1300-60gmNIR GigE, Basler) positioned above the arena centre, flush with the screen. Video recordings, as well as sensory and optogenetic stimulation protocols, were controlled with custom-written software (mantis64) in LabVIEW (2015 64-bit, National Instruments). A PCIe-6351 board (National Instruments) was used to trigger and synchronise all signals. Sensory stimuli were delivered when the mouse entered a predefined 'threat area' at the opposite end of the arena to the shelter (~15–20 × 20 cm). A 35 mm Petri dish was positioned in the threat area to enrich the environment and encourage exploration. Animals were allowed to explore the arena for a minimum of 7 min at the beginning of the recording, and a behavioural session typically lasted between 30 min and 2 hours.

Sensory stimulation. Escape was elicited using both auditory and visual stimuli. A frequency-modulated upsweep from 17 to 20 kHz over 3 s was used as an auditory stimulus, and delivered at different intensities ranging from 60 to 90 dB above the threat area via an ultrasound speaker (L60, Pettersson). The visual stimulus consisted of a sequence of 3 or 5 dark, linearly expanding circles (visual angle min–max: 2.6–47°, linear expansion: 118° / s over 260 ms, size maintained for 260 ms at the maximum visual angle) with an inter-spot interval of 500 ms. The contrast of the spots ranged between a Weber contrast of 60, 75 and 98 % (from low to high, converted to percentage from the respective negative Weber fractions), and was adjusted by changing the intensity of the circle against the grey screen which was maintained at a constant luminosity (see above). Visual stimuli were predominantly used and interleaved with auditory stimuli to prevent habituation. The stimulus intensity was adjusted based on the escape probability of each mouse in each session, and the experimental paradigm (as described in the results).

Behavioural data analysis. Mouse coordinates were extracted from the behavioural videos using DeepLabCut (75). Behavioural events were scored and annotated manually, including for escape: (1) first stimulus-reaction response after sensory stimulation (usually an ear movement followed by a brief startle), (2) beginning of the shelter-directed head-turn followed by a full-body turn, (3) start of run towards shelter, (4) escape termination. A successful escape was defined as a sequence of events (1) – (4), with an *uninterrupted* run into the shelter within 6 s of the end of a sensory stimulus (7.5 s for freely-moving calcium recordings to account for head-mounted microscope weight). The escape probability for a given stimulus is the fraction of trials which led to an escape to the shelter and was calculated for each animal separately. The peak velocity of the escape was determined by analysing the speed trace from the initiation of the escape until entry into the shelter.

Optogenetic experiments in freely moving animals.

Surgeries. For optogenetic activation experiments,

VGAT-Cre mice were injected with AAV-DIO-ChR2 into the dPAG (1–3 injections per hemisphere with a combined volume of 40–150nl per side; ML: ±0.4 to 0.45, AP: -4.6 to -4.9, DV: -2.2). Dual optic fibres with 60° conical tips (400 μm diameter, 1.2 mm apart, DFC_400/430-0.48_3.5mm_GS1.2_C60; Doric Lenses Inc.) were implanted above the centre of the injection sites (AP: -4.6 to -4.9, DV: -1.9 to -2.0). Behavioural testing was performed 13–65 days after surgery (mean = 31.1 ± 4.3 days), with 2–5 sessions per animal. For optogenetic inactivation experiments, VGAT-Cre mice were injected with AAV-DIO-iChloC into the dPAG (1–2 injections per hemisphere with a combined volume of 100–180nl per side; ML: ±0.4 to 0.45, AP: -4.6 to -4.9, DV: -2.2). Dual optic fibres with 400 μm thickness were implanted as described above (AP: -4.6 to -4.9, DV: -1.9). In two experiments, a single optic fibre (200-μm diameter, MFC-SMR; Doric Lenses Inc.) was implanted medially (ML: 0.0, AP: -4.6 to -4.9, DV: -1.9).

Behavioural testing. Behavioural testing was performed 12–40 days after surgery (mean = 23.8 ± 3.8 days), with 1–5 sessions per animal. Dual or single fibre-optic magnetic patch cords were attached to a fibre-optic rotary joint (FRJ_1×2i_FC-2FC_0.22 or FRJ1×1, both Doric Lenses Inc.) to allow unhindered movement. Light was delivered by a 472 nm laser module (Stradus, Vortran) with direct analogue modulation of laser power. Mice were presented with visual or auditory stimuli that elicited escape, and laser-on trials were interleaved with laser-off trials. Laser-on movement controls were performed without sensory stimulation. ChR2 stimulation was delivered at a frequency of 50 Hz for 3 or 5 s (pulse duration: 5 ms, laser intensity: 18.5–30 mW per hemisphere/optic fibre). Continuous blue-light stimulation of iChloC was delivered for 3 or 5 s at laser intensities of 0.03–0.3 mW using a neutral density filter (optical density: 0.3, Thorlabs) fitted in front of the laser output. Laser intensities were calibrated during exploratory movement to ensure no escape-like movements were elicited with too high laser power. To test for a bidirectional behavioural effect of VGAT neuron activation and inhibition, two separate behavioural paradigms were used to adjust the initial escape probability by changing the intensity of the visual (contrast) and auditory (dB) stimuli. Assuming a decrease in escape probability with VGAT neuron activation, we adjusted the sensory stimuli for each mouse during this experimental set to yield a high escape probability of ~ 80%. Conversely, hypothesising an increase of escape probability with VGAT neuron inactivation, we used low intensity stimuli to achieve a baseline escape probability of ~ 20–40%. To test the effect of optogenetic activation of VGAT neurons on escape termination, the laser was triggered manually immediately after the onset of stimulus-evoked escape runs. To test whether inactivation of VGAT neurons would lead to an increase in the escape distance, we moved the shelter away from the end wall into the middle of the arena. The shelter is open front and back to allow animals to pass through. The animals were allowed to explore the new shelter location, before a sensory stimulus was delivered to elicit escape. If the escape was to the new

shelter location, indicating that the animal had memorized the new shelter location, we proceeded to interleave the sensory stimulation with concomitant iChloC activation trials.

Data analysis. Behavioural tracking data was annotated and analysed as described above. The effect of optogenetic activation of VGAT[+] neurons on escape termination is calculated as the normalised distance from the shelter (0 = shelter entrance, 1 = far end of arena) at which the animal stopped its escape run. An average of all trials per tested animal is reported.

Calcium imaging in freely moving animals.

Surgeries and acquisition. Calcium imaging experiments were performed as previously described (31). Briefly, a miniaturised head-mounted fluorescence microscope (Model L, Doric Lenses Inc., Canada) was used to image the genetically encoded calcium sensor GCaMP6s in VGAT neurons of the dPAG. A volume of 100 nl AAV2/1-hsyn-DIO-GCaMP6s was injected into the dPAG (AP: -0.45 to -0.7, ML: +0.25–0.3, DV: -2.2). A 500 μm diameter GRIN lens imaging cannula (Model L, Doric Lenses Inc.) with an 80 μm working distance was slowly inserted, with the final tip position ~100 μm above the centre of the viral injection site. At least 19 days after surgery, the microscope was attached to the snap-in cannula implant on the mouse's head without anaesthesia and the mouse placed in the behavioural arena. Behavioural recordings were performed only after ensuring a clear field-of-view (FOV) was present (mean first session = 35.1 ± 5.3 days, mean last session = 62.2 ± 9.3 days after surgery; range = 19–104 days, N = 8 mice). Data were acquired with a resolution of 630 × 630 pixels at 10–20 Hz, with an excitation power of 150–300 μW and with a gain of 1 or 2. Sensory stimulation was delivered as described above.

Data analysis. Fluorescence movies were downsampled from 630 × 630 to 126 × 126 pixels and then flatfield corrected. To obtain an estimate of the movie's flatfield, we calculated the average intensity across time and applied a gaussian blur to the resulting image. The flat-field was centred around zero by subtracting its mean and this image was then subtracted from all frames of the movie. The movies were cropped to remove vignetting at the edges of the recorded FOV. The open-source analysis pipeline CaImAn (76) was used for motion correction and component extraction and curation. Briefly, movies were motion corrected using the NoRMCorre algorithm that corrects non-rigid motion artifacts, and source extraction was performed using constrained non-negative matrix factorization to extract components with overlapping spatial footprints. The extracted components were manually curated (merged, rejected or added) after visual inspection and the components' spatial shapes saved as cell masks. These masks were used to extract the background-subtracted (rolling-ball, 10px radius) raw fluorescence signal over time from the motion-corrected videos, and linearly interpolated to 30 Hz to match the behaviour camera frame-rate. Imaging sessions were almost always intermittent (mean 16 days between sessions with 1–3 sessions per 8 animals, range 1–72 days, one animal with three

consecutive days), and precluding FOV registration between imaging sessions and repeated cell identification. Each FOV was therefore treated as an independent sample. Delta F/F and Z-scored activity was used as the basis of all further analysis. Traces with a Z-score above 1.96 at any time during the recording session were considered active neurons.

To analyse VGAT$^+$ neuron activity in the freely-moving animal in the absence of threat, single neuron activity traces acquired during the baseline period of each session (before the first threat stimulus, mean duration 548 s, range 339–1046 s, n = 209 neurons, from 17 sessions in 8 animals) were Z-scored and Savitzky-Golay filtered (2 s window, polyorder = 2, *scipy*). Events were then detected using the *scipy.find_peaks* function (prominence = 0.3) to calculate the mean event rate per mouse. Neurons were further sorted by whether they were active during escape or not (see methods below), and whether events were coincident with animal movement, defined as the centre-of-mass of the animal exceeding 2 cm/s. For quantifying locomotion, movement periods were defined by timepoints when the animal's speed was greater than 1.5 cm/s and stationary periods as less than 1.5 cm/s while the animal was outside the shelter in the arena, in the baseline period before the first threat stimulus. A locomotion modulation index (LMI) and significance tests were calculated after Henschke et al [54]. The LMI was defined as the difference between the mean F/F$_0$ during locomotion (R$_L$) and stationary (R$_s$) periods, normalised by the sum, for each neuron as LMI = (R$_L$ − R$_s$)/(R$_L$ + R$_s$). To determine significance for the modulation indices, bootstrapping with replacement was used to estimate the error for each neuron. The data was binned into 1 s periods of continuous locomotion or stationary periods. The mean ΔF/F$_0$ was taken for each bin and regarded as a single sample. The average correlation between consecutive samples was found to be R = 0.57. We then randomly selected a reduced number of samples (43% = 1-R, to reflect the fact that samples are not completely independent) of ΔF/F$_0$ with replacement from the original set of samples, repeating this 1000 times to obtain 95% confidence intervals for individual significance tests for each neuron. If a neuron's 95% confidence interval was significantly different from an LMI of 0 and its absolute LMI value exceeded 0.2, indicating >50% change in F/F$_0$ between locomotion and stationary states, it was considered significantly locomotion responsive.

To identify neurons active during escape trials, Z-scored activity traces were first baselined by subtracting the mean value of 500 ms preceding stimulus onset. For each neuron, the distribution of activity slopes for each trial event (linear regression to a 660 ms window of activity starting at the following event onsets: *stimulus, first stimulus-reaction response, escape turn, escape run, shelter entry, escape stop*) over all trials in a session, was compared to the distribution of activity slopes in the 660 ms baseline periods preceding stimulus onset (two-tailed t-test with Bonferroni correction, P = 0.05) to determine whether there was a statistically significant change in signal for any event in the stimulus-escape sequence for that neuron, followed by a manual inspection and curation. 39 neurons showed a motion artefact (sharp

symmetrical downward deflection) as the microscope cables contacted the roof of the shelter during escape running, which was removed by cubic interpolation 500 ms either side of the artefact trough. To compare neural activity across all escape-to-shelter trials and all animals, where the sequence of behavioural events is stereotyped but inter-event intervals are variable, the median inter-event intervals for each behavioural event relative to stimulus onset was calculated and the median of these distributions was used to construct a template with a 2 s window added after escape-end. Z-scored calcium traces were then linearly interpolated to the template and the mean calcium trace calculated for each escape-active neuron (n = 123) for escape trials and fail-to-escape trials. As experiments were acquired at 10–20 Hz, all traces were identically low-pass FFT filtered to remove high-frequencies that was present in a subset of sessions. The mean calcium over time traces of all neurons (escape trails only) were transformed by principal component analysis (PCA) and K-means clustering was performed on the first 3 principal components (cumulatively explaining 86% of the variance) using n = 2 clusters, indicated as optimal by silhouette scores and elbow method. The timing of calcium activity relative to escape end was calculated using the time of peak activity over the entire mean time warped trace for each neuron.

Histological quantification of virus expression and fibre location. For *in vitro* electrophysiological recordings, the expression of virus was visually confirmed for each slice using 490 nm or 568 nm LED illumination (for green and red fluorescent proteins, respectively) prior to experimentation on the upright microscope. Slices from animals with clear infection of the deeper layers of the superior colliculus were discarded. For histological confirmation of fibre placement and injections site of *in vivo* calcium imaging and optogenetic experiments, mice were anaesthetized with pentobarbital (400–800 mg/kg, final volume: 0.1 ml per 10 g, diluted 1:1 in sterile 0.9 % NaCl solution) and decapitated. To remove optic fibres and GRIN lenses without damage to the brain, the dorsal skull was cut in a circle around the cemented fibres and was carefully removed together with the fibres. Brains were then extracted from the base of the skull and fixed in 0.1 M PBS containing 4% PFA overnight at 4º C. Slices of 100 μm thickness were cut in 0.1 M PBS on a HM650V vibratome (Microm), mounted using SlowFade Gold Antifade Mountant with DAPI (ThermoFisher) and imaged on a wide-field fluorescence microscope (AxioImager, Zeiss). The placement of optic fibres and GRIN lenses was assessed based on their tract and tip location. Tip locations are illustrated in the respective sections of the Paxinos mouse brain atlas [77] (see Figures S4 and S7). The expression of viruses was visually confirmed. For AAV2/2-DIO-ChR2 injections, only animals with expression localised to the dPAG were used. Animals with expression in the deeper layers of the SC were excluded. For AAV-DIO-iChloC injections (AAV2/1 or AAV2/9), experiments were included based on fibre tip location in the dPAG. Animals with tip locations outside the caudal dPAG (< -4.6 mm or > -5 mm

from bregma) or tip locations not aligned with the virus injection were excluded.

Chronic silicon probe recordings *in vivo*.

Surgery. Mice were anaesthetised in an isoflurane induction chamber (3–4 % isoflurane, 1 L/min) before being transferred to a nose cone (2 % isoflurane, 1 L/min). Surgical procedures were performed under anaesthesia using isoflurane (1.5–2 %, 1 L/min). Protective gel was applied to the eyes (Puralube Vet Ointment), analgesia was given subcutaneously (Metacam 25 µl/10g) and the temperature was maintained at 36º Celsius using a heating pad and temperature probe. The mouse was then secured on a stereotaxic frame (Angle Two, Leica Biosystems. After incision, the skull was roughened and a small craniotomy was made away from the site of implantation and a ground pin inserted. Next a ~1 mm diameter craniotomy was made behind lambda using a dental drill (Osada Electric, Japan) to expose the transverse sinus. Connective tissue and dura were removed and the vessel pushed forwards. A silicon probe (4-shank Neuropixels 2.0, IMEC)(78), coated with DiI (Invitrogen), was slowly inserted at a 30º angle, ~1 µm / s for 3 mm using a micromanipulator (Luigs and Neumann). Once in place, the probe was attached to the skull using UV-curing dental cement (RelyX Unicem 2 Automix, 3M), reinforced with dental cement (Super Bond C&B, Sun Medical). The ground was then connected to the pin and the implant reinforced with dental cement. A protective casing was attached and cemented in place using UV-curing cement. Following surgery, the mouse was returned to its home cage to recover on a heat pad. Following recovery, the mouse was briefly anaesthetised and the headstage connected and fixed in place.

Data acquisition and visual stimulation. We used the same behavioural arena and testing protocols as described previously (4). Briefly, each mouse was transferred in their home cage to the experimental room and given at least 5 min to acclimatise to the room under low light conditions, before being temporarily anesthetised to connect the Neuropixels interface cable to the headstage. A rotary joint was used to allow cable de-rotation during recording (AHRJ-OE_1x1_PT__FC_24, Doric Lenses). Mice were allowed to explore their home cage to acclimatise to the attachment and recover from anaesthesia, before being transferred to the behavioural arena (red Perspex with a white opaque floor, L:50 cm × W:20 cm × H:28 cm) with a red dome (10 cm diameter) or rectangular shelter (10 cm x 20 cm x 10 cm, both red Perspex). For chronic silicon probe recordings, data were acquired at 30 kHz using Neuropixels probes connected to a PXIe-1000 card (IMEC) housed in a PCIe-1071 chassis together with a PXIe-8398 remote control module connected to a PCIe-8398 host interface card (National Instruments). Data were acquired using SpikeGLX software. The 30 Hz TTL signal to trigger camera frame acquisition was recorded on the probe synchronisation channel.

Data pre-processing and analysis. Electrophysiological data were pre-processed using SpikeInterface (79) to implement phase shift correction, global common median refer-

encing and bandpass filtering (300–6000Hz) and run spike sorting using Kilosort 2.5 (78, 80). Units were manually curated using Phy (https://github.com/cortex-lab/phy) to exclude noise and multiunit activity from further analysis. Each unit included in the analysis passed manual curation, had a signal to noise ratio greater than 2, an inter-spike interval violations ratio less than 0.15 and an amplitude cut-off proportion less than 0.1. Baseline spontaneous firing rates were measured from a 180 second window during free exploration of the behavioural arena.

Histological quantification of probe tracks. To determine the exact location of probe tracks and single unit positions in the brain, a 3D whole-brain dataset was obtained at the end of each experiment using serial section microscopy. Mice were anaesthetised with pentobarbital (final volume: 100–150 µl, diluted 1:3 in saline), transcardially perfused with 10–20 ml of 0.1 M PBS followed by 10 ml of 4% PFA (diluted in PBS). Brains were retrieved and fixed overnight in 4 % PFA before being transferred to PBS. Brains were then mounted in 5% agar, glued to a microscope slide and transferred to a serial section two-photon microscope (81, 82). The microscope was controlled using ScanImage (v5.6, Vidrio Technologies) with BakingTray, a custom software wrapper for setting imaging parameters (github.com/SainsburyWellcomeCentre/BakingTray, doi.org/10.5281/zenodo.3631609). The agar block was submerged in 50 mM PBS and the brain was imaged using a 920nm laser (Mai Tai eHP DS, Spectra-Physics). Dichroic mirrors and bandpass filters were used to separate red, green and blue (background channel) signals, detected using three PMT channels (R10699, Hamamatsu Photonics with DHPCA-100 amplifiers, Femto Messtechnik). The entire brain was imaged with 2–4 optical sections per 50 µm physical section. Images were assembled following acquisition using StitchIt (github.com/SainsburyWellcomeCentre/StitchIt, zenodo.org/badge/latestdoi/57851444) and brains were then registered to a standardised atlas containing the individual columns of the PAG (83). DiI tracks were segmented using Brainreg-Segment (84, 85) and the depth of units along the probe used to determine their coordinates in atlas space. Only units located in the dPAG were included in the analysis.

General data analysis. Data analysis was performed using Python 3.6 and 2.7 using custom-written routines, unless stated otherwise. Code will be made available upon request. Data are reported as mean ± SEM unless indicated otherwise. Statistical comparisons were made as stated in the text. Data was tested for normality, and parametric or non-parametric tests were chosen accordingly and as stated in the text. Statistical tests were made in SciPy Stats and Prism 9 (GraphPad). In all cases, $p < 0.05$ was the statistical threshold. No statistical methods were used to predetermine sample sizes, and are similar to those reported in previous publications (31, 51). The experimenters were not blinded to the conditions of the experiment during data acquisition and analysis.

Supplementary Figures

Figure S1. Electrophysiological properties of VGAT⁺ dPAG neurons. A. Confocal stack of a biocytin-filled VGAT⁺ neuron in the dPAG. Arrows indicate the neuron's orientation: D – dorsal, V – ventral, M – medial, L – lateral. **B.** Left: Membrane potential traces of a VGAT⁺ dPAG neuron in response to current injections of different amplitudes. Right panel: summary graph of the current-firing rate relationship of VGAT⁺ dPAG neurons (shaded area is s.e.m; n = 29 cells, N = 8 mice). **C.** The mean resting membrane potential (-66.31 ± 0.69 mV), input resistance (0.55 ± 0.033 GΩ) and membrane time constant (25.61 ± 1.79 ms) were calculated for the same neurons as in B. **D.** Electrical stimulation of the deeper layers of the superior colliculus (dSC; left panel: schematic of the experimental configuration) evokes excitatory postsynaptic currents in all tested VGAT⁺ dPAG neurons with a mean peak amplitude of -37.07 ± 5.78 pA (peak amplitude of first pulse; middle panel) and a mean paired pulse ratio of 1.2 ± 0.11 (right panel) (n = 13 cells, N = 4 mice). Box-and-whisker plots show median, IQR and range, as well as mean (+) and individual data points.

Figure S2. Anatomical location of loose cell-attached recorded cells in acute coronal slices of the dPAG. A. Schematic of a sagittal brain section indicating the rostrocaudal extend of recorded neurons in the PAG (left panel), and an image of an acute coronal midbrain slice with the pipette at its recording location in the dPAG (right panel). Recordings were made from identified VGAT[+] (filled circles, dark and light red) and VGluT2[+] neurons (filled circles, dark and light blue; see legend). **B.** Superimposition of individual recorded cells along the rostrocaudal axis of the PAG, coordinates are in mm and from bregma. Mouse brain images adapted from (Paxinos and Franklin, 2001).

Figure S3. Firing properties of VGAT⁺ dPAG neurons and IPSCs *in vitro*, as well as firing rates *in vivo*. A. Example ISI histogram of a regularly firing VGAT⁺ neuron recorded *in vitro*. **B.** Summary histogram of the mean CV_2 of all VGAT⁺ dPAG cells recorded in control conditions. **C.** Example trace of ChR2-evoked IPSCs recorded in a presumptive dPAG excitatory neuron (stimulation protocol: 1ms duration, 5 pulses, 20Hz) before (blue trace) and after application of 10 μM CNO (orange trace) to silence VGAT⁺ dPAG neurons. **D.** Summary graph of the normalised evoked IPSC amplitude in control conditions and after 10 μM CNO application (control: n = 5 cells; 10 μM CNO: n = 4 cells; N = 2 mice). **E.** Example image of a coronal midbrain slice with the tracks of a 4-shank Neuropixels 2.0 probe made visible through DiI staining of the shanks prior to insertion into the brain. **F.** Example firing of a single unit recorded in the dPAG *in vivo* during exploration with the mean spike rate (histogram) over time shown for 10 randomly sampled 1 s time intervals from within a 60 s time window. The grey dotted line indicates the mean spike rate (19.96 Hz) for the same neuron. **G.** Average spike rate of all dPAG single units recorded *in vivo* (note that the y-axis is log scale). Each plotted data point (black filled circles) is the mean firing rate of a single unit. Box-and-whisker plots show median, IQR and range, as well as mean (+) and individual data points.

Figure S4. Gradient index lens placements in the dPAG for *in vivo* calcium imaging recordings using a miniaturised microscope in freely moving mice. A. PFA-fixed, coronal section stained with DAPI showing a recovered GRIN lens track in the dPAG (top, scale bar = 500 μm) and a zoom-in on the imaged region of interest with GCaMP6s-expressing VGAT[+] neurons (bottom, scale bar = 100 μm, dashed line = approximate imaging plane). **B.** Placements of recovered GRIN lenses along the rostrocaudal axis of the PAG (N = 7 out of 8 experiments). The images in A are from animal BF118.2.

(figure legend, see following page)

(Previous page.) **Figure S5. A small fraction of VGAT$^+$ calcium neurons show activity modulation by movement during exploration, escape-active neurons belong to two clusters present across animals, and are active during spontaneous escapes. A.** Single-neuron calcium signal (top) and animal's speed (bottom) over time, showing calcium events (pink lines) detected using the filtered signal (black line). **B.** Distribution of cells' locomotion modulation index of activity during exploration for populations that are inactive and active at escape. Dark coloured dots denote significantly modulated cells, determined by bootstrapping with resampling (95 % confidence interval significantly different from zero). Escape-active population contains a larger fraction of significantly modulated cells (8/123 and 7/123 cells, positive and negatively-modulated respectively) than the inactive population (1/76 and 2/76). **C.** All escape trials from one animal from one example session, showing animal's position along the long-axis of the rectangular arena over time, aligned to stimulus onset. **D.** Heatmap of the Z-scored, single trial calcium signal of one example neuron during the escape trials shown in C, with the peak calcium signal aligned to stimulus-onset, startle-response, escape turn, escape run and escape stop. **E.** and **F.**, same as C and D for another animal, session and neuron. **G.** PC scores for mean Z-scored, time-warped signal during escape for each neuron, coloured by K-means cluster ID (n = 123 neurons, N = 8 animals). **H.** Histogram of neurons assigned to clusters in G for each mouse. **I.** Mean time-warped signal for each neuron (same data as in Fig. 2G) for each mouse. **J.** Distribution of cells' locomotion modulation index of activity for the two escape-active clusters. Dark coloured dots denote significantly modulated cells, determined by bootstrapping with resampling (95 % confidence interval significantly different from zero). Both clusters contain a similar fraction of positively and negatively modulated cells (cluster 1, 14/62 and 3/62 cells; cluster 2, (4/61 and 4/61), positive and negative modulation index respectively). Negatively modulated cells in cluster 1 have a significantly lower modulation index than those in cluster 2 (P = 0.022, two-tailed t-test). **K.** Mean cluster population activity and animal speed trace for escape trials with slow peak run speed (top; mean peak speed 17.4 ± 1.3 cm/s) and fast running events (i.e., darting) during exploration by the same mice (bottom; mean peak speed, 21.8 ± 1.1 cm/s), aligned to the onset of running. Cluster 1 activity rises significantly during escape running but not exploratory running, while cluster 2 activity changes are not detected in either behaviour (*cluster 1, slow escapes*, mean signal -0.42 ± 0.22 Z and 0.22 ± 0.21 Z in the 2 s before and after run onset respectively, two-tailed paired t-test, P = 0.047, n = 12 neurons, 11 trials, N = 4 mice; *cluster 2, slow escapes*, mean signal -0.04 ± 0.26 Z before and 0.4 ± 0.16 Z after, two-tailed paired t-test, P = 0.16, n = 18 neurons, 10 trials, N = 4 mice; *cluster 1, runs during exploration*, mean signal -0.06 ± 0.14 Z before and -0.11 ± 0.15 Z after, two-tailed paired t-test, P = 0.80; n = 18 neurons, 66 trials, N = 4 mice; *cluster 2, runs during exploration,* mean signal -0.08 ± 0.10 Z before and -0.03 ± 0.10 Z after, two-tailed paired t-test, P = 0.72; n = 26 neurons, 60 trials, N = 4 mice). **L.** Mean cluster population activity and animal speed trace aligned to the onset of spontaneous escapes to the shelter in the absence of sensory stimulation (cluster 1, mean signal 0.05 ± 0.08 Z and 0.47 ± 0.08 Z in the 2 s before and after escape onset respectively, two-tailed t-test p < 0.001 , n = 51 neurons, 74 trials, N = 7 mice; cluster 2, mean signal 0.16 ± 0.08 Z before and 0.23 ± 0.07 Z after, two-tailed t-test p = 0.55, n = 55 neurons, 69 trials, N = 7 mice).

Figure S6. Population activity during escape is not significantly correlated with metrics of escape vigour. A. and **B.** Single trial mean peak F/F during escape for cluster 1 (**A.**) and 2 (**B.**) against maximum speed of the escape trial. **C.** and **D.** Area under curve of F/F during escape for cluster 1 (**C.**) and 2 (**D.**) against average acceleration from escape onset to time of maximum speed during escape. **E.** and **F.** Area under curve of F/F during escape for cluster 1 (**E.**) and 2 (**F.**) against average deceleration from time of maximum speed to escape termination. N = 100 trials, 8 animals; Pearson's r and p-values are indicated for each measure in the respective panel.

Figure S7. Optogenetic manipulation of VGAT⁺ dPAG neurons does not alter speed of movements during exploratory behaviour. **A.** Schematic of the experimental design and optogenetic stimulation protocols (ChR2: 50 Hz, 3 s; iChloC: continuous, 3 s). **B.** Coronal slice stained with DAPI (blue) with bilateral expression of AAV2/2-DIO-ChR2-mCherry (red) localised to the dPAG and optic fibre tracks. Scale bar = 1 mm. **C.** Bilateral optic fibre placements along the rostrocaudal axis of the dPAG are shown for ChR2 (blue, top; N = 5 mice) and iChloC experiments (orange, bottom; N = 4 mice). Rostrocaudal position from Bregma, based on Paxinos and Franklin. **D.** Example speed raster plot during optogenetic stimulation of VGAT⁺ dPAG neurons during exploratory movement. Grey bar on the right side of the histogram indicates no escapes. Laser duration is shown in blue. **E.** Summary graph of mean speed before and during laser stimulation (ChR2; N = 5 mice, n = 92 trials). **F.** Animals escape to threat stimuli when optogenetic stimulation of VGAT⁺ dPAG neurons precedes and ends immediately before threat stimulation (N = 4 mice, P = 0.44, Mann-Whitney test between probability of escape when laser precedes vs no laser condition). **G.** The same as D, for optogenetic inhibition. Laser duration shown in orange. **H.** Mean speed before and during laser stimulation for optogenetic inhibition (iChloC; N = 4 mice, n = 108 trials).

Figure S8. Optogenetic inhibition of VGAT$^+$ neurons during escape increases total escape distance. A. Left: Moving the shelter location in the arena away from the arena wall leads to sensory-evoked escapes to the new shelter location (indicated in light red). Control trials (Laser OFF) of one mouse during escape to shelter with normalised tracking coordinates over time (top panel) and the corresponding speed traces (bottom panel). Right: Same as A, for Laser ON trials with light-activation of iChloC after escape onset. The same animal displays continued movement after reaching the shelter (black arrow). **B.** Summary plot of the total escape distance for Laser OFF trials (10 trials) and Laser ON trials (13 trials) showing a significant increase during light-activation of iChloC (N = 2 mice). Graph is showing the mean ± s.e.m and individual trials, separated by mouse.