Tgif1-deficiency impairs cytoskeletal architecture in osteoblasts by suppressing PAK3

signaling

Simona Bolamperti[1], Hiroaki Saito[1,2,3], Sarah Heerdmann[1], Eric Hesse[1,2,3,#], Hanna

Taipaleenmäki[1,2,3,#,*]

[1]Molecular Skeletal Biology Laboratory, Department of Trauma, Hand and Reconstructive

Surgery, University Medical Center Hamburg-Eppendorf, Martinistraße 52, 20246 Hamburg,

Germany

[2]Institute of Musculoskeletal Medicine, University Hospital, LMU Munich, Fraunhoferstraße 20,

82152 Planegg-Martinsried, Germany

[3]Musculoskeletal University Center Munich, University Hospital, LMU Munich, Fraunhoferstraße

20, 82152 Planegg-Martinsried, Germany

[#]These authors contributed equally to this work

[*]Correspondence: hanna.taipaleenmaeki@med.uni-muenchen.de

Summary

Osteoblast adherence to bone surfaces is important for remodeling of the bone tissue. This study

demonstrates that deficiency of TG-interacting factor 1 (Tgif1) in osteoblasts results in altered cell

morphology, reduced adherence to collagen type I-coated surfaces, and impaired migration

capacity. Tgif1 is essential for osteoblasts to adapt a regular cell morphology and to efficiently

adhere and migrate on collagen type I-rich matrices *in vitro*. Furthermore, Tgif1 acts as

transcriptional repressor of p21-activated kinase 3 (PAK3), an important regulator of focal

adhesion formation and osteoblast spreading. Absence of Tgif1 leads to increased PAK3

expression, which impairs osteoblast spreading. Additionally, Tgif1 is crucial for osteoblast

recruitment and activation of bone surfaces in the context of bone regeneration and in response to

parathyroid hormone 1-34 (PTH 1-34) treatment *in vivo*. These findings provide important novel

insights in the regulation of the cytoskeletal architecture of osteoblasts.

Keywords: osteoblast spreading, Tgif1, PTH, bone regeneration, PAK3

1 **Introduction**

2 Bone remodeling is a highly coordinated and continuously ongoing process involving osteoblasts,

3 osteoclasts, and osteocytes, that interact with each other and function in concert to maintain bone

4 mass and preserve the integrity of the skeletal system (Baron and Hesse, 2012). This process is

5 essential for bone development, growth, maintenance and repair as well as for the functionality of

6 pharmacological interventions (Kenkre and Bassett, 2018). Osteoblasts, derived from

7 mesenchymal stem cells, play a pivotal role in synthesizing and depositing collagen type I-rich

8 bone matrix (Ponzetti and Rucci, 2021). To achieve this, osteoblasts undergo changes in cell shape

9 and migrate on bone surfaces to lay down matrix at places where new bone needs to be formed,

10 necessitating the regulation of cell movement and shape acquisition (Jones and Boyde, 1977;

11 Eleniste et al., 2014; Thiel et al., 2018).

12 Migration of osteoblasts involves movement from their origin to specific locations within the

13 bone tissue and occurs during both embryonic development and adult bone remodeling. Multiple

14 factors influence osteoblast migration, including growth factors, cytokines, signaling pathways

15 like bone morphogenetic proteins (BMPs) and transforming growth factor-beta (TGF-β) as well

16 as interactions with other cells and the extracellular matrix (Thiel et al., 2018). These signals not

17 only promote migration but also regulate the expression of adhesion molecules, cytoskeletal

18 rearrangements, and cellular contractility.

19 Osteoblasts exhibit various morphological shapes based on their differentiation stage and the

20 local microenvironment, ranging from elongated spindle-like morphology to cuboidal appearance

21 (Tsuji et al., 2022). The cytoskeleton, primarily composed of actin filaments, microtubules, and

22 intermediate filaments, plays a central role in determining cell shape. Actin filaments, among other

23 cytoskeletal elements, are involved in cell polarization, protrusion formation, and contractility

24 during migration (Schaks et al., 2019; Tang and Gerlach, 2017; Etienne-Manneville, 2004).

25 Signaling molecules, including Rho GTPases such as RhoA, Rac1, and Cdc42, regulate the

26 cytoskeleton by controlling actin polymerization and organization, contributing to the formation

of structures like lamellipodia, filopodia, and stress fibers (Schaks et al., 2019). These structures are essential for cell migration, enabling attachment to the extracellular matrix and generating the necessary force for movement. Integrins, transmembrane proteins mediating cell-extracellular matrix interactions, also play a critical role in osteoblast migration and cell shape regulation. They anchor cells to the matrix, transmit signals regulating migration and cytoskeletal dynamics, and activate intracellular pathways affecting cell shape, migration speed, and adhesion strength (Kechagia et al., 2019; SenGupta et al., 2021; Hood and Cheresh, 2002; Thiel et al., 2018). Thus, osteoblast migration and cell shape acquisition are interconnected processes crucial for bone development, growth, remodeling, and repair. Further investigation of the mechanisms underlying osteoblast migration and cell shape regulation is therefore important to better understand bone mass maintenance, treatment of bone-related disorders, and bone repair.

Fractures occur in the context of high-energy trauma but also due to bone fragility and represent a severe damage of bone integrity, requiring subsequent tissue regeneration. In response to fracture, inflammatory cells including neutrophils and macrophages are recruited to the site, releasing signaling molecules such as cytokines and growth factors. These molecules create a favorable environment for osteoblast migration and bone formation (Einhorn and Gerstenfeld, 2015). Osteoblast precursors derived from mesenchymal stem cells, or the periosteum migrate alongside sprouting vessels towards the fracture gap and differentiate into mature bone-forming osteoblasts (Maes et al., 2010). Once osteoblasts have reached the fracture site, production and deposition of new bone matrix is initiated to facilitate fracture repair (Maes et al., 2010; Einhorn and Gerstenfeld, 2015). During this process, chemotactic signals released by surrounding tissues and cells, including platelet-derived growth factor (PDGF), transforming growth factor-beta (TGF-β), and bone morphogenetic proteins (BMPs) guide osteoblast migration (Dirckx et al., 2013; Einhorn and Gerstenfeld, 2015; Thiel et al., 2018). Furthermore, migration of osteoblasts is facilitated by the dynamic rearrangement of the cytoskeleton, primarily composed of actin filaments (Thiel et al., 2018). Structures like lamellipodia and filopodia, enriched with actin,

enable osteoblasts to extend and protrude in the direction of migration, interacting with the extracellular matrix and providing necessary traction (Casati et al., 2015; Jafari et al., 2019).

Movement of osteoblasts is an important but often underestimated component of the pharmacologically induced gain in bone mass in response to the treatment with bone anabolic drugs. Teriparatide, a recombinant form of the first 34 amino acids of human parathyroid hormone (rhPTH1-34; hereafter PTH), is a bone anabolic drug used to treat severe osteoporosis, characterized by low bone mineral density and an increased fracture risk (Neer et al., 2001; Black and Rosen, 2016). PTH treatment stimulates bone formation by increasing the number, differentiation and activity of osteoblasts (Pettway et al., 2008; Silva et al., 2011; Ogita et al., 2008). In addition, PTH promotes the recruitment of osteoblast precursor cells and activates quiescent lining cells on bone surfaces, transforming them into active matrix-forming osteoblasts (Nishida et al., 1994; Dobnig and Turner, 1995; Kim et al., 2012). Furthermore, PTH increases the migration of mesenchymal precursor cells (Lv et al., 2020) and augments bone remodeling (Saito et al., 2019), which attracts osteoblasts (Dirckx et al., 2013). During osteoblast migration, cytoskeletal dynamics promotes actin filament remodeling and the formation of filopodia and lamellipodia (Lomri and Marie, 1990). These structures are essential for cell attachment to the extracellular matrix during migration. By enhancing cytoskeletal dynamics, PTH facilitates osteoblast movement (Thiel et al., 2018). PTH also increases the expression and activation of integrins, which mediate interactions between cells and the extracellular matrix (Gronthos et al., 2001; Kaiser et al., 2001). These combined pharmacological effects of PTH treatment result in an increase in bone formation and bone mineral density, which improves bone strength and ultimately reduces fracture risk (Neer et al., 2001; Black and Rosen, 2016).

Recently, we reported the important role of the homeodomain protein TG-interacting factor 1 (Tgif1) in osteoblast differentiation, activity and bone formation(Saito et al., 2019). Furthermore, Tgif1 was identified as PTH target gene, and in the absence of Tgif1, PTH treatment failed to increase bone mass in mice(Saito et al., 2019). These findings demonstrate the importance of Tgif1

as novel regulator of bone remodeling and emphasizes its essential involvement in mediating the

full bone anabolic effect of PTH treatment. Building upon these observations, we investigated the

role of Tgif1 in osteoblast morphology, adherence and migration. Our findings demonstrate that

Tgif1-deficient osteoblasts display an altered morphology, reduced adherence to collagen type I-

coated surfaces, impaired migration capacity and decreased spreading compared to control cells.

These deficits are associated with a compromised formation of focal adhesions and an upregulated

expression of p21-activated kinase 3 (PAK3). Mechanistically, Tgif1 exerts control over PAK3

expression through transcriptional repression. Thus, elevated PAK3 levels contribute to the

observed defects in osteoblast spreading of Tgif1-deficient cells. By using translational

approaches, we demonstrate that in the absence of Tgif1, the activation of bone surfaces by

osteoblasts in response to bone repair and PTH treatment is diminished. Furthermore, we

uncovered that PTH facilitates osteoblast spreading via Tgif1-PAK3 signaling.

Collectively, these findings increase the knowledge on processes governing osteoblast

morphology, adherence, and migration. These novel insights are important to better understand

mechanisms underlying bone remodeling and repair as well as the pharmacological effects of PTH

treatment.

Results

Tgif1-deficiency alters osteoblast morphology, adherence and migration

Recently, we reported that in Tgif1-deficient mice the number and activity of osteoblasts are

reduced, which contributes to a low turnover bone remodeling (Saito et al., 2019). Further

histological examination revealed that bone surfaces were scarcely occupied by small and flat

osteoblasts in mice bearing a germline deletion of Tgif1 (Fig. 1A) or a deletion of Tgif1 targeted

to osteoblasts (Fig. 1B) compared to control littermates. This finding suggested that Tgif1-

deficient osteoblasts might, in addition to an impaired bone matrix-producing capacity (Saito et

al., 2019) be compromised in acquiring a physiological osteoblast morphology. This deficit could

contribute to the low turnover bone phenotype of Tgif1-deficient mice.

Inactive and early-stage osteoblasts are rather small and flat in their morphological appearance

but enlarge and adopt a more cuboidal shape upon activation. Active osteoblasts produce

extracellular matrix and have the capacity to migrate to sites of bone remodeling and repair (Dirckx

et al., 2013). These features require the ability to adhere to surfaces and to change cell morphology

to facilitate cell migration. In support of our hypothesis that Tgif1-deficiency compromises

morphological plasticity of osteoblasts, calvarial- and long bone-derived osteoblasts obtained from

Tgif1-deficient mice were impaired in their ability to attach to collagen type I-coated surfaces

throughout a 4-hour time-course (Fig. 1C-E). To determine the ability of Tgif1-deficient

osteoblasts to migrate, we performed live cell video microscopy of osteoblasts seeded on collagen

type I-coated surfaces. This analysis revealed that osteoblasts lacking Tgif1 were impaired in their

migration capacity reflected by a shorter track length, a reduced velocity and a less straight and

therefore more meandering track path (Fig. 1 F-H, Fig. S1A). Thus, these *in vivo* and *in vitro* data

indicate that Tgif1 is important for osteoblasts to adhere and migrate on collagen type I-rich

surfaces. To confirm these findings in a 3-dimensional functional context, we obtained long bone

osteoblasts from Tgif1-defcient mice and control littermates and seeded them on a collagen type I

matrix placed over a porous membrane (Fig. 1I). In support of the previous observations,

osteoblasts lacking Tgif1 adhered less to collagen type I matrix and were impaired in their

capacities to migrate into the matrix compared to control cells (Fig. 1J). The experiment also

confirmed that Tgif1-deficient osteoblasts were smaller in size and rather flat in morphology (Fig.

1K). Collectively, these observations demonstrate that Tgif1 is indispensable for osteoblasts to

adapt a regular cell morphology and to adhere and migrate on collagen type I-rich matrices *in vitro*

and on bone surfaces *in vivo*.

Tgif1-deficient osteoblasts are impaired in spreading and in forming focal adhesions

Adherence on osseous matrices and migration alongside bone surfaces towards sites of remodeling

or repair are crucial functional features of osteoblasts (Dirckx et al., 2013; Thiel et al., 2018). Upon

adherence and migration, osteoblasts undergo morphological changes and spread on bone surfaces,

thereby decreasing their sphericity, a measure of how closely an object resembles the round shape

of a sphere (Cruz-Matías et al., 2019).

To determine the ability of Tgif1-deficient osteoblasts to spread, calvarial osteoblasts were

isolated from *Tgif1$^{-/-}$* mice and control littermates and subject to spreading on collagen type I-

coated slides for 60 minutes in the presence of Calcein-AM, a dye that is fluorescent in living cells.

Compared to control cells, *Tgif1$^{-/-}$* osteoblasts were significantly impaired in their ability to spread

during the experiment (Fig. 2A, B).

Cell spreading is associated with the assembly and disassembly of actin filaments to form cell

protrusions. Cell protrusions facilitate anchoring to the extracellular matrix via focal adhesion

complexes (Cronin and Demali, 2021). To determine whether Tgif1 is implicated in actin filament-

and focal adhesion assembly, *Tgif1$^{-/-}$* and control osteoblasts were stained for phalloidin to

visualize actin filaments and Paxillin, a major component of focal adhesions. Consistently, after

60 minutes of culture on collagen type I-coated surfaces, Tgif1-deficient osteoblasts were less

spread and therefore had a greater sphericity (Fig. 2C, D). Furthermore, Tgif1-deficient osteoblasts

that underwent spreading formed less cellular processes and focal adhesions compared to control

cells (Fig. 2C, E).

To confirm the findings obtained from osteoblasts bearing a germline deletion of *Tgif1*, we

transiently silenced *Tgif1* in cells of the OCY454 cell line, a cell model system resembling motile

osteocytes and mature osteoblasts (Spatz et al., 2015), using siRNA (Fig. S1B). Consistent with

the findings made by genetic deletion of *Tgif1*, siRNA-mediated silencing of Tgif1 in OCY454

cells resulted in an impaired cell spreading with a consecutive higher cell sphericity and a decrease

in focal adhesion formation (Fig. 2F-H).

PAK3 expression is increased in the absence of Tgif1 and impairs focal adhesion formation and

osteoblast spreading

To investigate the molecular mechanisms underlying the impaired ability of $Tgif1^{-/-}$ osteoblasts to

undergo morphological changes, expression and activation of the major focal adhesion

components including FAK, Integrin β1, Src, talin, paxillin, p38 and LRG5 was quantified during

cell spreading. However, no major changes were identified in the expression or activation of these

molecules in the absence of Tgif1 (Fig. S2A).

Next, we examined the localization of Tgif1 to determine a possible interaction of Tgif1 with

components of focal adhesions or actin filaments. Immunocytochemistry revealed that Tgif1 is

predominantly localized in the nucleus and to some extend in the cytoplasm (Fig. S2B). However,

cytoplasmic Tgif1 did not interact with focal adhesions, because no co-localization of eGFP-Tgif1

neither with paxillin nor with talin (Fig. S2B) was observed.

Given the predominantly nuclear localization of Tgif1 (Fig. S2B) and its established function

as transcriptional repressor (Saito et al., 2019)(Wotton et al., 1999), we investigated whether

absence of Tgif1 might alter the expression of genes involved in actin cytoskeletal assembly and

re-arrangement as prerequisite of cell adhesion and migration. First, we quantified the expression

of members of the cell division control protein 42 homolog (Cdc42), which are small GTPases of

the Rho family that participate in the control of multiple cellular functions including cell migration

and cell morphology (Hirsch et al., 2001). However, expression of none of the Cdc42 family

members Cdc42es2, Cdc42ep1, Cdc42ep2 or Cdc42ep4 was changed in the absence of Tgif1

during osteoblast adhesion and spreading (Fig. S3 A-D). These data indicate that the impaired

spreading of Tgif1-deficient osteoblasts is unlikely caused by Cdc42 family members and that

other factors are implicated in this process. We therefore quantified the expression of p21-

activated-kinase (PAK) family members, who have been shown to play a role in actin nucleation

in neurons (Kreis and Barnier, 2009). While expression of PAK 1, 2 and 4 was unchanged in

Tgif1-deficient osteoblasts during adhesion and spreading (Fig. S3 E-G), expression of PAK3 was

strongly increased in Tgif1-deficient osteoblasts compared to control cells at the mRNA and

protein level prior to and 60 min after adhesion (Fig. 3 A and B). Confirming this observation *in*

vivo, expression of PAK3 mRNA showed a trend towards an increase in bones from *Tgif1*$^{-/-}$ mice

compared to control littermates (Fig. 3C). These findings suggest that the impaired adhesion and

spreading of Tgif1-deficient osteoblasts might be related to a deregulated abundance of PAK3.

To further investigate the regulation of PAK3 expression by Tgif1, we performed an *in silico*

analysis of the PAK3 promoter and identified 3 putative Tgif1 binding sites (Fig. S4A). Since only

one site was species-conserved between rat and mouse, we verified the binding of Tgif1 to this

site using chromatin immunoprecipitation. Indeed, Tgif1 bound to the predicted promoter binding

site and to a site of the RAR promoter as positive control (Zhang et al., 2009), but not to a DNA

sequence lacking predicted Tgif1 binding site as negative control (Fig. 3D), suggesting a

regulation of the PAK3 promoter activity by Tgif1. To address this question, Tgif1 expression was

silenced in OCY454 osteoblast-like cells using siRNA followed by transfection of the cells with a

2.3kb fragment of the rat PAK3 promoter upstream of the luciferase gene. Silencing of Tgif1 lead

to a significant increase in luciferase activity compared to control (Fig. 3E) demonstrating that

Tgif1 is a transcriptional repressor of the PAK3 promoter.

To determine if an elevated abundance of PAK3 due to Tgif1-deficiency impairs osteoblast

spreading, PAK3 expression was silenced in Tgif1-deficient OCY454 cells using siRNA (Fig.

S4B). Indeed, attenuating PAK3 expression restored the Tgif1 deficiency-mediated impaired cell

spreading (Fig. 3F). Since cells are less spherical upon spreading, PAK3 silencing reduced the

Tgif1 deficiency-dependent gain in cell sphericity and restored the number of focal adhesions per

cell that were reduced by Tgif1 deficiency to the level of control (Fig. 3 G and H). Together, these

findings demonstrate that in the absence of Tgif1, the PAK3 promoter is de-repressed, leading to

an increased transcriptional activity and PAK3 expression. A higher abundance of PAK3 in turn

suppresses focal adhesion formation and osteoblast spreading.

Tgif1 expression increases during osteoblast spreading via activation of ERK1/2 and AP1

signaling pathways

To further investigate the role of Tgif1 during osteoblast spreading, we quantified Tgif1 expression

during this process. Both, Tgif1 mRNA and protein expression were significantly increased within

20 minutes of spreading with a further increase after 60 minutes in calvarial osteoblasts (Fig. 4A

and B) and in OCY454 cells (Fig. 4C and D). To determine if the increase in Tgif1 abundance is

transcriptionally regulated, osteoblasts were treated with the transcription inhibitor Actinomycin-

D during spreading, which fully prevented the increase in Tgif1 mRNA expression (Fig. 4E).

Since binding of cells to collagen I-coated surfaces activates the ERK1/2 pathway (Emerson et

al., 2009; Fincham et al., 2000), we investigated the potential implication of ERK1/2 signaling in

the increase in Tgif1 gene expression during osteoblast spreading. Experimentally, osteoblasts

were treated with an ERK1/2 inhibitor, which greatly attenuated ERK1/2 phosphorylation after 20

min and 60 min of osteoblast spreading (Fig. 4F). Although ERK1/2 inhibition prevented the

increase of Tgif1 mRNA expression during the first 20 min of osteoblast spreading, it failed to

suppress Tgif1 mRNA expression 60 min after initiation of osteoblast spreading (Fig. 4G). This

finding suggests that the increase in Tgif1 mRNA expression at early stages of osteoblast spreading

is mediated by ERK1/2 signaling while it is independent of ERK1/2 signaling at later stages of

osteoblast spreading.

To elucidate the molecular mechanisms underlying the activation of Tgif1 transcription at later

stages of osteoblast spreading, we performed progressive truncations of the 2.2 kb fragment of the

human Tgif1 promoter (Saito et al., 2019). While the 2.2 kb and 1.7 kb fragments of the human

Tgif1 promoter were fully activated after 60 minutes of osteoblast spreading, no activation of the

1.2 kb and 0.5 kb fragments was observed (Fig. 4H), suggesting that the regulatory promoter

region that is activated during osteoblast spreading must be located within the 500 bp between the

1.7 kb and the 1.2 kb promoter fragments.

Recently we reported the presence of an AP1 binding site in this region of the Tgif1 promoter

(Saito et al., 2019), suggesting that AP1 signaling might be implicated in this process. Since the

role of AP1 signaling in the context of cell spreading has not been fully elucidated, we first

determined the activation of an AP1-responsive promoter element during osteoblast spreading.

The findings demonstrate that AP1-signaling is activated 60 minutes after initiation of osteoblast

spreading (Fig. 4I), indicating a potential implication of AP1 signaling during advanced stages of

osteoblast spreading. To further test this hypothesis, osteoblasts were transfected either with a

plasmid encoding the wild-type Tgif1 promoter bearing an AP1 binding site or with a plasmid in

which the AP1 binding site was disabled by site-specific mutation (Saito et al., 2019). Although

the wild-type promoter was fully activated after 60 min of osteoblast spreading, the Tgif1 promoter

bearing the disabled AP1 binding site was not activated (Fig. 4J). These findings suggest that AP1

signaling is involved in the activation of Tgif1 gene expression during later stages of osteoblast

spreading.

Tgif1 deficiency leads to fewer and less active osteoblasts during bone regeneration

Bone has a regenerative capacity to heal upon fracture. In fulfilling this function, osteoblast

adherence and spreading are crucial features since osteoblasts migrate to sites of bone repair for

subsequent matrix production and regeneration (Thiel et al., 2018; Dirckx et al., 2013). Fracture healing involves various cell types and cellular processes (Einhorn and Gerstenfeld, 2015). One important component is the activation of periosteal cells to migrate alongside vessels that sprout into the fracture zone (Maes et al., 2010). Periosteal cells become active osteoblasts that produce a cartilage template, which then ossifies and forms new trabeculae to consolidate the fracture gap (Duchamp De Lageneste et al., 2018). Since this process involves adherence and spreading of osteoblasts, we hypothesized that Tgif1 might be important in the context of bone repair. To test this hypothesis, we performed open mid-shaft tibia fractures in 10-weeks old adult male mice bearing a germline deletion of Tgif1 and control littermates, followed by fracture stabilization using an intramedullary pin. Three weeks after fracture, newly formed trabeculae were densely occupied by cuboidal matrix-producing osteoblasts in control animals (Fig. 5 A-E). In contrast, in mice bearing a germline deletion of Tgif1, the callus was less ossified and the trabeculae in the repair zone were sparsely occupied by fewer, thinner and less-active osteoblasts (Fig. 5 A-E). These findings demonstrate that Tgif1 is important for bone regeneration. In the absence of Tgif1, osteoblasts are greatly impaired in the ability to migrate into the repair zone, to attach and spread on surfaces and to produce mineralized bone matrix.

Activation of bone surfaces by PTH is attenuated in the absence of Tgif1 in osteoblasts

Intermittent administration of PTH augments bone formation, leading to a remodeling-based increase in bone mass, bone mineral density and a consecutive decrease in fracture rate in humans (Neer et al., 2001; Taipaleenmäki et al., 2022; Baron and Hesse, 2012). This anabolic function requires multiple alterations at the cellular level. For instance, PTH induces the differentiation of mesenchymal precursor cells and reverts quiescent bone lining cells into active osteoblasts (Nishida et al., 1994; Dobnig and Turner, 1995; Kim et al., 2012). Furthermore, PTH increases the number and bone-forming capacity of mature osteoblasts (Mizoguchi and Ono, 2021; Jilka et al., 2009).

Recently, we identified Tgif1 as a PTH target gene in osteoblasts that is necessary for the full

PTH-mediated bone mass accrual (Saito et al., 2019). In addition, the site of the Tgif1 promoter

known to be activated by PTH signaling (Saito et al., 2019), is identical with the site identified

here that is activated by AP1 signaling during osteoblast adhesion (Fig. 4J). In this study we also

uncovered that Tgif1 is a physiological regulator of osteoblast adherence, spreading and migration.

Collectively, this evidence let us to propose that Tgif1 might also be implicated in the PTH-

mediated bone surface activation.

To determine whether Tgif1 is implicated in the PTH-mediated activation of bone surfaces by

osteoblasts, we performed histomorphometric analysis of bone sections from mice bearing a

germline deletion of Tgif1 and control littermates that were injected with PTH or vehicle control.

Consistent with the findings reported by others and in support of our hypothesis, PTH treatment

greatly increased the number of active osteoblasts and induces the acquisition of a cuboidal shape

and consequently the percentage of active bone surfaces (Kousteni and Bilezikian, 2008; Tam et

al., 1982), which all occurred at a much lesser extent in mice lacking Tgif1 (Fig. 6A and B). To

determine whether this effect is osteoblast-autonomous, we performed histomorphometric analysis

of bones obtained from mice in which Tgif1 deletion was targeted to mature osteoblasts using the

Dmp1-promoter (Bivi et al., 2012). Indeed, while PTH treatment induced osteoblast recruitment,

adaptation of a cuboidal morphology of matrix-producing osteoblasts and a profound activation of

bone surfaces by osteoblasts in control animals, these effects were much less pronounced in mice

lacking Tgif1 in osteoblasts (Fig. 6C and D). These findings therefore demonstrate that PTH-

dependent bone surface activation is attenuated in the absence of Tgif1 in osteoblasts.

PTH facilitates spreading of osteoblasts via Tgif1-PAK3 signaling

The observation that Tgif1 is crucial for osteoblast adherence, spreading and migration as well as

for the PTH-mediated activation of bone surfaces suggests that PTH may facilitate the spreading

of osteoblasts and therefore the decrease in cell sphericity via Tgif1. To test this hypothesis,

OCY454 cells were treated with PTH or vehicle control during spreading. Immunofluorescence staining and quantification of cell sphericity revealed that PTH supports osteoblast spreading (Fig. 7A and B). Furthermore, PTH treatment of cells in which Tgif1 expression was silenced failed to induce the spreading of osteoblasts (Fig. 7A and B), demonstrating that the PTH-mediated osteoblast spreading indeed depends on Tgif1.

Since PAK3 expression is increased in the absence of Tgif1 and impairs focal adhesion formation and osteoblast spreading, we investigated the possibility that PTH decreases PAK3 gene expression via Tgif1. Indeed, PAK3 mRNA expression was reduced upon PTH treatment (Fig. 7C). In addition, PTH treatment increased Tgif1 expression and reduced PAK3 protein abundance in control cells but not in osteoblasts in which Tgif1 expression was silenced by siRNA (Fig. 7D). Next, we examined osteoblast spreading upon PTH treatment in the absence of Tgif1 alone or in combination with loss of PAK3. Immunofluorescence staining and quantification of cell sphericity revealed that silencing Tgif1 expression impaired osteoblast spreading induced by PTH treatment, which was restored by the concomitant silencing of PAK3 expression (Fig. 7E and F). Furthermore, lack of both, Tgif1 and PAK3 prevented PTH-induced decrease in cell sphericity. These findings indicate that an increased abundance of PAK3 in response to Tgif1-deficiency impairs PTH-induced spreading of osteoblasts and that PTH facilitates osteoblast spreading via Tgif1-PAK3 signaling.

In summary, our results reveal that Tgif1 suppresses PAK3 expression in osteoblasts via the ERK1/2 and AP1 signaling pathways, which is crucial for osteoblasts to form focal adhesions, to spread on bone surfaces, and to migrate. This mechanism is implicated in bone regeneration and in the pharmacological effects of PTH treatment and is therefore of translational relevance.

Discussion

This study revealed that Tgif1-deficient osteoblasts exhibit an altered morphology, reduced adherence to collagen type I-coated surfaces, impaired migration capacity, and decreased spreading compared to control cells. These defects in Tgif1-deficient osteoblasts are associated with compromised focal adhesion formation and increased expression of PAK3. The findings further demonstrate that Tgif1 regulates PAK3 expression through transcriptional repression and that elevated PAK3 abundance contributes to the impaired osteoblast spreading observed in Tgif1-deficient cells. Additionally, Tgif1 is shown to participate in the activation of bone surfaces in the context of bone regeneration and PTH treatment since in the absence of Tgif1, bone regeneration- and PTH-induced activation of bone surfaces is attenuated. Mechanistically, PTH promotes osteoblast spreading via Tgif1-PAK3 signaling, with PAK3 expression being reduced by PTH treatment in a Tgif1-dependent manner. Overall, this study emphasizes the importance of Tgif1 in regulating osteoblast morphology, adherence and migration through the modulation of PAK3 expression, providing novel and translationally relevant mechanistic insights into the regulation of the cytoskeletal architecture of osteoblasts.

Both, bone repair and PTH treatment are frequent and clinically relevant events. Fractures often occur in the context of high energy accidents, however, bone fragility syndromes such as osteoporosis also cause fractures upon minimal or no trauma (Malluche et al., 2013; Löfman et al., 2009). Bone regeneration, a complex process involving osteoblast recruitment, matrix production, and tissue regeneration, is compromised in Tgif1-deficient mice. Activation of bone surfaces by osteoblasts, a critically important aspect of bone repair, is significantly reduced in the absence of Tgif1. These results indicate that Tgif1 is crucial for osteoblast recruitment to the site of repair and subsequent bone surface activation, which extends the established function of Tgif1 beyond the regulation of osteoblast differentiation and activity.

Albeit mechanistically distinct, bone repair and PTH treatment have in common that osteoblasts are an important component for the successful execution of bone regeneration and the PTH

pharmacological effects. In particular, the guided spatial relocation of activated osteoblasts is a critical feature to exert the respective functions at the right sites in a three-dimensional context. Since formation of new bone, regardless of whether in the cortical- or trabecular compartment, occurs in a timely but also in a spatially controlled manner to yield structurally supportive newly formed bone tissue.

Various types of cells and tissues of the local microenvironment produce a plethora of signaling factors, which are induced by fractures or PTH treatment. However, little is known about signaling pathways in osteoblasts that receive and integrate these diverse external cues to facilitate the adherence, spreading and movement of osteoblasts in a coordinated manner. In this context, this study identified the functional importance of PAK3 in Tgif1-deficient osteoblasts. PAK3 expression was increased in the absence of Tgif1, leading to impaired focal adhesion formation and reduced osteoblast spreading. Silencing PAK3 expression restored osteoblast spreading in Tgif1-deficient cells, demonstrating that PAK3 plays a critical role in mediating the effects of Tgif1 on osteoblast morphology. This work therefore contributes to the better understanding of osteoblast dynamics by identifying Tgif1 and PAK3 as downstream signaling regulators that are important for osteoblasts to fulfill motility. Furthermore, the work underscores the physiological relevance of spatial osteoblast dynamics as integral component of bone remodeling and maintenance of skeletal integrity.

Tgif1 recently emerged as a key determinant of osteoblast activity, bone mass maintenance and essential component of the bone anabolic PTH signaling pathway (Haider et al., 2018; Saito et al., 2019). This study also focused on investigating the role of PTH in regulating osteoblast spreading and revealed that PTH treatment facilitated osteoblast spreading via Tgif1-PAK3 signaling. PTH treatment reduced PAK3 expression and increased Tgif1 expression, leading to enhanced osteoblast spreading. Silencing Tgif1 expression impaired PTH-induced osteoblast spreading, which was restored by simultaneously silencing PAK3 expression. These new findings on PTH-Tgif1-PAK3 signaling provide novel insights into the function of PTH treatment of bone

remodeling and the implication of Tgif1 in this process. As part of being required for bone mass accrual in response to PTH treatment (Saito et al., 2019), Tgif1 also contributes to the PTH-induced osteoblast recruitment and subsequent movement to sites where new bone needs to be formed. This adds a spatial component to a functional feature and demonstrates the complexity of bone remodeling and PTH anabolic actions.

In summary, these findings highlight the pivotal role of Tgif1 in regulating osteoblast morphology, adherence, and migration. It also reveals the involvement of PAK3 in mediating the effects of Tgif1 on osteoblast spreading. The dysregulation of these processes due to Tgif1 deficiency has profound implications for bone remodeling, bone regeneration, and the pharmacological effects of PTH. Reaching a better understanding of the intricate interplay between PTH, Tgif1 and PAK3, provides valuable insights into the molecular mechanisms governing osteoblast function and could have translational relevance in bone-related disorders and therapies targeting bone health.

Materials and methods

Animal models. Mice with germ-line deletion of Tgif1 or loxP-flanked Tgif1 loci have been

reported previously (Shen and Walsh, 2005). To delete Tgif1 in mature osteoblasts and osteocytes,

mice expressing the Cre recombinase under the control of the 8kb fragment of the murine Dentin

matrix protein 1 (Dmp1-Cre^{Tg}) (Bivi et al., 2012) promoter were crossed with mice in which exons

2 and 3 of the Tgif1 gene were flanked by loxP sites (Tgif1$^{fl/+}$). Since no bone phenotype was

observed in *Dmp1-Cre$^+$*; *Tgif1$^{+/+}$* mice, *Dmp1-Cre$^-$; Tgif1$^{fl/fl}$* mice were used as controls (Saito et

al., 2019). In anabolic studies, parathyroid hormone (PTH 1-34; 100 µg/kg of body weight,

Biochem) was administered in 8-week-old male mice intraperitoneally 5 times a week for 3 weeks.

To induce conditions of bone healing, 10-week-old male mice of the genotypes *Tgif1$^{+/+}$* or *Tgif1$^-$*

$^{/-}$ were subject to an open fracture of the tibial midshaft, which was stabilized by an intramedullary

pin (Schindeler et al., 2008). Three weeks later, mice were sacrificed with subsequent histological

examination of the bone tissue. All animal experiments were approved by the local authorities.

Histology and histomorphometry. Mouse tibiae were collected and fixed in 3.7% PBS-buffered

formaldehyde. For histomorphometric analysis, tibiae were embedded in methylmethacrylate.

Toluidine blue staining was performed on 4 µm sagittal sections. Quantitative bone

histomorphometric measurements were performed on von Kossa and Toluidine blue stained

sections using the OsteoMeasure system (OsteoMetrics). For the quantification of quiescent

surfaces (Qs) only surfaces covered by flat cells of intense blue staining without osteoid deposition

were considered.

Isolation of primary osteoblasts and cell culture. To obtain calvarial osteoblasts, mouse

calvariae were dissected from 1–3 days old neonatal mice and digested sequentially 5 times for 25

minutes in α-MEM (Gibco) containing 0.1% collagenase (Roche) and 0.2% dispase (Roche). Cells

obtained from fractions 2–5 were combined according to the genotype and expanded in α-MEM

containing 10% FBS (Gibco), 100 U/ml penicillin, and 100 µg/ml streptomycin (P/S, Thermo Fisher Scientific). Long bone osteoblasts were isolated from femora and tibiae of 8-week-old mice. After removing muscles in sterile PBS, bone marrow was flushed, and bones were cut in small pieces. Bone pieces were digested with 0.1% collagenase for 2 hours at 37 °C and plated in α-MEM containing 10% FBS and P/S. The OCY454 cell line (RRID:CVCL_UW31) was kindly provided by Dr. Paola Divieti Pajevic (Department of Molecular and Cell Biology, Boston University, Boston, USA) and cultured in α-MEM with 10% FBS, 100 U/ml penicillin, and 100 µg/ml streptomycin at 33°C to permit proliferation and maintain osteoblast phenotype, and at 37°C for the experiments.

3D cell culture. Long bone osteoblasts were isolated as described above. Bone pieces were digested with 0.1% collagenase for 2 hours at 37 °C and plated in α-MEM containing 10% FBS and P/S. Once confluent, outgrowing osteoblasts were seeded on Cellmatrix Type 1A (Nitta Geratin) placed onto a Millicell® membrane (Sigma-Aldrich) in α-MEM with 10% FBS, 100 U/ml penicillin, and 100 µg/ml streptomycin, as reported previously (Uchihashi et al., 2013). After 24 hours, the membrane/gel/cells complexes were fixed in 1,5% formaldehyde, embedded in OCT and frozen. 10 µm sections were cut and stained with Hematoxylin/Eosin and for immunocytochemistry with phalloidin and DAPI as described below.

Cell spreading and migration assays. For adhesion and spreading assays, cells were incubated in α-MEM with 1% FBS for 4h and removed from the culture flasks by gentle trypsinization (Trypsin-EDTA 0.05%, Thermo Fisher Scientific). Cells were left to adhere for 20, 40, 60, 120 or 240 minutes on Collagen I-coated 8 well-slides ($5*10^3$ cells/well) or plates ($5*10^5$ cells/well). For qPCR, control cells were left in suspension. All spreading assays were performed in serum-free conditions at 37°C in duplicates or triplicates, which were combined prior to RNA or protein analysis. For the quantification of adhered cells, cells were fixed and stained with hematoxylin or

incubated with Calcein–AM 2 µM (Thermo Fisher Scientific) for 60 minutes. For migration

assays, cells were plated at a density of 7500 cells /cm^2 on Collagen I-coated 6-well plates in

culture medium. Cells adhered for 2-3 hours before the medium was changed to α-MEM

containing 0.1% FBS. Live cell migration assay was captured using the Improvision LiveCell

Spinning Disk microscope. For live cell imaging, 5 regions of interest/well were selected at time

point 0 and images were captured every 10 minutes for 16 hours. Analysis of migration track

length, velocity and meandering index was performed using the software Volocity 6.

Cell treatment. For the inhibition of gene transcription, cells were incubated for 15 minutes with

5 µM of Actinomycin-D (Sigma). For the inhibition of ERK1/2 signaling, cells were incubated for

20 minutes with 1 µg/ml SCH772984 (Selleckchem) (Morris et al., 2013). For the *in vitro* PTH

treatment, cells were incubated with 100 nM PTH (1-34, Biochem) in α-MEM containing 1% FBS

for 4 hours prior to the spreading assays.

RNA isolation and gene expression analysis. Total RNA was isolated using the RNeasy plus kit

(Invitrogen) according to manufacturer's instruction. cDNA was synthesized from total RNA

using the ProtoScript First Strand cDNA Synthesis Kit (NEBioLabs). Quantitative real-time PCR

was performed using SYBR Green (BioRad). The fold induction of each target gene was calculated

using the comparative CT method ($2^{-\Delta\Delta Ct}$). The C_t of each gene of interest was first normalized to

the housekeeping gene GAPDH (ΔC_t), and then to the control of the experiment (vehicle or DMSO

treatment) to calculate the $\Delta\Delta C_{t.}$ The oligonucleotide sequences used were the following:

mTgif1	**For** GCAGACACACCTGTCCACACTA
	Rev GGAATGAAATGGGCTCTCTTCT

mPAK3	**For** CTGAGCAATGGGCACGACTA
	Rev AGCCAAAGGAGGTTCCGAAG
mPAK1	**For** TTAGCCGAATCCAGCCTGTCA
	Rev TGGTGTTTCTCATCGGAGGG
mPAK2	**For** GCGGTTGTTCCGCTTCC
	Rev AATTATGAAACAGCCAGAGAGGA
mPAK4	**For** GCGCCAAGCCGATGAGTAA
	Rev CTGAGATCTCCACCCGCTTC
mCdc42ep1	**For** ACGACACGAGGTCTCCACTC
	Rev CCGGGCATCAGCTTTGATTG
mCdc42ep2	**For** CGCTCCTCAAGCTTCTCAACTC
	Rev AAGGCCAACAAAGACAGGGT
mCdc42ep4	**For** GAGGCCTGGCCCTCTGATT
	Rev GCACTGATCATCTCGGCTGT
mCdc42ep5	**For** TCCTCCTGTCCCTACAATCCC
	Rev CCGGCATCTGCACCAGGAATA
mCdc42se1	**For** GAGACTCAGAAAGGGCCCCA
	Rev AGATGTGGAATCCTGCGGCT
mCdc42se2	**For** GCTTGGTGTGTGGAGATCCT
	Rev TCAACACTGGCCCTATCGTC
mGAPDH	**For** TGCACCACCAACTGCTTAG
	Rev GGATGCAGGGATGATGTTC

Immunocytochemistry. Upon adherence of calvarial osteoblasts or OCY454 cells on Collagen I

(Roche)-coated surfaces of 8-well chambers, cells were fixed in 1.5% formaldehyde and immune-

stained with rabbit anti-Paxillin antibody (1:150, Abcam) or mouse anti-Talin antibody (1:100,

Sigma), followed by an incubation with a 488-Alexa Fluor conjugated secondary antibody (1:500,

Thermo Fisher Scientific). Hoechst 33258 (Thermo Fisher Scientific) was used at a 1:10000

dilution for nuclear staining. 633-Alexa Fluor conjugated phalloidin (Thermo Fisher Scientific)

was used at a 1:20 dilution to stain actin. Cover slips were mounted with Fluoromount G (Southern

Biotech).

Image acquisition and quantification. Images were acquired using the Zeiss Apotome or the

confocal Leica SP5 and SP8 microscopes with 20X or 63X objectives. Images for the analysis of

focal adhesions were quantified with the Image J software as described previously (Horzum et al.,

2014). Images for 3-dimensional (3D) quantification were elaborated by IMARIS software

(BITPLANE, Oxford). The Z-stacks obtained by confocal microscopy were rebuilt by IMARIS

and the parameters of interest were quantified after removal of the background noise. The value

of the sphericity for each cell was provided automatically by the software after cell selection.

Quantification of images was performed using raw data. Representative images shown have been

optimized equally and only for brightness and contrast.

Immunoblot analysis. Samples were lysed in low salt RIPA buffer pH 7.5 (50 mM Tris base, 150

mM NaCl, 0.5% Nonidet P-40, 0.25 % Sodium deoxycholate,) supplemented with complete

protease and phosphatase inhibitors (Roche). Lysates were denaturated in Laemmli Sample Buffer

(BioRad), resolved by SDS-PAGE and transferred onto nitrocellulose membranes (GE

Healthcare). Membranes were incubated with the following primary antibodies: rabbit anti-Tgif1

antibody (Abcam, 1:500), rabbit anti-PAK3 antibody (Cell Signalling, 1:300), rabbit anti-phospho

ERK1/2 antibody (Cell Signalling, 1:1000), mouse anti-ERK1/2 antibody (Cell Signalling,

1:1000), mouse anti-tubulin antibody (Sigma, 1:2000), mouse anti-actin antibody (Millipore,

1:5000), rabbit anti-integrin β1 (Cell Signaling, 1:500), rabbit-anti FAK (Cell Signaling, 1:500),

rabbit-anti p-FAK (Cell Signaling, 1:500), rabbit-anti paxillin (Cell Signaling, 1:500), rabbit-anti

p-S83 paxillin (Cell Signaling, 1:500), mouse anti-src (Cell Signaling, 1:500), rabbit anti-phospho

src (Cell Signaling, 1:500), rabbit anti-p38 (Cell Signaling, 1:500), rabbit anti-phospho p38 (Cell

Signaling, 1:500), rabbit-anti LRG5 (Cell Signaling, 1:500). Membranes were washed with TBST

and incubated either with anti-mouse or anti-rabbit IgG horseradish peroxidase-conjugated

secondary antibody (Promega, 1:10000). Chemiluminescence images were captured by the

BioRad ChemiDoc detection system.

Gene silencing. OCY454 cells (10^6 cells/reaction) were transfected with the small interfering

RNAs (siRNAs) ON-TARGET plus mouse Tgif1 SMART pool (Dharmacon) or ON-TARGET

plus non-targeting control pool (scramble) using NEON electroporation (Invitrogen). ON-

TARGET plus mouse PAK3 SMART pool siRNAs or scrambled were transfected 24 hours later

to the same cells using Lipofectamine 3000 (Life Technologies). The final siRNA concentration

was 70 nM/sample.

DNA constructs and luciferase assays. The wild-type human Tgif1 promoter luciferase reporter

constructs, and the Tgif1 promoter luciferase reporter construct in which the AP1 binding site was

mutated have been reported previously (Saito et al., 2019). OCY454 cells were transfected with

the human Tgif1 promoter luciferase reporter constructs, or the 6X-TRE-luciferase reporter (6X-

TRE-luc; Clontech) to analyze AP-1 activity (Sabatakos et al., 2008), or the (-2329/149) rat PAK3

promoter, provided by Dr. Virna D. Leaner (Institute of Infectious Disease and Molecular

Medicine, University of Cape Town, Cape Town, South Africa) (Holderness Parker et al., 2013),

along with the humanized Renilla reporter plasmid (1:10 ratio) using Lipofectamine 3000.

Luciferase assays were performed using the Dual Luciferase Reporter Gene Assay System

(Promega) according to the instructions provided by the manufacturer. Firefly luciferase activity

was normalized to Renilla luciferase activity.

Promoter analysis and chromatin immunoprecipitation. Analysis of putative Tgif binding sites

on the (-2329/149) rat PAK3 promoter was performed using the online platform ALGGEN-

PROMO. Chromatin immunoprecipitation experiments were performed using the A/G MAGNA

CHIP kit (Millipore) according to the manufacturer's instruction. Briefly, $10*10^6$ cells of the

OCY454 cell line were plated onto $15cm^2$ dishes and incubated at 37°C for 5 days. Cells were then

fixed with 1% formaldehyde for 10 min, followed by 5 min of quenching using glycin. After

chromatin isolation, DNA was sheared in fragments ranging between 100 and 500 bp in length

through 15 cycles of high frequency sonication using a BioraptorPlus. Immunoprecipitations were

performed using 7 μg of rabbit polyclonal anti-Tgif1 antibody (Santa Cruz Biotechnology) or anti-

rabbit ChIP grade IgG (Abcam). The amount of pulled-down DNA was quantified by qPCR. ChIP-

qPCR data were first normalized to the input of each precipitation ($2^{(Ct\ 100\%\ input-\ Ct\ sample)}$), and then

to the relative IgG sample. The gene RARα was used as positive control (Zhang et al., 2009). The

oligonucleotides used to amplify the Tgif1 binding site and as negative control were the following:

BS # 1	**For**	CCAGGAAAATCCTGTTATGCTTCC	**Neg**	**For**	GAAATTCTGTGTTGGCCGCA
	Rev	GGGTTAAAACAGTAGCTACATCCC		**Rev**	TCAGCACCTACAATTCTGACCA

Statistical analysis. Statistical analyses were performed using the statistical package Prism v 9.00

(GraphPad Software, San Diego, CA, USA). Statistically significant differences were determined

using the unpaired T-test for comparing two groups. ANOVA followed by Bonferroni or Tuckey

Multiple Comparison Test was used to compare more than two groups. The repeated measures

ANOVA, estimated margins means test was used to compare the differences of two genotypes

over time.

Figure legends

Figure 1. Loss of Tgif1 reduces osteoblast size *in vivo* and impairs osteoblast adhesion and migration *in vitro*. (**A**) Representative images of the femora from 12-week-old *Tgif1*$^{+/+}$ (n=4) and *Tgif1*$^{-/-}$ (n=4) mice stained with Toluidine blue. (**B**) Femora of 12-week-old *Dmp1-Cre*$^{-}$*;Tgif1*$^{fl/fl}$ and *Dmp1-Cre*$^{+}$*;Tgif1*$^{fl/fl}$ mice stained with toluidine blue. (**A, B**) Arrows indicate osteoblasts. Scale bars indicate 100 µM. (**C, D**) Osteoblasts isolated from calvariae of neonatal *Tgif1*$^{+/+}$ and *Tgif1*$^{-/-}$ mice upon adherence on Col-I coated surfaces for 1, 2 and 4 hours, fixation and staining with toluidine blue. (**C**) Representative images of Toluide blue-stained cells after 2 hours of adhesion. (**D**) Quantification of adherent *Tgif1*$^{+/+}$ and *Tgif1*$^{-/-}$ calvarial osteoblasts after 1, 2, 4 hours of adhesion on Col-I-coated surfaces. (**E**) Osteoblasts isolated from long bones of 8-week-old *Tgif1*$^{+/+}$ and *Tgif1*$^{-/-}$ mice upon adherence on Col-I coated surfaces for 1, 2 and 4 hours, fixation and staining with toluidine blue. Quantification of adherent cells at indicated time points. (**F-H**) Migration of calvarial osteoblasts obtained from neonatal *Tgif1*$^{+/+}$ and *Tgif1*$^{-/-}$ mice was analyzed using live cell imaging. Quantification of (**F**) track length, (**G**) migration velocity and (**H**) meandering index of *Tgif1*$^{+/+}$ and *Tgif1*$^{-/-}$ calvarial osteoblasts. (**I**) Long bones were harvested from *Tgif1*$^{+/+}$ and *Tgif1*$^{-/-}$ mice. Osteoblasts were isolated from bone chips and spread on Col-I matrices placed on porous membranes for 48h. Membranes were frozen, cut and stained with (**J**) H&E or (**K**) phalloidin (yellow) and DAPI (blue). Scale bars indicate 50 µm. n = minimum of 3 independent experiments with technical duplicates. Unpaired t-test, *p<0.05, **p<0.01, ***p<0.001 vs. *Tgif1*$^{+/+}$.

Figure 2. Tgif1-deficient osteoblasts are impaired to spread and form focal adhesions
(**A, B**) Calvarial osteoblast obtained from *Tgif1*$^{+/+}$ and *Tgif1*$^{-/-}$ mice adhered on Col-I coated slides for 20, 40 and 60 minutes. (**A**) Cell spreading was visualized by Calcein-AM. (**B**) The number of round (non-spread) cells was counted. Scale bar indicates 100 µm. n=6 independent experiments; Repeated Measures ANOVA, Estimated margins means test, *p<0.05 between genotypes. (**C**)

Tgif1$^{+/+}$ and *Tgif1*$^{-/-}$ calvarial osteoblasts were allowed to adhere on Col-I coated slides for 60

minutes. Focal adhesion formation was visualized by paxillin staining (green) and actin

cytoskeleton by phalloidin (magneta). Scale bars indicate 100 μm (lower magnification) or 10 μm

(single cells). **(D)** Analysis of single cell sphericity after 3D reconstruction using IMARIS and **(E)**

quantification of focal adhesions using Image J. **(F-H)** Tgif1 was silenced in OCY454 cells using

siRNA and cells were allowed to adhere on Col-I coated slides for 60 minutes. **(F)** Focal adhesion

formation was visualized by Paxillin staining (green) and cell protrusions by phalloidin (magenta).

(G) Analysis of cell sphericity after 3D reconstruction and **(H)** quantification of focal adhesions.

Unpaired t-test, ***p<0.001, **p<0.01 vs. *Tgif1*$^{+/+}$ (D, E) and scr (G, H).

Figure 3. Silencing of PAK3 restores cell spreading and focal adhesion formation in Tgif1-

deficient cells. (A) *PAK3* mRNA expression in OCY454 cells transfected with siRNA targeting

Tgif1 (siTgif1) or scrambled control (scr) siRNA before (0) and after 20 and 60 minutes of

spreading on Col-I coated slides. n=9, 1-way ANOVA with Tukey's multiple comparison test,

*p<0.05, ***p<0.001. **(B)** Representative images of immunoblots demonstrating the abundance

of PAK3 expression upon silencing of Tgif1. Actin was used as a control. **(C)** *PAK3* mRNA

expression in tibiae of 12-week-old *Tgif1*$^{+/+}$ (n=6) and *Tgif1*$^{-/-}$ (n=6) mice. **(D)** Tgif1 binding to

the predicted site of the PAK3 promoter in OCY454 cells analyzed by ChIP and quantified as fold

enrichment to the relative IgG control. Negative and positive (RAR) controls were used as

indicated. n=8, unpaired t-test with Welch's correction, *p<0.05, **p<0.01 vs. respective IgG

control. **(E)** Tgif1-deficient (siTgif1) or control (scr) OCY454 cells were transfected with renilla

plasmid and a pGL3 plasmid (EV) or a pGL3 plasmid containing a 2.3 kb fragment of the rat

PAK3 promoter upstream of the luciferase gene. The promoter activity was quantified using

luciferase assays and presented as relative luciferase activity (luciferase/renilla). 1-way ANOVA,

Tukey's multiple comparison test, ***p<0.001. **(F-H)** OCY454 cells were transfected alone or in

combinations with siTgif1 for 48h, siPAK3 for 24 h and scrambled (scr) control. **(F)** Cells were

allowed to adhere on Col-I coated slides for 60 minutes. Formation of focal adhesions was

visualized by paxillin staining (green) and actin cytoskeleton by phalloidin staining (magenta).

Scale bars indicate 100 µm (two upper rows) or 10 µm (two lower rows). **(G)** Quantification of

cell sphericity using IMARIS. **(H)** Quantification of the number of mature focal adhesions per cell

using the Image J software. n=4 independent experiments in which individual cells were analyzed.

1-way ANOVA, Tukey's multiple comparison test, **p<0.01 ***p<0.001.

Figure 4. Tgif1 expression is increased during cell spreading via ERK and AP1 signaling

pathways. (A) Quantification of *Tgif1* mRNA expression and **(B)** representative images of

immunoblots of Tgif1 expression in calvarial osteoblasts before (0 min) and after 20 and 60

minutes of spreading. Actin was used as control. **(C)** *Tgif1* mRNA expression and **(D)**

representative images of immunoblots of Tgif1 expression in OCY454 cells before (0 min) and

after 20 and 60 minutes spreading on Col-I coated slides. Actin was used as control. **(E)** OCY454

cells were treated with DMSO as control or with 5 µM Actinomycin D for 15 minutes prior to

adherence on Col-I coated slides for 20 and 60 minutes. *Tgif1* mRNA was quantified before (0

min) and during spreading. **(F)** Representative immunoblot images demonstrating the efficiency

of ERK1/2 inhibitor SCH772984 (1µg/ml) to prevent ERK1/2 phosphorylation (n=4). **(G)**

Quantification of *Tgif1* mRNA expression during spreading after 15 minutes pre-treatment with

DMSO (control) or with the ERK inhibitor SCH772984. **(H)** Quantification of the Tgif1 promoter

activity during cell spreading. OCY454 cells were transfected with a luciferase reporter plasmid

encoding a 2.2 kb fragment of the Tgif1 promoter or progressive truncations thereof (1.7, 1.2 and

0.5 kb) along with a plasmid encoding renilla firefly as control. Upon spreading for 60 minutes,

promoter activity was quantified using a dual luciferase reporter gene assay and presented as

normalized luciferase activity (luciferase/renilla). **(I)** Quantification of the AP1 transcriptional

activity during cell spreading. OCY454 cells were transfected with a 6X-TRE-luciferase reporter

to determine AP-1 activity along with a plasmid encoding renilla firefly as control. Upon spreading

for 60 minutes, promoter activity was quantified using a dual luciferase reporter gene assay and

presented as normalized luciferase activity (luciferase/renilla). **(J)** Quantification of the activity of

the wild-type (wt) Tgif1 2.2 kb promoter and of the same fragment bearing a mutant AP1 binding

site (AP1 mut) during cell spreading. OCY454 cells were transfected either with a luciferase

reporter plasmid encoding a wild-type (wt) 2.2 kb fragment of the Tgif1 promoter or with the same

promoter in which the AP1 binding site has been mutated (mut) along with a plasmid encoding

renilla firefly as control. Upon spreading for 60 minutes, the promoter activity was quantified

using a dual luciferase reporter gene assay and presented as normalized luciferase activity

(luciferase/renilla). 1-way ANOVA, Tukey's multiple comparisons test, $*p<0.05$, $**p<0.01$,

$***p<0.001$ vs. respective control.

Figure 5: Tgif1 deficiency causes a reduced number and activity of osteoblasts during bone

repair. **(A)** Representative images of Toluidine blue-stained sections of bones from male *Tgif1*$^{+/+}$

(n=9) and *Tgif1*$^{-/-}$ mice (n=5) that received an open fracture of the midshaft tibia at 10 weeks of

age followed by bone healing for 3 weeks upon intramedullary stabilization. Arrows indicate

osteoblasts. Scale bars indicate 100 µm (left panels) and 50 µm (right panels). **(B - E)** Histological

sections were used to quantify the histomorphometric parameters **(B)** bone volume per callus

volume (BV/Cal.V), **(C)** osteoid surface per bone surface (OS/BS), **(D)** number of osteoblasts per

bone surface (N.Ob/BS) and **(E)** osteoblast surface per bone surface (Ob.S/BS). Unpaired t-test,

$*p<0.05$, $**p<0.01$ vs. *Tgif1*$^{+/+}$.

Figure 6. PTH is impaired in activating quiescent bone surfaces in *Tgif1*-deficient mice. (A)

Toluidine blue staining of tibiae from 12-week-old *Tgif1*$^{+/+}$ and *Tgif1*$^{-/-}$ mice treated with vehicle

(veh) or PTH. Quiescent surfaces are indicated by red arrows. Representative images are shown.

Scale bar indicates 100 µm. **(B)** Quantification of the percentage of quiescent surfaces per bone

surface (BS). *Tgif1*$^{+/+}$, veh n=10; *Tgif1*$^{+/+}$, PTH n=10, *Tgif1*$^{-/-}$, veh n=8; *Tgif1*$^{-/-}$, PTH n=10. (**C**)

Toluidine blue staining of tibiae from 12-week-old *Dmp1-Cre$^-$;Tgif1$^{fl/fl}$* and *Dmp1-Cre$^+$;Tgif1$^{fl/fl}$*

mice treated with vehicle (veh) or PTH. Quiescent surfaces are indicated by red arrows.

Representative images are shown. Scale bar indicates 100 µm. **(D)** Quantification of the percentage

of quiescent surfaces per bone surface (BS). *Dmp1-Cre$^-$;Tgif1$^{fl/fl}$* , veh n=8; *Dmp1-Cre$^-$;Tgif1$^{fl/fl}$*,

PTH n=8, *Dmp1-Cre$^+$;Tgif1$^{fl/fl}$*, veh n=8; *Dmp1-Cre$^+$;Tgif1$^{fl/fl}$* , PTH n=8. 1-way ANOVA with

Tukey's multiple comparison test, *$p < 0.05$, **$p < 0.01$.

Figure 7. PTH promotes cell spreading via Tgif1-PAK3 signaling. (A) OCY454 cells were

transfected with siRNA against Tgif1 (siTgif1) or scramble control siRNA (scr) and treated with

vehicle (veh) or PTH for 4h. Cells were allowed to adhere on Col-I coated slides for 60 minutes.

The cytoskeleton was visualized by phalloidin staining (magenta). Scale bars indicate 100 µm

(upper panel) or 10 µm (lower panel). **(B)** Cell sphericity was quantified using IMARIS software.

1-way ANOVA with Tukey's multiple comparisons test was applied, **$p < 0.01$, ***$p < 0.001$. **(C)**

PAK3 mRNA expression after 4 hours of PTH treatment. Unpaired t-test, **$p < 0.01$ vs. veh. **(D)**

Representative image of immunoblots demonstrating PAK3 and Tgif1 protein abundance upon

PTH treatment in cells transfected with siTgif1 or scr. Actin was used as a loading control. **(E)**

OCY454 cells were transfected with scr or Tgif1 siRNA for 48h and PAK3 or scr siRNA for 24 h

and treated with vehicle (veh) or PTH for 4h. Cells adhered on Col-I coated slides for 60 minutes.

The cytoskeleton was visualized by phalloidin staining (magenta). Scale bar indicates 50 µm. **(F)**

Cell sphericity was quantified using IMARIS software. n=4, 1-way ANOVA, Tukey's multiple

comparison test, *$p < 0.05$, ***$p < 0.001$.

Supplementary Figures

Fig. S1: (A) Migration of calvarial osteoblasts obtained from neonatal *Tgif1*$^{+/+}$ (n=20) and *Tgif1*$^{-}$

$^{/-}$ (n=20) mice was analyzed using live cell imaging. The direction and distance of individual cell

migration is visualized.

(B) Representative image of an immunoblot demonstrating the expression of Tgif1 in OCY454

cells transfected with scrambled control siRNA (Scr) or siRNA against Tgif1 (siTgif1). Actin was

used as loading control.

Fig. S2: Tgif1-deficiency does not alter the abundance or activation of the FA components

and Tgif1 does not co-localize with FA complexes. (A) Osteoblasts adhered on Col-I coated

slides for 0, 20 or 60 minutes. Representative images of immunoblots demonstrating the

abundance of p-FAK, FAK, p-p38, p38, p-Paxillin, Paxillin, p-src, src, Integrin β1, LRG5 and

Talin. Actin was used as loading control. n=4 independent experiments. **(B)** Osteoblasts over-

expressing eGFP-Tgif1 or eGFP adhered on Col-I coated plates for 60 minutes. Cells were stained

for actin (magenta), nuclei (blue), Paxillin (upper panels) or Talin (lower panels, both yellow).

Fig. S3: Lack of Tgif1 does not alter the expression of CEPs, PAK1, PAK2 or PAK4. OCY454

cells were transfected with a siRNA targeting Tgif1 (siTgif1) or scramble control (scr) prior to

adherence on Col-I coated slides for 20 & 60 minutes. Gene expression was quantified by qPCR

for **(A)** Cdc42es2, **(B)** Cdc42ep1, **(C)** Cdc42ep2, **(D)** Cdc42ep4, **(E)** PAK1, **(F)** PAK2 and **(G)**

PAK4. 1-way ANOVA, Tukey's Multiple Comparison Test.

Fig. S4: (A) Schematic of the conserved predicted Tgif binding site on the rat and mouse PAK3

promoters. **(B)** OCY454 cells were transfected with siRNA targeting Tgif1 (siTgif1) for 48h and

with siRNA targeting PAK3 (siRNA) for 24h alone and in combination alongside with the

respective scrambled controls (scr) resulting in four groups: scr (scr; scr), siPAK3 (scr; siPAK3),

siTgif1 (scr; siTgif1) and siPAK3+siTgif1 (siPAK3; siTgif1). Representative images of

immunoblots demonstrate the abundance of PAK3 and Tgif1. Actin was used as loading control.

n=3 independent experiments.

Acknowledgements

We are grateful to A. Gasser und S. Schroeder as well as A.V. Failla and B. Zobiak of the UKE

Microscopy Imaging Facility (Umif) for conceptual and technical support; G. Arndt and P.

Missberger for mouse husbandry. We thank C. Walsh for providing Tgif1-deficient mice, S.J.

Brandt for providing the Tgif1 reporter constructs, P. Divieti Pajevic for kindly contributing the

Ocy454 cell line and V.D. Leaner for providing the rat PAK3 promoter. S. Bolamperti was

supported by a postdoctoral fellowship from the European Calcified Tissue Society

(ECTSABBF2015_0019). H. Saito was supported by a postdoctoral fellowship from the Japanese

Society for the Promotion of Science. E. Hesse received funding from the Deutsche

Forschungsgemeinschaft (HE 5208/2-1, HE 5208/2-3 and HE 5208/3-1), the AO-Foundation (S-

13-73H) and the European Commission (PCIG10-GA-2011-303722). H. Taipaleenmäki received

funding from the Deutsche Forschungsgemeinschaft (TA 1154/1-1).

References

Baron, R., and E. Hesse. 2012. Update on bone anabolics in osteoporosis treatment: rationale, current status, and perspectives. *J Clin Endocrinol Metab*. 97:311–25. doi:10.1210/jc.2011-2332.

Bivi, N., K.W. Condon, M.R. Allen, N. Farlow, G. Passeri, L.R. Brun, Y. Rhee, T. Bellido, and L.I. Plotkin. 2012. Cell autonomous requirement of connexin 43 for osteocyte survival: consequences for endocortical resorption and periosteal bone formation. *J Bone Miner Res*. 27:374–389. doi:10.1002/JBMR.548.

Black, D.M., and C.J. Rosen. 2016. Clinical Practice. Postmenopausal Osteoporosis. *N Engl J Med*. 374:254–62. doi:10.1056/NEJMCP1513724.

Casati, L., F. Celotti, P. Negri-Cesi, M.C. Sacchi, P. Castanoy, and A. Colciago. 2015. Platelet derived growth factor (PDGF) contained in Platelet Rich Plasma (PRP) stimulates migration of osteoblasts by reorganizing actin cytoskeleton. *https://doi.org/10.4161/19336918.2014.972785*. 8:595–602. doi:10.4161/19336918.2014.972785.

Cronin, N.M., and K.A. Demali. 2021. Dynamics of the Actin Cytoskeleton at Adhesion Complexes. *Biology 2022, Vol. 11, Page 52*. 11:52. doi:10.3390/BIOLOGY11010052.

Cruz-Matías, I., D. Ayala, D. Hiller, S. Gutsch, M. Zacharias, S. Estradé, and F. Peiró. 2019. Sphericity and roundness computation for particles using the extreme vertices model. *J Comput Sci*. 30:28–40. doi:10.1016/J.JOCS.2018.11.005.

Dirckx, N., M. Van Hul, and C. Maes. 2013a. Osteoblast recruitment to sites of bone formation in skeletal development, homeostasis, and regeneration. *Birth Defects Res C Embryo Today*. 99:170–91. doi:10.1002/bdrc.21047.

Dirckx, N., M. Van Hul, and C. Maes. 2013b. Osteoblast recruitment to sites of bone formation in skeletal development, homeostasis, and regeneration. *Birth Defects Res C Embryo Today*. 99:170–191. doi:10.1002/BDRC.21047.

Dobnig, H., and R.T. Turner. 1995. Evidence that intermittent treatment with parathyroid hormone increases bone formation in adult rats by activation of bone lining cells. *Endocrinology*. 136:3632–3638. doi:10.1210/ENDO.136.8.7628403.

Duchamp De Lageneste, O., A. Julien, R. Abou-Khalil, G. Frangi, C. Carvalho, N. Cagnard, C. Cordier, S.J. Conway, and C. Colnot. 2018. Periosteum contains skeletal stem cells with high bone regenerative potential controlled by Periostin. *Nat Commun*. 9. doi:10.1038/S41467-018-03124-Z.

Einhorn, T.A., and L.C. Gerstenfeld. 2015. Fracture healing: mechanisms and interventions. *Nat Rev Rheumatol*. 11:45–54. doi:10.1038/NRRHEUM.2014.164.

Eleniste, P.P., S. Huang, K. Wayakanon, H.W. Largura, and A. Bruzzaniti. 2014. Osteoblast differentiation and migration are regulated by dynamin GTPase activity. *Int J Biochem Cell Biol*. 46:9–18. doi:10.1016/J.BIOCEL.2013.10.008.

Emerson, L.J., M.R. Holt, M.A. Wheeler, M. Wehnert, M. Parsons, and J.A. Ellis. 2009. Defects in cell spreading and ERK1/2 activation in fibroblasts with lamin A/C mutations. *Biochim Biophys Acta*. 1792:810–821. doi:10.1016/J.BBADIS.2009.05.007.

Etienne-Manneville, S. 2004. Actin and Microtubules in Cell Motility: Which One is in Control? *Traffic*. 5:470–477. doi:10.1111/J.1600-0854.2004.00196.X.

Fincham, V.J., M. James, M.C. Frame, and S.J. Winder. 2000. Active ERK/MAP kinase is targeted to newly forming cell–matrix adhesions by integrin engagement and v-Src. *EMBO J*. 19:2911–2923. doi:10.1093/EMBOJ/19.12.2911.

Gronthos, S., P.J. Simmons, S.E. Graves, and P. G. Robey. 2001. Integrin-mediated interactions between human bone marrow stromal precursor cells and the extracellular matrix. *Bone*. 28:174–181. doi:10.1016/S8756-3282(00)00424-5.

Haider, M.-T., H. Saito, E. Hesse, and H. Taipaleenmäki. 2018. Breast Cancer Bone Metastases are Attenuated in a Tgif1-deficient Bone Microenvironment. *J Bone Miner Res 32*.

Hirsch, D.S., D.M. Pirone, and P.D. Burbelo. 2001. A new family of Cdc42 effector proteins, CEPs, function in fibroblast and epithelial cell shape changes. *J Biol Chem*. 276:875–883. doi:10.1074/JBC.M007039200.

Holderness Parker, N., H. Donninger, M.J. Birrer, and V.D. Leaner. 2013. p21-Activated Kinase 3 (PAK3) Is an AP-1 Regulated Gene Contributing to Actin Organisation and Migration of Transformed Fibroblasts. *PLoS One*. doi:10.1371/journal.pone.0066892.

Hood, J.D., and D.A. Cheresh. 2002. Role of integrins in cell invasion and migration. *Nature Reviews Cancer 2002 2:2*. 2:91–100. doi:10.1038/nrc727.

Horzum, U., B. Ozdil, and D. Pesen-Okvur. 2014. Step-by-step quantitative analysis of focal adhesions. *MethodsX*. doi:10.1016/j.mex.2014.06.004.

Jafari, A., A. Isa, L. Chen, N. Ditzel, W. Zaher, L. Harkness, H.E. Johnsen, B.M. Abdallah, C. Clausen, and M. Kassem. 2019. TAFA2 Induces Skeletal (Stromal) Stem Cell Migration Through Activation of Rac1-p38 Signaling. *Stem Cells*. 37:407–416. doi:10.1002/STEM.2955.

Jilka, R.L., C.A. O'Brien, A.A. Ali, P.K. Roberson, R.S. Weinstein, and S.C. Manolagas. 2009. Intermittent PTH stimulates periosteal bone formation by actions on post-mitotic preosteoblasts. *Bone*. 44:275–286. doi:10.1016/J.BONE.2008.10.037.

Jones, S.J., and A. Boyde. 1977. The migration of osteoblasts. *Cell Tissue Res*. 184:179–193. doi:10.1007/BF00223067/METRICS.

Kaiser, E., M. Sato, J.E. Onyia, and S. Chandrasekhar. 2001. Parathyroid hormone (1–34) regulates integrin expression in vivo in rat osteoblasts. *J Cell Biochem*. 83:617–630. doi:10.1002/JCB.1256.

Kechagia, J.Z., J. Ivaska, and P. Roca-Cusachs. 2019. Integrins as biomechanical sensors of the microenvironment. *Nature Reviews Molecular Cell Biology 2019 20:8*. 20:457–473. doi:10.1038/s41580-019-0134-2.

Kenkre, J.S., and J.H.D. Bassett. 2018. The bone remodelling cycle. *Ann Clin Biochem*. 55:308–327. doi:10.1177/0004563218759371.

Kim, S.W., P.D. Pajevic, M. Selig, K.J. Barry, J.Y. Yang, C.S. Shin, W.Y. Baek, J.E. Kim, and H.M. Kronenberg. 2012. Intermittent parathyroid hormone administration converts quiescent lining cells to active osteoblasts. *J Bone Miner Res*. 27:2075–2084. doi:10.1002/JBMR.1665.

Kousteni, S., and J.P. Bilezikian. 2008. The cell biology of parathyroid hormone in osteoblasts. *Current Osteoporosis Reports 2008 6:2*. 6:72–76. doi:10.1007/S11914-008-0013-9.

Kreis, P., and J.V. Barnier. 2009. PAK signalling in neuronal physiology. *Cell Signal*. 21:384–393. doi:10.1016/J.CELLSIG.2008.11.001.

Löfman, O., I. Hallberg, K. Berglund, O. Wahlström, L. Kartous, A.M. Rosenqvist, L. Larsson, and G. Toss. 2009. Women with low-energy fracture should be investigated for osteoporosis. *New Pub: Medical Journals Sweden*. 78:813–821. doi:10.1080/17453670710014608.

Lomri, A., and P.J. Marie. 1990. Changes in cytoskeletal proteins in response to parathyroid hormone and 1,25-dihydroxyvitamin D in human osteoblastic cells. *Bone Miner*. 10:1–12. doi:10.1016/0169-6009(90)90045-H.

Lv, Z., A. Muheremu, X. Bai, X. Zou, T. Lin, and B. Chen. 2020. PTH(1-34) activates the migration and adhesion of BMSCs through the rictor/mTORC2 pathway. *Int J Mol Med*. 46:2089–2101. doi:10.3892/ijmm.2020.4754.

Maes, C., T. Kobayashi, M.K. Selig, S. Torrekens, S.I. Roth, S. Mackem, G. Carmeliet, and H.M. Kronenberg. 2010. Osteoblast precursors, but not mature osteoblasts, move into developing and fractured bones along with invading blood vessels. *Dev Cell*. 19:329–344. doi:10.1016/J.DEVCEL.2010.07.010.

Malluche, H.H., D.S. Porter, H. Mawad, M.C. Monier-Faugere, and D. Pienkowski. 2013. Low-energy fractures without low T-scores characteristic of osteoporosis: a possible bone matrix disorder. *J Bone Joint Surg Am*. 95. doi:10.2106/JBJS.L.01281.

Mizoguchi, T., and N. Ono. 2021. The diverse origin of bone-forming osteoblasts. *J Bone Miner Res*. 36:1432–1447. doi:10.1002/JBMR.4410.

Morris, E.J., S. Jha, C.R. Restaino, P. Dayananth, H. Zhu, A. Cooper, D. Carr, Y. Deng, W. Jin, S. Black, B. Long, J. Liu, E. DiNunzio, W. Windsor, R. Zhang, S. Zhao, M.H. Angagaw, E.M. Pinheiro, J. Desai, L. Xiao, G. Shipps, A. Hruza, J. Wang, J. Kelly, S. Paliwal, X. Gao, B.S. Babu, L. Zhu, P. Daublain, L. Zhang, B.A. Lutterbach, M.R. Pelletier, U. Philippar, P. Siliphaivanh, D. Witter, P. Kirschmeier, W. Robert Bishop, D. Hicklin, D. Gary Gillil, L. Jayaraman, L. Zawel, S. Fawell, and A.A. Samatar. 2013. Discovery of a novel ERK inhibitor with activity in models of acquired resistance to BRAF and MEK inhibitors. *Cancer Discov*. doi:10.1158/2159-8290.CD-13-0070.

Neer, R.M., C.D. Arnaud, J.R. Zanchetta, R. Prince, G.A. Gaich, J.-Y. Reginster, A.B. Hodsman, E.F. Eriksen, S. Ish-Shalom, H.K. Genant, O. Wang, D. Mellström, E.S. Oefjord, E. Marcinowska-Suchowierska, J. Salmi, H. Mulder, J. Halse, A.Z. Sawicki, and B.H. Mitlak. 2001. Effect of Parathyroid Hormone (1-34) on Fractures and Bone Mineral Density in Postmenopausal Women with Osteoporosis. *https://doi.org/10.1056/NEJM200105103441904*. 344:1434–1441. doi:10.1056/NEJM200105103441904.

Nishida, S., A. Yamaguchi, T. Tanizawa, N. Endo, T. Mashiba, Y. Uchiyama, T. Suda, S. Yoshiki, and H.E. Takahashi. 1994. Increased bone formation by intermittent parathyroid hormone administration is due to the stimulation of proliferation and differentiation of osteoprogenitor cells in bone marrow. *Bone*. 15:717–723. doi:10.1016/8756-3282(94)90322-0.

Ogita, M., M. Therese Rached, E. Dworakowski, J.P. Bilezikian, and S. Kousteni. 2008. Differentiation and Proliferation of Periosteal Osteoblast Progenitors Are Differentially Regulated by Estrogens and Intermittent Parathyroid Hormone Administration. doi:10.1210/en.2008-0369.

Pettway, G.J., J.A. Meganck, A.J. Koh, E.T. Keller, S.A. Goldstein, and L.K. McCauley. 2008. Parathyroid hormone mediates bone growth through the regulation of osteoblast proliferation and differentiation. *Bone*. 42:806–818. doi:10.1016/J.BONE.2007.11.017.

Ponzetti, M., and N. Rucci. 2021. Osteoblast Differentiation and Signaling: Established Concepts and Emerging Topics. *International Journal of Molecular Sciences 2021, Vol. 22, Page 6651*. 22:6651. doi:10.3390/IJMS22136651.

Sabatakos, G., G.C. Rowe, M. Kveiborg, M. Wu, L. Neff, R. Chiusaroli, W.M. Philbrick, and R. Baron. 2008. Doubly truncated FosB isoform (Δ2ΔFosB) induces osteosclerosis in transgenic mice and modulates expression and phosphorylation of Smads in osteoblasts independent of intrinsic AP-1 activity. *Journal of Bone and Mineral Research*. doi:10.1359/jbmr.080110.

Saito, H., A. Gasser, S. Bolamperti, M. Maeda, L. Matthies, K. Jähn, C.L. Long, H. Schlüter, M. Kwiatkowski, V. Saini, P.D. Pajevic, T. Bellido, A.J. van Wijnen, K.S. Mohammad, T.A. Guise, H. Taipaleenmäki, and E. Hesse. 2019a. TG-interacting factor 1 (Tgif1)-deficiency attenuates bone remodeling and blunts the anabolic response to parathyroid hormone. *Nat Commun*. 10:1354. doi:10.1038/s41467-019-08778-x.

Schaks, M., G. Giannone, and K. Rottner. 2019. Actin dynamics in cell migration. *Essays Biochem*. 63:483–495. doi:10.1042/EBC20190015.

Schindeler, A., A. Morse, L. Harry, C. Godfrey, K. Mikulec, M. McDonald, J.A. Gasser, and D.G. Little. 2008. Models of tibial fracture healing in normal and Nf1-deficient mice. *J Orthop Res*. 26:1053–60. doi:10.1002/jor.20628.

SenGupta, S., C.A. Parent, and J.E. Bear. 2021. The principles of directed cell migration. *Nature Reviews Molecular Cell Biology 2021 22:8*. 22:529–547. doi:10.1038/s41580-021-00366-6.

Shen, J., and C.A. Walsh. 2005. Targeted disruption of Tgif, the mouse ortholog of a human holoprosencephaly gene, does not result in holoprosencephaly in mice. *Mol Cell Biol*. 25:3639–3647. doi:10.1128/MCB.25.9.3639-3647.2005.

Silva, B.C., A.G. Costa, N.E. Cusano, S. Kousteni, and J.P. Bilezikian. 2011. Catabolic and anabolic actions of parathyroid hormone on the skeleton. *J Endocrinol Invest*. 34:801–810. doi:10.3275/7925.

Spatz, J.M., M.N. Wein, J.H. Gooi, Y. Qu, J.L. Garr, S. Liu, K.J. Barry, Y. Uda, F. Lai, C. Dedic, M. Balcells-Camps, H.M. Kronenberg, P. Babij, and P.D. Pajevic. 2015. The Wnt Inhibitor Sclerostin Is Up-regulated by Mechanical Unloading in Osteocytes in Vitro. *J Biol Chem*. 290:16744–16758. doi:10.1074/JBC.M114.628313.

Taipaleenmäki, H., H. Saito, S. Schröder, M. Maeda, R. Mettler, M. Ring, E. Rollmann, A. Gasser, C. Haasper, T. Gehrke, A. Weiss, S.K. Grimm, and E. Hesse. 2022. Antagonizing microRNA-19a/b augments PTH anabolic action and restores bone mass in osteoporosis in mice. *EMBO Mol Med*. 14. doi:10.15252/EMMM.202013617.

Tam, C.S., J.N.M. Heersche, T.M. Murray, and J.A. Parsons. 1982. Parathyroid Hormone Stimulates the Bone Apposition Rate Independently of Its Resorptive Action: Differential Effects of Intermittent and Continuous Administration. *Endocrinology*. 110:506–512. doi:10.1210/ENDO-110-2-506.

Tang, D.D., and B.D. Gerlach. 2017. The roles and regulation of the actin cytoskeleton, intermediate filaments and microtubules in smooth muscle cell migration. *Respiratory Research 2017 18:1*. 18:1–12. doi:10.1186/S12931-017-0544-7.

Thiel, A., M.K. Reumann, A. Boskey, J. Wischmann, R. von Eisenhart-Rothe, and P. Mayer-Kuckuk. 2018. Osteoblast migration in vertebrate bone. *Biological Reviews*. 93:350–363. doi:10.1111/BRV.12345.

Tsuji, N., T. Sakamoto, K. Hoshi, and A. Hikita. 2022. Spatiotemporal Analysis of Osteoblast Morphology and Wnt Signal-Induced Osteoblast Reactivation during Bone Modeling in Vitro. *JBMR Plus*. 6. doi:10.1002/JBM4.10689.

Uchihashi, K., S. Aoki, A. Matsunobu, and S. Toda. 2013. Osteoblast migration into type I collagen gel and differentiation to osteocyte-like cells within a self-produced mineralized matrix: A novel system for analyzing differentiation from osteoblast to osteocyte. *Bone*. doi:10.1016/j.bone.2012.09.001.

Wotton, D., R.S. Lo, L.A.C. Swaby, and J. Massagué. 1999. Multiple modes of repression by the Smad transcriptional corepressor TGIF. *J Biol Chem*. 274:37105–37110. doi:10.1074/JBC.274.52.37105.

Zhang, H., N. Li, Y. Tang, W. Wu, Q. Zhang, and Z. Yu. 2009a. Negative functional interaction of retinoic acid and TGF-beta signaling mediated by TG-interacting factor during chondrogenesis. *Cell Physiol Biochem*. 23:157–164. doi:10.1159/000204104.

Zhang, H., N. Li, Y. Tang, W. Wu, Q. Zhang, and Z. Yu. 2009b. Negative functional interaction of retinoic acid and TGF-β signaling mediated by TG-interacting factor during chondrogenesis. *Cellular Physiology and Biochemistry*. doi:10.1159/000204104.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

Tgif1+/+

Tgif1-/-

B

BV/Cal.V

Tgif1+/+ *Tgif1-/-*

*

C

OS/BS (%)

Tgif1+/+ *Tgif1-/-*

**

D

N.Ob/BS (/mm)

Tgif1+/+ *Tgif1-/-*

**

E

Ob.S/BS (%)

Tgif1+/+ *Tgif1-/-*

**

Figure 6

Figure 7