Whole Genome Sequencing and Assembly of the House Sparrow, *Passer domesticus*

Vikas Kumar*[1], Gopesh Sharma*[2,7], Sankalp Sharma*[2,3], Samvrutha Prasad[4]*, Toral Vaishnoi[5], Dalia Vishnudasan[4], Gopinathan Maheswaran [1], Kaomud Tyagi [1], Inderjeet Tyagi [1], Shailesh Desai[7], Kavi Kishor PB[6], Gyaneshwer Chaubey[7] and Prashanth Suravajhala [2,4]*#

1. Zoological Survey of India, Kolkata, 700053
2. Bioclues.org, India
3. NIMS University, Jaipur,
4. Amrita School of Biotechnology, Amrita Vishwa Vidyapeetham, Clappana PO 690525, Kerala, India
5. Unipath speciality Labs, Ahmedabad, Gujarat
6. Department of Genetics, Osmania University, Hyderabad 50007, India
7. Cytogenetics Lab, Department of Zoology, Banaras Hindu University, Varanasi, UP, India
8. Department of Biosciences, Manipal University Jaipur, Jaipur

*Equal contributing authors
Correspondence: prash@bioclues.org and pbkavi@yahoo.com

Abstract

The common house sparrow, *Passer domesticus* is a small bird belonging to the family Passeridae. Here, we provide high-quality whole genome sequence data along with assembly for the house sparrow. The final genome assembly was made using Shovill/SPAdes/MASURCA/BUSCO which consisted of contigs spanning 268193 bases coalescing 922 MB. We employed rigorous statistical thresholds to check the coverage as the Passer genome showed considerable similarity to *Gallus gallus* (chicken) and *Taeniopygia guttata* (Zebra finch) genomes providing a functional annotation. This new annotated genome assembly will be a valuable resource as a reference for comparative and population genomic analyses of passerine, avian, and vertebrate evolution.

Keywords: Passerines, house sparrow, next generation sequencing, evolution, adaptation

Significance

Avian evolution has been of great interest in the context of extinction. Annotating the genomes such as passerines would be of significant interest as we could understand the behavior/foraging traits and further explore their evolutionary landscape. In this work, we provide a full genome sequence of Indian house sparrow, *viz. Passer domesticus* which will serve as a useful resource in understanding the adaptability, evolution, geography, allee effects and circadian rhythms.

Introduction

Over the last 12 years, a number of bird reference genomes have been studied, providing phylogenetic relationships among the avians [1-9]. The bird genome project [B10K] has been of great significance in providing major scientific breakthroughs in phylogenetics [10,11]. With more than 1200 species, comprising 13% of all known world's avian, India is blessed considering avian diversity. Regrettably, India is ranked 3rd in having rare and threatened avian species in the world [12]. House Sparrow was introduced to India via Europe from North Africa and Eurasia by the ancient Romans. Sparrows are found in a variety of habitats, including grasslands, forests, deserts, and agricultural areas, as well as in urban areas, such as parks and gardens. They are omnivorous, typically feeding on insects, spiders, worms, seeds, fruits, and grains, and are mainly seed eaters but feed their young ones with insects and other

invertebrates; during the breeding periods, they prefer areas with a high abundance of invertebrates[13]. Over the years, there has been a tremendous decline in their population worldwide, and therefore understanding causal mechanisms through which urbanization affects their population remains limited because of numerous perplexing factors, what is believed to be due to rapid urbanization along with deforestation, lack of cavity nesting, along with absence of hedges in modern landscaping. Many hypotheses have been proposed explaining the house sparrow population decline, for example increased predation by domestic cats or sparrow hawks (*Accipiter nisus)*, cleaner streets providing reduced foraging opportunities, competition for food from other urban species, loss of nesting sites, particularly under the eaves and in the roofs of houses, pollution/air quality, both in terms of immediate toxicity and indirect toxicity through the food supply, increased use of pesticides in parks and gardens, disease transmission[14] and finally, the Allee effect [48]. *Passer domesticus,* between 1970-1990 had an extremely large breeding population with over 63 million pairs, and has been declining over the years in the European Union, including Turkey and Netherlands. Analyzing house sparrow habitat characteristics on a fine scale in almost 200 census sites with census data from 2003-2017, was one of the most complex field studies in Urban Paris, with a dramatic decline of ~89% of the species over the study period observed [15,16]. In India, a sharp decline in the house sparrow population was observed across Mumbai, Bengaluru, Hyderabad, and other major cities with certain places in India having over 70% decline[17]. Typically the lifespan of the house sparrow is 3 - 5 years in the wild and only about 20 percent of young live past their first year. Cold weather and food availability decide the longevity of Sparrows. *Plasmodium relictum*, a parasite infection, has also affected house sparrow demography across suburban London where sparrows have declined by 71% since 1995[18].

The genome sequencing efforts have been yielding contentious debate heralding short and long read chemistry over the last few years. While the peacock genome has yielded results [19], we have earlier contemplated asking questions on the Passer genome sequencing [20]. The avian genome project[21] [https://b10k.genomics.cn/ last accessed on September 20, 2024] representing several orders is aimed to resolve the phylogeny and collate data for testing hypotheses, understanding extinction and speciation of the birds besides demographic events, their roles of drift and selection in the divergence process [22]. Recently, Magallanes-Alba et al.have done a rapid genome functional annotation pipeline anchored to the house sparrow genome re-annotation and provided transcripts [23]. Our current genome sequencing of house sparrow is based on a muscle of the bird followed by assembly, annotation employing *in silico* approaches using a myriad of tools that could serve as a valuable resource to understand passerines' evolution, bio/ethno/geography and demography [24].

Methods
Sample collection and genome assembly
A bird wing (muscular tissue) was extracted from a house sparrow male which was found dead in the lawns of the Zoological Survey of India, Kolkata, and frozen immediately in liquid nitrogen. The sample was handled by Unipath Labs for DNA extraction, Illumina sequencing and the paired end raw reads were obtained by us for further processing. We performed genome assembly using Shovill (version 1.1.0) /SPAdes (3.15.4) [25]. Initially, we mapped *Gallus gallus* and *Taeniopygia guttata* and the alignment was not successful as upon further alignment using bowtie2, we found 99% mapped reads which led us to conduct downstream scaffolding. The gene completeness for Passer was assessed through Benchmarking Universal Single-Copy Orthologs (Busco version 5.5.0) [26] by using the orthologous genes in the *Gallus gallus* [chicken] genome. We performed a *de novo* assembly rather than a reference-based assembly with a specific set of nucleotide sequences used to represent an organism's genome [27]. In addition to BUSCO, we employed three tools for assembling the sample, *viz.* Megahit [28],

Maryland Super-Read Celera Assembler (MaSuRCA) [29] and Spades. A summary of methods can be found in Figure 1

Assembly graph construction: In order to systematically traverse the sequence space and find read overlaps which are crucial for contig assembly, we performed statistical analysis of the assembly to identify the best assembly among the three assemblers as discussed previously, to select the best contigs for further downstream analysis. Statistics of the assembly was done using QUAST to identify the N50 statistics [55]. The ensuing gaps were closed using a command line tool called gap closer, which allowed us to close 3150 gaps constituting the contigs present in our assembly. Prediction of genes in the genome was performed using Augustus [30], an *ab initio* hidden Markov model(HMM)-based gene prediction tool. The scaffolds of the SPAdes were rendered as the input for the Augustus tool against trained species-specific datasets, *viz. Gallus gallus* as a reference for alignment. The obtained GFF file was parsed to get predicted coding sequence and amino acid fasta sequences.

Figure 1: An overview of methodology employed for annotation and assembly of *Passer domesticus*.

Mitochondrial genome assembly: We extracted the mitochondrial genome from the reads and the assembly was performed using GetOrganelle toolkit which includes a number of scripts and libraries[31]. We further employed WGS read data for manipulating and disentangling assembly graphs, and generated reliable organelle genomes, accompanied by labeled assembly graphs and visualized using Bandage[32]. The prediction of the mitochondrial gene from the genome and annotation of the genome was performed using the Mitochondrial genome annotation Server (MITOS2) [33]. It uses BLAST searches with previously annotated protein sequences to predict the protein coding genes and further annotates the tRNAs apart from rRNAs present in the genome. We further performed a BLAST to identify the D-loop, a non-coding region in Mitochondria which acts as the promoter for both light and heavy chains and it is an important feature in mitochondria. We were able to acquire the order of the gene, gene names,start and end of the genes and obtained the values of the intergenic region between genes to construct the mitochondrial DNA. The mitochondrial DNA was constructed and visualized using

GenomeVX [34]. For ascertaining codon usage and Ka/Ks, we employed relative synonymous codon usage (RSCU) analysis using Molecular Evolutionary Genetics Analysis Version 11 [35]

Phylogenetic analysis: The phylogenetic tree construction was performed using MEGA11 followed by reaching consensus with 1Qq tree [36] and MAFFT [37] as we considered species of interest *Passer domesticus: Passer montanus , Passer_domesticus, Passer_ammodendri, Petronia_petronia, Pyrgilauda_blanfordi, Montifringilla_adamsi, Fringilla_polatzeki, Anthus_cervinus ,Motacilla_alba, Emberiza_fucata, Spizella_passerina, Agelaius_phoeniceus, Dives_dives, Euphagus_cyanocephalus, Quiscalus_quiscula, Chrysomus_icterocephalus, Pseudoleistes_guirahuro, Molothrus_badius, Gymnomystax_mexicanus ,streptopelia_orientalis_voucher_zjbj2* (outgroup).   After bootstrapping, we constructed a tree using the unweighted pair group method with arithmetic mean (UPGMA)with a gap penalty of -400 and a gap extension of 0.00. For phylogenetic tree construction, we chose Kimura as the substitution model with bootstrapping set to 100 and the final tree with MAFFT was constructed, While the scoring matrix was given as BLOSUM62 AND 200 PAM, the Jukes-cantor model was set as the substitution model and neighbor joining (NJ) for tree construction wherein bootstrapping was set to 100. For validating the tree inferred to understand the reliability and robustness, we employed a statistical method called the bootstrapping method. From the bootstrapped phylogenetic tree, we concluded that a node is well supported if it remained unchanged after 95 out of 100 iterations of removing one character and resampling our tree; a bootstrap value of 95% indicates this.Using the Mega X tool, we constructed a tree using the UPGMA as an algorithm for the MSA method with a gap penalty of -400 and a gap extension of 0.00. For phylogenetic tree construction, the substitution model chosen was kimura-2 parameter model and method used was maximum likelihood method and bootstrapping values were put to 100 and a tree was obtained. Then finally a tree with MAFFT alignment was constructed, where the MSA and the tree construction was performed with a penalty score rendered as 1.53. The sequences were aligned using progressive method with the tree algorithm as a default parameter.

Genome annotation, comparison and statistics
Gene annotation was performed with the protein sequence obtained from gene prediction to annotate the genes. Repetitive regions were identified and masked prior to gene prediction using Repeat Modeler [38], a *de novo* transposable element [TE] family identification and modeling package with Repeat Scout tool embedded for identifying the boundaries.[39]. We used the Shovill contigs that were generated as an input with 1391 sequences [1696224 bp] for downstream analyses and the resulting library was later checked with *Gallus gallus* repeat libraries. We compared and checked the annotation using Red [REpeat Detector: (Version 2018.09.10)], a rapid tool for detecting repeats *de novo* on the genomic scale[40].As we searched for prediction of incomplete genes at the sequence boundaries, we also aimed to predict complete genes. The Red GFF file was used as Augustus produced the fasta files containing predicted coding sequences and that were used for Pfamscan searches[Supplementary Table 3a, 3b, 3c) [41]. The resulting Augustus predictions were used as an input with Pfam-A HMM library manually downloaded in stockholm format. We also queried characteristic active site residues if any between the overlaps belonging to the same clans and further checked for functional annotation and domains using InterProScan [42] in inferring the Gene Ontology terms. The *Acanthisitta chloris* genome was obtained from CNGB for comparing the contigs with similarity [43]. Batch Entrez, Smith-Waterman algorithm using UVA FASTA (version 36.3.8i May, 2023) from local searches were made for predicting the proteins [https://fasta.bioch.virginia.edu/fasta/fasta_list.html last accessed on April 1, 2024]. GffCompare was employed to compare predicted sequences at different levels of granularity, thereby annotating the sequences based on their overlaps or proximity to reference annotation transcripts [44]. Sensitivity and precision metrics were computed with GTF file as an input for generating annotated files, yielding the "super-locus" for measuring the accuracy with true positives with other features like true negatives [TN] and false positives [FP] : Sensitivity = TP/[TP+FN] and Precision = TP/[TP+FP]. Finally, RefMap and

TMAP files [4] were obtained measuring the reference transcript that either fully or partially matches a transcript from the GTF and those columns in the file describing the most closely matching reference transcript respectively.

Comparative divergence time estimation
In order to estimate the divergence among selected clade and species, we used BEAST V2.0 [50], where Jmodeltest 2 [51] to decide evolutionary model based on BIC, where best suggested model were HKY I+G. we used mutation rate 0.018 substitution per site per Millions to estimate divergence [52]. The log files were checked using Tracer ensuring that ESS values were more than 200, and final trees were annotated with Treeannotator. Time divergences were estimated using data from all 20 taxa representing seven different families.

Results and Discussion

SPAdes achieved better genome assembly statistics when compared to Masurca and Megahit

We obtained the paired end files of 14 GB each which underwent quality check clearance wherein the Phred quality score was improved to 30 after FastP trims which were then assembled. SPAdes achieved better genome assembly statistics when compared to Masurca and Megahit [Table 1]. While the number of contigs were consistent between the Megahit and SPAdes, Masurca yielded a smaller number of contigs as it is not referenced for bird or avian genomes. However, the GC% was found to be consistent with all the three tools attributing to an average of 771 bp. On the other hand, N50 was comparatively better for SPAdes to the other two. In summary, *Passer domesticus* genome size was attributed to 922 MB taking SPAdes into consideration.

Tools>>>	Megahit	SPAdes	MaSuRCA
contigs	1018705	1000366	608253
Length	866 MB	922 MB	830 MB
Largest contig	83946	87277	78946
N50	1027	1343	740
N90	434	399	400

N's	0.00	0.00	0.00
GC %	41.74	41.93	41.49%

Table 1 : Assembly features extracted from three different tools, *viz.* Megahit, SPAdes and Masurca. From all the results the one with better N50 and avg value was the SPAdes

Gene prediction yielded 24152 genes across as many as 45634 transcripts

When genes were predicted using Augustus, and mapped from the BWA-mem of *Gallus gallus,* a considerable similarity of ca. 80.3 % was achieved further attributing to 24152 genes, 38972 introns and 45634 transcripts. The GFF file produced was then parsed to check downstream functional annotation tools using KOG, non redundant (NR) and Uniprot databases. The consensus hits were searched with "uniq sort" to find the most occurring species and considered the 10 best hits. We observed that NR yielded better results with *Passer montanus* occurring the most with a count of 7308, 1365 *Stutzerimonas stutzeri*, 939 *Melospiza melodia maxima*, 788 *Pyrgilauda ruficollis*, 780 *Hirundo rustica rustica*, 715 *Chloebia gouldiae*, 679 *Limosa lapponica baueri*, 652 *Onychostruthus taczazanowskii*, 457 *Lonchura striata domestica* and 319 *Motacilla alba.* What remains intriguing was that we detected the presence of bacterial sequences among the hits which was quite surprising. We argue that this could be due to the contamination caused during sequencing or any horizontal gene transfer events that might have happened over the years or birds could include a lot of symbiotic or pathogenic bacteria which can be part of their microbiota. We also observed the presence of many hypothetical proteins indicating that they are known unknown regions implying the need to annotate the genome significantly. There were the presence of many genes which are important for the adaptations, vocal learning, circadian rhythms, allee effects etc. BMAL1, clock genes were found to be associated in our list which are known to maintain the circadian rhythms of the species. We also found TLR4 which helps in identifying pathogens and initiation of immune responses besides HBAA ,HBB, HBA genes that are known for oxygen transport which helps in adaptation.

Mitochondrial genome yielded a genome of size 16,804 bp

Gene prediction and annotation predicted a total of 37 genes from which 13 genes were protein coding genes and 22 were tRNAs, 2 rRNAs and the non-coding region which is the D-loop was also identified.A labeled mitogenome was produced using GenomeVX online platform (Figure 2). The relative synonymous codon usage (RSCU) analysis of the codons was calculated in which the lowest preferences are AGA (R), AGG(R), ACG (T) [Table 2]. While they are less frequently used due to the lower level of those codon specific tRNAS, they might slow down the translation of proteins efficiently thus avoiding the use of these codons in efficient protein translation. We observed different patterns of selective pressure by analyzing the Ka/Ks ratios for the specified genes with strong purifying selection een in the majority of the genes, which have Ka/Ks ratios much less than 1, including cox1 (0.003), Cox3 (0.012), Nad2 (0.017), Nad3 (0.02), cob (0.02), NAD1 (0.034), Nad4l (0.035), Nad5 (0.03), and Nad6 (0.048). Given that the proteins these genes encode are essential to the organism's survival and functionality and that harmful mutations are progressively eliminated, it is likely that these genes are highly conserved. The Atp6 gene, on the other hand, appears to be under positive selection, indicating that natural selection may be favoring modifications in the protein product of this gene that may be advantageous with a Ka/Ks ratio of 1.03

[Table 3; Figure 2]. While the scoring matrix was given as BLOSUM62 AND 200 PAM, the Jukes-cantor model was set as the substitution model and NJ for tree construction wherein bootstrapping was set to 100 (Figure 3).

Figure 2 : Labeled mitogenome representing the 37 genes including the protein coding genes tRNAs,rRNAs and the non coding region which is the control region or the d- loop

Codon	RSCU	Codons	RSCU	Codons	RSCU	Codons	RSCU
UUU(F)	0.27	UCU(S)	0.53	UAU(Y)	0.44	UGU(C)	0.26
UUC(F)	1.73	UCC(S)	2.08	UAC(Y)	1.56	UGC(C)	1.74
UUA(L)	0.47	UCA(S)	1.83	UAA(*)	0.28	UGA(*)	2.69

UUG(L)	0.18	UCG(S)	0.35	UAG(*)	0.03	UGG(W)	1
CUU(L)	0.34	CCU(P)	0.45	CAU(H)	0.32	CGU(R)	0.49
CUC(L)	1.23	CCC(P)	1.42	CAC(H)	1.68	CGC(R)	1.15
CUA(L)	3.08	CCA(P)	1.84	CAA(Q)	1.64	CGA(R)	3.7
CUG(L)	0.71	CCG(P)	0.29	CAG(Q)	0.36	CGG(R)	0.49
AUU(I)	0.52	ACU(T)	0.58	AAU(N)	0.31	AGU(S)	0.14
AUC(I)	1.61	ACC(T)	1.71	AAC(N)	1.69	AGC(S)	1.07
AUA(I)	0.86	ACA(T)	1.67	AAA(K)	1.7	AGA(R)	0.08
AUG(M)	1	ACG(T)	0.04	AAG(K)	0.3	AGG(R)	0.08
GUU(V)	0.83	GCU(A)	0.48	GAU(D)	0.29	GGU(G)	0.38
GUC(V)	0.93	GCC(A)	2.22	GAC(D)	1.71	GGC(G)	0.95
GUA(V)	1.64	GCA(A)	1.19	GAA(E)	1.47	GGA(G)	1.9
GUG(V)	0.61	GCG(A)	0.11	GAG(E)	0.53	GGG(G)	0.77

Table 2: The RSCU values provided for the mitochondrial genome with the codons CUA (L), GCC(A), CGA(R) having highest values indicate more frequent occurrence. They might be preferred due to the abundance of tRNA contributing to efficient protein translation and protein stability reflecting evolutionary adaptation for protein synthesis. The codons that are translated efficiently are preferred in highly expressed genes

Genes	Ka/ks
cox2	0.81

atp6	1.03
atp8	0.429
cox3	0.012
nad3	0.02
nad4l	0.035
nad5	0.03
nad4	0.13
cob	0.02
nad6	0.048
nad1	0.034
nad2	0.017
cox1	0.003

Table 3 : ka/ks values indicating all the genes except atp6 have values less than one indicating they are undergoing strong purifying selection whereas the atp6 undergoes a positive selection with a value greater than one.

Avian genomes are ca. 70% smaller than mammalian genomes and Passerines being small are thought to maintain the contiguity that we have shown as compared to other vertebrates birds. We are limited by the amount of genomic attributes coming from non-coding regions, nevertheless, we argue that this could be because of the large number of hypothetical proteins which demand accurate annotation. While many genomes show syntenic, we are indefatigable to sequence more genomes that could provide us insights on adaptation, evolutionary mechanisms, substitutions, and importantly the circadian rhythms or electromagnetic effects that these birds might be associated with, thus drifting away from the urban life.

An approximate 25% of the genome has sequences matched to bacteria

The Passer genome showed considerable similarity [ca.25%] with bacterial sequences indicating that a large number of gut microbiota may have evented horizontal gene transfer [HGT] over the years [45]. This could be attributed to the bird's foraging on insects and small grains with bacteria inhabited in them could exist with endosymbionts and their invertebrate hosts, particularly insects. We argue that these could in principle be largely associated with modern evolutionary theory promoting such gene transfer. In addition, the alignment between these genomes is indicative of inversions/transversions which is in agreement with bird genomes [46]. The 710 contigs were mapped to the GenBank yielding 152 hypothetical proteins indicating that a large amount of proteins emerged from the known unknown regions, implying the need to annotate the genome significantly {SupplementaryTable 1). Among these, 252 sequences from protein domains match to animals while the rest match to bacteria. We observed a large number of Kelch domains in addition to a PHD finger 1 found in Histone-lysine N-methyltransferase

2C (KMT2C) and 2D (KMT2D) protein that is largely conserved. Also termed myeloid/lymphoid or mixed-lineage leukemia protein 3 (MLL3), it is associated with circadian factors contributing to genome-scale circadian transcription. These observations could indicate that a large number of genes possibly have undergone divergent evolution or the genes might have been lost and found through speciation. This fragmented genome is also indicative of BUSCO statistics from avian genomes that were reported earlier in Chaffinch genomes [47].

(a)

(b)

Figure 3 (a) :Phylogenetic tree representing all the 20 species, in which the passer domesticus is closely related to Passer montanus, *Passer ammodendri* ,Petronia petronia , *Pyrigilauda blandford*i and *Montifringilla adamsi* with a highly supported bootstrap values ie,100 where as all the other species such as *Motacilla alba ,Anthus cervinus* all belongs to the branches with bootstrapping value is 78 which is mediocre. The *streptopelia_orientalis_voucher_zjbj2* shows the most distant species in the tree which is the outgroup species with a low supported bootstrap value 53. (b)

Our annotation level of sequence representation is scanty but diverse as we found increased gene counts in comparison to *Gallus gallus* [53]. However, a large fraction of our genes remain unmapped and undiscovered in the Chicken genome. We deem that ours is a draft genome, nonetheless, small species such as Passerines still have no assigned function and their demand for evidence can only be classified through experimentation. While we attempted to create a good quality *de novo* based reference genome, the final assembly encompassing 922 MB is in agreement with the Elgvin et al. [54] works of a similar sized genome albeit the medium-density linkage map and order, the assembly scaffolds into chromosomes was meticulously done across several samples which in our case we couldn't as the sample acquisition has been a major limitation. Nevertheless, the genes were mapped to their reference genome for downstream analysis [Table 4].

S.No	Gene Name	NCBI accession	Contigs mapped
1	APC	NC_087512.1:52445657-52540315	NODE_14189_length_3730_cov_2.598686
2	HSDL2	NC_087512.1:69765571-69788616	NODE_20005_length_3394_cov_4.411215
3	ETFA	NC_087487.1:960048-989096	NODE_45445_length_2642_cov_2.323977
4	NOTCH2	NC_087480.1:32766603-32849060	NODE_2951_length_5290_cov_3.101669
5	NOTCH1	NC_087491.1:10550409-10592339	NODE_82034_length_2127_cov_3.214634
6	GLI3	NC_087474.1:51942133-52148229	NODE_5580_length_4619_cov_3.460810
7	WNT4	NC_087495.1:277607-294952	NODE_126693_length_1756_cov_2.892198
8	TIMP2	NC_087493.1:1552785-1561149	NODE_92905_length_2020_cov_2.762738
9	RBP1	NC_087484.1:25002122-25019513	NODE_19406_length_3423_cov_2.725344
10	PRK11A	NC_087512.1:13609738-13635725	NODE_48977_length_2574_cov_3.450941
11	GNAI1	NC_087478.1:9772700-9805360	NODE_5061_length_4724_cov_3.114698
12	CRHR1	NC_087500.1:437197-467780	NODE_32621_length_2944_cov_3.164632
13	ZFAND5	NC_087512.1:62564205-62578406	NODE_86240_length_2083_cov_4.130608
14	SECISBP2	NC_087512.1:42294093-	NODE_2582_length_5428_cov_

		42319272	3.587927
15	REEP5	NC_087512.1:52407038-52426127	NODE_48707_length_2579_cov_3.693046
16	MIA3	NC_087476.1:85960932-85989020	NODE_12474_length_3848_cov_3.843543
17	A2M	NC_087475.1:87627899-87657608	NODE_583_length_7451_cov_3.871576
18	KCNN2	NC_087512.1:68344752-68419555	NODE_16813_length_3560_cov_3.043928
19	BNC2	NC_087512.1:33461143-33796687	NODE_5333_length_4668_cov_3.558048

Tabel 4:comparison of genes that were mapped to the reference genome.

Seven superfamily domains are conserved between *Passer domesticus and Acanthisitta chloris*
The final set of contigs obtained for assembly were mapped against NCBI taxa [40] even as it matches to 14947 contigs with 769 of them making 100% query coverage with various hits from taxa. Amongst the superfamily members, intermediate filament protein, BASP1: Brain acid soluble protein 1, glycine rich LPXTG-anchored collagen-like adhesin and collagen with keratin and some hypothetical protein-FTsK translocase domains. What remains intriguing is the presence of probable chromatin-remodeling complex ATPase chain which is possibly associated with Allee effects that occur when individual fitness suffers at low population size or density [48]. We also argue that the genetic architecture and community-wide admixture could provide insights into their evolution [49]. On the other hand, keeping in view of the sample characteristics as a measure of genome completeness, we also mined the long read data from Oxford nanopore technologies [data not shown]. While the aforementioned statistics in the form of N50, N90, BUSCO are optimized for short reads, we were short of full potential in analyzing the genomic reads generated from long reads. Furthermore, the sequences generated need to be polished to overcome the mis-assemblies and have accurate scaffolding for generating more suitable short reads. On divergence, Columbidae appeared to have diverged early compared to others and was considered an outgroup in this analysis. Our sequenced sample, the house sparrow belonged to the family Passeridae. The divergence and expansion of major families began around 7-9 million years ago. Our studied sample formed a subclade with the Eurasian tree sparrow (P. m. saturatus) and Saxual sparrow (P. ammodendri), having a Time to Most Recent Common Ancestor (TMRCA) of 4.4 million years ago. Furthermore, our analysis yielded a TMRCA of 2.9 million years ago for *Passer domesticus* and P. m. saturates (Figure 4).

Figure 4: A time-calibrated tree with 20 selected taxa using complete mitochondrial DNA sequences The tree was generated using Beast v2.5 with 20 million MCMC runs. Each respective family is represented by a different color, and the node labels indicate the age in million years.

Conclusions

We provide here a high-quality whole genome sequencing and assembly for the house sparrow, *Passer domesticus* which could serve us to ask a large number of questions from genomics. Our annotation serves as a useful resource to check adaptation, divergence and speciation. The orthologous annotation and protein mapping, bacterial correlates show that the HGT events may not be ruled out.

Acknowledgments: PS gratefully acknowledges Department of Forestry, Governments of Rajasthan and India, Zoological Survey of India, and Birla Institute of Scientific Research for providing the apriori permission for collection of bird's sample.

Authors' contributions: PS ideated the project with GC and PBK. VK,SS, SP, AT, GS, DV and PS contributed equally to the work. All the other authors worked towards improvement of tables and figures. PS, GC and PBK proofread the manuscript.

Data Availability: The genome assembly has been deposited at NCBI under BioProject ID: PRJNA1027087.

Competing interests: None

References:
1. Zhang, G., Parker, P., Li, B., Li, H. and Wang, J., 2012. The genome of Darwin's finch [*Geospiza fortis*]. *GigaScience*. Database. http://dx.doi.org/10.5524/100040

2. Jarvis, E.D., Mirarab, S., Aberer, A.J., Li, B., Houde, P., Li, C., Ho, S.Y., Faircloth, B.C., Nabholz, B., Howard, J.T. and Suh, A., 2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science*, 346(6215), 1320-1331.
3. Poelstra, J.W., Vijay, N., Bossu, C.M., Lantz, H., Ryll, B., Müller, I., Baglione, V., Unneberg, P., Wikelski, M., Grabherr, M.G. and Wolf, J.B., 2014. The genomic landscape underlying phenotypic integrity in the face of gene flow in crows. *Science*, 344(6190), 1410-1414.
4. Frankl-Vilches, C., Kuhl, H., Werber, M., Klages, S., Kerick, M., Bakker, A., de Oliveira, E.H., Reusch, C., Capuano, F., Vowinckel, J. and Leitner, S., 2015. Using the canary genome to decipher the evolution of hormone-sensitive gene regulation in seasonal singing birds. *Genome Biology*, 16, 1-25.
5. Friis, G., Fandos, G., Zellmer, A.J., McCormack, J.E., Faircloth, B.C. and Milá, B., 2018. Genome□wide signals of drift and local adaptation during rapid lineage divergence in a songbird. *Molecular Ecology*, 27(24), 5137-5153.
6. Louha, S., Ray, D.A., Winker, K. and Glenn, T.C., 2020. A high-quality genome assembly of the North American Song Sparrow, *Melospiza melodia*. G3: Genes, Genomes, Genetics, 10(4), 1159-1166.
7. Peñalba, J.V., Deng, Y., Fang, Q., Joseph, L., Moritz, C. and Cockburn, A., 2020. Genome of an iconic Australian bird: High□quality assembly and linkage map of the superb fairy□wren (*Malurus cyaneus*). *Molecular Ecology Resources*, 20(2), 560-578.
8. Ducrest, A.L., Neuenschwander, S., Schmid□Siegert, E., Pagni, M., Train, C., Dylus, D., Nevers, Y., Warwick Vesztrocy, A., San□Jose, L.M., Dupasquier, M. and Dessimoz, C., 2020. New genome assembly of the barn owl (*Tyto alba alba*). *Ecology and Evolution*, 10(5), 2284-2298.
9. Wang, W., Wang, F., Hao, R., Wang, A., Sharshov, K., Druzyaka, A., Lancuo, Z., Shi, Y. and Feng, S., 2020. First de novo whole genome sequencing and assembly of the bar-headed goose. *PeerJ*, 8, 8914.
10. Zhang, G., 2015. Bird sequencing project takes off. *Nature*, 522(7554), 34-34.
11. Feng, S., Stiller, J., Deng, Y., Armstrong, J., Fang, Q.I., Reeve, A.H., Xie, D., Chen, G., Guo, C., Faircloth, B.C. and Petersen, B., 2020. Dense sampling of bird diversity increases power of comparative genomics. *Nature*, 587(7833), 252-257.
12. Rahmani, A.R., 2008. Flight to extinction. Spectrum, *The Tribune*.
13. JD, S.S., 2003. The decline of the House Sparrow: a review. *Brit. Birds*, 96, 439-446.
14. Crick, H.Q., Robinson, R.A., Appleton, G.F., Clark, N.A. and Rickard, A.D., 2002. Investigation into the causes of the decline of starlings and house sparrows in Great Britain. *BTO Research Report*, 290, 1-305.
15. De Coster, G., De Laet, J., Vangestel, C., Adriaensen, F. and Lens, L., 2015. Citizen science in action—Evidence for long-term, region-wide House Sparrow declines in Flanders, Belgium. *Landscape and Urban Planning*, 134, 139-146.
16. Mohring, B., Henry, P.Y., Jiguet, F., Malher, F. and Angelier, F., 2021. Investigating temporal and spatial correlates of the sharp decline of an urban exploiter bird in a large European city. *Urban Ecosystems*, 24, 501-513.
17. Gulati, V., 2005. House sparrow on verge of extinction. *Tribune News Service*, Chandigarh, India.
18. Dadam, D., Robinson, R.A., Clements, A., Peach, W.J., Bennett, M., Rowcliffe, J.M. and Cunningham, A.A., 2019. Avian malaria-mediated population decline of a widespread iconic bird species. *Royal Society Open Science*, 6(7), 182197.
19. Dhar, R., Seethy, A., Pethusamy, K., Singh, S., Rohil, V., Purkayastha, K., Mukherjee, I., Goswami, S., Singh, R., Raj, A. and Srivastava, T., 2019. De novo assembly of the Indian blue peacock (*Pavo cristatus*) genome using Oxford Nanopore technology and Illumina sequencing. *Gigascience*, 8(5), 038.

20. Suravajhala, P. and Acharya, K., 2014. WHITHER GENOME SEQUENCING OF *PASSER*? *ECB*, 3(11), 1086-1087.

21. The Bird 10 genome project: https://b10k.genomics.cn/ last accessed on July 8, 2024.

22. Sendell-Price, A.T., Ruegg, K.C. and Clegg, S.M., 2020. Rapid morphological divergence following a human-mediated introduction: The role of drift and directional selection. *Heredity*, 124(4), 535-549.

23. Magallanes-Alba, M.E., Baricalla, A., Rego, N., Brun, A., Karasov, W.H. and Caviedes-Vidal, E., 2023. Rapid genome functional annotation pipeline anchored to the house sparrow (*Passer domesticus*, Linnaeus 1758) genome reannotation. *Biology Methods and Protocols*, 8(1), 013.

24. Illera, J.C., Rando, J.C., Rodriguez☐Exposito, E., Hernández, M., Claramunt, S. and Martín, A., 2018. Acoustic, genetic, and morphological analyses of the Canarian common chaffinch complex *Fringilla coelebs* ssp. reveals cryptic diversification. *Journal of Avian Biology*, 49(12).

25. Seemann, T., 2017. Shovill: faster SPAdes assembly of Illumina reads. *Tilgjengelig fra Tilgjengelig fra*.

26. Seppey, M., Manni, M. and Zdobnov, E.M., 2019. BUSCO: assessing genome assembly and annotation completeness. *Gene Prediction: Methods and Protocols*, 227-245.

27. Li D, Liu CM, Luo R, Sadakane K, Lam TW. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics. 2015 May 15;31(10):1674-6. doi: 10.1093/bioinformatics/btv033. Epub 2015 Jan 20. PMID: 25609793.

28. Kitts PA, Church DM, Thibaud-Nissen F, Choi J, Hem V, Sapojnikov V, Smith RG, Tatusova T, Xiang C, Zherikov A, DiCuccio M, Murphy TD, Pruitt KD, Kimchi A. Assembly: a resource for assembled genomes at NCBI. Nucleic Acids Res. 2016 Jan 4;44(D1):D73-80. doi: 10.1093/nar/gkv1226. Epub 2015 Nov 17. PMID: 26578580; PMCID: PMC4702866.

29. Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. The MaSuRCA genome assembler. Bioinformatics. 2013 Nov 1;29(21):2669-77. doi: 10.1093/bioinformatics/btt476. Epub 2013 Aug 29. PMID: 23990416; PMCID: PMC3799473.

30. Stanke, M. and Waack, S., 2003. Gene prediction with a hidden Markov model and a new intron submodel. *Bioinformatics-Oxford*, 19(2), 215-225.

31. Jin, JJ., Yu, WB., Yang, JB. *et al.* GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of organelle genomes. *Genome Biol* 21, 241 (2020). https://doi.org/10.1186/s13059-020-02154-5

32. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de novo genome assemblies. Bioinformatics. 2015 Oct 15;31(20):3350-2. doi: 10.1093/bioinformatics/btv383. Epub 2015 Jun 22. PMID: 26099265; PMCID: PMC4595904.

33. Donath A, Jühling F, Al-Arab M, Bernhart SH, Reinhardt F, Stadler PF, Middendorf M, Bernt M. Improved annotation of protein-coding genes boundaries in metazoan mitochondrial genomes. Nucleic Acids Res. 2019 Nov 18;47(20):10543-10552. doi: 10.1093/nar/gkz833. PMID: 31584075; PMCID: PMC6847864.

34. Conant GC, Wolfe KH. GenomeVx: simple web-based creation of editable circular chromosome maps. Bioinformatics. 2008 Mar 15;24(6):861-2. doi: 10.1093/bioinformatics/btm598. Epub 2008 Jan 28. PMID: 18227121

35. Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Mol Biol Evol. 2021 Jun 25;38(7):3022-3027. doi: 10.1093/molbev/msab120. PMID: 33892491; PMCID: PMC8233496.

36. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015 Jan;32(1):268-74. doi: 10.1093/molbev/msu300. Epub 2014 Nov 3. PMID: 25371430; PMCID: PMC4271533.

37. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002 Jul 15;30(14):3059-66. doi: 10.1093/nar/gkf436. PMID: 12136088; PMCID: PMC135756.

38. Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C. and Smit, A.F., 2020. RepeatModeler2 for automated genomic discovery of transposable element families. Proceedings of the National Academy of Sciences, 117(17), 9451-9457.

39. Bourque, G., Burns, K.H., Gehring, M., Gorbunova, V., Seluanov, A., Hammell, M., Imbeault, M., Izsvák, Z., Levin, H.L., Macfarlan, T.S. and Mager, D.L., 2018. Ten things you should know about transposable elements. Genome Biology, 19, 1-12.

40. Girgis, H.Z., 2015. Red: an intelligent, rapid, accurate tool for detecting repeats de-novo on the genomic scale. BMC Bioinformatics, 16, 1-19

41. Wang, L., Park, H.J., Dasari, S., Wang, S., Kocher, J.P. and Li, W., 2013. CPAT: Coding-Potential Assessment Tool using an alignment-free logistic regression model. Nucleic Acids Research, 41(6), 74.

42. Jones, P., Binns, D., Chang, H.Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G. and Pesseat, S., 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics, 30(9), 1236-1240.

43. Zhang, G., Li, B., Li, C., Gilbert, M.T.P., Jarvis, E.D., 2014. The Avian Genome Consortium; Wang J Genomic data of the Rifleman (Acanthisitta chloris). GigaScience Database. http://dx.doi.org/10.5524/101015

44. Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M. J., Salzberg, S. L., Wold, B. J., & Pachter, L. (2010). Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. Nature Biotechnology, 28(5), 511â515. doi:10.1038/nbt.1621

45. Dunning Hotopp JC. Horizontal gene transfer between bacteria and animals. Trends Genet. 2011 Apr;27(4):157-63. doi: 10.1016/j.tig.2011.01.005. Epub 2011 Feb 18. PMID: 21334091; PMCID: PMC3068243.

46. Stoltzfus, A. and Norris, R.W., 2016. On the causes of evolutionary transition: transversion bias. *Molecular Biology and Evolution*, 33(3), 595-602.

47. Recuerda, M., Vizueta, J., Cuevas-Caballé, C., Blanco, G., Rozas, J. and Milá, B., 2021. Chromosome-level genome assembly of the common chaffinch (Aves: *Fringilla coelebs*): a valuable resource for evolutionary biology. *Genome Biology and Evolution*, 13(4), 034.

48. Gascoigne, J.C., 2003. A theoretical and experimental study of Allee effects. The College of William and Mary.

49. Enbody, E.D., Sendell-Price, A.T., Sprehn, C.G., Rubin, C.J., Visscher, P.M., Grant, B.R., Grant, P.R. and Andersson, L., 2023. Community-wide genome sequencing reveals 30 years of Darwin's finch evolution. *Science*, 381(6665), 6218.

50. Bouckaert R, Vaughan TG, Barido-Sottani J, et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. PLoS Comput Biol. 2019;15(4):e1006650. Published 2019 Apr 8. doi:10.1371/journal.pcbi.1006650.

51. Darriba, D., Taboada, G., Doallo, R. et al. jModelTest 2: more models, new heuristics and parallel computing. Nat Methods 9, 772 (2012). https://doi.org/10.1038/nmeth.2109.

52. Lerner HR, Meyer M, James HF, Hofreiter M, Fleischer RC. Multilocus resolution of phylogeny and timescale in the extant adaptive radiation of Hawaiian honeycreepers. *Curr Biol*. 2011;21(21):1838-1844. doi:10.1016/j.cub.2011.09.039

53. Warren WC, Hillier LW, Tomlinson C, Minx P, Kremitzki M, Graves T, Markovic C, Bouk N, Pruitt KD, Thibaud-Nissen F, Schneider V, Mansour TA, Brown CT, Zimin A, Hawken R, Abrahamsen M, Pyrkosz AB, Morisson M, Fillon V, Vignal A, Chow W, Howe K, Fulton JE, Miller MM, Lovell P, Mello CV, Wirthlin M, Mason AS, Kuo R, Burt DW, Dodgson JB, Cheng HH. A New Chicken

Genome Assembly Provides Insight into Avian Genome Structure. G3 (Bethesda). 2017 Jan 5;7(1):109-117. doi: 10.1534/g3.116.035923.

54. Elgvin TO, Trier CN, Tørresen OK, Hagen IJ, Lien S, Nederbragt AJ, Ravinet M, Jensen H, Sætre GP. The genomic mosaicism of hybrid speciation. Sci Adv. 2017 Jun 14;3(6):e1602996. doi: 10.1126/sciadv.1602996.

55. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome assemblies. Bioinformatics. 2013 Apr 15;29(8):1072-5. doi: 10.1093/bioinformatics/btt086. Epub 2013 Feb 19. PMID: 23422339; PMCID: PMC3624806.