An Alzheimer's disease-associated common regulatory variant in PTK2B has causal effects on microglial function

Erica Bello[1,2,3*], Kathleen Long[1], Sho Iwama[1], Juliette Steer[1,2], Sarah Cooper[1,2], Kaur Alasoo[1,4], Natsuhiko Kumasaka[1,5], Jeremy Schwartzentruber[1,2], Nikolaos I Panousis[1], Andrew Bassett[1,2,*]

[1] Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SA, UK
[2] OpenTargets, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SA, UK
[3] Milner Therapeutics Institute, Jeffrey Cheah Biomedical Centre, University of Cambridge, Cambridge, CB2 0AW, UK
[4] Institute of Computer Science, University of Tartu, Tartu, Estonia
[5] National Center for Child Health and Development, Tokyo, Japan

*To whom correspondence should be addressed. Tel: +44(0)1223 494933; Email: ab42@sanger.ac.uk, eb956@cam.ac.uk

Abstract

Genome-wide association studies (GWAS) are revealing an ever growing number of genetic associations with disease, but identifying and functionally validating the causal variants underlying these associations is very challenging and has only been done for a vanishingly small number of variants. Here we validate a causal single nucleotide polymorphism (SNP) associated with an increased risk of Alzheimer's disease (AD) in an intronic enhancer of the *PTK2B* gene, by engineering it into human induced pluripotent stem cells (hiPSCs). Upon differentiation to macrophages and microglia, this variant shows effects on chromatin accessibility of the enhancer and gene expression changes of the PTK2B gene, and results in global changes to the transcriptome and phenotype of these cells. Expression of interferon gamma responsive genes including chemokine transcripts and their protein products are altered, and chemotaxis of the resulting microglial cells is affected. This variant thus causes disease-relevant transcriptomic and phenotypic changes, and we propose that it acts by altering microglia reactivity, consistent with the role of these cells in progression of AD.

Introduction

Alzheimer's disease (AD) is the most common form of dementia, and has a strong genetic component estimated to be 60-80%[1]. AD is characterised by the presence of β-amyloid-containing plaques (Aβ) and intracellular neurofibrillary tangles of hyperphosphorylated tau, accompanied by neuroinflammation, neuronal, and synapse loss[2]. Multiple GWAS studies have collectively identified more than 80 genetic loci associated with disease[3–7], the vast majority of which do not affect protein coding sequence. Due to linkage disequilibrium at these loci, it is difficult to identify the causal variants. Multiple computational methods have been developed to prioritise variants that cause disease including statistical fine mapping, co-localisation with expression or chromatin accessibility quantitative trait loci (eQTL, caQTL), overlap with enhancer regions identified by e.g. assay for transposase accessible chromatin (ATAC) or transcription factor binding in disease relevant cell types. However, none of these methods are able to unambiguously assign causal variants and, given the vanishingly small number of truly validated variants, it is difficult to benchmark their effectiveness. Experimental methods have also been developed to identify or prioritise variants such as massively parallel reporter assays (MPRA[8]) and genome engineering-based interrogation of enhancers (GenIE[9], GenIE-ATAC[10]) but these are limited to understanding the effects of variants on transcription or chromatin accessibility and do not directly demonstrate the downstream effects of these polymorphisms. Genome engineering of isogenic human cell models, such as differentiated hiPSCs differing in a single variant, provide an attractive means for understanding the role of specific SNPs on cell function and have been successful in understanding the role of strong effect familial mutations[11,12,13]. However, such systems have yet to be successfully applied to more than a handful of common genetic variants[14,15,16] and their effects are often confounded by variation between edited clones and during differentiation[17].

Neuroinflammation has been shown to be a crucial factor in many neurodegenerative disorders, especially AD[18]. Microglia are resident immune cells of the central nervous system and perform various functions under both homeostatic and disease conditions [19]. Microglia play important roles in synaptic remodelling, synaptic pruning and phagocytosis of dead neurons[20]. Microglia have been implicated in AD by human genetics studies because of specific expression of AD-related genes in these cells[21–23], colocalisation of microglial eQTLs with AD GWAS hits[24,25] and implication of rare causal variants in microglial-specific genes, such as TREM2[26] and CD33[26,27]. Nevertheless, the exact role of microglia in AD pathogenesis is still unclear. Aβ and tau aggregates are able to activate microglia and elicit an inflammatory response that can have two functions: it can be neuroprotective by increasing Aβ or tau clearance, but it can also contribute to Aβ and tau production and spreading and induce neurodegeneration and synaptic loss[28,29]. Microglia are known to be involved in clearance of aggregated proteins such as Aβ[30] and imbalanced microglial activation or insufficient microglial phagocytic capacity can accelerate the

accumulation of extracellular Aβ[31]. However, depletion of microglia in amyloid precursor
protein-transgenic mouse models was found to have little effect on plaque formation or
maintenance. Microglia have also been found to potentially contribute to tauopathy by
spreading tau protein via exosomes across brain regions[32] as well as its uptake and
degradation[29].   Moreover, sustained microglia activation appears to contribute to
neurodegeneration by impacting synaptic plasticity and causing neuronal damage[33,34]. As
well as microglia, other myeloid cell types, such as infiltrating monocytes[35,36], choroid
plexus[37] and perivascular macrophages[38–40], have also been found to play a role in AD
pathogenesis.

PTK2B is a non-receptor tyrosine kinase that is expressed in neurons, microglia, astrocytes,
monocytes and tissue-resident macrophages, and has diverse physiological and
pathological roles including cell adhesion, cell migration, inflammatory responses, tumor
invasiveness[41], neuronal development and plasticity[42,43]. PTK2B is a member of the focal
adhesion kinase (FAK) subfamily and after autophosphorylation it recruits Src-family
kinases[44] and interacts with multiple partners to activate various downstream signalling
pathways, such as the MAPK/ERK pathway[42]. In neurons, PTK2B can be activated following
stimulation of glutamate receptors[45], neuronal depolarization[46] and increased intracellular
free Ca2+[47].   In neurons PTK2B is implicated in synaptic plasticity[48] and neurite
outgrowth[49]. In non-neuronal cells, such as osteoclasts[50], macrophages[51–53] and
monocytes[54], PTK2B is involved in cell migration and regulation of actin cytoskeleton
downstream of integrins. In monocytes, the ROS-sensitive calcium channel TRPM2 can
activate the PTK2B/ERK pathway, leading to nuclear translocation of nuclear factor-kB and
increased chemokine production[55]. This pathway is activated by Aβ in microglia, where
PTK2B is autophosphorylated and generates a positive feedback loop to further activate
TRPM2[56]. PTK2B has been implicated in AD through a number of GWAS studies[4,6,57–64]. As
well as being involved in response to Aβ, PTK2B co-localizes with hyperphosphorylated Tau
in the brain of AD patients and in a mouse model, and it may contribute to Tau
phosphorylation[65,66]. Evidence suggests that genetic deletion of Ptk2b rescues a number
of Aβ-associated phenotypes, such as memory impairment, synapse loss and impaired
synaptic plasticity, in APP/PS1 animal models of AD[67]. Data collected from hemizygous
PS19 (MAPT P301S, 1N4R) transgenic mice crossed with Ptk2b[−/−] animals, points to a
protective role for Ptk2b against Tau phosphorylation and Tau-induced behavioral deficits,
in part through the suppression of LKB1 and p38 MAPK[68].
Several lines of evidence point to PTK2B as an important player in the pathophysiology of
AD but much remains to be learnt about the exact role of PTK2B in AD pathogenesis.

Results
Prioritisation of a putative Alzheimer's disease causal variant at the PTK2B locus
GWAS have identified two independent genetic associations at the PTK2B-CLU locus[57]. The
signal over the PTK2B gene colocalises with both an eQTL and caQTL in hiPSC-derived

macrophages (Fig 1a)[69]. Re-analysis of publicly available datasets[25,69,70] shows that the risk
haplotype is linked to both a downregulation of PTK2B gene expression in hiPSC-derived
macrophages (Fig 1b and 1c) and an increase in chromatin accessibility at an enhancer
within intron 6 of the PTK2B gene (Fig 1d). Interestingly, in this case there is a single variant
in the credible set (rs28834970) that lies within the ATAC peak that is changed in
accessibility in the caQTL analysis, strongly suggesting that this variant is causal (C being
the risk allele). Note that there is some inconsistency in published literature on the
directionality of these effects[69,25,70], but this data has been reanalysed in conjunction with
the authors, and in additional hiPSC-derived macrophage datasets so we are confident of
the directionality of these changes described here. This confirmed that the C allele of
rs28834970 decreases expression of *PTK2B* in hiPSC-derived macrophages (Fig 1b and Fig
1c, whereas in monocytes and one study of primary microglia, the C allele of rs28834970
is associated with increased expression of *PTK2B* (Supp Fig 1a)[69].

Characterisation of rs28834970 in hiPSC-derived macrophages and microglia

In a recent study this variant was found to be associated with an increase in PTK2B
expression only in blood of AD patients and not in any brain region examined[71]. Given the
important role of PTK2B in myeloid cells and the evidence suggesting this variant modifies
expression of PTK2B in this cell type, we decided to investigate the role of this variant in
macrophages and microglia. We edited the C allele of rs28834970 into a well characterised
hiPSC line (KOLF2_C1) that was homozygous for the T allele at this position (and thus the
non-risk haplotype) using CRISPR-mediated homology directed repair[72] (Suppl Fig 1b and
Methods). This allows us to unambiguously study the role of rs28834970 independently
of the other variants in linkage disequilibrium. We subsequently analysed three
independent homozygous (C/C) clones and two unedited (T/T) clones, one of which had
been through the editing process, and the second being the parental line. These were
differentiated in three biological replicates into either macrophages[73,74] or microglia[75,76]
followed by a number of assays including transcriptomics (RNAseq), chromatin
accessibility (ATACseq) and quantitative digital RT-PCR (dd-qRT-PCR) to measure gene
expression (Fig 1e). Differentiation was highly reproducible between the two different
genotypes by FACS for macrophage markers CD14, CD16 and CD206 (Supp Fig 2a) and
microglia markers CD11b and CD45 (Supp Fig 2b). This was further validated by analysis of
macrophage and microglial marker gene expression which were mostly consistent with the
differentiation trajectory and did not cluster by genotype (Supp Fig 2c). Similarly, there did
not appear to be genotype-dependent change in expression of canonical markers of
macrophage or microglial activation (Supp Fig 2d).

rs28834970 reduces PTK2B expression by creating a novel CEBPB binding site in hiPSC-derived macrophages and microglia

155   We then analysed the effect of the rs28834970 allele on *PTK2B* expression and function of
156   the intronic enhancer element. Consistent with the prediction from the eQTL analysis in

hiPSC-derived macrophages, both RNAseq and digital droplet qPCR (dd-qPCR) analysis showed that expression of *PTK2B* was modestly reduced by introduction of the C allele in macrophages (RNAseq: p=0.019 and logFC=-0.2, ddqPCR: p=0.0045 and logFC=-0.39) (Fig 2a) and microglia (RNAseq: p=0.06 logFC=-0.57, ddqPCR: p=0.22 and logFC=-0.71) (Fig 2b). The effect was consistent between RNAseq and dd-qPCR, and across microglia and macrophages, but was variable in magnitude. This is consistent with the small effect size in the eQTL analysis (Fig 1b and Fig 1c), and is likely a feature of the majority of common variants that would be expected to have only small or context-specific effects on target gene expression[16].

Given that an enhancer can have long range effects and mapping of chromosomal interactions at this locus in microglia had shown that enhancers harboring AD-risk variants interacted with active promoters of not only PTK2B and CLU, but also TRIM35 and CHRNA2[77], we also analysed expression of genes in a 600 kb window around rs28834970 (Supp Fig 3a). The effect on PTK2B expression was the most consistent change in microglia and macrophages in this window, but there was additionally a slight upregulation of CLU expression specifically in macrophages (logFC=0.72 p-value=0.03) which was not observed in microglia (Supp Fig 3b). This implies that in macrophages, there may be an effect of the rs28834970 enhancer on CLU expression as well as PTK2B. We also analysed changes in splicing of the PTK2B gene and saw only small changes between the two genotypes (Supp Fig 3c), none of which affected splicing of the exons near to the variant.

We next analysed the effect of rs28834970 on chromatin accessibility, which showed that the C allele created a novel region of accessible chromatin in both macrophages and microglia (Fig 2c) which was consistent with the expectations from the caQTL analysis in hiPSC-derived macrophages (Fig 1d). Analysis of chromatin accessibility within a 600 kb window around the SNP showed that there was no change in most peaks, and the only statistically significant change consistent between macrophages and microglia was the new peak over rs28834970 (macrophages: p=1.65e-16 logFC=2.71, microglia: p=4.19e-12 logFC=2.77) (Supp Fig 3d). Peaks within the CHRNA2 and CLU genes were significantly reduced in microglia, and an intergenic peak was increased in macrophages, but these were not consistent between the two cell types, or with the expression changes of these genes (Supp Fig 3d and Supplementary Table 1).

Further analysis of the sequence around rs28834970 showed that the C allele increased the probability of binding of transcription factors CEBPA and CEBPB (Fig 2d). We thus analysed CEBPB binding to chromatin by CUT&RUN, which showed an increased binding for the C allele of rs28834970 (p=0.04 logFC=2.09)(Fig 2c), consistent with CEBPB binding being responsible for the change in chromatin accessibility. Indeed, in a genome-wide analysis, this was the only region that showed a significant change in CEBPb binding between the C/C and T/T alleles at rs28834970 (Fig 2e)(Supplementary Table 2). CEBPA

198 and CEBPB have previously been shown to have both activator and repressor functions[78],
199 and thus we propose that this novel binding event represses expression of *PTK2B*.
200
201 Taken together, these results validate that rs28834970 within the PTK2B locus is the causal
202 variant and acts through introducing a novel CEBPB binding site that increases chromatin
203 accessibility at the intronic enhancer, and results in repression of PTK2B expression.
204
rs28834970 causes strong effects on the transcriptome including chemokine expression
We next analysed the effects of editing rs28834970 on genome-wide gene expression in
macrophages and microglia. Principal component analysis showed that although there was
a strong clone and batch effect, the two genotypes were well separated by the second
principal component (Fig 3a). Similar results were seen with analysis of global chromatin
accessibility, where genotypes were separated by the second principal component (Supp
Fig 4a). Analysis of the differentially expressed genes between genotypes showed 690
downregulated and 428 upregulated genes in macrophages, and 760 downregulated and
328 upregulated in microglia (Fig 3a) (Supplementary Table 3). Importantly, there was a
significant (p<0.0001) overlap in the differentially expressed genes between the two cell
types (Supp Fig 4b), further corroborating that we were identifying true effects of the
genotype, not simply variation between edited clones or differentiation artefacts.
217
We performed gene set enrichment analysis of the differentially expressed genes relative
to all genes expressed in the respective cell type. This showed that a number of gene
ontology pathways relevant to microglial or macrophage function were enriched in the
downregulated gene set including chemokine-mediated signalling, cellular response to
interferon gamma (IFNγ) and migration and chemotaxis (Fig 3b, Fig 3c). These pathways
were frequently overlapping between microglia and macrophages (Supp Fig 4c), again
supporting that these changes were genotype-dependent. Conversely, there were no
obviously relevant gene ontology (GO) enrichments in the upregulated gene set (Supp Fig
4d) which may be due to the smaller number of genes analysed.
227
Further analysis of the downregulated genes confirmed that there was considerable
overlap between microglia and macrophages across the different GO categories (Fig 3d).
Of note, we found that the Macrophage Receptor With Collagenous Structure (*MARCO*)
gene was among the top 3 downregulated genes in both microglia and macrophages with
the C/C allele (Supp Fig 4e). Given their role in microglial and macrophage function, we
focused on chemokines, MHC complex proteins and surface receptors and validated their
expression by dd-qPCR. All of the chosen genes showed some level of downregulation as
expected in microglia, and all except one (HLA-DQA1) were downregulated in
macrophages (Fig 3e), with effect sizes generally consistent between dd-qPCR and RNAseq.
237
238

rs28834970 causes changes in chemokine release and migration

Guided by the consistent effects on chemokine and migration-related gene expression in both macrophages and microglia, we next analysed the release of 13 chemokines using a multiplexed assay on microglial cell supernatants. In unstimulated cells, we saw significant reduction in production of 10 of the 13 chemokines tested with the C/C allele at rs28834970, consistent with the results of the transcriptomic analysis (Fig 4a). CCL22 was unexpectedly upregulated, and CXCL2 and CXCL5 were unchanged (Fig 4a).

Given that several of these differentially expressed chemokines are produced by microglia in response to stimuli such as IFNγ and lipopolysaccharide (LPS), we also analysed whether their production was altered in microglia harbouring the C/C allele in the presence of these stimuli. As expected, both treatments increased production of most (10 of 12) of the chemokines (Supp Fig 5a and Supp Fig 5b). CCL1 production was increased by LPS stimulation in microglia harbouring the C/C allele but not in microglia with the T/T allele (Supp Fig 5b). Similar to what was observed in unstimulated cells, production of most of the chemokines was reduced in the C/C allele at rs28834970 when microglia were stimulated with IFNγ, but the magnitude of the reduction was greatly reduced, such that only 5 of the 12 chemokines were significant (Fig 4b). A similar response was seen upon stimulation with LPS (Supp Fig 5c). Taken together, these data suggest that the C/C allele results in a lower basal chemokine release, and in the presence of stimuli such as IFNγ, this reduction becomes less marked.

We further investigated the effect of rs28834970 on microglial migration and chemotaxis using a transwell assay in the presence or absence of IFNγ and with or without a chemoattractant relevant to microglia, complement 5a (C5a). Microglia harbouring the unedited (T/T) allele showed chemotaxis towards C5a that was inhibited by IFNγ treatment (Fig 4c). Microglia harbouring the C/C alleles at rs28834970 showed no significant chemotaxis towards C5a or inhibition of migration following IFNγ treatment (Fig 4c). Taken together, these data suggest that microglia harbouring the C/C allele are less sensitive to the chemoattractant C5a and to the inhibitory effect of IFNγ on migration.

In summary, we show that the C allele of rs28834970 lowers chemokine release by microglia, reduces chemotaxis towards C5a and alters their response to IFNγ stimulation, perhaps by altering microglia reactivity.

Discussion

One of the major challenges in interpreting GWAS is to understand the genes and causal variants underlying the genetic association with disease. The use of isogenic cell lines generated by CRISPR in model systems such as hiPSC differentiated cell types offer one way to dissect the role of individual variants in cellular phenotypes independently of other variants in linkage disequilibrium. However, common regulatory variants that are

identified in GWAS have rarely been linked to changes in cellular 'omics or phenotypes. Here we identify a likely causal variant from comparison of Alzheimer's disease GWAS, eQTL and caQTL studies and engineer this using CRISPR/Cas9 into hiPSCs to generate isogenic cell lines differing in the single variant. We show that the C/C allele of rs28834970 likely introduces a novel CEBPB binding site in an intron of PTK2B that results in transcriptional repression of PTK2B expression. Even though the downregulation of PTK2B caused by the variant does not reach statistical significance in our models, it is consistent with previous reports that this variant causes a decrease in PTK2B expression in monocytes[79] and macrophages[25,69,70] (Fig 1b and c). This could be due to the small effect size making it difficult to study with a limited number of isogenic cell clones or compensatory regulatory mechanisms that may buffer expression levels. Nevertheless, we found that the C/C allele at rs28834970 caused a large transcriptional response and resulting phenotypic changes, including reduction in chemokine release and decreased chemotaxis towards C5a. Downregulation of genes involved in cellular response to IFNγ and the lack of inhibition of migration following IFNγ treatment, are phenotypes consistent with the downregulation of PTK2B in cells harbouring the C/C allele. Following IFNγ receptor stimulation, PTK2B has been shown to bind to JAK2 and contribute to the activation of STAT1 and MAPK pathways[80]. Moreover, inhibition of PTK2B has been shown to decrease GSK-3β-mediated IFNγ signalling[80]. We propose that these phenotypic changes are the result of a change in microglia reactivity, causing them to respond differently to stimulations such as IFNγ, and this may impact the ability of the microglia to respond to aggregated proteins such as Aβ plaques and cellular damage in the brain of Alzheimer's disease patients. Indeed several pro-inflammatory chemokines, such as CCL2, CCL5 and CXCL1, which are downregulated in microglia with the C/C allele both in naive state and when stimulate with IFNγ, are produced by microglia in response to Aβ and are important for Aβ-induced microglia chemotaxis[81–83]. We believe this study demonstrates the power of isogenic cell lines to identify and validate individual variants, including common regulatory SNPs identified from GWAS.

References

1.   Gatz, M. *et al.* Role of genes and environments for explaining Alzheimer disease. *Arch. Gen. Psychiatry* **63**, (2006).

2.   Knopman, D. S. *et al.* Alzheimer disease. *Nature Reviews Disease Primers* **7**, 1–21 (2021).

3.   The complex genetic architecture of Alzheimer's disease: novel insights and future directions. *eBioMedicine* **90**, 104511 (2023).

4.   Bellenguez, C. *et al.* New insights into the genetic etiology of Alzheimer's disease and related dementias. *Nat. Genet.* **54**, (2022).

5.   Kunkle, B. W. *et al.* Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Aβ, tau, immunity and lipid processing. *Nat. Genet.* **51**, 414–430 (2019).

6.   Schwartzentruber, J. *et al.* Genome-wide meta-analysis, fine-mapping and integrative prioritization implicate new Alzheimer's disease risk genes. *Nat. Genet.* **53**, (2021).

7.   Ghoussaini, M. *et al.* Open Targets Genetics: systematic identification of trait-associated
genes using large-scale genetics and functional genomics. *Nucleic Acids Res.* **49**, D1311–D1320
(2021).

8.   Klein, J. C. *et al.* A systematic evaluation of the design and context dependencies of massively
parallel reporter assays. *Nat. Methods* **17**, 1083–1091 (2020).

9.   Cooper, S. E., Schwartzentruber, J., Bello, E., Coomber, E. L. & Bassett, A. R. Screening for
functional transcriptional and splicing regulatory variants with GenIE. *Nucleic Acids Res.* **48**, e131
(2020).

10.   Cooper, S., Schwartzentruber, J., Coomber, E. L., Wu, Q. & Bassett, A. Screening for functional
regulatory variants in open chromatin using GenIE-ATAC. *Nucleic Acids Res.* (2023)
doi:10.1093/nar/gkad332.

11.   Kwart, D. *et al.* A Large Panel of Isogenic APP and PSEN1 Mutant Human iPSC Neurons Reveals
Shared Endosomal Abnormalities Mediated by APP β-CTFs, Not Aβ. *Neuron* **104**, (2019).

12.   Penney, J., Ralvenius, W. T. & Tsai, L.-H. Modeling Alzheimer's disease with iPSC-derived brain
cells. *Mol. Psychiatry* **25**, 148–167 (2019).

13.   Moore, S. *et al.* APP Metabolism Regulates Tau Proteostasis in Human Cerebral Cortex
Neurons. *Cell Rep.* **11**, 689–696 (2015).

14.   APOE4 Causes Widespread Molecular and Cellular Alterations Associated with Alzheimer's
Disease Phenotypes in Human iPSC-Derived Brain Cell Types. *Neuron* **98**, 1141–1154.e7 (2018).

15.   Hall-Roberts, H. *et al.* TREM2 Alzheimer's variant R47H causes similar transcriptional
dysregulation to knockout, yet only subtle functional phenotypes in human iPSC-derived
macrophages. *Alzheimers. Res. Ther.* **12**, 151 (2020).

16.   Soldner, F. *et al.* Parkinson-associated risk variant in distal enhancer of α-synuclein modulates
target gene expression. *Nature* **533**, 95–99 (2016).

17.   Volpato, V. *et al.* Reproducibility of Molecular Phenotypes after Long-Term Differentiation to
Human iPSC-Derived Neurons: A Multi-Site Omics Study. *Stem Cell Reports* **11**, 897–911 (2018).

18.   Labzin, L. I., Heneka, M. T. & Latz, E. Innate Immunity and Neurodegeneration. *Annu. Rev.*
Med. **69**, (2018).

19.   Wang, H. Microglia Heterogeneity in Alzheimer's Disease: Insights From Single-Cell
Technologies. *Front. Synaptic Neurosci.* **13**, 773590 (2021).

20.   Li, Q. & Barres, B. A. Microglia and macrophages in brain homeostasis and disease. *Nat. Rev.*
Immunol. **18**, 225–242 (2017).

21.   Li, Y. *et al.* Genomics of Alzheimer's disease implicates the innate and adaptive immune
systems. *Cell. Mol. Life Sci.* **78**, 7397–7426 (2021).

22.   Nott, A. & Holtman, I. R. Genetic insights into immune mechanisms of Alzheimer's and
Parkinson's disease. *Front. Immunol.* **14**, 1168539 (2023).

23.   Tansey, K. E., Cameron, D. & Hill, M. J. Genetic risk for Alzheimer's disease is concentrated in
specific macrophage and microglial transcriptional networks. *Genome Med.* **10**, (2018).

24.   Lopes, K. P. *et al.* Genetic analysis of the human microglial transcriptome across brain regions,
aging and disease pathologies. *Nat. Genet.* **54**, (2022).

25.   Young, A. M. H. *et al.* A map of transcriptional heterogeneity and regulatory variation in
human microglia. *Nat. Genet.* **53**, (2021).

26.   Ulland, T. K. & Colonna, M. TREM2 — a key player in microglial biology and Alzheimer disease.
Nat. Rev. Neurol. **14**, 667–675 (2018).

27.   Zhao, L. CD33 in Alzheimer's Disease – Biology, Pathogenesis, and Therapeutics: A Mini-

Review. *Gerontology* **65**, 323–331 (2018).

28. Novoa, C. *et al.* Inflammation context in Alzheimer's disease, a relationship intricate to define. *Biol. Res.* **55**, 1–18 (2022).

29. Vogels, T., Murgoci, A.-N. & Hromádka, T. Intersection of pathological tau and microglia at the synapse. *Acta Neuropathologica Communications* **7**, 1–25 (2019).

30. Rogers, J., Strohmeyer, R., Kovelowski, C. J. & Li, R. Microglia and inflammatory mechanisms in the clearance of amyloid β peptide. *Glia* **40**, 260–269 (2002).

31. Hansen, D. V., Hanson, J. E. & Sheng, M. Microglia in Alzheimer's disease. *J. Cell Biol.* **217**, (2018).

32. Asai, H. *et al.* Depletion of microglia and inhibition of exosome synthesis halt tau propagation. *Nat. Neurosci.* **18**, (2015).

33. Hong, S. *et al.* Complement and microglia mediate early synapse loss in Alzheimer mouse models. *Science* **352**, (2016).

34. Miao, J. *et al.* Microglia in Alzheimer's disease: pathogenesis, mechanisms, and therapeutic potentials. *Front. Aging Neurosci.* **15**, 1201982 (2023).

35. Monoranu, C. M. *et al.* Is There Any Evidence of Monocytes Involvement in Alzheimer's Disease? A Pilot Study on Human Postmortem Brain. *Journal of Alzheimer's disease reports* **5**, (2021).

36. Malm, T., Koistinaho, M., Muona, A., Magga, J. & Koistinaho, J. The role and therapeutic potential of monocytic cells in Alzheimer's disease. *Glia* **58**, 889–900 (2010).

37. Gião, T., Teixeira, T., Almeida, M. R. & Cardoso, I. Choroid Plexus in Alzheimer's Disease-The Current State of Knowledge. *Biomedicines* **10**, (2022).

38. Park, L. *et al.* Brain Perivascular Macrophages Initiate the Neurovascular Dysfunction of Alzheimer Aβ Peptides. *Circ. Res.* **121**, (2017).

39. De Schepper, S. *et al.* Perivascular cells induce microglial phagocytic states and synaptic engulfment via SPP1 in mouse models of Alzheimer's disease. *Nat. Neurosci.* **26**, (2023).

40. Munawara, U. *et al.* Hyperactivation of monocytes and macrophages in MCI patients contributes to the progression of Alzheimer's disease. *Immun. Ageing* **18**, 1–25 (2021).

41. Zhu, X., Bao, Y., Guo, Y. & Yang, W. Proline-Rich Protein Tyrosine Kinase 2 in Inflammation and Cancer. *Cancers* **10**, (2018).

42. Beck, T. N., Nicolas, E., Kopp, M. C. & Golemis, E. A. Adaptors for disorders of the brain? The cancer signaling proteins NEDD9, CASS4, and PTK2B in Alzheimer's disease. *Oncoscience* **1**, 486 (2014).

43. de Pins, B., Mendes, T., Giralt, A. & Girault, J.-A. The Non-receptor Tyrosine Kinase Pyk2 in Brain Function and Neurological and Psychiatric Diseases. *Front. Synaptic Neurosci.* **13**, 749001 (2021).

44. Day, R. N., Day, K. H. & Pavalko, F. M. Direct visualization by FRET-FLIM of a putative mechanosome complex involving Src, Pyk2 and MBD2 in living MLO-Y4 cells. *PLoS One* **16**, (2021).

45. Bartos, J. A. *et al.* Postsynaptic clustering and activation of Pyk2 by PSD-95. *J. Neurosci.* **30**, (2010).

46. Siciliano, J. C., Toutant, M., Derkinderen, P., Sasaki, T. & Girault, J. A. Differential regulation of proline-rich tyrosine kinase 2/cell adhesion kinase beta (PYK2/CAKbeta) and pp125(FAK) by glutamate and depolarization in rat hippocampus. *J. Biol. Chem.* **271**, (1996).

47. Lev, S. *et al.* Protein tyrosine kinase PYK2 involved in Ca(2+)-induced regulation of ion channel and MAP kinase functions. *Nature* **376**, (1995).

48.   Girault, J. A., Costa, A., Derkinderen, P., Studler, J. M. & Toutant, M. FAK and PYK2/CAKbeta in
the nervous system: a link between neuronal activity, plasticity and survival? *Trends Neurosci.* **22**,
(1999).
49.   Ivankovic-Dikic, I., Grönroos, E., Blaukat, A., Barth, B. U. & Dikic, I. Pyk2 and FAK regulate
neurite outgrowth induced by growth factors and integrins. *Nat. Cell Biol.* **2**, (2000).
50.   Kong, L. *et al.* Integrin-associated molecules and signalling cross talking in osteoclast
cytoskeleton regulation. *J. Cell. Mol. Med.* **24**, (2020).
51.   Miao, L., Xin, X., Xin, H., Shen, X. & Zhu, Y.-Z. Hydrogen Sulfide Recruits Macrophage Migration
by Integrin β1-Src-FAK/Pyk2-Rac Pathway in Myocardial Infarction. *Sci. Rep.* **6**, 1–10 (2016).
52.   Okigaki, M. *et al.* Pyk2 regulates multiple signaling events crucial for macrophage morphology
and migration. *Proc. Natl. Acad. Sci. U. S. A.* **100**, (2003).
53.   Du, Q.-S. *et al.* Inhibition of PYK2-induced actin cytoskeleton reorganization, PYK2
autophosphorylation and focal adhesion targeting by FAK. *J. Cell Sci.* **114**, 2977–2987 (2001).
54.   Das, P. *et al.* A PKCβ-LYN-PYK2 Signaling Axis Is Critical for MCP-1-Dependent Migration and
Adhesion of Monocytes. *J. Immunol.* **206**, (2021).
55.   Yamamoto, S. *et al.* TRPM2-mediated Ca2+ influx induces chemokine production in
monocytes that aggravates inflammatory neutrophil infiltration. *Nat. Med.* **14**, 738 (2008).
56.   Alawieyah, S. M. S., Sim, J. A., Neubrand, V. E. & Jiang, L. H. A critical role of TRPM2 channel
in Aβ42 -induced microglial activation and generation of tumor necrosis factor-α. *Glia* **66**, (2018).
57.   Lambert, J.-C. *et al.* Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for
Alzheimer's disease. *Nat. Genet.* **45**, 1452–1458 (2013).
58.   Beecham, G. W. *et al.* Genome-wide association meta-analysis of neuropathologic features of
Alzheimer's disease and related dementias. *PLoS Genet.* **10**, (2014).
59.   Jiao, B. *et al.* Polygenic Analysis of Late-Onset Alzheimer's Disease from Mainland China. *PLoS*
One **10**, (2015).
60.   Li, Y. Q. *et al.* Common variant in PTK2B is associated with late-onset Alzheimer's disease: A
replication study and meta-analyses. *Neurosci. Lett.* **621**, (2016).
61.   Wang, X. *et al.* Genetic Determinants of Survival in Patientswith Alzheimer's Disease. *J.*
Alzheimers. Dis. **45**, (2015).
62.   Nettiksimmons, J., Tranah, G., Evans, D. S., Yokoyama, J. S. & Yaffe, K. Gene-based aggregate
SNP associations between candidate AD genes and cognitive decline. *Age* **38**, (2016).
63.   Lawingco, T. *et al.* Genetic variants in glutamate-, Aβ-, and tau-related pathways determine
polygenic risk for Alzheimer's disease. *Neurobiol. Aging* **101**, (2021).
64.   Rajabli, F. *et al.* Multi-ancestry genome-wide meta-analysis of 56,241 individuals identifies
LRRC4C, LHX5-AS1 and nominates ancestry-specific loci PTPRK , GRB14 , and KIAA0825 as novel risk
loci for Alzheimer's disease: the Alzheimer's Disease Genetics Consortium. *medRxiv : the preprint*
server for health sciences (2023) doi:10.1101/2023.07.06.23292311.
65.   Dourlen, P. *et al.* Functional screening of Alzheimer risk loci identifies PTK2B as an in vivo
modulator and early marker of Tau pathology. *Mol. Psychiatry* **22**, (2017).
66.   Köhler, C., Dinekov, M. & Götz, J. Active glycogen synthase kinase-3 and tau pathology-related
tyrosine phosphorylation in pR5 human tau transgenic mice. *Neurobiol. Aging* **34**, (2013).
67.   Salazar, S. V. *et al.* Alzheimer's Disease Risk Factor Pyk2 Mediates Amyloid-β-Induced Synaptic
Dysfunction and Loss. *J. Neurosci.* **39**, (2019).
68.   Brody, A. H. *et al.* Alzheimer risk gene product Pyk2 suppresses tau phosphorylation and
phenotypic effects of tauopathy. *Mol. Neurodegener.* **17**, 1–33 (2022).

69.   Alasoo, K. *et al.* Shared genetic effects on chromatin and gene expression indicate a role for
enhancer priming in immune response. *Nat. Genet.* **50**, 424–431 (2018).

70.   Panousis, N. I. *et al.* Gene expression QTL mapping in stimulated iPSC-derived macrophages
provides insights into common complex diseases. *bioRxiv* 2023.05.29.542425 (2023)
doi:10.1101/2023.05.29.542425.

71.   Tan, M.-S. *et al.* Associations of Alzheimer's disease risk variants with gene expression,
amyloidosis, tauopathy, and neurodegeneration. *Alzheimers. Res. Ther.* **13**, (2021).

72.   Bruntraeger, M., Byrne, M., Long, K. & Bassett, A. R. Editing the Genome of Human Induced
Pluripotent Stem Cells Using CRISPR/Cas9 Ribonucleoprotein Complexes. *Methods Mol. Biol.* **1961**,
153–183 (2019).

73.   van Wilgenburg, B., Browne, C., Vowles, J. & Cowley, S. A. Efficient, long term production of
monocyte-derived macrophages from human pluripotent stem cells under partly-defined and fully-
defined conditions. *PLoS One* **8**, e71098 (2013).

74.   Douthwaite, H., Arteagabeitia, A. B. & Mukhopadhyay, S. Differentiation of Human Induced
Pluripotent Stem Cell into Macrophages. *Bio Protoc* **12**, e4361 (2022).

75.   Brownjohn, P. W. *et al.* Functional Studies of Missense TREM2 Mutations in Human Stem Cell-
Derived Microglia. *Stem Cell Reports* **10**, 1294–1307 (2018).

76.   Washer, S. J. *et al.* Single-cell transcriptomics defines an improved, validated monoculture
protocol for differentiation of human iPSC to microglia. *Sci. Rep.* **12**, 19454 (2022).

77.   Nott, A. *et al.* Brain cell type-specific enhancer-promoter interactome maps and disease - risk
association. *Science* **366**, (2019).

78.   Bégay, V., Baumeier, C., Zimmermann, K., Heuser, A. & Leutz, A. The C/EBPβ LIP isoform
rescues loss of C/EBPβ function in the mouse. *Sci. Rep.* **8**, 1–10 (2018).

79.   Ryan, K. J. *et al.* A human microglia-like cellular model for assessing the effects of
neurodegenerative disease gene variants. *Sci. Transl. Med.* **9**, (2017).

80.   Takaoka, A. *et al.* Protein tyrosine kinase Pyk2 mediates the Jak-dependent activation of MAPK
and Stat1 in IFN-γ, but not IFN-α, signaling. *EMBO J.* **18**, 2480–2488 (1999).

81.   El Khoury, J. B. *et al.* CD36 Mediates the Innate Host Response to β-Amyloid. *J. Exp. Med.* **197**,
1657–1666 (2003).

82.   Lee, J. K., Schuchman, E. H., Jin, H. K. & Bae, J.-S. Soluble CCL5 Derived from Bone Marrow-
Derived Mesenchymal Stem Cells and Activated by Amyloid β Ameliorates Alzheimer's Disease in
Mice by Recruiting Bone Marrow-Induced Microglia Immune Responses. *Stem Cells* **30**, 1544–1555
(2012).

83.   Zhang, K. *et al.* CXCL1 Contributes to β-Amyloid-Induced Transendothelial Migration of
Monocytes in Alzheimer's Disease. *PLoS One* **8**, (2013).

84.   Slaymaker, I. M. *et al.* Rationally engineered Cas9 nucleases with improved specificity. *Science*
351, 84–88 (2016).

85.   Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler transform.
Bioinformatics **26**, (2010).

86.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res. **44**, W160–W165 (2016).

87.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2013).

88.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

89. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).
90. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2009).
91. Raudvere, U. *et al.* g:Profiler: a web server for functional enrichment analysis and conversions
of gene lists (2019 update). *Nucleic Acids Res.* **47**, W191–W198 (2019).
92. Buenrostro, J., Wu, B., Chang, H. & Greenleaf, W. ATAC-seq: A Method for Assaying Chromatin
Accessibility Genome-Wide. *Curr. Protoc. Mol. Biol.* **109**, 21.29.1 (2015).
93. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078
(2009).
94. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, 1–9 (2008).
95. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).
96. Yu, G., Wang, L. G. & He, Q. Y. ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. *Bioinformatics* **31**, (2015).
97. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357.
98. Meers, M. P., Tenenbaum, D. & Henikoff, S. Peak calling by Sparse Enrichment Analysis for
CUT&RUN chromatin profiling. *Epigenetics Chromatin* **12**, 1–11 (2019).
99. McQuin, C. *et al.* CellProfiler 3.0: Next-generation image processing for biology. *PLoS Biol.* **16**,
e2005970 (2018).
100. Coetzee, S. G., Coetzee, G. A. & Hazelett, D. J. motifbreakR: an R/Bioconductor package for
predicting variant effects at transcription factor binding sites. *Bioinformatics* **31**, 3847 (2015).
101. Haenseler, W. *et al.* A Highly Efficient Human Pluripotent Stem Cell Microglia Model Displays
a Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response. *Stem Cell Reports*
8, 1727–1742 (2017).
102. Vaughan-Jackson, A. *et al.* Differentiation of human induced pluripotent stem cells to
authentic macrophages using a defined, serum-free, open-source medium. *Stem Cell Reports* **16**,
3093 (2021).
103. Jurga, A. M., Paleczna, M. & Kuter, K. Z. Overview of General and Discriminating Markers of
Differential Microglia Phenotypes. *Front. Cell. Neurosci.* **14**, 544457 (2020).
104. Murray, P. J. *et al.* Macrophage activation and polarization: nomenclature and experimental
guidelines. *Immunity* **41**, (2014).

Methods

Human induced pluripotent stem cells (hiPSC) culture

KOLF2_C1 (https://hpscreg.eu/cell-line/WTSIi018-B-1) is a male hiPSC line generated as part of the HipSci project (Cambridgeshire 1 NRES REC Reference 09/H0304/77, HMDMC 14/013) that has been subcloned from a single cell to ensure homogeneity of the population. KOLF2_C1 cells were cultured in TeSR E8 medium (StemCell Technologies) on Synthemax (Corning) (final amount 2 µg/cm^2) and routinely passaged 1:10 every 5 days using Gentle Cell Dissociation Reagent (StemCell Technologies).

Engineering rs28834970 C/C allele in hiPSCs

The C/C risk allele at variant rs28834970 was engineered in hiPSCs by nucleofection of ribonucleoprotein (RNP) complex containing full-length chemically modified synthetic guide RNA and eSpCas9, along with a ssODN repair template, as previously described[72].

Briefly, eSpCas9[84] was expressed and purified from Escherichia coli using a His-tag. KOLF2_C1 cells were cultured ahead of nucleofection in TeSR E8 medium (StemCell Technologies) on Synthemax (Corning) (final amount 5 µg/cm^2). eSpCas9 (5µl, 20 µg) was mixed with full-length chemically modified guide RNA (AGTGGAATTGTACAACACTG) (Synthego) (5 µl, 225 pmol) at room temperature for 20 min for RNP complexes to form, followed by addition of the ssODN repair template (CTGGCAAGACTAATCTACTTTCTATTTTTATGGATCTGCCTTTTCTGGTCATTCCATATAAGTGGAATTGCA CAACACTGTGGCCTTTCGCGACGGCTGCTTTCACTTAGCACAATGTTTTGAAACTTCCTCCATGTTGT) (5 µl, 500 pmol) just before the nucleofection. Cells were detached using Accutase (StemCell Technologies) and dissociated into a single cell suspension. 1×10^6 cells were resuspended in P3 buffer (Lonza), mixed with the RNP/template complex and nucleofected using Lonza 4D-Nucleofector with program CA137. After nucleofection cells were plated onto a 10 cm dish coated with Synthemax (5 µg/cm^2) with TeSR-E8 supplemented with CloneR (StemCell Technologies). Cells were cultured to 80% confluency, detached using Accutase and dissociated into single cells for subcloning. 5000 cells were plated in a 10 cm dish and expanded, and colonies were picked for genotyping and freezing. Colonies were screened for introduction of the homozygous C allele by high throughput sequencing of an amplicon spanning the rs28834970 variant using an Illumina MiSeq instrument. Final cell lines were validated using Sanger sequencing. Clones used in downstream assays include: three homozygous (C/C) clones (named C8, C11 and D3), one unedited (T/T) picked clone (B11) and the parental KOLF2_C1 line.

hiPSC differentiation to macrophages

iPSCs were differentiated to macrophages as previously described[73,74]. Briefly, KOLF2_C1 cells were cultured in Essential E8 medium (Gibco) on Vitronectin (Gibco, 100x) and split onto a feeder layer of Mouse Embryonic Fibroblasts (MEFs) (Merck) using Gentle Cell Dissociation Reagent (StemCell Technologies). hiPSCs were cultured on the feeder layer in hiPSC base medium, containing Advanced DMEM F12 (Gibco), KnockOut Serum Replacement (Gibco), 2mM Glutamax (Gibco), 100 U/ml penicillin/streptomycin (Gibco), 0.055 mM 2-Mercaptoethanol (Sigma) and 4ng/ml of Recombinant Human FGF (R&D Systems), until the colonies were sufficiently large. HiPSC colonies were lifted using a solution of equal amounts of Collagenase type IV (ThermoFisher Scientific) and Dispase type II (ThermoFisher Scientific) for 15 minutes at 37°C. Detached colonies were plated in 10 cm low adherent dishes in hiPSC base medium without FGF to form Embryoid Bodies (EBs). After five days, EBs were harvested and plated on gelatin coated 10 cm dishes in EB medium, containing X-Vivo 15 medium (Lonza), 2 mM Glutamax (Gibco), 100 U/ml penicillin/streptomycin (Gibco), 0.055mM β-mercaptoethanol (Sigma), 50 ng/ml rhM-CSF (R&D Systems) and 25 ng/ml rhIL-3 (R&D Systems). Macrophage precursor cells could be routinely harvested from dishes containing EBs every 6/7 days for up to 3/4 months. Precursors were differentiated for 7 days by plating 15,000 cells/cm^2 in macrophage media, containing RPMI 1640 medium (Gibco), 10% FBS (Thermo Fisher

585 Scientific), 2 mM Glutamax (Gibco), 100 U/ml penicillin/streptomycin (Gibco) and 100 ng/ml rhM-
586 CSF (R&D Systems).
587
HiPSCs differentiation to microglia
HiPSCs were differentiated to microglia as previously described[75]. Briefly, KOLF2_C1 cells were
cultured in Essential E8 medium (Gibco) on Synthemax (Corning) (final amount 5 µg/cm^2). Two days
after passaging, cells were detached using Accutase (StemCell Technologies) and dissociated into
a single cell suspension. 10,000 hiPSCs were plated into each well of a 96-well plate in EB medium,
containing Essential E8 medium (Gibco), 10 uM Y-27632 (Dihydrochloride)(StemCell technologies),
50 ng/ml rhBMP-4 (Perprotech), 20 ng/ml rhSCF (Perprotech) and 50 ng/ml rhVEGF (Perprotech).
The 96-well plate was span down at 300xg for 3 minutes and incubated for 4/5 days for EB
formation. EBs were then transferred to tissue culture dishes coated in 0.1% gelatin (Sigma) and
cultured in Microglia Precursors media, containing X-Vivo 15 medium (Lonza), 2 mM Glutamax
(Gibco), 100 U/ml penicillin/streptomycin (Gibco), 0.055 mM β-mercaptoethanol (Sigma), 100
ng/ml rhM-CSF (Perprotech) and 25 ng/ml rhIL-3 (Cell Guidance Systems). Microglia precursor cells
could be routinely harvested from dishes containing EBs every 6/7 days for up to 3 months.
Precursors were differentiated for 7 days by plating at a density of 180,000 cells/cm^2 in microglia
media, containing RPMI 1640 medium (Gibco), 10% FBS (Thermo Fisher Scientific), 2 mM Glutamax
(Gibco), 100 U/ml penicillin/streptomycin (Gibco), 10 ng/ml rhGM-CSF (Perprotech) and 100 ng/ml
rhIL-34 (Perprotech).
605
Fluorescence activated cell sorting analysis
Microglia or macrophages were detached using Accutase (StemCell Technologies) and washed
twice in PBS. 1x10^5 cells were incubated with Human TruStain FcX (Fc Receptor Blocking Solution)
(Biolegend) for 30 minutes at 4°C in the dark. Microglia were stained with 5 ul each of PE-anti
human CD45 antibody clone HI30 (Biolegend) and APC-anti human CD11b antibody clone ICRF44
(Biolegend) for 30 minutes at 4°C. Macrophages were instead stained with 5 ul of either APC/Cy7-
anti human CD16 antibody clone 3G8 (Biolegend), APC-anti human CD206 antibody (BD
Biosciences) or PE-anti human CD14 antibody (BD Biosciences). Isotype controls were stained with
either 0.2 mg/ml APC-mouse IgG1 k(BD Biosciences), 0.2 mg/ml of PE-mouse IgG2A (BD
Biosciences) or 0.2 mg/ml APC/Cy7 mouse IgG1 k (BD Biosciences). Cells were then washed three
times in FACS buffer (5% FBS in D-PBS) and resuspended in 200 μl of DAPI solution (1 ug/ml in PBS)
and analysed. All samples were analysed using a BD LSRll instrument. An unpaired t-test was used
to test the difference between samples from cells with the T/T and the C/C allele at rs28834970.
619
RNA extraction
TRI Reagent (Zymo) was added direct on cell layers previously washed in PBS and mixed. RNA was
extracted using the Direct-zol RNA Miniprep kit (Zymo) following the manufacturer's protocol. The
optional in-column DNase digest was performed, and RNA was eluted in 50 µl DNAse/RNAse-free
water. We made cDNA from 1 µg RNA using Superscript IV (Thermo-Fisher) according to the
manufacturer's protocol.
626
627 **Reverse transcription**
628 1 µg RNA was reverse transcribed using Superscript IV (Thermo-Fisher) and random hexamers,
629 according to the manufacturer's protocol.
630
RNA sequencing of hiPSC-derived macrophages and microglia
Transcriptome libraries for hiPSC-derived macrophages and microglia were generated with the
Illumina TruSeq stranded RNAseq kit (polyA) and all samples were sequenced using Illumina HiSeq
with 50 million mapped reads on average for each sample.

Reads were mapped to the human genome (GRCh38/hg38) using the BWA software[85] and coverage
tracks were visualised on the UCSC genome browser using bigwig files, made with bamCoverage[86]
and normalised by Counts per Million (CPM). The number of reads per transcript were counted
using the featureCounts software[87] and a custom R script was used to calculate Transcripts per
Million (TPM). For RNAseq analysis of macrophages, TPMs were corrected for batch effects using
limma function limma::removeBatchEffect[88]. For microglia, all samples were differentiated and
sequenced in the same batch. Raw read counts per transcript were used to analyse differential
gene expression between cells with the T/T and the C/C allele at rs28834970, using the package
DESeq2[89]. DESeq2 was also used to generate PCA plots of variance stabilizing transformation (vst)
transformed counts. To investigate whether the rs28834970 variant had an effect on PTK2B
splicing, we used EdgeR[90] to calculate differential exon usage between microglia harbouring the
C/C and the T/T allele. Gene ontology (GO) analysis of differential expressed genes was carried out
using g:Profiler package in R[91].

ATACseq of hiPSC-derived macrophages and microglia
hiPSC-derived macrophages and microglia were detached using a solution of 4 mg/ml Lidocaine
hydrochloride monohydrate (Sigma) and 5mM EDTA (ThermoFisher scientific) for 15 minutes at
37°C. ATACseq was performed as previously described[92]. Briefly cells were lysed using sucrose
buffer containing 10 mM Tris-Cl pH 7.5, 3 mM $CaCl_2$, 2 mM $MgCl_2$, 0.32 M sucrose and
permeabilised using 0.5% Triton-X-100 (Sigma). Nuclei were recovered by spinning at 450xg for 5
minutes at 4°C and tagmentation was performed using Illumina Tagment DNA TDE1 Enzyme and
Buffer Kits for 30 minutes at 37°C37C. 5 μl of TDE1 enzyme was used per 100,000 cells. MinElute
PCR Purification Kit buffer PB (Qiagen) was added to stop the reaction and tagmented DNA was
purified and eluted in 10 μl of EB. The whole volume was PCR amplified and dual indexed Illumina
adapters were added (primer sequences from[92]). Amplified libraries were gel purified and
sequenced using an Illumina Novaseq 6000 with 30 million reads on average per sample.
Reads were mapped to the human genome (GRCh38/hg38) using the BWA software[85] and
duplicate reads were removed using picard tools MarkDuplicates. Coverage tracks were visualised
on the UCSC genome browser using bigwig files, made with bamCoverage[86] and normalised by
Counts per Million (CPM). The number of mapped reads per sample was determined and all files
were subsampled to the number of reads of the lowest sample, using samtools view[93]. ATACseq
peaks were called using MACS2[94] with flags --shift -100 --extsize 200 for window extension. A
merged file of all called peaks in all samples was created and the number of reads under each peak
was counted in all samples using bedtools coverageBed function[95]. Genome-wide analysis of
differential chromatin accessibility between cells harbouring the C/C and the T/T allele at
rs28834970 was performed using DESeq2[89]. DESeq2 was also used to generate PCA plots of
variance stabilizing transformation (vst) transformed counts. Peaks were annotated to genes using
CHIPseeker functiomannotatePeak[96].

CUT & RUN assay
Microglia were harvested using a solution of 4mg/ml Lidocaine hydrochloride monohydrate
(Sigma) and 5mM EDTA (ThermoFisher Scientific) for 15 minutes at 37°C37C. CUT&RUN for CEBPβ
was performed using CUTANA CUT&RUN Kit (EpiCypher) according to the manufacturer's
instructions. 400,000 microglia were used per sample and permeabilization was optimised using
Digitonin (EpiCypher) 0.01%. 1ug of C/EBP beta Antibody (H-7) sc-7962 (Santa Cruz Biotechnology)
was used per sample and one control sample was treated in parallel with 1 μl of CUTANA Rabbit
IgG CUT&RUN Negative Control Antibody. 0.5ng of E. coli Spike-in DNA (EpiCypher) was added to
each sample to normalise reads to control for experimental variability and sequencing depth.
Around 5 ng of purified CUT&RUN-enriched DNA was used to prepare NGS libraries using the
NEBNext Ultra II DNA Library Prep Kit (NEB) with NEBNext Multiplex Oligos for Illumina (Dual Index

Primers Set 1) (NEB). All libraries were sequenced 150bp paired end using an Illumina Novaseq
6000 with 15 million reads on average per sample.
Reads were trimmed to exclude Illumina adapter sequences using trim_galore and mapped to the
human genome (GRCh38/hg38) using the bowtie2 software[97]. The number of human mapped
reads per sample was determined and all files were subsampled to the number of reads of the
lowest sample, using samtools view[93]. Trimmed reads were also mapped to the E. coli genome
using bowtie2 and reads mapped to the human genome were normalised to spiked-in E.coli DNA
using a previously published script (https://github.com/Henikoff/Cut-and-
[Run/blob/master/spike_in_calibration.csh](https://github.com/Henikoff/Cut-and-Run/blob/master/spike_in_calibration.csh)) integrated into a custom bash script
(https://github.com/ericabello/PTK2B_rs28834970/blob/main/Cut_Run_CEBPb_micro/cmds_Cut
[Run_CEBPb_micro.sh](https://github.com/ericabello/PTK2B_rs28834970/blob/main/Cut_Run_CEBPb_micro/cmds_Cut_Run_CEBPb_micro.sh)). Coverage tracks were visualised on the UCSC genome browser using bigwig
files made from E.coli normalised bedgraph files of fragments 1-1000bp. Bigwigs were made using
UCSC bedGraphToBigWig. Peaks in all samples containing the CEBPβ antibody were called using
the SEACR software[98] and the IgG negative control sample was used as input to identify the
threshold value at which the percentage of target versus IgG signal is maximized. A merged file of
all CEBPβ peaks in all samples was created and the number of reads under each peak was counted
in all samples using bedtools coverageBed function[95]. Genome-wide analysis of differential CEBPβ
binding between microglia harbouring the C/C and the T/T allele at rs28834970 was performed
using DESeq2[89].
704
705 **Digital droplet qPCR**
706 50 ng of cDNA per sample was mixed with 10 µl of ddPCR Supermix for probes (no dUTP) (Biorad),
707 1 µl of FAM-tagged TaqMan assay for detection of the gene of interest and 1 µl of VIC-tagged
708 Taqman assay for a housekeeping gene control. The housekeeping control gene was chosen to
709 match the level of expression of the gene of interest in each cell type.
710
711 Taqman assays list (ThermoFisher Scientific), containing probes and primers:
712 *Macrophages*

gene	FAM probe ID	control gene	VIC control probe ID
CCL1	Hs00171072_m1	TBP	Hs00427620_m1
CCL8	Hs00271615_m1	SDHA	Hs00188166_m1
CCL13	Hs00234646_m1	GAPDH	Hs99999905_m1
CCL5	Hs99999048_m1	TBP	Hs00427620_m1
CCL2	Hs00234140_m1	UBC	Hs00824723_m1
TRIM35	Hs01093979_m1	TBP	Hs00427620_m1
CLU	Hs00156548_m1	TBP	Hs00427620_m1
SCARA3	Hs00939871_m1	TBP	Hs00427620_m1
PTK2B	Hs01559703_m1	SDHA	Hs00188166_m1
HLA-DQA1	Hs03007426_mH	TBP	Hs00427620_m1
HLA-F	Hs01587840_m1	TBP	Hs00427620_m1

714 *Microglia*

gene	FAM probe ID	control gene	VIC control probe ID
CCL1	Hs00171072_m1	HPRT1	Hs02800695_m1
CCL22	Hs00171080_m1	TBP	Hs00427620_m1
CCL8	Hs00271615_m1	TBP	Hs00427620_m1
CXCL6	Hs00237017_m1	TBP	Hs00427620_m1
CCL13	Hs00234646_m1	UBC	Hs00824723_m1
CXCL2	Hs00601975_m1	TBP	Hs00427620_m1
IL18	Hs01038788_m1	TBP	Hs00427620_m1
CXCL3	Hs00171061_m1	TBP	Hs00427620_m1
CXCL1	Hs00236937_m1	SDHA	Hs00188166_m1
CXCL5	Hs00171085_m1	SDHA	Hs00188166_m1
CCL5	Hs99999048_m1	TBP	Hs00427620_m1
CCL2	Hs00234140_m1	GAPDH	Hs99999905_m1
TRIM35	Hs01093979_m1	TBP	Hs00427620_m1
CLU	Hs00156548_m1	TBP	Hs00427620_m1
SCARA3	Hs00939871_m1	TBP	Hs00427620_m1
PTK2B	Hs00169444_m1	TBP	Hs00427620_m1
HLA-DQA1	Hs03007426_mH	TBP	Hs00427620_m1
HLA-F	Hs01587840_m1	TBP	Hs00427620_m1

Droplets were generated for each reaction using a QX200 Droplet Generator (Biorad) and
transferred to 96 well plates. Sealed plates were incubated in a thermocycler and cDNA was PCR
amplified with the following steps: 95°C for 10 min then 95°C for 30 sec and 60°C for 1 min for 40
cycles and finally 98°C for 10 min. The resulting FAM and VIC fluorescence was read using a QX200
Droplet Reader (Biorad) and the fraction of positive droplets in the sample was determined.
Poisson statistics were used to determine the absolute number of copies per μl in each sample for
both target and housekeeping control genes. The number of copies of target gene per sample was
normalised by the control gene for each reaction and an unpaired t-test was used to test the
difference between samples from cells with the T/T and the C/C allele at rs28834970.

725
726

Microglia stimulation and Luminex assay

hiPSC-derived microglia at day 6 of differentiation were stimulated with IFNγ (R&D Systems) 200 ng/ml or Lipopolysaccharide (LPS) (Invitrogen) 100 ng/ml for 24 hours. Culture plates were spun at 4,000 rpm for 20 min at 4°C and media collected. Harvested media was diluted 1:20 for unstimulated controls and 1:250 for stimulated samples. Luminex assay was performed using ProcartaPlex Human Basic Kit (Invitrogen) and a panel of ProcartaPlex Human Simplex (Invitrogen) beads, according to the manufacturer's recommendations. Plates were read using a Luminex MAGPIX instrument. Media from cells with the T/T and C/C allele were ran on the same plate, allowing direct comparison of each protein concentration across genotypes. An unpaired t-test was used to test the difference in concentration (pg/ml) between different samples. The propagation of error for log fold change values of chemokine concentration between unstimulated and stimulated microglia was calculated using the Python package uncertainties (https://pythonhosted.org/uncertainties/).

Microglia stimulation and migration assay

HiPSCs-derived microglia at day 7 of differentiation were stimulated with IFNγ (R&D Systems) 100 ng/ml for 72 hours. At day 10, cells were detached using a solution of 4 mg/ml Lidocaine hydrochloride monohydrate (Sigma) and 5 mM EDTA (Gibco) and 5.5×10^4 cells in 100 μl macrophage media were seeded onto each transwell (PET with 5 µm pores, Sarstedt) in a 24-well plate. Cells were incubated for 15 minutes to settle and 600 μl of macrophage media containing 3 nM human recombinant C5a was added beneath the transwells. The plate was incubated for 10 hours to allow cell migration. The transwell was then rinsed in PBS, transferred to a fresh 24-well plate, and fixed with 4% paraformaldehyde for 20 min at RT. Cells on either side of the transwell membrane were stained with NucBlue (Invitrogen) and imaged with an EVOS FL Auto automated microscope (Thermo Fisher) set up to take images of each transwell in full (top image). The transwells were swabbed with a cotton wool bud to remove cells on the top surface, leaving behind only migrated cells, transferred into a fresh plate and imaged again with the same settings (bottom image). Cell counting was performed with CellProfiler 3.0 software[99] and the percentage of migrated cells per transwell was calculated as: (no. cells in bottom image) ÷ (no. cells in top image) × 100. Stimulations were performed in triplicate and the average for each sample across wells was taken. An unpaired t-test was used to test the difference between different samples.

Data availability

Raw sequencing data is available at ENA https://www.ebi.ac.uk/ena/browser/search with accession number ERP024123 for ATACseq and Cut&Run data and ERP024128 for RNAseq data. A description of the sample IDs can be found in Supplementary Table 4.

All analysis scripts are available at https://github.com/ericabello/PTK2B_rs28834970.

Acknowledgements

This work was funded by OpenTargets [OTAR037] and Wellcome [206194]. We thank the Sanger Institute flow cytometry core facility for flow cytometry analysis. Addgene plasmids, etc. We acknowledge Scientific Operations at the Sanger Institute for support in next generation sequencing and quality control. We thank members of the Bassett lab for helpful discussions and support.

Author contributions

A.B. and E.B. conceived the work and designed experiments, K.L. made the hiPSC isogenic cell lines, E.B., S.I., J.S. executed the experiments, E.B., K.A., N.K., J.S., N.I.P. analysed data, E.B. and A.B. wrote the manuscript with input from all other authors.

Competing interests

A.B. is a founder of EnsoCell therapeutics. N.I.P. was an employee of GSK at the time the manuscript was submitted. J.S. was an employee of Illumina at the time the manuscript was submitted.

Materials and Correspondance

All correspondence and requests for materials should be addressed to Andrew Bassett (ab42@sanger.ac.uk) or Erica Bello (eb956@cam.ac.uk).

Figures and legends

Figure 1

791 **Figure 1 - The rs28834970 variant in PTK2B is predicted to modulate an intronic enhancer**
792 A) *PTK2B* gene structure and Manhattan plots for Alzheimer's Disease (AD) GWAS hits (top
793 panel), eQTL in naïve iPSC-derived macrophages (second panel) and caQTL in naïve iPSC-
794 derived macrophages (bottom panel). P-values replotted from[69].

B) Boxplots showing the expression of the *PTK2B* gene stratified by the rs28834970
genotype in iPSC-derived macrophages. Data replotted from[25]. The y-axis shows
normalised expression levels (log TPM value) and each dot shows the expression level of a
single sample.

C) Boxplots showing the expression of the *PTK2B* gene stratified by the rs28834970
genotype in naïve iPSC-derived macrophages. Data replotted from[70]. The y-axis shows
normalised expression levels (log TPM value) and each dot shows the expression level of a
single sample.

D) ATAC-seq fragment coverage in iPSC-derived macrophages stratified by the rs28834970
genotype (top panel) and CEBPβ ChIP-seq fragment coverage and peaks in primary human
macrophages (middle panel). Data replotted from[69]. The bottom panel shows the structure
of the *PTK2B* gene.

E) Overview of the experimental design. Created with BioRender.com.

Figure 2

808
Figure 2 - Isogenic edited cell lines show that rs28834970 effects *PTK2B* expression
through effects on chromatin accessibility and CEBPβ binding at the intronic enhancer in
iPSC-derived macrophages and microglia.

A) Boxplots showing *PTK2B* expression in macrophages harbouring the T/T or C/C allele at
rs28834970 measured by RNA-seq (top) and droplet digital q-PCR (ddqPCR) (bottom). Top
plot: TPM= transcripts per million. Bottom plot: Relative concentration = copies/ul relative
to housekeeping gene (SDHA), n=3 and **p<0.01, one-way paired ANOVA.
B) Boxplots showing *PTK2B* expression in microglia harbouring the T/T or C/C allele at
rs28834970 measured by RNA-seq (top) and droplet digital q-PCR (ddqPCR) (bottom). Top
plot: TPM= transcripts per million. Bottom plot: Relative concentration = copies/ul relative
to housekeeping gene (TBP), n=3 and **p<0.01, one-way paired ANOVA. ns= non-
significant
C) Coverage tracks of cells with T/T (orange) and C/C (blue) allele at rs28834970. The tracks
are: ATAC-seq in macrophages (upper track), ATAC-seq in microglia (second track),
CUT&RUN for CEBPβ in microglia (third track) and IgG negative control for CUT&RUN
(bottom track). The rs28834970 variant in *PTK2B* intron is highlighted in yellow.
D) Effect of rs28834970 SNP on transcription factor binding probability. Position
probability matrices are shown for CEBPβ (top) and CEBPα (middle) as well as the position
of the binding motifs on the *PTK2B* sequence (bottom). The plot was generated using
motifbreakerR[100].
E) Volcano plot showing fold changes of CEBPβ binding between microglia harbouring the
C/C and the T/T allele for all detectable peaks, measured by CUT&RUN. Blue dots indicate
peaks with no significant change while red dots represent significantly increased peaks
(logFC>1 and p<0.05). The only significantly increased CEBPβ peak found spans the
rs28834970 variant in the *PTK2B* intron.
834

Figure 3

835
Figure 3 - The rs28834970 variant affects global gene expression in iPSC-derived
macrophages and microglia

A) Results of differential expression analysis of macrophages (upper panels) and microglia
(lower panels) harbouring the C/C allele versus the corresponding iPSC-derived cells
harbouring the T/T allele at rs28834970. Left panels show Principal component analysis
(PCA) of transcriptomes of cells differentiated from two hiPSC clones with the T/T allele
and three clones with the C/C allele. Right panels are a volcano plot showing the log2 fold-
change (FC) between cells with the C/C and the T/T allele and corresponding p values for
all transcribed genes. Significantly upregulated genes (logFC>0.5 and p<0.05) are shown as
red dots, significantly downregulated genes (logFC<-0.5 and p<0.05) are shown as blue
dots and non-significant genes as black dots.
B) Heatmaps showing relative expression (z-scores) of DE genes found in gene ontology
(GO) analysis in macrophages with the T/T (orange) and the C/C (purple) allele at
rs28834970. GO pathways identified are also annotated.
C) Heatmaps showing relative expression (z-scores) of DE genes found in gene ontology
(GO) analysis in microglia with the T/T (orange) and the C/C (purple) allele at rs28834970.
GO pathways identified are also annotated.
D) Dot plot showing logFC of downregulated genes belonging to the top eight GO pathways
in macrophages and microglia.
E) Bar graph showing logFC of downregulated chemokines and HLA genes in macrophages
(left) and microglia (right) measured by RNAseq and ddqPCR. ddqPCR: n=3 *p<0.05,
**p<0.01, ***p<0.001, ****p<0.0001 by one-way paired ANOVA.
858

Figure 4 - The rs28834970 variant causes changes in chemokine production and migration

in iPSC-derived microglia

A) Concentration of downregulated chemokines in unstimulated microglia with the C/C

allele at rs28834970 relative to the concentration in cells with the T/T allele, measured by

Luminex assay. Results are shown as the mean ± SEM of independent biological replicates

(n=3). ns= non-significant, * p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, unpaired t-test.

B) Concentration of downregulated chemokines in microglia with the C/C allele at

rs28834970 relative to the concentration in cells with the T/T allele, stimulated with IFNγ.

Results are shown as the mean ± SEM of independent biological replicates (n=3). ns= non-

significant, * p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, unpaired t-test.

C) Bar graph showing the percentage of migrated microglia harbouring the T/T allele (left)

or the C/C allele (right) either unstimulated or stimulated with IFNγ, and with or without

the chemoattractant C5a. Results are shown as the mean ± SEM of independent biological

replicates (n=3). ns= non-significant, * p<0.05, **p<0.01, ***p<0.001, unpaired t-test.

874