**Zinc stimulation of phytoplankton in a low carbon dioxide, coastal Antarctic environment:**

**evidence for the Zn hypothesis**

Riss M. Kell[1,+], Adam V. Subhas[1], Nicole L. Schanke[2], Lauren E. Lees[3], Rebecca J. Chmiel[1],

Deepa Rao[1], Margaret M. Brisbin[1], Dawn M. Moran[1], Matthew R. McIlvin[1], Francesco

Bolinesi[4], Olga Mangoni[4], Raffaella Casotti[5], Cecilia Balestra[6], Tristan J. Horner[1], Robert B.

Dunbar[7], Andrew E. Allen[8,9], Giacomo R. DiTullio[2], Mak A. Saito[1*]

[1]Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution,

Woods Hole, MA, 02543, USA

[2]Hollings Marine Laboratory, College of Charleston, Charleston, SC, 29424, USA

[3]Department of Ecology and Evolutionary Biology, University of California at Irvine, Irvine,

CA, 92697, USA

[4]Department of Biology, Università degli Studi di Napoli Federico II, Complesso di Monte

Sant'Angelo, Via Cinthia 21, 80126, Napoli, Italy

[5]Department of Integrative Marine Ecology, Stazione Zoologica Anton Dohrn, Villa Comunale,

80121, Naples, Italy

[6]National Institute of Oceanography and Applied Geophysics, 34010 Sgonico (TS), Italy

[7]Doerr School of Sustainability, Stanford University, Stanford, CA, 94305, USA

[8]Microbial & Environmental Genomics, J. Craig Venter Institute, San Diego, CA, 92093, USA

[9]Scripps Institution of Oceanography, Integrative Oceanography Division, University of

California, San Diego, CA, 92037, USA

*Correspondence to:* Mak A. Saito (msaito@whoi.edu)

[+]Formerly published under Riss Kellogg; now affiliated with Gloucester Marine Genomics

Institute, Gloucester MA, 01930

**Abstract.**

The ocean acts as a carbon sink, absorbing carbon from the atmosphere and resulting in

substantial uptake of anthropogenic $CO_2$ emissions. As biological processes in the oceans such as

net primary production (NPP) contribute significantly to this sink, understanding how they will

shift in response to increasing atmospheric $CO_2$ is necessary to project future ocean carbon

storage capacity. Macronutrient and micronutrient resource limitation within the oceans regulates

NPP, and while some micronutrients such as zinc (Zn) are present at very low concentrations,

their ability to limit NPP has remained unclear. Zn is a key micronutrient used by phytoplankton

for a multitude of metabolic functions, yet there have been few observations of its influence on

natural oceanic phytoplankton populations. In this study, we observed Zn limitation of growth in

the natural phytoplankton community of Terra Nova Bay, Antarctica, in addition to primary iron

(Fe) limitation. Shipboard incubation experiments amended with Zn and Fe resulted in

significantly higher chlorophyll *a* content and dissolved inorganic carbon drawdown compared

to Fe addition alone. Zn and Fe stress response proteins detected in incubation and

environmental biomass provided independent verification of algal co-stress for these

micronutrients. We consider total biomass and low surface ocean $pCO_2$ as potential drivers of

environmental Zn stress. This study definitively establishes that Zn limitation can occur in the

modern oceans, opening up new possibility space in our understanding of nutrient regulation of

NPP through geologic time, and we consider the future of oceanic Zn limitation in the face of

climate change.

**1 Introduction**

Primary productivity in the oceans is a key component of the global carbon cycle and is

largely controlled by the availability of nitrogen (N), phosphorus (P), and iron (Fe). Yet there is

increasing evidence that other micronutrients such as zinc (Zn), cobalt (Co), and vitamin $B_{12}$ can

also influence phytoplankton productivity, often as secondary limiting nutrients after N, P, or Fe

are added (Moore et al. 2013). Zn can be particularly scarce in the photic zone (Bruland 1980;

Jakuba et al. 2012) where total dissolved Zn ($dZn_T$) can be below 0.2 nM in seawater due to

biological uptake and complexation by organic ligands (Bruland 1989; Lohan et al. 2002; Baars

and Croot 2011; Middag et al. 2019), which further lowers Zn bioavailability (Sunda and

Huntsman 2000; Saito et al. 2008; Lhospice et al. 2017). Marine eukaryotic algae and

copiotrophic bacteria possess a large metabolic demand for Zn that is on par with that of Fe

(Sunda and Huntsman 2000; Mazzotta et al. 2021) and polar phytoplankton in particular

demonstrate an enhanced Zn demand based on genomic and laboratory analyses (Ye et al. 2022).

Despite the scarcity of bioavailable Zn in the surface ocean and its high cellular demand,

relatively few experimental studies have examined the ability of Zn addition to stimulate natural

phytoplankton communities (**Supplementary Table 1**). These results have been variable with

findings that include negative results (Scharek et al. 1997; Coale et al. 2003; Ellwood 2004),

slight Zn stimulatory results (Crawford et al. 2003), a "very small increase" relative to controls

in an unreplicated experiment (Coale et al. 2003), Zn stimulation within Fe and Si uptake

experiments (Franck et al. 2003), Zn primary and secondary limitation in the North Pacific in an

unreplicated experiment (Jakuba et al. 2012), secondary Zn limitation after primary Si limitation

in the Costa Rica Dome (Dreux Chappell et al. 2016), and enhanced Zn uptake rates under low

$pCO_2$ (Xu et al. 2012). Whether due to the early negative results, the few positive findings, or a

general prejudice against considering additional factors in controlling marine productivity, it is

our experience that there is currently no broad community recognition that zinc limitation is a

process that could affect primary productivity in any region of the oceans, leaving the original

'zinc hypothesis' unresolved (Morel et al. 1994).

In juxtaposition, laboratory studies have unequivocally demonstrated that marine

phytoplankton can easily be Zn-limited in culture, and that Zn stress is exacerbated by low $CO_2$

due to an inability to synthesize the metalloenzyme carbonic anhydrase and resultant carbon co-

limitation (Morel et al. 1994; Buitenhuis et al. 2003; Sunda and Huntsman 2005). In this study,

we reconcile these perspectives with a comprehensive, multipronged study of the natural

phytoplankton assemblage in Terra Nova Bay (TNB), Antarctica, documenting evidence of Fe

and Zn stress in a low $pCO_2$ coastal environment.

**2 Results**

**2.1 Biogeochemical characterization of Terra Nova Bay**

Terra Nova Bay (TNB), Antarctica, was temporally sampled for over one month (January

9 – February 18, 2018) during the 2017-2018 CICLOPS expedition (**Fig. 1a; Supplementary**

**Table 2**) to concurrently characterize the natural progression of the phytoplankton bloom and

biogeochemical changes in the water column (Kell et al. 2024). Surface waters within TNB had

low (~200 µatm) seawater $pCO_2$ (**Fig. 1b**) which contrasted with measurements >400 µatm

further from the study site (**Fig. 1c**). A large phytoplankton bloom was present as indicated by

high (> 3000 ng $L^{-1}$) chlorophyll fluorescence concentrations in January that waned into

February (**Fig. 1d**), characteristic of Antarctic polynya environments. This phytoplankton

community initially consisted of a mixed assemblage of both diatoms as indicated by

fucoxanthin (fuco, **Fig. 1e**) and *Phaeocystis* as verified by shipboard microscopy and as

indicated by 19'-hexanoyloxyfucoxanthin (19'-hex, **Fig. 1f**). Surface fucoxanthin concentrations

>200 ng L$^{-1}$ were observed at the late TNB stations (**Fig. 1e**) while 19'-hex decreased to ~20 ng

L$^{-1}$ (**Fig. 1f**), indicating that the stations sampled in late February were dominated by diatoms

rather than *Phaeocystis*. This was consistent with historical observations of phytoplankton

succession patterns in TNB (DiTullio and Smith 1996; Smith et al. 2006; Mangoni et al. 2019).

Pronounced and progressively deepening total dissolved Zn (dZn$_T$) depletion over time was

observed, with dZn depleted down to an average of $0.82 \pm 0.47$ nM at 10 m over all TNB

stations (**Fig. 1g**). The Zn-depleted area of the water column progressively deepened to 50 m and

then to ~100 m throughout January and February **(Fig. 1g),** indicative of strong Zn uptake and

export from the euphotic zone. Total Zn uptake ($\rho$Zn, measured concurrently using a stable

isotope tracer method) (Kell et al. 2024) was highest in the shallow euphotic zone in early

January and waned into February (**Fig. 1h**), following trends seen in chlorophyll fluorescence

(**Fig. 1d**) and 19'-hex (**Fig. 1f**). This $\rho$Zn trend was consistent with laboratory studies

demonstrating the substantial Zn requirements of both diatoms and *Phaeocystis antarctica* (Saito

and Goepfert 2008; Kellogg et al. 2020). Across all TNB stations, total dissolved Fe (dFe$_T$) in the

upper 50 m remained below 1 nM (**Fig. 1i**) as observed previously in this region (Fitzwater et al.

2000). In the Ross Sea, dissolved iron (dFe) has previously been demonstrated to be the primary

limiting nutrient for phytoplankton growth (Martin et al. 1990; Coale et al. 2003; Sedwick et al.

2011).

**2.2 Biogeochemical characterization of the incubation study site**

Within TNB station, station 27 (referred to as the "experimental site" herein) was chosen

for the multifactor shipboard incubation experiment (**Fig. 1a,b**; red star). This site harbored a

coastal bloom and was biologically and chemically characterized as having high *in situ*

chlorophyll *a* levels (maximum of 3259 ng $L^{-1}$ at 30 m; **Fig. 1j**) and was comprised of diatoms as

indicated by fucoxanthin and *Phaeocystis* as indicated by 19'-hex (**Fig. 1k**). A decrease in

surface total dissolved inorganic carbon ($DIC_T$; 2181 µmol $kg^{-1}$ at 15 m compared to the deep

water (200-1065 m) average of $2224 \pm 2.1$ µmol $kg^{-1}$**, Fig. 1l**) was also observed. Within the

water column, dZn demonstrated a pronounced decrease from 5.1 nM at 50 m to 0.9 nM at 10 m,

representing an 82% decrease (and a 76% decrease comparing the minimum $dZn_T$ value at 10 m

to the average deepwater (210 – 1000 m) concentration of 3.9 nM $\pm$ 0.4; **Fig. 1m**), consistent

with prior observations of surface dZn depletion in this region (Fitzwater et al. 2000).

Observations of rapid Zn uptake (46 pmol $L^{-1}$ $d^{-1}$ at 10 m) at the experimental site (**Fig.**

**1n**) likely contributed to this surface depletion, as Zn uptake rates of this magnitude are of the

appropriate scale to induce the multi-nanomolar surface water depletion during the austral spring

and summer season (Kell et al. 2024). Consistent with high macronutrient abundance in this

region, surface macronutrient concentrations (nitrate and nitrite, phosphate, and silicic acid) were

partially depleted at the experimental site with 64%, 46%, and 29% decreases in N+N, P, and Si,

respectively, comparing 10 m and average deep water (200 – 1000 m) values (**Fig. 1o**).

**2.3 Evidence for Zn stimulation of phytoplankton: experimental site shipboard incubations**

A multifactor incubation experiment was conducted using surface waters collected at the

experimental site by trace metal clean fish sampler (7 m) fed into a shipboard cleanroom to

examine controls on net primary productivity, with triplicate treatments of Zn amended (+Zn; 2

nM as $ZnCl_2$), Fe amended (+Fe; 1 nM as $FeCl_2$), and Fe and Zn amended (+Fe+Zn) incubations,

in addition to unamended controls. Addition of Fe alone (+Fe) resulted in significantly higher chl

*a* content compared to controls ($p$ = 9.5e-5) after six days (T6) (**Fig. 2a**), demonstrating primary

Fe limitation as observed previously in the Ross Sea (Martin et al. 1990; Mangoni et al. 2019).

However, addition of Zn alone (+Zn) also resulted in significantly higher chl *a* content compared

to the controls ($p = 0.011$), implying that a subset of the incubated phytoplankton population

benefitted from the addition of Zn alone, without additional Fe, and may thus have been

experiencing primary Zn limitation (**Fig. 2a**). The combined addition of +Fe+Zn resulted in the

highest average chl *a* content among all treatments at T6, with $4.5 \pm 0.21$ µg L$^{-1}$ compared to 3.9

$\pm 0.35$ µg L$^{-1}$ and $2.8 \pm 0.15$ µg L$^{-1}$ achieved by +Fe and +Zn alone, respectively, demonstrating

additive co-limitation between Zn and Fe (Sperfeld et al. 2016). The Zn stock solution was

analyzed to confirm these results were not caused by inadvertent iron contamination (see

Methods). Significant differences in seawater chemistry were also observed within these

incubations over time, with larger decreases in DIC$_T$ in all metal treatments compared to the

control (-12.7 µmol kg$^{-1}$ for +Fe ($p = 5.3e-6$), -8.2 µmol kg$^{-1}$for +Zn ($p = 5e-5$), and -18.5 µmol

kg$^{-1}$ for +Fe+Zn ($p = 2.2e-16$); **Fig. 2b**). The decrease in DIC$_T$ observed with +Fe+Zn was

significantly larger than that achieved with +Fe alone ($p = 4.4e-3$; **Fig. 2b**). Statistically

significant differences in measured parameters among treatments are summarized in

**Supplementary Table 3.**

Further consistent with the observed Zn stimulation of biomass in the incubations, the

largest decreases in macronutrient (P and N+N) concentrations in these incubations at T6 were

observed in the +Fe+Zn treatment (**Supplementary Figure 1a,b**), as was the largest increase in

particulate organic carbon (POC; **Supplementary Figure 1c**). POC collected from the +Zn and

+Fe+Zn incubations was characterized by larger C:N atomic ratios (5.9 and 6.2, respectively)

compared to the +Fe and T6 control (5.2 and 5.3; **Supplementary Figure 1d**). Significantly

higher bacterial abundances in both +Fe ($p = 9.1e-4$) and +Fe+Zn ($p = 6.3e-4$) treatments relative

to the T6 control (**Supplementary Figure 1e**) indicated the alleviation of bacterial Fe limitation,

consistent with prior reports (Obernosterer et al. 2015; Fourquez et al. 2020; Sun et al. 2021).

At the conclusion of the incubation experiments, biomass was collected via filtration and

proteomic analysis was performed (see Methods) and analyzed for biomarkers of Zn and Fe

stress. We detected both algal Fe- and Zn-stress proteins, which provided an independent line of

evidence corroborating the results described above (**Fig. 2c**). This included the detection of the

Zn/Co responsive protein ZCRP-A (a putative Zn chaperone) (Kellogg et al. 2022a) as a

biomarker of Zn stress as well as the iron starvation-induced proteins ISIP1A, ISIP2A and ISIP3

(ISIPs) as biomarkers of Fe stress (**Supplementary Table 4**). The ISIPs represent a group of

unrelated proteins that are upregulated under Fe limitation in various algal species. ISIP1

proteins are responsible for endocytosis of siderophore-bound iron, ISIP2 proteins are involved

in $Fe^{3+}$ uptake, and ISIP3 has been suggested to act as an Fe storage protein (Allen et al. 2008;

Behnke and LaRoche 2020). RUBISCO abundance within each treatment is shown in **Fig. 2c** as

a proxy for the potential phytoplankton production. Within the T6 incubation biomass, there was

an increased abundance of ISIPs in the control and +Zn treatment, and a decrease in ISIP protein

abundance within the +Fe and +Fe+Zn treatments, consistent with primary Fe limitation and the

expected response to Fe addition (**Fig. 2c**). ISIPs were taxonomically assigned to diatoms,

*Phaeocystis*, and dinoflagellates (**Fig. 2d**). The strongest expression of ZCRP-A protein was

detected in the +Fe treatment (**Fig. 2c,d**) indicative of Fe addition driving the community

towards increased Zn stress. Notably, ZCRP-A was still detected in the +Fe+Zn treatment (**Fig.

2c,d**), implying that the added Zn was unable to completely satiate Zn demand as phytoplankton

biomass increased (as indicated by the increase in chl *a* at T6, **Fig. 2a**), despite added Zn (2 nM)

being double that of added Fe (1 nM). Sequence analysis of the contigs identified as ZCRP-A

homologs in these incubations revealed that all contigs contained one or more canonical

conserved motifs found in COG0523 family proteins such as ZCRP-A (**Supplementary Figure**

**2**). Coupled with evidence from prior laboratory studies (Kellogg et al. 2022a), this provides

further support for the role of ZCRP-A in responding to Zn scarcity. ZCRP-A proteins were

taxonomically assigned to chlorophytes, dinoflagellates, and *Phaeocystis*, with the detection of

*Phaeocystis* ZCRP-A only in the +Fe treatment (**Fig. 2d**). The detection of ZCRP-A attributed to

*Phaeocystis*, but the nondetection of ZCRP-A attributed to diatoms, implies that diatoms present

in these incubations (as indicated by diatom RUBISCO; **Fig. 2d**) were outcompeting *Phaeocystis*

for Zn. Our observations of Fe and Zn biomarkers shifting in abundance in response to their

respective metal treatment provides independent evidence for Zn/Fe co-limitation.

**2.4 Taxonomic characterization of incubation results**

To characterize the phytoplankton species responding to metal amendment, we measured

phytoplankton pigments within the shipboard incubations over time, which revealed a diverse

taxonomic response to metal amendments. Measured pigments included fucoxanthin (fuco), 19'-

hexanoyloxyfucoxanthin (19'-hex), prasinoxanthin (prasino), chlorophyll *b* (chl *b*), and

chlorophyll c3 (chl c3). Fuco is produced by both diatoms and by *Phaeocystis* under certain

conditions, while 19'-hex and chl c3 are indicative of *Phaeocystis* in the Southern Ocean

(DiTullio et al. 2007). Fuco:19'-hex ratios significantly increased in the +Fe ($p = 4.2e-4$) and

+Fe+Zn treatments ($p = 2.7e-3$) (**Supplementary Figure 3a**) due to no significant change in

fuco (**Supplementary Figure 3b**) and decreased 19'-hex (**Supplementary Figure 3c**) relative to

the T6 control. *Phaeocystis* contributions to total fuco concentrations are typically minimal at the

low Fe levels of the Ross Sea, though *Phaeocystis* can revert to making fuco rather than 19'-hex

when released from Fe limitation (DiTullio et al. 2007), as was evident in these incubations by

decreased 19'-hex:chl c3 ratios within the +Fe and +Fe+Zn treatments (**Supplementary Figure**

**3d**). *Phaeocystis* therefore likely contributed to total fuco by responding to Fe addition. Notably,

significant decreases in both fuco:chl *a* and 19'-hex:chl *a* (**Supplementary Figure 3e,f**) in all

treatments compared to the T6 control indicated that other phytoplankton groups contributed to

chl *a* (**Fig. 2a**) without contributing fuco nor 19'hex. Increases in chl *b* (**Supplementary Figure**

**3g**) and prasinoxanthin (**Supplementary Figure 3h**) suggest that small green algae such as

chlorophytes and prasinophytes also responded to +Fe and +Zn independently, consistent with

the detection of chlorophyte ZCRP-A in these incubations (**Fig. 2d**). Photosynthetic efficiency of

photosystem II (Fv/Fm) significantly increased with +Fe (p = 0.011) and with +Fe+Zn (p =

0.0036) at T4 (day 4) compared to T4 controls, but did not significantly increase with +Zn alone,

implying Fv/Fm may not be useful as a diagnostic for Zn stress and that caution should be used

in interpreting its signals universally (**Supplementary Figure 3i**). No significant difference in

Fv/Fm was observed among treatments at T6. Selective zooplankton grazing on small diatoms

and solitary *Phaeocystis* cells may have played a role in affecting phytoplankton biomass and the

observed pigment:chl *a* ratios. For instance, higher ratios of phaeophytin:total phaeopigments

were observed in +Fe and +Zn amended incubations (**Supplementary Figure 4**) which may

reflect grazing on solitary *Phaeocystis* cells, as high phaeophytin:total phaeopigments ratios were

previously observed in *Phaeocystis* dominated waters of the Ross Sea (DiTullio and Smith

1996).

**2.5 Detection of Zn- and Fe-stress protein biomarkers in the water column**

Metaproteomic and metatranscriptomic analyses of biomass within the water column at

the experimental site provided additional confirmation of the incubation results, as we detected

Zn- and Fe- stress-response proteins present within the water column, which were therefore

naturally present without influence from incubation conditions. In addition to ISIPs and ZCRP-

A, we detected ZCRP-B (a putative membrane-tethered Zn-binding protein) (Kellogg et al.

2022a), Zrt/Irt-like (ZIP) Zn transporters (which are known to be used by marine phytoplankton

for uptake of $Zn^{2+}$ and other divalent metal cations (Allen et al. 2008; Milner et al. 2013; Bender

et al. 2018)), and θ (theta) and δ (delta) carbonic anhydrases (CAs). θ-CAs with $Zn^{2+}$

coordination sites have been documented in diatoms (Jensen et al. 2020), including the polar

diatom *Chaetoceros neogracile* RS19 (Kellogg et al. 2022b), but no studies to date have

investigated enzyme activity nor efficiency with $Co^{2+}$ or $Cd^{2+}$. In contrast, δ-CA (i.e,

*Thalassiosira weissflogii* TWCA1) is known to function with either $Co^{2+}$ or $Zn^{2+}$ as a cofactor

(Lane and Morel 2000b) conferring metabolic flexibility when $Zn^{2+}$ is scarce.

Both proteins and transcripts of Zn and Fe stress biomarkers (ZCRP-A and ISIPs) were

observed throughout the water column at the experimental site. RUBISCO, ZCRP-A, and ISIP

protein spectral counts were most abundant at the surface and decreased with depth within the

within the 3μm size fraction (**Fig. 3a-c**), consistent with the depletion of trace metals in the

photic zone due to high-biomass bloom conditions. ZCRP-A was detected in both 3 and 51 μm

filter pore-size fractions (**Fig. 3b**) and was predominantly attributed to *Phaeocystis* and the

diatom genus *Chaetoceros* in the euphotic zone, and predominantly to *Phaeocystis* and the

diatom genus *Pseudo-nitzschia* in the mesopelagic zone (**Fig. 3i, Supplementary Figure 5a**).

The presence of *Phaeocystis* below the photic zone is consistent with prior observations of rapid

export of *Phaeocystis* cells (DiTullio et al. 2000). Throughout the water column, ISIPs were

predominantly attributed to *Phaeocystis* and to the diatom genera *Fragilariopsis, Chaetoceros,*

and *Pseudo-nitzschia* (**Fig. 3i, Supplementary Figure 5b**). We note that ZCRP-B, a protein also

found to be upregulated in marine diatoms under low Zn/Co and characterized as a putative

membrane-tethered Zn/Co protein ligand (Kellogg et al. 2022a) was most abundant in the 0.2 μm

fraction throughout the water column (**Fig. 3d**). As ZCRP-B shares ~30% similarity to the

bacterial ABC-type nickel transporter component NikA, spectral counts within the bacterial 0.2

μm fraction most likely reflect true bacterial NikA. BLAST analysis of all ZCRP-B contigs

confirmed that all ZCRP-B hits across all size fractions corresponded to bacteria (**Fig. 3i**).

ZIPs were almost solely detected in the 51 μm fraction, likely due to the capture of

abundant *Phaeocystis* colonies and chain-forming diatoms (**Fig. 3e; Supplementary Figure 5c**).

The assignment of the majority of ZCRP-A and ISIP proteins to *Phaeocystis* in the upper water

column provides additional evidence that *Phaeocystis* was likely Zn/Fe co-limited at the study

site, consistent with incubation results (**Fig. 2d**). The increased abundance of diatom θ-CA and

δ-CA proteins within the water column (**Fig. 3f,g**), as well as transcripts for the diatom Cd

carbonic anhydrase CDCA, which can replace $Zn^{2+}$ with $Cd^{2+}$ as the catalytic cofactor (Lane and

Morel 2000a), in the 3 μm and 51 μm fractions at 200 m (**Fig. 3h**) was indicative of a sinking,

prior diatom bloom event (Subhas et al. 2019). θ-CA and δ-CA were predominantly

taxonomically assigned to the diatom genera *Chaetoceros* and *Pseudo-nitzschia*, respectively,

while CDCA transcripts belonged to the diatom genera *Chaetoceros* and *Corethron* (**Fig. 3i**).

The presence of θ-CA, but lack of δ-CA, assigned to *Chaetoceros* is consistent with proteomic

analysis of the polar diatom *Chaetoceros neogracile* RS19 grown in culture under Zn limiting

conditions (Kellogg et al. 2022b).

**2.6 Zn:P ratios of the surface seawater at the experimental site**

A third independent line of evidence for the nutritional influence of Zn scarcity on TNB

phytoplankton was obtained from *in situ* cellular stoichiometry. Particulate Zn:P ratios (Zn:P)

analyzed from biomass collected at the surface of this experimental station were consistent with

ratios from Zn-limited culture studies. Particulate Zn:C ratios reported previously in Zn-limiting

culture studies of the diatom *Thalassiosira pseudonana* (Sunda and Huntsman 2005) were

converted to Zn:P ratios using the Redfield ratio (Redfield 1958) (**Supplementary Table 5**). We

then compared these ratios and associated growth rates with particulate Zn:P measured within

biomass collected at 10, 25, 50 and 100 m at the experimental site. At each of these surface

depths, Zn:P measured at the experimental site was ~ 2E-4 mol:mol, which, in comparison to

cultured diatom Zn:P ratios, fell within the range of severely Zn-limited growth rates

(**Supplementary Figure 6**), again demonstrating the propensity for Zn-limited growth in this

region and corroborating the incubation results.

**3 Discussion**

Antarctic waters are generally considered to not be prone to Zn limitation, given that high

(> 1nM) dZn concentrations are typically observed in surface Southern Ocean waters (Coale et

al. 2003).  However, we observed multiple independent lines of evidence from both the field

incubation experiment (chlorophyll, DIC, Zn and Fe biomarker proteins) and contextual

environmental biogeochemical data of the water column at the incubation site (dZn, Zn uptake

rates, pigments, cellular Zn:P stoichiometry, metaproteomic, and metatranscriptomic analyses)

demonstrating that phytoplankton within Terra Nova Bay of the Ross Sea, Antarctica, were

experiencing Zn and Fe nutritional stress.

Multiple factors could be considered as potential drivers in the creation of Zn-limiting

conditions in the field, including Zn demand imposed by total biomass and the species

comprising this biomass. The phytoplankton bloom observed during this expedition was

comprised primarily of diatoms and *Phaeocystis*, consistent with previous Ross Sea seasonal

blooms (Smith et al. 2006; Arrigo et al. 2012; Mangoni et al. 2019), and which contributed to the

observed high Zn uptake rates and thus surface Zn depletion (Kell et al. 2024), resulting in

nutrient-like dZn profiles throughout TNB.

Our field observation of Zn limitation was made in an environment characterized by

diminished $pCO_2$, which we consider as a factor potentially driving Zn stress. We observed a

substantial drawdown of surface seawater $pCO_2$ to 221 µatm at the incubation site (a ~45%

decrease compared to offshore waters in the Ross Sea measured during the same time frame; **Fig.**

**1b**). Laboratory studies have unequivocally demonstrated that marine phytoplankton can easily

be Zn-limited in culture due to their large Zn requirement, and that this effect is exacerbated at

low $pCO_2$ (Morel et al. 1994; Sunda and Huntsman 2005) due to the use of Zn as a required

catalytic cofactor within carbonic anhydrase (CA) metalloenzymes (Sunda and Huntsman 2005).

CAs catalyze the reversible dehydration of $HCO_3^-$ to $CO_2$, the substrate required by the carbon

fixing enzyme RUBISCO. As $HCO_3^-$ constitutes about 90% of the dissolved inorganic carbon

(DIC) pool in the surface ocean, sufficient CA activity prevents carbon stress in marine

phytoplankton by ensuring adequate $CO_2$ supply to RUBISCO. It has therefore been

hypothesized that the combination of high biomass and resulting low $CO_2$ may cause severe Zn

depletion that may limit algal growth rates due to lack of Zn and thus reduced CA activity, and

thus reduced availability of carbon for photosynthesis (Morel et al. 1994; Sunda and Huntsman

2005). This Zn-C limitation relationship is referred to as 'biochemically dependent co-

limitation', in which the availability of one nutrient is essential for the acquisition or utilization

of another nutrient, especially at low concentrations (Saito et al. 2008).

To explore this in the context of our field observations, using the available quantitative

constraints on Zn and $CO_2$ co-limitation thresholds available from the literature (see Methods),

we estimated that the threshold for Zn-$CO_2$ limitation in culture synthesized across many alga

occurs at 259 μatm $pCO_2$. We then compared this laboratory-determined Zn/C limitation

threshold estimate to both the *in situ* 221 μatm $pCO_2$ measured at our field study site, and to the

historical, global trend in surface ocean $pCO_2$ (**Fig. 4a,b**). Global surface ocean $pCO_2$ levels are

rapidly rising above both the laboratory-estimated 259 μatm $pCO_2$ Zn/C limitation threshold and

our field observation value of 221 μatm (Jiang et al. 2023) (**Fig. 4a,b**). Though only a fraction of

the modern-day surface ocean is currently at ≤ 250 ppm $pCO_2$ (predominantly comprised of

polar regions; **Fig. 4c**), this represents a large decrease in oceanic extent compared to only 100

years ago (**Fig. 4d**). Even though this may move the majority of oceanic regions farther from Zn

and C limitation thresholds, there continue to be highly productive and episodic coastal blooming

events that induce significant $pCO_2$ drawdown (Harrison et al. 2018; Dai et al. 2022). These

coastal regions are increasingly recognized as being disproportionally significant contributors to

global ocean carbon export (with respect to their area), particularly at the high latitudes (Harrison

et al. 2018; Dai et al. 2022), and will hence continue to be prone to Zn stress at low $CO_2$ as we

have observed. Many other coastal regions have been observed to experience depressed $CO_2$

such as the Amundsen Sea (Tortell et al. 2012), Amazon River plume (Valerio et al. 2021), the

west Florida Shelf (Robbins et al. 2018), the East China Sea (Shim et al. 2007), the Northern

Gotland Sea (Schneider and Müller 2018), and Monterey Bay, California (Chavez et al. 2018) to

name a few examples.

## 4 Conclusions

Given the great challenge of conducting Zn manipulation experiments without contamination, we did not try to manipulate $pCO_2$ as an additional experimental treatment. Instead, we actively sought out a low $pCO_2$ environment for the study site, building on prior laboratory studies and a cadmium (Cd) $pCO_2$ field study (Cullen et al. 1999). The interaction of Zn (and Cd and Co) with $CO_2$ is an important area of future research, particularly in coastal environments. With the continuing rise in atmospheric and surface ocean $pCO_2$ levels, broader changes in the biogeochemical cycling of Zn and other bioavailable trace metals will likely occur within the oceans, influencing NPP and thus total ocean carbon storage capacity. These low $pCO_2$ conditions environments that routinely occur in numerous coastal environments globally should be further examined for Zn effects in addition to carbon uptake dynamics in different temperature environments (Tortell et al. 2008; Dai et al. 2022). While there are elaborate biochemical capabilities available to many marine algae for dealing with Zn scarcity (Kellogg et al. 2022a), our results suggest that the geographic extent of possible Zn/C co-limiting environments may further decrease in the coming decades with rising anthropogenic $CO_2$ emissions. Despite this, the biochemical demand for Zn in marine organisms remains substantial, with cellular demand rivaling that of Fe. The multitude of metabolic functions requiring Zn, including but not limited to carbonic anhydrase activity, implies the need for further exploration of Zn influences on primary productivity in a changing ocean environment.

## 5 Materials and Methods

### 5.1 Study area and sample collection

Sample collection occurred during the CICLOPS (Cobalamin and Iron Co-limitation of

Phytoplankton Species) expedition (NBP18-01) aboard the RVIB *Nathaniel B. Palmer*,

December 11, 2017 – March 3, 2018 in the Amundsen Sea and Ross Sea of the Southern Ocean

(**Fig. 1a**). Station metadata is provided in **Supplementary Table 2.** All stations were assumed to

be representative of TNB during this temporal study (as evident in the total dissolved metal,

macronutrient, and chlorophyll *a* datasets). Water samples for dissolved trace metal analyses

were collected using trace metal sampling protocols described previously (Cutter and Bruland

2012). A trace metal clean rosette suspended on a Kevlar line and equipped with twelve 8L X-

Niskin bottles (Ocean Test Equipment) was used to collect seawater at depths ranging from 10 –

600 m. Niskin bottles were transported to a positive-pressure trace metal clean shipboard van for

filtration upon surfacing. Total fluorescence on the vertical profiles was measured using an ECO

chlorophyll fluorometer (Wet Labs) equipped to the rosette. The rosette also included

instrumentation for measuring conductivity and temperature (Sea-Bird Electronics).

**5.2 Preparation of plasticware**

Polyethylene and polycarbonate sampling and incubation bottles were rigorously cleaned

to remove trace metal contaminants before use. Bottles were rinsed with 18.2 Ω Milli-Q water

(Millipore), soaked for 72h in <1% Citranox detergent, rotated, soaked for an additional 72h, and

then rinsed five times with Milli-Q water. Bottles were then filled with 10% HCl (Baker instar-

analyzed) by volume and soaked for a minimum of one week, rotated, and soaked for another

week. Bottles were then rinsed five times with dilute acid (HCl, pH 2) and stored double-bagged

in plastic zip bags. All cleaning work was conducted in a Class 100 clean room.

**5.3 Underway seawater pCO$_2$ partial pressure**

Surface water pCO$_2$ measurements were conducted aboard the RVIB *Nathaniel B.*

*Palmer* using an underway method consisting of an air-water equilibrator and IR CO$_2$ analyzer

developed and operated by the Lamont-Doherty Earth Observatory (LDEO) group (Takahashi et

al. 2020). A complete data report and sensor list are

available: https://service.rvdata.us/data/cruise/NBP1801/doc/NBP1801DATA.pdf (last access:

14 December 2024) (https://www.rvdata.us/, last access: 14 December 2024).

**5.4 TDIC and POC measurements**

Total alkalinity (TA) and dissolved inorganic carbon (DIC) were measured on CTD and

incubations samples in near real-time aboard the NBP. Dissolved inorganic carbon (DIC) and

total alkalinity (TA) samples were collected following the protocols of Dickson et. al

2007(Dickson et al. 2007). DIC analyses were conducted within ~4 h of collection. We acidified

1.25 mL of sample using an automated custom-built injection and bubble stripping system

coupled to an infrared gas analyzer (LICOR LI7000) and integrated the infrared absorption

signal versus time for each stripped gas sample to yield a total mass of CO$_2$. Each sample was

analyzed in triplicate or greater. Since microbubbles regularly formed as samples warmed

between sample acquisition and DIC analysis, every integration curve was visually inspected and

those curves that exhibited evidence for bubbles were rejected. Certified reference materials

(Dickson CRM batch 169) were analyzed between every 3 to 4 unknowns. The estimated

precision based upon unknowns (>860 samples run in triplicate) and CRM replicates (n $=$ 738)

was $\pm$ 2.0 $\mu$mol kg$^{-1}$ ($\pm$1 SD). As described by DeJong et al. 2015(DeJong et al. 2015), analyses

for TA on filtered samples were completed within ~12 h of collection by using a potentiometric

titrator (Metrohm 855 Robotic Titrosampler). The estimated precision based on replicate

analyses of CRMs (n = 195) was ± 2.6 µmol kg$^{-1}$ (±1 SD).

**5.5 Analysis of historical atmospheric and surface ocean pCO$_2$ trends**

Decadal surface ocean pCO$_2$ reconstructions from Jiang et al. 2023 (Jiang et al. 2023)

were downloaded, binned by decade, and plotted using the 'violinplot' in MATLAB.

Atmospheric pCO$_2$ data was assembled from the running Mauna Loa record (Keeling et al.

1976), and from measurements made on Antarctic firn ice (Etheridge et al. 1996).

**5.6 Calculation of Zn-and pCO$_2$ co-limitation of phytoplankton thresholds**

There are few experimental measurements of Zn- and pCO$_2$-co-limitation, either in the

lab or *in situ*. This study documented Zn of a natural phytoplankton assemblage in the field at a

pCO$_2$ of ~220 ppm. In the literature, several models exist to interpret co-limitation (Buitenhuis et

al. 2003; Saito et al. 2008). For this study we chose to use the biochemically dependent co-

limitation model for growth rate (µ):

$$\mu = V_{max} \frac{[CO_{2(aq)}]}{\varphi K_s + [CO_{2(aq)}]}, \quad (1)$$

427 where $V_{max}$ is the maximum growth rate, [CO$_{2(aq)}$] is the aqueous CO$_2$ concentration of the

428 growth medium in micromoles per kilogram of seawater, $K_s$ is the half-saturation constant in

429 micromoles per kilogram of seawater, and φ is a Zn-dependent growth term:

$$\varphi = \frac{[dZn] + K_{s,Zn}}{[dZn]} \quad (2)$$

Here, the dissolved Zn concentration in the growth medium [dZn] is modified by a Zn-dependent

saturation constant ($K_{s,Zn}$). Few studies have enough experimental data to robustly establish a

kinetic relationship between [dZn] and [$CO_{2(aq)}$], so we compiled several estimates for these

terms from the literature. Diatom growth rates under $pCO_2$ limitation were taken from Riebesell

et al. (1993) (Riebesell et al. 1993). Reported pH and temperature measurements for each

treatment, a total alkalinity of 2300 μmol kg$^{-1}$, and a salinity of 35 were used to calculate

aqueous $CO_2$ concentrations using CO2SYSv3.2.1.(Sharp et al. 2023). Reported $V_{max}$ and $K_s$ for

*D. brightwellii, T. punctigera,* and *R. alata* were 1.46, 1.30, and 0.93 d$^{-1}$, and 1.4, 1.2, and 2.1

μmol kg$^{-1}$, respectively. Values for coccolithophore growth ($K_s$ = 0.97 μmol kg$^{-1}$, $V_{max}$ = 4.7 d$^{-1}$)

were taken from Krumhardt et al. (2017) (Krumhardt et al. 2017). A value for $K_{s,Zn}$ of 300 pmol

L$^{-1}$ was taken from Buitenhuis et al. (2003). This value is for the coccolithophore *E. huxleyii*

generated under varying $CO_2$ and Zn conditions, and Zn response growth curves under single

$CO_2$ conditions (ambient) are similar to other diatoms like *T. pseudonana*(Sunda and Huntsman

1995). The value of 300 pmol L$^{-1}$ appears high, but is tied to the functional form of the

biochemically co-limitation equation(Buitenhuis et al. 2003; Saito et al. 2008). Based on the

same dataset and different models for co-limitation, Buitenhuis et al. (2003) arrived at $K_{s,Zn}$

values of ranging from 38 pmol L$^{-1}$ to 300 pmol L$^{-1}$. They calculated Zn-limitation alone, at $CO_2$-

replete conditions, of 19 pmol L$^{-1}$. Thus, the chosen value of $K_{s,Zn}$ is not a reflection of high Zn

demand but determined by the functionality of biochemical co-limitation by Zn and C.

To calculate φ, a surface ocean Zn concentration of 50 pmol L$^{-1}$ was assumed (Bruland

1980; Wyatt et al. 2014). While these concentrations reflect total dissolved Zn, the relationship

between bioavailable free Zn and dZn, especially in the field, remains unclear. Eq. 1 was then

used to calculate effective $CO_2$ concentrations (and thus $pCO_2$ values) at which growth is halved,

or in other words, $\mu = 0.5V_{max}$. We note that this calculation is distinct from the $CO_2$ half-

saturation constant because of the co-limitation by Zn. The median $pCO_2$ threshold for 50%

growth from the three diatom species was 278 ppm. Including coccolithophores decreases the

median to 259 ppm. These values are slightly higher than the *in situ* evidence for Zn limitation at

220 ppm presented in the present study. Our results cannot be considered as an upper bound for

Zn-$CO_2$ limitation, but serve as evidence for growth limitation under those specific

environmental conditions.

**5.7 Analyses of total dissolved metals using isotope dilution**

The analysis of total dissolved metals for this expedition has been described previously

(Kell et al. 2024). Briefly, seawater collected shipboard by pressure-filtering X-Niskin bottles

through an acid-washed 142 mm, 0.2 µM polyethersulfone Supor membrane filter (Pall) within 3

hours of rosette recovery using high purity (99.999%) $N_2$ gas and stored at 4ºC. All sample

collection occurred shipboard within an on-deck trace metal clean van. Samples were acidified to

pH 1.7 with high purity HCl (Optima) within 7 months of collection and were stored acidified at

room temperature for over 1 year prior to analysis.

Quantification of total dissolved Fe, Mn, Ni, Cu, Zn, and Cd was performed using isotope

dilution. Acidified seawater samples were spiked with a stable isotope spike solution artificially

enriched in $^{57}$Fe, $^{61}$Ni, $^{65}$Cu, $^{67}$Zn, and $^{110}$Cd (Oak Ridge National Laboratory). Concentrations

and spike ratios were verified by ICP-MS using a multi-element standard curve (SPEX

CertiPrep). Preconcentration of spiked seawater samples for total dissolved metal analysis was

performed using the automated solid phase extraction system seaFAST-pico (Elemental

Scientific) in offline concentration mode with an initial volume of 15mL and elution volume of

500μL (Rapp et al. 2017; Wuttig et al. 2019). Following preconcentration, multielemental

quantitative analysis was performed using an iCAP-Q inductively coupled plasma-mass

spectrometer (ICP-MS) (Thermo Scientific). Concentrations of Mn, Fe, Ni, Cu, Zn and Cd were

determined using a six-point external standard curve of a multi-element standard (SPEX

CertiPrep), diluted to range from 1-10 ppb in 5% nitric acid. An indium standard (SPEX

CertiPrep) was similarly added to these standard stocks, diluted to range 1-10 ppb. Instrument

injection blanks consisted of 5% nitric acid in Milli-Q. Standard curve $R^2$ values were $\geq 0.98$ for

all metals monitored. Method accuracy and precision were assessed using the 2009

GEOTRACES coastal surface seawater (GSC) standard (n = 8; **Supplementary Table 6**), which

produced values consistent with consensus results (Kell et al. 2024).

**5.8 Macronutrient, pigment, and Fv/Fm analyses**

Seawater for macronutrient (silicate, phosphate, nitrate, and nitrite) analyses were filtered

through 0.2 μm pore-size Supor membrane filters and frozen at sea in acid-washed 60-mL high-

density polyethylene bottles until analysis. Macronutrient analyses were conducted by nutrient

autoanalyzer (Technicon Autoanalyzer II) by Joe Jennings at Oregon State University. The

chemotaxonomic distribution of phytoplankton pigments was determined using HPLC as

described previously (DiTullio et al. 2003). Photosynthetic efficiency of photosystem II (Fv/Fm)

was measured using a Phyto PAM phytoplankton analyzer (Walz, Effeltrich, Germany) as

described previously (Schanke et al. 2021).

**5.9 Bacterial abundance**

One ml samples for heterotrophic prokaryotes abundance (HPA) analysis were fixed for

10 min with a mix of paraformaldehyde and glutaraldehyde (1% and 0.05% final concentration,

respectively), frozen in liquid nitrogen and stored at $-80°C$ until analysis. After thawing,

samples were stained with SYBR Green (Invitrogen Milan, Italy) using $10^{-3}$ dilution of stock

solution for 15 min at room temperature. Cell concentrations were assessed using a FACSVerse

flow cytometer (BD BioSciences Inc., Franklin Lakes, USA) equipped with a 488 nm Ar laser

and standard set of optical filters. FCS Express software was used for analyzing the data and HP

were discriminated from other particles on the basis of scatter and green fluorescence from

SYBR Green (Balestra et al. 2011).

**5.10 ICP-MS analysis and Zn uptake rates using $^{67}$Zn**

$^{67}$Zn stable isotope uptake experiments were performed to quantify the movement of

dissolved Zn to the particulate phase (in units of pmol $L^{-1} d^{-1}$), see (Kell et al. 2024). Briefly,

unfiltered seawater was collected using the trace metal rosette over a depth range of 10 – 600 m

into 250mL trace metal clean polycarbonate bottles. Bottles were spiked with $^{67}$Zn such that the

total added (spiked) concentration of Zn was 2 nM. Immediately after spiking, incubation bottles

were sealed, inverted to mix, and transferred to flow-through on-deck incubators for 24hr.

Biomass was collected after 24hr by vacuum filtering each incubation sample at 34.5 kPa (5 psi)

onto an acid-cleaned 3μm pore-size acrylic copolymer Versapore filter (Pall) mounted on an

acid-cleaned plastic filtration rig. Sample filters were retrieved from storage at -80°C, removed

from cryovials using plastic acid-washed forceps, and transferred into trace metal clean 15 mL

PFA vials with 4 mL of 5% $HNO_3$ (Optima) containing a 1 ppb Indium (In) internal standard.

Filters were digested for ~3.5h at 140°C using a HotBlock® heating block (Environmental

Express, USA) before they were removed and discarded. After evaporating the remaining

solution to just dryness, the residue was resuspended in 2 mL of 5% $HNO_3$ (Optima) by light

vortexing. Process blank filters were digested and processed as sample filters were. This

experiment was also carried out using $^{110}Cd$ as a tracer of Cd uptake in separate incubation

bottles (data not shown here). Digests were analyzed in duplicate by ICP-MS using a Thermo

ICAP-Q plasma mass spectrometer calibrated to a multi-element standard curve (Spex Certiprep)

over a range of 1 – 20 ppb. Natural Cd and Zn isotope abundances of the standards were

assumed to calculate concentrations of $^{110}Cd$, $^{111}Cd$, $^{114}Cd$, $^{67}Zn$, $^{66}Zn$, and $^{68}Zn$. Total Zn uptake

(pmol $L^{-1}$ $d^{-1}$) was calculated using particulate $^{67}Zn$ and total water column dZn measurements as

described previously(Cox et al. 2014). The particulate metal measurements captured

contributions from the active transport of metal into cells, nonspecific metal adsorption to cell

surfaces, metal adsorption to non-living particulate organic matter, and metal adsorption to

particulate inorganic matter, though we expect active transport into cells to dominate the

measured particulate isotopic signal due to the high abundance of actively growing autotrophic

cells in the photic zone observed in Southern Ocean during austral summer. Particulate Zn:P

measurements were calculated using particulate Zn measured on Cd-spiked filters and thus do

not include any pZn contribution from Zn tracer addition. Particulate phosphorus concentrations

were measured by ICP-MS simultaneously and were calibrated to a standard curve ranging from

100 to 3200 ppb using a 1 ppm certified P stock (Alfa Aesar Specpure). All SPEX and P

standard curves had $R^2$ values > 0.99. The Zn stock solution used in the incubation experiments

was similarly analyzed by ICP-MS to confirm that the stock was not Fe contaminated— this

analysis showed that less than 2.3 pM (which was near the instrument blank level for this

analysis) of iron was added for every 2 nM of zinc, far less than needed to stimulate

phytoplankton to the extent observed in our experiments.

**5.11 Shipboard incubation experiments**

Incubation experiments were conducted at station 27 (-74.9870°N, 165.8898°E). Raw

surface seawater was pumped directly into a cleanroom container van, collected into acid-

cleaned 50L carboys, and dispensed into acid-washed 1L polycarbonate bottles using a trace

metal sampling system with acid-washed polypropylene tubing and a teflon diaphragm pump.

Incubation bottles were first rinsed with seawater then filled. Seawater was collected at 16:05

UTC. Triplicate incubation bottles were amended with +Fe (1 nM), +Zn (2 nM) and +Fe+Zn,

sealed, and placed into a flow-through on-deck incubator with light screens that shaded the

incubator to 20% percent ambient surface irradiance. Incubations were sampled at 0, 48, 96, and

144 hours (corresponding to T0, T2, T4, and T6 timepoints) for analysis by filtering onto GFF

filters for chlorophyll (all time points, biological triplicates), pigment analyses (T6, biological

triplicates), and proteomic analyses (T6, pooled biological triplicates). Chlorophyll was extracted

immediately, otherwise samples were frozen at -80°C until further analyses, with pigment and

protein samples kept in -80°C freezers, liquid nitrogen dewars, or dry ice coolers at all times

during transport back to the laboratories. All amendments and sampling were conducted in a

positive-pressure, clean room van with laminar flow hoods and plastic sheeting to minimize

trace-metal contamination.

**5.12 Metaproteomic analysis**

Water column metaproteomic biomass was collected onto 0.2, 3, and 51 µm pore-size

filters using in-situ battery operated McLane pumps. Incubation metaproteomic samples were

collected onto GFF filters. Filters were frozen at -80°C and stored until laboratory extraction.

Frozen filters were thawed and extracted as described previously (Hughes et al. 2014). Briefly,

filters were extracted using SDS detergent to solubilize both membrane and soluble proteins and

subjected to a 10-min heating incubation at 95 °C. Protein material was subsequently purified,

alkylated, reduced and digested with trypsin (mass spectrometry grade; Promega Corporation) at

a trypsin:protein ratio of 1:20 and peptides were purified using C18 tips (Pierce C18 Tips, 100-µl

bed; Thermo Fisher Scientific). Purified peptides were diluted to 0.1 µg µl$^{-1}$ and 20 µl (2 µg) was

injected onto a Dionex UltiMate 3000 RSLCnano LC system (Thermo Fisher Scientific) with an

additional RSLCnano pump run in online two-dimensional active modulation mode coupled to a

Thermo Fusion Orbitrap mass spectrometer.

A translated metatranscriptome (see below) was used as a reference protein database and

peptide spectra matches were performed using the SEQUEST algorithm within Proteome

Discoverer v.2.1 (Thermo Fisher Scientific) with a fragment tolerance of 0.6 Da and parent

tolerance of 10 ppm. Identification criteria consisted of a peptide threshold of 95% (false

discovery rate (FDR) = 0.1%) and protein threshold of 99% (1 peptide minimum, FDR = 0.8%)

in Scaffold v.5 (Proteome Software) resulting in 5,387 proteins identified in the incubation

experiment and 27,924 proteins identified in the water column. To avoid double-counting mass

spectra, exclusive spectral counts were used for the downstream proteomic analysis. Exclusive

spectral counts were normalized using the normalized spectral abundance factor (NSAF)

calculation (Zhang et al. 2010) to allow for a comparison of protein abundance across samples

while remaining consistent with the metatranscriptomic procedure, see Cohen et al. 2021 for

details. Counts associated with redundant ORFs (sharing identical taxonomic and functional

assignments) were summed together. The stand-alone command line application BLAST+ from

the National Center for Biotechnology Information (NCBI) was used to identify proteins of

interest in the metaproteomic data. Metaproteomes were BLAST searched (E = 5e-5) against the

known sequences of proteins of interest acquired from annotated proteomic databases

**(Supplementary Table 4)** and combined with further annotation data based on contig ID (see

below).

**5.13 Metatranscriptomic analysis**

RNA sequencing was performed using the Illumina HiSeq platform. Transcriptomic

assemblies were generated for biomass collected using McLane pumps filtered through 0.2, 3,

and 51 μm pore-size filters. In order to enrich metatranscriptomic libraries derived from 0.2 μm

filters in prokaryotic transcripts and libraries derived from 3 μm and 51 μm filters in eukaryotic

transcripts, 0.2 μm libraries were generated from total rRNA-depleted mRNA and 3 μm and 51

μm libraries were generated from polyA mRNA. Total RNA was extracted from 0.2 μm, 3 μm,

and 51μm filters using Macherey-Nagel a NucleoMag RNA kit (Macherey-Nagel GmbH &

Co.KG). Cleared lysate was loaded into a 96 deep-well plate and put on an epMotion 5075 TMX

liquid handler to complete the RNA extraction following the Machery-Nagel standard protocol.

For 3 μm and 51 μm samples with total RNA greater than 1 μg, 800 ng of total RNA was used

for preparing poly A libraries with an Illumina Stranded mRNA Prep Ligation kit (Illumina),

following the manufacturer's protocol. For the 3 μm and 51 μm samples with total RNA less

than 1 μg, 20 ng of total RNA was used as input for the SMART-Seq v4 Ultra Low Input RNA

kit (Takara Bio USA. Inc), which converts poly(A) RNA to full-length cDNA using a modified

oligo (dT) primer with simultaneous cDNA amplification. The resulting double-stranded cDNA

was then fragmented using a Covaris E210 system with the target size of 300bp. Libraries were

prepared from fragmented double-stranded cDNA using an Illumina Stranded mRNA Prep

Ligation kit (Illumina). For RNA obtained from 0.2 μm filters, ribosomal RNA was removed

using a riboPOOL Seawater Kit (Galen Laboratory Supplies, North Haven, Connecticut, USA).

The riboPOOL Seawater Kit is a customized mixture of Removal Solutions: Pan-Prokaryote

riboPOOL, Pan-Plant riboPOOL and Pan-Mammal in a ratio of 6:1:1. The rRNA-depleted total

RNA was used for cDNA synthesis by Ovation RNA-Seq System V2 (TECAN, Redwood City,

USA). Double stranded cDNA was then prepared for the libraries using an Illumina Stranded

mRNA Prep Ligation kit (Illumina). Ampure XP beads (Beckman Coulter) were used for final

library purification. Library quality was analyzed on a 2200 TapeStation System with an Agilent

High Sensitivity DNA 1000 ScreenTape System (Agilent Technologies, Santa Clara, CA, USA).

Resulting libraries were subjected to paired-end Illumina sequencing via NovaSeq S4.

The input paired-end fastq sequences are trimmed of sequencing adapters, primers and

low quality bases by using either BLASTN (NCBI, v2.2.25) (Altschul et al. 1990) or

trimmomatic, v0.36 (Bolger et al. 2014). The trimmed paired and unpaired sequences were then

depleted of rRNA sequences with riboPicker v0.4.3 (Schmieder et al. 2012). The command-line

program clc_assembler, v5.2.1 (Qiagen) was used to assemble processed sequences into contigs

and ORFs were identified by FragGeneScan, v1.31 (Rho et al. 2010). The trimmed sequences

were mapped to the predicted ORFs using the command-line program clc_mapper, v5.2.1

(Qiagen) to generate mapped raw read counts for each ORF. The raw counts were normalized

initially to RPKM values, to account for variations in inter-sample sequencing depth and the

ORF sequence length (Mortazavi et al. 2008). The RPKM values were subsequently converted to

TPM (transcripts per million) units for estimation of the relative RNA abundance among samples

(Li and Dewey 2011). The ORFs were annotated for putative function by several programs in

parallel using BLASTP against PhyloDB, hidden Markov models PFAM and TIGRFAM by

HMMER, v3.3.2 (Eddy 2011), KEGG Ortholog HMM by kofamscan, v1.3.0 (Aramaki et al.

2020), and transmembrane HMM by TMHMM (Krogh et al. 2001). Additional annotations were

generated by similarity searches using BLASTP to transporter (PhyloDB), organelle (PhyloDB)

and KOG (Tatusov et al. 2003) databases. The ORFs are assigned to the best taxonomic

species/group as determined by LPI (Lineage Probability Index) analysis (Podell and

Gaasterland 2007). The final list of curated ORFs was generated by removing ORFs with low

mapping coverage ($<$ 50 reads total over all samples) and with no BLAST hits and no known

domains.

**5.14 Statistical analysis and data visualization**

ANOVA and Dunnett tests were performed using MATLAB 2019a. Statistics are

summarized in **Supplementary Table 3**. Figures were made using matplotlib (version 3.5.0),

Ocean Data View (version 5.5.2), Excel (2019), and RStudio (version 1.3.1093). Color palettes

used in Ocean Data View section plots (https://doi.org/10.5281/zenodo.1243862) are inverse

"roma" for trace metal concentrations, "thermal" for Zn and Cd uptake rates, and "algae" for

chlorophyll fluorescence (Crameri 2023).

**Data Availability**

CICLOPS (NBP18-01) conductivity–temperature–depth (CTD) hydrography data

including pressure, temperature, total dissolved oxygen, conductivity, fluorescence, and beam

transmission (https://doi.org/10.1575/1912/bco-dmo.783911.1) and total dissolved metal, Zn and

Cd uptake rate, macronutrient, and pigment datasets are available through the NSF Biological

and Chemical Oceanography Data Management Office (BCO-DMO) repository

(https://doi.org/10.7284/907753). Underway $pCO_2$ data collected during cruise NBP1801 are

available through R2R at https://doi.org/10.7284/139318. The mass spectrometry global

proteomics data for CICLOPS bottle incubations and water column analyses have been deposited

with the ProteomeXchange Consortium through the PRIDE repository under the project name

"Zinc-iron co-limitation of natural marine phytoplankton assemblages in coastal Antarctica" with

project accession number PXD037056

(https://www.ebi.ac.uk/pride/archive/projects/PXD037056). This data is accessible for review by

using the following login information: username reviewer_pxd037056@ebi.ac.uk, password:

lFdOUoEb. The translated transcriptome used for spectrum to peptide matching has been

deposited in the National Center for Biotechnology Information sequence read archive under

BioProject accession no. PRJNA890306

(https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA890306) and RNA-Seq BioSample

accession nos. SAMN31286421-SAMN31286522

(https://www.ncbi.nlm.nih.gov/biosample/?term=SAMN31286421).

**Acknowledgements**

We thank the captain, crew, marine technicians and science party of RVIB Nathaniel B.

Palmer for their support and contributions to the success of the NBP18-01 cruise. We thank

Natalie Cohen for assistance and training with the SeaFAST. We thank Veronique Oldham for

assistance with trace metal sampling. This work was funded by the National Science Foundation

**Author contributions statement**

All authors contributed to data acquisition and analysis. RMK, NLS, LEL, RJC, DR, DMM,

MRM, FB, RBD, GRD and MAS implemented the shipboard incubation study. RMK, MMB,

RJC, DR, TJH, and AVS contributed to the formal analysis. NLS, LEL, FB, OM, RF, and GRD

contributed to pigment datasets and interpretations. CB provided bacterial abundance data. RBD

contributed DIC and *in situ* $pCO_2$ data. AVS contributed analyses and discussion regarding

historical $pCO_2$ data and Zn-C growth limitation estimates. AEA contributed the

metatranscriptome reference database used for proteomic analyses. RMK and MAS wrote the

original draft. RMK, MAS, GRD, AVS, TJH, and RBD contributed to review and editing. All

authors approved the final submitted manuscript.

**Competing interests statement**

697 The authors declare no competing interests.

**Figures**

707
708

**Figure 1**. **Temporal biogeochemistry of Terra Nova Bay and characterization of the**
**experimental site at Station 27.** (**a**) Sampling locations over the Ross Sea shelf in Terra Nova
Bay, Antarctica. (**b**) Location of station 27 (red star) and surrounding seawater $pCO_2$ measured
over a three-day transit northwards represented in color scale. (**c**) $pCO_2$ measured over time
within TNB during the three-day transit shown in (b). The vertical red line denotes the $pCO_2$
level at the time of initial seawater collection at station 27. (**d**) Total chlorophyll fluorescence,
(**e**) fucoxanthin (fuco), (**f**) 19' hexanoyloxyfucoxanthin (19'-hex), (**g**) total dissolved Zn, (**h**) total
Zn uptake rates, and (**i**) total dissolved Fe measured in the upper 250 m represented on a color
scale. Station data is presented in order of sampling date, from the earliest (Stn 22, early January)
to the latest (Stn 79, late February). The data gap between January 13-23 occurred when the ship
was unable to sample due to icebreaking duties for the McMurdo Station resupply ship. Stations

indicated in (a) are those where the trace metal rosette (TMR) was deployed; pigment data was supplemented with additional TNB stations using a CTD (Table S2). Depth profiles of (**j**) chlorophyll a, (**k**) the pigments fuco and 19'hex, (**l**) total dissolved inorganic carbon (DICT), (**m**) total dissolved Zn, (**n**) total Zn uptake rates, and (**o**) the macronutrients nitrate+nitrite (N+N), phosphate (P), and silicate (Si) at the study site. Panels (**d**),(**g**),(**h**) and (**i**) were originally presented in Kell et al. (2024) and are reprised here to introduce the environmental context of the study site.

750
**Figure 2. Evidence for Zn co-limitation with Fe in bottle incubations.** (**a**) Chlorophyll *a* and
(**b**) total DIC (DICT) at T0 (day 0) and in each treatment at T6 (day 6). Significant differences
among groups were found using one-way ANOVA and post-hoc Dunnett test (*** $p < 0.001$, **
$p < 0.01$, * $p < 0.05$). Error bars are the standard deviation of biological triplicates (n=3).
Individual data points are overlaid (white circles). (**c**) Heatmap of row-scaled exclusive protein
spectral counts (normalized total spectra) showing relative protein abundance in each treatment.
The dendrogram shows similarity in spectral abundance among samples based on Euclidean
distance and hierarchical clustering. Color gradients represent low (yellow) to high (blue) protein
expression. Ribulose-1,5-biphosphate carboxylase/oxygenase (RUBISCO), zinc/cobalt
responsive protein A (ZCRP-A), and iron starvation induced proteins (ISIP1, ISIP2A, ISIP3) are
shown. (**d**) Taxonomies assigned to RUBISCO, ZCRP-A, and ISIP proteins in each treatment at
T6. Counts (normalized total spectra) assigned to each taxa are shown. ISIPs are the combined
spectral counts of ISIP1A, ISIP2A and ISIP3.

**Figure 3. Metatranscriptomic and metaproteomic detection of Zn- and Fe-related proteins of interest at the experimental site**. Depth profiles of summed NSAF-normalized protein spectral counts of (**a**) RUBISCO, (**b**) ZCRP-A, (**c**) iron starvation induced proteins (ISIPs), (**d**) ZCRP-B, (**e**) ZIP, (**f**) Theta CA, and (**g**) Delta CA detected from proteomic analysis of each filter size fraction (0.2, 3 and 51μm). (**h**) TPM-normalized transcript read counts of CDCA. (**i**) Stacked pie charts depicting relative community composition for proteins of interest for euphotic (< 200 m) and mesopelagic (≥ 200 m) depths. The outer rings show community composition based on NSAF-normalized protein spectral counts while the inner rings are TPM-normalized transcript read counts. Protein and transcript counts plotted in (**i**) were summed across all size fractions. ISIPs are the combined spectral counts of ISIP1A, ISIP2A and ISIP3.

790
**Figure 4**. **The partial pressure of CO₂ (pCO₂) and associated phytoplankton responses from**
**this study and the literature**. (**a**) Pre-industrial and decadal surface ocean pCO₂ reconstructions
plotted as violin plots, with a running black line through the median values. The atmospheric
curve is a composite of ice core data (dashed yellow line (Etheridge et al. 1996)) and the Mauna
Loa record (solid yellow line (Keeling et al. 1976)). An estimated thresholds for zinc-limited
growth is plotted as the median of previous laboratory results (259 μatm, dark green line; see
Methods), and is compared to the *in situ* results of this study (220 μatm, light green line). (**b**)
Data in (a) plotted as a histogram comparing preindustrial and modern (2010) pCO₂ values, with
the same pCO2 levels indicated. (**c**) Global map of surface ocean pCO₂ plotted using
GLODAPv2.2022 data (Lauvset et al. 2022). (**b**) Percentage of the ocean surface less than 250
μatm pCO₂ as a function of time. Surface ocean pCO₂ reconstructions taken from Jiang et al.
2023.

**References**

Allen, A. E. and others. 2008. Whole-cell response of the pennate diatom Phaeodactylum

tricornutum to iron starvation. Proc. Natl. Acad. Sci. **105**: 10438–10443.

doi:10.1073/pnas.0711370105

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment

search tool. J. Mol. Biol. doi:10.1016/S0022-2836(05)80360-2

Aramaki, T., R. Blanc-Mathieu, H. Endo, K. Ohkubo, M. Kanehisa, S. Goto, and H. Ogata.

2020. KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive

score threshold A. Valencia [ed.]. Bioinformatics **36**: 2251–2252.

doi:10.1093/bioinformatics/btz859

Arrigo, K. R., K. E. Lowry, and G. L. van Dijken. 2012. Annual changes in sea ice and

phytoplankton in polynyas of the Amundsen Sea, Antarctica. Deep Sea Res. Part II Top.

Stud. Oceanogr. **71–76**: 5–15. doi:10.1016/j.dsr2.2012.03.006

Baars, O., and P. L. Croot. 2011. The speciation of dissolved zinc in the Atlantic sector of the

Southern Ocean. Deep Sea Res. Part II Top. Stud. Oceanogr. **58**: 2720–2732.

doi:10.1016/j.dsr2.2011.02.003

Balestra, C., L. Alonso-Sáez, J. M. Gasol, and R. Casotti. 2011. Group-specific effects on coastal

bacterioplankton of polyunsaturated aldehydes produced by diatoms. Aquat. Microb.

Ecol. doi:10.3354/ame01486

Behnke, J., and J. LaRoche. 2020. Iron uptake proteins in algae and the role of Iron Starvation-

Induced Proteins (ISIPs). Eur. J. Phycol. **55**: 339–360.

doi:10.1080/09670262.2020.1744039

Bender, S. J. and others. 2018. Colony formation in Phaeocystis antarctica: connecting molecular

mechanisms with iron biogeochemistry. Biogeosciences **15**: 4923–4942. doi:10.5194/bg-

15-4923-2018

Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics **30**: 2114–2120. doi:10.1093/bioinformatics/btu170

Bruland, K. W. 1980. Oceanographic distributions of cadmium, zinc, nickel, and copper in the

North Pacific. Earth Planet. Sci. Lett. **47**: 176–198. doi:10.1016/0012-821X(80)90035-7

Bruland, K. W. 1989. Complexation of zinc by natural organic ligands in the central North

Pacific. Limnol. Oceanogr. **34**: 269–285. doi:10.4319/lo.1989.34.2.0269

Buitenhuis, E. T., K. R. Timmermans, and H. J. W. de Baar. 2003. Zinc-bicarbonate colimitation

of Emiliania huxleyi. Limnol. Oceanogr. **48**: 1575–1582. doi:10.4319/lo.2003.48.4.1575

Chavez, F. P., J. Sevadjian, C. Wahl, J. Friederich, and G. E. Friederich. 2018. Measurements of

pCO2 and pH from an autonomous surface vehicle in a coastal upwelling system. Deep

Sea Res. Part II Top. Stud. Oceanogr. **151**: 137–146. doi:10.1016/j.dsr2.2017.01.001

Coale, K. H., X. Wang, S. J. Tanner, and K. S. Johnson. 2003. Phytoplankton growth and

biological response to iron and zinc addition in the Ross Sea and Antarctic Circumpolar

Current along 170°W. Deep Sea Res. Part II Top. Stud. Oceanogr. **50**: 635–653.

doi:10.1016/S0967-0645(02)00588-X

Cox, A. D., A. E. Noble, and M. A. Saito. 2014. Cadmium enriched stable isotope uptake and

addition experiments with natural phytoplankton assemblages in the Costa Rica

Upwelling Dome. Mar. Chem. **166**: 70–81. doi:10.1016/j.marchem.2014.09.009

Crameri, F. 2023. Scientific colour maps.doi:10.5281/ZENODO.1243862

Crawford, D. W. and others. 2003. Influence of zinc and iron enrichments on phytoplankton

growth in the northeastern subarctic Pacific. Limnol. Oceanogr. **48**: 1583–1600.

doi:10.4319/lo.2003.48.4.1583

Cullen, J. T., T. W. Lane, F. M. M. Morel, and R. M. Sheerell. 1999. Modulation of cadmium

uptake in phytoplankton by seawater $CO_2$ concentration. Nature. doi:10.1038/46007

Cutter, G. A., and K. W. Bruland. 2012. Rapid and noncontaminating sampling system for trace

elements in global ocean surveys. Limnol. Oceanogr. Methods.

doi:10.4319/lom.2012.10.425

Dai, M. and others. 2022. Carbon Fluxes in the Coastal Ocean: Synthesis, Boundary Processes,

and Future Trends. Annu. Rev. Earth Planet. Sci. **50**: 593–626. doi:10.1146/annurev-

earth-032320-090746

DeJong, H. B., R. B. Dunbar, D. Mucciarone, and D. A. Koweek. 2015. Carbonate saturation

state of surface waters in the Ross Sea and Southern Ocean: controls and implications for

the onset of aragonite undersaturation. Biogeosciences **12**: 6881–6896. doi:10.5194/bg-

12-6881-2015

Dickson, A. G., C. L. Sabine, and J. R. Christian,. 2007. Guide to best practices for ocean $CO_2$

measurement. North Pacific Marine Science Organization.

DiTullio, G. R. and others. 2000. Rapid and early export of Phaeocystis antarctica blooms in the

Ross Sea, Antarctica. Nature. doi:10.1038/35007061

DiTullio, G. R., N. Garcia, S. F. Riseman, and P. N. Sedwick. 2007. Effects of iron concentration

on pigment composition in Phaeocystis antarctica grown at low irradiance.

Biogeochemistry **83**: 71–81. doi:10.1007/s10533-007-9080-8

DiTullio, G. R., M. E. Geesey, A. Leventer, and M. P. Lizotte. 2003. Algal pigment ratios in the

Ross Sea: Implications for Chemtax analysis of Southern Ocean data, p. 35–51. *In*.

DiTullio, G. R., and W. O. Smith. 1996. Spatial patterns in phytoplankton biomass and pigment

distributions in the Ross Sea. J. Geophys. Res. Oceans **101**: 18467–18477.

doi:10.1029/96JC00034

Dreux Chappell, P., J. Vedmati, K. E. Selph, H. A. Cyr, B. D. Jenkins, M. R. Landry, and J. W.

Moffett. 2016. Preferential depletion of zinc within Costa Rica upwelling dome creates

conditions for zinc co-limitation of primary production. J. Plankton Res. **38**: 244–255.

doi:10.1093/plankt/fbw018

Eddy, S. R. 2011. Accelerated Profile HMM Searches W.R. Pearson [ed.]. PLoS Comput. Biol.

**7**: e1002195. doi:10.1371/journal.pcbi.1002195

Ellwood, M. J. 2004. Zinc and cadmium speciation in subantarctic waters east of New Zealand.

Mar. Chem. doi:10.1016/j.marchem.2004.01.005

Etheridge, D. M., L. P. Steele, R. L. Langenfelds, R. J. Francey, J.-M. Barnola, and V. I.

Morgan. 1996. Natural and anthropogenic changes in atmospheric $CO_2$ over the last

1000 years from air in Antarctic ice and firn. J. Geophys. Res. Atmospheres **101**: 4115–

4128. doi:10.1029/95JD03410

Fitzwater, S. E., K. S. Johnson, R. M. Gordon, K. H. Coale, and W. O. Smith. 2000. Trace metal

concentrations in the Ross Sea and their relationship with nutrients and phytoplankton

growth. Deep-Sea Res. Part II Top. Stud. Oceanogr. **47**: 3159–3179. doi:10.1016/S0967-

0645(00)00063-1

Fourquez, M., M. Bressac, S. L. Deppeler, M. Ellwood, I. Obernosterer, T. W. Trull, and P. W.

Boyd. 2020. Microbial Competition in the Subpolar Southern Ocean: An Fe–C Co-

limitation Experiment. Front. Mar. Sci. **6**: 776. doi:10.3389/fmars.2019.00776

Franck, V., K. Bruland, D. Hutchins, and M. Brzezinski. 2003. Iron and zinc effects on silicic

acid and nitrate uptake kinetics in three high-nutrient, low-chlorophyll (HNLC) regions.

Mar. Ecol. Prog. Ser. **252**: 15–33. doi:10.3354/meps252015

Harrison, C. S., M. C. Long, N. S. Lovenduski, and J. K. Moore. 2018. Mesoscale Effects on

Carbon Export: A Global Perspective. Glob. Biogeochem. Cycles **32**: 680–703.

doi:10.1002/2017GB005751

Hughes, C. S., S. Foehr, D. A. Garfield, E. E. Furlong, L. M. Steinmetz, and J. Krijgsveld. 2014.

Ultrasensitive proteome analysis using paramagnetic bead technology. Mol. Syst. Biol.

**10**: 757. doi:10.15252/msb.20145625

Jakuba, R. W., M. A. Saito, J. W. Moffett, and Y. Xu. 2012. Dissolved zinc in the subarctic

North Pacific and Bering Sea: Its distribution, speciation, and importance to primary

producers. Glob. Biogeochem. Cycles. doi:10.1029/2010GB004004

Jensen, E. L., S. C. Maberly, and B. Gontero. 2020. Insights on the Functions and

Ecophysiological Relevance of the Diverse Carbonic Anhydrases in Microalgae. Int. J.

Mol. Sci. **21**: 2922. doi:10.3390/ijms21082922

Jiang, L. and others. 2023. Global Surface Ocean Acidification Indicators From 1750 to 2100. J.

Adv. Model. Earth Syst. **15**: e2022MS003563. doi:10.1029/2022MS003563

Keeling, C. D., R. B. Bacastow, A. E. Bainbridge, C. A. Ekdahl, P. R. Guenther, L. S.

Waterman, and J. F. S. Chin. 1976. Atmospheric carbon dioxide variations at Mauna Loa

Observatory, Hawaii. Tellus Dyn. Meteorol. Oceanogr. **28**: 538.

doi:10.3402/tellusa.v28i6.11322

Kell, R. M. and others. 2024. High metabolic zinc demand within native Amundsen and Ross sea

phytoplankton communities determined by stable isotope uptake rate measurements.

Biogeosciences **21**: 5685–5706. doi:10.5194/bg-21-5685-2024

Kellogg, R. M. and others. 2022a. Adaptive responses of marine diatoms to zinc scarcity and

ecological implications. Nat. Commun. 2022 131 **13**: 1–13. doi:10.1038/s41467-022-

29603-y

Kellogg, R. M., M. R. McIlvin, J. Vedamati, B. S. Twining, J. W. Moffett, A. Marchetti, D. M.

Moran, and M. A. Saito. 2020. Efficient zinc/cobalt inter-replacement in northeast Pacific

diatoms and relationship to high surface dissolved Co : Zn ratios. Limnol. Oceanogr. **65**:

2557–2582. doi:10.1002/lno.11471

Kellogg, R. M., D. M. Moran, M. R. McIlvin, A. V. Subhas, A. E. Allen, and M. A. Saito.

2022b. Lack of a Zn/Co substitution ability in the polar diatom *Chaetoceros neogracile*

RS19. Limnol. Oceanogr. **67**: 2265–2280. doi:10.1002/lno.12201

Krogh, A., B. Larsson, G. von Heijne, and E. L. L. Sonnhammer. 2001. Predicting

transmembrane protein topology with a hidden markov model: application to complete

genomes11Edited by F. Cohen. J. Mol. Biol. **305**: 567–580. doi:10.1006/jmbi.2000.4315

Krumhardt, K. M., N. S. Lovenduski, M. D. Iglesias-Rodriguez, and J. A. Kleypas. 2017.

Coccolithophore growth and calcification in a changing ocean. Prog. Oceanogr. **159**:

276–295. doi:10.1016/j.pocean.2017.10.007

Lane, T. W., and F. M. Morel. 2000a. A biological function for cadmium in marine diatoms.

Proc. Natl. Acad. Sci. U. S. A. **97**: 4627–31. doi:10.1073/pnas.090091397

Lane, T. W., and F. M. M. Morel. 2000b. Regulation of Carbonic Anhydrase Expression by

Zinc, Cobalt, and Carbon Dioxide in the Marine Diatom Thalassiosira weissflogii. Plant

Physiol. **123**: 345–352. doi:10.1104/pp.123.1.345

Lauvset, S. K. and others. 2022. GLODAPv2.2022: the latest version of the global interior ocean

biogeochemical data product. Earth Syst. Sci. Data **14**: 5543–5572. doi:10.5194/essd-14-

5543-2022

Lhospice, S. and others. 2017. Pseudomonas aeruginosa zinc uptake in chelating environment is

primarily mediated by the metallophore pseudopaline. Sci. Rep. **7**: 17132.

doi:10.1038/s41598-017-16765-9

Li, B., and C. N. Dewey. 2011. RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. BMC Bioinformatics **12**: 323. doi:10.1186/1471-

2105-12-323

Lohan, M. C., P. J. Statham, and D. W. Crawford. 2002. Total dissolved zinc in the upper water

column of the subarctic North East Pacific. Deep Sea Res. Part II Top. Stud. Oceanogr.

**49**: 5793–5808. doi:10.1016/S0967-0645(02)00215-1

Mangoni, O., M. Saggiomo, F. Bolinesi, M. Castellano, P. Povero, V. Saggiomo, and G. R.

DiTullio. 2019. Phaeocystis antarctica unusual summer bloom in stratified antarctic

coastal waters (Terra Nova Bay, Ross Sea). Mar. Environ. Res.

doi:10.1016/j.marenvres.2019.05.012

Martin, J. H., S. E. Fitzwater, and R. M. Gordon. 1990. Iron deficiency limits phytoplankton

growth in Antarctic waters. Glob. Biogeochem. Cycles **4**: 5–12.

doi:10.1029/GB004i001p00005

Mazzotta, M. G., M. R. McIlvin, D. M. Moran, D. T. Wang, K. D. Bidle, C. H. Lamborg, and M.

A. Saito. 2021. Characterization of the metalloproteome of Pseudoalteromonas (BB2-

AT2): biogeochemical underpinnings for zinc, manganese, cobalt, and nickel cycling in a

ubiquitous marine heterotroph. Metallomics **13**. doi:10.1093/mtomcs/mfab060

Middag, R., H. J. W. De Baar, and K. W. Bruland. 2019. The Relationships Between Dissolved

Zinc and Major Nutrients Phosphate and Silicate Along the GEOTRACES GA02

Transect in the West Atlantic Ocean. Glob. Biogeochem. Cycles **33**: 63–84.

doi:10.1029/2018GB006034

Milner, M. J., J. Seamon, E. Craft, and L. V. Kochian. 2013. Transport properties of members of

the ZIP family in plants and their role in Zn and Mn homeostasis. J. Exp. Bot. **64**: 369–

381. doi:10.1093/jxb/ers315

Moore, C. M. and others. 2013. Processes and patterns of oceanic nutrient limitation. Nat.

Geosci. **6**: 701–710. doi:10.1038/ngeo1765

Morel, F. M. M., J. R. Reinfelder, S. B. Roberts, C. P. Chamberlain, J. G. Lee, and D. Yee. 1994.

Zinc and carbon co-limitation of marine phytoplankton. Nature **369**: 740–742.

doi:10.1038/369740A0

Mortazavi, A., B. A. Williams, K. McCue, L. Schaeffer, and B. Wold. 2008. Mapping and

quantifying mammalian transcriptomes by RNA-Seq. Nat. Methods **5**: 621–628.

doi:10.1038/nmeth.1226

Obernosterer, I., M. Fourquez, and S. Blain. 2015. Fe and C co-limitation of heterotrophic

bacteria in the naturally fertilized region off the Kerguelen Islands. Biogeosciences **12**:

1983–1992. doi:10.5194/bg-12-1983-2015

Podell, S., and T. Gaasterland. 2007. DarkHorse: a method for genome-wide prediction of

horizontal gene transfer. Genome Biol. **8**: R16. doi:10.1186/gb-2007-8-2-r16

Rapp, I., C. Schlosser, D. Rusiecka, M. Gledhill, and E. P. Achterberg. 2017. Automated

preconcentration of Fe, Zn, Cu, Ni, Cd, Pb, Co, and Mn in seawater with analysis using

high-resolution sector field inductively-coupled plasma mass spectrometry. Anal. Chim.

Acta. doi:10.1016/j.aca.2017.05.008

Redfield, A. C. 1958. The Biological Control of Chemical Factors in the Environment. Am. Sci.

doi:10.2307/27828530

Rho, M., H. Tang, and Y. Ye. 2010. FragGeneScan: predicting genes in short and error-prone

reads. Nucleic Acids Res. **38**: e191–e191. doi:10.1093/nar/gkq747

Riebesell, U., D. A. Wolf-Gladrow, and V. Smetacek. 1993. Carbon dioxide limitation of marine

phytoplankton growth rates. Nature **361**: 249–251. doi:10.1038/361249a0

Robbins, L. L. and others. 2018. Spatial and Temporal Variability of $p$CO$_2$ , Carbon Fluxes, and

Saturation State on the West Florida Shelf. J. Geophys. Res. Oceans **123**: 6174–6188.

doi:10.1029/2018JC014195

Saito, M. A., and T. J. Goepfert. 2008. Zinc-cobalt colimitation of *Phaeocystis antarctica*.

Limnol. Oceanogr. **53**: 266–275. doi:10.4319/lo.2008.53.1.0266

Saito, M. A., T. J. Goepfert, and J. T. Ritt. 2008. Some thoughts on the concept of colimitation:

Three definitions and the importance of bioavailability. Limnol. Oceanogr. **53**: 276–290.

doi:10.4319/lo.2008.53.1.0276

Schanke, N. L., F. Bolinesi, O. Mangoni, C. Katlein, P. Anhaus, M. Hoppmann, P. A. Lee, and

G. R. DiTullio. 2021. Biogeochemical and ecological variability during the late summer–

early autumn transition at an ice-floe drift station in the Central Arctic Ocean. Limnol.

Oceanogr. doi:10.1002/lno.11676

Scharek, R., M. A. Van Leeuwe, and H. J. W. De Baar. 1997. Responses of Southern Ocean

phytoplankton to the addition of trace metals. Deep-Sea Res. Part II Top. Stud. Oceanogr.

doi:10.1016/S0967-0645(96)00074-4

Schmieder, R., Y. W. Lim, and R. Edwards. 2012. Identification and removal of ribosomal RNA

sequences from metatranscriptomes. Bioinformatics **28**: 433–435.

doi:10.1093/bioinformatics/btr669

Schneider, B., and J. D. Müller. 2018. Biogeochemical transformations in the Baltic Sea:

observations through carbon dioxide glasses, Springer.

Sharp, J. D., D. Pierrot, M. P. Humphreys, J.-M. Epitalon, J. C. Orr, E. R. Lewis, and D. W. R.

Wallace. 2023. CO2SYSv3 for MATLAB.doi:10.5281/ZENODO.3950562

Shim, J., D. Kim, Y. C. Kang, J. H. Lee, S.-T. Jang, and C.-H. Kim. 2007. Seasonal variations in

pCO2 and its controlling factors in surface seawater of the northern East China Sea. Cont.

Shelf Res. **27**: 2623–2636. doi:10.1016/j.csr.2007.07.005

Smith, W. O., A. R. Shields, J. A. Peloquin, G. Catalano, S. Tozzi, M. S. Dinniman, and V. A.

Asper. 2006. Interannual variations in nutrients, net community production, and

biogeochemical cycles in the Ross Sea. Deep-Sea Res. Part II Top. Stud. Oceanogr.

doi:10.1016/j.dsr2.2006.02.014

Sperfeld, E., D. Raubenheimer, and A. Wacker. 2016. Bridging factorial and gradient concepts

of resource co-limitation: towards a general framework applied to consumers J. Grover

[ed.]. Ecol. Lett. **19**: 201–215. doi:10.1111/ele.12554

Subhas, A. V., J. F. Adkins, S. Dong, N. E. Rollins, and W. M. Berelson. 2019. The carbonic

anhydrase activity of sinking and suspended particles in the North Pacific Ocean. Limnol.

Oceanogr. **65**: 637–651. doi:10.1002/lno.11332

Sun, Y., P. Debeljak, and I. Obernosterer. 2021. Microbial iron and carbon metabolism as

revealed by taxonomy-specific functional diversity in the Southern Ocean. ISME J. **15**:

2933–2946. doi:10.1038/s41396-021-00973-3

Sunda, W. G., and S. A. Huntsman. 1995. Cobalt and zinc interreplacement in marine

phytoplankton: Biological and geochemical implications. Limnol. Oceanogr. **40**: 1404–

1417. doi:10.4319/lo.1995.40.8.1404

Sunda, W. G., and S. A. Huntsman. 2000. Effect of Zn, Mn, and Fe on Cd accumulation in

phytoplankton: Implications for oceanic Cd cycling. Limnol. Oceanogr. **45**: 1501–1516.

doi:10.4319/lo.2000.45.7.1501

Sunda, W. G., and S. A. Huntsman. 2005. Effect of $CO_2$ supply and demand on zinc uptake and

growth limitation in a coastal diatom. Limnol. Oceanogr. **50**: 1181–1192.

doi:10.4319/lo.2005.50.4.1181

Takahashi, T., S. C. Sutherland, and A. Kozyr. 2020. Global Ocean Surface Water Partial

Pressure of CO2 Database (LDEO Database Version 2019): Measurements Performed

During 1957-2019 (NCEI Accession 0160492). NOAA Natl. Cent. Environ. Inf.

Tatusov, R. L. and others. 2003. The COG database: an updated version includes eukaryotes.

BMC Bioinformatics **4**: 41. doi:10.1186/1471-2105-4-41

Tortell, P. D., M. C. Long, C. D. Payne, A.-C. Alderkamp, P. Dutrieux, and K. R. Arrigo. 2012.

Spatial distribution of pCO2, $\Delta O_2/Ar$ and dimethylsulfide (DMS) in polynya waters and

the sea ice zone of the Amundsen Sea, Antarctica. Deep Sea Res. Part II Top. Stud.

Oceanogr. **71–76**: 77–93. doi:10.1016/j.dsr2.2012.03.010

Tortell, P. D., C. Payne, C. Gueguen, R. F. Strzepek, P. W. Boyd, and B. Rost. 2008. Inorganic

carbon uptake by Southern Ocean phytoplankton. Limnol. Oceanogr. **53**: 1266–1278.

doi:10.4319/lo.2008.53.4.1266

Valerio, A. M., M. Kampel, N. D. Ward, H. O. Sawakuchi, A. C. Cunha, and J. E. Richey. 2021.

CO2 partial pressure and fluxes in the Amazon River plume using in situ and remote

sensing data. Cont. Shelf Res. **215**: 104348. doi:10.1016/j.csr.2021.104348

Wuttig, K. and others. 2019. Critical evaluation of a seaFAST system for the analysis of trace

metals in marine samples. Talanta. doi:10.1016/j.talanta.2019.01.047

Wyatt, N. J., A. Milne, E. M. S. Woodward, A. P. Rees, T. J. Browning, H. A. Bouman, P. J.

Worsfold, and M. C. Lohan. 2014. Biogeochemical cycling of dissolved zinc along the

GEOTRACES South Atlantic transect GA10 at 40°S. Glob. Biogeochem. Cycles **28**: 44–

56. doi:10.1002/2013GB004637

Xu, Y., D. Shi, L. Aristilde, and F. M. M. Morel. 2012. The effect of pH on the uptake of zinc

and cadmium in marine phytoplankton: Possible role of weak complexes. Limnol.

Oceanogr. doi:10.4319/lo.2012.57.1.0293

Ye, N. and others. 2022. The role of zinc in the adaptive evolution of polar phytoplankton. Nat.

Ecol. Evol. **6**: 965–978. doi:10.1038/s41559-022-01750-x

Zhang, Y., Z. Wen, M. P. Washburn, and L. Florens. 2010. Refinements to Label Free Proteome

Quantitation: How to Deal with Peptides Shared by Multiple Proteins. Anal. Chem. **82**:

2272–2281. doi:10.1021/ac9023999

1073

**Supplementary information for "Zinc stimulation of phytoplankton in a low carbon dioxide, coastal Antarctic environment: evidence for the Zn hypothesis"**

1076

1077 **Supplementary Table 1. Summary of zinc incubation experiments in marine environments.**
1078 Note that all experiments focus on phytoplankton populations with the exception of Fukada et al
1079 (2000) and Mazzotta et al. (2021) which focus on bacterial heterotrophy. This list represents our
1080 knowledge of the literature but may be incomplete.

1081

| Study | Location | Zn Treatment Replication | pCO$_2$ Data | Dissolved Environmental Zn Data | Zn Response |
|---|---|---|---|---|---|
| Scharek et al. 1997[1] | Southern Ocean | Zn n=2, control n=1 | No | No | Negative/"very small" |
| Fukada et al., 2000[2] | North Pacific | Integrated water column leucine aminopeptidase activity (0-100m) | No | Yes | Leucine aminopeptidase activity positively correlated with dZn |
| Cochlan et al. 2002[3] | Western Ross Sea | None (n=1 +Zn, +Zn+Fe) | No | No | Increase in specific uptake of NO$_3^-$ |
| Coale et al. 2003[4] | Ross Sea, Southern Ocean | None (n=1 +Zn, +Zn+Fe) | No | Yes | Negative/"very small" |
| Crawford et al. 2003[5] | Northeast Pacific Ocean | Triplicate (n=3) | Yes | No | "Slightly but significantly" altered Chl, nitrate and phosphate |
| Franck et al. 2003[6] | Costa Rica Dome | Triplicate (n=3) | No | No | Secondary Zn limitation after Fe in diatom microscopy data |
| Ellwood 2004[7] | Subantarctic | Duplicates (n=2) | No | Yes | Negative |
| Cullen et al. 1999; Cullen and Sherrell 2005[8,9] | Coastal California | Duplicates (n=2) | No | No | Enhanced Cd:P in low pCO$_2$ treatments, decrease in Cd:P with Zn addition |
| Jakuba et al. 2012[10] | North Pacific | Singlicate with timepoints sacrificed | No | Yes | Primary, no additive effect with iron. |
| Dreux Chappell et al. 2016[11] | Costa Rica Dome | Triplicate (n=3) | No | Yes | Secondary limitation with Si |
| Sharma et al. 2020[12] | Eastern Arabian Sea | Triplicate (n=3) | Yes | No | Negative/ "insignificant effects" |
| Mazzotta et al. 2021[13] | Equatorial Pacific | Quintuplicate (n=5) sediment trap incubations | No | Yes in separate study (Cohen et al., 2021) | Enhancement of alkaline phosphatase activity with Zn addition in sediment trap samples |
| This study | Terra Nova Bay, Ross Sea | Triplicate (n=3) | Yes | Yes | Primary and Secondary limitation. Independent validation by Zn biomarkers. |

1082  **Supplementary Table 2**. Station metadata for the NBP18-01 cruise. Stations at which total Zn
1083  uptake rates were determined are indicated by asterisks (*).
1084

| Station | Latitude (°N) | Longitude (°E) | Sampling Date (yyyy-mm-dd hh:mm) |
|---------|---------------|----------------|----------------------------------|
| 4* | -72.751 | -116.001 | 2017-12-30 01:23 |
| 10 | -73.054 | -129.988 | 2018-01-03 03:39 |
| 11* | -74.047 | -133.764 | 2018-01-03 19:58 |
| 15* | -75.864 | -151.918 | 2018-01-05 14:51 |
| 20* | -76.714 | 179.819 | 2018-01-08 02:00 |
| 22* | -75.013 | 165.358 | 2018-01-09 15:56 |
| 25 | -75.293 | 163.914 | 2018-01-11 01:26 |
| 27* | -74.987 | 165.890 | 2018-01-11 16:05 |
| 29* | -76.001 | 172.997 | 2018-01-16 03:00 |
| 31 | -77.295 | 175.390 | 2018-01-17 04:39 |
| 32* | -76.750 | 172.000 | 2018-01-17 19:19 |
| 34 | -77.147 | 168.503 | 2018-01-23 22:59 |
| 35* | -76.231 | 168.769 | 2018-01-26 20:17 |
| 41* | -74.833 | 165.002 | 2018-01-29 00:41 |
| 46* | -74.742 | 165.287 | 2018-01-31 21:32 |
| 50 | -74.741 | 165.488 | 2018-02-02 21:17 |
| 52* | -75.000 | 164.005 | 2018-02-03 21:43 |
| 57* | -74.879 | 164.482 | 2018-02-06 20:12 |
| 60 | -74.959 | 164.739 | 2018-02-08 20:06 |
| 62* | -74.999 | 169.491 | 2018-02-09 19:27 |
| 67* | -76.454 | 167.919 | 2018-02-11 19:19 |
| 70 | -74.744 | 170.374 | 2018-02-13 21:25 |
| 72* | -74.800 | 164.395 | 2018-02-14 22:28 |
| 76* | -74.799 | 164.597 | 2018-02-16 20:15 |
| 78 | -74.696 | 164.796 | 2018-02-17 20:34 |
| 79* | -74.757 | 164.356 | 2018-02-18 19:29 |

1085

1086

1087

1088

1089

1090
1091
1092
1093

**Supplementary Table 3**. Summary of ANOVA statistics for T6 incubation treatments
comparing measured parameters. Significant differences among groups were found using one-
way ANOVA and post-hoc Dunnett test (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, . $p < 0.1$). Chl $a$,
chlorophyll $a$; $DIC_T$, total dissolved inorganic carbon; chl $b$, chlorophyll $b$; Prasino,
prasinoxanthin; fuco, fucoxanthin; 19'hex, 19'-hexanoyloxyfucoxanthin; chl c3, chlorophyll c3.
NA, no statistically significant difference. All parameters were measured in biological triplicate.
1100

| Treatment (T6) | Chl $a$ | $DIC_T$ | Chl $b$ | Prasino | Fuco | 19'hex | Fuco:hex | Hex:Chl c3 | Bacterial abundance |
|---|---|---|---|---|---|---|---|---|---|
| +Fe vs Ctrl | $p$ = 9.5e-5 (***) | $p$ = 5.3e-6 (***) | $p$ = 4.0e-3 (**) | $p$ = 2.9e-2 (*) | NA | $p$ = 8e-3 (**) | $p$ = 4.2e-4 (***) | $p$ = 2.0e-4 (***) | $p$ = 9.1e-4 (***) |
| +Zn vs Ctrl | $p$ = 1.1e-2 (*) | $p$ = 5.0e-6 (***) | $p$ = 8.0e-2 (.) | $p$ = 7.4e-3 (**) | NA | NA | NA | $p$ = 0.02520 (*) | NA |
| +FeZn vs Ctrl | $p$ = 1.3e-7 (***) | $p$ = 2.2e-16 (***) | $p$ = 4.0e-4 (***) | $p$ = 5.7e-2 (.) | NA | NA | $p$ = 2.7e-3 (**) | $p$ = 1.9e-4 (***) | $p$ = 6.3e-4 (***) |
| +FeZn vs +Fe | $p$ = 3.4e-2 (*) | $p$ = 4.4e-3 (**) | NA | NA | NA | $p$ = 6.0e-2 (.) | NA | NA | NA |

1101

1102

1103 **Supplementary Table 4.** Representative proteins of interest, reference organism and IDs.

| Protein of interest | Reference organism | Protein ID |
|---|---|---|
| ZCRP-A | *Thalassiosira pseudonana* CCMP1335 | 3054 (JGI Thaps3[*]) |
| ZCRP-B | *Thalassiosira pseudonana* CCMP1335 | 938 (JGI Thaps3_bd[**]) |
| RUBISCO | *Phaeodactylum tricornutum* CCMP632 | AAF07200.1 (NCBI) |
| ISIP1A | *Thalassiosira oceanica* CCMP1005 | K0RCT3 (Uniprot) |
| ISIP2A | *Phaeodactylum tricornutum* CCMP632 | B7FYL2 (Uniprot) |
| ISIP3 | *Phaeodactylum tricornutum* CCMP632 | B7G4H8 (Uniprot) |
| ZIP | *Phaeodactylum tricornutum* CCMP632 | 46780 (JGI Phatr2[†]) |
| CDCA | *Thalassiosira pseudonana* CCMP1335 | 25840 (JGI Thaps3[*]) |

[*]Joint Genome Institute (JGI) Thaps3 database
(https://mycocosm.jgi.doe.gov/Thaps3/Thaps3.home.html)
[**]Joint Genome Institute Thaps3_bd database
(https://mycocosm.jgi.doe.gov/Thaps3_bd/Thaps3_bd.home.html
[†]Joint Genome Institute CCAP 1055/1 v2.0 Phatr2, all models database
(https://mycocosm.jgi.doe.gov/Phatr2/Phatr2.home.html)

**Supplementary Table 5**. Particulate Zn:P ratios measured in *T. pseudonana* cultured at various
concentrations of Zn (log [Zn′]) (Sunda and Huntsman 2005) compared to particulate Zn:P ratios
measured at Station 27 of research cruise NBP18-01. Zn:C measurements by Sunda and
Huntsman 2005 were converted to Zn:P using the Redfield ratio. Data is plotted in
Supplementary Figure 6. Zn′ refers to the sum of inorganic Zn metal complexes.
+Particulate P data measured directly in this study
*Particulate P from Sunda 2005 data was estimated by converting particulate C measurements to
P using the Redfield ratio (106C:1P).

1126

| Particulate Zn:P ratios at Station 27 | | | | | | | |
|---|---|---|---|---|---|---|---|
| Depth (m) | | | Particulate Zn (mol) | | Particulate P (mol)+ | Zn:P (mol:mol) | |
| 100 | | | 1E-11 | | 8E-08 | 1E-04 | |
| 50 | | | 1E-11 | | 1E-07 | 1E-04 | |
| 25 | | | 3E-11 | | 2E-07 | 2E-04 | |
| 10 | | | 4E-11 | | 2E-07 | 2E-04 | |
| Cellular Zn:P ratios of cultured *T. pseudonana* from Sunda and Huntsman 2005, Table 1. | | | | | | | |
| Experiment | Log [Zn'] | pH | Particulate Zn (mol) | Particulate C (mol) | Particulate P (mol)* | Zn:P (mol:mol) | Growth rate (d⁻¹) |
| 1 | -12.05 | 8.2 | 3.3E-05 | 22 | 0.21 | **2E-04** | **0.1** |
| 1 | -11.47 | 8.2 | 5.2E-05 | 15 | 0.14 | 4E-04 | 1.23 |
| 1 | -10.87 | 8.2 | 1.2E-04 | 15 | 0.14 | 9E-04 | 1.62 |
| 1 | -10.36 | 8.2 | 1.5E-04 | 15 | 0.14 | 1E-03 | 1.76 |
| 1 | -9.82 | 8.2 | 1.8E-04 | 15 | 0.14 | 1E-03 | 1.76 |
| 1 | -11.96 | 9 | 3.8E-05 | 15 | 0.14 | 3E-04 | 0.51 |
| 1 | -11.38 | 9 | 8.5E-05 | 15 | 0.14 | 6E-04 | 1.12 |
| 1 | -10.78 | 9 | 2.0E-04 | 15 | 0.14 | 1E-03 | 1.28 |
| 1 | -10.27 | 9 | 4.1E-04 | 15 | 0.14 | 3E-03 | 1.42 |
| 1 | -9.73 | 9 | 5.0E-04 | 15 | 0.14 | 4E-03 | 1.49 |
| 2 | -11.82 | 8.2 | 4.9E-05 | 15 | 0.14 | 3E-04 | 0.64 |
| 2 | -11.82 | 8.2 | 4.0E-05 | 15 | 0.14 | 3E-04 | 0.79 |
| 2 | -10.87 | 8.2 | 1.3E-04 | 15 | 0.14 | 9E-04 | 1.45 |
| 2 | -9.82 | 8.2 | 2.0E-04 | 15 | 0.14 | 1E-03 | 1.51 |
| 2 | -11.71 | 9 | 5.8E-05 | 15 | 0.14 | 4E-04 | 0.73 |
| 2 | -11.71 | 9 | 5.8E-05 | 15 | 0.14 | 4E-04 | 0.78 |
| 2 | -10.76 | 9 | 1.9E-04 | 15 | 0.14 | 1E-03 | 1.39 |
| 2 | -10.76 | 9 | 2.1E-04 | 15 | 0.14 | 1E-03 | 1.38 |
| 2 | -9.71 | 9 | 4.3E-04 | 15 | 0.14 | 3E-03 | 1.44 |
| 2 | -9.71 | 9 | 5.0E-04 | 15 | 0.14 | 4E-03 | 1.42 |
| 4 | -11.82 | 8.2 | 2.8E-05 | 18.1 | 0.21 | 3E-04 | 0.72 |
| 4 | -11.32 | 8.2 | 5.5E-05 | 15.2 | 0.13 | 4E-04 | 1.45 |
| 4 | -10.82 | 8.2 | 1.3E-04 | 14.7 | 0.14 | 9E-04 | 2 |
| 4 | -10.32 | 8.2 | 2.2E-04 | 15.1 | 0.14 | 2E-03 | 2.03 |
| 4 | -9.82 | 8.2 | 2.6E-04 | 15.6 | 0.14 | 2E-03 | 2.04 |

1127
1128

1129

1130

1131

1132

1133 **Supplementary Table 6**. Reference seawater comparisons using the 2009 GEOTRACES coastal
1134 surface seawater (GSC) standard.

| Metal | This study (n = 8) (nM) | GEOTRACES GSC consensus (nM) |
|-------|------------------------|------------------------------|
| Fe | $1.6 \pm 0.23$ | $1.6 \pm 0.12$ |
| Zn | $1.4 \pm 0.23$ | $1.5 \pm 0.10$ |
| Cd | $0.4 \pm 0.01$ | $0.4 \pm 0.02$ |
| Cu | $1.3 \pm 0.05$ | $1.1 \pm 0.15$ |
| Ni | $4.2 \pm 0.07$ | $4.5 \pm 0.21$ |
| Mn | $2.1 \pm 0.37$ | $2.2 \pm 0.08$ |

1135

1136
1137 **Supplementary Figures**
1138
1139

1140
1141
**Supplementary Figure 1. Additional parameters measured in shipboard bottle incubations**.
Time course decreases in (**a**) phosphate and (**b**) nitrite + nitrate. (**c**) total POC of biomass in each
treatment at T6. (**d**) the atomic carbon:nitrogen (C:N) ratio of biomass in each treatment at T6,
and (**e**) bacterial abundance. Significant differences among groups were found using one-way
ANOVA and post-hoc Dunnett test (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$). Error bars are the
standard deviation of biological triplicates (n=3) with individual data points overlaid (white

circles). Macronutrients, POC, and PON were measured in singlicate (n=1) from pooled
biological triplicates.
1150
1151

a.

b.

| orf_id | Hit | E value | % ID |
|---|---|---|---|
| contig_652592_49_1335_- | Phaeodactylum_tricornutum_ZCRPA | 4.75E-162 | 56.266 |
| contig_104831_3_980_- | Phaeodactylum_tricornutum_ZCRPA | 6.51E-110 | 50.311 |
| contig_947870_2_1288_- | Phaeodactylum_tricornutum_ZCRPA | 5.00E-86 | 49.488 |
| contig_104832_3_716_- | Phaeodactylum_tricornutum_ZCRPA | 5.25E-80 | 48.954 |
| contig_120421_1_351_- | Phaeodactylum_tricornutum_ZCRPA | 1.62E-32 | 51.818 |
| contig_343121_503_1144_+ | Phaeodactylum_tricornutum_ZCRPA | 3.24E-23 | 60.526 |
| contig_120413_3_335_- | Phaeodactylum_tricornutum_ZCRPA | 5.54E-22 | 41.964 |
| contig_120390_752_1060_- | Phaeodactylum_tricornutum_ZCRPA | 7.76E-13 | 41.111 |

1152

1153

**Supplementary Figure 2. Sequence alignments of ZCRP-A peptides detected in T6**
**incubation biomass. (a)** Sequence alignment of the *Phaeodactylum tricornutum* ZCRP-A
protein compared to all ZCRP-A proteins detected in T6 incubation biomass. Alignment was
generated using the MUSCLE algorithm with default parameters within MEGA11. Four
conserved GTPase (G1/Walker A, G2/SwitchI, CXCC metal binding, and G3/Walker B) are
labeled. (**b**) E values and % identities of the identified proteins with significant sequence
similarity to *P. tricornutum* ZCRP-A aligned above.

**Supplementary Figure 3. Pigment analysis of T6 incubations**. (**a**) ratio of fucoxanthin: 19'-
Hex, (**b**) fucoxanthin, (**c**) 19-hexanoyloxyfucoxanthin (19'-Hex), (**d**) ratio of 19'-Hex: chl c3, (**e**)
ratio of fucoxanthin: chlorophyll *a*, (**f**) ratio of 19'-Hex: chlorophyll *a*, (**g**) chlorophyll *b*, (**h**)
prasinoxanthin, and (**i**) maximum quantum efficiency (Fv/Fm) among treatments at T4.
Significant differences among groups were found using one-way ANOVA and post-hoc Dunnett
test (*** p < 0.001, ** p < 0.01, * p < 0.05, . p < 0.1). Data with error bars are presented as mean
values ± the standard deviation of biological triplicates (n=3) with individual data points overlaid
(white circles).

1181
1182 **Supplementary Figure 4.** Phaeopigments measured in T6 incubation biomass. Abundances of
1183 (**a**) phaeophorbide and (**b**) phaeophytin, and (**c**) the ratio of phaeophytin: total phaeopigments
1184 (the sum of phaeophorbide and phaeophytin). Error bars are the standard deviation of biological
1185 triplicates (n=3).

1196
1197 **Supplementary Figure 5. Depth profiles for proteins of interest at station 27 categorized by**
1198 **genus.** Depth profiles of NSAF-normalized protein spectral counts of (**a**) ZCRP-A, (**b**), iron
1199 starvation induced proteins (ISIPs), and (**c**) ZIPs summed by genus. Proteins assigned to an
1200 individual genus were summed across all size fractions (0.2, 3 and 51 μm). ISIPs are the
1201 combined spectral counts of ISIP1A, ISIP2A and ISIP3.

1214

1215

1216

1217

**Supplementary Figure 6. Comparison of particulate Zn: particulate P ratios measured in the water column at the experimental site to those ratios measured in culture studies of a Zn-limited diatom.** Ratio of particulate Zn (pZn) and particulate phosphorus (pP) (pZn:pP) measured in the upper water column at the study site (station 27; vertical dashed lines) compared to pZn:pP measurements in Zn limitation studies of the diatom *Thalassiosira pseudonana* in culture by Sunda and Huntsman 2005.

1224

1244
**Supplementary Information References**

1.    Scharek, R., Van Leeuwe, M. A. & De Baar, H. J. W. Responses of Southern Ocean

phytoplankton to the addition of trace metals. *Deep-Sea Res. Part II Top. Stud.*

*Oceanogr.* (1997) doi:10.1016/S0967-0645(96)00074-4.

2.    Fukuda, R. *et al.* East—west gradient in ectoenzyme activities in the subarctic Pacific:

Possible regulation by zinc. *Limnol. Oceanogr.* **45**, 930–939 (2000).

3.    Cochlan, W. P., Bronk, D. A. & Coale, K. H. Trace metals and nitrogenous nutrition of

Antarctic phytoplankton: experimental observations in the Ross Sea. *Deep Sea Res. Part*

*II Top. Stud. Oceanogr.* **49**, 3365–3390 (2002).

4.    Coale, K. H., Wang, X., Tanner, S. J. & Johnson, K. S. Phytoplankton growth and

biological response to iron and zinc addition in the Ross Sea and Antarctic Circumpolar

Current along 170°W. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **50**, 635–653 (2003).

5.    Crawford, D. W. *et al. Influence of Zinc and Iron Enrichments on Phytoplankton Growth*

*in the Northeastern Subarctic Pacific Limitation of Biomass and Primary Production of*

*Phyto-Plankton by Dissolved Iron (Fe) in High-Nutrient, Low-Chlo*. *Limnol. Oceanogr*

vol. 48 1583–1600 (2003).

6.    Franck, V., Bruland, K., Hutchins, D. & Brzezinski, M. Iron and zinc effects on silicic

acid and nitrate uptake kinetics in three high-nutrient, low-chlorophyll (HNLC) regions.

*Mar. Ecol. Prog. Ser.* **252**, 15–33 (2003).

7.    Ellwood, M. J. Zinc and cadmium speciation in subantarctic waters east of New Zealand.

*Mar. Chem.* (2004) doi:10.1016/j.marchem.2004.01.005.

8.    Cullen, J. T., Lane, T. W., Morel, F. M. M. & Sheerell, R. M. Modulation of cadmium

uptake in phytoplankton by seawater CO2 concentration. *Nature* (1999)

doi:10.1038/46007.

9.    Cullen, J. T. & Sherrell, R. M. Effects of dissolved carbon dioxide, zinc, and manganese

on the cadmium to phosphorus ratio in natural phytoplankton assemblages. *Limnol.*

*Oceanogr.* **50**, 1193–1204 (2005).

10.    Jakuba, R. W., Saito, M. A., Moffett, J. W. & Xu, Y. Dissolved zinc in the subarctic

North Pacific and Bering Sea: Its distribution, speciation, and importance to primary

producers. *Glob. Biogeochem. Cycles* (2012) doi:10.1029/2010GB004004.

11.    Dreux Chappell, P. *et al.* Preferential depletion of zinc within Costa Rica upwelling dome

creates conditions for zinc co-limitation of primary production. *J. Plankton Res.* **38**, 244–

255 (2016).

12.    Sharma, D. *et al.* Impacts of Zn and Cu enrichment under ocean acidification scenario on

a phytoplankton community from tropical upwelling system. *Mar. Environ. Res.* **155**,

104880 (2020).

13.    Mazzotta, M. G. *et al.* Characterization of the metalloproteome of Pseudoalteromonas

(BB2-AT2): biogeochemical underpinnings for zinc, manganese, cobalt, and nickel

cycling in a ubiquitous marine heterotroph. *Metallomics* **13**, (2021).

14.    Sunda, W. G. & Huntsman, S. A. Effect of CO2 supply and demand on zinc uptake and

growth limitation in a coastal diatom. *Limnol. Oceanogr.* **50**, 1181–1192 (2005).

15.    Lauvset, S. K. *et al.* GLODAPv2.2022: the latest version of the global interior ocean

biogeochemical data product. *Earth Syst. Sci. Data* **14**, 5543–5572 (2022).

16.    Jiang, L. *et al.* Global Surface Ocean Acidification Indicators From 1750 to 2100. *J. Adv.*

*Model. Earth Syst.* **15**, e2022MS003563 (2023).

1290