From sequence to ecology: siderophore-receptor coevolution algorithm

predicts bacterial interactions in complex communities

Shaohua Gu[1,2,3], Zhengying Shao[3], Yuanzhe Shao[2], Shenyue Zhu[3], Di Zhang[1], Richard Allen[4],

Ruolin He[1], Jiqi Shao[1], Guanyue Xiong[1], Zeyang Qu[1], Alexandre Jousset[3], Ville-Petri Friman[5],

Zhong Wei[3*], Rolf Kümmerli[4*], Zhiyuan Li[1,2*]

[1] Center for Quantitative Biology, Academy for Advanced Interdisciplinary Studies, Peking

University, Beijing, 100871, China

[2] Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies,

Peking University, Beijing, 100871, China

[3] Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, Key lab of organic-based

fertilizers of China, Nanjing Agricultural University, Nanjing, P R China

[4] University of Zurich, Department of Quantitative Biomedicine, Winterthurerstr. 190, 8057

Zurich, Switzerland

[5] University of Helsinki, Department of Microbiology, 00014, Helsinki, Finland

[*]Corresponding authors (email: weizhong@njau.edu.cn; rolf.kuemmerli@uzh.ch;

zhiyuanli@pku.edu.cn)

Abstract

Unlocking the secrets of microbial interactions through genomics is pivotal for advancing

microbial ecology. In most ecosystems, the scarcity of iron makes iron-mediated interactions a

central theme in shaping microbial communities. Bacteria have evolved diverse strategies,

including the production of siderophores—diverse secondary metabolites—to scavenge iron

from their surroundings. Here, we use bioinformatic tools to predict siderophore

iron-interaction networks among 1928 *Pseudomonas* strains from sequence data. Our

approach uses coevolution analysis to group siderophore synthetase clusters and receptors

used for uptake into key-lock pairs. Through a mix of computational analyses and

experimental validation, we reconstruct *Pseudomonas* iron-interaction networks across a

spectrum of habitats, from soil to water, plants, and human-related environments and reveal

substantial differences in network structure and connectivity across habitats. Altogether, our

sequence-to-interaction mapping tool empowers researchers to decode bacterial ecology in

complex microbiomes, setting the stage for novel interventions to engineer microbiome

functionality.

Introduction

Microbial communities populate all ecosystems on earth from terrestrial to aquatic environments, impacting human health, agriculture, and industry [1-3]. The dynamics and functioning of these communities are shaped by complex interactions between microorganisms [4,5]. As the number of sequenced microbial genomes continues to grow exponentially [6,7], there is a growing interest in predicting microbial interaction networks based on the genomic data. Such efforts potentiate major advances, as many microorganisms cannot be cultured in the laboratory [8], while their roles in natural communities can potentially be inferred through sequence-to-interaction mapping. Currently, sequence-to-interaction mapping approaches primarily focus on metabolic interactions, with Genome-scale Metabolic Models (GEMs) serving as the primary tool for establishing the pan-reactome of microbial communities [9,10]. These methods infer metabolic reactions from the genome annotation of enzymes, and then reconstruct a flux model to understand how microorganisms take up essential nutrients and release metabolic byproducts into the environment [11-13].

Despite the significance of primary nutrients, like carbon and nitrogen, there is increasing evidence that secondary metabolism also plays a major role in shaping microbial interactions [14,15]. Nearly all microbes actively synthesize compounds to fulfill a diverse set of functions, including resource scavenging, motility, attack of and defense against competitors, and communication [16,17]. These compounds, referred to as "secondary metabolites," were previously considered non-essential for microbial growth in a laboratory setting, but have since been shown to be critical for competitiveness in natural environments [14,15,18,19]. However,

genome data-based sequence-to-interaction mapping have rarely been applied to secondary

metabolism, although this could provide fundamentally new insights into microbial community

assembly rules.

Here, we developed a secondary metabolite sequence-to-interaction approach focusing

on iron-scavenging siderophores, one of the most prevalent classes of microbial secondary

metabolites [20]. Iron is a critical nutrient for microbial survival, as it is used as a catalytic group

in enzymes guiding key biological processes such as respiration and replication [21].

However, the concentration of bioavailable iron is typically below the level required for

microbial survival in most habitats [21-23]. In response to iron limitation, nearly all bacteria

produce siderophores, a chemically diverse class of low molecular weight compounds that

efficiently chelate iron from the environment [24,25]. Siderophores are typically diffusible and able

to chelate iron over a broad physical range [26]. Once iron is bound, the complex is recognized

and taken up by specific receptors in microbial cell membranes [25]. Given their diffusible nature,

siderophores mediate a range of social interactions. For bacteria possessing receptors

capable of recognizing the iron-siderophore complex, siderophores act as a public good

promoting cooperation between individuals [25,27]. When bacteria possess receptors for

siderophore uptake but do not produce siderophores themselves they can act as cheaters by

exploiting the public goods secreted by others [24,25]. Finally, for bacteria unable to recognize

and uptake a specific siderophore, the iron-siderophore complex restricts access to iron and

intensifies iron competition [25,28]. Consequently, siderophore-mediated interactions can have an

important impact on microbial community composition and dynamics [29-32], yet the prediction of

such interactions from sequence data is an unsolved challenge.

We have previously developed bioinformatic pipelines to predict the chemical structure of

siderophores and to identify receptors from sequence data. Here, we apply our data mining

approach to infer how receptor and siderophore had co-evolved and to find matching

siderophore-receptor pairs to predict interaction networks in bacterial communities. We focus

on *Pseudomonas* spp. (1928 strains) featuring 188 predicted variants of pyoverdine (their

main siderophore) and 94 groups of FpvA receptors (siderophore receptors for pyoverdines)[33].

We developed a Co-evolution Pairing Algorithm that revealed nearly 50 unique lock-key

groups, where groups of FpvA receptors emerge as specific "locks" that recognize

corresponding pyoverdines as the "key". Validation experiments yielded prediction accuracies

of over 90%. Based on the predicted lock-key pairs, we reconstructed the iron-interaction

networks among the 1928 *Pseudomonas* strains. We noticed that network topologies differ

fundamentally between different ecological habitats (soil, plant, water, human). Taken together,

our work provides a robust sequence-to-interaction mapping tool to predict social interaction

networks mediated by secondary metabolites in complex microbial communities.

Results

Section 1: Three classes of pyoverdine strategies in *Pseudomonas* and the lock-key
(receptor-synthetase) principle of co-evolution

In our recent work, we developed bioinformatic pipelines to predict pyoverdine molecule

structures and receptors based on the genome sequence data. Among 1928 nonredundant

Pseudomonas strains, we identified 188 chemically different pyoverdine types and 94 different

receptor groups[33]. In this work, we seek to understand how different *Pseudomonas* strains

interact through these pyoverdines and receptors.

We first compared all the strains according to phylogeny, ecological habitat, and

pyoverdine function. At the phylogenetic level, our data set features a large diversity of

Pseudomonas species, whereby *P. aeruginosa* (28.7%), *P. fluorescens* (7.0%), *P. syringae*

(6.0%), and *P. putida* (2.2%) were the most abundant ones (Figure 1a). The strains originated

from a diverse set of habitats, including humans (21.2%), soils (13.6%), plants (12.1%), and

water (6.4%), although the origin of many strains (39.5%) is unknown (Figure 1a).

To assess diversity at the pyoverdine functional level, we checked for the absence or

presence of pyoverdine synthesis clusters and counted the number of FpvA receptors per

strain. We found three basic types of pyoverdine-utilization strategies (Figure 1b).

"Single-receptor producers" are the most common type (985 strains, 51.1%) and refer to

strains with one pyoverdine synthesis locus and one FpvA receptor gene. "Multi-receptor

producers" are the second most common type (679 strains, 35.2%) and refer to strains with

one pyoverdine synthesis cluster and multiple receptor genes. "Non-producers" are the least

common type (264 strains, 13.7%) and refer to strains that lack the pyoverdine synthesis

cluster but contain at least on receptor gene. While each strain possesses on average two to

three FpvA receptor genes, no strain carries more than one pyoverdine synthesis cluster. This

observation is in line with the expected high costs of pyoverdine synthetase, which is based on

a series of gigantic modular enzymes known as NPRS (non-ribosomal peptide synthetases)[34].

Figure 1 Classification of *Pseudomonas* strains and elucidation of the co-evolution

between pyoverdine synthetase and receptors. a. Phylogenetic relationship among the 1928

Pseudomonas strain based on the concatenated alignment of 400 single-copy conserved genes. Starting

from inside, colors in the first ring distinguish the five most prevalent species, with "Others" representing

the remaining less abundant species. Colors in the second ring distinguish the four most prevalent

sources of isolation. In the third ring, claret and blank regions cover strains with complete pyoverdine

synthetase clusters and strains without synthetase gene clusters, respectively. In the fourth blue ring, the

bar height indicates the number of FpvA receptors present in each strain. **b.** Strains can be classified into

three types by scoring the presence/absence of a synthetase cluster and counting the number receptors

in each genome: (i) single-receptor producers containing one pyoverdine synthetase cluster and one

FpvA receptor gene; (ii) multi-receptor producers containing one pyoverdine synthetase cluster and

several FpvA receptor genes; and (iii) Non-producers lacking synthetase gene but containing at least one

receptor gene. **c.** Heatmap visualizing distances between feature sequences of the FpvA receptors and

the pyoverdine synthetase clusters and between FpvA features sequences and phylogenetic genes

among the 986 single-receptor producers. The hierarchical clustering of the strains is based on the FpvA

feature sequences for all three heatmaps. The black squares on the heatmaps denote the five major

FpvA groups. Three of these groups correspond to the receptors found among *P. aeruginosa* strains and

are labelled with black text. **d.** Scheme of our approach to predict lock-key interactions between

pyoverdines and receptors from sequence data. S-labels refer to the respective results section of our

work.

Based on these findings, we hypothesize that in each single-receptor producer, the sole

receptor should recognize its self-produced pyoverdine. Consequently, when the synthetase

structure is altered, the receptor should correspondingly change sequences to preserve their

matching relationship. This implies that synthetase and receptor pairs engage in molecular

co-evolution. To test this hypothesis, we focused on the 986 single-receptor producers and

calculated the degree of covariation between sequence distances matrices of the receptor, the

synthesis cluster, and the 400 conserved phylogenetic genes. For the receptors (FpvA) and

the synthesis cluster, we used the feature sequences that are most predictive of receptor

specificity and pyoverdine molecular structure[33].

In support of the co-evolution hypothesis, we found a strong correlation between the

distance matrixes of the receptors and the synthesis clusters (Pearson's r=0.89), a correlation

that is much stronger than between the receptor and the phylogeny matrix (Pearson's r=0.51)

(Figure 1c). Notably, we observed strong clustering patterns in the sequence space of the

receptors, forming distinctive blocks that closely match the clustering patterns of their

corresponding synthesis clusters. Using our receptor clustering pipeline[33], we identified 17

receptor groups among the 986 single-receptor producers. Importantly, three out of the 17

receptor groups represent the FpvA receptors characteristic of the human pathogen *P.*

aeruginosa (text marked in Figure 1c left panel, as type I-IV FpvAs), and their associated

synthetase groups were known to produce pyoverdines that these receptors could selectively

uptake[24,35]. These analyses support the hypothesis that cognate receptors and synthesis

genes have co-evolved in single-receptor producers, resulting in one-to-one "lock-key"

relationships: Each group of receptors, characterized by similar sequence features, acts as a

"lock" specifically recognizes pyoverdines (the "key") produced a corresponding group of

synthetases.

To be able to reconstruct the iron interaction network in *Pseudomonas* communities, we

now need to uncover the lock-key groups in multi-receptor producers and to match the

receptors of any strain to the synthetase groups of all producers in a community. To achieve

this, we followed a five-step approach, which is summarized in Figure 1d. In section 2, we

systematically studied the evolution and diversification of the pyoverdine synthesis clusters

and receptors in 1928 *Pseudomonas* strains. In section 3, we used this knowledge to develop

an unsupervised algorithm based on coevolutionary principles, in order to allocate synthesis

clusters and receptors into lock-key groups. This analysis yields a complete prediction map

regarding the groups of pyoverdines each strain is capable of producing and utilizing. In

section 4, we use two experimental systems to validate the predicted siderophore-mediated

interactions in model *Pseudomonas* communities. In section 5, we finally construct the

predicted pyoverdine-mediated interaction networks for soil, plant, water, and

human-associated *Pseudomonas* communities.

Section 2: Concurrent diversification patterns of pyoverdine synthetases and receptors
reveal multi-receptor producers as evolutionary hubs.

We observed that the structural diversity of pyoverdines was much higher when strains carried

multiple receptors (108 unique structures among 678 strains) compared to when they carried a

single receptor (47 unique structures among 986 strains). Only 33 structures were predicted to

be produced by both receptor type strains. This indicates that pyoverdine diversity is tightly

linked with the apparently different iron-acquisition strategies of multi-receptor and

single-receptor producers. Here, we explore this diversity in detail and ask how do pyoverdine

NRPSs evolve. The 188 pyoverdine structures varied in their length, chirality, and substrate

composition, agreeing with the notion that recombination dominates NRPS evolution [36]. We

developed the "NRPS tracer" algorithm, which calculates the minimal number of editing events

required to transform one pyoverdine product into another. Our algorithm builds on the double

Figure 2 Diversifications of pyoverdine synthetases and receptors reveal multi-receptor producers as evolutionary hubs. a. Network of all the 188 known pyoverdine

structures connected by one-step evolutionary events. Stripes of colored squares show the amino-acid

sequence of pyoverdines with black-framed boxes representing D-type amino acids. Colored circles

(nodes) indicate the different FpvA receptor groups found among single-receptor producers. Empty

nodes depict pyoverdines exclusively found among multi-receptor producers. The bold orange line shows

a representative path of pyoverdine diversification. The positioning of pyoverdine stripes and nodes has

no specific meaning. The lower right panel show the scheme of six different recombination events in the

evolution of pyoverdine NRPS clusters. Grey and colored boxes indicate conserved and recombination

events, respectively. Black-framed boxes indicate recombination involving E domain alterations (e.g.

affecting L- vs. D-conformation). **b.** Frequency of edge types and recombination events in the network

(right panel). There are four edge types (left panel): ET1 – edges connecting pyoverdine structures from

single-receptor producers sharing the same receptor group, ET2 – edges connecting pyoverdine

structures from single-receptor producers featuring different receptor groups, ET3 – edges connecting

pyoverdine structures that only appeared in multi-receptor producers and these appeared at least one

time in single-receptor producers, ET4 – edges connecting pyoverdine structures exclusively occurring in

multi-receptor producers. **c.** The 43 largest FpvA receptor groups with more than 10 members (sorted by

group size) and their frequency among single-receptor producers, multi-receptor producers, and

non-producers. **d.** Silhouette index analysis on the compactness of all receptor groups in single-receptor

producers (left panel), multi-receptor producers (middle panel) and non-producers (right panel). Colors

represent all the 17 receptor groups found among single-receptor producers. All other receptor groups

are shown in black. The dashed vertical lines represent the average of the Silhouette index across all the

receptor groups within each strain class. **e.** The observed and expected connectivity value between

different strains types in the sequence similarity network of all 4547 FpvAs receptors (Figure S4).

To keep analysis tractable, we focused on pairs of pyoverdine structures that can

transform from one to the other by a single recombination event and for which the number of

NRPS modules involved in the transformation is ≤ 3 (corresponding to the maximum number

of amino acids a NRPS enzyme can incorporate into the pyoverdine backbone). We identified

216 such single-event transformations. They connect 148 of the 188 pyoverdine structures into

multiple sub-networks (Figure 2a). While many sub-networks are small the five largest connect

70 out of 188 structures and 1016 out of 1664 strains (Figure S1). The most frequent genetic

recombination events were domain and subdomain symmetry replacement (40%), insertions

and deletions (indel, 35%) and asymmetry replacement (16%), while duplication events,

E-domain indels and rearrangements (total 9%) were much rarer (Figure 2a, the lower right

panel).

We then asked whether pyoverdine structures from the same sub-network tend to share

similar receptors. To address this question, we allocated nodes to each structure and filled

them with the colors of the corresponding receptor groups found in single-receptor producers

(17 receptor groups, Figure 1c). Thus, we had 80 colored nodes for the 47 structures

exclusively found in single-receptor producers and the 33 structures occurring in both producer

types. The 108 nodes of structures that only occurred in multi-receptor producers were left

empty (Figure 2a). We then counted the number of cases in which two nodes were connected

by the same vs. different colored receptors (Figure 2b, split according to recombination type).

We found that the great majority of edges connecting nodes of single-receptor producers were

of the same color (97%), while different-color connections were rare (3%). This result strongly

supports the molecular co-evolution hypothesis. By contrast, structures that only occur in

multi-receptor producers (empty node) were as often connected to colored nodes (57 events)

than to other empty nodes (54 events). This indicates that multi-receptor producers can

connect sub-networks of different structures from single-receptor producers (Figure 2a, bold

orange line and Figure S2). Altogether, our analysis reveals that multi-receptor producers

follow different iron-uptake strategies and have undertaken different evolutionary trajectories in

terms of pyoverdine diversification (Figure 2). The latter implies that multi-receptor producers

(and probably also non-producers) should also differ in the FpvA receptors they possess, a

question we address in the subsequent section.

The previously identified 4547 FpvAs genes cluster into 94 distinct groups based on their

feature sequences. For our analysis, we focus on the 43 largest groups comprising more than

10 members (Figure 2c). We found that FpvA receptors of the three strain types were

unequally distributed across the 43 groups (Figure 2c). Receptors of single-receptor producers

were restricted to 14 out of the 43 groups (32.6%). Conversely, receptors of multi-receptor

producers occurred in almost all groups (95.3%). Similarly, receptors from non-producers were

also found in many groups (83.7%) with two distinct groups containing exclusively

non-producer receptors. This simple frequency analysis shows that lock-key relationships from

single-receptor producers are not sufficient to recover the whole iron interaction network

because multi-receptor producers and non-producers possess a much more diverse FpvA

receptor repertoire than single-receptor producers.

This notion receives further support when comparing the compactness of receptor groups

across the three strain types (Figure 2d). We observed that receptors from single-receptor

producers tend to connect more compactly (mean silhouette index = 0.96±0.16), while

receptors from multi-receptor producers (mean silhouette index = 0.78±0.19) and

non-producers (mean silhouette index = 0.79±0.20) were more dispersed in the sequence

space (Figure S3). This observation indicates that receptors from single-receptor producers

are more conserved, whereas receptors from multi-receptor producers harbor greater variation.

The distinct and conserved nature of receptors from single-receptor producers also emerges

when conducting a network analysis with all 4547 FpvAs using their feature sequences (Figure

S4). When focusing on the shortest distances between receptor groups, we noticed that none

of the 17 receptor groups from single-receptor producers are connected (Figure 2e). In

contrast, connections are disproportionately enriched among receptor groups of multi-receptor

producers and non-producers, covering 94% and 50% of all the detected 2195 shortest

between-group distances, respectively (Figure 2e). These results indicate that evolutionary

trajectories of receptors differ between the three strain types. While single-receptor producers

harbor conserved and evolutionarily distinct receptors, receptors of multi-receptor producers

and non-producers are much more diverse and evolutionarily connected.

Taken together, our results suggest that multi-receptor producers are the main reservoir

for siderophore and receptor diversification. These strain types should thus be able to take up

several pyoverdine types and are thus expected to form denser interaction networks at the

ecological level.

Section 3: Matching synthetases and receptors in sequence space results in 47
lock-key groups.

Next, we aim is to establish a lock-key receptor-pyoverdine interaction map across all three

strain types. A first task in this process is to identify receptors in multi-receptor producers that

are used to take up the self-produced pyoverdine. The first intuitive solution is to check for

receptors proximate to the pyoverdine synthetase. We can refer to single-receptor producers

for the proximity threshold, where 95.6% (943 of 986) of FpvA receptors locate within 20 kb

distance from the pyoverdine synthetase cluster. When applying this proximity threshold to

multi-receptor producers we can identify putative self-receptors in 87.1% (591 out of 678) of

the strains (Solution 1). An alternative approach is to use the lock-key pairs identified for

single-receptor producers and check whether similar pairs occur in multi-receptor strains

(Solution 2). However, as shown in Figure 2c, this approach only works for the 17 receptor

groups found in single-receptor producers and could be applied to 68.7% (466 of 678)

multi-receptor strains. Even when combining the Solutions 1 and 2, more than half of the

receptor groups could not be paired to any pyoverdine synthetase.

We thus developed an unsupervised learning algorithm, termed Co-evolution Pairing

Algorithm (Solution 3), which matches the feature sequence of the synthetase cluster in each

strain with its receptors by searching for the set of synthetase-receptor combinations that

maximizes co-evolutionary association. Among all multi-receptor producers, there are 678

synthetases and 2812 receptors in total. First, considering that NRPS pathways mainly evolve

by large genetic rearrangement like recombination, we used the synthetase feature sequences

(concatenated Amotif4-5 regions with consideration of recombination, See Method for details)

to build the 678x678 synthetase distance matrix (Figure 3a). We then picked a random

receptor as putative self-receptor for each multi-receptor producer and used the receptor

features sequences (168 Pro to 295 Ala) to calculate the corresponding 678x678 receptor

distance matrix. Subsequently, we calculated co-evolution coefficient cr, defined as the

Pearson's correlation coefficient between the two matrices (see Method for details). The initial

random self-receptor assignment resulted in poor co-evolution coefficients. We thus

introduced an iterative optimization process, during which putative self-receptors were shuffled

within each multi-receptor producer. We discarded iterations that decreased *cr* values and

continued with those that increased *cr* values until an optimization plateau was reached

(Figure 3b, *cr* = 0.84). We predicted the self-receptor of all multi-receptor producers based on

the final assignment.

Figure 3 Developing an unsupervised algorithm to identify self-receptor in

multi-receptor producers and establishing the lock-key pairs of synthetase and

receptor subgroups. **a.** The flowchart of the Co-evolution Pairing Algorithm (Solution 3) that

matches the synthetase in each strain to its "self-receptor", by an unsupervised learning scheme that

optimizes co-evolutionary strength between the sequence distance matrices of pyoverdine synthetase

and matched receptors. The mean correlation coefficients (r) between the two matrices before and after

the optimization were shown. **b.** The correlation coefficient (r) and stability of the algorithm were

examined by multiple rounds of learning (exp 1 to 50, with final r recorded in the brackets). **c.** The

consistency of self-receptors identified by different solutions. The consistency is calculated by comparing

the self-receptor found by the multi-receptor producer based on the two supervised solutions (Solution 1

and 2) and one unsupervised algorithm (Solution 3). **d.** Lock-key pairs connecting the sequence spaces

of synthases with their self-receptors in both single-receptor producers and multi-receptor producers. The

colored (use the same color code as Figure 2a) and black lines represent groups with single-receptor

producer and without single-receptor producer, respectively.

We then checked for consistency in self-receptor identification across the three solutions

(Figure 3c). Solution 1 and Solution 2 can be classified as supervised machine learning, and

they yield high levels of consistency (99.5% across 433 strains). The unsupervised Solution 3

also shares high consistency with Solution 1 (93.7%, for 591 strains) and Solution 2 (94.4%,

for 466 strain). These high degrees of consistency legitimate all three solutions, with Solution 3

having the advantage of being applicable to all strains.

With the help of our co-evolution pairing algorithm, we could predict and allocate a

self-receptor to each of the 1664 pyoverdine-producing strains, segregating into 47 distinct

lock-key groups (Figure 3d and Figure S5). All single-receptor producers and 84.4%

multi-receptor producers belong to 17 receptor groups, while the remaining 15.6%

self-receptors found among multi-receptor producers segregate into 30 additional receptor

groups (Figure S5). This leaves us with 2883 FpvA receptors among multi-receptor producers

and non-producers that are not self-receptors and could be used to uptake pyoverdines

produced by other strains, representing cheating. Indeed, when mapping these 2883

non-self-receptors to the co-evolution matrix, we found that 2703 receptors associated with

one of the 47 lock-key groups, suggesting that they can take up the corresponding

non-self-produced pyoverdine. There were only 180 FpvA receptors (4.0%) that could not be

linked to any of the 47 lock-key groups. They belong to rare receptor groups (Figure S5) that

are either erroneously identified as FpvA receptors or match rare pyoverdine structures not

covered by the 1664 producers in our dataset. Taken together, we can now predict

iron-interactions between strains based on the siderophores they produce and the receptors

they possess for siderophore utilization.

Section 4: The receptor-synthetase lock-key pairs successfully predict
pyoverdine-mediated interactions in experiments.

We conducted two validation experiments to show that our bioinformatic lock-key approach

reliably predicts pyoverdine-mediated interactions with accuracy > 90%. The first validation

was conducted with a *Pseudomonas* community from the Nanjing (China) collection originally

isolated from the tomato rhizosphere [31]. We chose the genomes of 24 independent strains and

subjected them to our previously developed bioinformatic pipelines to predict pyoverdine

molecular structures and to identify all FpvA receptors[33]. These 24 strains included 4

single-receptor producers, 16 multi-receptor producers and 4 non-producers (Figure S6). We

applied our Co-evolution Pairing Algorithm to identify the self-receptors of all 20 producers

(Solution 3) and verified that all the predicted self-receptors were within the 20 KB genome

proximity threshold to the synthetase genes (Solution 1). Subsequently, we assigned the

self-receptors to the previously identified 47 lock-key paring groups in our full database, and

found that 13 pairing groups occur among the 20 producer strains (Figure S7). Combining this

information with the knowledge on non-self-receptors occurring in each strain, we predicted

the pyoverdine-mediated interactions between the 24 strains in our community (Figure 4a).

Figure 4 The receptor-synthetase lock-key pairings inferred from sequence data reliably

predict experimentally observed pyoverdine-mediated iron interactions. a. Predicted vs.

observed iron-interaction network among the 24 experimental strains. Each circular node represents an

experimental strain. Green, yellow, and red circular nodes represent single-receptor producers,

multi-receptor producers and non-producers, respectively. Hexagons represent the predicted 13 lock-key

receptor-pyoverdine groups. Edges from strain nodes to lock-key nodes represent pyoverdine production,

while edges from lock-key nodes to strain nodes represent utilization. Green (pyoverdine non-usable)

and pink (pyoverdine usable) edges depict cases in which experimental observations match

bioinformatically predicted interactions. Blue edges depict incorrectly predicted pyoverdine interactions.

The pyoverdine groups that appeared at least once in single-receptor producers are shown as colored

hexagones with the color of the respective receptor group, whereas the pyoverdine groups exclusively

secreted by the multi-receptor producer are represented by grey hexagons. **b-c.** Predicted vs. observed

iron-interaction networks based on data from a previous study carried out in Zurich lab. The predicted

interactions were inferred by the algorithms presented in this study, while the experimental data is taken

from Table S2 of Figueiredo et al. (DOI: 10.1111/ele.13912).

For the experimentally validation, we first confirmed that the 20 producers can indeed

produce pyoverdine under iron-limited conditions, while the 4 non-producers cannot (Figure

S6). We then followed a modified version of our previously established protocols to calculate

the net effect pyoverdine has on the growth of other strains (GE_{Pyo}), while controlling for the

effects of other metabolites in the supernatant [31]. In principle, $GE_{Pyo} > 0$ indicates

pyoverdine-mediated facilitation. However, because there is substantial experimental variation

between experimental replicates, we increased a threshold value of $GE_{Pyo} > 0.05$ and

classified values above this threshold as positive interactions, where the receiving strain can

use the respective pyoverdine for iron acquisition (interaction type 1). Conversely, $GE_{Pyo} \leq$

0.05 values were classified as neutral or negative interactions, where the receiving strain

cannot use the respective pyoverdine for iron acquisition (interaction type 0). This approach

allowed us to infer an experimental pyoverdine-mediated interaction network (Figure 4a and

Figure S8), in which 90% of the observed interactions matched (based on the sign) the

predicted interactions from sequence data.

The second experimental validation involved strains from the Zurich (Switzerland)

collection, isolated from soil and freshwater habitats [25]. In this case, we used published

experimental data from the literature [38]. The focus of this earlier study was to test whether the

opportunistic human pathogen *P. aeruginosa* PAO1 can invade natural soil and pond

communities based on its ability to use pyoverdine from the natural isolates. We used data

from all the strains for which genome sequences were available (PAO1 and 33 natural

isolates), to establish pyoverdine-mediated interaction networks (Figure 4b-c). We then

applied our bioinformatic pipelines as explained for the Nanjing collection and found a high

level of consistency (94%) between the predicted and observed pyoverdine-mediated

interaction in pairwise cultures (Figure 4b-c).

The high consistency between observed and predicted interactions among both the

Nanjing and the Zurich strain collection demonstrates that siderophore-mediated microbial

interactions can be predicted based on genome-sequence analysis alone using the lock-key

relationship between receptor and synthetase genes.

Section 5: Pyoverdine interaction networks differ across habitats

We then applied the lock-key pairing approach to our full data set to reconstruct the

pyoverdine-interaction network among all the 1928 *Pseudomonas* strains (Figure S9). To keep

traceability in such an enormous network, we allocated strains into behavioral groups sharing

the same "pyoverdine-interaction behavior". Specifically, strains that produce the same

pyoverdine type and can utilize the same repertoire of pyoverdines are allocated to the same

behavior group and represented as (circular) nodes in the network. There were 407 such

behavioral nodes. We included a second type of (hexagonal) nodes in the network and they

represent the 47 different receptor-pyoverdine groups identified based on the lock-key

algorithm (Figure 3). Edges from behavior nodes to lock-key nodes represent pyoverdine

production, while edges from lock-key nodes to behavior nodes represent utilization. Overall,

the network featured 307 production edges and 1788 utilization edges. This network analysis

does not only reveal that certain lock-key nodes are much more densely connected to

behavior nodes than others, it also reveals a high connectivity between behavior groups and

the widespread ability of strains to use non-self-produced pyoverdines, indicative of potential

cheating.

Figure 5 *Pseudomonas* iron-interaction networks vary across habitats. The predicted

iron-interaction networks mediated by pyoverdines among *Pseudomonas* isolates from soil (**a**), plant (**b**),

water (**c**) and human (**d**) habitats. The interaction networks were built using the Cytoscape's yFiles

circular layout. Circular nodes represent behavioral groups (i.e., strains that produce the same

pyoverdine type and utilize the same repertoire of pyoverdines) with node size being proportional to the

number of strains that exhibit this behavior. Green, yellow, and red circular nodes represent

single-receptor producers, multi-receptor producers and non-producers, respectively. Hexagonal nodes

represent the lock-key receptor-pyoverdine groups with node size being proportional to the number of

strains utilizing this siderophore. The pyoverdine groups once appeared in single-receptor producers

were highlighted with the receptors' group colors (colored hexagons) and the pyoverdine groups only

secreted by the multi-receptor producer in the database represented by grey hexagons. Edges from

behavior nodes to lock-key nodes represent pyoverdine production, while edges from lock-key nodes to

behavior nodes represent utilization, with the same color of the pyoverdine node.

One biological limitation of the above analysis is that most of the strains would never meet

in nature as they were isolated from different environments. That is why we created separate

networks for strains isolated from soil (262 strains), plant (234), water (124), and

human-derived (409) habitats. We found that strain type frequencies varied fundamentally

between the four habitats (Figure 5). For example, in the soil-derived strains, there were 56.9%

multi-receptor producers, 27.5% single-receptor producers and 15.7% non-producers (Table

S1). In contrast, there were only 10.0% multi-receptor producers and 4.0% non-producers, but

86.1% single-receptor producers among human-derived strains. These differences in strategy

frequencies affected network topology and connectivity. Particularly, the number of behavior

groups was higher for soil (130, 0.50 = scaled relative to number of strains), plant (97, 0.41),

and water (70, 0.56) habitats than for human-related habitats (41, 0.10). Similarly, the

distribution of the 47 pyoverdine lock-key groups differed across habitats, with more groups

occurring in soil (29, 0.11 = scaled relative to number of strains), plant (25, 0.11), and water (25,

0.20) habitats compared human-derived habitats (18, 0.04). These differences affected the

number of utilization edges, which is higher in soil (446, 0.25= scaled relative to 1788

utilization edges), plant (274, 0.15) and water (237, 0.13) habitats than in human-derived

habitats (88, 0.05). Important to note is also that many behavior groups are unique in each

habitat: soil (80, 19.7%), plant (56, 13.0%), water (43, 10.6%), and human (26, 6.4%), and that

only 8 behavior groups (Figure S10a) and 11 lock-key groups are shared by all four habitats

(Figure S10b). This latter finding suggests divergent evolution of pyoverdine-interaction

networks across habitats. Taken together, these results show that our algorithms can predict

pyoverdine-interaction networks in natural communities and reveal key differences between

habitats.

Discussion

Predicting interactions between microbes from sequence data offers exciting opportunities for

understanding the ecology and evolution of microbiomes. While sequence-to-interaction

mapping has predominantly been carried out for primary metabolism involving resource

consumption, conversion, and cross-feeding, there are few approaches to reconstruct

microbial interactions based on secondary metabolites (antibiotics, toxins, siderophores,

surfactants) [39-41]. In our paper, we offer such an approach by developing a bioinformatic

approach to infer iron-interaction networks mediated by pyoverdines (a class of siderophores)

within communities of *Pseudomonas* bacteria. To achieve our goal, we analyzed patterns of

pyoverdine and pyoverdine-receptor evolution and their co-evolution from sequence data for

1928 strains and reconstructed strain interaction networks for soil, plant, water, and

human-derived habitats. Our experimentally validated approach provides a roadmap on how

to perform sequence-to-interaction mapping for secondary metabolism, and to obtain new

insights on the effects of these compounds on community assembly and dynamics.

Our approach underscores several challenges associated with secondary metabolites, to

which we offer solutions. The first challenge is that secondary metabolites are often built

through complicated synthesis machineries like NRPS and PKS, such that the chemical

structure of the metabolite can hardly be directly inferred from sequencing data. We solved this

challenge in our first paper[33], in which we developed an approach based on feature sequences

that allowed us to infer the chemical structure of 188 pyoverdines produced by the strains in

our data set. The second challenge is to identify pyoverdine receptor genes among the many

different types of siderophore receptor genes each strain possesses. We also solved this

challenge in our first paper[33], by again using feature sequences to identify 4547 pyoverdine

receptor genes belonging to 94 groups. The third challenge is to pair pyoverdines to matching

receptors within and across strains. For this purpose, we developed an unsupervised learning

algorithm (called Co-evolution Paring Algorithm), which yielded 47 synthetase-receptor

lock-key pairs and allowed us to reconstruct iron-interaction networks.

Although co-evolution analyses are a widely used computational tool, employed in diverse

areas ranging from ab initio protein structure to host-pathogen interaction predictions [42], we

could not use existing algorithms, such as DCA, SCA, and Evoformer [43-45]. The reason is that

these classical site-based co-evolution methods depend on paired sequences between which

the degree of covariation is quantified. We faced two levels of complexity in this context. First,

the existence of multi-receptor producer strains impeded direct assignments of

synthetase-receptor pairs, and a major part of our algorithm is therefore devoted to the

identification of the correct receptor for the self-produced pyoverdine of a strain. Second, due

to the variation in the number of pyoverdine synthetase modules, site-based covariation

quantification is not applicable, and whole-sequence alignment also performs poorly in

characterizing co-evolution due the complexity of pyoverdine synthesis. Therefore, we had to

define reasonable distance metrics that capture the signatures of co-evolution and then apply

the Co-evolution Pairing Algorithm to maximize coevolutionary strength between synthetases

and receptors. Our new pipeline has the potential to be applied to many other microbial traits.

For example, microbial membrane receptors co-evolve with phages[46,47], and pairing phages

with the receptor they utilize for infection could provide insights into host-pathogen

co-evolution.

While our work provides a novel framework for efficient sequence-based prediction of

siderophore-mediated microbial interactions, there are a number of limitations that require

careful consideration. First, our approach assumes that sequence similarity indicates

functional similarity, i.e., similar gene sequences can be put into "behavioral groups" that

produce and uptake the same pyoverdine. In our first paper, we show that this is a reasonable

assumption when feature (but not when full) sequences are used. Second, we assume

discrete lock-key relationships between paired groups of pyoverdine synthetases and

receptors. While it is certainly true that self-receptors have high affinity for the pyoverdine the

strain produces, it is also known that receptors can be promiscuous and take up other

pyoverdine types although at lower efficiencies[35]. Receptor promiscuity is not covered by our

approach. Third, we assume that each producer strain must have at least one self-receptor

that recognizes its own pyoverdine. This assumption is reasonable and empirically well

documented[48]. Fourth, our approach focusses entirely on the presence/absence of pyoverdine

synthetases and receptors, and does not consider regulation. This comes with two

shortcomings that affect our prediction accuracy: (i) there are strains that have an intact

pyoverdine synthetase machinery but are functionally non-producers[24]. These non-producers

are not detected by our approach, which can lead to errors in interaction predictions. (ii)

Strains can vary considerably in the amount of pyoverdine they produce with production levels

influencing the strength of interactions, a parameter that cannot be measured with our

approach. However, despite these limitations our empirical verification experiments reveal an

accuracy of 90% and 94% of correctly predicted interactions.

While the primary goal of our work was to establish methods for sequence-to-interaction

mapping, our results already yielded several new biological insights on iron-interaction

networks in pseudomonads. First, we identified three different iron-acquisition strategies

(single-receptor producers, multi-receptor producers, non-producers) that can co-exist and

appear to follow different evolutionary trajectories. Specifically, single-receptor producers

showed relatively low structural diversity in pyoverdines and their receptor groups, suggesting

relatively conserved iron-uptake strategies. In contrast, pyoverdine and receptor diversity is

much higher among multi-receptor producers, identifying them as the evolutionary hub for both

pyoverdine and receptor diversification. Second, we found that the above findings have direct

consequences for iron-network topologies because multi-strain producers (featuring high

pyoverdine and receptor diversity) can connect sub-networks into larger interaction networks.

Indeed, soil-, plant-, and water-associated habitats, dominated by multi-receptor producers,

featured highly connected iron-interaction networks. Conversely, iron-interaction networks

were small and fragmented in human-derived habitats dominated by single-receptor producers.

This latter finding potentially stems from a (relatively recent) ecological expansion of *P.*

aeruginosa from environmental habitats to humans and other hosts. This ecological expansion

could have facilitated diversification, as evidenced by the diverse array of *P. aeruginosa* strains

in our data set. However, it appears that pyoverdine synthetase and receptors were little

affected by the diversification and remained conserved.

In conclusion, we succeeded to develop a sequence-to-interaction mapping approach for

siderophores that has high potential to deliver new insights into species and strain interaction

networks in bacterial communities. Given that iron is a key trace element that is limited in most

environments, siderophore-mediated interactions are an ideal entry point for secondary

metabolite analysis from sequence data. While we focused on *Pseudomonas* strains, we know

that siderophore-mediated interactions occur across the species boundaries. For example. *P.*

aeruginosa possesses receptors to take up enterobactin produced by *Enterobacteriaceae* spp.

and schizokinen produced by. *Ralstonia solanacearum* [49]. Thus, the next step would be to

apply our concepts to more diverse bacterial communities to derive microbiome-level

iron-interaction maps. Moreover, pyoverdines represent a particularly complex group of

secondary metabolites, such that our evolution-guided approach should be easily

customizable to simpler groups of secondary metabolites.

Data Availability

The source code and parameters used are available in the supplementary material.

Acknowledgements

We thank Xuedi Huang for the insight on pyoverdine receptor analysis.

Funding

This work was supported by the National Key Research and Development Program of China

(No. 2021YFF1200500, 2021YFA0910700), National Natural Science Foundation of China

(No. 42107140, No. 32071255, No.41922053), National Postdoctoral Program for Innovative

Talents (No. BX2021012). R.K. is supported jointly by a grant from the Swiss National Science

Foundation no. 310030_212266. V-P.F. is supported jointly by a grant from UKRI, Defra, and

the Scottish Government, under the Strategic Priorities Fund Plant Bacterial Diseases

program (BB/T010606/1), Research Council of Finland, and The Finnish Research Impact

Foundation.

Author contributions

Shaohua Gu performed the majority of computational analysis in this research and drafted the

manuscript. Zhengying Shao and Shenyue Zhu performed the experiment of testing

pyoverdine-mediated interaction between *Pseudomonas* strains. Yuanzhe Shao built the

NRPStracer pipeline to infer patterns of diversification in pyoverdine synthetase. Di Zhang

developed unsupervised co-evolution pairing algorithm for identifying self-receptor. Ruolin He,

Jiqi Shao, Guanyue Xiong, Zeyang Qu assisted in cleaning up the codes. Alexandre Jousset

and Ville-Petri Friman offered insightful comments and assisted in revising and writing of the

manuscript. Rolf Kümmerli and Zhong Wei oversaw the project, designed experiments and

revised the manuscript. Zhiyuan Li conceptualized, oversaw the project and revised the

manuscript.

Competing interests

The authors declare no competing interests.

References

1   Cavicchioli, R. *et al.* Scientists' warning to humanity: microorganisms and climate

598   change. *Nature Reviews Microbiology* **17**, 569-586, doi:10.1038/s41579-019-0222-5

599   (2019).

600   2   Rani, N., Sangwan, P., Joshi, M., Sagar, A. & Bala, K. in *Microbial Wastewater*

601   *Treatment*   (eds Maulin P. Shah & Susana Rodriguez-Couto)   83-102 (Elsevier,

602   2019).

603   3   Wang, X. *et al.* Phage combination therapies for bacterial wilt disease in tomato.

604   *Nature Biotechnology* **37**, 1513-1520, doi:10.1038/s41587-019-0328-3 (2019).

605   4   Kuramitsu Howard, K., He, X., Lux, R., Anderson Maxwell, H. & Shi, W. Interspecies

606   Interactions within Oral Microbial Communities. *Microbiology and Molecular Biology*

607   *Reviews* **71**, 653-670, doi:10.1128/MMBR.00024-07 (2007).

608   5   Konopka, A., Lindemann, S. & Fredrickson, J. Dynamics in microbial communities:

609   unraveling mechanisms to identify principles. *The ISME Journal* **9**, 1488-1495,

610   doi:10.1038/ismej.2014.251 (2015).

611   6   Handelsman, J. Metagenomics: Application of Genomics to Uncultured

612   Microorganisms. *Microbiology and Molecular Biology Reviews* **68**, 669-685,

613   doi:10.1128/MMBR.68.4.669-685.2004 (2004).

614   7   Almeida, A. *et al.* A new genomic blueprint of the human gut microbiota. *Nature* **568**,

615   499-504, doi:10.1038/s41586-019-0965-1 (2019).

616   8   Schloss, P. D. & Handelsman, J. Metagenomics for studying unculturable

617   microorganisms: cutting   the   Gordian   knot.   *Genome   Biology*   **6**,   229,

doi:10.1186/gb-2005-6-8-229 (2005).

9   Gu, C., Kim, G. B., Kim, W. J., Kim, H. U. & Lee, S. Y. Current status and applications

of genome-scale metabolic models. *Genome Biology* **20**, 121,

doi:10.1186/s13059-019-1730-3 (2019).

10   Colarusso, A. V., Goodchild-Michelman, I., Rayle, M. & Zomorrodi, A. R.

Computational modeling of metabolism in microbial communities on a genome-scale.

Current Opinion in Systems Biology **26**, 46-57, doi:10.1016/j.coisb.2021.04.001

(2021).

11   Pacheco, A. R., Moel, M. & Segrè, D. Costless metabolic secretions as drivers of

interspecies interactions in microbial ecosystems. *Nature Communications* **10**, 103,

doi:10.1038/s41467-018-07946-9 (2019).

12   Gude, S. & Taga, M. E. Multi-faceted approaches to discovering and predicting

microbial nutritional interactions. *Current Opinion in Biotechnology* **62**, 58-64,

doi:10.1016/j.copbio.2019.08.005 (2020).

13   Louca, S. & Doebeli, M. Calibration and analysis of genome-based models for

microbial ecology. *eLife* **4**, e08208, doi:10.7554/eLife.08208 (2015).

14   Penn, K. *et al.* Genomic islands link secondary metabolism to functional adaptation in

marine Actinobacteria. *The ISME Journal* **3**, 1193-1203, doi:10.1038/ismej.2009.58

(2009).

15   Scherlach, K. & Hertweck, C. Mediators of mutualistic microbe–microbe interactions.

Natural Product Reports **35**, 303-308, doi:10.1039/C7NP00035A (2018).

16   Netzker, T. *et al.* Microbial communication leading to the activation of silent fungal

secondary metabolite gene clusters. *Front Microbiol* **6**, 299, doi:10.3389/fmicb.2015.00299 (2015).

Demain, A. L. & Fang, A. The natural functions of secondary metabolites. *Adv Biochem Eng Biotechnol* **69**, 1-39, doi:10.1007/3-540-44964-7_1 (2000).

Gao, H., Weitao, T. & He, Q. Coping with the environment: how microbes survive environmental challenges. *Int J Microbiol* **2011**, 379519, doi:10.1155/2011/379519 (2011).

Wolf, D. M., Vazirani, V. V. & Arkin, A. P. Diversity in times of adversity: probabilistic strategies in microbial survival games. *Journal of Theoretical Biology* **234**, 227-253, doi:10.1016/j.jtbi.2004.11.020 (2005).

He, R. *et al.* Knowledge-guided data mining on the standardized architecture of NRPS: Subtypes, novel motifs, and sequence entanglements. *PLOS Computational Biology* **19**, e1011100, doi:10.1371/journal.pcbi.1011100 (2023).

Andrews, S. C., Robinson, A. K. & Rodríguez-Quiñones, F. Bacterial iron homeostasis. *FEMS Microbiology Reviews* **27**, 215–237, doi:10.1016/s0168-6445(03)00055-x (2003).

Boyd, P. W. & Ellwood, M. J. The biogeochemical cycle of iron in the ocean. *Nat. Geosci.* **3**, 675-682, doi:10.1038/ngeo964 (2010).

Emerson, D., Roden, E. & Twining, B. The microbial ferrous wheel: iron cycling in terrestrial, freshwater, and marine environments. *Frontiers in Microbiology* **3**, doi:10.3389/fmicb.2012.00383 (2012).

Butaitė, E., Baumgartner, M., Wyder, S. & Kümmerli, R. Siderophore cheating and

cheating resistance shape competition for iron in soil and freshwater Pseudomonas

communities. *Nature Communications* **8**, 414, doi:10.1038/s41467-017-00509-4

(2017).

25 Kramer, J., Özkaya, Ö. & Kümmerli, R. Bacterial siderophores in community and host

interactions. *Nature Reviews Microbiology* **18**, 152-163,

doi:10.1038/s41579-019-0284-4 (2020).

26 Leventhal, G. E., Ackermann, M. & Schiessl, K. T. Why microbes secrete molecules to

modify their environment: the case of iron-chelating siderophores. *Journal of The*

Royal Society Interface **16**, 20180674, doi:10.1098/rsif.2018.0674 (2019).

27 West, S. A., Diggle, S. P., Buckling, A., Gardner, A. & Griffin, A. S. The Social Lives of

Microbes. *Annual Review of Ecology, Evolution, and Systematics* **38**, 53-77,

doi:10.1146/annurev.ecolsys.38.091206.095740 (2007).

28 Miethke, M. & Marahiel Mohamed, A. Siderophore-Based Iron Acquisition and

Pathogen Control. *Microbiology and Molecular Biology Reviews* **71**, 413-451,

doi:10.1128/MMBR.00012-07 (2007).

29 Seyoum, Y., Baye, K. & Humblot, C. Iron homeostasis in host and gut bacteria – a

complex interrelationship. *Gut Microbes* **13**, 1874855,

doi:10.1080/19490976.2021.1874855 (2021).

30 Schaible, U. E. & Kaufmann, S. H. E. Iron and microbial infection. *Nature Reviews*

Microbiology **2**, 946-953, doi:10.1038/nrmicro1046 (2004).

31 Gu, S. *et al.* Competition for iron drives phytopathogen control by natural rhizosphere

microbiomes. *Nature Microbiology* **5**, 1002–1010, doi:10.1038/s41564-020-0719-8

(2020).

32   Gu, S. *et al.* Siderophore-Mediated Interactions Determine the Disease

Suppressiveness of Microbial Consortia. *mSystems* **5**, e00811-00819,

doi:10.1128/mSystems.00811-19 (2020).

33   Shaohua, G. *et al.* From sequence to molecules: Feature sequence-based genome

mining uncovers the hidden diversity of bacterial siderophore pathways. *bioRxiv*,

2023.2010.2030.564663, doi:10.1101/2023.10.30.564663 (2023).

34   Bozhüyük, K. A. J. *et al.* De novo design and engineering of non-ribosomal peptide

synthetases. *Nature Chemistry* **10**, 275-281, doi:10.1038/nchem.2890 (2018).

35   Meyer, J. M. *et al.* Use of siderophores to type pseudomonads: the three

Pseudomonas aeruginosa pyoverdine systems. *Microbiology (Reading)* **143 (Pt 1)**,

35-43, doi:10.1099/00221287-143-1-35 (1997).

36   Baunach, M., Chowdhury, S., Stallforth, P. & Dittmann, E. The Landscape of

Recombination Events That Create Nonribosomal Peptide Diversity. *Mol Biol Evol* **38**,

2116-2130, doi:10.1093/molbev/msab015 (2021).

37   Bohnenkämper, L., Braga, M. D. V., Doerr, D. & Stoye, J. Computing the

Rearrangement Distance of Natural Genomes. *Journal of Computational Biology* **28**,

410-431, doi:10.1089/cmb.2020.0434 (2020).

38   Figueiredo, A. R. T., Özkaya, Ö., Kümmerli, R. & Kramer, J. Siderophores drive

invasion dynamics in bacterial communities through their dual role as public good

versus public bad. *Ecology Letters* **25**, 138-150, doi:10.1111/ele.13912 (2022).

39   Tsilimigras, M. C. B. & Fodor, A. A. Compositional data analysis of the microbiome:

706  fundamentals, tools, and challenges. *Annals of Epidemiology* **26**, 330-335,

707  doi:10.1016/j.annepidem.2016.03.002 (2016).

708 40 Weiss, S. *et al.* Correlation detection strategies in microbial data sets vary widely in

709  sensitivity and precision. *ISME J* **10**, 1669-1681, doi:10.1038/ismej.2015.235 (2016).

710 41 Faust, K. & Raes, J. Microbial interactions: from networks to models. *Nature Reviews*

711  *Microbiology* **10**, 538-550, doi:10.1038/nrmicro2832 (2012).

712 42 Chen, H. M., Guo, W., Shen, J., Wang, L. & Song, J. N. Structural Principles Analysis

713  of Host-Pathogen Protein-Protein Interactions: A Structural Bioinformatics Survey.

714  *Ieee Access* **6**, 11760-11771, doi:10.1109/Access.2018.2807881 (2018).

715 43 Greensmith, J., Feyereisl, J. & Aickelin, U. The DCA: SOMe comparison. *Evolutionary*

716  *Intelligence* **1**, 85-112, doi:10.1007/s12065-008-0008-6 (2008).

717 44 Mirjalili, S. SCA: A Sine Cosine Algorithm for solving optimization problems.

718  *Knowledge-Based Systems* **96**, 120-133, doi:10.1016/j.knosys.2015.12.022 (2016).

719 45 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*

720  **596**, 583-589, doi:10.1038/s41586-021-03819-2 (2021).

721 46 Hampton, H. G., Watson, B. N. J. & Fineran, P. C. The arms race between bacteria

722  and their phage foes. *Nature* **577**, 327-336, doi:10.1038/s41586-019-1894-8 (2020).

723 47 Kortright, K. E., Chan, B. K., Koff, J. L. & Turner, P. E. Phage Therapy: A Renewed

724  Approach to Combat Antibiotic-Resistant Bacteria. *Cell Host & Microbe* **25**, 219-232,

725  doi:10.1016/j.chom.2019.01.014 (2019).

726 48 Bodilis, J. *et al.* Distribution and evolution of ferripyoverdine receptors in

727  Pseudomonas aeruginosa. *Environ Microbiol* **11**, 2123-2135,

728   doi:10.1111/j.1462-2920.2009.01932.x (2009).

729   49   Kümmerli, R. Iron acquisition strategies in pseudomonads: mechanisms, ecology, and

730   evolution. *Biometals* **36**, 777-797, doi:10.1007/s10534-022-00480-8 (2023).

731