Siderophore synthetase-receptor gene coevolution reveals habitat- and

pathogen-specific bacterial iron interaction networks

Shaohua Gu[1,2,3], Zhengying Shao[3], Zeyang Qu[1], Shenyue Zhu[3], Yuanzhe Shao[2], Di Zhang[1],

Richard Allen[4], Ruolin He[1], Jiqi Shao[1], Guanyue Xiong[1], Alexandre Jousset[3], Ville-Petri

Friman[5], Zhong Wei[3#], Rolf Kümmerli[4#], Zhiyuan Li[1,2#]

[1] Center for Quantitative Biology, Academy for Advanced Interdisciplinary Studies, Peking

University, Beijing, 100871, China

[2] Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies,

Peking University, Beijing, 100871, China

[3] Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, Key lab of organic-based

fertilizers of China, Nanjing Agricultural University, Nanjing, P R China

[4] University of Zurich, Department of Quantitative Biomedicine, Winterthurerstr. 190, 8057

Zurich, Switzerland

[5] University of Helsinki, Department of Microbiology, 00014, Helsinki, Finland

[#]Corresponding authors (email: weizhong@njau.edu.cn; rolf.kuemmerli@uzh.ch;

zhiyuanli@pku.edu.cn)

Abstract

Predicting bacterial social interactions from genome sequences is notoriously difficult. Here,

we developed bioinformatic tools to predict whether secreted iron-scavenging siderophores

stimulate or inhibit the growth of community members. Siderophores are chemically diverse

and can be stimulatory or inhibitory depending on whether bacteria possess or lack

corresponding uptake receptors. We focused on 1928 representative *Pseudomonas* genomes

and developed a co-evolution algorithm to match all encoded siderophore synthetases to

corresponding receptor gene groups with >90% accuracy based on experimental validation.

We derived community-level iron interaction networks to show that selection for

siderophore-mediated interactions differs across habitats and lifestyles. Specifically, dense

networks of siderophore sharing and competition were observed among environmental

(soil/water/plant) strains and non-pathogenic species, while only fragmented networks

occurred among human-derived strains and pathogenic species. Altogether, our

sequence-to-ecology approach empowers the analyses of social interactions among

thousands of bacterial strains and uncovers ways for targeted intervention to microbial

communities.

35

Introduction

Microbial communities populate all ecosystems on earth from terrestrial to aquatic environments, impacting human health, agriculture, and industry [1-3]. The dynamics and functioning of these communities are shaped by complex and unexplored interactions between microorganisms [4,5]. As the number of sequenced microbial genomes continues to grow [6,7], there is enormous interest in developing approaches to predict microbial interaction networks based on genomic data. Such efforts are essential to obtain complete insights into community functioning as many microorganisms cannot be cultured in the laboratory [8], while their ecological roles could still be inferred through sequence-to-interaction mapping. Currently, sequence-to-interaction mapping approaches primarily focus on metabolic interactions, with Genome-scale Metabolic Models (GEMs) serving as the primary tool for establishing the pan-reactome of microbial communities [9,10]. These methods infer metabolic reactions from the genome annotation of enzymes, and then reconstruct a flux model to understand how microorganisms take up essential nutrients and release metabolic byproducts into the environment [11-13].

Despite the significance of primary metabolism, there is increasing evidence that also other secreted compounds synthesized through secondary metabolism [14], play a major role in shaping microbial interactions [15,16]. Nearly all microbes actively synthesize and secrete compounds to fulfill a diverse set of functions, including communication, resource scavenging, motility, and attack of and defense against competitors [17]. Many of these secreted compounds were previously considered non-essential for microbial growth in laboratory settings, but have since been shown to be critical for competitiveness in natural environments [15,16]. However,

sequence-to-interaction mapping has rarely been applied to secreted compounds, particularly

because the synthesis and mode of action of secondary metabolites is challenging to predict.

Here, we developed a secondary metabolite sequence-to-interaction approach focusing

on iron-scavenging siderophores, one of the most prevalent and diverse classes of microbial

secondary metabolites [18]. Iron is critical for microbial growth and survival, because of its

importance as catalytic group in enzymes guiding key biological processes such as respiration

and DNA replication [19]. However, the concentration of bioavailable iron is typically below the

required level in most habitats [19-21], and upon iron limitation, nearly all bacteria produce

siderophores that efficiently chelate iron from insoluble environmental stocks [22,23].

Siderophores are typically diffusible and able to chelate iron over a broad physical range [24].

Once iron is bound, the complex is recognized and taken up by specific receptors embedded

into the bacterial cell membrane [23]. Diffusible siderophores mediate several types of social

interactions. They can cooperatively be shared among contributing bacterial strains with

matching receptors for the uptake of the iron-siderophore complex [23,25]. Siderophore can also

be deployed as competitive agents to limit access to iron if co-occurring strains lack matching

receptors [23,26]. Finally, siderophores can be exploited by cheaters bacteria that have receptors

for siderophore uptake but do not pay the cost of producing siderophores themselves [22,23].

Consequently, while siderophore-mediated interactions have important impacts on microbial

community dynamics and functions [27-30], we still poorly understand how these interactions

scale up at the network level in environmental and in host-associated bacterial populations.

The aim of our study was to infer how receptors and siderophores have co-evolved and

use this information to develop algorithms that identify matching siderophore-receptor pairs

that predict interaction networks in bacterial communities based on sequence data. We

previously used the genome sequences of 1928 *Pseudomonas* strains to develop

bioinformatic pipelines that allowed us to predict the chemical structure of 188 pyoverdines

(the main siderophores of this genus) and to identify 4547 FpvA-receptor genes (segregating

into 94 groups) involved in pyoverdine uptake [31]. Here, we capitalize on this work to develop a

Co-evolution Pairing Algorithm to match the pyoverdines (key) and receptors (lock) into 47

unique lock-key groups and over 90% of these predicted interactions could be validated

experimentally *in vivo*. Using the predicted lock-key pairs, we then reconstructed

siderophore-mediated iron interaction networks among all *Pseudomonas* strains. We found

that network complexity was high among strains isolated from soil-, water-, and plant-derived

habitats, whereas complexity was lower among strains isolated from human-associated

habitats. We further noticed that interaction networks among pathogenic species were small

and loose with few pyoverdine interactions existing between strains. The opposite was the

case for strains from environmental habitats. Taken together, the developed

sequence-to-interaction mapping tool can accurately predict social interaction networks

mediated by siderophores in complex bacterial communities. Our findings suggest that

selection for social interactions varies across habitats and lifestyles, thus providing novel

insights into community functions and connectivity.

Results

Three classes of pyoverdine strategies in *Pseudomonas* strains and the lock-key

(receptor-synthetase) principle of co-evolution

Our data set consists of 1928 nonredundant *Pseudomonas* strains, producing a total of 188

chemically different pyoverdine types and featuring 94 different receptor groups according to

our recently developed bioinformatic prediction tools[31]. We first explored the diversity of strains

in terms of phylogeny, ecological habitat, and pyoverdine strategies. At the phylogenetic level,

our data set included a diverse set of *Pseudomonas* species, where *P. aeruginosa* (28.7%), *P.*

fluorescens (7.0%), *P. syringae* (6.0%), and *P. putida* (2.2%) were the most abundant ones

(Figure 1a). The strains originated from diverse habitats, including human-derived habitats

(21.2%), soils (13.6%), plants (12.1%), and water (6.4%), although the origin of many strains

(39.5%) is unknown (Figure 1a).

To assess the diversity of pyoverdine production and uptake strategies, we analyzed the

absence or presence of pyoverdine synthesis clusters and counted the number of FpvA

pyoverdine receptors per strain. We found three types of pyoverdine-utilization strategies

(Figure 1b). "Single-receptor producers" were the most common type (985 strains, 51.1%) and

refer to strains with one pyoverdine synthesis locus and one FpvA receptor gene.

"Multi-receptor producers" were the second most common type (679 strains, 35.2%) and

includes strains with one pyoverdine synthesis cluster but multiple FpvA receptor genes.

"Non-producers" were the least common type (264 strains, 13.7%) and refer to strains that

lack the pyoverdine synthesis cluster but contain at least one receptor gene. While strains can

possess multiple FpvA receptor genes, no strain carried more than one pyoverdine synthesis

cluster. This observation is in line with the expected high costs of pyoverdine synthesis, which

is based on a series of gigantic modular enzymes known as NPRS (non-ribosomal peptide

synthetases)[32].

124
Figure 1 Classification of *Pseudomonas* strains and elucidation of the co-evolution
between pyoverdine synthetases and receptors. a. Phylogenetic relationship among the 1928

Pseudomonas strain based on the concatenated alignment of 400 single-copy conserved genes. Starting
from inside, colors in the first ring distinguish the five most prevalent species, with "Others" representing
the remaining less abundant species. Colors in the second ring distinguish the four most prevalent
sources of isolation. In the third ring, claret and blank regions cover strains with complete pyoverdine
synthetase clusters and strains without synthetase gene clusters, respectively. In the fourth blue ring, the
bar height indicates the number of FpvA receptors present in each strain. **b.** Strains can be classified into
three types by scoring the presence/absence of a synthetase cluster and counting the number receptors
in each genome: (i) single-receptor producers containing one pyoverdine synthetase cluster and one
FpvA receptor gene; (ii) multi-receptor producers containing one pyoverdine synthetase cluster and
several FpvA receptor genes; and (iii) Non-producers lacking synthetase gene but containing at least one
receptor gene. **c.** Heatmap visualizing distances between feature sequences of the FpvA receptors and
the pyoverdine synthetase clusters and between FpvA features sequences and phylogenetic genes
among the 986 single-receptor producers. In all three heatmaps, the hierarchical clustering of the strains
follows the one used for the FpvA feature sequences (left panel). The black squares on the heatmaps
denote the five major FpvA groups. Three of these groups correspond to the receptors found among *P.*
aeruginosa strains and are labelled with black text. **d.** The 43 largest FpvA receptor groups with more
than 10 members (sorted by group size) and their frequency among single-receptor producers,
multi-receptor producers, and non-producers.
145

Based on these findings, we hypothesized that in each single-receptor producer, the sole

receptor present should recognize the self-produced pyoverdine to ensure fitness benefits

when faced with iron limitations. Consequently, synthetase and receptor pairs should reflect

molecular co-evolution, where mutational alterations in the synthetase structure should select

for corresponding changes in the receptor sequences to preserve the lock-key relationship and

efficient iron uptake. To test this hypothesis, we focused on the 986 single-receptor producers

and calculated the degree of covariation between sequence distances matrices of the receptor,

the synthesis cluster, and 400 conserved genes. For the receptors (FpvA) and the synthesis

cluster, we used the feature sequences that are most predictive of receptor specificity and

pyoverdine molecular structure, as identified in our previous work[31]. In support of the

co-evolution hypothesis, we found a strong correlation between the distance matrixes of the

receptors and the synthesis clusters (Pearson's r=0.88), a correlation that is much stronger

than between the receptor and the phylogeny matrix (Pearson's r=0.51) (Figure 1c). Notably,

we observed strong clustering patterns in the sequence space of the receptors, forming

distinctive blocks that closely match with the clustering patterns of their corresponding

pyoverdine synthesis clusters. Using our receptor clustering pipeline[31], we identified 17

receptor groups among the 986 single-receptor producers. Importantly, three out of the 17

receptor groups represent the FpvA receptors found in the human pathogen *P. aeruginosa*

(labelled as GI, GII and GIII+IV in Figure 1c left panel) for which the selective uptake of the

corresponding pyoverdines has been demonstrated[22,33]. These analyses strongly indicate that

cognate receptors and synthesis genes have co-evolved in single-receptor producers,

resulting in one-to-one "lock-key" relationships.

Co-evolutionary lock-key groups cannot directly be inferred for multi-receptor producers

because there is currently no method to distinguish the "self-receptor" responsible for

absorbing the self-produced pyoverdine from the other FpvA receptors responsible for the

uptake of heterologous pyoverdines produced by other strains. Moreover, receptor diversity

seems to be much larger among multi-receptor than among single-receptor producers. Indeed,

when focusing on the 43 (out of the 94) receptor groups with more than 10 members, we found

that single-receptor producers covered only 14 of these groups (32.6%), while multi-receptor

producers had a much more diverse receptor coverage (41 groups, 95.3%) (Figure 1d).

Non-producers had a similarly broad receptor coverage (34 groups, 79.1%). We also found

that single-receptor producers tend to connect more compactly (mean silhouette index =

0.96±0.16) than multi-receptor producers (mean silhouette index = 0.78±0.19) and

non-producers (mean silhouette index = 0.79±0.20) in the sequence space (Figure S1 and S2),

suggesting that receptors from single-receptor producers might evolutionarily be more

conserved than receptors from non-producers and multi-receptor producers. Altogether, these

observations imply that single- and multi-receptor producers take on different roles in iron

interaction networks.

Development of the Co-evolution Pairing Algorithm and its experimental validation to

predict iron interaction networks

The aim of this section is to establish a lock-key receptor-pyoverdine interaction map across all

three strain types. A first task in this process is to identify receptors in multi-receptor producers

that are used to take up the self-produced pyoverdine. A first intuitive approach was to check

for receptors proximate to the pyoverdine synthetase genes (Solution 1), while an alternative

approach was to use the lock-key pairs identified for single-receptor producers and analysis

whether similar pairs occur in multi-receptor strains (Solution 2). Even after completing these

two solutions, more than half of the receptor groups could not be paired with any pyoverdine

synthetases. Specifically, solution 1 identifies putative self-receptors in 87.1% (591 out of 678)

of the multi-receptor producers, while the solution 2 could be applied to only 68.7% (466 of 678)

of multi-receptor strains.

We thus developed an unsupervised learning algorithm, termed "Co-evolution Pairing

Algorithm" (Solution 3), which searches for the set of synthetase-receptor combinations that

maximizes co-evolutionary association based on feature sequence distance between

synthetase and receptor (Figure 2a-b, see Method 4 for details). We then checked for

consistency in self-receptor identification across the three solutions (Figure 2c). Solution 1 and

Solution 2 yield high levels of consistency (99.5% across 433 strains). The unsupervised

Solution 3 shares high consistency with Solution 1 (93.7%, across 591 strains) and Solution 2

(94.4%, across 466 strains), indicating that all three solutions are legitimate, with Solution 3

having the advantage of being applicable to all strains.

Predicted VS Real siderophore-mediated interactions

Figure 2 Unsupervised co-evolutionary algorithm to establish a lock-key pair map for
pyoverdine synthetase and receptor groups and its experimental validation. a. Flowchart
depicting the Co-evolution Pairing Algorithm (Solution 3) to match the synthetase in each strain to its
"self-receptor", based on an unsupervised learning scheme that optimizes co-evolutionary strength
(correlation coefficients) between the feature sequence distance matrices of synthetases and matched
receptors. The mean correlation coefficients (cr) between the two matrices before and after the
optimization process are shown. **b.** The correlation coefficient (cr) and stability of the algorithm were
examined by multiple rounds of learning (exp 1 to 50). **c.** Consistency of identifying the same receptor as
self-receptor across three different solutions. Solution 1: receptor is within 20,000 bp proximity to the
synthetase. Solution 2: the lock-key pairs of single-receptor producers are applied to multi-receptor
producers. Solution 3: unsupervised co-evolution pairing algorithm. **d.** Predicted lock-key pairs
connected in sequence space: 1664 synthases (bottom left) linked to the 4547 receptors (top right).
Arrows depict the 47 lock-key links between synthetase and receptor groups. The colored (use the same
color code as Figure 1c) and black/ grey shaded lines represent groups with single-receptor producers
and without single-receptor producers, respectively. **e.** Predicted vs. observed iron interaction network
among the 24 experimental strains. Each circular node represents an experimental strain and line colors
stand for single-receptor producers (green), multi-receptor producers (yellow) and non-producers (red).
Hexagons represent the predicted 13 lock-key receptor-pyoverdine groups. Edges from strain nodes to
lock-key nodes represent pyoverdine production, while edges from lock-key nodes to strain nodes
represent utilization. Green (non-usable pyoverdine) and pink (usable pyoverdine) edges depict cases in
which experimental observations match bioinformatically predicted interactions. Blue edges depict
incorrectly predicted pyoverdine interactions. The pyoverdine groups that are produced by at least one
single-receptor producer are shown as colored hexagones with the color of the respective receptor group
(Fig. 1c), whereas grey hexagons depict the pyoverdine groups that are exclusively produced by
multi-receptors. **f+g.** Predicted vs. observed iron interaction networks based on data from a previous
study carried out in the Zurich lab. The predicted interactions were inferred by the algorithms presented
in this study, while the experimental data is taken from Table S2 of Figueiredo et al. [34].

Our Co-evolution Pairing Algorithm, allocated the self-receptors of the 1664

pyoverdine-producing strain into 47 distinct lock-key groups based on the receptor feature

distance (Figure 2d and Figure S3). Most self-receptors belonged to 17 lock-key groups

(single-receptor producers: 986 = 100%, multi-receptor producers: 572 = 84.4%), while the

remaining self-receptors of multi-receptor producers (106 = 15.6%) segregated into 30

additional receptor groups (Figure S3). Out of the total 4547 FpvA genes detected, we

identified 2883 receptors that are not self-receptors and thus possibly serve as "cheating

receptors" to take up heterologous pyoverdines produced by other strains. Most of these

cheating receptors (2703 = 93.8%) also segregated into the 47 lock-key groups, confirming

that they could be used to exploit at least one of the 188 produced pyoverdines. The remaining

cheating receptors (180 = 6.2%) could not be linked to any of the 47 lock-key groups,

suggesting the existence of rare receptor groups that presumably match rare pyoverdine

structures not covered by our dataset (Figure S3).

By combining co-evolution pairing algorithm and the lock-key pairing, we predicted

siderophore-mediated iron interaction networks, based on pyoverdines that were produced

and could be taken up by corresponding receptors by community members. We conducted

two experiments to validate predicted interactions. For the first validation, we used a

Pseudomonas community from the Laboratory of rhizosphere microbial ecology (LorMe) in

Nanjing (China), which was originally isolated from the tomato rhizosphere [29]. We included 24

independent strains and subjected their genomes to our bioinformatic pipelines to predict

pyoverdine structures, to find all FpvA receptors[31], to identify self-receptors, and to allocate

pyoverdines and receptors into lock-key groups. We found that these 24 strains included 4

single-receptor producers, 16 multi-receptor producers and 4 non-producers (Figure S4) and

that their self-receptors could be allocated to 13 lock-key groups (Figure S5). With this

information, we predicted the pyoverdine-mediated iron interaction network between the 24

strains (Figure 2e). For the experimentally validation, we confirmed the pyoverdine production

status of the 20 putative producers and the 4 non-producers (Figure S4). We then followed a

modified version of our previously established protocols to calculate the net effect pyoverdine

has on the growth of other strains (GE_{Pyo}), while controlling for the effects of other metabolites

in the supernatant [29]. This approach allowed us to obtain an experimentally derived

pyoverdine-mediated iron interaction network (Figure 2e and Figure S6). We found that 90% of

the observed interactions (whether strains are stimulated or inhibited by the pyoverdines of

others) matched the computationally predicted interactions from sequence data.

The second experimental validation involved strains from the Zurich (Switzerland)

collection, isolated from soil and freshwater habitats [23]. In this case, we used published

experimental data from the literature [34]. The focus of this earlier study was to test whether the

opportunistic human pathogen *P. aeruginosa* PAO1 can invade natural soil and pond

communities based on its ability to use pyoverdine produced by the natural isolates. We used

data from all the strains for which genome sequences were available (PAO1 and 33 natural

isolates), to establish pyoverdine-mediated iron interaction networks (Figure 2f-g). We then

applied our bioinformatic pipelines as explained for the Nanjing collection and found a high

level of consistency (94%) between the predicted and observed pyoverdine-mediated iron

interactions (Figure 2f-g). Altogether, the two validation experiments demonstrate that

siderophore-mediated microbial interactions can accurately be predicted based on

genome-sequence analysis using the lock-key relationship between receptor and synthetase

genes.

Pyoverdine-mediated iron interaction networks vary across habitats

Following the successful validation, we applied the lock-key pairing methodology to our full

data set to construct the pyoverdine-mediated iron interaction network among all 1928

Pseudomonas strains (Figure 3a). To keep traceability in such an enormous network, we

allocated strains into microbial siderophore functional groups, defined as strains that produce

the same pyoverdine types and can utilize the same repertoire of pyoverdines. Overall, the

network featured 407 microbial siderophore functional groups, 47 different lock-key

receptor-pyoverdine groups, 307 production edges and 1788 utilization edges.

Figure 3 *Pseudomonas* iron interaction networks vary across habitats. a. The predicted

iron interaction networks mediated by pyoverdines among 1928 *Pseudomonas* strains. Circular nodes

represent functional groups (i.e., strains that produce the same pyoverdine type and utilize the same

repertoire of pyoverdines) with node size being proportional to the number of strains within this functional
group. Line colors of circular nodes represent single-receptor producers (green), multi-receptor
producers (yellow) and non-producers (red). Hexagonal nodes represent lock-key pyoverdine groups
with node size being proportional to the number of strains utilizing the corresponding pyoverdine. The
hexagons of the 17 pyoverdine groups found among single-receptor producers are highlighted with the
corresponding receptor group colors (as shown in Fig. 1c), while the pyoverdine groups that are
exclusively found among multi-receptor producers are depicted by grey hexagons. Edges from circular to
hexagonal nodes represent pyoverdine production, while edges from hexagonal to circular nodes
represent pyoverdine utilization (with edge color matching the color of the functional group). **b.** The in-
and **c.** the out-degree distribution of the pyoverdine nodes. The in- and out- degrees are defined by the
number of edges pointing towards (representing production) or originating from (representing utilization)
a pyoverdine node. **d-g.** Iron interaction networks of strains isolated from soil, plant, water and human
habitats. The color bars on the right of each panel show the proportion of single-receptor producers
(green), multi-receptor producers (yellow) and non-producers (red). Node symbols and colors are the
same as in panel **a.**

Our iron interaction network can be considered as a special version of a bipartite network,

characterized by two types of nodes (microbial functional groups and pyoverdine groups) and

two types of directional edges (utilization and production). In ecological bipartite networks such

as those associated with pollination and food webs, topology plays a crucial role for ecological

functions and community assemblies [35]. The topological metrics of our network reveals

significant heterogeneity. Specifically, the in- and out-degree distribution of pyoverdine nodes

is heavy-tailed, indicating "hub" siderophores that are either produced or utilized more

extensively than others (Figure 3b-c). This high degree of heterogeneity suggests non-random

network structure, likely influenced by specific eco-evolutionary forces (*e.g.,* fierce competition

for iron). Indeed, we observed higher than expected yet moderate modularity (Qb=0.51,

compared to Qb=0.41 in a randomized network) and nestedness (NODF=0.15, compared to

NODF=0.04 in a randomized network). These values differ from mutualistic networks that

typically feature high nestedness (*e.g.,* pollination networks), while antagonistic networks are

typically characterized by high modularity (*e.g.,* herbivory and host-parasite networks) [35,36].

Taken together, the iron interaction network may represent a novel type of ecological bipartite

network and stands out as one of the largest reported to date.

Next, we created separate networks for strains isolated from soil (262 strains), plant (234),

water (124), and human-derived (409) habitats. We found that frequencies of the three

pyoverdine strategies and network topologies varied fundamentally between the four habitats

(Figure 3d-g). For example, among the soil-derived strains, there were 56.9% multi-receptor

producers, 27.5% single-receptor producers and 15.7% non-producers (Table S1). In contrast,

there were only 10.0% multi-receptor producers and 4.0% non-producers, but 86.1%

single-receptor producers among human-derived strains. Regarding network topologies, we

observed that the number of microbial functional groups was higher for soil (130, value scaled

to number of strains = 0.50), plant (97, 0.41), and water (70, 0.56) habitats than for

human-related habitats (41, 0.10). Moreover, many functional groups (60.7%) exclusively

occurred in a single habitat: soil (80, 23.7%), plant (56, 16.6%), water (43, 12.7%), and human

(26, 7.7%), whereas only 8 functional groups (Figure S7a) and 11 pyoverdine groups (Figure

S7b) were present in all four habitats. Altogether, these results indicate that the

eco-evolutionary forces shaping the iron interaction networks differ substantially across

habitats.

Pyoverdine-mediated iron interaction networks vary between pathogenic and

non-pathogenic species

Pseudomonas spp. do not only populate a variety of habitats but can also display diverse

lifestyles. The most prominent division in lifestyles occur between pathogenic and

non-pathogenic species. Here, we explore whether iron interaction networks differ between

these two lifestyles. We allocated strains to pathogenic and non-pathogenic species groups

and predicted the iron interaction networks for all species with more than five strains (Fig. 4a

and Table S2). The most abundant pathogenic species were the human pathogen *P.*

aeruginosa and the plant pathogen *P. syringae*, while the most abundant non-pathogenic

environmental species were *P. fluorescens* and *P. putida*, of which many are neutral or even

beneficial for hosts.

Figure 4. Network properties differ between pathogenic and non-pathogenic

***Pseudomonas* spp. a.** The pyoverdine-mediated iron interaction network for pathogenic (top row) and non-pathogenic (bottom row) species with more than five strains in our database. Symbols and color codes match the ones in Figure 3a. **b-c.** The difference of the percentage of multi-receptor producers and the number of the microbial functional group between different species lifestyles. Pathogenic species are colored red, non-pathogenic species are colored blue, and species that do not have clear role in pathogenicity are colored gray. **d.** The difference of diversity quantified by the entropy of their functional group frequencies between different species lifestyles. **e.** Counts of annotated receptors (x-axis) and annotated synthetase (y-axis) for each lock-key group. Lock-key groups are colored by as these in Figure 3a. The grey dotted line represents 0.1 and 0.9 ratios of the synthetase count verse receptor count (termed as "Syn/Rec"). **f.** The difference of the weighted Syn/Rec ratio for each species between different species lifestyles. **g.** The percentage of multi-receptor producers (y-axis) and the weighted Syn/Rec ratio (x-axis) for each species clearly separates pathogenic and non-pathogenic strains (segregation illustrated by the dashed line). The size of the dot represents its diversity quantified by the entropy of their functional group frequencies. Same color coding as that in **b**.

We observed multiple differences in network topologies between the two lifestyles (Figure 4a). First, strains of pathogenic species were mostly single-receptor producers or non-producers (Figure 4a-b), while strains of non-pathogenic species were primarily multi-receptor producers (Figure 4a-b). Second, the diversity of siderophore functional groups was much lower in pathogenic compared to non-pathogenic species (Figure 4c and Table S2). Consequently, the complexity of iron interaction networks (quantified by the entropy of their functional groups) was lower in pathogenic than in non-pathogenic species (Figure 4c). For example, *P. aeruginosa* (the most abundant species in our dataset, 554 strains), has a simple interaction network with only three functional groups, while *P. fluorescens* (135 strains) has a complex interaction network with 13 functional groups (Table S2). Finally, we calculate the Syn/Rec value defined as the ratio of synthetase and receptor groups present in a lock-key group (Figure 4e). In each lock-key group, we calculated the Syn/Rec ratio as the count of synthetase divided by the count of receptors. A Syn/Rec ratio near one indicates that there are no cheating receptors in this lock-key group, *i.e.*, this pyoverdine is exclusive to its producers

and cannot be utilized by strains not producing it. Conversely, a Syn/Rec ratio near zero

means that most receptors in this lock-key group are cheating receptors, and the

corresponding pyoverdine is more sharable and exploitable.

We found that the Syn/Rec ratio was much lower in non-pathogenic than in pathogenic

species (Figure 4f). Furthermore, Syn/Rec ratio differed substantially across lock-key groups

(Figure 4e). For example, the lock-key groups 39 (purple hexagon, presents in *P. aeruginosa*)

and 35 (cyan, presents in several plant pathogens, Fig. 4a) had high Syn/Rec ratios of 0.93

and 0.89, respectively. In contrast, the lock-key group 94 (light green, presents in *P.*

fluorescens) had a much lower Syn/Rec ratio of 0.20. Collectively, the proportion of

multi-receptor producers and the preference for producing shareable or exclusive

siderophores clearly differentiated between pathogenic and non-pathogenic species on the

two-dimensional plane (Figure 4g). Altogether, these results show that non-pathogenic species

form open networks dominated by shareable pyoverdine lock-key groups, whereas pathogenic

species form more closed networks dominated by exclusive pyoverdine lock-key groups.

Modeling the relationship between pyoverdine utilization strategies and community

dynamics

In order to explore the connection between pyoverdine utilization strategies of individual

strains and the resulting community dynamics, we build simple ecological models of

siderophore-mediated ecological competition (Figure 5a) [37]. Our model revealed that different

single-receptor producers cannot co-exist, illustrating the competitive exclusion principle [37] and

matching our observation that a high proportion of single-receptor producers were associated

with simple network structures (Figure 4a). In contrast, multi-receptor producers were more

likely to coexist based on our model (Figure 5b), mirroring our observation that a high

proportion of multi-receptor producers correlated with increased network diversity and

complexity (Figure 3 and Figure 4a). Finally, non-producers could not exist in the absence of

producers in our models.

411
412
Figure 5. Mathematical model exploring the relationship between pyoverdine utilization

413
strategies and community dynamics. a. Schematic diagram of the ecological iron competition

414
model. Each microbe i can produce siderophores of type j with resource budget α_i, and obtain iron by

415
absorbing siderophore-iron complexes through corresponding receptors (fraction of receptors denoted as

416
v_{ij}). Growth rate is proportional to the total amount of absorbed iron and the fraction of resources

417
allocated to primary metabolism, $1 - \alpha_i$. Various strains with distinct strategies were considered,

418
including single-receptor producers (green), multi-receptor producers (orange), and non-producers (red),

419
differing in their α_{ij} and v_{ij} values. 20 strains are assembled into communities, with diverse initial

420
inoculation densities. They then compete in a chemostat-like model until reaching a steady state. **b.**

421
Distribution of the number of species at steady state, for communities initiated with a single strategy type:

422
single-receptor producers (green), multi-receptor producers (yellow), and non-producers (red). **c.**

423
Schematic diagram depicting the invasion scenario together with the different possible outcomes.

424
Putative invaders are introduced at low frequency into a local community at steady state. **d.** Examples

425
illustrating common invasion dynamics. Top panel: a local community with three species (various blue

426
shades) is invaded by a single-receptor producer (purple). Middle panel: a local community is invaded by

a multi-receptor producer. Bottom panel: summary statistics showing the proportion of cases, in which
invaders completely wiped out the local community as a function of their pyoverdine utilization strategy. **e.**
The likelihood of a single-receptor producer to invade a local community of multi-receptor producers for
four different iron interaction networks. Scenarios from top to bottom show: (1) no overlap in siderophore
utilization between the local community and the invader; (2) multi-receptor producer from the local
community can cheat on invader's siderophore; (3) invader and one multi-receptor producer share the
same siderophore; (4) scenario combining (2) and (3). Schematics are simplifications and do not reflect
the actual number of species in the local community.

To delve deeper into the connection between pathogenicity and pyoverdine utilization

strategies, we simulated the consequences of invasion by introducing a strain ("invader") with

a higher resource budget (available for intrinsic growth and siderophore production) into a

steady-state community ("local community") (Figure 5c). Across numerous parameters sets,

we consistently observed that invasions by single-receptor producers and non-producers

tended to disrupt local communities (Figure 5d). Successful single-receptor producers typically

wiped out the local community. Conversely, successful multi-receptor producers integrated

themselves into and co-existed with the local community (Figure 5d). In brief, the more severe

consequences of invasion provide a plausible explanation for the association between certain

strategies and pathogenicity.

Invasion probability further depended on the pyoverdine lock-key relationship between the

invader (here modelled as a single-receptor producer, *e.g.,* matching a pathogenic lifestyle)

and the local community (Figure 5e). The probability of successful invasion was close to zero if

the local community could cheat on the invader's siderophore, which is more likely when the

invader produces siderophores with low Syn/Rec ratio. Conversely, invasion probability was

high if the invader could produce an exclusive siderophore, or if one of the local community

members produced the same siderophore as the invader (*i.e.*, enabling cheating of the

invader). Taken together, our modelling results reveal that the relative proportion of single- and

multi-receptor producers and their interactions via siderophores has fundamental

consequences for community diversity and network complexity.

Discussion

Predicting interactions between microbes from sequence data offers exciting opportunities for

understanding microbiome assembly and stability, and may lay the foundation of

biotechnological and medical microbiome interventions. While sequence-to-interaction

mapping has predominantly been carried out for primary metabolism involving resource

consumption, conversion, and cross-feeding, there are few approaches to reconstruct

microbial interactions based on secondary metabolites (antibiotics, toxins, siderophores,

surfactants) [38-40]. In our paper, we offer such an approach by developing a co-evolution

inspired computational approach to infer iron interaction networks mediated by pyoverdines (a

class of iron-scavenging siderophores) within communities of *Pseudomonas* bacteria.

Pyoverdines can both promote (through molecule sharing) and inhibit (through iron blocking)

the growth of other strains, depending on a molecular lock-key (receptor-pyoverdine)

mechanism[23,29]. Our co-evolution pairing algorithm managed to pair 188 pyoverdines types

and 4547 receptors into 47 lock-key groups. Our experimentally validated approach allowed us

to reconstruct the iron interaction network of 1928 *Pseudomonas* strain. We found intriguing

network differences between habitats (soil, plant, water, human-derived habitats) and between

microbial lifestyles (pathogenic and non-pathogenic). Large and highly connected

siderophore-mediated iron interaction networks occurred among non-pathogenic

environmental strain, whereas small and disconnected network dominated among pathogenic

strains. These results suggest that selection pressures shaping bacterial interaction networks

differ fundamentally between habitats and lifestyles.

Our sequence-to-ecology approach underscores several challenges associated with the

reconstruction of interactions driven by secondary metabolites. The first challenge is that the

chemical structures of secondary metabolites are often difficult to infer from sequencing data

because the metabolites are produced by non-ribosomal peptide and polyketide synthesis

pathways. The second challenge is to identify pyoverdine receptor genes among the many

different types of siderophore receptor genes each strain possesses. We solved these

challenges in a previous paper[31], where we developed approaches based on feature

sequences that allowed us to infer the chemical structure of 188 pyoverdines and identify 4547

pyoverdine receptor genes. The third challenge was to pair pyoverdines to matching receptors

within and across strains. While co-evolution analyses are a widely used computational tool,

employed in diverse areas ranging from ab initio protein structure to host-pathogen interaction

predictions [41], we could not use existing algorithms, such as DCA, SCA, and Evoformer [42-44].

The reason is that these classical site-based co-evolution methods depend on paired

sequences between which the degree of covariation is quantified, yet the existence of

multi-receptor producers impeded direct assignments of synthetase-receptor pairs. For this

purpose, we developed an unsupervised learning algorithm (called Co-evolution Paring

Algorithm), which yielded 47 receptor-synthetase lock-key pairs. This step was essential to

reconstruct iron interaction networks. Our new pipeline has the potential to be applied to

several other microbial traits. For example, microbial membrane receptors co-evolve with

phages[45,46], and pairing phages with the receptor they utilize for infection could provide

insights into host-pathogen co-evolution and thus bacteria-host interaction networks in natural

communities.

Our sequence-to-interaction mapping together with the mathematical model yielded

several new biological insights into the ecology and evolution of microbial iron interaction

networks. First, multi-receptor producers seem to be the glue of iron interaction networks.

They harbor a large diversity of pyoverdine and receptor types, which can connect many other

single- and multi-receptor producers and foster the formation of large and highly connected

networks. Second, networks of non-pathogenic species and communities in natural soil and

water habitats were large and highly interconnected. These networks were dominated by

multi-receptor producers and were more open to invasion. In other words, new strains could

easily integrate themselves into existing communities without much disturbance to the local

community. This suggests that selection for siderophore and receptor diversity is particularly

high in species-rich habitats. Third and in contrast to the above, networks of pathogenic

species and communities in human-derived habitats were small, scattered and had low

complexity. They were dominated by single-receptor producers with a compromised repertoire

of pyoverdine and receptor types. Single-receptor producers were predicted to be the invaders

with severe consequence, having a high potential to disrupt and displace the local community.

Utilization of exclusive siderophore groups increases the chances of successful invasion.

These results suggest that selection favors exclusive siderophores in pathogens as successful

invasion is a key aspect of their lifestyle. Fourth, strains with a strict cheating strategy

(non-producers) occur but are relatively rare (13.6%). However, it is important to note that our

approach yields a conservative non-producer estimate because regulatory non-producers that

have a synthetase cluster but do not express it also occur[22] but cannot be detected with our

approach. Given that non-producers are fully dependent on siderophore producers, it is

intuitive to understand that they are (i) more common in environmental habitats featuring many

multi-receptor producers with a rich pyoverdine repertoire, and (ii) best at invading

multi-receptor communities.

Our insights on pyoverdine utilization strategies and their consequences for community

dynamics reveal ways of how siderophore-mediated interactions could be leveraged for

biotechnological applications [47]. In this context, there is great interest in using probiotic strains

in agriculture to protect crops from infections by bacterial plant pathogens. There is increasing

evidence that siderophore-mediated interactions play a key role in this process[29,30]. For

example, it was shown that plant-beneficial *Chryseobacterium* strains use their siderophores

to suppress the plant-pathogen *Ralstonia solanacearum* [48]. The approach has recently been

extended to human pathogens, which were found to be suppressed by exclusive siderophores

from environmental *Pseudomomnas* spp. [47]. While these studies explored

siderophore-interactions across the species boundaries without clear knowledge on the

specific receptor setup, we here propose a strategic two-pronged design approach. First,

single-receptor producers with exclusive siderophores should be designed and used as

probiotics to competitively exclude pathogens in simple communities. Second, multi-receptor

producers with an exclusive siderophore against the pathogen and cheating receptors utilizing

the pathogen's siderophores should be designed and used as probiotics in more complex

communities. They could integrate themselves without major disruption of the local community,

yet still competitively exclude the pathogen via the exclusive siderophore.

In conclusion, we succeeded in developing a sequence-to-interaction mapping approach

for siderophores that has a high potential to deliver new insights into microbial ecology. Given

that iron is a key trace element that is limited in most environments, siderophore-mediated

interactions are an ideal entry point for secondary metabolite analysis from sequence data.

While we focused on *Pseudomonas* strains, we know that siderophore-mediated interactions

occur across the species boundaries. For example. *P. aeruginosa* possesses receptors to take

up enterobactin produced by *Enterobacteriaceae* spp. and schizokinen produced by. *Ralstonia*

solanacearum [49]. Thus, the next step would be to apply our concepts to more diverse bacterial

communities to derive microbiome-level iron-interaction maps that could further guide rational

designs for biotechnological microbiota interventions.

Data Availability

The source code and parameters used are available in the supplementary material.

Acknowledgements

We thank Xuedi Huang for the insight on pyoverdine receptor analysis.

Funding

This work was supported by the National Key Research and Development Program of China

(No. 2021YFF1200500, 2021YFA0910700), National Natural Science Foundation of China

(No. 42107140, No. 32071255, No.41922053, No.T2321001), National Postdoctoral Program

for Innovative Talents (No. BX2021012). R.K. is supported jointly by a grant from the Swiss

National Science Foundation no. 310030_212266. V-P.F. is supported jointly by Research

Council of Finland, Novo Nordisk Fonden and The Finnish Research Impact Foundation.

Author contributions

Shaohua Gu performed the majority of computational and experimental data analysis in this

research and drafted the manuscript. Zhengying Shao and Shenyue Zhu performed the

experiment of testing pyoverdine-mediated interaction between *Pseudomonas* strains. Zeyang

Qu performed the mathematical model exploring the relationship between pyoverdine

utilization strategies and community dynamics. Di Zhang developed unsupervised co-evolution

pairing algorithm for identifying self-receptor and Richard Allen offered the insight on this

algorithm. Yuanzhe Shao, Ruolin He, Jiqi Shao, Guanyue Xiong assisted in cleaning up the

codes. Alexandre Jousset and Ville-Petri Friman offered insightful comments and assisted in

revising and writing of the manuscript. Rolf Kümmerli and Zhong Wei oversaw the project,

designed experiments and revised the manuscript. Zhiyuan Li conceptualized, oversaw the

project and revised the manuscript.

Competing interests

The authors declare no competing interests.

References

1   Cavicchioli, R. *et al.* Scientists' warning to humanity: microorganisms and climate

585   change. *Nature Reviews Microbiology* **17**, 569-586 (2019).

2   Wang, X. *et al.* Phage combination therapies for bacterial wilt disease in tomato.

Nature Biotechnology **37**, 1513-1520 (2019).

3   Sonnert, N. D. *et al.* A host–microbiota interactome reveals extensive transkingdom

connectivity. *Nature*, doi:10.1038/s41586-024-07162-0 (2024).

4   Kuramitsu Howard, K., He, X., Lux, R., Anderson Maxwell, H. & Shi, W. Interspecies

Interactions within Oral Microbial Communities. *Microbiology and Molecular Biology*

Reviews **71**, 653-670 (2007).

5   Konopka, A., Lindemann, S. & Fredrickson, J. Dynamics in microbial communities:

unraveling mechanisms to identify principles. *The ISME Journal* **9**, 1488-1495 (2015).

6   Handelsman, J. Metagenomics: Application of Genomics to Uncultured

Microorganisms. *Microbiology and Molecular Biology Reviews* **68**, 669-685 (2004).

7   Almeida, A. *et al.* A new genomic blueprint of the human gut microbiota. *Nature* **568**,

499-504 (2019).

8   Schloss, P. D. & Handelsman, J. Metagenomics for studying unculturable

microorganisms: cutting the Gordian knot. *Genome Biology* **6**, 229 (2005).

9   Gu, C., Kim, G. B., Kim, W. J., Kim, H. U. & Lee, S. Y. Current status and applications

of genome-scale metabolic models. *Genome Biology* **20**, 121 (2019).

10   Colarusso, A. V., Goodchild-Michelman, I., Rayle, M. & Zomorrodi, A. R.

Computational modeling of metabolism in microbial communities on a genome-scale.

Current Opinion in Systems Biology **26**, 46-57 (2021).

11   Pacheco, A. R., Moel, M. & Segrè, D. Costless metabolic secretions as drivers of

interspecies interactions in microbial ecosystems. *Nature Communications* **10**, 103

(2019).

12   Gude, S. & Taga, M. E. Multi-faceted approaches to discovering and predicting

microbial nutritional interactions. *Current Opinion in Biotechnology* **62**, 58-64 (2020).

13   Louca, S. & Doebeli, M. Calibration and analysis of genome-based models for

microbial ecology. *eLife* **4**, e08208 (2015).

14   Bozhüyük, K. A. J. *et al.* Evolution-inspired engineering of nonribosomal peptide

synthetases. *Science* **383**, eadg4320 (2024).

15   Penn, K. *et al.* Genomic islands link secondary metabolism to functional adaptation in

marine Actinobacteria. *The ISME Journal* **3**, 1193-1203 (2009).

16   Scherlach, K. & Hertweck, C. Mediators of mutualistic microbe–microbe interactions.

Natural Product Reports **35**, 303-308 (2018).

17   Vining, L. C. in *Ciba Foundation Symposium 171 - Secondary Metabolites: their*

Function and Evolution Novartis Foundation Symposia 184-198 (2007).

18   He, R. *et al.* Knowledge-guided data mining on the standardized architecture of NRPS:

Subtypes, novel motifs, and sequence entanglements. *PLOS Computational Biology*

19, e1011100 (2023).

19   Andrews, S. C., Robinson, A. K. & Rodríguez-Quiñones, F. Bacterial iron homeostasis.

FEMS Microbiology Reviews **27**, 215–237 (2003).

20   Boyd, P. W. & Ellwood, M. J. The biogeochemical cycle of iron in the ocean. *Nat.*

Geosci. **3**, 675-682 (2010).

21   Emerson, D., Roden, E. & Twining, B. The microbial ferrous wheel: iron cycling in

terrestrial, freshwater, and marine environments. *Frontiers in Microbiology* **3**,

doi:10.3389/fmicb.2012.00383 (2012).

22 Butaitė, E., Baumgartner, M., Wyder, S. & Kümmerli, R. Siderophore cheating and

cheating resistance shape competition for iron in soil and freshwater Pseudomonas

communities. *Nature Communications* **8**, 414 (2017).

23 Kramer, J., Özkaya, Ö. & Kümmerli, R. Bacterial siderophores in community and host

interactions. *Nature Reviews Microbiology* **18**, 152-163 (2020).

24 Leventhal, G. E., Ackermann, M. & Schiessl, K. T. Why microbes secrete molecules to

modify their environment: the case of iron-chelating siderophores. *Journal of The*

Royal Society Interface **16**, 20180674 (2019).

25 West, S. A., Diggle, S. P., Buckling, A., Gardner, A. & Griffin, A. S. The Social Lives of

Microbes. *Annual Review of Ecology, Evolution, and Systematics* **38**, 53-77 (2007).

26 Miethke, M. & Marahiel Mohamed, A. Siderophore-Based Iron Acquisition and

Pathogen Control. *Microbiology and Molecular Biology Reviews* **71**, 413-451 (2007).

27 Seyoum, Y., Baye, K. & Humblot, C. Iron homeostasis in host and gut bacteria – a

complex interrelationship. *Gut Microbes* **13**, 1874855 (2021).

28 Schaible, U. E. & Kaufmann, S. H. E. Iron and microbial infection. *Nature Reviews*

Microbiology **2**, 946-953 (2004).

29 Gu, S. *et al.* Competition for iron drives phytopathogen control by natural rhizosphere

microbiomes. *Nature Microbiology* **5**, 1002-1010 (2020).

30 Gu, S. *et al.* Siderophore-Mediated Interactions Determine the Disease

Suppressiveness of Microbial Consortia. *mSystems* **5**,

10.1128/msystems.00811-00819 (2020).

31 Shaohua, G. *et al.* From sequence to molecules: Feature sequence-based genome

mining uncovers the hidden diversity of bacterial siderophore pathways. *bioRxiv*,

2023.2010.2030.564663, doi:10.1101/2023.10.30.564663 (2023).

32 Bozhüyük, K. A. J. *et al.* De novo design and engineering of non-ribosomal peptide

synthetases. *Nature Chemistry* **10**, 275-281 (2018).

33 Meyer, J. M. *et al.* Use of siderophores to type pseudomonads: the three

Pseudomonas aeruginosa pyoverdine systems. *Microbiology (Reading)* **143 (Pt 1)**,

35-43 (1997).

34 Figueiredo, A. R. T., Özkaya, Ö., Kümmerli, R. & Kramer, J. Siderophores drive

invasion dynamics in bacterial communities through their dual role as public good

versus public bad. *Ecology Letters* **25**, 138-150 (2022).

35 Thébault, E. & Fontaine, C. Stability of Ecological Communities and the Architecture of

Mutualistic and Trophic Networks. *Science* **329**, 853-856 (2010).

36 Bascompte, J., Jordano, P., Melián, C. J. & Olesen, J. M. The nested assembly of

plant–animal mutualistic networks. *Proceedings of the National Academy of Sciences*

100, 9383-9387 (2003).

37 Jiqi, S., Yinxiang, L., Jingyuan, L., Shaohua, G. & Zhiyuan, L. Siderophore Piracy

Promotes Dynamical Coexistence in Microbial Community. *bioRxiv*,

2023.2011.2021.568182, doi:10.1101/2023.11.21.568182 (2023).

38 Tsilimigras, M. C. B. & Fodor, A. A. Compositional data analysis of the microbiome:

fundamentals, tools, and challenges. *Annals of Epidemiology* **26**, 330-335 (2016).

39 Weiss, S. *et al.* Correlation detection strategies in microbial data sets vary widely in

sensitivity and precision. *ISME J* **10**, 1669-1681 (2016).

40   Faust, K. & Raes, J. Microbial interactions: from networks to models. *Nature Reviews*

Microbiology **10**, 538-550 (2012).

41   Chen, H. M., Guo, W., Shen, J., Wang, L. & Song, J. N. Structural Principles Analysis

of Host-Pathogen Protein-Protein Interactions: A Structural Bioinformatics Survey.

Ieee Access **6**, 11760-11771 (2018).

42   Greensmith, J., Feyereisl, J. & Aickelin, U. The DCA: SOMe comparison. *Evolutionary*

Intelligence **1**, 85-112 (2008).

43   Mirjalili, S. SCA: A Sine Cosine Algorithm for solving optimization problems.

Knowledge-Based Systems **96**, 120-133 (2016).

44   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*

596, 583-589 (2021).

45   Hampton, H. G., Watson, B. N. J. & Fineran, P. C. The arms race between bacteria

and their phage foes. *Nature* **577**, 327-336 (2020).

46   Kortright, K. E., Chan, B. K., Koff, J. L. & Turner, P. E. Phage Therapy: A Renewed

Approach to Combat Antibiotic-Resistant Bacteria. *Cell Host & Microbe* **25**, 219-232

(2019).

47   Vera, V. *et al.* Antimicrobial activity of iron-depriving pyoverdines against human

opportunistic pathogens. *bioRxiv*, 2023.2007.2018.549568,

doi:10.1101/2023.07.18.549568 (2023).

48   Rehm, K. *et al.* Chryseochelins—structural characterization of novel citrate-based

siderophores produced by plant protecting Chryseobacterium spp. *Metallomics* **15**,

mfad008 (2023).

49   Kümmerli, R. Iron acquisition strategies in pseudomonads: mechanisms, ecology, and

evolution. *Biometals* **36**, 777-797 (2023).

699