Bioenergetic mapping of 'healthy microbiomes' via compound processing potential

imprinted in gut and soil metagenomes

Craig Liddicoat[a,*], Robert A. Edwards[a], Michael Roach[a], Jake M. Robinson[a], Kiri Joy Wallace[b], Andrew D.

Barnes[b], Joel Brame[a], Anna Heintz-Buschart[c], Timothy R. Cavagnaro[a], Elizabeth A. Dinsdale[a], Michael P.

Doane[a], Nico Eisenhauer[d,e], Grace Mitchell[b,f], Bibishan Rai[b], Sunita Ramesh[a], Martin F. Breed[a]

[a]College of Science and Engineering, Flinders University, Bedford Park, South Australia, Australia

[b]Environmental Research Institute, University of Waikato, Hamilton, Aotearoa New Zealand

[c]Swammerdam Institute for Life Sciences, University of Amsterdam, 1098 XH Amsterdam, The

Netherlands

[d]German Centre for Integrative Biodiversity Research (iDiv), 04103 Leipzig, Germany

[e]Institute of Biology, Leipzig University, 04103 Leipzig, Germany

[f]Manaaki Whenua – Landcare Research, Hamilton, Aotearoa New Zealand

Corresponding author: Craig Liddicoat; craig.liddicoat@flinders.edu.au

ORCIDs: CL: 0000-0002-4812-7524, RAE: 0000-0001-8383-8949, MR: 0000-0003-1488-5148, JMR:

0000-0001-8108-3271, KJW: 0000-0003-1236-6846, ADB: 0000-0002-6499-381X, JB: 0000-0002-1852-

9987, AHB: 0000-0002-9780-1933, TRC: 0000-0002-9922-5677, EAD: 0000-0002-2177-203X, MPD:

0000-0001-9820-2193, NE: 0000-0002-0371-6720, GM: 0000-0002-2839-7981, SR: 0000-0003-2230-

4737, MFB: 0000-0001-7810-9696

Author Contributions: CL, RAE, MFB designed research; CL, MR, JMR, KJW, ADB, AHB, MPD, NE,

GM, BR performed research; CL, RAE, JMR, KJW, ADB, JB, AHB, TRC, EAD, MPD, NE, SR, MFB

analyzed data; CL drafted the manuscript; all authors reviewed and edited the final version.

Preprint Server: bioRxiv.org (ID: BIORXIV/2023/565728) [CC BY-NC-ND]

Competing Interest Statement: The authors declare no competing interest.

Keywords: Metagenomics, compound processing potential, healthy microbiome, human health,

bioenergetic mapping, van Krevelen diagram

ABSTRACT

Despite mounting evidence of their importance in human health and ecosystem functioning, the definition and measurement of 'healthy microbiomes' remain unclear. More advanced knowledge exists on health associations for compounds used or produced by microbes. Environmental microbiome exposures (especially via soils) also help shape, and may supplement, the functional capacity of human microbiomes. Given the synchronous interaction between microbes, their feedstocks, and micro-environments, with functional genes facilitating chemical transformations, there exists an intriguing opportunity to examine microbiomes in terms of their capacity to process compounds relevant to human health. Here we integrate functional genomics and biochemistry frameworks to derive new quantitative measures of the potential for human gut and environmental soil metagenomes to process major compound classes (e.g., lipids, carbohydrates) and selected biomolecules (e.g., vitamins, short-chain fatty acids) linked to human health. Metagenome functional potential profile data were translated into a universal compound mapping 'landscape' based on bioenergetic van Krevelen mapping of function-level meta-compounds and corresponding functional relative abundances, reflecting imprinted genetic capacity of microbiomes to metabolize an array of different compounds. We show that measures of 'compound processing potential' associated with human health and disease (examining atherosclerotic cardiovascular disease, colorectal cancer, type 2 diabetes and anxious-depressive behavior case studies), and displayed seemingly predictable shifts along gradients of ecological disturbance in plant-soil ecosystems (three case studies). Ecosystem quality explained 60–92% of variation in soil metagenome compound processing potential measures in a post-mining restoration case study dataset. With growing knowledge of the varying proficiency of environmental microbiota to process human health associated compounds, we might design environmental interventions or nature prescriptions to modulate our exposures, thereby advancing microbiota-oriented approaches to human health. Compound processing potential offers a simplified, integrative approach for applying metagenomics in ongoing efforts to understand and quantify the role and linkages of microbiota in environmental- and human-health.

1. Introduction

Microbial communities (microbiota), their feedstocks (substrates, nutrients) and environmental conditions (e.g., pH, redox potential, temperature, moisture, salinity) work in concert to drive microbially-mediated reactions essential to fueling life on Earth (Averill et al., 2022). Microbiomes (microbiota, genetic material and metabolites) are critical to the health and functioning of humans (Banerjee and van der Heijden, 2022; Gilbert et al., 2018; Round and Mazmanian, 2009) and ecosystems (Delgado-Baquerizo et al., 2020; Sokol et al., 2022). However, clear definitions of what constitutes a 'healthy microbiome' remain elusive (Eisenstein, 2020). More advanced knowledge exists on health associations for the compounds used or produced by microbes. Because microbes, their feedstocks and micro-environments interact synchronously—using functional genes to facilitate chemical transformations—this presents an intriguing opportunity to examine microbiomes through their potential to process compounds associated with human health.

Human microbiomes are supplemented and partly shaped by environmental microbial exposures (Banerjee and van der Heijden, 2022; Gilbert et al., 2018). Therefore, the involvement of environmental microbiomes in processing human health-associated compounds is also of interest. Transfer of environmental microbiota to humans may help supplement important functional capacity, protective microbiota, and immune-signaling agents, particularly in infants, but also in adults who have depleted microbiota due to antibiotic use, poor diet, lifestyle or other health incidents (Banerjee and van der Heijden, 2022; Brame et al., 2021; Flandroy et al., 2018; Roslund et al., 2022). We know that microbiota are shaped by the resources they utilize and the environments they inhabit (Fierer, 2017; Gilbert et al., 2018). If the functional composition of microbiota varies predictably along environmental gradients, then through design, management, and behavior we should be able to modulate our exposure to health-promoting versus disease-associated microbes. Soils, in particular, can represent a rich source of microbial diversity with potential to support human health (Banerjee and van der Heijden, 2022; Li et al.; Liddicoat et al., 2018; Liddicoat et al., 2020; Ottman et al., 2018; Sun et al., 2023). Microbiota in plant-soil systems are shaped by

macro-scale factors including climate, soil characteristics, vegetation composition, diversity, land use and

management (Delgado-Baquerizo et al., 2018; Fierer, 2017). With the prospect of cost-effectively

encouraging health-promoting microbes, it is frequently asked, "What type of environment is best?" Yet,

the attributes of health-promoting environmental microbiomes, including potential functional overlaps with

human microbiomes, remain understudied.

Microbes typically operate as a community (Barton and Northup, 2011) where many taxa may be

highly efficient at performing certain functions, but they lack functional capacity for stand-alone survival

(Zengler and Zaramela, 2018). Complex cross-feeding and resource sharing in the extracellular space

(Daisley et al., 2021), and involvement of multiple taxa in functional pathways, suggest that community-

scale functional profiles (rather than specific microbial taxa) underpin the health-supporting capacity of

microbiota. Indeed, it is this community-scale complexity that has hindered progress towards clear

definitions of a healthy microbiome. Nevertheless, researchers want to better understand the assembly and

structure of health-promoting microbiomes to improve the course of microbiome-associated diseases.

Disease-associated microbiota are often characterized by a loss of diversity and overgrowth by

opportunistic pathogens (Belizario and Napolitano, 2015; Round and Mazmanian, 2009), but it may be

unclear whether they are the facilitators or followers of disease.

Many microbiome-associated diseases are linked to bioenergetic mechanisms (Daisley et al., 2021),

with oxidation-reduction (redox) potential recognized as a key factor shaping microbial communities. The

healthy anaerobic gut favors obligate anaerobes, whereas dysbiosis is often accompanied by increased

oxygenation of the colonic epithelium and expanded abundance of oxygen-tolerant facultative anaerobic

bacteria (Litvak et al., 2018; Rivera-Chávez et al., 2017). Oxygen is a highly electronegative element

important in shaping electrochemical gradients, biochemical reactions, and gene expression (Wilson et al.,

2020). Oxygen content varies in different types of organic matter (i.e., microbial feedstocks). Yet, the

interrelationship between bioenergetic (electrochemical) status of compounds, microbial environments,

microbiota development and human health appears to receive insufficient attention. In soils, redox potential

varies with weather, vegetation, land use, management, drainage, organic-content, vicinity to roots, soil

characteristics, and microbial activity (Hinsinger et al., 2009; Husson, 2013). At the molecular level, redox

potential shapes what kind of molecules can be made and how energy is stored. Therefore, a compound-

oriented examination of healthy microbiomes might capitalize on available knowledge linking compounds

with human health, while also considering deterministic influences of bioenergetic (electrochemical)

context.

Here, we examine the functional potential of gut and soil microbiota from a compound processing

potential (CPP) viewpoint, to assess patterns in human health and disease, and with gradients of ecosystem

maturity (or quality). Such an approach might discern health- versus disease-promoting microbiotas from

the types of compounds they are attuned to consuming or producing (reflecting microbial feedstocks and

metabolites respectively). Previous metabolome prediction frameworks (e.g., Diener et al., 2020; Garza et

al., 2018; Mallick et al., 2019; Neveu et al., 2023) rely variously on supplementary metabolome training

datasets, microorganism-specific genome-scale metabolic models, taxonomic abundance estimates, and

assumed environments (e.g., human gut), to model short-term reaction rates. Here, we wanted to exploit

community-scale compound-oriented information that might be embedded within metagenomes, regardless

of taxa present. As functional potential profiling from whole genome sequencing (or shotgun

metagenomics) does not directly measure functions performed, we characterized microbiota-linked CPP

from DNA sequencing, without direct measurement of compounds. Our conceptual interest was to

characterize the distribution of available community-scale genetic capacity of microbiomes to process an

array of compound classes and specific biomolecules (compounds produced by living organisms), which

may be present over a period of time and with variation in feedstocks and environmental conditions,

consistent with dynamic influences that have shaped existing microbiota structure. We premised that the

ease of transformation between health-associated compounds and other compounds that closely resemble

them will depend on stoichiometric and energetic similarities, microbiota functional diversity, and

environmental conditions.

We utilized a framework that integrates information about compounds, bioenergetics, and

environmental conditions. The complexity-reducing van Krevelen (vK) coordinate space offers a simplified

and intuitive bioenergetic framework for approximate mapping of compounds based on their carbon (C),

oxygen (O) and hydrogen (H) content, while also reflecting energy density and principal axes that explain

microbial distribution and chemical speciation (Fig. 1 and Fig. S1) (D'Andrilli et al., 2015; Wu et al., 2018).

Compounds are mapped into vK space using their O:C and H:C molar ratios (x- and y-axis respectively).

We surmised this framework could offer an exhaustive and intuitive mapping space to summarize the nature

of microbiota-mediated functional reactions in a way that reflects the collective mean or dominant

compound properties, reaction stoichiometries, and potential overlaps between dietary or environmental

substrates, and key health-associated compounds.

Specifically, we combined SUPER-FOCUS functional profiling annotations (Silva et al., 2015), the

comprehensive ModelSEED (Seaver et al., 2020) functional-biochemistry database system and vK

coordinate mapping to assign functional potential relative abundances from human gut and soil sample

metagenomes to overall mean function-level vK coordinates. This approach effectively mapped every

SUPER-FOCUS function (where feasible via available corresponding database information), based on one

or more associated chemical reactions, to a function-level 'meta-compound' represented in the two-

dimensional vK space (Fig. S2, Methods). We aimed to: (1) investigate measures of microbiota CPP

imprinted in human gut and soil metagenomes; (2) test for differences in human health and disease, and in

disturbed, restored and natural ecosystems; and (3) identify compounds linked to human health with

differing potential metabolism (i.e., CPP) by environmental microbiota due to differences in ecosystem

quality. We hypothesized this universal bioenergetic compound mapping approach might identify CPP

profiles, and possible overlaps in human and environmental metagenomes, that could inform the definition

and future shaping of 'healthy microbiomes'. We were also keen to explore whether CPP measures might

enhance the interpretability and accessibility of metagenomics data to aid hypothesis building and

prioritizing future research.

Fig. 1. Van Krevelen (vK) coordinate space (adapted from D'Andrilli et al., 2015), displaying major compound classes (purple zones and text), key gradients (grey axes and text), focus biomolecules examined in this study (large dots), and additional example health-associated biomolecules, dietary or environmental substrates (legend) (see Tables S1, S2). vK zones were adapted from (Wu et al., 2018). Key gradients include oxidation-reduction (x-axis), hydrogenation-dehydrogenation (y-axis), hydration-condensation (top-right to bottom-left), methylation-demethylation (top-left to bottom-right) and increasing energy content (towards top-left).

2. Materials and Methods

2.1. Case study datasets

We assessed metagenomics samples from four human health and disease datasets, comprising

atherosclerotic cardiovascular disease (ACVD) (Jie et al., 2017), colorectal cancer (Zeller et al., 2014), type

2 diabetes (T2D) (Forslund et al., 2015) and problem (anxious-depressive) behaviors in children (Flannery

et al., 2020); and three environmental soil datasets from ecological restoration and disturbed versus natural

plant-soil systems (Barnes et al., 2020; Bissett et al., 2016; Sun and Badgley, 2019) (Table 1; further detail

in Appendix A Supporting Information).

Table 1. Description of case study metagenome datasets.

Case study focus	Main comparison variable or diagnosis groups (and sample/subject numbers[a])	Metagenome functional profile characteristics (sample mean ± s.d.)[b]	Source data reference; Country of origin
Human gut			
Atherosclerotic cardiovascular disease (ACVD)	ACVD (n_F = 53, n_M = 157) or normal healthy (n_F 101, n_M = 69). Total n = 380.	Functions[0] n = 21,117 (9301 ± 3523) Functions[vK] n = 10,829 (5051 ± 1843) Total fxn rel abun[vK] = 63.2 ± 3.5 % vK coordinates n = 2535 (1509 ± 502)	SRA accession PRJEB21528 (Jie et al., 2017); China
Colorectal cancer	Colorectal cancer (n_F = 24, n_M = 29) or normal healthy (n_F = 33, n_M = 27). Total n = 113.	Functions[0] n = 12,896 (2809 ± 1924) Functions[vK] n = 6932 (1750 ± 1117) Total fxn rel abun[vK] = 66.7 ± 2.7 % vK coordinates n = 1954 (711 ± 362)	SRA accession PRJEB6070 (Zeller et al., 2014); France
Type 2 diabetes (T2D) and impaired glucose tolerance (IGT)	T2D with no Metformin treatment (T2D Met-, n = 33), T2D with Metformin (T2D Met+, n = 20), IGT (n = 49), or normal healthy (n = 43). Subjects are females only. Total n = 145.	Functions[0] n = 19,099 (9916 ± 2393) Functions[vK] n = 9916 (4490 ± 874) Total fxn rel abun[vK] = 64.4 ± 2.2 % vK coordinates n = 2393 (1431 ± 202)	SRA accession PRJEB1786 (Forslund et al., 2015); Sweden
Problem behaviors in children	First principal component (PC1) of anxious-depressive problem behaviors, examined as either numeric scores (n_F =	Functions[0] n = 20,599 (10282 ± 2481) Functions[vK] n = 10,322 (5507 ± 1224) Total fxn rel abun[vK] = 54.9 ± 1.3 %	SRA accession PRJNA496479 (Flannery et al.,

	20, n_M = 17); or high/low PC1 groups[c] (high PC1 n_F = 10, n_M = 8; Low PC1 n_F = 10, n_M = 9). Total n = 37.	vK coordinates n = 2459 (1645 ± 281)	2020); United States
Soils			
People Cities and Nature (PCaN) urban forest ecosystem restoration	Soil samples spanned young to old revegetation age, and remnant sites (treated as ordinal variables). Data were separated into pH-based groups: strongly acidic (pH < 4.5, 10-40 yr old, remnant, n = 8); and acidic-neutral soils (4.5 < pH < 7, 11-48 yr old, n = 10). Total n = 18.	Functions[0] n = 36,324 (27,969 ± 1011) Functions[vK] n = 18,197 (14,690 ± 396) Total fxn rel abun[vK] = 61.7 ± 0.3 % vK coordinates n = 3302 (2965 ± 49)	Aotearoa Genomic Data Repository project AGDR00045 (Barnes et al., 2020); Aotearoa New Zealand
Post-mining forest ecosystem restoration	Soil samples spanned revegetation ages of 6, 12, 22, 31 years, and unmined (UM) samples (treated as ordinal variables). Comprising five age-based groups, each with three replicates. Total n = 15.	Functions[0] n = 30,125 (20,328 ± 767) Functions[vK] n = 15,576 (11,051 ± 334) Total fxn rel abun[vK] = 61.9 ± 0.3 % vK coordinates n = 3076 (2537 ± 54)	MG-RAST project mgp16379 (Sun and Badgley, 2019); United States
Australian Microbiome Initiative (AMI) disturbed versus natural	Disturbed (n = 29) or natural (n = 55) soils, comprising temperate climate zone, surface (0-10cm) with 7.5–45% clay content (i.e., avoiding very sandy and very clayey soils). Total n = 84.	Functions[0] n = 37,335 (22,959 ± 1172) Functions[vK] n = 18,551 (12,212 ± 553) Total fxn rel abun[vK] = 60.8 ± 1.0 % vK coordinates n = 3326 (2700 ± 69)	AMI Data Portal [Data accessed Sep 2022] (Bissett et al., 2016) [d]; Australia

178　Notes: [a]F = females, M = males. [b]Number of functions[0] from initial SUPER-FOCUS profiling, versus

179　functions[vK] (fxn) with available compound information for mapping to vK coordinates. [c]PC1 of problem

180　behaviors were analyzed as numeric values for visualizing and assessing CPP_{class} and $CPP_{density}$ data; and

181　high vs. low PC1 groups for CPP_{ASALR} data. High PC1 values represent more problematic behavior. [d]AMI

182　Data portal URL: https://data.bioplatforms.com/organization/australian-microbiome.

2.2. *Compound processing potential (CPP) mapping approach*

CPP measures were derived via the following steps (Fig. 2; further detail in Appendix A Supporting Information):

a. Shotgun metagenomics raw sequences were accessed, and bioinformatic steps were run on Flinders University DeepThought high-performance computing facility (Flinders_University, 2021).

b. Raw sequence data were inspected using FastQC (v0.11.9; Andrews, 2018) and quality control trimming performed using Fastp (v0.23.2; Chen et al., 2018).

c. Functional potential profiles were derived from good quality read 1 sequences using SUPER-FOCUS (Silva et al., 2015) functional annotation software, linked to the Diamond sequence aligner (v0.9.19; Buchfink et al., 2021) and version 2 100% identity-clustered reference database (100_v2; https://github.com/metageni/SUPER-FOCUS/issues/66). Where subjects/samples were represented by multiple sequence files, the combined SUPER-FOCUS outputs were normalized so that the total functional relative abundances summed to 100% in each subject/sample.

d. Every SUPER-FOCUS function (output row) was translated to one or more corresponding chemical reaction(s) using a purpose-built R-script algorithm based on ModelSEED database lookup tables (from https://github.com/ModelSEED/ModelSEEDDatabase; accessed 10 Aug 2022). The algorithm sought matches based on either: full matching of functional hierarchies (using subsystem-class, -subclass, -name and -role); detection of EC number; or matching of SUPER-FOCUS function name within ModelSEED lookup tables for reactions (reaction name or alias), subsystems (role), or reaction-pathways (external reaction name).

e. Every chemical reaction was converted to mean vK coordinates (O:C and H:C molar ratios), considering all C-containing reaction input and product compounds and weighted according to reaction stoichiometry (see example in Fig. 2.). Compounds not containing C were ignored due to undefined O:C and H:C ratios. Data for compounds were based on Hill system chemical formulae in protonated form.

f.   Overall mean vK coordinates were calculated for each SUPER-FOCUS function via averaging one or

more linked chemical reactions. The majority of SUPER-FOCUS functional relative abundances

were successfully mapped to vK coordinates (sample ranges 52-84%, means 55-67%, Table 1,

Table S3).

g.   In vK coordinate space, we analyzed the spatial assignment of functional relative abundances to derive

the following CPP data types:

1) CPP_{class}: summed functional relative abundances mapping to major compound classes (based on

215   pre-defined zones from Wu et al., 2018) (see Fig. 1, Table S1).

216   2) CPP_{ASALR}: to address large variation in CPP_{class} values across case studies, we implemented a

217   first-pass normalization aiming to account for microbial activity levels, here termed amino sugar

218   adjusted log ratio (ASALR) data. Amino sugars have previously been used as a biomarker of

219   microbial residue turnover (Hu et al., 2018) as they are major components of bacterial and

220   fungal cell walls (peptidoglycan and chitin). CPP_{class} values were divided by the CPP_{class} value

221   for amino sugars, followed by a variance-stabilizing log10-transformation.

222   3) $CPP_{density}$: captured the density of functional relative abundances within radial buffers of varying

223   proximity (radii of 0.05, 0.1, 0.15, 0.2, 0.25 vK units) to selected focus biomolecules (Fig. 1,

224   Table S2, Appendix A Supporting Information). These comprised short-chain fatty acids:

225   acetate, propionate, and butyrate; vitamins: riboflavin (B2), cobalamin (B12), pyridoxal 5'-

226   phosphate (B6), folate (B9), menaquinone (K2); and glutamate and pyruvate. Functional relative

227   abundances within radial buffers were summed then divided by the respective area in vK units^2.

228   4)   Compound-associated vK coordinates: Each SUPER-FOCUS functional row was translated to

229   corresponding vK coordinates (as used for the above spatial assignments). Further

230   supplementary analyses were undertaken at the vK coordinate level, including network analyses

231   (in the T2D case study), differential abundance analysis (in the problem behavior case study),

232   and weighted-mean vK coordinate analysis within major compound zones.

Fig. 2. Flow chart of compound processing potential (CPP) analyses. The example reaction shows

calculation of mean O:C and H:C molar ratios (e.g., 1.833, 0.5) based on stoichiometry of all carbon-

containing compounds. Mean O:C and H:C values are then averaged across one or more chemical reactions

per function, to derive function-level vK coordinates. Each function-level mapping into vK space represents

a conceptual 'meta-compound' based on the collective elements and stoichiometry of all C-containing

compounds involved in all reactions corresponding to a single microbiota-mediated function.

 2.3. Data visualization and statistical analyses

Details of visualization and statistical testing using standard approaches are provided in Appendix A

Supporting Information.

3. Results

3.1. Human health and disease

Gut metagenome CPP_{class} (Figs. S4, S8, S12, S18, Table S3) and $CPP_{density}$ (Figs. S5-S6, S9-S10, S13-S14,

S19-S20, Table S4) data produced strong associations in ACVD and colorectal cancer compared to normal

subjects (detailed below). Many patterns observed in CPP_{class} data were reinforced in the putative activity-

normalized CPP_{ASALR} measures (Fig. 3, Table S5), and this transformed data format showed stabilized

variance across case study datasets. Interestingly, across all CPP_{class}, $CPP_{density}$, and CPP_{ASALR}

measurements, when associations were found in both sexes they were always in the same direction (Tables

S3-S5). In the T2D (female only) and problem behavior case studies, we observed far fewer relationships in

the coarse CPP_{class}, CPP_{ASALR} or biomolecule-focused $CPP_{density}$ data. Therefore, we pursued network

analyses and differential abundance analyses respectively in these case studies, as illustrative examples of

more detailed supplementary analyses. Weighted mean vK-coordinate analyses produced striking

associations in ACVD (Fig. S7 Tables S6-S7), but weaker effects in other case studies (Figs. S11, S15, S21,

Tables S8-S12). The statistical test results described below are detailed in Appendix A.

Fig. 3. Amino sugar-adjusted log ratio compound processing potential (CPP$_{ASALR}$), representing putative microbiota activity-normalized values, in normal healthy and diseased female (F) and male (M) subjects for atherosclerotic cardiovascular disease (ACVD), colorectal cancer, type 2 diabetes (T2D) with and without Metformin treatment (met +/-), impaired glucose tolerance (IGT), and high and low first principal component (PC1) problem behavior values. For visualization purposes outlying values are not shown. However, statistical tests were based on all data (Table S5). Sample sizes are detailed in Table 1. Tests for differences are performed within a single sex. In T2D data, groups not sharing a letter are different.

ACVD associated with CPP_{class} values, compared to normal subjects, in the form of increased

potential metabolism of lipids (in females – F, and in males – M), proteins (M), condensed aromatics (F, M)

lignin (F, M), tannins (M), other hydrated compounds (F, M), other demethylated compounds (M), and near

vitamin K2-vitamin B12 compounds (F); contrasting with decreased potential metabolism of carbohydrates

(F, M) and other condensed compounds (M). Despite all samples initially summing to 100% total functional

relative abundances, there was less complete conversion to identifiable reactions and total sum CPP_{class} data

in females only for ACVD compared to normal cases. Many of these patterns were reinforced in the

CPP_{ASALR} data, with ACVD notably associated in both sexes with increased potential metabolism of lipids,

condensed aromatics, lignin, and other hydrated compounds; but decreased potential metabolism of

carbohydrates. For $CPP_{density}$ measures within 0.05 vK unit radii, ACVD associated with increased potential

metabolism of propionate (F), vitamin B12 (F), vitamin B6 (M), vitamin B9 (F, M), vitamin K2 (F, M) and

pyruvate (M); but decreased potential metabolism of acetate (M) and glutamate (F, M).

Colorectal cancer also associated prominently with CPP_{class} values, specifically increased potential

metabolism of lipids (M), amino sugars (F, M), condensed aromatics (M), lignin (M), other condensed

compounds (F, M) and near vitamin K2-vitamin B12 compounds (M); but decreased potential metabolism

of proteins (F, M) and tannins (F, M). Total sum CPP_{class} data were reduced in male colorectal cancer cases

compared to normal. CPP_{ASALR} values in colorectal cancer subjects of both sexes were associated with

decreased potential metabolism of proteins, lignin, tannins, and other demethylated compounds. For

$CPP_{density}$ measures within 0.05 vK unit radii, colorectal cancer associated with increased potential

metabolism of vitamin B2 (M), vitamin B12 (F, M), vitamin B9 (F, M), and pyruvate (M). Weighted mean

vK-coordinates varied between colorectal cancer and normal subjects with different centroids in condensed

aromatics (F), lignin (M), and tannins (M); and different beta-dispersion or spread in proteins (F, M) and

condensed aromatics (F, M).

In the female-only T2D case study results, we found CPP_{class} values decreased for other methylated

compounds in IGT compared to normal subjects and decreased for near vitamin K2-vitamin B12

compounds in IGT compared to normal, with a reduced total sum CPP_{class} in normal compared to other

subjects. No associations were found with $CPP_{density}$ values. CPP_{ASALR} values were decreased for near

vitamin K2-vitamin B12 compounds in IGT compared to normal and T2D Met+ subjects. From analysis of

weighted mean vK-coordinates among diagnosis groups, a difference in centroids was found only in the

compound class of proteins. Lacking interpretable results at the compound class or focus biomolecule

levels, further investigation was performed via network analysis (Appendix A Supporting Information) on

20 subjects in each diagnosis group based on inferred correlations between functional relative abundances

mapped to unique vK-coordinates. Network analyses were based on commonly observed vK coordinates

(present in at least 60% of samples and minimum 2% sum functional relative abundance across samples).

Network diagrams (Fig. 4) and structure dendograms (Fig. S16) display a transition in their complexity,

number of nodes and fraction of negative edges (interactions) from simplest in normal and IGT subjects to

most complex in T2D Met+ and T2D Met- subjects. Comparing network characteristics for the four groups

(normal, IGT, T2D Met-, T2D Met+; n = groups of 20) to a bootstrapped (B = 1000) density distribution of

randomly resampled networks (n = 20, drawn from the same pool of 80 subjects) (Fig. S17, Table S17) we

found normal healthy subjects had the lowest fraction of negative edges and the highest degree

centralization. Untreated disease, T2D Met-, had the lowest closeness centralization (graph-level inverse of

average geodesic distance between nodes); and borderline significant results for the highest fraction of

negative edges (negative correlations between vK-coordinates), lowest betweenness centralization (graph-

level centrality based on broker positions connecting others), and lowest mean distance (average path length

between nodes). In short, normal subjects appear to have far less correlations between vK-coordinates

(fewer nodes / vertices), and for the nodes and links that are present they are largely positively correlated

and highly interlinked. Whereas T2D Met- (untreated disease) is characterized by a much larger number of

negatively correlated vK-coordinate nodes, which on average have shorter links, and are less well

connected-up across the whole network.

Fig. 4. Network diagrams based on commonly observed vK coordinates for female subjects (n = groups of 20) with diagnoses: (a) normal, (b) impaired glucose tolerance (IGT), (c) type 2 diabetes without Metformin (T2D Met-), and (d) type 2 diabetes with Metformin (T2D Met+). Nodes are located according to unique compound-associated vK coordinates, with size reflecting node degree (number of linked significant correlations). Links between nodes display positive (aqua color) and negative (red color) correlations (p ≤ 0.05).

Problem behaviors displayed few significant associations in CPP_{class} values: increasing PC1 of problem behaviors associated with increased potential metabolism of lipids (M), amino sugars (F), other demethylated compounds (M), and near vitamin K2-vitamin B12 compounds (F); but decreased potential metabolism of carbohydrates (M). For $CPP_{density}$ values, patterns were found in males only: increasing PC1 of problem behaviors associated with increased potential metabolism of vitamin B6 (M) and vitamin B9 (M); but decreased potential metabolism of acetate (M), vitamin B12 (M) and glutamate (M). For CPP_{ASALR} values, the comparison was made between groups of high PC1 versus low PC1 of problem behaviors (for consistent display with other case studies in Fig. 3). High PC1 values associated with increased potential metabolism of other methylated compounds (F); and decreased potential metabolism of carbohydrates (M). Weighted mean vK-coordinates between high PC1 and low PC1 of problem behaviors showed a difference in beta-dispersion within condensed aromatics for females only. To explore this dataset in more detail, we identified differentially abundant functions and vK-coordinates (i.e., aggregated functional relative abundances via bioenergetic mapping) in high PC1 versus low PC1 subjects, separately within each sex. In females, 22 differentially abundant functions, compared to only 2-vK coordinates (with 3 corresponding functions), were identified (Figs. S22, S24, Table S18, S20). In males, 6 functions compared to 2 vK-coordinates (with 8 corresponding functions) were identified (Figs. S23, S25, Table S19, S21). Not all functions could be mapped into vK coordinate space. Interestingly, there was no overlap in functions identified directly versus indirectly (from aggregation into vK-coordinates). This means that differential abundance analysis using vK-coordinates can provide entirely different focal areas for investigation compared to the standard function-level analysis. From the vK-coordinate level analysis, high PC1 (compared to low PC1) females exhibited increased potential metabolism of Uridine phosphorylase (EC 2.4.2.3) (fxn_14491) involved in pyrimidine conversions; and decreased aldehyde lyases dihydroneopterin phosphate phosphatase and dihydroneopterin aldolase (EC 4.1.2.25) (fxn_12938; fxn_12942). High PC1 males exhibited increased phosphoenolpyruvate carboxykinase (GTP) (EC 4.1.1.32) (fxn_2926) associated with pyruvate metabolism; and also increases in multiple functions (fxn_821; fxn_2958; fxn_2973;

fxn_12703; fxn_12705; fxn_12786; fxn_12788) all involving alcohol dehydrogenase (EC 1.1.1.1) and

acetaldehyde dehydrogenase (EC 1.2.1.10), with or without pyruvate-formate-lyase deactivase – involved in

degradation of aromatic compounds, biphenyl, tryptophan, and pyruvate metabolism. Results and

visualizations for the standard function-level analysis are included for comparison but are not discussed

further.

3.2. Plant-soil systems

We found remarkable consistency in many observed patterns across the environmental soil case studies.

Results reported here are for relative trends with increasing ecosystem maturity (i.e., older revegetation and

natural samples) in CPP_{class} (Figs. S26, S30, S33, Table S3), $CPP_{density}$ (Figs. S27-S28, S31-S32, S34-35,

Table S4), and CPP_{ASALR} (Figure 5, Table S5). Below, we highlight trends found in at least two of three

case studies.

In CPP_{class} data we observed: increased potential metabolism of lipids (post-mining, AMI) and

condensed aromatics (post-mining, AMI); but decreased potential metabolism of proteins (post-mining,

AMI), carbohydrates (post-mining, AMI, with marginal indications in both PCaN soil groups), lignin (post-

mining, AMI), other methylated compounds (post-mining, AMI), and other hydrated compounds (AMI,

PCaN acidic-neutral soils). Mixed or isolated results included potential metabolism: either decreased (AMI)

or increased (PCaN strongly acidic soils) for tannins; decreased for other demethylated compounds (AMI);

increased for other condensed compounds (AMI); and decreased (post-mining, PCaN acidic-neutral soils)

or increased (PCaN strongly acidic soils) for near vitamin K2-vitamin B12 compounds. Total CPP_{class}

compounds appeared to be less well characterized and mapped to functional reactions in AMI natural

compared to disturbed samples, but more well characterized in older revegetation (compared to younger

revegetation) within the PCaN strongly acidic soils.

CPP_{ASALR} results reinforced many patterns observed in the CPP_{class} data: we observed increased

potential metabolism of lipids (post-mining, AMI, marginal in PCaN strongly acidic soils) and condensed

aromatics (post-mining, AMI); but decreased potential metabolism of proteins (AMI, marginal in post-

mining), carbohydrates (post-mining, AMI, PCaN acidic-neutral soils), and other methylated compounds

(post-mining, AMI). Isolated results included, potential metabolism: increased for other condensed

compounds (AMI); but decreased for lignin (AMI), tannins (AMI), other hydrated compounds (AMI), and

other demethylated compounds (AMI). Differing from CPP_{class} results, for CPP_{ASALR} potential metabolism

of near vitamin K2-vitamin B12 compounds increased (post-mining).

$CPP_{density}$ results were also quite consistent across case studies: we observed increased potential

metabolism of vitamin B9 (AMI, marginal in post-mining and PCaN strongly acidic soils) and vitamin K2

(post-mining, AMI, marginal in PCaN strongly acidic); but decreased potential metabolism of acetate (post-

mining, AMI, PCaN acidic-neutral soils), propionate (AMI), vitamin B2 (post-mining, marginal in PCaN

strongly acidic soils), vitamin B12 (post-mining, AMI, PCaN strongly acidic soils), vitamin B6 (post-

mining, AMI), glutamate (post-mining, AMI, PCaN acidic-neutral soils), and pyruvate (AMI). While

butyrate $CPP_{density}$ values were low and indistinguishable across all samples using 0.05 vK unit radial

buffers, for an example comparison we tested near butyrate $CPP_{density}$ using a larger 0.1 vK unit buffer in the

AMI soils and found increased levels in natural compared to disturbed soils (Fig. S36).

Post-mining and AMI samples displayed significant and mostly consistent directional shifts in

weighted mean vK-coordinate centroids across all compound classes (vK mapping zones) considered, while

PCaN acidic to neutral soils also displayed shifts within carbohydrates and condensed aromatics (Fig. 6,

Fig. S29, Tables S13-S16). Ecosystem maturity (/quality) explained 60–92% of the variation in weighted

mean vK-coordinates in post-mining soils (Fig. 6a, Table S15). The following general patterns emerged

with increasing ecosystem maturity: lipids became more reduced (lower oxygen content), proteins became

more hydrated, amino sugars became more dehydrogenated or condensed, carbohydrates became more

condensed (post-mining, PCaN acidic-neutral) or reduced (AMI), condensed aromatics became more

condensed, lignin showed mixed trends (dehydrogenation in post-mining, reduction in AMI), and tannins

became more demethylated. Except for carbohydrates and amino sugars, these trends largely represented an

outward extension of sample profile mapping into vK coordinate space with older ecosystems.

Fig. 5. Amino sugar-adjusted log ratio compound processing potential (CPP$_{ASALR}$), representing putative

microbiota activity-normalized values, from People Cities and Nature (PCaN) urban restoration, post-

mining restoration soils, and Australian Microbiome Initiative (AMI) disturbed versus natural soils. Linear

trends were used for visualization purposes. However, Kendall's tau correlation tests (suited to ordinal data) were applied. Sample sizes are detailed in Table 1.

Fig. 6. Weighted-mean van Krevelen coordinates within select compound classes display significant shifts with maturity of plant-soil ecosystems. Patterns are from (a) post-mining forest ecosystem restoration soil samples (n = age-based groups of 3), and (b) AMI disturbed (n = 29) vs natural (n = 55) soil samples (b). PERMANOVA and beta-dispersion results are in Table S15-S16.

3.3. Consistency in bioenergetic 'topography' mapping

CPP$_{density}$ plots visualizing local polynomial regression fitting ('loess') smoothed vK mapping profiles for

increasing radii buffer areas displayed striking consistency of form within each compound class, with

alternative expressions in either gut (Figs. S5, S9, S13, S19) or soils (Figs. S27, S31, S34).

4. Discussion

We show a meaningful bioenergetic basis to the development and differentiation of community-scale

microbiomes. Our CPP metrics quantified putative shifts in a microbial community's relative proficiency to

process different types of compounds across gradients of human- and environmental-health. We found

significant patterns of CPP association at varying resolutions: within major classes of compounds, near

focus biomolecules, and for unique vK coordinates. In the environmental soil metagenomes, we observed

surprising consistency in CPP profiles, which suggests they may link to plant-soil system conditions in

coherent and predictable ways. Our findings align with the notion that microbiota are shaped by the

bioenergetic status of prevailing substrates and micro-environments, and this information is simultaneously

recorded in their metagenomes. From a methodological perspective, our compound-focused bioenergetic

mapping approach demonstrates new pathways for assessing and interpreting microbial systems, capable of

supporting ongoing efforts to define healthy microbiomes. For example, aggregating functions via vK-

coordinates can provide entirely different focal areas for investigation compared to standard function-level

analyses.

4.1. Patterns in human health

4.1.1. Atherosclerotic cardiovascular disease (ACVD)

We found strong links in both sexes between ACVD and increased potential gut microbial metabolism of

lipids, condensed aromatics, lignin, other hydrated compounds, vitamin B9 and vitamin K2; and decreased

potential metabolism of carbohydrates and glutamate. Globally, ACVD is the leading cause of death, with

established treatment involving lipid-lowering statins, while human gene editing approaches to disrupt lipid

processing are being trialed (Li and Wu, 2022; Vaduganathan et al., 2022).

Links between high-fat diets and ACVD are well established (Law, 2000). Gut microbiota can

contribute to ACVD by metabolizing the dietary lipid phosphatidylcholine, with subsequent production of

harmful trimethylamine oxide (Albenberg and Wu, 2014). Polycyclic aromatic hydrocarbons are known

risk factors in ACVD (Burstyn et al., 2005; Mallah et al., 2021). Lignin (different to lignan) is a complex

and ubiquitous structural plant polymer, considered predominantly insoluble fiber, with content ranging

from 1-2 g/100g in vegetables, fruits and cereals, up to 30-40 g/100g in nut shells and stone fruit kernels

(Tao et al., 2020). Lignin inhibits the enzymatic activity of α-glucosidase, delaying carbohydrate digestion

and absorption, with potential for low post-meal blood sugar levels (Chen et al., 2015). Acute hypoglycemia

(low blood sugar) can trigger cardiac events (Frier et al., 2011), with greater adverse risks in subjects with

significant comorbidities (e.g., T2D, ACVD). Our finding for decreased carbohydrates in ACVD might

align with these impacts on blood sugar. Alternatively, we speculate that result might be symptomatic of a

more Westernized diet (high in animal protein, sugar, starch, and fat – and lower in carbohydrate content

than a plant-rich diet; Albenberg and Wu, 2014).

Excessive vitamin B9 (folate) is associated with ACVD risk via a non-linear u-shaped dose-

response relationship (Xu et al., 2023). Non-linear u-shaped dose-response relationships are common in

biological systems (i.e., hormesis, or deficiency-sufficiency-toxicity) (Calabrese et al., 2007). Similarly, our

results linking increased vitamin K2 with ACVD appear contrary to recent opinion (Essa et al., 2021),

although non-linear u-shaped dose-responses have also been observed (Bellinge et al., 2021). Vitamin K2 is

commonly found in fermented foods which are less common in Western diets (McFarlin et al., 2017). Here,

the ACVD case study was based on Chinese subjects whose diets potentially contained higher quantities of

fermented foods including vitamin K2. Possibly, these subjects were more susceptible to adverse effects if

excessive levels of vitamin K2 were reached.

Our results linking reduced glutamate with ACVD appear contrary to findings from large US cohort

studies which found higher glutamate levels, and lower glutamine:glutamate ratios, correlated with

increased ACVD risk (Ma et al., 2017; Zheng et al., 2016). Dietary proteins are a major source of glutamate

(Loï and Cynober, 2022). However, some ethnic populations may have inadequate protein in their diets (Liu

et al., 2013). Interestingly, in female adults from rural western China with inadequate (and largely plant-

derived) protein intake, increasing animal protein associated with reduced risk of hypertension (Liu et al.,

2013). Together, these observations suggest that a u-shaped dose-response may also operate for animal-

based proteins, glutamate, and ACVD risk.

4.1.2.   Colorectal cancer

In the colorectal cancer subjects (from France) we found consistent associations across CPP metrics in both

sexes with increased potential metabolism of amino sugars, vitamin B12 and B9; and decreased potential

metabolism of proteins and tannins. Colon cancer has been associated with low dietary fiber, low fruit and

vegetable consumption, and high red meat consumption (Law, 2000). Our results were consistent with

reports for anticancer activity, including protective effects against colorectal cancer, from some tannins or

polyphenols (e.g., components in green and black tea, resveratrol in red wine and grapes) (Alam et al.,

2018). Unfortunately, our data did not distinguish between animal- and plant-based protein. However, red

meat is widely consumed in France, with 41% of males and 24% of females consuming above guideline

levels (Mota et al., 2021). Amino sugars are sugar molecules with at least one hydroxyl group substituted

by an amino group. In biological systems, they are formed by catalytic activity acting on amino acids

(glutamate, glutamine – building blocks of protein) to transfer an amino functionality to a sugar phosphate

or sugar nucleotide (Skarbek and Milewska, 2016). Therefore, both glucose (sugar) and amino acids

contribute to amino sugar formation. Meanwhile, metabolism of both glucose and amino acids plays a key

role in colorectal cancer development (Zhang et al., 2023). Possibly, our finding of increased potential

metabolism of amino sugars with colorectal cancer may reflect dysregulated activity of glucose and amino

acids with the product of their interaction (amino sugars) recorded by the gut microbiome. Consistent with

our findings, vitamin B9 (folate or folic acid) and vitamin B12 supplementation have been associated with

increased risk of colorectal cancer (Oliai Araghi et al., 2019).

4.1.3.   Type 2 diabetes (T2D)

In the T2D case study, our lack of clear findings linked to major compound classes or focus biomolecules

seems consistent with reports that T2D is a complex, multifaceted, highly heterogeneous polygenic disease

with uncertain etiology (Pearson, 2019). We found untreated (Met-) T2D exhibited an anomalous and

complex CPP network, including a high number of negative correlations (indicating negative feedbacks)

ranging widely across vK coordinate space (i.e., covering a spectrum of compounds and bioenergetic

status). Diagnosis of T2D is based on elevated blood glucose, primarily arising from insulin resistance and

inadequate insulin secretion (Forslund et al., 2015). However, a clear diagnostic test for T2D is lacking,

except by exclusion of other causes (Pearson, 2019). A range of factors including genetics, dietary habits,

sedentary lifestyle, and gut microbiota are involved in disease development (Forslund et al., 2015).

Possibly, with more detailed examination, diagnostic relationships (e.g., correlations, ratios) might be

uncovered in relative abundance patterns of compound-associated vK-coordinates underpinning the

anomalous T2D Met- network.

4.1.4.   Problem behaviors

In the problem behavior case study, different compound associations were observed in female and male

children. Increased problem behavior (higher PC1) in females associated with increased potential

metabolism of amino sugars, other methylated compounds, and near vitamin K2-vitamin B12 compounds.

From differential abundance analysis, high PC1 females exhibited decreased dihydroneopterin aldolase—an

enzyme involved in converting dihydroneopterin (a molecule involved in folate biosynthesis) into other

compounds (Deng et al., 2000). Excessive serum levels of dihydroneopterin have been associated with

major depression (Kusunoki et al., 1999). Possibly, in case study subjects, reduced levels of

dihydroneopterin-degrading enzyme have promoted accumulation of dihydroneopterin in association with

problem behaviors. High PC1 females also exhibited increased uridine phosphorylase, an enzyme involved

in pyrimidine metabolism that converts uridine to uracil (Cao and Pizzorno, 2004), therefore possibly

degrading uridine levels in those subjects. Uridine is linked to energy metabolism and glutamate-mediated

excitatory neurotransmission in the brain, and supplemental uridine treatments have been used to reduce

depressive symptoms in adolescents (Douglas G. Kondo, 2011).

In males, increased PC1 associated with increased potential metabolism of lipids, vitamins B6 and

B9; and decreased potential metabolism of carbohydrates, acetate, vitamin B12, and glutamate. Vitamins

B12 (cobalamin) and B9 (folate) are recognized precursors involved in forming key neurotransmitters

dopamine, noradrenaline (norepinephrine), and serotonin (Hutto, 1997). These three neurotransmitters

occur in the vicinity of vitamin B12 and K2 in vK coordinate space. Vitamin B12 deficiency has been

associated with depressive disorders in older subjects (Henning Tiemeier et al., 2002). Glutamate is the

major excitatory neurotransmitter of the healthy mammalian brain, and an abundant free amino acid

important in multiple metabolic pathways, which requires regulation at optimal levels in extracellular fluids

(Zhou and Danbolt, 2014). High PC1 males also exhibited increased phosphoenolpyruvate carboxykinase—

an enzyme involved in cataplerosis, or removal of intermediate 4- and 5-carbon compounds from the TCA

cycle (Owen et al., 2002; Yang et al., 2009). These intermediates are removed because they cannot be fully

oxidized for energy metabolism within the TCA cycle, but are converted elsewhere to glucose, fatty acids

or amino acids (Owen et al., 2002). High PC1 males also exhibited increases in alcohol dehydrogenase,

acetaldehyde dehydrogenase, and pyruvate-formate-lyase deactivase, variously involved in pyruvate

metabolism, degradation of aromatics and biphenyl, and tryptophan catabolism. Key processes of energy

metabolism involving glucose, lipids, protein and the TCA cycle (via keystone molecules pyruvate, acetyl-

CoA, and glutamate) have been implicated in major depressive disorder, although precise pathways of

pathogenesis are still unclear (Gu et al., 2021).

4.2. Patterns in plant-soil systems

Our findings point to generalizable patterns with older ecosystems for increasing microbial CPP associated with lipids, condensed aromatics, vitamin B9, and vitamin K2; and decreasing CPP associated with proteins, carbohydrates, lignin, other methylated compounds, other hydrated compounds, acetate, vitamin B12, vitamin B6, and glutamate. Drivers of shifting CPP in soils are expected to include changing: (1) composition of biota and biotic materials including plants, organic debris, and re-assembly of invertebrate and microbial communities, and (2) soil abiotic conditions due to plant-soil feedbacks (e.g., pH, nutrients, organic carbon content, temperature, moisture regime) (van der Putten et al., 2016; Wardle et al., 2004). This includes macro-environmental influences with development of vegetation structure and canopy cover (e.g., shading, rainfall interception, altered drainage). CPP values also likely reflect a dynamic balance between resource availability and use by microbiota.

For example, we might expect greater accumulation of lignin in soils of older ecosystems due to plant inputs such as dead roots, bark, leaf litter, and other structural plant residues. However, we observed reduced CPP for lignin in these sample types. Fungi are major lignin degraders (Janusz et al., 2017) and fungal communities vary with ecosystem disturbance and abiotic conditions (Rodriguez-Ramos et al., 2021). Interestingly, our results were counter to expectations for elevated fungal decomposition of lignin in older ecosystems. Reforestation with native mixed-species can produce higher levels of recalcitrant soil organic matter (Cunningham et al., 2015) (e.g., humic acid which is hard to decompose and maps to lignin in vK space). Our CPP metrics are relative and compositional (based on functional relative abundances summing to a maximum of 100%), so it may be that in relative terms, the metabolic foci of microbiota are shifted to processing other materials. Or possibly, structural plant materials may be more accessible for degradation in disturbed (e.g., agricultural) soil environments, depending on plant residue management, nutrient availability and other factors.

We expect some CPP quantities are driven primarily by plant material inputs. For example, soils from more mature ecosystems in temperate climates, represented in samples from AMI and post-mining (in

the Appalachian Plateau, southwestern Virginia USA; Avera et al., 2015; Sun and Badgley, 2019),

displayed a positive relationship with CPP for lipids and condensed aromatics. These two compound classes

are represented in plant-based essential oils and volatile, aromatic organic compounds. Oils are found in

high densities in much of the fire-adapted Australian flora (unlike New Zealand flora) (Bowman et al.,

2012). Increased CPP for lipids might also arise due to increased density of energy storage linked to

primary production, or more active plant signaling in response to abiotic stress (Hou et al., 2016). High

levels of lipids and condensed aromatics in mature ecosystem soils could also be a result of increasing plant

investment into defensive mechanisms via antimicrobial essential oils (Hammer et al., 1999), and volatile

and aromatic secondary defense compounds induced by herbivory (typically by invertebrates) (Erb and

Kliebenstein, 2020).

Shifting weighted mean vK coordinates across many compound classes (in AMI and post-mining)

suggests broad changes in the composition of microbial substrates with more mature ecosystems. The

changing composition of the microbiota itself may contribute to this. Carbohydrates are of interest due to

the potential contribution of plant-based material to human diet, and CPP_{class} values for carbohydrates were

consistently assigned the largest sum of functional relative abundances in the human gut samples. With

more mature ecosystems, CPP for carbohydrates decreased in relative terms, but weighted vK coordinates

suggest carbohydrate CPP shifts towards favoring processing materials with reduced oxygen content per

unit of carbon. There is likely to be global variation in environmental soil CPP driven by soil abiotic factors

and changing biota (vegetation and animals), previously outlined.

4.3. Potential environment-human health links

This work opens new avenues for investigating environment-human health connections as environments

vary in their production of human health-associated compounds. Moreover, varying environmental

microbiota exposures may supply modulating CPP profiles for colonizing or transient impacts to human

microbiomes (e.g., skin, airway, gut), which are intimately linked to our health. We show CPP patterns

imprinted in environmental soil metagenomes are linked with the maturity (or quality) of plant-soil systems

and abiotic factors such as soil pH. We demonstrate that CPP 'landscapes' (or the range of meta-

compounds) involved in functions that can be performed by soil and gut metagenomes show a large degree

of overlap with similar distributions of functional relative abundances across vK space (Fig. S2). We also

show a similar spectrum of CPP measures that are impacted by ecosystem quality (e.g., increased potential

metabolism of lipids, condensed aromatics, vitamin B9, vitamin K2; and decreased potential metabolism of

proteins, carbohydrates, lignin, acetate, vitamin B12, vitamin B6, and glutamate) are also variously linked

to human health and disease. However, we urge caution in attempting to directly translate CPP trends in

plant-soil environments to infer possible implications for human health. We stress that non-linear, u-shaped

dose-response relationships, as discussed earlier (Calabrese et al., 2007), are common and relevant in the

context of dietary and environmental exposure-human health links. With potential for u-shaped dose-

response effects (i.e., deficiency, sufficiency, toxicity), we might expect environmental microbiota CPP to

have varying influences in different geographic/cultural populations. Also, the gut represents a more tightly

controlled micro-environment (redox, pH, etc.) unsuited to many environmental microbes.

Example evidence for potential environment-human transfer of microbial CPP comes from

Endomicrobia species found in oral microbiota of indigenous peoples from central Australia (Handsley-

Davis et al., 2022). Endomicrobia species provide energetic advantage for cellulose digestion in the guts of

termites and wood-eating insects—and transfer to humans has occurred likely through use of termites and

termite mounds in traditional food and medicine (Handsley-Davis et al., 2022). Speculatively, our results

suggest that if broad supplementation of human microbiota CPP capacity is required (spanning a range of

health-supporting biomolecules), this may require exposure to multiple types of environments.

Alternatively, to address particular microbiota-associated disease states, specific environment types may

provide more targeted microbiota CPP supplementation. As knowledge builds of the varying proficiency of

environmental microbiota to process human health associated compounds in different qualities of

ecosystems, we might design environmental interventions or nature prescriptions to modulate human-

microbial exposures, thereby advancing microbiota-oriented approaches to human health.

4.4. Limitations

There are important limitations in this study. CPP metrics do not measure actual compounds; rather, they

quantify conceptual 'meta-compounds' or assemblies of elements based on function-level summary mean

O:C and H:C ratios consistent with compounds of interest. Quantification occurred via mapping into vK

space and aggregating functional relative abundances into major compound classes, near focus

biomolecules, or at unique vK coordinates, to assess CPP structural profiles of metagenomes. Conceptually,

mean function-level attributes (O:C and H:C ratios) represent a mid-point of chemical transformation

mediated by microbiota (i.e., interpreted as 'X% of functions were involved in processing

compound/biomolecule type Y'). Stoichiometry rules determine that reaction inputs and products will have

balanced O, H, and C atomic counts. However, different mean O:C and H:C ratios can arise due to

uncounted O and H atoms in non-C containing species (O:C, H:C values become undefined). For future

work, the CPP mapping algorithm could be readily adjusted to separately target reaction inputs, or products,

or individual chemical species.

Our coarse compound classes did not distinguish (for example) plant versus animal proteins or high-

fiber versus low-fiber carbohydrates. Finer-resolution vK mapping zones would increase the precision of

results. We could not discern CPP differences for butyrate between sample types using 0.05 vK radii.

Butyrate is often present at low concentrations in the gut compared to other SCFAs, with rapid consumption

by colonocytes (den Besten et al., 2013). The volatility of butyrate (Wagner et al., 2017) may make it

susceptible to loss from soils. Our butyrate $CPP_{density}$ profiles spanning large to small vK radii may depict

source-sink dynamics. Compound mapping using O, H, and C content enable exhaustive and

compartmented mapping within vK coordinate space, however this represents a simplified, imperfect

approach. Multi-element compound mapping would offer increased precision (Rivas-Ubach et al., 2018)

and may be developed to provide exhaustive and compartmented mapping across a range of compound

types. CPP measures used here were relative, not absolute. Mapping of SUPER-FOCUS functions was

incomplete (sample functional relative abundances ranged from 52-84%, with means 55-67%). This may be

improved with future algorithm refinement.

Further work is required to refine our first-pass ASALR normalization, to examine CPP-disease

links in wider geographic/cultural populations, and to explore other potential explanatory variables not

considered in our analyses. Like many microbiome studies, our analyses do not permit causal insight to

interpret whether increased or decreased CPP may facilitate or follow disease. For example, excessive CPP

may produce metabolites at toxic levels, or degrade substrates leading to deficiency. Reduced CPP

measures might correspond to dietary deficiencies or suppression of functional pathways due to

dysregulated environmental conditions. Nonetheless, observed CPP trends may assist hypothesis-building

and prioritizing mechanistic research. We suggest that future work might address these limitations.

5. Conclusions

Our assessment of microbial metagenomes using the compound processing potential framework offers a

new approach to examine potential overlapping functional capacities between human and environmental

microbiomes. Our findings reveal a meaningful bioenergetic basis to the development and differentiation of

microbiomes in human health and disease, and with variation in ecosystem quality. We show that

ecosystem quality can shape environmental microbiota in seemingly predictable ways, with varying

proficiency to process a range of compounds that are associated with human health and disease—as

measured by the functional potential relative abundances imprinted in their collective DNA. This work

informs the ongoing development of microbiota-oriented approaches in human health, including the

potential for nature-based solutions to help address global healthcare challenges.

Data availability

All data used in this study are available from repositories listed in Table 1. Code used to support this study

is available from: https://github.com/liddic/compound_potential

Acknowledgments

We acknowledge the authors of previously published metagenomics datasets used here which enabled

comparison of diverse samples spanning human and environmental health. We acknowledge First Nations

Peoples and traditional custodians of the lands from which the case study datasets originate. This work was

supported by the People, Cities and Nature research program (Aotearoa New Zealand Ministry of Business,

Innovation and Employment, grant UOWX2101), Marsden Fund Fast-Start grant funding (Royal Society Te

Apārangi), funding from the German Research Foundation iDiv ([DFG]–FZT 118, 202548816; Ei 862/29-

1; Ei 862/31-1; Ei 862/27-1; HE 8266/4-1), and the Australian Research Council (grant numbers

LP190100051, LP190100484). We acknowledge the contribution of the Australian Microbiome consortium

in the generation of data used in this publication. The Australian Microbiome Initiative is supported by

funding from Bioplatforms Australia and the Integrated Marine Observing System (IMOS) through the

Australian Government's National Collaborative Research Infrastructure Strategy (NCRIS), Parks Australia

through the Bush Blitz program funded by the Australian Government and BHP, and the CSIRO.

References

Alam MN, Almoyad M, Huq F. Polyphenols in Colorectal Cancer: Current State of Knowledge including Clinical Trials and Molecular Mechanism of Action. BioMed Research International 2018; 2018: 4154185.

Albenberg LG, Wu GD. Diet and the Intestinal Microbiome: Associations, Functions, and Implications for Health and Disease. Gastroenterology 2014; 146: 1564-1572.

Andrews S. FastQC - a quality control application for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc, 2018.

Avera BN, Strahm BD, Burger JA, Zipper CE. Development of ecosystem structure and function on reforested surface-mined lands in the Central Appalachian Coal Basin of the United States. New Forests 2015; 46: 683-702.

Averill C, Anthony MA, Baldrian P, Finkbeiner F, Hoogen Jvd, Kiers T, et al. Defending Earth's terrestrial microbiome. Nature Microbiology 2022; 7: 1717-1725.

Banerjee S, van der Heijden MGA. Soil microbiomes and one health. Nature Reviews Microbiology 2022.

Barnes A, Eisenhauer N, Wallace KJ, Rai B, Mitchell G, Heintz-Buschart A. People Cities and Nature (PCaN) pilot metagenomics dataset. In: Waikato Uo, editor, Aotearoa New Zealand, 2020.

Barton LL, Northup DE. Microbial Ecology: Wiley-Blackwell, 2011.

Belizario JE, Napolitano M. Human microbiomes and their roles in dysbiosis, common diseases, and novel therapeutic approaches. Frontiers in Microbiology 2015; 6.

Bellinge JW, Dalgaard F, Murray K, Connolly E, Blekkenhorst LC, Bondonno CP, et al. Vitamin K Intake and Atherosclerotic Cardiovascular Disease in the Danish Diet Cancer and Health Study. Journal of the American Heart Association 2021; 10: e020551.

Bissett A, Fitzgerald A, Meintjes T, Mele PM, Reith F, Dennis PG, et al. Introducing BASE: the Biomes of Australian Soil Environments soil microbial diversity database. GigaScience 2016; 5: 21.

Bowman DMJS, Murphy BP, Burrows GE, Crisp MD. Fire regimes and the evolution of the Australian biota. CSIRO Publishing, Collingwood, Vic, 2012, pp. 27-47.

Brame JE, Liddicoat C, Abbott CA, Breed MF. The potential of outdoor environments to supply beneficial butyrate-producing bacteria to humans. Science of The Total Environment 2021; 777: 146063.

Buchfink B, Reuter K, Drost H-G. Sensitive protein alignments at tree-of-life scale using DIAMOND. Nature Methods 2021; 18: 366-368.

Burstyn I, Kromhout H, Partanen T, Svane O, Langård S, Ahrens W, et al. Polycyclic Aromatic Hydrocarbons and Fatal Ischemic Heart Disease. Epidemiology 2005; 16: 744-750.

Calabrese EJ, Bachmann KA, Bailer AJ, Bolger PM, Borak J, Cai L, et al. Biological stress response terminology: Integrating the concepts of adaptive response and preconditioning stress within a hormetic dose–response framework. Toxicology and Applied Pharmacology 2007; 222: 122-128.

Cao D, Pizzorno G. Uridine phosophorylase: an important enzyme in pyrimidine metabolism and fluoropyrimidine activation. Drugs of Today 2004; 40: 431-443.

Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 2018; 34: i884-i890.

Chen Y, Liu Y, Li X, Zhang J, Li G. Lignin Interacting with α-glucosidase and its Inhibitory Effect on the Enzymatic Activity. Food Biophysics 2015; 10: 264-272.

Cunningham SC, Cavagnaro TR, Mac Nally R, Paul KI, Baker PJ, Beringer J, et al. Reforestation with native mixed-species plantings in a temperate continental climate effectively sequesters and stabilizes carbon within decades. Global Change Biology 2015; 21: 1552-1566.

D'Andrilli J, Cooper WT, Foreman CM, Marshall AG. An ultrahigh-resolution mass spectrometry index to estimate natural organic matter lability. Rapid Communications in Mass Spectrometry 2015; 29: 2385-2401.

Daisley BA, Koenig D, Engelbrecht K, Doney L, Hards K, Al KF, et al. Emerging connections between gut microbiome bioenergetics and chronic metabolic diseases. Cell Reports 2021; 37: 110087.

Delgado-Baquerizo M, Reich PB, Trivedi C, Eldridge DJ, Abades S, Alfaro FD, et al. Multiple elements of soil biodiversity drive ecosystem functions across biomes. Nature Ecology & Evolution 2020; 4: 210-220.

Delgado-Baquerizo M, Reith F, Dennis PG, Hamonts K, Powell JR, Young A, et al. Ecological drivers of soil microbial diversity and soil biological networks in the Southern Hemisphere. Ecology 2018; 99: 583-596.

den Besten G, van Eunen K, Groen AK, Venema K, Reijngoud D-J, Bakker BM. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. Journal of Lipid Research 2013; 54: 2325-2340.

Deng H, Callender R, Dale GE. A Vibrational Structure of 7,8-Dihydrobiopterin Bound to Dihydroneopterin Aldolase. Journal of Biological Chemistry 2000; 275: 30139-30143.

Diener C, Gibbons SM, Resendis-Antonio O. MICOM: Metagenome-Scale Modeling To Infer Metabolic Interactions in the Gut Microbiota. mSystems 2020; 5: 10.1128/msystems.00606-19.

Douglas G. Kondo Y-HS, Tracy L. Hellem, Kristen K. Delmastro, Eun-Kee Jeong, Namkug Kim, Xianfeng Shi, and Perry F. Renshaw. Open-Label Uridine for Treatment of Depressed Adolescents with Bipolar Disorder. Journal of Child and Adolescent Psychopharmacology 2011; 21: 171-175.

Eisenstein M. The hunt for a healthy microbiome. Nature 2020; 577: S6-S8.

Erb M, Kliebenstein DJ. Plant Secondary Metabolites as Defenses, Regulators, and Primary Metabolites: The Blurred Functional Trichotomy. Plant Physiology 2020; 184: 39-52.

Essa H, Nicholas K, Jean-Pierre I, Leon JS, Anas S, Omar A, et al. Vitamin K2—a neglected player in cardiovascular health: a narrative review. Open Heart 2021; 8: e001715.

Fierer N. Embracing the unknown: Disentangling the complexities of the soil microbiome. Nat Rev Microbiol 2017; 15: 579-590.

Flandroy L, Poutahidis T, Berg G, Clarke G, Dao M-C, Decaestecker E, et al. The impact of human activities and lifestyles on the interlinked microbiota and health of humans and of ecosystems. Science of The Total Environment 2018; 627: 1018-1038.

Flannery JE, Stagaman K, Burns AR, Hickey RJ, Roos LE, Giuliano RJ, et al. Gut Feelings Begin in Childhood: the Gut Metagenome Correlates with Early Environment, Caregiving, and Behavior. mBio 2020; 11: e02780-19.

Flinders_University. DeepThought (HPC). Retrieved from https://doi.org/10.25957/FLINDERS.HPC.DEEPTHOUGHT, 2021.

Forslund K, Hildebrand F, Nielsen T, Falony G, Le Chatelier E, Sunagawa S, et al. Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. Nature 2015; 528: 262-266.

Frier BM, Schernthaner G, Heller SR. Hypoglycemia and Cardiovascular Risks. Diabetes Care 2011; 34: S132-S137.

Garza DR, van Verk MC, Huynen MA, Dutilh BE. Towards predicting the environmental metabolome from metagenomics with a mechanistic model. Nature Microbiology 2018; 3: 456-460.

Gilbert JA, Blaser MJ, Caporaso JG, Jansson JK, Lynch SV, Knight R. Current understanding of the human microbiome. Nature Medicine 2018; 24: 392.

Gu X, Ke S, Wang Q, Zhuang T, Xia C, Xu Y, et al. Energy metabolism in major depressive disorder: Recent advances from omics technologies and imaging. Biomedicine & Pharmacotherapy 2021; 141: 111869.

Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. Journal of Applied Microbiology 1999; 86: 985-990.

Handsley-Davis M, Kapellas K, Jamieson LM, Hedges J, Skelly E, Kaidonis J, et al. Heritage-specific oral microbiota in Indigenous Australian dental calculus. Evolution, Medicine, and Public Health 2022; 10: 352-362.

Henning Tiemeier, M.D. ,, H. Ruud van Tuijl, M.Sc. ,, Albert Hofman, M.D., Ph.D. ,, John Meijer, M.Sc. ,, Amanda J. Kiliaan, Ph.D. , and, Monique M.B. Breteler, M.D., Ph.D. Vitamin B12, Folate, and

Homocysteine in Depression: The Rotterdam Study. American Journal of Psychiatry 2002; 159:
2099-2101.
Hinsinger P, Bengough AG, Vetterlein D, Young IM. Rhizosphere: biophysics, biogeochemistry and
ecological relevance. Plant and Soil 2009; 321: 117-152.
Hou Q, Ufer G, Bartels D. Lipid signalling in plant responses to abiotic stress. Plant, Cell & Environment
2016; 39: 1029-1048.
Hu Y, Zheng Q, Zhang S, Noll L, Wanek W. Significant release and microbial utilization of amino sugars
and d-amino acid enantiomers from microbial cell wall decomposition in soils. Soil Biology and
Biochemistry 2018; 123: 115-125.
Husson O. Redox potential (Eh) and pH as drivers of soil/plant/microorganism systems: a transdisciplinary
overview pointing to integrative opportunities for agronomy. Plant and Soil 2013; 362: 389-417.
Hutto BR. Folate and cobalamin in psychiatric illness. Comprehensive Psychiatry 1997; 38: 305-314.
Janusz G, Pawlik A, Sulej J, Świderska-Burek U, Jarosz-Wilkołazka A, Paszczyński A. Lignin degradation:
microorganisms, enzymes involved, genomes analysis and evolution. FEMS Microbiology Reviews
2017; 41: 941-962.
Jie Z, Xia H, Zhong S-L, Feng Q, Li S, Liang S, et al. The gut microbiome in atherosclerotic cardiovascular
disease. Nature Communications 2017; 8: 845.
Kusunoki K, Ozaki N, Sawada M, Sato T, Hirano S, Narita T. Serum. Levels of Dihydroneopterin and
Soluble Cytokine Receptors in Major Depression. Pteridines 1999; 10: 24-26.
Law M. Dietary fat and adult diseases and the implications for childhood nutrition: an epidemiologic
approach. The American Journal of Clinical Nutrition 2000; 72: 1291s-1296s.
Li N, Zhang H, Bai Z, Jiang H, Yang F, Sun X, et al. Soil exposure accelerates recovery of the gut
microbiota in antibiotic-treated mice. Environmental Microbiology Reports; n/a.
Li Z-F, Wu N-Q. The Progression of Treatment for Refractory Hypercholesterolemia: Focus on the
Prospect of Gene Therapy. Frontiers in Genetics 2022; 13.
Liddicoat C, Bi P, Waycott M, Glover J, Breed M, Weinstein P. Ambient soil cation exchange capacity
inversely associates with infectious and parasitic disease risk in regional Australia. Science of The
Total Environment 2018; 626: 117-125.
Liddicoat C, Sydnor H, Cando-Dumancela C, Dresken R, Liu J, Gellie NJC, et al. Naturally-diverse
airborne environmental microbial exposures modulate the gut microbiome and may provide
anxiolytic benefits in mice. Science of The Total Environment 2020; 701: 134684.
Litvak Y, Byndloss MX, Bäumler AJ. Colonocyte metabolism shapes the gut microbiota. Science 2018;
362: eaat9076.
Liu R, Dang S, Yan H, Wang D, Zhao Y, Li Q, et al. Association between dietary protein intake and the
risk of hypertension: a cross-sectional study from rural western China. Hypertension Research
2013; 36: 972-979.
Loï C, Cynober L. Glutamate: A Safe Nutrient, Not Just a Simple Additive. Annals of Nutrition and
Metabolism 2022; 78: 133-146.
Ma W, Heianza Y, Huang T, Wang T, Sun D, Zheng Y, et al. Dietary glutamine, glutamate and mortality:
two large prospective studies in US men and women. International Journal of Epidemiology 2017;
47: 311-320.
Mallah MA, Mallah MA, Liu Y, Xi H, Wang W, Feng F, et al. Relationship Between Polycyclic Aromatic
Hydrocarbons and Cardiovascular Diseases: A Systematic Review. Frontiers in Public Health 2021;
9.
Mallick H, Franzosa EA, McLver LJ, Banerjee S, Sirota-Madi A, Kostic AD, et al. Predictive metabolomic
profiling of microbial communities using amplicon or metagenomic sequences. Nature
Communications 2019; 10: 3136.
McFarlin BK, Henning AL, Venable AS. Oral Consumption of Vitamin K2 for 8 Weeks Associated With
Increased Maximal Cardiac Output During Exercise. Altern Ther Health Med 2017; 23: 26-32.

Mota JDO, Guillou S, Pierre F, Membré J-M. Public health risk-benefit assessment of red meat in France:
Current consumption and alternative scenarios. Food and Chemical Toxicology 2021; 149: 111994.
Neveu V, Nicolas G, Amara A, Salek RM, Scalbert A. The human microbial exposome: expanding the
Exposome-Explorer database with gut microbial metabolites. Scientific Reports 2023; 13: 1946.
Oliai Araghi S, Kiefte-de Jong JC, van Dijk SC, Swart KMA, van Laarhoven HW, van Schoor NM, et al.
Folic Acid and Vitamin B12 Supplementation and the Risk of Cancer: Long-term Follow-up of the
B Vitamins for the Prevention of Osteoporotic Fractures (B-PROOF) Trial. Cancer Epidemiology,
Biomarkers & Prevention 2019; 28: 275-282.
Ottman N, Ruokolainen L, Suomalainen A, Sinkko H, Karisola P, Lehtimäki J, et al. Soil exposure modifies
the gut microbiota and supports immune tolerance in a mouse model. Journal of Allergy and
Clinical Immunology 2018; 143: 1198-1206.e12.
Owen OE, Kalhan SC, Hanson RW. The Key Role of Anaplerosis and Cataplerosis for Citric Acid Cycle
Function. Journal of Biological Chemistry 2002; 277: 30409-30412.
Pearson ER. Type 2 diabetes: a multifaceted disease. Diabetologia 2019; 62: 1107-1112.
Rivas-Ubach A, Liu Y, Bianchi TS, Tolić N, Jansson C, Paša-Tolić L. Moving beyond the van Krevelen
Diagram: A New Stoichiometric Approach for Compound Classification in Organisms. Analytical
Chemistry 2018; 90: 6152-6160.
Rivera-Chávez F, Lopez CA, Bäumler AJ. Oxygen as a driver of gut dysbiosis. Free Radical Biology and
Medicine 2017; 105: 93-101.
Rodriguez-Ramos JC, Cale JA, Cahill Jr JF, Simard SW, Karst J, Erbilgin N. Changes in soil fungal
community composition depend on functional group and forest disturbance type. New Phytologist
2021; 229: 1105-1117.
Roslund MI, Parajuli A, Hui N, Puhakka R, Grönroos M, Soininen L, et al. A Placebo-controlled double-
blinded test of the biodiversity hypothesis of immune-mediated diseases: Environmental microbial
diversity elicits changes in cytokines and increase in T regulatory cells in young children.
Ecotoxicology and Environmental Safety 2022; 242: 113900.
Round JL, Mazmanian SK. The gut microbiome shapes intestinal immune responses during health and
disease. Nature Reviews. Immunology 2009; 9: 313-323.
Seaver SMD, Liu F, Zhang Q, Jeffryes J, Faria JP, Edirisinghe JN, et al. The ModelSEED Biochemistry
Database for the integration of metabolic annotations and the reconstruction, comparison and
analysis of metabolic models for plants, fungi and microbes. Nucleic Acids Research 2020; 49:
D575-D588.
Silva GGZ, Green KT, Dutilh BE, Edwards RA. SUPER-FOCUS: a tool for agile functional analysis of
shotgun metagenomic data. Bioinformatics 2015; 32: 354-361.
Skarbek K, Milewska MJ. Biosynthetic and synthetic access to amino sugars. Carbohydrate Research 2016;
434: 44-71.
Sokol NW, Slessarev E, Marschmann GL, Nicolas A, Blazewicz SJ, Brodie EL, et al. Life and death in the
soil microbiome: how ecological processes influence biogeochemistry. Nature Reviews
Microbiology 2022.
Sun S, Badgley BD. Changes in microbial functional genes within the soil metagenome during forest
ecosystem restoration. Soil Biology and Biochemistry 2019; 135: 163-172.
Sun X, Liddicoat C, Tiunov A, Wang B, Zhang Y, Lu C, et al. Harnessing soil biodiversity to promote
human health in cities. npj Urban Sustainability 2023; 3: 5.
Tao J, Li S, Ye F, Zhou Y, Lei L, Zhao G. Lignin – An underutilized, renewable and valuable material for
food industry. Critical Reviews in Food Science and Nutrition 2020; 60: 2011-2033.
Vaduganathan M, Mensah GA, Turco JV, Fuster V, Roth GA. The Global Burden of Cardiovascular
Diseases and Risk. Journal of the American College of Cardiology 2022; 80: 2361-2371.
van der Putten WH, Bradford MA, Pernilla Brinkman E, van de Voorde TFJ, Veen GF. Where, when and
how plant–soil feedback matters in a changing world. Functional Ecology 2016; 30: 1109-1121.

Wagner AO, Markt R, Puempel T, Illmer P, Insam H, Ebner C. Sample preparation, preservation, and
storage for volatile fatty acid quantification in biogas plants. Engineering in Life Sciences 2017; 17:
132-139.
Wardle DA, Bardgett RD, Klironomos JN, Heikki S, van der Putten WH, Wall DH. Ecological Linkages
Between Aboveground and Belowground Biota. Science 2004; 304: 1629-1633.
Wilson JW, Shakir D, Batie M, Frost M, Rocha S. Oxygen-sensing mechanisms in cells. The FEBS Journal
2020; 287: 3888-3906.
Wu X, Wu L, Liu Y, Zhang P, Li Q, Zhou J, et al. Microbial Interactions With Dissolved Organic Matter
Drive Carbon Dynamics and Community Succession. Frontiers in Microbiology 2018; 9.
Xu J, Zhu X, Guan G, Zhang Y, Hui R, Xing Y, et al. Non-linear associations of serum and red blood cell
folate with risk of cardiovascular and all-cause mortality in hypertensive adults. Hypertension
Research 2023; 46: 1504-1515.
Yang J, Kalhan SC, Hanson RW. What is the metabolic role of phosphoenolpyruvate carboxykinase? J Biol
Chem 2009; 284: 27025-9.
Zeller G, Tap J, Voigt AY, Sunagawa S, Kultima JR, Costea PI, et al. Potential of fecal microbiota for
early-stage detection of colorectal cancer. Molecular Systems Biology 2014; 10: 766.
Zengler K, Zaramela LS. The social network of microorganisms — how auxotrophies shape complex
communities. Nature Reviews Microbiology 2018; 16: 383-390.
Zhang J, Zou S, Fang L. Metabolic reprogramming in colorectal cancer: regulatory networks and therapy.
Cell & Bioscience 2023; 13: 25.
Zheng Y, Hu FB, Ruiz-Canela M, Clish CB, Dennis C, Salas-Salvado J, et al. Metabolites of Glutamate
Metabolism Are Associated With Incident Cardiovascular Events in the PREDIMED
PREvención con DIeta MEDiterránea (PREDIMED) Trial. Journal of the American
Heart Association 2016; 5: e003755.
Zhou Y, Danbolt NC. Glutamate as a neurotransmitter in the healthy brain. Journal of Neural Transmission
2014; 121: 799-817.
906