Spatial conservation planning of forest genetic resources in a Mediterranean multi-refugial area

Elia Vajana[a,*], Marco Andrello[b], Camilla Avanzi[a], Francesca Bagnoli[a], Giovanni G. Vendramin[a], Andrea Piotti[a]

[a] Institute of Biosciences and BioResources (IBBR), National Research Council (CNR), 50019 Sesto Fiorentino (Firenze), Italy

[b] Institute for the study of Anthropic impacts and Sustainability in the marine environment, National Research Council, CNR-IAS, Rome, Italy

* Corresponding author. *E-mail address*: elia.vajana@ibbr.cnr.it. *Full postal address*: Institute of Biosciences and Bioresources, National Research Council of Italy, Via Madonna del Piano 10, 50019 Sesto Fiorentino (Firenze), Italy.

Abstract

The Kunming-Montreal Global Biodiversity Framework recognised the urgency of taking action to conserve intraspecific genetic diversity (IGD) as an insurance against habitat degradation and environmental change. Recent work suggests that 90-99% of IGD should be conserved to safeguard viability of future generations.

Here, we addressed such a conservation issue in three forest tree species in Italy: silver fir (*Abies alba* Mill.), Heldreich's pine (*Pinus heldreichii* H. Christ), and pedunculate oak (*Quercus robur* L.). We used microsatellite markers to measure IGD of 36 (*A. alba*), 15 (*P. heldreichii*) and 25 (*Q. robur*) natural sites, including several putative glacial refugia. We developed a spatial conservation planning (SCP) analysis to quantify the genetic irreplaceability of each site and identify the minimum set coverage ensuring IGD protection. Finally, we compared SCP results with the contributions to allelic diversity within and between sites, total allelic diversity and private allelic richness.

We found that between 44% and 73% of sites were required to conserve 90-99% of the alleles, and that this conservation effort held even when targeting lower percentages of alleles to protect (50-75%). Glacial refugia were often included in the minimum set coverage, confirming theoretical and biogeographical expectations. Finally, sites with high genetic irreplaceability were found to have higher private allelic richness on average. These results are discussed in the light of the biogeographic history of the species studied and the current policies for the conservation of forest genetic resources.

Keywords: Spatial conservation planning; forest genetic resources; intraspecific genetic diversity; *Abies alba*; *Pinus heldreichii*; *Quercus robur*.

1. Introduction

Intraspecific genetic diversity (IGD), encompassing genetic variation within and between populations, is a key component of ecosystem functioning as it underlies the species' capacity to adapt to the environment, survive environmental change and provide human society with ecosystem services (Robuchon et al., 2022). Particularly, IGD of forest trees (*aka* forest genetic resources) contributes to climate regulation, water filtration, nutrient cycling, wood production, and coastal and soil protection against erosion (Alberto et al., 2013; Gougherty et al., 2021; Panagos et al., 2015; Schweitzer et al., 2011). The preservation of forest genetic resources lies at the core of sustainable forest management and conservation (Fady et al., 2016). Yet, human activities inducing global warming, habitat fragmentation and overharvesting are accelerating biodiversity decline worldwide, with massive losses in genetic diversity and unpredictable impacts on the biosphere (Exposito-Alonso et al., 2022; Hoban et al., 2023). Consequently, between 19 and 66% of the whole amount of IGD on Earth is estimated to go lost in the next decades unless specific actions are undertaken to halt genetic erosion and sustain evolutionary rescue (Hoban et al., 2023, 2021).

Among the long-term goals for 2050, the Kunming-Montreal global biodiversity framework formally recognized the need of conserving IGD and the urgency of putting in place actions to protect 'the genetic diversity within and between populations of native, wild, and domesticated species to maintain their adaptive potential' (Convention on Biological Diversity, 2022; Hoban et al., 2023). To this aim, conservation of 90-99% of extant IGD is usually advocated together with guaranteeing an effective population size >500 so as to reduce genetic erosion and prevent 'future losses for all populations [...] regardless of past losses' (Frankham, 2022; Hoban et al., 2023).

Area-based conservation is among the approaches proposed to contrast genetic erosion (Hoban et al., 2023; Robuchon et al., 2021). However, budget constraints usually require choices to be made when attempting to preserve IGD with such a strategy (Marris, 2007; Vane-Wright et al., 1991; Weitzman, 1998). Then, scientifically based criteria were developed for setting spatial priorities for conservation, e.g., whether to prioritise genetically distinct vs. highly diverse populations (Fernandez-Fournier et al., 2021; Petit et al., 1998; Weitzman, 1993) or sites with endemic lineages rather than centres of radiation (Erwin, 1991; Robuchon et al., 2021). More recently, spatial conservation planning (SCP) was applied to optimise the selection of new conservation areas based on genetic information (Andrello et al., 2022; Nielsen et al., 2023a). In this case, spatially optimised solutions are found across the study area to minimise the costs required to achieve specific conservation objectives for IGD, a formulation of the SCP problem known as 'minimum set coverage' (Moilanen et al., 2009).

Here, we used SCP to identify areas that could maximise the protection of IGD in three keystone forest tree species characterised by a highly fragmented distribution in Italy – two mountainous conifers, i.e., silver fir (*Abies alba* Mil.) and Heldreich's pine (*Pinus heldreichii* H. Christ), and a broadleaved lowland tree, the pedunculate oak (*Quercus robur* L.). Palaeobotanical and genetic evidence proved that the Mediterranean region provided temperate trees with climatically favourable refugial areas during the glacial periods characterising the Quaternary (Petit et al., 2003). Most glacial refugia were located in the Iberian, Italian and Balkan peninsulas, with populations inhabiting these areas generally

showing high levels of genetic distinctiveness (Petit et al., 2003). This genetic feature makes
forest tree populations from former glacial refugia candidate sites for conserving forest
genetic resources, as they are likely to contain unique variation that may prove necessary to
foster adaptive potential in the face of climate change (Petit et al., 2005). Here, we focused
our analysis on a set of sites within a complex of refugial areas (Brewer et al., 2002; Petit et
al., 2002; Piotti et al., 2017), where fragmentation is likely pushing peripheral populations
towards an extinction vortex in concert with climate change (Bucci et al., 1997; Gibbs, 1997;
Maiorano et al., 2013; Ruosch et al., 2016; Thomas et al., 2002).

The goals of the present work were (i) to identify priority sites for conserving IGD in silver fir,
Heldreich's pine and pedunculate oak in a multi-refugial area, and (ii) to investigate the
consensus between priority sites identified through SCP and those found through other
genetic parameters, namely allelic diversity and its decomposition into within-site and
between-site contributions (Petit e al., 1998), and effective population size, which is related
to genetic load and adaptive potential and has been formally adopted by the Convention on
Biological Diversity (2022) to monitor intraspecific genetic diversity.

2. Materials and methods

2.1. Sampling and genotyping

We relied on genetic data from 36 natural populations of *A. alba* (seven of which were
specifically characterised for this study, resulting in 1769 trees in total, 49 on average per
site; **Supplementary Table 1**), 15 natural populations of *P. heldreichii* (515 trees, 34 on
average), and 25 natural populations of *Q. robur* (745 trees, 30 on average) to feed SCP
analyses (**Figure 1**). The large representativeness of IGD was guaranteed by densely
covering the local distribution of the species, and by including sampling sites from all known
putative glacial refugial areas.

Abies alba was genotyped at 16 nuclear short sequence repeats (nSSRs), resulting in 145
alleles (Piotti et al., 2017; Santini et al., 2018), *P. heldreichii* at 12 nSSRs (89 alleles;
Piovesan et al., 2023), and *Q. robur* at 16 nSSRs (245 alleles; Avanzi et al., 2023). The
details about genotyping and standard population genetic analyses can be found in the
above-mentioned references.

Figure 1. Sampling sites: *Abies alba* (green points), *Pinus heldreichii* (blue points), and *Quercus robur* (orange points). Approximate species distributions (Caudullo et al., 2017) are also shown in the background with the same colour scheme used to indicate sampling sites (note: *A. alba* and *Q. robur* distributions are rather imprecise representations of the actual distribution of these species in the study area). The inset map zooms over the Pollino National Park, which encircles the distribution of Heldreich's pine in Italy. Shades of grey indicate elevation.

2.2. Spatial conservation planning

The SCP analysis was run separately for each species, as there were no common sampling sites to help identify priority sites shared by more than one species. In each species, each allele scored was considered as a distinct conservation feature (see the 'AL' method in Andrello et al., 2022). The spatial distribution of each allele was coded as presence/absence in each site. The number of selected sites was considered as a simple proxy for total conservation cost (Paz-Vinas et al., 2018; von Takach et al., 2021). The objective was to find solutions for conserving low (50%), moderate (75%) and high (90%, 99% and 100%) proportions of protected alleles. These proportions of protected alleles were interpreted as levels of protected IGD (Paz-Vinas et al., 2018), and included the 90% objective proposed in the first draft of the post-2020 GBF (CBD, 2021), as well as the 99% objective proposed more recently (Frankham, 2022; Hoban et al., 2023).

The SCP was run under the 'minimum set coverage' formulation to minimise the total
conservation cost needed to achieve a given level of IGD protection (Andrello et al., 2022).
This was operationalized as finding the minimum set of sites ensuring that the given
proportion of alleles (i.e., 50%, 75%, 90%, 99% or 100%) was adequately protected. An
allele was considered as adequately protected if at least 30% of its occurrence sites were
included in the solution as an insurance against catastrophic events (Lefèvre et al., 2020;
Wilson et al., 2009). For the 50%, 75%, 90% and 99% IGD levels, we performed 100
independent replicates, each based on a random selection of alleles to be protected. We
calculated the genetic irreplaceability (GI) of each site as the number of times each site was
selected across the 100 replicates (Pressey et al., 1994). For the 100% IGD level, a single
analysis was performed as all the alleles were included by definition, which allowed the
prioritisation problem to find a unique, optimal set of sites needed to conserve all the alleles.
All SCP analyses were performed using the R package 'prioritizer' v. 7.2.2 (Hanson et al.,
2022) and solved within 10% optimality using the 'Gurobi' solver v. 9.5.2 (Gurobi
Optimization, LLC, 2023).

2.3. Relationship between genetic irreplaceability and genetic diversity parameters

We studied the relationship between GI and five genetic parameters commonly used to
prioritise populations for conservation: the relative site-specific contributions to (1) within-site
(A_S) and (2) between-site (D_A) allelic diversity (the mean number of alleles occurring in the
sites under study, and the mean number of alleles occurring in a site and absent in another,
respectively), and (3) total allelic diversity (A_T; the sum of A_S and D_A), as well as (4) private
allelic richness (*PAr*; the number of unique alleles in a given site) and (5) effective population
size (*Ne*; the size of a Wright-Fisher's idealised population displaying the same rate of
inbreeding or drift observed in the population under study, roughly corresponding to the
number of reproductive individuals contributing to next generations; Jamieson and Allendorf,
2012). Sites mostly contributing to A_T can be identified by their relative contribution to A_S and
D_A (Petit et al., 1998). A_T, A_S and D_A were calculated for all sites in each species with the
'metapop2' software (López-Cortegano et al., 2019). *PAr* was estimated with 'HP-RARE'
(Kalinowski, 2005). Rarefaction was used to account for uneven sample sizes in the
calculation of A_T, A_S, D_A, and *PAr* (El Mousadik and Petit, 1996). *Ne* was estimated using the
linkage disequilibrium method (Waples and Do, 2010) implemented in 'NeEstimator' v. 2.1
(Do et al., 2014) and screening out alleles with a frequency <2%.

The relationship between GI and genetic parameters was assessed based on the GI values
obtained when protecting 99% of alleles. Considering the distribution of GI values, sites were
classified to be at low/high GI, using a threshold of ≤20% (low GI) and ≥80% (high GI),
respectively. Associations between levels of GI and genetic parameters were tested through
either the Welch or Student's two sample *t*-test, depending on whether the variances of the
groups resulted to be unequal or equal based on the Levene's test. To reduce inflation in the
familywise type I error due to multiple testing, the Bonferroni correction was applied to the *p*-
values of the tests involving the same species. A test was deemed to be statistically
significant if its adjusted *p*-value was equal to or lower than the nominal significance
threshold (α) of 0.05. The lower bound of the confidence intervals around the *Ne* estimates
was used in the tests to avoid dealing with infinite *Ne* values. All operations were performed
in the R programming environment v. 4.3.2 (R Core Team, 2023).

3. Results

3.1. Spatial conservation planning

In all species, the number of sites needed for conserving 100% IGD was less than the total number of sampled sites, and was equal to 16 (44% of sampled sites) in *A. alba*, 11 (73%) in *P. heldreichii*, and 18 (72%) in *Q. robur*. For conserving lower levels of IGD, the median number of sites needed was equal or slightly lower with almost no variation among the replicates (**Table 1**).

Table 1. Median number of sites needed to conserve 50%, 75%, 90%, 99%, and 100% of IGD across 100 replicates in *Abies alba*, *Pinus heldreichii* and *Quercus robur*.

	Protection levels				
	50%	75%	90%	99%	100%
A. alba (36)	14 (±1)[a]	15 (ND)[b]	16 (ND)	16 (ND)	16[c]
P. heldreichii (15)	8 (±1)	10 (±1)	11 (ND)	11 (ND)	11
Q. robur (25)	15 (±1)	17 (±1)	18 (ND)	18 (ND)	18

[a] Median absolute deviation (MAD), i.e., an estimate of the variability around the number of sites in the replicates. Each MAD was calculated as the median of the absolute deviations from the number of sites. [b] No deviation from the number of sites observed. [c] No variation was associated with the 100% level as a unique analysis was performed in this scenario. The total number of sampled sites is given in brackets next to each species name (first column).

Regardless of the number of alleles to be protected in the SCP analysis, we found that ~98% of the replicates conserved at least 90% of all alleles present in the dataset (**Figure 2**). All replicates protected >90% of alleles present in the dataset when including 75% of alleles in the SCP analysis. Overall, the SCP analyses systematically ensured protection for higher numbers of alleles than those included as conservation features.

Figure 2. Variation in the percentage of alleles protected in at least 30% of their sites of occurrence as a function of levels of IGD secured (50%, 75%, 90%,99% and 100%). The proportion of alleles adequately protected is represented separately for each species.

For *A. alba*, a bimodal spatial pattern of GI appeared and four sites, two in the northern Apennines and two in the southern Apennines, were irreplaceable (i.e., GI=100%) when 90% of the alleles were conserved (**Figure 3a**). When targeting the 99% level, 11 sites became irreplaceable and were more evenly distributed across the Italian peninsula (**Figure 3b**). For *P. heldreichii*, 11 sites showed GI≥88% when conserving ≥90% of alleles, including one irreplaceable site on the southern margins (**Figure 3c**). When 99% of the alleles were conserved, all such 11 sites became irreplaceable (**Figure 3d**). Irreplaceable sites were located at short distances (<10 Km apart), and included a group of five close, geographically central sites, and three sites on the southern margins. For *Q. robur*, we found three irreplaceable sites even when conserving 75% of alleles (result not shown), including the southernmost site in Italy ('Bosco Pantano'). When protecting 90% of alleles (**Figure 3e**), four sites were found to be irreplaceable, including a new site in the Po valley (northern Italy). In this case, we found 18 sites with GI≥68%, and seven sites with GI≤27%. The same sites at higher GI were confirmed when conserving 99% of the alleles, when we observed a GI higher than 96% and lower than 5% in sites at high vs. low GI, respectively (**Figure 3f**).

210
211 **Figure 3.** Genetic irreplaceability (GI) of sampled sites in *A. alba* (**a,b**), *P. heldreichii* (**c,d**), and *Q. robur* (**e,f**), as
212 calculated over 100 replicates for different levels of IGD secured (i.e., 90% and 99% on the left and right
213 columns, respectively). The GI of each site is indicated by both the size of the dots and their colour (a smaller
214 size and blue tones indicate low GI, while a larger size and red tones indicate high GI).

3.2. Relationship between genetic irreplaceability and genetic diversity parameters

After multiple testing corrections, four out of the 15 tests used to assess the relationship between GI and genetic diversity parameters were statistically significant at a nominal α=0.05 (**Figure 4**). Particularly, *PAr* resulted to be associated with GI in all species, with the set of sites at higher GI showing a higher mean *PAr*. *PAr* displayed high variability in high GI sites (**Figure 4** and **Supplementary Table 2**). A_T, A_S and D_A were not associated with GI. On the contrary, *Ne* turned out to be significantly associated with GI in *Q. robur*, where high GI sites had lower *Ne* on average compared to low GI sites.

Figure 4. Distributions of site-specific contributions to within-site allelic diversity (A_S), between-site allelic diversity (D_A) and total allelic diversity (A_T), private allelic richness (*PAr*) and effective population size (*Ne*) in sites with low and high genetic irreplaceability (GI<20% and GI>80%, respectively). Darker boxes indicate statistically

227 significant differences between the means of the low and high GI sites after the Bonferroni correction for multiple
228 testing. *Ne* values are presented on a logarithmic scale for graphical purposes.

4. Discussion

We applied SCP to intraspecific genetic data covering the Italian distribution of three forest
tree species that are key elements of Mediterranean forest ecosystems (*Abies alba, Pinus*
heldreichii and *Quercus robur*). Overall, we found that between 44% and 73% of the
sampled sites, depending on the species, are needed to achieve adequate protection for all
the alleles present in the sites studied. We showed that even if the objective was protecting
fewer alleles (50% of alleles), the required number of sites did not change much. Moreover,
even when the objective was to protect fewer alleles, the selected sites could ensure
protection for a higher proportion of alleles. Finally, we found that private allelic richness (i.e.,
the number of alleles uniquely present in a given site) was the genetic parameter mostly
related to genetic irreplaceability, a measure of the importance of conserving a given site to
protect IGD. In the following, we discuss these results in the light of the conservation status
of silver fir, Heldreich's pine and pedunculate oak, as well as their life history traits and
biogeographic history.

4.1. Full protection can be achieved at 'little' extra cost

Our SCP analysis resulted in a rather costly solution to protect IGD to the suggested level of
99% (Frankham, 2022). In fact, the percentage of protected sites was never less than 44%
and included 16, 11 and 18 sites for silver fir, Heldreich's pine and pedunculate oak,
respectively. Nevertheless, a difference of only two to three sites separated the achievement
of half and full protection of alleles, suggesting that low levels of protection (50% of alleles)
are neither a parsimonious strategy for IGD conservation in these species, nor do they imply
substantial cost reduction. Yet, any additional economic effort to achieve the 99% protection
level (e.g., land acquisition and management) will depend critically on the available budget
and socio-political will (Margules and Pressey, 2000; Naidoo et al., 2006).

4.2. Private allelic richness is not a perfect proxy for genetic irreplaceability

To explore the relationship between different approaches to priority setting, we related the
genetic irreplaceability of a site to its relative contribution to within-site, between-site, and
total allelic diversity, as well as to its effective population size and private allelic richness.
These genetic parameters find wide application in conservation thanks to their sensitivity to
demographic fluctuations (López-Cortegano et al., 2019; Petit et al., 1998; Tapio et al.,
2006), and their capacity to comply with the principle of complementarity (Bonin et al., 2007)
and to inform allocation of economic resources (Von Takach et al., 2023). The significant,
positive correlation found between GI and *PAr* points to the spatial distribution of private
alleles as one of the main genetic features driving site selection in our SCP analysis
(**Supplementary Figure 1**). By relying on an SCP approach like the one used here,
selection for sites enriched in private alleles has also been observed in other taxa, including
freshwater fish species (Paz-Vinas et al., 2018) and marsupials (Von Takach et al., 2023).
However, such a correspondence between GI and *PAr* is not straightforward, as we found
that some sites with low *PAr* can show a high GI. Thus, we recommend caution when using
PAr as a perfect proxy for GI, as some highly irreplaceable sites could be ignored. We also

note that selecting sites with private alleles implies including all non-private alleles contained
in these sites in the final selection, which, depending on the spatial distribution of alleles and
their frequencies, may increase the proportion of alleles that are adequately protected
overall. Here, we never ended up in protecting less than 80% of the alleles, regardless of the
level of protection set prior to the analysis.

4.3. Towards improving the interpretation and calculation of genetic irreplaceability

The other genetic parameters tested were not related to genetic irreplaceability, except for
Ne in pedunculate oak, where high GI sites display a significantly lower Ne than low GI sites.
This finding may suggest the use of genetic irreplaceability in combination with Ne to
simultaneously locate priority sites for IGD conservation and evaluate their genetic health
status, especially when dealing with highly fragmented populations as in the case of
pedunculate oak in Italy (Avanzi et al., 2023). Rapidly shrinking populations in an area
characterised by a strong genetic structure may explain the occurrence of high private allelic
richness in sites with low Ne (see, for example, the case of the 'Foglino' and 'Bosco
Pantano' sites in central and southern Italy, where PAr is high and Ne ranges from 27 to 81;
Supplementary Table 2). On the other hand, new analyses with a broader species panel
would be needed to clarify the relationship between Ne and GI and to provide guidance on
how the genetic indicators adopted by the Convention on Biological Diversity (2022) can be
harmonised with the SCP approach (in our case, for example, less than half of the high GI
sites of each species had $Ne>500$, and our solutions often discarded sites with high Ne).

Our approach optimises the selection of sites so as to include at least 30% of the sites
where an allele is present as an insurance against random allele loss. While valid, this
approach ignores the potential relevance of within-site allele frequencies in calculating GI.
Taking allele frequencies into account would be important from a conservation point of view
because low frequency alleles are more prone to disappear due to the effects of genetic drift.
For example, alleles with within-site frequency ≤5% were evenly distributed between low and
high priority sites in the species under study (**Supplementary Figure 2**). The possible loss
of rare alleles due to lack of protection could be overcome by setting specific targets for
them, analogously to what is done when setting targets for species representation in
conservation prioritisation (e.g., Rodrigues et al., 2004), or by 'locking-in' sites with higher
proportions of rare alleles, so as to force their inclusion in the final selection of sites.

4.4. Glacial refugia are priority areas for conservation

The geographical range of the large majority of irreplaceable sites for *A. alba* corresponds to
the Apennine mountain range, and includes all refugia areas that were proposed for this
species based on both genetic and paleobotanical data (Magri et al., 2015; Piotti et al.,
2017): the Calabrian arc (southern Apennines) including four sites; the central Apennines,
including three sites located across two highly differentiated genetic lineages; and the
northern Apennines, including four sites. Despite fragmentation and documented declines in
abundance during the Holocene (Magri et al., 2015), such areas still maintain large
population sizes, which will likely protect them from future genetic erosion (Forester et al.,
2022; Kardos et al., 2021), and show high allelic diversity, local adaptation and growth vigour
in provenance studies (Kerr et al., 2015; Piotti et al., 2017). The higher occurrence of high GI
sites in the southern refugial area compared to the northern area may reflect high private

allelic richness and a long history of genetic isolation from the northernmost populations
(Piotti et al., 2017). Physical barriers (i.e., the Gran Sasso and Majella massifs) and possible
constraints to long-distance dispersal due to heavy seeds and pollen (Leonarduzzi et al.,
2016) may have acted in concert to prevent gene flow from the southern refugial area,
facilitating the development of spatial genetic structuring (Piotti et al., 2017).

High GI sites for *Q. robur* also generally corresponded to refugial areas, which include the
Pre-Alps, the southern slope of the Tuscan-Emilian Apennines, the Tyrrhenian coast and
some areas in southern Italy, where the species was likely abundant at the onset of the
Holocene (Magri et al., 2015). In particular, irreplaceable sites include six sites along the Po
River (northern Italy), five sites in central Italy (two of them in Tuscany) and the putative
refugium of 'Bosco Pantano' in the South. Contrary to what is observed in other refugial
areas, 'Bosco Pantano' is genetically depauperate, with extremely low allelic diversity and
effective population size (*Ne*<100) due to recent intensive human impact. The high
irreplaceability of this site may rather reflect its extremely large genetic distinctiveness and
the presence of private alleles (Avanzi et al., 2023). Similar to our findings, Beridze et al.
(2023) have also recently shown that sites within refugial areas of *Castanea sativa* deserve
high conservation priority.

4.5. Practical implications of SCP results

Based on comprehensive genetic datasets characterising neutral genetic diversity in a multi-
refugial area, we were able to identify priority sites for conserving IGD in *A. alba*, *P.
heldreichii* and *Q. robur*. The implications are manifold.

First, the genetic irreplaceability map (**Figure 3**) can make it possible to assess the current
level of legal protection of sites whose loss would reduce the IGD of the species. All *P.
heldreichii* sampled populations are within the Pollino National Park, whose perimeter was
originally designed to include and protect all Italian populations of this species. The majority
of *A. alba* and *Q. robur* populations are within National and Regional Parks, or public
Reserves. Nonetheless, some irreplaceable sites for *Q. robur* are located on private property
without any protection regime (e.g., the site with the highest genetic diversity in Italy; see
Avanzi et al., 2023). Our results can contribute to change the protection status of such
stands or to raise awareness among private owners toward a correct management of forest
genetic resources. Finally, if some of these priority sites were ultimately unprotectable,
replacements could be identified on the basis of their genetic irreplaceability, considering
complementary genetic indicators (e.g., *Ne*), or through additional genetic sampling
(Margules and Pressey, 2000).

Seed collection for *in situ* and *ex situ* conservation strategies as well as for reforestation
programs, the latter having been massively funded in Italy in the last years (Mariotti et al.,
2022), is usually carried out by the public sector in designated seed stands. Seed stands
were not identified based on their genetic characteristics but mostly relying on the
knowledge of local experts. Our results can help refine the Italian network of seed stands in
terms of optimising the number and geographical position of selected stands (Wei and Jiang,
2020). For *A. alba*, for instance, >50 seed stands are registered at the national level
(MASAF, 2023) but none is present in a relatively small genetic cluster found in central Italy
(where two irreplaceable sites were identified) or in the site with the highest genetic diversity

in Italy ('Terranova di Pollino' in Piotti et al., 2017), which we found to be irreplaceable
starting from the lowest protection level. For *Q. robur,* there are about 100 seed stands
registered, but none of them includes sites in central or southern Italy (the southernmost
seed stand is at 43° 21' latitude), where our SCP analysis identified six sites with high GI.
This means that the genetic diversity of populations at the rear edge of the species'
distribution, which might have a disproportionate value in boosting the adaptive potential of
the northernmost populations through gene flow or assisted migration (EUFORGEN, 2021),
is currently not represented in seed collection initiatives in Italy. Finally, *P. heldreichii* has no
registered seed stands so far, and our findings may be useful in defining an initial priority list.

The Pan-European network of genetic conservation units (Lefèvre et al., 2013;
http://portal.eufgis.org/) shows a better agreement with our SCP analyses. For instance, the
southernmost *Q. robur* population, which represents a unique genetic lineage and nowadays
consists of only 64 individuals (Avanzi et al., 2023), is recognized as a genetic conservation
unit but not as a seed stand. In addition, despite the absence of seed stands, a genetic
conservation unit for *P. heldreichii* was established in a 166-ha area of the Pollino National
Park that includes five of the irreplaceable sites identified in our analysis. Nonetheless, the
selection of genetic conservation units (for which a first, partial genetic characterisation is
currently underway in the framework of the EU-funded FORGENIUS project;
www.forgenius.eu) could greatly benefit from SCP studies such as the one presented here.

4.6. Conclusions

SCP proved to be an effective method of identifying conservation areas capable of
preserving the intraspecific allelic diversity of silver fir, Heldreich's pine and pedunculate oak.
In particular, we found that between 50% and 75% of sampled sites are required to protect
IGD in these species, which is a higher percentage than the one usually recommended to
protect habitats (30%; Convention on Biological Diversity, 2022). Importantly, we found that
these percentages hold for both low (50%) and more ambitious percentages of alleles to be
protected (>90%). Confirming theoretical expectations (Petit et al., 2003), glacial refugial
areas were frequently included in the networks of proposed protected areas. This result may
also apply to forest species for which SCP has not yet been undertaken.

Multifaceted applications of data and approaches presented here have been highlighted,
with particular reference to optimising the seed collection necessary to fuel ongoing
afforestation initiatives such as the commitment for planting at least 3 billion trees in the
European Union by 2030 (EC, 2020). SCP could provide forest conservationists with clear
guidance on which sites should be prioritised or added to the existing genetic conservation
unit network. In this regard, informed decision-making is key to implementing any action
aimed at achieving specific conservation targets (e.g., achieving 90-99% protection of
existing forest genetic resources; Frankham, 2022; Hoban et al., 2023), monitoring IGD over
time and flagging any need for active intervention (EUFORGEN, 2021).

Finally, we advocate that future research should focus on (i) describing the relative
performance of SCP compared to more traditional methods based on allelic diversity and
effective population size in different conservation scenarios, (ii) harmonising SCP objectives
with the principles for the conservation of intraspecific genetic diversity proposed by the
Convention on Biological Diversity (2022), and (iii) exploring the potential impact of sample

size and the set of genetic markers used on SCP performance. In particular, the extension of
SCP to genomic data should merit special attention to include both neutral and functional
variants in the analysis (Hanson et al., 2020; Nielsen et al., 2023b, 2020; Xuereb et al.,
2021), which promises to increase the effectiveness of genetic-informed conservation
measures.

Abbreviations

404 A_S: the site-specific contribution to within-site allelic diversity; D_A: the site-specific
405 contribution to between-site allelic diversity; A_T: the site-specific contribution to total allelic
406 diversity; GI: genetic irreplaceability; IGD: intraspecific genetic diversity; *Ne*: effective
407 population size; *PAr*: private allelic richness; SCP: spatial conservation planning.

Acknowledgments

409 E. Vajana is supported by the EU-funded project MedForAct (grant agreement ID
410 101107604). We thank all Parks and Reserves where sampling sites are located for
411 permissions and logistic support. The acknowledgements to specific persons and institutions
412 are reported in the articles where original data were published, indicated in the Materials &
413 Methods section.

Author contributions

Elia Vajana: Conceptualization; Data curation; Formal analysis; Visualization; Writing -
original draft. **Marco Andrello**: Conceptualization; Formal analysis; Methodology;
Supervision; Writing - review & editing. **Camilla Avanzi**: Conceptualization; Data curation;
Formal analysis; Visualization; Writing - review & editing. **Francesca Bagnoli**: Funding
acquisition; Writing - review & editing. **Giovanni G. Vendramin**: Funding acquisition;
Supervision; Writing - review & editing. **Andrea Piotti**: Conceptualization; Data curation;
Formal analysis; Funding acquisition; Supervision; Visualization; Writing - review & editing.

References

Alberto, F.J., Aitken, S.N., Alía, R., González-Martínez, S.C., Hänninen, H., Kremer, A.,
Lefèvre, F., Lenormand, T., Yeaman, S., Whetten, R., Savolainen, O., 2013.
Potential for evolutionary responses to climate change – evidence from tree
populations. Glob. Change Biol. 19, 1645–1661. https://doi.org/10.1111/gcb.12181
Andrello, M., D'Aloia, C., Dalongeville, A., Escalante, M.A., Guerrero, J., Perrier, C., Torres-
Florez, J.P., Xuereb, A., Manel, S., 2022. Evolving spatial conservation prioritization
with intraspecific genetic data. Trends Ecol. Evol. 37, 553–564.
https://doi.org/10.1016/j.tree.2022.03.003
Avanzi, C., Bagnoli, F., Romiti, E., Spanu, I., Tsuda, Y., Vajana, E., Vendramin, G.G., Piotti,
A., 2023. The latitudinal trend in genetic diversity and distinctiveness of *Quercus*
robur rear edge forest remnants calls for new conservation priorities. bioRχiv.
https://doi.org/10.1101/2023.10.30.564819
Beridze, B., Sękiewicz, K., Walas, Ł., Thomas, P.A., Danelia, I., Fazaliyev, V, Kvartskhava,
G., Sós, J., Dering, M., 2023. Biodiversity protection against anthropogenic climate
change: Conservation prioritization of *Castanea sativa* in the South Caucasus based
on genetic and ecological metrics. Ecol. Evol. 13(5), e10068.
Bonin, A., Nicole, F., Pompanon, F., Miaud, C., Taberlet, P., 2007. Population Adaptive

Index: a new method to help measure intraspecific genetic diversity and prioritize populations for conservation. Conserv. Biol. 21, 697–708. https://doi.org/10.1111/j.1523-1739.2007.00685.x

Brewer, S., Cheddadi, R., de Beaulieu, J.L., Reille, M., 2002. The spread of deciduous *Quercus* throughout Europe since the last glacial period. For. Ecol. Manag. 156, 27–48. https://doi.org/10.1016/S0378-1127(01)00646-6

Bucci, G., Vendramin, G.G., Lelli, L., Vicario, F., 1997. Assessing the genetic divergence of *Pinus leucodermis* Ant. endangered populations: use of molecular markers for conservation purposes. Theor. Appl. Genet. 95, 1138–1146. https://doi.org/10.1007/s001220050674

Caudullo, G., Welk, E., San-Miguel-Ayanz, J., 2017. Chorological maps for the main European woody species. Data Brief 12, 662–666. https://doi.org/10.1016/j.dib.2017.05.007

CBD, 2021. First draft of the post-2020 global biodiversity framework.

Convention on Biological Diversity (CBD/COP/DEC/15/4), 2022. Conference of the parties to the Convention of Biological Diversity (Fifteenth meeting - Part II) (No. Agenda item 9A). Montreal, Canada.

Do, C., Waples, R.S., D. Peel, D., Macbeth, G.M., Tillett, B.J., Ovenden, J.R., 2014. NEESTIMATOR v2: re-implementation of software for the estimation of contemporary effective population size (*Ne*) from genetic data. Mol. Ecol. Resour. 14, 209–214. https://doi.org/10.1111/1755-0998.12157

EC, 2020. EU Biodiversity Strategy for 2030. Bringing nature back into our lives, COM(2020) 380 final. Brussels.

El Mousadik, A., Petit, R.J., 1996. High level of genetic differentiation for allelic richness among populations of the argan tree [*Argania spinosa* (L.) Skeels] endemic to Morocco. Theor. Appl. Genet. 92, 832–839.

Erwin, T.L., 1991. An evolutionary basis for conservation strategies. Sci. New Ser. 253, 750–752.

EUFORGEN, 2021. Forest genetic resources strategy for Europe. European Forest Institute.

Exposito-Alonso, M., Booker, T.R., Czech, L., Gillespie, L., Hateley, S., Kyriazis, C.C., Lang, P.L.M., Leventhal, L., Nogues-Bravo, D., Pagowski, V., Ruffley, M., Spence, J.P., Arana, S.E.T., Weiß, C.L., Zess, E., 2022. Genetic diversity loss in the Anthropocene. Science 377, 1431–1435. https://doi.org/10.1126/science.1083264

Fady, B., Cottrell, J., Ackzell, L., Alía, R., Muys, B., Prada, A., González-Martínez, S.C., 2016. Forests and global change: what can genetics contribute to the major forest management and policy challenges of the twenty-first century? Reg. Environ. Change 16, 927–939. https://doi.org/10.1007/s10113-015-0843-9

Fernandez-Fournier, P., Lewthwaite, J.M.M., Mooers, A.Ø., 2021. Do We Need to Identify Adaptive genetic variation when prioritizing populations for conservation? Conserv. Genet. 22, 205–216. https://doi.org/10.1007/s10592-020-01327-w

Forester, B.R., Beever, E.A., Darst, C., Szymanski, J., Funk, W.C., 2022. Linking evolutionary potential to extinction risk: applications and future directions. Front. Ecol. Environ. fee.2552. https://doi.org/10.1002/fee.2552

Frankham, R., 2022. Evaluation of proposed genetic goals and targets for the Convention on Biological Diversity. Conserv. Genet. 23, 865–870. https://doi.org/10.1007/s10592-022-01459-1

Gibbs, J., 1997. Biotic and abiotic factors affecting the dying back of pedunculate oak *Quercus robur* L. Forestry 70, 399–406. https://doi.org/10.1093/forestry/70.4.399

Gougherty, A.V., Keller, S.R., Fitzpatrick, M.C., 2021. Maladaptation, migration and extirpation fuel climate change risk in a forest tree species. Nat. Clim. Change 11, 166–171. https://doi.org/10.1038/s41558-020-00968-6

Gurobi Optimization, LLC, 2023. Gurobi Optimizer Reference Manual.

Hanson, J., Schuster, R., Morrell, N., Strimas-Mackey, M., Edwards, B., Watts, M., Arcese, P., Bennett, J., Possingham, H., 2022. prioritizr: Systematic Conservation Prioritization in R.

Hanson, J.O., Marques, A., Veríssimo, A., Camacho-Sanchez, M., Velo-Antón, G., Martínez-Solano, Í., Carvalho, S.B., 2020. Conservation planning for adaptive and neutral evolutionary processes. J. Appl. Ecol. 57, 2159–2169. https://doi.org/10.1111/1365-2664.13718

Hoban, S., Bruford, M.W., da Silva, J.M., Funk, W.C., Frankham, R., Gill, M.J., Grueber, C.E., Heuertz, M., Hunter, M.E., Kershaw, F., Lacy, R.C., Lees, C., Lopes-Fernandes, M., MacDonald, A.J., Mastretta-Yanes, A., McGowan, P.J.K., Meek, M.H., Mergeay, J., Millette, K.L., Mittan-Moreau, C.S., Navarro, L.M., O'Brien, D., Ogden, R., Segelbacher, G., Paz-Vinas, I., Vernesi, C., Laikre, L., 2023. Genetic diversity goals and targets have improved, but remain insufficient for clear implementation of the post-2020 global biodiversity framework. Conserv. Genet. https://doi.org/10.1007/s10592-022-01492-0

Hoban, S., Campbell, C.D., da Silva, J.M., Ekblom, R., Funk, W.C., Garner, B.A., Godoy, J.A., Kershaw, F., MacDonald, A.J., Mergeay, J., Minter, M., O'Brien, D., Vinas, I.P., Pearson, S.K., Pérez-Espona, S., Potter, K.M., Russo, I.-R.M., Segelbacher, G., Vernesi, C., Hunter, M.E., 2021. Genetic diversity is considered important but interpreted narrowly in country reports to the Convention on Biological Diversity: Current actions and indicators are insufficient. Biol. Conserv. 261, 109233. https://doi.org/10.1016/j.biocon.2021.109233

Jamieson, I.G., Allendorf, F.W., 2012. How does the 50/500 rule apply to MVPs? Trends Ecol. Evol. 27, 578–584. https://doi.org/10.1016/j.tree.2012.07.001

Kalinowski, S.T., 2005. hp-rare 1.0: a computer program for performing rarefaction on measures of allelic richness. Mol. Ecol. Notes 5, 187–189.

Kardos, M., Armstrong, E.E., Fitzpatrick, S.W., Hauser, S., Hedrick, P.W., Miller, J.M., Tallmon, D.A., Funk, W.C., 2021. The crucial role of genome-wide genetic variation in conservation. Proc. Natl. Acad. Sci. 118, e2104642118. https://doi.org/10.1073/pnas.2104642118

Kerr, G., Stokes, V., Peace, A., Jinks, R., 2015. Effects of provenance on the survival, growth and stem form of European silver fir (*Abies alba* Mill.) in Britain. Eur. J. For. Res. 134, 349–363. https://doi.org/10.1007/s10342-014-0856-9

Lefèvre, F., Alia, R., Bakkebø Fjellstad, K., Graudal, L., Oggioni, S.D., Rusanen, M., Vendramin, G.G., Bozzano, M., 2020. Dynamic conservation and utilization of forest tree genetic resources: indicators for *in situ* and *ex situ* genetic conservation and forest reproductive material. European Forest Genetic Resources Programme (EUFORGEN). European Forest Institute.

Lefèvre, F., Koskela, J., Hubert, J., Kraigher, H., Longauer, R., Olrik, D.C., Schüler, S., Bozzano, M., Alizoti, P., Bakys, R., Baldwin, C., Ballian, D., Black-Samuelsson, S., Bednarova, D., Bordács, S., Collin, E., De Cuyper, B., De Vries, S.M.G., Eysteinsson, T., Frýdl, J., Haverkamp, M., Ivankovic, M., Konrad, H., Koziol, C., Maaten, T., Notivol Paino, E., Öztürk, H., Pandeva, I.D., Parnuta, G., Pilipovič, A., Postolache, D., Ryan, C., Steffenrem, A., Varela, M.C., Vessella, F., Volosyanchuk, R.T., Westergren, M., Wolter, F., Yrjänä, L., Zariņa, I., 2013. Dynamic conservation of forest genetic resources in 33 European countries. Conserv. Biol. 27, 373–384. https://doi.org/10.1111/j.1523-1739.2012.01961.x

Leonarduzzi, C., Piotti, A., Spanu, I., Vendramin, G.G., 2016. Effective gene flow in a historically fragmented area at the southern edge of silver fir (*Abies alba* Mill.) distribution. Tree Genet. Genomes 12, 95. https://doi.org/10.1007/s11295-016-1053-4

López-Cortegano, E., Pérez-Figueroa, A., Caballero, A., 2019. metapop2: Re-implementation of software for the analysis and management of subdivided populations using gene and allelic diversity. Mol. Ecol. Resour. 19, 1095–1100. https://doi.org/10.1111/1755-0998.13015

López-Cortegano, E., Pouso, R., Labrador, A., Pérez-Figueroa, A., Fernández, J., Caballero, A., 2019. Optimal management of genetic diversity in subdivided populations. Front. Genet. 10, 843. https://doi.org/10.3389/fgene.2019.00843

Magri, D., Agrillo, E., Di Rita, F., Furlanetto, G., Pini, R., Ravazzi, C., Spada, F., 2015. Holocene dynamics of tree taxa populations in Italy. Rev. Palaeobot. Palynol. 218, 267–284. https://doi.org/10.1016/j.revpalbo.2014.08.012

Maiorano, L., Cheddadi, R., Zimmermann, N.E., Pellissier, L., Petitpierre, B., Pottier, J., Laborde, H., Hurdu, B.I., Pearman, P.B., Psomas, A., Singarayer, J.S., Broennimann, O., Vittoz, P., Dubuis, A., Edwards, M.E., Binney, H.A., Guisan, A., 2013. Building the niche through time: using 13,000 years of data to predict the effects of climate change on three tree species in Europe. Glob. Ecol. Biogeogr. 22, 302–317. https://doi.org/10.1111/j.1466-8238.2012.00767.x

Margules, C.R., Pressey, R.L., 2000. Systematic conservation planning. Nature 405.

Mariotti, B., Mezzalira, G., Allasia, E., Fazio, F., Fiorentin, R., Maltoni, A., Marchetti, M., Matteucci, G., Mori, P., Motta, R., Piotti, A., Rositi, A., Sabatti, M., Tognetti, R., Salbitano, F., 2022. The Italian forest nursery sector at the crossroads: challenges and strategies. For. - Riv. Selvic. Ed Ecol. For. 19, 85–94. https://doi.org/10.3832/efor4193-019

Marris, E., 2007. Conservation priorities: What to let go. Nature 450, 152–155.

MASAF, 2023. *Cartografia dei materiali di base per la produzione di materiali forestali di moltiplicazione*.

Moilanen, A., Possingham, H.P., Polasky, S., 2009. A mathematical classification of conservation prioritization problems, in: Moilanen, A., Wilson, K.A., Possingham, H.P. (Eds.), Spatial Conservation Prioritization: Quantitative Methods and Computational Tools. Oxford University Press, Oxford, UK.

Naidoo, R., Balmford, A., Ferraro, P., Polasky, S., Ricketts, T., Rouget, M., 2006. Integrating economic costs into conservation planning. Trends Ecol. Evol. 21, 681–687. https://doi.org/10.1016/j.tree.2006.10.003

Nielsen, E.S., Hanson, J.O., Carvalho, S.B., Beger, M., Henriques, R., Kershaw, F., Von Der Heyden, S., 2023a. Molecular ecology meets systematic conservation planning. Trends Ecol. Evol. 38, 143–155. https://doi.org/10.1016/j.tree.2022.09.006

Nielsen, E.S., Beger, M., Henriques, R., Von Der Heyden, S., 2023b. Integrating environmental, evolutionary, and socioeconomic vulnerability to future-proof coastal conservation planning. Biol. Conserv. 286, 110302. https://doi.org/10.1016/j.biocon.2023.110302

Nielsen, E.S., Beger, M., Henriques, R., Von Der Heyden, S., 2020. A comparison of genetic and genomic approaches to represent evolutionary potential in conservation planning. Biol. Conserv. 251, 108770. https://doi.org/10.1016/j.biocon.2020.108770

Panagos, P., Borrelli, P., Poesen, J., Ballabio, C., Lugato, E., Meusburger, K., Montanarella, L., Alewell, C., 2015. The new assessment of soil loss by water erosion in Europe. Environ. Sci. Policy 54, 438–447. https://doi.org/10.1016/j.envsci.2015.08.012

Paz-Vinas, I., Loot, G., Hermoso, V., Veyssière, C., Poulet, N., Grenouillet, G., Blanchet, S., 2018. Systematic conservation planning for intraspecific genetic diversity Proceedings of the Royal Society B, 20172746. http://dx.doi.org/10.1098/rspb.2017.2746

Petit, R.J., Abdelhamid, E.M., Odile, P., 1998. Identifying populations for conservation on the basis of genetic markers. Conservation biology 12, 844–855.

Petit, R.J., Aguinagalde, I., de Beaulieu, J.-L., Bittkau, C., Brewer, S., Cheddadi, R., Ennos, R., Fineschi, S., Grivet, D., Lascoux, M., Mohanty, A., Müller-Starck, G., Demesure-Musch, B., Palme, A., Martín, J.P., Rendell, S., Vendramin, G.G., 2003. Glacial refugia: hotspots but not melting pots of genetic diversity. Science 300, 1563–1565. https://doi.org/10.1126/science.1083264

Petit, R.J., Brewer, S., Bordács, S., Burg, K., Cheddadi, R., Coart, E., Cottrell, J., Csaikl, U.M., van Dam, B., Deans, J.D., Espinel, S., Fineschi, S., Finkeldey, R., Glaz, I., Goicoechea, P.G., Jensen, J.S., König, A.O., Lowe, A.J., Madsen, S.F., Mátyás, G., Munro, R.C., Popescu, F., Slade, D., Tabbener, H., de Vries, S.G.M., Ziegenhagen, B., de Beaulieu, J.-L., Kremer, A., 2002. Identification of refugia and post-glacial colonisation routes of European white oaks based on chloroplast DNA and fossil

pollen evidence. For. Ecol. Manag. 156, 49–74. https://doi.org/10.1016/S0378-
1127(01)00634-X

Petit, R.J., Hampe, A., Cheddadi, R., 2005. Climate changes and tree phylogeography in the
Mediterranean. Taxon 54, 877. https://doi.org/10.2307/25065474

Piotti, A., Leonarduzzi, C., Postolache, D., Bagnoli, F., Spanu, I., Brousseau, L., Urbinati, C.,
Leonardi, S., Vendramin, G.G., 2017. Unexpected scenarios from Mediterranean
refugial areas: disentangling complex demographic dynamics along the Apennine
distribution of silver fir. J. Biogeogr. https://doi.org/10.1111/jbi.13011

Piovesan, G., Rita, A., Biondi, F., Baliva, M., Borghetti, M., Brunetti, M., De Vivo, G., Di
Filippo, A., Dinella, A., Gentilesca, T., Maugeri, M., Palli, J., Piotti, A., Saba, E.P.,
Ripullone, F., Schettino, A., Vendramin, G.G., 2023. Bell-shaped tree-ring responses
to air temperature drive productivity trends in long-lived mountain Mediterranean
pines. Sci. Total Environ. 890, 164103.
https://doi.org/10.1016/j.scitotenv.2023.164103

Pressey, R.L., Johnson, I.R., Wilson, P.D., 1994. Shades of irreplaceability: towards a
measure of the contribution of sites to a reservation goal. Biodivers. Conserv. 3, 242–
262. https://doi.org/10.1007/BF00055941

R Core Team, 2023. R: A language and environment for statistical computing. R Found.
Stat. Comput. Vienna Austria.

Robuchon, M., Pavoine, S., Véron, S., Delli, G., Faith, D.P., Mandrici, A., Pellens, R.,
Dubois, G., Leroy, B., 2021. Revisiting species and areas of interest for conserving
global mammalian phylogenetic diversity. Nat. Commun. 12, 3694.
https://doi.org/10.1038/s41467-021-23861-y

Robuchon, M., Silva, J. da, Dubois, G., Gumbs, R., Hoban, S., Laikre, L., Owen, N.R.,
Perino, A., 2022. Conserving species' evolutionary potential and history:
opportunities under the new post-2020 global biodiversity framework (preprint).
Preprints. https://doi.org/10.22541/au.166568699.98478640/v1

Rodrigues, A.S.L., Akçakaya, H.R., Andelman, S.J., Bakarr, M.I., Boitani, L., Brooks, T.M.,
Chanson, J.S., Fishpool, L.D.C., Da Fonseca, G.A.B., Gaston, K.J., Hoffmann, M.,
Marquet, P.A., Pilgrim, J.D., Pressey, R.L., Schipper, J., Sechrest, W., Stuart, S.N.,
Underhill, L.G., Waller, R.W., Watts, M.E.J., Yan, X., 2004. Global Gap Analysis:
Priority Regions for Expanding the Global Protected-Area Network. BioScience 54,
1092.

Ruosch, M., Spahni, R., Joos, F., Henne, P.D., Van Der Knaap, W.O., Tinner, W., 2016.
Past and future evolution of *Abies alba* forests in Europe – comparison of a dynamic
vegetation model with palaeo data and observations. Glob. Change Biol. 22, 727–
740. https://doi.org/10.1111/gcb.13075

Santini, F., Andrisano, T., Leonardi, S., Ciaschetti, G., Labriola, M., Vendramin, G.G., Piotti,
A., 2018. Tracking the origin of silver fir plantations along the boundary between
different genetic clusters in central Apennines: Implications for their management.
For. Ecol. Manag. 408, 220–227. https://doi.org/10.1016/j.foreco.2017.10.045

Schweitzer, J.A., Fischer, D.G., Rehill, B.J., Wooley, S.C., Woolbright, S.A., Lindroth, R.L.,
Whitham, T.G., Zak, D.R., Hart, S.C., 2011. Forest gene diversity is correlated with
the composition and function of soil microbial communities. Popul. Ecol. 53, 35–46.
https://doi.org/10.1007/s10144-010-0252-3

Tapio, I., VäRv, S., Bennewitz, J., Maleviciute, J., Fimland, E., Grislis, Z., Meuwissen,
T.H.E., Miceikiene, I., Olsaker, I., Viinalass, H., Vilkki, J., Kantanen, J., 2006.
Prioritization for conservation of northern European cattle breeds based on analysis
of microsatellite data. Conserv. Biol. 20, 1768–1779. https://doi.org/10.1111/j.1523-
1739.2006.00488.x

Thomas, F.M., Blank, R., Hartmann, G., 2002. Abiotic and biotic factors and their
interactions as causes of oak decline in Central Europe. For. Pathol. 32, 277–307.
https://doi.org/10.1046/j.1439-0329.2002.00291.x

Vane-Wright, R.I., Humphries, C.J., Williams, P.H., 1991. What to protect?—Systematics
and the agony of choice. Biol. Conserv. 55, 235–254. https://doi.org/10.1016/0006-

3207(91)90030-D
von Takach, B., Penton, C.E., Murphy, B.P., Radford, I.J., Davies, H.F., Hill, B.M., Banks,
S.C., 2021. Population genomics and conservation management of a declining
tropical rodent. Heredity 126, 763–775. https://doi.org/10.1038/s41437-021-00418-9
von Takach, B., Sargent, H., Penton, C.E., Rick, K., Murphy, B.P., Neave, G., Davies, H.F.,
Hill, B.M., Banks, S.C., 2023. Population genomics and conservation management of
the threatened black-footed tree-rat (*Mesembriomys gouldii*) in northern Australia.
Heredity 130, 278–288. https://doi.org/10.1038/s41437-023-00601-0
Waples, R.S., Do, C., 2010. Linkage disequilibrium estimates of contemporary N_e using
highly variable genetic markers: a largely untapped resource for applied conservation
and evolution. Evol. Appl. 3, 244–262. https://doi.org/10.1111/j.1752-
4571.2009.00104.x
Wei, X., Jiang, M., 2020. Meta-analysis of genetic representativeness of plant populations
under *ex situ* conservation in contrast to wild source populations. Conserv. Biol. 35,
12–23. https://doi.org/10.1111/cobi.1361
Weitzman, M.L., 1998. The Noah's ark problem. Econometrica 66, 1279–1298.
Weitzman, M.L., 1993. What to preserve? An application of diversity theory to crane
conservation. Q. J. Econ. 108, 157–183.
Wilson, K.A., Cabeza, M., Klein, C.J., 2009. Fundamental concepts of spatial conservation
prioritization, in: Moilanen, A., Wilson, K.A., Possingham, H.P. (Eds.), Spatial
Conservation Prioritization: Quantitative Methods and Computational Tools. Oxford
University Press, Oxford, UK.
Xuereb, A., D'Aloia, C.C., Andrello, M., Bernatchez, L., Fortin, M., 2021. Incorporating
putatively neutral and adaptive genomic data into marine conservation planning.
Conserv. Biol. 35, 909–920. https://doi.org/10.1111/cobi.13609