Circadian coupling of mitochondria in a deep-diving mammal

Running title: Circadian mitochondria in a diving mammal

Chiara Ciccone[1], Fayiri Kante[1,2], Lars P. Folkow[1], David G. Hazlerigg[1], Alexander C. West[1],

Shona H. Wood[1*]

[1] Arctic Seasonal Timekeeping Initiative (ASTI), Arctic Chronobiology and Physiology

research group, Department of Arctic and Marine Biology, UiT – The Arctic University of

Norway, Tromsø NO-9037, Norway.

[2] Current address: Department of Biological Chemistry, Weizmann Institute of Science,

Rehovot 7610001, Israel.

[*]Corresponding author: shona.wood@uit.no

Keywords: diving mammals, hooded seal, biological clock, circadian, hypoxia, mitochondrial

respiration

SUMMARY STATEMENT

A functional clockwork and circadian variation in mitochondrial complex I activity is
demonstrated in skin fibroblasts from the deep diving hooded seal.

ABSTRACT

Regulation of mitochondrial oxidative phosphorylation is essential to match energy supply to
changing cellular energy demands, and to cope with periods of hypoxia. Recent work
implicates the circadian molecular clock in control of mitochondrial function and hypoxia
sensing. Since diving mammals experience intermittent episodes of severe hypoxia, with diel
patterning in dive depth and duration, it is interesting to consider circadian - mitochondrial
interaction in this group. Here we demonstrate that the hooded seal (*Cystophora cristata*), a
deep diving Arctic pinniped, shows strong daily patterning of diving behaviour in the wild.
Cultures of hooded seal skin fibroblasts exhibit robust circadian oscillation of the core clock
genes *per2* and *arntl*. In liver tissue collected from captive hooded seals, expression of *arntl*
was some 4-fold higher in the middle of night than in the middle of the day. To explore the
clock-mitochondria relationship, we measured the mitochondrial oxygen consumption in
synchronized hooded seal skin fibroblasts and found a circadian variation in mitochondrial
activity, with higher coupling efficiency of complex I coinciding with the trough of *arntl*
expression. These results open the way for further studies of circadian - hypoxia interactions
in pinnipeds during diving.

INTRODUCTION

The ability of animals to predict and prepare for daily changes in environmental demands relies on the presence of an intrinsic physiological system: the circadian clock. The molecular basis of the circadian clock in mammals is a transcription-translation negative feedback loop (TTFL). The positive limb of this loop depends on the dimerization of the transcriptional activators CLOCK and ARNTL (also known as BMAL1), while the negative limb depends on the formation of a transcriptionally repressive complex containing PER and CRY (Buhr and Takahashi, 2013). Circadian dynamics emerge through the autoregulatory effects of PER and CRY: the CLOCK/ARNTL complex activates transcription of PER and CRY which, after dimerization, translocate into the nucleus and repress their own transcription. This creates a loop with a period of around 24-hours and therefore called *circadian*.

One of the roles of the circadian clock is to coordinate the metabolic transcription network, presumably in order to optimize mitochondrial metabolism to the daily changes (Neufeld-Cohen et al., 2016; Turek et al., 2005). In mice, mitochondrial oxidative phosphorylation (OXPHOS) oscillates on a daily manner and in concomitance with the expression of some rate-limiting mitochondrial enzymes (Neufeld-Cohen et al., 2016). The OXPHOS rate depends on the activity of a group of enzymatic complexes located on the inner mitochondrial membrane (IMM) and together referred to as Electron Transport System (ETS). These complexes (from I to IV) are responsible for the creation of an electrochemical gradient of protons across the IMM and transport of electrons through it. At the level of complex IV, oxygen (O_2) is needed to act as electrons final acceptor, while the proton gradient is used by an ATP synthase (also indicated as complex V) to generate ATP (Chance and Williams, 1956). Variations in the activity of these complexes generate variations in mitochondrial OXPHOS and ATP production.

In mice with a dysfunctional circadian clock, the daily cycle of OXPHOS is abolished, suggesting that there is an intrinsic dependence of the mitochondrial respiration complexes on the circadian clockwork (Neufeld-Cohen et al., 2016). Deletion of *arntl* in C2C12 myotubes causes a reduction of both OXPHOS and extracellular acidification rate (an indicator of glycolytic rate), but deletion in embryonic stem cells increases OXPHOS while reducing glycolytic rate (Ameneiro et al., 2020; Peek et al., 2017). This suggests there is tissue specificity in the effect of "circadian clock knockout" on mitochondrial responses.

O_2, the final electron acceptor of the ETS, constitutes a fundamental factor in mitochondrial

regulation: hypoxia (O_2 deficiency) leads to a switch to anaerobic glycolysis from OXPHOS-

based metabolism (Kim et al., 2006; Semenza et al., 1994; Wang and Semenza, 1993). In

mice, interactions between the hypoxia inducible factor 1α (HIF-1α) and the circadian clock

optimize the mitochondrial response during hypoxic episodes (Adamovich et al., 2017;

Manella et al., 2020; Peek et al., 2017). However, these studies have only been conducted in

species that do not normally experience severe hypoxia (e.g., mouse and rat). It is therefore of

interest to investigate circadian – mitochondrial interactions in a diving mammal, which

through its behaviour frequently experiences intermittent episodes of severe hypoxia.

Diving mammals have evolved numerous hypoxia coping strategies (e.g., Blix, 2018; Davis,

2014), including enhanced O_2 storage and carrying capacity (with blood volume and

hemoglobin and myoglobin levels being higher than in other species), which combines with

strict O_2 economy by way of cardiovascular adjustments (bradycardia and peripheral

vasoconstriction) and hypometabolism (Blix et al., 1983; Burns et al., 2007; Lenfant et al.,

1970; Scholander, 1940). Nevertheless, they do still experience severe hypoxia, particularly in

connection with deep dives (Meir et al., 2009; Qvist et al., 1986).

There are several reports of seasonal and daily variations in diving behaviour of different

pinnipeds species (Bennett et al., 2001; Folkow and Blix, 1999; Nordøy et al., 2008;

Photopoulou et al., 2020). Among them, the hooded seal (*Cystophora cristata*) is known for

its remarkable diving capacity (Folkow and Blix, 1999): dives exceeding durations of 1 hour

and depths of 1600 meters have been recorded (Andersen et al., 2013; Vacquie-Garcia et al.,

2017), making this species a good research model for hypoxia tolerance and oxidative stress

in mammals (e.g., Fabrizius et al., 2016; Folkow et al., 2008; Geiseler et al., 2016; Geßner et

al., 2020; Hoff et al., 2016; Hoff et al., 2017; Mitz et al., 2009; Vázquez-Medina et al., 2011).

The hooded seal dive duration and depth are reportedly greater during the day than at night-

time (Andersen et al., 2013; Folkow and Blix, 1999), suggesting that there may be a daily

rhythm in exposure to severe hypoxia. However, the circadian molecular clock is

uncharacterised in pinnipeds, and mitochondrial activity has not been measured in this

species.

Here, we provide the first molecular characterisation of the circadian clock in the hooded seal

and explore the clock-mitochondria relationship, showing that mitochondrial activity relates

to circadian clock phase. These results open the way for further studies of circadian and

(diving-induced) hypoxia interactions in pinnipeds.

| 105 |
| 106 |

MATERIALS AND METHODS

Diving behaviour analysis

Data collected by the Norwegian Polar institute in 2007 and 2008 as part of the MEOP

(Marine Mammals Exploring the Ocean Pole to Pole) programme were provided by Dr. M.

Biuw and used for diving behaviour analysis based on dive data for 12 adult and 8 juvenile

hooded seals of both sexes (Vacquie-Garcia et al., 2017, 10.21334/npolar.2017.881dbd20, file

name: dive_2007.txt, dive_2008.txt). Data collection was funded by the Norwegian Research

Council (grant number 176477/S30) and the Norwegian Polar Institute. Conductivity-

Temperature-Depth Satellite relay Data Loggers (CTD-SRDLs) (Sea Mammal Research Unit,

University of St Andrews) were glued to the fur on the back of the neck of the seals. Data

were collected and transmitted for every 6-hour period and included the following variables:

dive duration, time of dive end, maximum depth, location (latitude and longitude), and time

spent at surface after the dive. Knowing dive duration, time of dive end and time spent at

surface, it was possible to infer the starting time for every single dive. Starting time was taken

to represent the start of a 'diving event', which then formed the basis for analyses of diurnal

changes in dive duration and depth.

Hooded seal pups perform shallower and shorter dives than adults (Folkow et al., 2010). Our

analyses therefore focused on the 12 adults (3 males and 9 females) that were represented in

the data set.

After having evaluated the dive duration distribution, we decided to analyse only dives

between 2 minutes and the 95% percentile of the maximum dive duration of each individual.

This allowed us to exclude outliers from further analysis.

All the following analysis were done in RStudio (4.2.1 version). To verify the presence of a

daily oscillation in the hooded seal diving behaviour, a cosine curve with a 24-h period was

fitted to the data through the function *cosinor.lm* in the package *cosinor*. To test whether the

cosine model significantly represented the data, we used the *cosinor.detect* function in the

package *cosinor2*. Calculations were repeated for each seal and for each month separately.

The function *cosinor* in the package *card* was used to define the Midline Statistic of Rhythm

(MESOR): this is a circadian rhythm-adjusted mean which gives an estimate of the average

value of an oscillating variable. The MESOR was used to identify different trends in diving duration throughout the year. Using the information about latitude and longitude, it was possible to calculate time of sunrise and sunset for each day of every month through the *sunriset* function in the *maptools* package.

To compare dive durations at different times of day we binned the data by hour and calculated the hourly mean dive duration.

Animals used for tissue sampling

Hooded seals (*Cystophora cristata*) were captured in their breeding colonies on the pack ice of the Greenland Sea, at ~71°N and ~019° W, during a research/teaching cruise with the R/V Helmer Hanssen in late March 2018, under permits of relevant Norwegian and Greenland authorities. Six live seals were brought to the Department of Arctic and Marine Biology (AMB) at UiT – The Arctic University of Tromsø, Norway, where they were maintained in a certified research animal facility (approval no. 089 by the Norwegian Food Safety Authority (NFSA)). Seals were euthanized (in 2019, as juveniles, at age ~10 mo.) for purposes other than the present study, in accordance with a permit issued by NFSA (permit no. 12268): the seals were sedated by intramuscular injection of zolazepam/tiletamine (Zoletil Forte Vet., Virbac S.A., France; 1.5 – 2.0 mg per kg of body mass), then anaesthetized using an endotracheal tube to ventilate lungs with 2 – 3 % isoflurane (Forene, Abbott, Germany) in air and, when fully anaesthetized, they were euthanized by exsanguination via the carotid arteries.

During another research cruise in March 2019, additionally 6 weaned pups were captured (NFSA; permit no. 19305) and brought to AMB, UiT. Before culling and tissue sampling (February 2020, at age 11-12 mo.), the animals were exposed to 18 days of 12 h light and 12 h darkness (12L;12D). In this experiment, we defined Zeitgeber Time 0 (ZT0) as the time when lights went on. Euthanasia was performed as explained above at ZT6 (light phase) for three seals and at ZT18 (dark phase) for the other three.

Tissue sampling

Skin biopsies were collected postmortem from the seal captured in March 2018 between the digits of the hind flipper with a biopsy punch (6 mm, 33-36-10, Miltex Inc., York) and processed for culturing of fibroblasts as described below. Samples from kidneys and liver were collected from the 6 seals held under 12L;12D light conditions. Tissues were minced

with a sterile scalpel blade and placed in 4 ml of RNAlater (AM7021, Thermo Fisher) in a

1:10 ratio. Samples were stored at 4°C for 24 h and then moved to -20°C until mRNA

extraction.

Culture of hooded seal skin fibroblasts

Skin biopsies were processed as described elsewhere (Du and Brown, 2021). Briefly, biopsies

were placed in collection medium (Dulbecco's Modified Eagle Medium (DMEM, D5796,

Sigma) + 50 % Fetal Bovine Serum (FBS, F7524, Sigma) + 1% penicillin-streptomycin (Pen-

Strep, P4458, Sigma)) and then moved into 6-well-plates containing digestion medium

(DMEM + 10% FBS + 1% amphotericin B (A2942, Sigma)) and 10% liberase (Roche, 05 401

119 001) for 8-9 hours at 37° C/5% CO_2. Tissue biopsies and digestion medium were then

transferred to a tube containing warm Dulbecco's Phosphate Buffered Saline (PBS, D8537,

Sigma), plates were rinsed again with PBS to collect all the tissue fragments. Samples were

centrifuged at 200 g for 5 minutes. The pellet was resuspended in culture medium (DMEM +

20% FBS + 0.1% gentamycin (15710, Invitrogen)) and placed in fresh 6-well-plates.

Fragments were overlaid with a Millicell Cell Culture insert (Cat. PICMORG50 Millipore):

1.5 ml of culture medium was added to the interior of the insert and 0.5 ml to the exterior.

Plates were incubated for about 2 weeks at 37° C/5% CO_2. Medium was changed every 3-4

days.

When cells covered approximately 80% of the plates surface (80% confluency), they were

trypsinised (Trypsin-EDTA solution, T4049, Sigma), resuspended in culture media and

centrifuged at 400 g for 5 minutes. Pelleted cells were then suspended in 3 ml of freezing

medium per T75 culture flask (culture medium + 10% Dimethyl sulfoxide (DMSO, D5879,

Sigma)) and progressively cooled in a cryocooler overnight at - 80° C. Cells were then moved

to liquid nitrogen for long-term storage.

A few days before the start of the experimental procedures (temperature cycling and

respirometry) cells were thawed at 37°C for 1-2 minutes and then added to 10 ml of pre-

warmed medium (DMEM + 20% FBS + 1% Pen-Strep). The solution was centrifuged at 300

g for 5 minutes. The pellet was resuspended in 12 ml of medium and transferred to a T75

culture flask. When cells reached 80 % confluency, they were trypsinised and re-plated into

three different plates (1 T175, 1 T75 and 1 6-well-plate) for each sampling timepoint.

Temperature cycling treatment

Cells were exposed to five consecutive 24h-cycles of temperature alternations between 12-h
at 36.5°C and 12-h at 39.5°C. In all the temperature cycling experiments, we defined ZT0 as
the time of the transition from 36.5 to 39.5°C. Cells were collected from the sixth cycle with
4-hour intervals over 2 cycles (Fig. S1A). For mRNA, 6-well plates were washed with PBS
and directly frozen on dry ice before storage at - 80°C. For the mitochondrial O_2 consumption
measurements, cells grown in T175 flasks were sampled with an 8-hour interval, thus giving
data around 1 full 24h-cycle with only one repetition for each sampling timepoint (Fig. S1A).
The presence of phenol red in the culture medium was used as standard pH indicator in the
cultures: if changes in pH were detected, the medium was changed at the end of the third
cycle.

In a second experiment, cells were kept under temperature cycle for 3 cycles. Cells were
sampled over a 48-hour period, at 36.5°C for the first 12 hours and at 39.5°C for the
remaining 36 hours (Fig. S1B).

mRNA extraction, cDNA conversion

For both tissues and cells, RNA was extracted using RNeasy mini kit (74104, Qiagen)
following the manufacturer's instructions. Tissue samples (~100mg) were thawed then
inserted in a low bind tube containing 600 µl RLT buffer + 6 µl β-mercaptoethanol with a
metal bead. The tube was shaken in a Tissue Lyser for 6 minutes at 20 shakes/second. Cell
plates were taken out of the - 80°C freezer and the contents of each well was collected in 350
µl of RLT buffer before being transferred to a QIAshredder column (79654, Qiagen). The
column was centrifugated for 2 minutes at full speed and 350 µl of 70% ethanol were added
to the lysate. Lysates were treated with DNase I (79254, Qiagen) and heated to 37°C for 10
minutes. Enzymatic reaction was stopped by adding EDTA at a final concentration of 0.05 M
in each tube at 75°C for 10 minutes. Samples were then treated with cDNA buffer and enzyme
reverse transcriptase, heated for 1 hour at 37°C and cooled down to 4°C. The cDNA samples
were then stored at - 20°C until further use.

qPCR

Seal gene sequences for *arntl*, *per2* and housekeeping genes *tbp* and *ppib* were identified by
aligning in BLAST dog (*Canis lupus*) genes sequences against the Weddell seal
(*Leptonychotes weddellii*) genome
(https://www.ncbi.nlm.nih.gov/assembly/GCF_000349705.1/). Primer specificity was
confirmed by cloning PCR amplicons into Zero Blunt TOPO vector and sanger dideoxy

sequencing (BigDye). Only primers with efficiency above 90% and a single product were
used (Table 1). Reagents from the Promega GoTaq qPCR Master Mix kit (A6001, Promega)
were used to perform qPCR. The Bio-Rad manager software was used to control the Bio-Rad
CFX Connect Real_Time PCR system. The qPCR template used started with 2 minutes at
95°C. It then repeated 39 temperature cycles, each of: 15 seconds at 95°C, 15 seconds at 57°C
and 1 minute at 60°C, where fluorescence was measured. It finished with 5 seconds at 65°C
and then a final constant temperature of 95°C. The cycle threshold (CT) values of each gene
were analysed using the $\Delta\Delta$CT ($2^{-\Delta\Delta Ct}$) method (Livak and Schmittgen, 2001) against
housekeeping genes *ppib* and *tbp*.

Mitochondrial oxygen consumption measurements

Mitochondrial oxygen consumption was measured with two O2k high-resolution
respirometers (Oroboros Instruments, Innsbruck, Austria) and data were recorded in real-time
using the Oroboros DatLab software (Oroboros Instruments).

Due to the duration of the respiration measurement protocol, and limited availability of
Oroboros respirometers, it was possible to process samples only six times over 48 hours. After
exposure to temperature cycling treatment, cells were collected by trypsinisation, resuspended
in culture medium, and centrifuged at 300 g for 5 minutes. Pellet was then resuspended in the
respiration medium MiR05 (0.5 mM EGTA, 3 mM $MgCl_2$, 60 mM K-lactobionate, 20 mM
taurine, 10 mM KH_2PO_4, 20 mM HEPES, 110 mM sucrose, 1g/L BSA; pH = 7). Cells were
added to the Oxygraph chambers at a concentration of 0.6×10^6 cells/ml with a final volume of
2.1 ml in each chamber.

After measurement of basal respiration (Routine), cells were permeabilised with digitonin (3
µg/ml). Different mitochondrial states were assessed by injecting the following chemicals in
the chambers: pyruvate, malate, ADP, glutamate, succinate, cytochrome c, rotenone,
oligomycin and carbonyl cyanide p-trifluoro-methoxyphenyl hydrazone (FCCP). Optimal
concentration of both digitonin and FCCP were determined in pilot experiments (data not
shown). Table 2 gives an overview of the chemicals injected and the respiratory states
measured.

Each experiment was done with four replicate respiratory chambers, except for ZT13 and
ZT21, which have three replicates. The fourth replicate was excluded because the cells did not
permeabilize, therefore the mitochondria did not respond to the treatments.

The *leak state* (or *state 4*) is defined as the mitochondrial respiration in the presence of fuel

substrates and in the absence of ADP. It reflects proton leak across the inner membrane that

does not result in ATP production (Chance and Williams, 1956; Gnaiger, 2009; Perry et al.,

2013). It can be used to determine the coupling efficiency of the different ETS enzymes

through the respiratory control ratio (RCR), measured as OXPHOS/LEAK (P/L), where

OXPHOS (also referred to as *state 3*) corresponds to mitochondrial respiration in the presence

of substrates and saturating ADP (Chance and Williams, 1956; Gnaiger, 2009). In the

literature, the RCR is preferably expressed as OXPHOS coupling efficiency $(1 - RCR^{-1})$, with

values between 0 and 1, where 1 corresponds to fully coupled mitochondria (Doerrier et al.,

2018; Gnaiger, 2020). Accordingly, we calculated complex I coupling efficiency using this

formula:

272
$$1 - (^{PM - L}/_{PM - P})\ \text{(Eq. 1)}$$

273 Where PM-L corresponds to the complex I *leak state* and PM-P to complex I OXPHOS (Table

274 2).

Lentiviral vector construction

Circadian clock gene activity was recorded using the vectors pLV6-Bmal-luc (Addgene

plasmid #68833) and pLV6-Per2-luc. The latter consisted of the pLV6 backbone and a mouse

per2 promoter with adjacent luciferase sequence contained in the pGL3 basic E2 vector

(Addgene plasmid 48747). To ligate the Per2:luciferase reporter with pLV6 backbone we

designed a restriction cloning approach shown to be efficient in large plasmids using the

QuickChange Lightning Site-Directed Mutagenesis (SDM) kit (Agilant, 210518) according to

manufacturer's instructions (Zhang and Tandon, 2012). Briefly, the SDM kit was used to

create an extra BamHI cutting site 5' on the pLV6-Bmal-luc vector. Mutated plasmid was

transformed with stable competent *E. coli* (NEB, C3040H) according to the manufacture's

manual. Cells were grown on LB agar plates (40g/l LB-agar (Sigma-Aldrich, L3147), 100

µg/ml Ampicillin (Sigma-Aldrich, A9393)) and single clones were cultivated in LB broth

overnight (25g LB (Sigma-Aldrich, L3522), 100 µg/ml Ampicillin). We used a Qiagen

miniprep kit (Qiagen, 27104) to extract plasmid DNA and the mutation was confirmed

through whole-plasmid sequencing (Plasmidsaurus, https://www.plasmidsaurus.com/). After

the mutation was detected, the pLV6-Bmal-luc vector was digested with BamHI (NEB,

R3136S) then treated with quick CIP (NEB, M0525S) to avoid self-ligation. The digest was

run on an 1% Agarose gel (TAE buffer, 0.25 µg/ml EtBr) and the the QIAquick Gel extraction

kit (Qiagen, 28704) was used to extract the pLV6 backbone according to the manufacturer's

instructions. Since the Per2 promoter and luciferase in the pGL3-Per2-luc vector is already

flanked by BamHI cutting sites, the pGL3-Per2-luc vector was directly digested with BamHI

for Per2:luciferase sequence extraction.

Finally, the pLV6 backbone and the Per2:luciferase fragment were ligated with T4 DNA

ligase (NEB, M0202S) according to the manufacture's manual. Vectors extracted from

transformed single colonies were sequenced by Plasmidsaurus and checked for correct

ligation and sequence. The resulting vector is referred to as pLV6-Per2-luc.

Lentiviral vectors were constructed according to (Du and Brown, 2021). Briefly, HEK293T

cells were incubated with pMD2G (#12259, Addgene), psPAX2 (#12260, Addgene), and the

two transfer vectors. Supernatant was collected 24 and 48 hours after incubation and

transfection and filtered through a 0.2 µm filter. Filtered supernatant was aliquoted and frozen

at -80°C until use.

Generation of a stable cell line expressing lentiviral vector and bioluminescence recordings

Two ml of medium containing lentivirus were added to a T25 flask containing hooded seal

fibroblasts at 50% confluency, with 3 ml of DMEM + 20% FBS. Cells were incubated

overnight and then the media was changed to clean cells from the virus. After 3 days, medium

was changed to DMEM + 20% FBS + 10 µg/ml blasticidin (R21001, Gibco) selection

antibiotic (cells were passaged in a T75 flask at this point). From the T75 flask, 10 3.5-cm-

wells were seeded and cultured in DMEM + 20% FBS + 10 µg/ml blasticidin. One week

before starting the temperature cycling, medium was changed to recording medium (DMEM

without phenol red + 0.47% $NaCO_3$ + 1% HEPES buffer + 0.25 % Pen-Strep + 5% FBS + 10

µg/ml blasticidin + 0.1 mM luciferin) (Yamazaki and Takahashi, 2005). Prior to the

experiment, medium was again changed, and wells were sealed with parafilm and placed in a

Photon multiplier tube (Lumicycle, Hamamatsu) for recording.

On a separate experiment, after a 2-days incubation at 37° C/5% CO_2 in recording medium,

transfected cells were synchronized with dexamethasone (100 nM, D4902, Sigma) for 30 min.

Thereafter, cells were washed twice with PBS and medium was changed to normal recording

medium. Cells were then sealed and placed in the Photon multiplier tube for recording.

Statistical analysis

All statistical analyses were performed in RStudio using one-way ANOVA and post
comparison Tukey HSD test. The analysis of circadian oscillations for clock genes mRNA
expression was done using JTK cycle, a non-parametric algorithm designed to detect cyclic
patterns in datasets with regular intervals between measures (Hughes et al., 2010). All data are
represented as mean ± standard error (SEM). Periods of the photon multiplier tube recordings
were analysed by fitting a damped sine wave in GraphPad Prism 8 (version 8.0.2). Principal
Component Analysis (PCA) of ZTs was performed across the 7 measured mitochondrial states
using the *prcomp* function in RStudio. Ellipses were drawn with a 0.80 confidence interval.

RESULTS

Seasonal and daily variations in hooded seal diving behaviour

We used the data collected between 2007 and 2008 by the Norwegian Polar Institute (see Vacquie-Garcia et al., 2017) to perform a high time resolution analysis of hooded seal diving behaviour. Initial analysis revealed that the maximum dive duration was 87.25 minutes for males (at ~09:00 in January 2007, adult male Mj98) and 53.25 minutes for females (at ~10:00 in April 2008, adult female F175). For most seals, cosinor analysis of diving duration data showed that there is a significant relationship between dive duration and time of the day with a 24-h period throughout the year (Table 3). Figure 1 shows this relationship for a male adult hooded seal (ID: M171, August 2007 – April 2008). While dives were observed in the night, the average dive duration was consistently higher during daytime (05:00 – 16:00) for most of the months, and very few night-time dives were longer than 1 hour (Fig. 2). Therefore, by investigating the hourly distribution of diving durations, we were able to confirm the presence of a diurnal pattern in the hooded seal diving behaviour (Fig. 1, 2).

Despite the limited amount of data, we also wished to investigate seasonal changes. We observed that, in accordance to previous data (Folkow and Blix, 1999), the mean value based on the daily distribution of values (MESOR) shows that dives were longer during the winter than in the summer (Fig.S2). Specifically, hourly mean dive duration was particularly short in July and August (Fig. 2) but increased again in September. In October, there was a clear pattern of shorter dives at evening and night and longer dives during the day (Fig. 2), which was persistent throughout all the winter months, until February. In March, dive duration decreased and dives were also less frequent, most likely linked to behavioural changes in connection with the breeding season (Kovacs et al., 1996).

Circadian molecular clock function in hooded seal fibroblasts and tissues

We derived primary skin fibroblasts from hooded seals and cultured them under a 24-h temperature cycle, to synchronise the cells (Buhr et al., 2010). Measuring the endogenous mRNA abundance of the clock genes *arntl* and *per2*, we show a significant oscillation with a period of 24 hours (JTK cycle, $p<0.0001$) and antiphase rhythms to one another: *arntl* expression peaking in middle of the low temperature phase and *per2* expression peaking in the middle of the high temperature phase (Fig. 3A). To determine whether these oscillations persist in constant conditions, we synchronised the cells using a temperature cycle and then

held temperature constant for 36 hours. Both *arntl* and *per2* endogenous mRNA levels
continued to oscillate in anti-phase (Fig. 3B).

We then generated lentiviral reporters for *arntl* and *per2* to stably transduce our primary
fibroblasts and allow real time monitoring of clock gene expression for multiple days. After
synchronisation with dexamethasone, we observed significant oscillations of approximately
24 hours for both *arntl* and *per2* (R^2 = 0.96 and 0.95, respectively) and persistent oscillations
for 3 days post synchronisation (Fig. 3C). These results are consistent with a functional
molecular clockwork in hooded seal fibroblasts.

We also measured endogenous *arntl* mRNA expression in liver and kidney samples collected
in the mid-light (ZT6) and mid-dark phase (ZT18) from captive hooded seals held on a
LD12:12 light-dark cycle. *Arntl* expression in the liver showed a significant time-of-day
effect, with expression at ZT18 being some four-fold higher than at ZT6 ($p<0.05$; Fig. 3D).

Circadian clock phase relationship with the metabolic capacity of hooded seal mitochondria

To investigate whether there is a time-of-day variation in mitochondrial respiration, we used a
O2k high-resolution respirometers (Oroboros) substrate-uncoupler-inhibitor titration (SUIT)
protocol (see Table 2) and primary skin fibroblasts from the hooded seal. We sampled at 6
different zeitgeber times (ZT) across a 48-h temperature cycling experiment (Fig. S1A) and
found significant differences between timepoints (one-way ANOVA analysis; Fig. S3). We
used PCA analysis to integrate values for the 7 parameters measured by the respirometers for
each individual sample taken at each time-point. This generated 2 principal components (PC1
and PC2), which together accounted for over 90% of the overall variance for these 7
parameters across the study as a whole (Fig. 4A). There was wide ZT-dependent variation,
and we speculated that this variation was related to predicted phases for *arntl* and *per2*
expression. Based on the endogenous mRNA expression (Fig. 3A) we focused on ZT5 (low
arntl, high *per2* expression) and ZT17 (high *arntl*, low *per2* expression), which separated
more clearly on the PCA plot (Fig. 4A). The mitochondrial state that differed the most
between ZT5 and ZT17 was the *leak state* at complex I (PM-L in Table 2). We therefore
calculated complex I coupling efficiency using Eq. 1 across all the ZTs.

This revealed the presence of a time effect on complex I efficiency ($p<0.01$; fig. 4B), with
lowest values at ZT21 (0.46 ± 0.017). High coupling efficiencies (close to 1) indicate highly

coupled mitochondria (Doerrier et al., 2018; Gnaiger, 2020) and changes in coupling

efficiency at complex I suggest a time of day effect on mitochondrial function.

To provide independent verification of the relationship between mitochondrial activity and

clock phase we collected samples from virally transduced *arntl:*luciferase-reporter-expressing

hooded seal fibroblasts at phases of peak and trough *arntl* expression and processed them for

respirometry analysis (Fig. 4C). Calculation of complex I coupling efficiency revealed a

lower oxidative phosphorylation activity at complex I at peak *arntl* expression (0.40 ± 0.024)

than at trough (0.52 ± 0.017)($p<0.001$; Fig. 4D). Therefore, when *arntl* is at its peak, complex

I is not as efficiently coupled as when *arntl* is at its trough, indicating a clock-dependent

variation of complex I coupling efficiency. These difference in coupling efficiency likely

relate to a switch from an OXPHOS-based metabolism to a less oxidative metabolism.

DISCUSSION

The present study confirms that under natural conditions hooded seal diving behaviour is highly time of day dependent, and further demonstrates that under laboratory conditions, hooded seal cells show robust circadian characteristics, which are associated with cyclical changes in mitochondrial activity. Daily physiological and molecular changes are a key factor for determining the ability of different species to overcome the challenges imposed by a continuously changing environment. For the hooded seal the biggest challenge probably is the restricted access to oxygen during long and deep dives.

We report that dives were consistently longer during the day and shorter at night and that there was a 24-h period in the diving duration pattern. Such daily variations are most likely related to similar patterns in the behaviour of hooded seal prey (e.g., by redfish (*Sebastes spp.*) and squid (*Gonatus spp.*)), which can display diurnal vertical migrations as their prey, in turn, migrate according to the photic state in the water column (Kristensen, 1984; Torsvik et al., 1995). Even though the day-night trend is quite consistent, there is some seasonal variation, as previously reported (Andersen et al., 2013). These variations relate in part to the breeding season (late March) and the moulting season (June/July), during which adult hooded seals spend more time hauled out (Folkow and Blix, 1999; Folkow et al., 1996; Kovacs et al., 1996; Øritsland, 1959). However, dives also tend to generally be deeper and longer in winter than in summer (Folkow & Blix, 1999). The time-of-day dependent diving behaviour led us to investigate whether a circadian clock – mitochondria coupling might be present in hooded seal cells.

Here, we show that circadian changes in *arntl* expression relate to a switch from an OXPHOS-based metabolism to a less oxidative metabolism, via regulation of complex I. We propose that through this regulatory system, the ability to spend time underwater searching for food will be maximised when it is needed the most - i.e., during the day, when the seals forage species are at greater depths.

In support of this we show, by two independent methods, that primary hooded seal skin fibroblasts display antiphase circadian oscillations of the clock genes *arntl* and *per2*, a result consistent with other characterisations of the mammalian circadian clock (Buhr and Takahashi, 2013). These oscillations are circadian because they persist under constant conditions (temperature, in this study) and with a period of approximately 24 hours (Dunlap et

al., 2004). We also show that the activity of complex I varies according to time-of-day and to
the circadian clock phase defined by *arntl*.

Diurnal oscillations in complex I activity have been documented in mice liver (Jacobi et al.,
2015; Neufeld-Cohen et al., 2016) and in human skeletal muscle (van Moorsel et al., 2016). In
the human cell line HepG2, *arntl* expression regulates complex I activity by a process of
acetylation and deacetylation through the NAMPT-NAD-SIRT1/3 machinery (Cela et al.,
2016).

Finally, we show a time-of-day difference in *arntl* gene expression in the hooded seal liver,
consistent with previous recordings in freshly isolated mice livers (Jacobi et al., 2015;
Manella et al., 2020). However, we observed no difference in the kidneys, but we were
limited to only two time points so no strong conclusions regarding tissue-specific circadian
rhythmicity can be drawn. We wanted to consider those organs specifically, because they are
known to undergo substantial reduction in their arterial blood supply (by >80%) – and, hence,
in the supply of blood-borne O_2 - in connection with long-duration diving in seals (e.g. Blix et
al., 1983; Zapol et al., 1979). Therefore, the liver and kidney appear to be particularly
hypoxia-/ischemia-tolerant (e.g., Hochachka et al., 1988), from which follows that a
potentially tissue-dependent circadian clock phasing and interaction with mitochondria may
be expected to be found in these particular tissues. However, further studies are needed to
investigate this.

In summary, one of the major functions of the circadian clock is to modulate metabolic
responses through regulation of mitochondrial activity (for a review see: Puig et al., 2018).
Exposure to hypoxia has been documented to have an effect on both mitochondrial activity
and circadian clock function (Adamovich et al., 2017; Kim et al., 2006; Manella et al., 2020;
Papandreou et al., 2006). Specifically, it appears that the interaction between HIF-1α and
ARNTL is at the basis of the clock-modulated hypoxia-response (Peek et al., 2017). In mice,
during hypoxia, the dimer HIF-1 is responsible for the switch from a more OXPHOS-based
metabolism towards anaerobic glycolysis (Kim et al., 2006; Semenza et al., 1994; Wang and
Semenza, 1993). Furthermore, changes in O_2 levels reset the circadian clock through the
interaction of the clock genes with the hypoxia inducible factor 1α (HIF-1α) (Adamovich et
al., 2017). Potentially, our observation of a switch to less oxidative metabolism at daytime in
the seal fibroblast is an anticipatory/adaptive switch in prospect of more severely hypoxic
conditions due to deeper and longer day-time diving. Our study lays the groundwork for

further investigations into the roles of HIF-1, the circadian clock and mitochondrial

metabolism in diving mammals.

ACKNOWLEDGMENTS

The authors would like to thank the animal technicians of the Arctic Chronobiology and Physiology group for all the help with the animal handling: Hans Lian, Hans-Arne Solvang, and Renate Thorvaldsen. We would also like to thank Dr. Martin Biuw for his help in the initial stages of the dive data analyses and Dr. Daniel Appenroth for his work on the lentiviral reporter.

COMPETING INTERESTS

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Conceptualization: S.H.W., A.C.W, D.G.H., C.C.; Methodology: A.C.W., F.K., C.C.; Software: C.C.; Validation: F.K., C.C.; Formal analysis: C.C., D.G.H., A.C.W.; Investigation: C.C., F.K., A.C.W., S.H.W.; Resources: S.H.W., L.P.F.; Data curation: C.C.; Writing-original draft: C.C., D.G.H.; Writing – review and editing: D.G.H., C.C., S.H.W., A.C.W., L.P.F., F.K.; Visualization: C.C., D.G.H., A.C.W.; Supervision: L.P. F., S.H.W., A.C.W., D.G.H.; Project administration: S.H.W., L.P.F; Funding acquisition: S.H.W.

FUNDING

The work was supported by grants from the Tromsø ForskningsStiftelse (TFS) starter grant TFS2016SW and the TFS infrastructure grant (IS3_17_SW) awarded to S.H.W., the Arctic Seasonal Timekeeping Initiative (ASTI) grant and UiT strategic funds support to D.G.H., S.H.W., & A.C.W.

 DATA AVAILABILITY

Code used for the analysis and data are available at https://github.com/ShonaWood/SealClock

REFERENCES

Adamovich, Y., Ladeuix, B., Golik, M., Koeners, M. P. and Asher, G. (2017). Rhythmic
Oxygen Levels Reset Circadian Clocks through HIF1α. *Cell Metab.* **25**, 93–101.

**Ameneiro, C., Moreira, T., Fuentes-Iglesias, A., Coego, A., Garcia-Outeiral, V.,
Escudero, A., Torrecilla, D., Mulero-Navarro, S., Carvajal-Gonzalez, J. M.,
Guallar, D., et al.** (2020). BMAL1 coordinates energy metabolism and differentiation of
pluripotent stem cells. *Life Sci. Alliance* **3**, 1–15.

**Andersen, J. M., Skern-Mauritzen, M., Boehme, L., Wiersma, Y. F., Rosing-Asvid, A.,
Hammill, M. O. and Stenson, G. B.** (2013). Investigating annual diving behaviour by
hooded seals (cystophora cristata) within the northwest atlantic ocean. *PLoS One* **8**,.

Bennett, K. A., McConnell, B. J. and Fedak, M. A. (2001). Diurnal and seasonal variations
in the duration and depth of the longest dives in southern elephant seals (Mirounga
leonina): Possible physiological and behavioural constraints. *J. Exp. Biol.* **204**, 649–662.

Blix, A. S. (2018). Adaptations to deep and prolonged diving in phocid seals. *J. Exp. Biol.*
221,.

Blix, A. S., Elsner, R. and Kjekshus, J. K. (1983). Cardiac output and its distribution
through capillaries and A□V shunts in diving seals. *Acta Physiol. Scand.* **118**, 109–116.

Buhr, E. D. and Takahashi, J. S. (2013). Molecular components of the mammalian
circadian clock. *Handb. Exp. Pharmacol.* **217**, 3–27.

Buhr, E. D., Yoo, S.-H. and Takahashi, J. S. (2010). Temperature as a Universal Resetting
Cue for Mammalian Circadian Oscillators. *Science (80-.).* **330**, 379–385.

Burns, J. M., Lestyk, K. C., Folkow, L. P., Hammill, M. O. and Blix, A. S. (2007). Size
and distribution of oxygen stores in harp and hooded seals from birth to maturity. *J.
Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **177**, 687–700.

**Cela, O., Scrima, R., Pazienza, V., Merla, G., Benegiamo, G., Augello, B., Fugetto, S.,
Menga, M., Rubino, R., Fuhr, L., et al.** (2016). Clock genes-dependent acetylation of
complex I sets rhythmic activity of mitochondrial OxPhos. *Biochim. Biophys. Acta -
Mol. Cell Res.* **1863**, 596–606.

Chance, B. and Williams, R. (1956). The respiratory chain and oxidative phosphorylation.

Adv. Enzymol. Relat. Areas Mol. Biol. **17**,.

Davis, R. W. (2014). A review of the multi-level adaptations for maximizing aerobic dive duration in marine mammals: From biochemistry to behavior. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **184**, 23–53.

Doerrier, C., Garcia-souza, L. F., Krumschnabel, G., Gnaiger, E. and Wohlfarter, Y. (2018). *High-Resolution FluoRespirometry and OXPHOS Protocols for Human Cells, and Isolated Mitochondria.*

Du, N.-H. and Brown, S. A. (2021). Measuring Circadian Rhythms in Human Cells.pp. 53–67.

Dunlap, J. C., Loros, J. J. and DeCoursey, P. J. (2004). *Chronobiology: Biological timekeeping.* Sinauer Associates.

Fabrizius, A., Hoff, M. L. M., Engler, G., Folkow, L. P. and Burmester, T. (2016). When the brain goes diving: Transcriptome analysis reveals a reduced aerobic energy metabolism and increased stress proteins in the seal brain. *BMC Genomics* **17**,.

Folkow, L. P. and Blix, A. S. (1999). Diving behaviour of hooded seals (Cystophora cristata) in the Greenland and Norwegian seas. *Polar Biol.* **22**, 61–74.

Folkow, L. P., Mårtensson, P. E. and Blix, A. S. (1996). Annual distribution of hooded seals (Cystophora cristata) in the Greenland and Norwegian seas. *Polar Biol.* **16**, 179–189.

Folkow, L. P., Ramirez, J. M., Ludvigsen, S., Ramirez, N. and Blix, A. S. (2008). Remarkable neuronal hypoxia tolerance in the deep-diving adult hooded seal (Cystophora cristata). *Neurosci. Lett.* **446**, 147–150.

Folkow, L. P., Nordøy, E. S. and Blix, A. S. (2010). Remarkable development of diving performance and migrations of hooded seals (Cystophora cristata) during their first year of life. *Polar Biol.* **33**, 433–441.

Geiseler, S. J., Larson, J. and Folkow, L. P. (2016). Synaptic transmission despite severe hypoxia in hippocampal slices of the deep-diving hooded seal. *Neuroscience* **334**, 39–46.

Geßner, C., Stillger, M. N., Mölders, N., Fabrizius, A., Folkow, L. P. and Burmester, T. (2020). Cell Culture Experiments Reveal that High S100B and Clusterin Levels may

Convey Hypoxia-tolerance to the Hooded Seal (Cystophora cristata) Brain. *Neuroscience* **451**, 226–239.

Gnaiger, E. (2009). Capacity of oxidative phosphorylation in human skeletal muscle. New perspectives of mitochondrial physiology. *Int. J. Biochem. Cell Biol.* **41**, 1837–1845.

Gnaiger, E. (2020). *Mitochondrial Pathways and Respiratory Control. An introduction to OXPHOS Analysis*. 5th ed. Innsbruck: Bioenergetics Communications 2020.

Hochachka, P. W., Castellini, J. M., Hill, R. D., Schneider, R. C., Bengtson, J. L., Hill, S. E., Liggins, G. C. and Zapol, W. M. (1988). Protective metabolic mechanisms during liver ischemia: Transferable lessons from long-diving animals. *Mol. Cell. Biochem.* **84**, 77–85.

Hoff, M. L. M., Fabrizius, A., Folkow, L. P. and Burmester, T. (2016). An atypical distribution of lactate dehydrogenase isoenzymes in the hooded seal (Cystophora cristata) brain may reflect a biochemical adaptation to diving. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **186**, 373–386.

Hoff, M. L. M., Fabrizius, A., Czech-Damal, N. U., Folkow, L. P. and Burmester, T. (2017). Transcriptome analysis identifies key metabolic changes in the hooded seal (Cystophora cristata) brain in response to hypoxia and reoxygenation. *PLoS One* **12**, 1–21.

Hughes, M. E., Hogenesch, J. B. and Kornacker, K. (2010). JTK-CYCLE: An efficient nonparametric algorithm for detecting rhythmic components in genome-scale data sets. *J. Biol. Rhythms* **25**, 372–380.

Jacobi, D., Liu, S., Burkewitz, K., Gangl, M. R., Mair, W. B., Lee, C., Unluturk, U., Li, X., Kong, X., Hyde, A. L., et al. (2015). Hepatic Bmal1 Regulates Rhythmic Mitochondrial Dynamics and Promotes Metabolic Fitness Article Hepatic Bmal1 Regulates Rhythmic Mitochondrial Dynamics and Promotes Metabolic Fitness. *Cell Metab.* **22**, 709–720.

Kim, J. W., Tchernyshyov, I., Semenza, G. L. and Dang, C. V. (2006). HIF-1-mediated expression of pyruvate dehydrogenase kinase: A metabolic switch required for cellular adaptation to hypoxia. *Cell Metab.* **3**, 177–185.

Kovacs, K. M., Lydersen, C., Hammill, M. and Lavigne, D. M. (1996). Reproductive effort

of male hooded seals (Cystophora cristata): Estimates from mass loss. *Can. J. Zool.* **74**, 1521–1530.

Kristensen, T. (1984). Biology of the squid Gonatus fabricii (Lich- tenstein, 1818) from West Greenland waters. *Medd Grùn Biosci* **13**, 17.

Lenfant, C., Johansen, K. and Torrance, J. D. (1970). Gas transport and oxygen storage capacity in some pinnipeds and the sea otter. *Respir. Physiol.* **9**, 277–286.

Livak, K. J. and Schmittgen, T. D. (2001). Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2 C T Method. *METHODS* **25**, 402–408.

Manella, G., Aviram, R., Bolshette, N., Muvkadi, S., Golik, M., Smith, D. F. and Asher, G. (2020). Hypoxia induces a time- And tissue-specific response that elicits intertissue circadian clock misalignment. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 779–786.

Meir, J. U., Champagne, C. D., Costa, D. P., Williams, C. L. and Ponganis, P. J. (2009). Extreme hypoxemic tolerance and blood oxygen depletion in diving elephant seals. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **297**, 927–939.

Mitz, S. A., Reuss, S., Folkow, L. P., Blix, A. S., Ramirez, J. M., Hankeln, T. and Burmester, T. (2009). When the brain goes diving: glial oxidative metabolism may confer hypoxia tolerance to the seal brain. *Neuroscience* **163**, 552–560.

Neufeld-Cohen, A., Robles, M. S., Aviram, R., Manella, G., Adamovich, Y., Ladeuix, B., Nir, D., Rousso-Noori, L., Kuperman, Y., Golik, M., et al. (2016). Circadian control of oscillations in mitochondrial rate-limiting enzymes and nutrient utilization by PERIOD proteins. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E1673–E1682.

Nordøy, E. S., Folkow, L. P., Potelov, V., Prischemikhin, V. and Blix, A. S. (2008). Seasonal distribution and dive behaviour of harp seals (Pagophilus groenlandicus) of the White Sea-Barents Sea stock. *Polar Biol.* **31**, 1119–1135.

Øritsland, T. (1959). The hooded seal (in Norwegian). *Fauna Oslo* **12**, 70–90.

Papandreou, I., Cairns, R. A., Fontana, L., Lim, A. L. and Denko, N. C. (2006). HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab.* **3**, 187–197.

Peek, C. B., Levine, D. C., Cedernaes, J., Taguchi, A., Kobayashi, Y., Tsai, S. J., Bonar,

N. A., McNulty, M. R., Ramsey, K. M. and Bass, J. (2017). Circadian Clock

Interaction with HIF1α Mediates Oxygenic Metabolism and Anaerobic Glycolysis in

Skeletal Muscle. *Cell Metab.* **25**, 86–92.

Perry, C. G. R., Kane, D. A., Lanza, I. R. and Neufer, P. D. (2013). Methods for assessing

mitochondrial function in diabetes. *Diabetes* **62**, 1041–1053.

Photopoulou, T., Heerah, K., Pohle, J. and Boehme, L. (2020). Sex-specific variation in

the use of vertical habitat by a resident Antarctic top predator: Sex-specific variation in

diving. *Proc. R. Soc. B Biol. Sci.* **287**,.

Puig, L. S., Valera-alberni, M., Cantó, C. and Pillon, N. J. (2018). Circadian Rhythms and

Mitochondria⎕: Connecting the Dots. **9**, 1–8.

Qvist, J., Hill, R. D., Schneider, R. C., Falke, K. J., Liggins, G. C., Guppy, M., Elliot, R.

L., Hochachka, P. W. and Zapol, W. M. (1986). Hemoglobin concentrations and blood

gas tensions of free-diving Weddell seals. *J. Appl. Physiol.* **61**, 1560–1569.

Scholander, P. F. (1940). Experimental investigations on the respiratory function in diving

mammals and birds. *I kommisjon hos Jacob Dybwad* **20**,.

Semenza, G. L., Roth, P. H., Fang, H. M. and Wang, G. L. (1994). Transcriptional

regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1. *J. Biol.*

Chem. **269**, 23757–23763.

Torsvik, N., Mortensen, S. and Nedreaas, K. (1995). Fishery biology (in Norwegian).

Landbruksforlaget, Aurskog.

Turek, F. W., Joshu, C., Kohsaka, A., Lin, E., Ivanova, G., McDearmon, E., Laposky,

A., Losee-Olson, S., Easton, A., Jensen, D. R., et al. (2005). Obesity and metabolic

syndrome in circadian Clock mutant nice. *Science (80-.).* **308**, 1043–1045.

Vacquie-Garcia, J., Lydersen, C., Biuw, M., Haug, T., Fedak, M. A. and Kovacs, K. M.

(2017). Hooded seal Cystophora cristata foraging areas in the Northeast Atlantic

Ocean—Investigated using three complementary methods. *PLoS One* **12**, e0187889.

van Moorsel, D., Hansen, J., Havekes, B., Scheer, F. A. J. L., Jörgensen, J. A., Hoeks, J.,

Schrauwen-Hinderling, V. B., Duez, H., Lefebvre, P., Schaper, N. C., et al. (2016).

Demonstration of a day-night rhythm in human skeletal muscle oxidative capacity. *Mol.*

Metab. **5**, 635–645.

Vázquez-Medina, J. P., Olguín-Monroy, N. O., Maldonado, P. D., Santamaría, A.,

Königsberg, M., Elsner, R., Hammill, M. O., Burns, J. M. and Zenteno-Savín, T.

(2011). Maturation increases superoxide radical production without increasing oxidative

damage in the skeletal muscle of hooded seals (Cystophora cristata). *Can. J. Zool.* **89**,

206–212.

Wang, G. L. and Semenza, G. L. (1993). General involvement of hypoxia-inducible factor 1

in transcriptional response to hypoxia. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 4304–4308.

Yamazaki, S. and Takahashi, J. S. (2005). Real-Time Luminescence Reporting of Circadian

Gene Expression in Mammals. *Methods Enzymol.* **393**, 288–301.

Zhang, G. and Tandon, A. (2012). Quantitative assessment on the cloning efficiencies of

lentiviral transfer vectors with a unique clone site. *Sci. Rep.* **2**,.

TABLES

Table 1. Primers sequences used for qPCR: efficiency calculated at 57° C.

Gene	Sequence	Efficiency
arntl F arntl R	GTGAAGTCTATGGAGTATGTG GTTCTTGTGGTAAATATGCC	98.34%
per2 F per2 R	CCATAGAAGAGATTGAGAGTG TTGGTCAGAGATGTACAGG	93.84%
ppib F ppib R	CATCTATGGTGAACGCTTC TAACGGTCGTGATAAAGAAC	89.32%
tbp F tbp R	GAAGAAAGTGAACATCATGG GTAAGGCATCATTGGACTG	92.98%

F=forward; R=reverse.   656

673 **Table 2.** List of chemicals used during the oxygen consumption measurements and relative

674 mitochondrial state.

Chemical	Cat. No. and Supplier	Final Concentration	State	Description
-	-	-	Routine	Basal respiration without addition of any substrate.
Digitonin	D5628, Sigma	3 µg/ml	-	Cell permeabilization
Pyruvate	P2256, Sigma	5 mM	Leak (PM – L)	Leak state at Complex I in the absence of ADP (inactive ATP synthase): non-phosphorylating state where protons leaking through complex I are not used for ATP production.
Malate	M1000, Sigma	2 mM		
ADP	117105, Calbiochem	2.5 mM	OXPHOS (PM – P)	Oxidative phosphorylation through complex I with pyruvate and malate as substrates
Glutamate	G1626, Sigma	10 mM	OXPHOS (CI – P)	Oxidative phosphorylation through complex I with pyruvate, malate and glutamate as substrates
Succinate	S2378, Sigma	10 mM	OXPHOS (CI + CII – P)	Oxidative phosphorylation through complex I and complex II with pyruvate, malate, glutamate, and succinate as substrates
Cytochrome c	C7752, Sigma	10 µM	-	Mitochondrial integrity test
Rotenone	R8875, Sigma	0.5 µM	OXPHOS (CII – P)	Oxidative phosphorylation through complex II after inhibition of complex I
Oligomycin	O4876, Sigma	10 nM	Leak (CII – L)	Leak state at complex II after inhibition of ATP synthase: non phosphorylating state
FCCP	C2920, Sigma	1 µM	Uncoupled state (CII – E)	Uncoupled state measuring maximal mitochondrial capacity

Table 3. P values for the fitted cosine model for each seal and month calculated through cosinor.detected function in the cosinor2 package in RStudio.

ID	Year	Month	P value	ID	Year	Month	P value
Mj98	2007	7	6.13E-05	F115	2008	3	2.85E-24
		8	1.27E-23			4	9.22E-114
		9	0.483512			5	3.34E-25
		10	3.31E-36			6	0.2485375
		11	2.83E-17	F119	2008	3	6.80E-13
		12	0.259391			4	5.57E-150
	2008	1	4.27E-09			5	5.59E-14
		2	0.0001599			6	0.0219391
		3	3.66E-16	F147	2008	3	1.04E-19
		4	1.06E-07			4	4.17E-39
		5	2.59E-07			5	9.68E-30
M171	2007	8	1.74E-06			6	6.50E-10
		9	2.88E-06	F154	2008	3	1.76E-22
		10	9.10E-93			4	4.20E-23
		11	1.51E-19			5	0.0001276
		12	1.29E-37			6	0.5792379
	2008	1	4.58E-26	F175	2008	3	2.77E-22
		2	1.31E-30			4	9.78E-36
		3	0.0621116			5	3.50E-14
		4	1.97E-28			6	0.904318
		5	3.78E-05	F180	2008	3	3.91E-20
F150	2008	3	1.30E-15			4	5.87E-14
		4	3.58E-08			5	0.0002414
		5	0.0215091			6	1.47E-08
		6	0.0729425	M225	2008	3	0.7984863
F110	2008	3	0.0003611			4	5.74E-07
		4	0.0746835			5	4.31E-21
		5	0.0725687			6	0.100202
		6	0.0531834	F93	2008	3	0.4504113
						4	2.06E-49
						5	8.77E-24
						6	0.5173444

685   *Non-significant values (p>0.05) are indicated by grey colour.*

 FIGURE LEGENDS

Figure 1. Hourly and daily variations in diving durations in an adult male hooded seal

(ID: M171) from the Northeast Atlantic population. Data were collected between August

2007 and April 2008 (data provided by M. Biuw: 10.21334/npolar.2017.881dbd20). Only

dives lasting between 2 min and the 95 percentiles of maximal duration are included. Each

point represents a diving event, the red line represents the cosine curve generated through

cosinor.lm in RStudio. Each panel corresponds to a different month and the respective p

values are indicated in the upper left. Blue and white areas represent day and night hours

calculated in relation to position (latitude and longitude) and time of year.

Figure 2. Hourly mean dive duration (minutes) throughout the year for 12 adult hooded

seals. Data from 2007 are presented in red (2 adult males); data from 2008 in black (9 adult

females and 1 adult male). Only dives lasting between 2 min and 95 percentiles of maximal

duration were included in the analysis. Data were binned by hour and the hourly mean dive

duration was calculated.

Figure 3. Clock genes expression profile in synchronised hooded seals fibroblasts and

tissues. A) mRNA expression of core clock genes *arntl* and *per2* in hooded seal fibroblasts

synchronised by temperature cycle (T°C), as represented in the upper panel. *Arntl* and *per2*

mRNAs show a 24-h period and are in antiphase. P values and phases were calculated through

JTK cycle analysis (*n*=4). **B)** mRNA expression of core clock genes *arntl* and *per2* in hooded

seal fibroblasts synchronised by temperature cycling and then left at constant temperature

(upper panel). *Arntl* and *per2* mRNAs maintain a 24-h period and their antiphasic relationship

also in constant conditions. P values and phases were calculated through JTK cycle analysis

(*n*=4). **C)** Photon multiplier tube recordings of hooded seal skin fibroblasts transfected with

arntl:luciferase and *per2*:luciferase. Cells were synchronised with dexamethasone. **D)** *Arntl*

mRNA expression in hooded seal kidney and liver tissue, sampled in mid-light phase (ZT6)

and mid-dark phase (ZT18). *$p<0.05$ (One-way ANOVA analysis; *n*=3 except liver ZT18

where *n*=2). Data are expressed as mean ± SEM.

Figure 4. Relationship between clock gene expression rhythms and mitochondrial

respiration in hooded seal fibroblast cultures. A) PCA analysis across 7 mitochondrial

states (*Leak* (PM-L, CII-L), *OXPHOS* (PM-P, CI-P, CI +CII-P, CII-P), *Uncoupled state* (CII-

E)). There is strong clustering at ZT5 and ZT17. Ellipses were drawn with a 0.80 confidence

interval (*n*=4 except at ZT13, 21 where *n*=3). **B)** Complex I coupling efficiency (calculated as

1-(PM-L/PM-P)) at all the ZTs where oxygen consumption measurements were made.

Coupling efficiency shows a time-dependent variation (one-way ANOVA: $p<0.01$; $n=4$ except

ZT13, 21 where $n=3$) **C)** Photon multiplier tube recordings of hooded seal skin fibroblasts

transfected with *arntl*:luciferase and synchronised by temperature cycling (36.5 – 39.5°C), as

indicated at the top . Recordings from the first 5 days were used to calculate expected peak

(red arrow) and through (green arrow). **D)** Complex I coupling efficiency calculated as in **B,**

at the expected peak and through indicated in **C**. ***$p<0.001$ (One-way ANOVA analysis,

$n=4$). Data are expressed as mean ± SEM.

Figure 1

A

Figure 2

Figure 3

Figure 4