1 **Title:**

2 **Single cell RNA-seq reveals protracted germ line X chromosome reactivation dynamics**

3 **directed by a PRC2 dependent mechanism**

13 **Authors:**

14 Yaqiong Liu [1], Xianzhong Lau [3], Munusamy Prabhakaran[3], Carlos M Abascal Sherwell

15 Sanchez[1], Daniel Snell[4] and Mahesh Sangrithi[1,2*]

Affiliations:

[1] King's College London, Centre for Gene Therapy and Regenerative Medicine, School of

Basic & Medical Biosciences, Faculty of Life Sciences and Medicine, London, UK

[2] King's College London, School of Life Course and Population Sciences, Faculty of Life

Sciences and Medicine, London, UK

[3] KK Women's and Children Hospital, Div. of Obstetrics and Gynaecology, Singapore

[4] Francis Crick Institute, London, UK

[*] Lead Contact: mahesh.sangrithi@kcl.ac.uk

Abstract

Initiating soon after PGC specification, female germ cells undergo reactivation of the silenced X chromosome during genome wide reprogramming. However, the kinetics and dynamics of XCR *in vivo* have remained poorly understood. To address this here we perform a global appraisal of XCR using high-dimensional techniques. Using F_1 *B6 v CAST* mouse embryos, we perform a detailed assessment, applying single-cell RNA-seq and chromatin profiling on germ cells purified from E10.5 to E16.5. While scRNA-seq profile showed that male and female germ cells are transcriptionally indistinct at E11.5, they are sexually dimorphic by E12.5, diverging further through development to E16.5. With allelic resolution, we show that the reactivating X chromosome is only partly active at E10.5, then reactivates gradually and reaches near parity in output to the constitutively active X chromosome at ~E16.5 when developing oogonia are meiosis prophase I. Crucially, we show that sexually dimorphic dosage compensation patterns observed in germ cells, occur in tandem with an increase in the allelic proportion from the reactivating X chromosome. While *Xist* is extinguished from E10.5, the epigenetic memory of earlier XCI in female cells persists much longer, likely from self-sustained PRC2 complex (*Ezh2 / Eed / Suz12*) function. The reactivating X chromosome is enriched in the epigenetic silencing mark H3K27me3 at E13.5, which is removed by E16.5 permitting gene expression. Our findings link XCR, along with functional regulation of PRC2 in promoting female meiosis.

Introduction

The accurate transmission of genomic information over generations involves the complex regulation of chromatin in germ cells. This process, germ cell genome wide reprogramming (GWR), initiates soon after primordial germ cell (PGC) specification very early in mammalian embryonic development. In the mouse, PGCs arise from the proximal epiblast at ~E6.25 - 6.75. By around E7.25, around 30-40 PGCs are present in the developing embryo, and continue to divide mitotically increasing in numbers. From E8.5 – 9.5 early PGCs migrate along the hindgut endoderm and reach the developing gonadal primordia by ~E10.5. At approximately E11.5 gonadal sex determination occurs, where expression of the Y-chromosome encoded *Sry* transcription factor in the gonadal pre-Sertoli cells initiates male gonadal fate and formation of a testis (Koopman et al., 1990). Conversely in females, the absence of the Y chromosome instigates the formation of an ovary. Thereafter gonadal PGCs assume sex-specific developmental fates.

GWR initiates soon after PGC specification at E6.75 and is sustained cell autonomously. It involves re-expression of the pluripotency network of genes (including *Pou5f1* (aka. *Oct4*), *Sox2*, *Nanog*, *Prdm14*), and repression of the somatic differentiation program, global DNA demethylation and removal of parental imprints (Tang et al., 2016). PGC further undergo extensive chromatin changes and remodelling of histone marks. Specifically female PGCs start to reverse X chromosome inactivation (XCI) established earlier at E5.5, i.e. X-chromosome reactivation (XCR).

The timing and dynamics of germ cell XCR are not fully understood. Early studies surmised that reactivation of X-linked genes in PGCs mostly occurred after they reached the genital ridge (~E11.5) (de Napoles et al., 2007; Johnston, 1981; Kratzer and Chapman, 1981; Monk and McLaren, 1981; Tam et al., 1994). Landmark experiments performed in single germ cells obtained from embryos, using reverse transcription followed by polymerase chain reactivation (RT-PCR), *Xist* RNA-FISH and immunofluorescence microscopy, had indicated that gene expression from the reactivating X-chromosome initiates in nascent PGCs (Sugimoto and Abe, 2007). While limited by a small number of X-linked genes and cells being assayed, they showed that the process of XCR in PGCs commences earlier, during the pre-gonadal phase of development, progresses gradually and was incomplete at E14.5. Similar results were reported in recently in PGC-like cells (PGLCS) derived *in-vitro* from mouse embryonic stem (ES) cells, highlighting that incorrect rapid XCR kinetics had resulted in limited meiotic potential (Severino et al., 2022).

XCI in the embryo is enacted by expression of the long non-coding RNA *Xist* (Kay et al., 1993;

Penny et al., 1996). *Xist* directly represses transcription by evicting the RNA polymerase

transcription machinery, while also initiating a cascade of epigenetic processes including the

loss of H3K4me3, recruitment of histone deacetylases to remove activating histone marks

(e.g. H3K27ac), and the *Polycomb repressor complex* PRC1 / 2 complexes which deposit

silencing histone marks H2A119ubq and H3K27me3 respectively (reviewed in Loda et al.

(2022)). Later in XCI, CpG-islands (CGI) of genes subject to XCI are methylated, and the

inactivated X (*Xi*) is compacted forming heterochromatin. Expression of *Xist* RNA is

downregulated in female PGCs from around E7.75 - E9.5, and is extinguished by ~E11.5 (de

Napoles et al., 2007; Sangrithi et al., 2017; Sugimoto and Abe, 2007). While histone marks

are globally reorganized during PGC development, the extent to which PRC 1 / 2 related

repressive marks deposited as a specific consequence of XCI in the epiblast are remodelled

remains poorly understood. For instance global changes to H3K27me3 levels are known to

occur as part of PGC development, and these can be challenging to disentangle from more

specific regulation of the mark relating to XCI (Lowe et al., 2022; Saitou et al., 2012).

It has been shown previously that sexually dimorphic dosage compensation states arise

during germ cell GWR, which is dependent on the number of X-chromosomes present

(Sangrithi et al., 2017). During GWR, females (XX) show X chromosome to autosome (X:A)

expression ratios greater than 1, while this was below 1 in males (XY). Strikingly the elevated

X:A ratios in oogonia persists from E11.5 until entry into meiosis. Hence elevated X:A ratios in

females may be inherently linked to the sexually dimorphic developmental fates of germ cells,

including timely meiotic-entry or / and the re-establishment of transcription on the reactivating

X (Sangrithi and Turner, 2018). Indeed dosage differences in X-linked genes could promote

germ cell sexual dimorphism and meiosis entry and / or progression itself in females, via the

involvement of X-linked genes in these processes and possibly XCR itself.

In this article we perform a global analysis of XCR dynamics of XCR using single-cell RNA

sequencing on developing mouse germ cells *in vivo* during the course of GWR. We leverage

on a genetically tractable murine model that enables clear delineation of the constitutively

active and reactivating X chromosomes for females. Using this model we describe a precise

allele-specific map of XCR, that charts the overall kinetics of this classic epigenetic process,

but also the dynamics of individual X-linked genes. This study demonstrates that XCR in

female germ cells initiates before E10.5 and accelerates while a sexually dimorphic

transcriptome is established. We demonstrate that transcription from the reactivating X

chromosome reaches close to parity with the constitutively active X chromosome only at

~E16.5 when developing oogonia are in zygonema / early pachynema. *Xist*-dependent

silencing that is present in early PGCs, then gives way to persisting repression of X-linked
genes due to histone 3 lysine-27 trimethylation (H3K27me3) later in gonadal germ cells where
direct *Xist* activity has ceased. H3K27me3 marks remain at the transcription-start sites (TSSs)
and gene-bodies of repressed X genes on the reactivating X chromosome at meiotic entry
(pre-Leptonema) in E13.5 females germ cells, with germ line genes expressed thereafter
showing a dynamic reduction in H3K27me3 at E16.5. We posit XCR as a mechanism of
meiotic upregulation in females.

Results

Transcriptional divergence of mouse germ cells occurs after E12.5

We developed a model to examine germline XCR, by combining a classical F_1 genetics and
single-cell RNA-sequencing (scRNA-seq). Leveraging on an interspecific cross of *Mus*
castaneus (*CAST*) males with reference strain C57B6J (*B6*) female mice, we performed
single-cell RNA-seq on F_1 female ($X_{CAST}X_{B6}$) and male ($X_{B6}Y$) embryonic germ cells from E10.5
to E16.5 stages. The *CAST* strain is highly polymorphic in relation to the reference *C57B6NJ*
strain, on average containing up to 1 single-nucleotide polymorphism (SNP) for every 150bp
in the genome (Keane et al., 2011). We further introduced an *Xist*-null (*Xist∆*) allele into our
experimental strategy to skew X-inactivation toward the *CAST* X chromosome (Marahrens et
al., 1998). By crossing *Xist$_{+/∆}$* C57B6J (*B6*) females (carrying *Oct4-EGFP* transgene) and
CAST males, we derived F_1 female ($X_{CAST}X_{B6-Xist∆}$) and male ($X_{B6}Y$) embryos, from which
embryonic gonads were obtained daily from E10.5 to E16.5 (Yeom et al., 1996; Yoshimizu et
al., 1999). Single *EGFP* positive germ cells were sorted using fluorescence-activated cell
sorting (FACS) and processed to generate high-quality scRNA-seq libraries using the SMART-
seq2 protocol (Picelli et al., 2014). To distinguish allele-specific expression in F_1 embryos,
reads were first aligned to a reference genome with SNPs N-masked to minimize bias arising
during alignment (Degner et al., 2009). Reads were then further assigned specifically to each
of the parental (*B6* and *CAST*) genomes respectively to obtain allele-specific counts (**Figure**
1A; please refer to **Methods** for details) (Krueger and Andrews, 2016). Where possible we
sought to include at least 20 germ cells from at least two individual embryos at each time point
and sex. Following stringent quality control checks, we retain 681 single-cell transcriptomes
for further analysis (**Supplementary Figure 1A**). These include 333 male and 348 female
cells (**Supplementary Figure 1B**).

First, we performed dimensionality reduction on the dataset. The scRNA-seq profiles reveals
that male and female germ cells appear transcriptionally indistinct prior to E12.5 (**Figure 1B**).
Following gonadal sex determination at E11.5, sexually dimorphic transcriptomes are
apparent by E12.5, which diverge further with continued development to E16.5. We sexed our

samples based on the expression of Y-chromosome encoded genes in the data which is
typically only detected males, and concordant with PCR based genotyping that was performed
at the time of sample collection (**Supplementary Figure 1C**). We utilized partition graph
abstraction (PAGA) analysis to visualize connectivity and relatedness between cell groups in
an unbiased manner. The PAGA graph emphasizes the divergence of the male and female
germ line after E11.5, further lending weight that our data reliably captures the establishment
of sexually dimorphic transcriptomes in germ cells (**Figure 1C**).

We next performed rigorous differential expression (DE) testing to examine transcriptional
changes arising during the establishment of germ cell transcriptional identity (**Figure 1D**).
Expression of the *Tet* enzymes (*Tet1* and *Tet2*), that regulate DNA demethylation was
observed in both male and female PGCs at E11.5, consistent with germ cells reaching a
methylation base state by E13.5 (Hackett et al., 2013; Popp et al., 2010; Seisenberger et al.,
2012). Male germ cells then begin to express the *Lefty* genes (*Lefty1* and *Lefty2*), *Pitx2* and
the transcription factor *Etv1* that is important in prospermatogonial development. In females,
we observe *Phlda2*, along with transcription factors and *Gata2.* Consistent with the pre-
leptonema stage, female germ cells subsequently begin to upregulate meiotic genes at E13.5,
including *Rec8, Stra8, Prdm9, Spo11* and genes with roles in meiosis I prophase thereafter
(**Figure 1E** and **Supplementary Figure 1D**). Together these culminate in the establishment
of pro-spermatogonia and oogonia respectively by E16.5, confirmed with gene ontology (GO)
enrichment scoring (**Figure 1F**). We next turned our focus toward understanding germ cell
XCR in females.

Global X chromosome reactivation dynamics
In our experimental model, the *Xist*Δ allele skews XCI invariably toward the *CAST* X
chromosome. The *CAST* X-chromosome thus is always inactivated in all cells in the epiblast,
and hence will also be the X chromosome that undergoes XCR in germ cells. Hence
expression specific to the *CAST* X-chromosome in our F_1 germ cells can be used to accurately
chart XCR. We computed allele-level counts and calculated an allelic deviation score ('*d*
score'), as the ratio of B6 reads to the total number of informative allelic reads (i.e. *d = CAST*
/ (B6 + CAST) - 0.5) for each gene (Xu et al., 2017) (see **Methods**). A negative *d-score*
corresponds to a bias toward expression from *B6* alleles, while positive values corresponding
to a bias toward *CAST* alleles, with *d-scores* ~0 being indicative of equal allelic expression.
Instances where the *d*-scores are less than -0.4 or greater than 0.4 represent monoallelic
expression bias. To test the assumptions of our experimental model, we first turned to
inspecting allelic balance in E11.5 female gonadal somatic cells obtained as a control.
Consistent with the *CAST* X chromosome being inactivated invariably in somatic cells, we

$$\text{d-score} = \frac{\text{CAST}}{\text{B6} + \text{CAST}} - 0.5$$

Figure 1. Transcriptome profiling of mouse germ cells by scRNA-seq

A. Illustration of experimental design.

B. *tSNE* analysis of a total of 681 germ cell scRNA-seq data. Cells are coloured according to sex and developmental stages. The number of cells at each stage and sex are summarised in the **Supplementary Figure 1**.

C. Partition-based graph abstraction (PAGA) plot representing a graph of inferred connectivity among all cell clusters. Line thickness indicates the strength of cluster connections.

D. Heatmap of top 10 DE genes at different embryonic stages in female and male germ cells.

E. A tracks-based bar-plot depicting the expression levels of key genes involved in PGCs, DNA methylation regulation, prospermatogonial / oogonial development, and meiosis in females and males.

F. Top GO:BP (Gene Ontology for Biological Process) terms with *p*-values for different embryonic stage germ cells.

substantiate a strong expression bias toward alleles on the *B6* X chromosome (median d_{ChrX}

= -0.42) in E11.5 female somatic cells. (**Supplementary Figure 2A**). This affirms that our

approach is efficient in identifying allelic expression deviation expected in XCI.

Next, we turned to studying allelic expression globally in female germ cells. We computed

global *d*-scores for each chromosome in all cells in our dataset. In more detail, we see that

male germ cells have *d-scores* of ~-0.5 throughout, consistent with these cells only having a

B6 X chromosome. In females, we observe that E10.5 female PGCs had a *d-score* of -0.2 for

expression from the X chromosome, with a minimum of ~-0.4 seen in some cells (**Figure 2A**).

d-scores then rise in female germ cells thereafter, remaining below 0 until E15.5, and only

reaching close to parity (i.e. ~0) at E16.5. Overall the allelic bias observed indicates that a

number of genes on the *CAST* X chromosome are clearly silent at E10.5 in female PGCs,

which is likely due to the effects of earlier XCI. No allelic expression bias is detected on any

of the autosomes at all stages we examined (**Supplementary Figure 2B**). To corroborate

these observations further, we chart the expression of *Xist* RNA in our dataset. In keeping with

the predictions of XCI, *Xist* is strongly expressed in E11.5 female <u>somatic</u> cells (**Figure 2B**).

In comparison *Xist* only expresses at very low levels in E10.5 female PGCs and is

downregulated after this stage, consistent with previous studies using RNA FISH (de Napoles

et al., 2007; Sangrithi et al., 2017; Sugimoto and Abe, 2007).

Sexually dimorphic dosage compensation states emerge alongside the reactivation kinetics

of the X chromosome

We next turned to examining dosage compensation patterns in germ cells. In order to chart

global transcriptional output from the X chromosome in relation to the autosomes, we

computed X chromosome to autosome expression ratios (X:A) in each cell and plotted these

(**Figure 2C**). We observed that male germ cells consistently have a X:A < 1 over the course

of their development. In contrast, female germ cells show increasing X:A ratios, which range

above 1 from E13.5 to E16.5. These observations are consistent with previous studies that

male germ cells have low X:A ratios (X:A < 1), while females showed an excess of X-gene

dosage (X:A > 1) as they enter meiosis (Sangrithi et al., 2017). While X:A ratios in males and

females are similar at E10.5, an increase is evident in females thereafter, occurring

contemporaneously with XCR as shown earlier (see **Figure 2A**). At E10.5, females have an

X:A ratio of 0.82 and males 0.77. X:A ratios show maximal differences between females and

males from E14.5 to E16.5 (median values of 1.05 versus 0.66 respectively at E16.5).

Expression dynamics at individual X gene loci reveal XCR mainly occurs after E12.5

Following stringent filtering we chart the *d*-score for 281 representative X-linked genes subject

253
254

Figure 2. Sexually dimorphic kinetics of X chromosome reactivation and X dosage

compensation

A. Boxplot of mean *d*-scores showing allelic balance of chromosome X and 9 during germ cell

development in female and male. Each point represents a single cell.

B. The dot plot representing *Xist* expression levels in female somatic cells (highted in dashed

red square) and in female germ cells at different stages.

C. Violin plot representing the ratio of mean read counts of genes on chromosome X (dark

red) and 9 (blue) to the mean read counts of all genes across all autosomes for all germ cells

at different embryonic stages between female and male.

to XCR, expressing biallelically during female germ cell development (see **Methods**). These
are depicted as heatmaps, shown in relation to development age and in context of their
genomic location (**Figure 3Ai and ii**). Similarly we also detail their overall expression
alongside (**Figure 3Bi and ii**). Together these data depict both allelic balance and expression
dynamics for these genes expressing in the female germline from embryonic ages E10.5 to
E16.5. From these we surmise that around ~32% of assayed genes (91/281) express from
both alleles (i.e. d > -0.4) at E10.5 (**Figure 3C**). This increases to ~90% of genes expressing
biallelically at E16.5 (250/281). Crucially we discover that a significant portion of XCR (~40%)
occurs from E12.5 to E16.5, which is much later than previously appreciated (**Figure 3C**). A
number of genes only express later during this time course, with many notably peaking in
expression at E16.5.
280
Further analysis reveals gene expression occurring over the entire length of the X
chromosome during germline XCR, even at E10.5. As we observe very low levels of *Xist*
expression in E10.5 female PGCs, we examined if XCR could be linked to *Xist* entry sites, by
surveying *Xist* RAP data (**Supplementary Figure 3**) (Engreitz et al., 2013). Further to this
appraisal we find that neither proximity to *Xist* entry sites nor to the X-inactivation centre (XIC)
itself appear to have a significant impact on allelic expression from the reactivating X
chromosome in the female germline. This analysis indicates that XCR occurs in a locus-
specific manner.
289
In summary, three important conclusions emerge in this regard. First, we show that the
increase in X:A ratios seen during female germ cell development occurs contemporaneously
with increasing *d*-scores (i.e. to 0), demonstrating that the increase in expression from the X
chromosome is specifically due to biallelic gene expression. Second, allelic balance of the
reactivating *CAST* X chromosome and the constitutively active *B6* X reach close to parity at
E16.5 when developing oogonia are already in prophase I of meiosis, suggesting that XCR
may have an impact in meiosis. Thirdly the protracted reactivation dynamics of the silenced
CAST X chromosome, beyond *Xist* activity, specifically points toward epigenetic memory from
XCI persisting on this chromosome. In all, these results show that germline XCR begins in
early PGCs, then proceeds gradually, with most of germline XCR occurring after E11.5 and
during female meiosis. While *Xist* RNA expression is notably downregulated after E10.5 in
female PGCs, we hypothesise that silencing histone marks deposited as an effect of *Xist*
related PRC2 / 1 activity until this developmental age, persists on PGC chromatin (e.g.
H3K27me3 and / or H2A119Ubq deposited consequent to XCI in early PGCs).
304
305

Figure 3. The change in *d*-scores and expression levels of X chromosome linked genes
during female germline development from E10.5 to E16.5.
A. The heatmap plot of *d*-score showing the allelic balance of 281 X-linked genes; genes are
ordered based on (i) reactivation timing according to gestational age and (ii) the genomic
location. Five classical XCI 'escape' genes (*Ddx3x, Eif2s3x, Kdm5c, Kdm6a* and *Zfx*) are
highlighted in a red square.
B. The heatmap depicting the expression of 281 X-linked genes; genes are ordered based on
(i) reactivation timing according to gestational age and (ii) the genomic location.
C. The percentage of X-linked genes expressing biallelically (*d* > -0.4) at each stage.
316

H3K27me3 is enriched on the reactivating X chromosome

To investigate this possibility in greater detail, we next turn to examine how PRC2 dependent

H3K27me3 is allelically regulated during germline development. We first optimized low-input

c̲leavage u̲nder t̲arget and r̲elease u̲sing n̲uclease (CUT&RUN) for H3K27me3 and after

validating antibody specificity and reproducibility (**Supplementary Figure 4A and B**), perform

low-input CUT&RUN for H3K27me3 on FACS purified germ cell fractions obtained from E13.5

and E16.5 F_1 mouse ovaries (Skene and Henikoff, 2017).

H3K27me3 enrichment is seen broadly across the genome, and detected over promoters and

gene bodies in female germ cells (**Figure 4A, Supplementary Figure 4C**). Specifically we

see a strong enrichment signal over the *Hoxa* gene cluster, which is typically observed in germ

cells associated with repression of somatic program (Zheng et al., 2016) (**Supplementary**

Figure 4D). From this analysis, we notice a higher enrichment signal for H3K27me3 on the X

chromosomes versus the autosomes, more specifically on the *CAST* X chromosome (which

is subject to subject to XCI) compared to the *B6* X, clearly more evident at E13.5 versus E16.5

(**Figure 4A**). To confirm differences in H3K27me3 enrichment between the *CAST* and *B6*

hemigenomes at each age, we compute a *log2*-ratio of *CAST* over *B6* specific signals in E13.5

and E16.5 germ cells (**Supplementary Figure 4E**). In this manner, we demonstrate a greater

enrichment of this mark on the *CAST* X chromosome compared to the *B6* X at E13.5*,* which

we interpret as being due to epigenetic memory of earlier XCI. Crucially, we discover a

reduction in the H3K27me3 overall signal in E16.5 oogonia compared to E13.5 oogonia

(**Figure 4A**).

Since the reduction in H3K27me3 levels occurs over the same time frame as the increase in

X chromosome *d*-score and X:A ratios, we hypothesise that gene activation on the X occurs

following removal of this silencing mark. We took two steps to test this directly: 1) we first

performed DE analysis to identify genes activated between E13.5 to E16.5 in our RNA-seq

dataset (**TableS1**; see **Methods**), and 2) we computed the H3K27me3 signal from CUT&RUN

at these DE gene loci in oogonia at E13.5 versus E16.5. In this manner first we identify 76 DE

genes on the X chromosome, that increase in expression from E13.5 to E16.5; and 70 DE

genes on chromosome 9 were used as autosomal controls. Now returning to CUT&RUN, we

examine H3K27me3 levels specifically at these DE genes. Importantly a reduction of

H3K27me3 levels is evident at these genes, with the signal change being most pronounced

at gene promoters, transcription start sites (TSS) and gene bodies (**Figure 4B**). A number of

observations come to light. We detail this reduction at specific DE X-linked gene loci - including

Nxf2, *Ccnb3*, *Tktl1*, *Taf7l* and *Taf9b*, which have recognized functions in the germ cells and

associated with fertility phenotypes (Chotiner et al., 2022; Gura et al., 2020; Pan et al., 2009;
Rolland et al., 2011; Wang et al., 2001) (**Figure 4C**).

356
357
358

Figure 4. H3K27me3 enrichment remains on the reactivating X chromosome during oogonia development.

A. H3K27me3 enrichment signals for all genes on chromosome X and 9 (total), and shown for alleles on the *B6* and *CAST* hemigenomes at E13.5 and E16.5 plotted as heatmaps, with associated summary profiles plotted above. Regions 2kb upstream of transcription start sites (TSSs), gene body and transcription end sites (TESs) are shown. Chromosome X signal is shown coloured orange, and chromosome 9 in blue.

B. Genome tracks for H3K27me3 enrichment at E13.5 and E16.5 oogonia for representative genes.

C. H3K27me3 enrichment signal at DE genes between E13.5 and E16.5.

Overall, we evidence a greater reduction in signal between E13.5 and E16.5 occurring on the
X chromosome versus chromosome 9 (**Supplementary Figure 4F**). Looking specifically at
the *B6* and *CAST* hemigenomes, we demonstrate a greater reduction in H3K27me3 levels on
the reactivating *CAST* X chromosome versus the *B6* X chromosome (or any autosome) at
these gene loci (**Figure 4B & Supplementary Figure 4E,F**). These results confirm that
expression of these DE genes is specifically associated with the loss of H3K27me3 at
promoters, TSS and gene bodies.
377
XCR is dependent on Ezh2 activity
Previous studies have shown a vital role for H3K27me3 in regulating sexually dimorphic
germline development, regulated by component factors of the PRC2 complex, *Ezh2* and *Eed*
(Huang et al., 2021; Lowe et al., 2022). In order to further assess the role of H3K27me3 in this
context, we sought to examine data from *Ezh2*-mutant germ cells (**TableS2)** (Huang et al.,
2021). From our analysis we notice that a sizeable proportion of genes that are de-repressed
in E13.5 *Ezh2*-null oogonia reside on the X chromosome (Huang et al., 2021). Distinctly we
see that ~12.5% of all *Ezh2*-dependent DE genes are X-linked (121 genes with log2folchange
> 1; see **Methods**), indicating that *Ezh2* activity significantly regulates gene expression on the
X chromosome (**Figure 5A, Supplementary Figure 5B and D**). Again ~13.2% of DE genes
derepressed in E13.5 *Ezh2KO* prospermatogonia were X-linked (48 genes; **Supplementary**
Figure 5C and D). These data indicate that Ezh2 dependent H3K27me3 levels on germ cell
chromatin regulates the repression of germline genes, orchestrated in a sexually dimorphic
manner. We therefore hypothesise that at least some X chromosome genes subject to XCR
later (i.e. genes seen to express after E13.5 in our RNA-seq dataset) are directly regulated by
H3K27me3 levels. To verify, we first examined the expression of the 121 de-repressed X-
linked genes identified in *Ezh2*-null female oogonia, in our scRNA-seq dataset. This analysis
indeed confirms these genes are mostly expressed physiologically after E13.5, i.e. from E14.5
to E16.5 (**Figure 5B**). Thus, taken in the context of XCR, these data definitively show that
Ezh2-dependent H3K27me3 deposition is a functional requirement for the repression of these
X-linked genes in E13.5 oogonia, and this silencing mark is then likely reversed for their timely
expression observed thereafter. To test this we next turned to directly examine H3K27me3
levels at *Ezh2* regulated genes in oogonia. In keeping with XCR, we observe a greater
reduction in H3K27me3 signal at Ezh2-dependent genes on the *CAST* X chromosome
identified in our earlier analysis (**Figure 5C)**. In all, our data provide strong evidence that XCR
involves the removal of H3K27me3 at specific X-linked gene promoters, TSS and gene bodies
by meiosis prophase 1 at E16.5.
405
406

Figure 5. Role of *Ezh2* in X reactivation during oogonia development.

(A) The box plot representing the percentages of *Ezh2*-dependent de-repressed genes on each chromosome in the E13.5 oogonia. Chromosome X is shown coloured orange, and gray dotted line indicates the number of expected genes per chromosome if distributed randomly (~6-8 genes; cf. ~5-6%).

(B) Heatmap showing the expression levels of 121 *Ezh2*-related de-repressed X-linked genes in our scRNA-seq dataset.

(C) H3K27me3 enrichment signal for *Ezh2*-dependent genes in our CUT&RUN data, comparing the signal in B6, CAST, and both alleles between E13.5 and E16.5.

Discussion

In this article we present a highly resolved analysis of X chromosome reactivation kinetics and dynamics using single-cell SMART-Seq2 RNA seq and low-input CUT&RUN at matched time-points. Here our data provides important new insights into gene expression regulation, X chromosome dosage compensation and chromatin dynamics during XCR. While conventional thinking has been that XCR begins early and was mostly complete by the early gonadal phase of germ cells, from our extended work we show that XCR is more protracted than previously appreciated, extends into meiosis, and evidence implicating PRC2 complex (*Ezh2 / Eed / Suz12*) function regulating gene repression on the reactivating X chromosome.

The protracted kinetics of re-establishing transcription on the reactivating X chromosome is striking. We delineate how XCR progresses, and matches with non-canonical X:A dosage compensation patterns observed in the germline. During GWR in female germ cells, X:A ratios exceed 1 from gestational age E13.5. From allele-specific analysis we show that higher X:A ratios in female germ cells are coincident with biallelic gene expression, indicating that X chromosome output arises from the reactivating (*CAST*) X chromosome combined with expression from the constitutively active (*B6*) X chromosome. Our data further highlights a time-window at which near complete reactivation of the X occurs at E16.5 when female germ cells are known to be in late zygotene and early pachytene stage of meiosis.

One of the important findings in this study is the identification of a functional role for H3K27me3 levels in regulating gene expression from the reactivating X chromosome in female germ cells during this developmental time-window. H3K27me3 enrichment on the reactivating X chromosome and its extended presence beyond the expression of *Xist* RNA itself is interesting - notably while *Xist* is downregulated in female germ cells from E10.5, H3K27me3 deposited as a consequence of XCI earlier during PGC development persists, implicating epigenetic memory. This is most likely due to the "self-contained" activity of the PRC2 complex (*Ezh2 / Eed / Suz12*), extending beyond direct *Xist* activity (Yu et al., 2019). There is a precedence for the self-sustaining silencing activity of the PRC complexes on the inactive X (X_i), which has also been observed in trophoblast stem cells (Mak et al., 2002; Masui et al., 2023). Here we conclude that XCR does not complete with *Xist* downregulation, and gene silencing on the reactivating X chromosome persists in the absence of *Xist*, due the presence of this repressive histone mark. Self-contained PRC2 activity ensures stable maintenance of H3K27me3 mediated gene silencing, mitigating against cell-cycle dependent dilution of this mark. Female germ cells cycle mitotically until ~E13.5 at which point they commit to meiosis. In the absence of cell-cycle dependent dilutional effects beyond this stage in the female germ line, the loss of

464   H3K27me3 signal observed thus points to the existence of active processes to reverse
465   H3K27me3 silencing at these loci. In this regard, future studies could be aimed at identifying
466   and evaluating such mechanisms involved in reversing H3K27me3 mediated gene silencing
467   in the germline.
468
469   The presence of H3K27me3 on the reactivating X chromosome crucially also accounts for the
470   dynamic X:A dosage compensation pattern seen in XX germ cells. The relative overexpression
471   of X-genes seen from E13.5 in females, observed by X:A ratios ranging above 1, could be
472   attenuated by this silencing mark (cf. related observations made in the male by Lowe et al.
473   (2022)).
474
It is unexpected that the reactivating X chromosome remains enriched for H3K27me3 over the
constitutively active X in female germ cells in the context of meiosis. PRC2 is known to act as
a factor to maintain repression of distinct gene types: e.g. developmentally regulated genes,
protein-coding genes that are repressed (including germ line genes) and imprinted genes.
Thus, the control of PRC2 activity may be developmentally regulated to promote meiosis via
the timely expression of germline genes in a sex-specific manner. Three further points also
come up in this regard. First the asymmetric distribution of H3K27me3 on the X chromosome
at the point of meiotic entry at E13.5 merits further investigation (i.e. in chromosome pairing,
synapsis and segregation). Second the regulation of heterochromatin more generally during
female meiosis would also be of interest as it highlights possible roles for other silencing marks
on the inactive X (e.g. H2A119Ubq and H3K9me3). And third it of importance to understand
the mechanisms involved in the reversal of silencing on the X, to achieve expression.
Furthermore, from an evolutionary perspective it would be of interest to compare XCR in other
mammals such as humans to understand conservation and / or divergence in these
fundamental epigenetic processes.
490
Despite the well-known complex dynamics of XCR, notably important differences are apparent
with our *in vivo* appraisal of XCR and previous studies using *in vitro* derived PGC models of
XCR, where incomplete XCI had occurred in PGCLCs further limiting meiotic potential
(Severino et al., 2022). This was confirmed in a recent manuscript using a random XCI model
in early *in vivo* PGCs (Roidor *et al.* 2023), where PGCs had mostly maintained an inactive X
chromosome at E9.5, with a limited number of genes undergoing XCR thereafter. From the
experimental strategy employed in this study using F_1 female ($X_{CAST}X_{B6-Xist\Delta}$) embryos,
gestational ages covered, we are able to demonstrate many more genes subject to XCR. This
supports the finding that a number of genes are orchestrated to undergo XCR with important
functional roles therein in development of the female germ line.

In conclusion we present new evidence that challenge existing paradigms in X chromosome
biology, showing that XCR extends much later than previously appreciated in female germ
cells. Our work further connects PRC2 activity beyond a sex specific role in the germline, more
directly in regulating X chromosome activity in promoting female meiosis (Huang et al., 2021;
Lowe et al., 2022).
506
507
508

Supplementary Figure legend:

Supplementary Figure 1. Quality control of scRNA-seq data

(A)   Dot plot of total counts, number of genes expressed, and the percentage of mitochondrial reads in the scRNA-seq data in this study. Each dot represents a single cell.

(B)   Summary of cells represented in scRNA-seq dataset.

(C)   The expression level of Y-chromosome encoded genes across different embryonic stages between male and female.

(D)   The expression level of key genes across cells, visualised on the *t-SNE*.

Supplementary Figure 2. The allelic expression of chromosomes in female cells (relates to Figure 2)

(A)   The boxplot of *d*-scores of chromosome X and 9 in E11.5 female germ cells and somatic cell.

(B)   The boxplot of *d*-scores of all autosomes and chromosome X in oogonia from E10.5 to E16.5. Each dot represents a single cell sample.

Supplementary Figure 3 (relates to Figure 3)

The heatmap plot of *d*-scores for of 281 X-linked genes; genes are depicted based on the genomic location. The *Xist* RAP-seq signal is represented in blue in the figure, and putative *Xist* entry sites as red dots. *Xist* / XIC location is marked as shown (high signal for *Xist* over the XIC is not shown).

Supplementary Figure 4. CUT&RUN results (relates to Figure 4)

(A)   The H3K27me3 antibody verification: heatmap showing the H3K27me3 antibody specificity validated using SNAP-CUTANA K-MetStat Panel. Representative immunofluorescence image of α-H3K27me3 (magenta), α-Ddx4 (cyan) and endogenous GFP in an E13.5 ovary. White arrows indicate Barr bodies in somatic cells.

(B)   The heatmap representing the spearman correlation of reads counts generated from CUT&RUN samples.

(C)   The genome tracks showing H3K27me3 enrichment signal over *Hoxa* gene cluster in E13.5 and E16.5 oogonia.

(D)   The distribution of H3K27me3 peaks across functional annotations, including promoter, exons, introns, downstream, and distal intergenic regions in E13.5 and E16.5 germ cells.

(E)   The box plots showing the *log2*-ratio of H3K27me3 enrichment signal in *CAST* to *B6*
alleles in all the chromosomes. Chromosome X is highlighted in orange colour and 9 in blue
(relates to **Figure 4A**).

(F)   The *log2*-ratio of H3K27me3 enrichment depicting change in signal in E13.5 over
E16.5 germ cells at DE genes. Plots show higher enrichment at E13.5, especially at CAST
specific alleles (relates to **Figure 4C**).

Supplementary Figure 5. Analysis of *Ezh2*-null germ cells scRNA-seq (relates to Figure
5)

(A)   Principal component analysis (PCA) of RNA-seq data from *Ezh2*-null (conditional
knockout) and control male and female E13.5 gem cells.

(B)   Volcano plot showing the differentially expressed genes (adjusted p-value <0.05,
log2FC>1 or log2FC<-1, orange dots) between *Ezh2*-null and control female germ cells. The
X-linked DEGs were highlighted in dark blue (n=121).

(C)   Volcano plot showing the differentially expressed genes (adjusted P value <0.05,
log2FC>1 or log2FC<-1, orange dots) between *Ezh2*-null and control male germ cells. The X-
linked DEGs were highlighted in dark blue (n=48).

(D)   Summary of DE genes between *Ezh2*-null and Control mice.

Acknowledgements:
We thank Dr. James Turner (Francis Crick Institute) for enabling pilot experiments / sharing
relevant mouse lines, helpful discussions, critical reading of the manuscript. We thank
members of the Sangrithi lab for discussions and helpful feedback. Access to high-
performance computing resource for the Sangrithi lab is obtained from King's College London,
King's Computational Research, Engineering and Technology Environment (CREATE). MS is
supported by the Wellcome Trust (222052/Z/20/Z).

Contributions:
YL designed and performed all the experiments and maintained the mouse lines. YL and XZ
collected the samples. Bioinformatic analysis were performed by PM and supervised by MS.
CASS performed analysis of *Ezh2*-null data and immunofluorescence microscopy. DS
performed FACS. MS designed and supervised all experiments, wrote the manuscript,
maintained all university compliances, and attained funding for the experiments.

Data availability:

Single-cell RNA-seq data and CUT&RUN data will be publicly available as of the date of

publication in a peer-reviewed journal.

METHOD DETAILS:

Biological Samples and the Ethical Use of Animals

Use of experimental mice was undertaken in strict accordance with the Animals (Scientific

Procedures) Act 1986 subject to local ethical review and carried out under UK Home Office

license. Mice were maintained on a 12:12 hour light : dark cycle at 22 ± 2 °C, with food and

water available *ad libitum*. *Oct4-EGFP* mice are maintained on a C57B6 background and used

to mark / isolate fluorescently marked germ cells (Yeom et al., 1996; Yoshimizu et al., 1999).

*Xist*Δ mice are also maintained on a C57B6 background (Marahrens et al., 1997) and

maintained with the *Oct4-EGFP* line. *Mus castaneus* (*CAST / EiJ*; RRID*:IMSR_JAX:000928*)

were purchased from The Jackson Laboratory (JAX).

Experimental F_1 embryos were generated by crossing $Xist^{+/\Delta}$ (B6) *dams* also carrying *Oct4-*

EGFP transgene, and *CAST sires*, producing F_1 female ($XX^{Xist\Delta}$) and males (XY) that were

analyzed. Typically timed natural matings were set up at around 17:00 hrs, by placing female

mice into with the male. Females were checked for the presence of a vaginal plug indicating

a mating has occurred, with noon on the day the vaginal plug taken as E0.5 and was separated

from the male.

Genotyping

PCR genotyping was performed on extracted DNA. The primer information is presented in

TableS3.

Sample collection and processing

F_1 embryos were harvested daily at E10.5 to E16.5 from the date of the vaginal plug, with

further morphological verification at dissection. The embryonic urogenital complexes were

carefully removed under the stereomicroscopy, and washed in cold DPBS twice. For

transcriptomic and CUT&RUN, the embryonic gonads were separated from adjacent

mesonephroi and placed into 1.5 mL Eppendorf tube with 300uL DPBS.

FACS Isolation of germ cells

To obtain a single-cell suspension for FACS, isolated gonads were digested in the HBSS buffer

containing Collagenase Type 1 (200U/mL), Dispase II (2.4 U/mL), and $CaCl_2$ (1mM) at 37°C.

Tubes were shaken every 2-3 mins until the tissue fully dispersed. Enzymatic digestion was

then neutralisation with DMEM, containing 10% foetal bovine serum, followed by
centrifugation at 1000g, at 4°C for 4 minutes. The cell pellet was then resuspended in cold
FACS buffer (1% BSA in HBSS, with 25 mM HEPES, 1mM EDTA, and PI buffer) and filtered
through a 40μM strainer. EGFP fluorescence was used to isolate the fluorescently marked
germ cells through FACS. Live cells germ cells (PI negative, GFP positive) and somatic cells
(PI negative, GFP negative) were sorted by florescence-activated cell sorting (FACS) using
ARIA III flow cytometer (BD Bioscience).

For single cell RNA-seq, individual cells were sorted into 96-wells plates containing lysis buffer.
Plates were sealed and centrifuged immediately (700g, 10 seconds) following the sort and
stored at -80°C until needed. For CUT&RUN experiments, bulk sorting was performed to
collect purified germ cell and / or somatic cell fractions separately into 300μL of collection
buffer (5% BSA-HBSS) and used fresh.

SMART-Seq2 library preparation and sequencing
SMART-seq2 method was applied to generate the full-length cDNAs. Briefly, a single cell
(~0.3μL) was sorted into the 96 -wells plates containing 4 μL of lysis buffer (a ribonuclease
(RNase) inhibitor, dNTPs, and oligo-dT oligonucleotides). Cell lysis was performed by
incubating samples at 72 °C for 3 mins and put back on ice immediately. Then, reverse
transcription (RT) reaction was carried out by adding 0.5μL Maxima H-reverse transcriptase
(200 U / μL), 0.25μL RNA inhibitor (40 U / μl), 2μL Maxima-H RT buffer (200U / μL), 2μL
Betaine (5M), 0.06μL $MgCl_2$ (1M), 0.1μL TSO (100 μM), and nuclease-free water to make a
total of 10μl reaction. The reaction program of Maxima H reverse transcriptase was 90 mins
at 50°C, followed by 5 mins enzyme activation at 85°C; the reaction was then hold at 4°C. The
pre-amplification was then performed by adding 12.5μL KAPA HiFi HotStart, and 2.5μL
nuclease-free water. 20 PCR cycles were used, and the PCR cycle was set as follow: 98°C
for 3 mins, 20 cycles of 20s at 98°C, 15s at 67°C, and 6 mins at 72°C. The final elongation
was performed at 5 mins at 72°C. After purification with AMPure XP beads (Beckman Coulter)
(in a ratio of 0.7:1), the PCR product was quality checked (QC). Tagmentation of 0.3ng cDNA
was carried out by using the Illumina Nextera XT DNA sample preparation kit in a reaction
volume of 4 μL (2μL of tagmentation DNA buffer (TD, 2 x); 1μL of Amplicon tagmentation mix,
and 1μl of diluted cDNA), followed by stripping Tn5 transposase off with 1μL of NT buffer
added. Finally, amplification of adapter-ligated fragments was performed by adding 3μL
Nextera PCR master mix, and 2μL of Index primer combination. After purification with AMPure
XP beads (in the ratio of 0.6:1), the final cDNA library was quantified and QC-ed.

The concentration of PCR product was measured by using Qubit dsDNA assay kit (Invitrogen),
and size distribution was checked using an Agilent high-sensitivity chip (D1000) on 4200
TapeStation System. Overall, 741 scRNA-seq libraries were prepared and 2x150 bp paired-
end (PE) reads was performed on an Illumina Novaseq 6000 instrument. Where possible we
aimed to obtain and sequence a minimum of 24 cells from each embryo.
659
RNA-seq data analysis and processing
All processing and analyses were performed on the high-performance computing (HPC)
platform King's Computational Research, Engineering and Technology Environment
(CREATE). Fastq files containing reads from single-cell RNA seq libraries were subject to
quality control (fastqc), and trimmed for adapter content using Trimgalore. Using the SNPsplit
tool, SNPs positions for *C57B6NJ* and *CAST / EiJ* in the GRCm38.p6 reference mouse
genome (*Ensembl*), were first N-masked to eliminate mapping-bias between alleles (Krueger
and Andrews, 2016). An N-masked genome index was then generated using *STAR* (Dobin et
al., 2013; van de Geijn et al., 2015). Trimmed fastq files were subsequently aligned to the N-
masked reference using the *STAR* aligner.
670
671   Total gene-level expression counts were then generated using *featureCounts* (*Subread*) and
672   used in downstream analyses (Liao et al., 2014). Similarly allele-specific counts for
673   alignments specific to *B6* and *CAST* were generated using split `bam` files generated by
674   SNPsplit.
675
Total counts were imported into *SCANPY* for further specific analyses including normalization,
dimensionality reduction and statistical approaches (Wolf et al., 2018). Cells kept fulfilled the
following criteria - min_genes > 3500, max_genes < 10000, min_counts > 500000,
mitochondrial counts < 5%. Genes kept for analysis fulfilled the following criteria - min_cells =
5. Following this, 681 (out of 738) high-quality single-cells transcriptomes were retained for
further processing with each time point represented by a minimum of 20 cells derived from
each embryo (please see **Supplementary Table 1B**). *In toto* 681 cells and 21174 expressed
genes were analysed.
684
Allele-specific analyses.
Custom scripts in R were used to derive allelic-ratios and *d-scores* (*d*) from count matrices
output from *featureCounts*. *CAST* allelic ratios were calculated as *CAST / (B6 + CAST)*, and
d score = *CAST / (B6 + CAST) - 0.5)* was calculated for each gene and / or each cell. Reads
to the *CAST* X chromosome in male cells likely arising from technical artifacts of singular mis-
annotated SNPs, were not considered. For XCR analysis, genes with counts detected in at

least 2 or more cells, and 5 fragment counts or more at each gene locus were kept for XCR

analyses.

Cleavage Under Targets and Release Using Nuclease (CUT&RUN)

For each CUT&RUN reaction, typically ~5000 cells were used, pooled from different embryos

of identical genotype in the same litter. Here we applied a CUT&RUN method modified from

previous protocol (Derek Janssens, Steven Henikoff 2019. CUT&RUN: Targeted in situ

genome-wide profiling with high efficiency for low cell numbers. protocols.io

https://dx.doi.org/10.17504/protocols.io.zcpf2vn). Sorted cells were washed with Wash Buffer

(20 mM HEPES, pH7.5, 150 mM NaCl, 0.5 mM spermidine and a Roche complete tablet per

50 ml), and then bound to activated Concanavalin A (ConA) -coated magnetic beads. After 10

mins of binding, cells were permeabilized and incubated in 50µL Antibody Buffer (Wash buffer

containing 0.01% Digitonin and 0.5µL of specific primary antibody) at 4°C overnight on a

nutator. For the negative control, anti-Rabbit IgG was added instead of primary antibody. Next

day, after 2 washes with Digitonin buffer (Wash Buffer containing 0.01% Digitonin), beads

were resuspended in pAG / MNase buffer (50µL Digitonin buffer with 2.5µL of pAG / MNase

reagent (EpiCypher) and incubated at room temperature (10 mins) on a nutator for pAG-Mase

binding. After washing away pAG / MNase with Digitonin buffer, the chromatin digestion was

performed by adding 1µl of 100mM $CaCl_2$ to each reaction and incubating at 4 °C for 2 hours

on a nutator. Digestion was then stopped with the addition of 33µL Stop Buffer (340 mM NaCl,

20mM EDTA, 4mM EGTA, 50µg / mL RNase A, and 50µg / mL glycogen) and the cleaved

chromatin was released to supernatant by incubating at 37°C for 10 mins. DNA was then

purified using a DNA clean-up column (Monarch® PCR & DNA) following the manufacturer's

instructions, and used for library preparation (NEBNext® Ultra™ II DNA Library Prep Kit for

Illumina®). Specifically, the concentration of adaptor (reduced to 3µM) and the cycle number

of PCR amplification (15 cycles) were optimised in our experiment considering of the low input

of DNA. Finally, the clean-up of PCR product was carried out by using AMPure X beads

(Beckman Coulter) (in a ratio of 0.9:1). After measuring the quantity of library and checking

the size distribution, libraries were pooled and sequenced to produce 2 X 50bp PE sequencing

on Illumina NextSeq 2000 instrument.

We determined the specificity for the antibodies used, using the SNAP-CUTANA K-MetStat

Panel (EpiCypher, US). 2µL of 1:20 diluted solution was applied to ConA-immobilized cells

prior to the addition of antibody. As per the manufacturer's instructions, reads were matched

to the unique DNA barcodes in the panel and normalized to either on-target or total counts.

Antibody enrichment of the expected spike-in nucleosome target confirmed the integrity of the

H3K27me3 antibody used in our experiment, while the IgG control antibody showed no

preferential enrichment for any nucleosome as expected (**supplementary Figure 4A**). In

addition, we validated this antibody by immunostaining of H3K27me3 on E13.5 embryo

gonads (**supplementary Figure 4A**).

CUT&RUN data processing

In short, *fastq* files were pre-processed for quality and trimmed (min length >35). Reads were

aligned to the N-masked reference genome using *STAR* with spliced-alignments turned off.

Subsequent steps in the analyses were performed using the *deepTools* package (v3.5.0)

(Ramirez et al., 2016). Pearson correlations were computed and plotted to visualize cross-

replicate validity (*multiBamSummary*), before proceeding to merge bam files at each time-

point. Reads across the genome were counted in 10kb-size bins for each library, and effective

library sizes were calculated using *csaw*, with and the TMM method applied to compute

normalization factors (Lun and Smyth, 2014; Robinson et al., 2010). *Bigwig* files were

generated from the *bam* files using *bamCoverage* with settings "-binSize 100 --smoothLength

1000". Signals were further computed / visualized using deepTools' functionality (e.g.

computeMatrix, plotHeatmap), and to plot enrichment profiles relating to each timepoint.

Quantification and statistical analysis

Statistical analyses were performed in R statistical computing playform (https://www.r-

project.org/) or Python. Statistical tests and the cut-off values used for each analysis are

described in each methods subsection. $p < 0.05$ was considered significant for all tests. For the

biological replicated, in the scRNA-seq experiment, at least 2 embryos and 24 scRNA-seq

libraries / embryo were generated at each time-point. In the CUT&RUN experiment, 2

biological repeats were carried out at each time-point.

References:

Chotiner, J.Y., Leu, N.A., Xu, Y., and Wang, P.J. (2022). Recurrent pregnancy loss in mice lacking the X-linked Ccnb3 genedagger. Biology of reproduction *106*, 382-384.

de Napoles, M., Nesterova, T., and Brockdorff, N. (2007). Early loss of Xist RNA expression and inactive X chromosome associated chromatin modification in developing primordial germ cells. PLoS One *2*, e860.

Degner, J.F., Marioni, J.C., Pai, A.A., Pickrell, J.K., Nkadori, E., Gilad, Y., and Pritchard, J.K. (2009). Effect of read-mapping biases on detecting allele-specific expression from RNA-sequencing data. Bioinformatics *25*, 3207-3212.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Engreitz, J.M., Pandya-Jones, A., McDonel, P., Shishkin, A., Sirokman, K., Surka, C., Kadri, S., Xing, J., Goren, A., Lander, E.S.*, et al.* (2013). The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. Science *341*, 1237973.

Gura, M.A., Mikedis, M.M., Seymour, K.A., de Rooij, D.G., Page, D.C., and Freiman, R.N. (2020). Dynamic and regulated TAF gene expression during mouse embryonic germ cell development. PLoS genetics *16*, e1008515.

Hackett, J.A., Sengupta, R., Zylicz, J.J., Murakami, K., Lee, C., Down, T.A., and Surani, M.A. (2013). Germline DNA demethylation dynamics and imprint erasure through 5-hydroxymethylcytosine. Science *339*, 448-452.

Huang, T.C., Wang, Y.F., Vazquez-Ferrer, E., Theofel, I., Requena, C.E., Hanna, C.W., Kelsey, G., and Hajkova, P. (2021). Sex-specific chromatin remodelling safeguards transcription in germ cells. Nature *600*, 737-742.

Johnston, P.G. (1981). X chromosome activity in female germ cells of mice heterozygous for Searle's translocation T(X;16)16H. Genetical research *37*, 317-322.

Kay, G.F., Penny, G.D., Patel, D., Ashworth, A., Brockdorff, N., and Rastan, S. (1993). Expression of Xist during mouse development suggests a role in the initiation of X chromosome inactivation. Cell *72*, 171-182.

Keane, T.M., Goodstadt, L., Danecek, P., White, M.A., Wong, K., Yalcin, B., Heger, A., Agam, A., Slater, G., Goodson, M.*, et al.* (2011). Mouse genomic variation and its effect on phenotypes and gene regulation. Nature *477*, 289-294.

Koopman, P., Munsterberg, A., Capel, B., Vivian, N., and Lovell-Badge, R. (1990). Expression of a candidate sex-determining gene during mouse testis differentiation. Nature *348*, 450-452.

Kratzer, P.G., and Chapman, V.M. (1981). X chromosome reactivation in oocytes of Mus caroli. Proceedings of the National Academy of Sciences of the United States of America *78*, 3093-3097.

Krueger, F., and Andrews, S.R. (2016). SNPsplit: Allele-specific splitting of alignments
between genomes with known SNP genotypes. F1000Res *5*, 1479.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.

Loda, A., Collombet, S., and Heard, E. (2022). Gene regulation in time and space during X-
chromosome inactivation. Nat Rev Mol Cell Biol *23*, 231-249.

Lowe, M.G., Yen, M.R., Hsu, F.M., Hosohama, L., Hu, Z., Chitiashvili, T., Hunt, T.J., Gorgy, I.,
Bernard, M., Wamaitha, S.E.*, et al.* (2022). EED is required for mouse primordial germ cell
differentiation in the embryonic gonad. Dev Cell *57*, 1482-1495 e1485.

Lun, A.T., and Smyth, G.K. (2014). De novo detection of differentially bound regions for ChIP-
seq data using peaks and windows: controlling error rates correctly. Nucleic Acids Res *42*,
e95.

Mak, W., Baxter, J., Silva, J., Newall, A.E., Otte, A.P., and Brockdorff, N. (2002). Mitotically
stable association of polycomb group proteins eed and enx1 with the inactive x chromosome
in trophoblast stem cells. Current biology : CB *12*, 1016-1020.

Marahrens, Y., Loring, J., and Jaenisch, R. (1998). Role of the Xist gene in X chromosome
choosing. Cell *92*, 657-664.

Marahrens, Y., Panning, B., Dausman, J., Strauss, W., and Jaenisch, R. (1997). Xist-deficient
mice are defective in dosage compensation but not spermatogenesis. Genes Dev *11*, 156-
166.

Masui, O., Corbel, C., Nagao, K., Endo, T.A., Kezuka, F., Diabangouaya, P., Nakayama, M.,
Kumon, M., Koseki, Y., Obuse, C.*, et al.* (2023). Polycomb repressive complexes 1 and 2 are
each essential for maintenance of X inactivation in extra-embryonic lineages. Nat Cell Biol *25*,
134-144.

Monk, M., and McLaren, A. (1981). X-chromosome activity in foetal germ cells of the mouse.
J Embryol Exp Morphol *63*, 75-84.

Pan, J., Eckardt, S., Leu, N.A., Buffone, M.G., Zhou, J., Gerton, G.L., McLaughlin, K.J., and
Wang, P.J. (2009). Inactivation of Nxf2 causes defects in male meiosis and age-dependent
depletion of spermatogonia. Dev Biol *330*, 167-174.

Penny, G.D., Kay, G.F., Sheardown, S.A., Rastan, S., and Brockdorff, N. (1996). Requirement
for Xist in X chromosome inactivation. Nature *379*, 131-137.

Picelli, S., Faridani, O.R., Bjorklund, A.K., Winberg, G., Sagasser, S., and Sandberg, R.
(2014). Full-length RNA-seq from single cells using Smart-seq2. Nature protocols *9*, 171-181.

Popp, C., Dean, W., Feng, S., Cokus, S.J., Andrews, S., Pellegrini, M., Jacobsen, S.E., and
Reik, W. (2010). Genome-wide erasure of DNA methylation in mouse primordial germ cells is
affected by AID deficiency. Nature *463*, 1101-1105.

Ramirez, F., Ryan, D.P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dundar,
F., and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data
analysis. Nucleic Acids Res *44*, W160-165.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics *26*, 139-140.

Rolland, A.D., Lehmann, K.P., Johnson, K.J., Gaido, K.W., and Koopman, P. (2011).
Uncovering gene regulatory networks during mouse fetal germ cell development. Biology of
reproduction *84*, 790-800.

Saitou, M., Kagiwada, S., and Kurimoto, K. (2012). Epigenetic reprogramming in mouse pre-
implantation development and primordial germ cells. Development *139*, 15-31.

Sangrithi, M.N., Royo, H., Mahadevaiah, S.K., Ojarikre, O., Bhaw, L., Sesay, A., Peters, A.H.,
Stadler, M., and Turner, J.M. (2017). Non-Canonical and Sexually Dimorphic X Dosage
Compensation States in the Mouse and Human Germline. Dev Cell *40*, 289-301 e283.

Sangrithi, M.N., and Turner, J.M.A. (2018). Mammalian X Chromosome Dosage
Compensation: Perspectives From the Germ Line. Bioessays *40*, e1800024.

Seisenberger, S., Andrews, S., Krueger, F., Arand, J., Walter, J., Santos, F., Popp, C.,
Thienpont, B., Dean, W., and Reik, W. (2012). The dynamics of genome-wide DNA methylation
reprogramming in mouse primordial germ cells. Mol Cell *48*, 849-862.

Severino, J., Bauer, M., Mattimoe, T., Arecco, N., Cozzuto, L., Lorden, P., Hamada, N.,
Nosaka, Y., Nagaoka, S.I., Audergon, P.*, et al.* (2022). Controlled X-chromosome dynamics
defines meiotic potential of female mouse in vitro germ cells. EMBO J *41*, e109457.

Skene, P.J., and Henikoff, S. (2017). An efficient targeted nuclease strategy for high-resolution
mapping of DNA binding sites. Elife *6*.

Sugimoto, M., and Abe, K. (2007). X chromosome reactivation initiates in nascent primordial
germ cells in mice. PLoS genetics *3*, e116.

Tam, P.P., Zhou, S.X., and Tan, S.S. (1994). X-chromosome activity of the mouse primordial
germ cells revealed by the expression of an X-linked lacZ transgene. Development *120*, 2925-
2932.

Tang, W.W., Kobayashi, T., Irie, N., Dietmann, S., and Surani, M.A. (2016). Specification and
epigenetic programming of the human germ line. Nature reviews Genetics *17*, 585-600.

van de Geijn, B., McVicker, G., Gilad, Y., and Pritchard, J.K. (2015). WASP: allele-specific
software for robust molecular quantitative trait locus discovery. Nat Methods *12*, 1061-1063.

Wang, P.J., McCarrey, J.R., Yang, F., and Page, D.C. (2001). An abundance of X-linked genes
expressed in spermatogonia. Nat Genet *27*, 422-426.

Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene
expression data analysis. Genome Biol *19*, 15.

Xu, J., Carter, A.C., Gendrel, A.V., Attia, M., Loftus, J., Greenleaf, W.J., Tibshirani, R., Heard,
E., and Chang, H.Y. (2017). Landscape of monoallelic DNA accessibility in mouse embryonic
stem cells and neural progenitor cells. Nat Genet *49*, 377-386.

Yeom, Y.I., Fuhrmann, G., Ovitt, C.E., Brehm, A., Ohbo, K., Gross, M., Hubner, K., and Scholer,
H.R. (1996). Germline regulatory element of Oct-4 specific for the totipotent cycle of
embryonal cells. Development *122*, 881-894.

Yoshimizu, T., Sugiyama, N., De Felice, M., Yeom, Y.I., Ohbo, K., Masuko, K., Obinata, M.,
Abe, K., Scholer, H.R., and Matsui, Y. (1999). Germline-specific expression of the Oct-4/green
fluorescent protein (GFP) transgene in mice. Development, growth & differentiation *41*, 675-
684.

Yu, J.R., Lee, C.H., Oksuz, O., Stafford, J.M., and Reinberg, D. (2019). PRC2 is high
maintenance. Genes Dev *33*, 903-935.

Zheng, H., Huang, B., Zhang, B., Xiang, Y., Du, Z., Xu, Q., Li, Y., Wang, Q., Ma, J., Peng, X.*,
et al.* (2016). Resetting Epigenetic Memory by Reprogramming of Histone Modifications in
Mammals. Mol Cell *63*, 1066-1079.
876