Research Article

Using Extracellular miRNA Signatures to Identify Patients with LRRK2-Related

Parkinson's Disease

Luca Jannik Braunger[a]*, Felix Knab[a]* and Thomas Gasser[a,b]

[a]Hertie Institute for Clinical Brain Research, Department of Neurodegeneration,

University of Tuebingen, Tuebingen, Germany

[b]German Center for Neurodegenerative Diseases (DZNE), Tuebingen, Germany

*Authors contributed equally

Running title: LRRK2-Driven miRNA Patterns

Corresponding author:

Felix Knab, felix.knab@med.uni-tuebingen.de, 0049 7071 29 87608

Conflict of interest: The authors declare that they have no conflict of interest.

Abstract

Background: Mutations in the Leucine Rich Repeat Kinase 2 gene are highly relevant in both sporadic and familial cases of Parkinson's disease. Specific therapies are entering clinical trials but patient stratification remains challenging. Dysregulated microRNA expression levels have been proposed as biomarker candidates in sporadic Parkinson's disease.

Objective: In this proof-of concept study we evaluate the potential of extracellular miRNA signatures to identify LRRK2-driven molecular patterns in Parkinson's disease.

Methods: We measured expression levels of 91 miRNAs via RT-qPCR in ten individuals with sporadic Parkinson's disease, ten *LRRK2* mutation carriers and eleven healthy controls using both plasma and cerebrospinal fluid. We compared miRNA signatures using heatmaps and t-tests. Next, we applied group sorting algorithms and tested sensitivity and specificity of their group predictions.

Results: miR-29c-3p was differentially expressed between *LRRK2* mutation carriers and sporadic cases, with miR-425-5p trending towards significance. Individuals clustered in principal component analysis along mutation status. Group affiliation was predicted with high accuracy in the prediction models (sensitivity up to 89%, specificity up to 70%). miRs-128-3p, 29c-3p, 223-3p and 424-5p were identified as promising discriminators among all analyses.

Conclusions: LRRK2 mutation status impacts the extracellular miRNA signature measured in plasma and separates mutation carriers from sporadic Parkinson's disease patients. Monitoring LRRK2 miRNA signatures could be an interesting approach to test drug efficacy of LRRK2-targeting therapies. In light of small sample size, the suggested approach needs to be validated in larger cohorts.

44   **Keywords:** [Parkinson's disease, LRRK2, microRNA, miRNA, Biomarker]

Introduction

The majority of familial PD (fPD) cases is caused by mutations in the Leucine Rich Repeat Kinase 2 (*LRRK2*) gene [1,2], with the G2019S mutation being the most common [3]. Despite its apparent role in PD, LRRK2 expression levels in the CNS are relatively low compared to peripheral tissues such as the lung and the kidneys, as well as blood monocytes [4,5]. Increased kinase activity is thought to be the cause of the disturbed cellular homeostasis observable in cell models of PD [6]. Most importantly, LRRK2 was shown to play a role in sporadic cases, with non-coding variants in the *LRRK2* gene increasing the risk of PD [7]. Further, sporadic PD (sPD) and fPD cases show overlapping clinical features [8]. Studying fPD cases therefore offers the opportunity to better understand the molecular pathogenesis of both fPD and sPD cases. Targeting multiple molecular pathways in selected PD patients with an individually formulated combination of drugs could be a potential strategy for disease modification [9]. However, this strategy necessitates not only the discovery of particular druggable pathways, but also the careful selection of patients who are most likely to benefit from a given treatment, ideally at an early or even asymptomatic stage in the course of their disease. While aberrant LRRK2 activity represents such a molecular target and LRRK2 inhibitors are approaching clinical testing [10], [11], pre-selecting patients as well as monitoring target engagement and drug efficacy remain challenging. There are no established biomarkers available that can aid in the early detection of disease onset before significant neurodegeneration occurs.

Recently, microRNAs (miRNAs), small non-coding RNA molecules that are involved in the posttranscriptional regulation of gene expression [12], have been intensively studied as potential biomarkers for neurodegenerative diseases, including PD [13,14]. So far, most biomarker studies using miRNAs have focused on the differentiation of

sPD patients from healthy controls (HC), while in the present proof-of-concept study

we aimed to test how well extracellular miRNAs can be used to reliably differentiate

between sPD patients and *LRRK2* mutation carriers (LRRK2$_{MC}$). Since the distinction

between fPD and sPD is possible by using more stringent methods, such as sequencing,

in the long run we aim to identify individuals with relevant involvement of LRRK2-

dependent pathways among the sPD population. However, in a first step we decided to

define what constitutes a molecular LRRK2 fingerprint by comparing the extracellular

miRNA signatures of fPD and sPD cases. Given that LRRK2 was described to change

the cellular miRNAome [15], we hypothesized that mutations in the *LRRK2* gene also

alter the extracellular miRNA signature in LRRK2$_{MC}$.

Materials and Methods

Experimental Design

In a first step, we quantified the expression levels of 91 extracellular miRNAs from a cohort of ten sPD, ten fPD and eleven HC in both plasma and CSF using RT-qPCR (Fig. 1). After processing of raw Ct values and calculation of log2 fold change values (log2fc), as described in more detail in the respective method section, we performed t-tests to identify single miRNAs with the potential of discriminating between groups. We further applied group sorting algorithms such as principal component analysis (PCA), least absolute shrinkage and selection operator (LASSO) and random forest (RF) using the calibrated Ct values of all miRNAs that could be reliably detected throughout the cohort. The ability to differentiate between sPD patients and $LRRK2_{MC}$ with and without PD was assessed for each sorting algorithm by evaluating their sensitivity and specificity. Finally, we selected those miRNAs that had a significant impact on the performance of each model.

Cohort Design

For each $LRRK2_{MC}$, one sPD and one HC was added to the study cohort. To obtain homogenous groups, individuals were selected based on age, gender, and disease duration. As the histopathological complexity of PD likely increases over time and patients with long disease durations might no longer display pathologies specific to mutation status rather than showing features common to all PD patients, disease durations were kept as short as possible. The group of $LRRK2_{MC}$ initially included a PD patient carrying two *LRRK2* variants (N1437S, S1647). Given that these polymorphisms were not reported to be pathogenic, we finally decided to exclude this patient from the analysis. The final cohort therefore comprised three groups (Table 1):

Ten LRRK2$_{MC}$ (fPD patients: n=6 (*LRRK2* G2019S: n = 3; *LRRK2* G2019S/G1819:

n=1, *LRRK2* R1441C: n=1; *LRRK2* I2020T: n=1), asymptomatic G2019S mutation

carriers: n=4), ten sPD patients and eleven HCs. After performing PCA as described in

more detail in the respective method section, the data set of one of the asymptomatic

LRRK2$_{MC}$ appeared as a technical or biological outlier and was removed from the

plasma dataset before further analysis (Fig. S1). Clinical features are reported in Table

1. The researcher performing the experiments was blinded to the group annotation.

miRNA selection and qPCR panel design

We scanned the literature for miRNAs that were previously reported to be dysregulated

in the context of *LRRK2* mutations, functionally relevant to LRRK2 or generally

dysregulated in the context of PD. Next, we selected a total of 91 miRNAs to be

included in our customized qPCR panel (Table S1). Selection was based on a set of

criteria, including the number of relevant studies reporting dysregulation and a

preference for miRNAs assessed in CSF and plasma samples over miRNAs reported in

brain tissue or animal models. Additionally, miRNAs associated with LRRK2-linked

PD were given priority over those associated with sporadic PD or other

neurodegenerative diseases. For quality control purposes, we included a set of spike-

in controls. Those included UniSpike 2 and 4 from the QIAGEN RNA Spike-In Kit

(Cat. No.: 339390, QIAGEN, Hilden, Germany), which were used to monitor RNA

isolation efficacy, UniSpike 6 (included in the QIAGEN miRCURY LNA RT Kit) to

verify the efficacy of reverse transcription (RT) and UniSpike 3 (pre-applied primers

on the custom qPCR plates) for inter-plate calibration. Additionally, a blank spot

containing no primer was included and functioned as a negative control.

Collection and Storage of Biofluids from Study Participants

Blood and CSF samples were collected from individuals according to standardized

protocols. Within 90 minutes after collection from the individual, the collected material

was brought from the medical facilities to the Neuro-Biobank Tuebingen and

immediately processed. CSF and EDTA plasma were centrifuged at 2000g for 10

minutes. Samples were then transferred to a 15 ml Falcon tube, mixed thoroughly by

vortexing, and aliquoted to cryotubes. Finally, both CSF and plasma cryotubes were

frozen and stored at -80°C in the biobank. Before RNA isolation, samples were slowly

thawed on ice.

RNA isolation

Extracellular RNA was isolated from plasma and CSF using the QIAGEN miRNeasy

Serum/Plasma Advanced Kit (Cat. No: 217204, QIAGEN) and following the vendor's

instructions with slight modifications. Briefly, 200 µl of plasma or 400 µl of CSF were

used as a starting volume and subsequent reaction volumes were adjusted accordingly.

After addition of RPL buffer and incubation for three minutes, we added 1 µl of the

spike-in mix provided in the QIAGEN RNA Spike-In Kit (Cat. No.: 339390, QIAGEN)

to the tube, in order to monitor the efficacy of the RNA isolation. Finally, RNA elution

volume was reduced to 7 µl of RNase-free water.

Reverse Transcription

The RT reaction was performed using the QIAGEN miRCURY LNA RT Kit (Cat. No.:

339340, QIAGEN). The reaction mix included 2 µl of 5x miRCURY SYBR® Green

RT Reaction Buffer, 1 µl of 10x miRCURY RT Enzyme Mix, 4.5 µl of RNase-free

water, 0.5 µl of UniSpike 6 RNA spike-in and 2 µl of template RNA per sample. After

brief centrifugation, tubes were incubated at 42 °C for 60 minutes, followed by 5

minutes of incubation at 95 °C, which inactivated the RT reaction. Finally, samples

were diluted 1:40 using RNase-free water.

qPCR

qPCR was performed using QIAGEN miRCURY LNA miRNA Custom PCR Panels

(Catalog No.: 339330, QIAGEN) along with the QIAGEN miRCURY LNA SYBR

Green PCR Kit (Cat. No.: 339347, QIAGEN). The reaction mix included 515 µl of 2x

miRCURY SYBR® Green Master Mix, 115 µl of RNase-free water and 400 µl of the

cDNA template per sample. After pipetting 10 µl into each reaction-well, which already

contained the dried primers, the plates were sealed and vortexed for one minute. After

brief centrifugation, plates were incubated for ten minutes at room temperature to allow

primers to resolve. Finally, plates were vortexed for one minute and briefly centrifuged.

Amplification and quantification were performed using a LightCycler® 480 instrument

(Roche, Basel, Switzerland). The cycling program included a two-minute heat

activation step at 95 °C, followed by 45 cycles of denaturation at 95 °C for ten seconds

and annealing at 56 °C for 60 seconds. To ensure specificity of amplifications, melting

points of the amplified products were analyzed (Table S2 and S3).

Processing of raw data

Before applying statistical analysis, raw data were processed and Ct and fold-change

values (fc) were calculated (Fig. S2): First, raw data from the qPCR were converted

using the LC480Conversion software to make them readable for LinRegPCR (Version

11.0, [16]), which was used for calculating Ct values. Further processing and analysis

of data was performed in R (Version 4.2.2) using RStudio (Version 2023.03.0+386)

[17]. To account for variations between plates, the Ct values were normalized using an

interplate calibration factor. The calibration factor for each plate was calculated by

subtracting the mean Ct value of UniSpike 3 reactions from all plates from the mean Ct

value of all UniSpike 3 reactions in a given plate. All Ct values of the respective plate

were then calibrated by subtracting the respective calibration factor from all Ct values

in that plate. After calibration, Ct values > 40 were considered unspecific and excluded

from the dataset. If a miRNA was not specifically amplified in more than one

individual, it was removed completely from the analysis, after confirming that missing

values occurred equally over all groups (Fig. S3). This resulted in 58 remaining

miRNAs for the plasma dataset and eleven miRNAs for the CSF dataset.

Next, the fc and log2fc values were calculated using the $2^{-\Delta\Delta Ct}$ method (Fig. S4) [18].

In a first step, for each individual we independently calculated the mean Ct by adding

all Ct values of every quantified miRNA and dividing it by the number of quantified

miRNAs, which resulted in a $mean_{Ct}$ for each individual. We then computed ΔCt for

each miRNA and individual by subtracting the $mean_{Ct}$ of the respective individual from

every miRNA quantified in that individual. This resulted in multiple ΔCt_{miR-n} per

individual, where n represents the specific miRNA. The entire group of healthy controls

was used as a reference group. Subsequently, for each miRNA we calculated the mean

of ΔCt_{miR-n}, only using ΔCt_{miR-n} from the HCs ($mean\ \Delta Ct_{miR-n-healthy}$). Next, $\Delta\Delta$Ct for each

miRNA and individual was determined by subtracting the $mean\ \Delta Ct_{miR-n-healthy}$ from the

patient's ΔCt_{miR-n}. Finally, the fold change was obtained using the following formula:

fold change $= 2^{-\Delta\Delta Ct}$. To achieve a linear scale with symmetry around zero, log2fc was

used for the heatmaps and t-tests. Data from the HCs were used only as a reference for

calculating the fold change values. Since the focus of this study was put on the

identification of a LRRK2-driven molecular patterns and the discrimination between

fPD and sPD cases, in the subsequent analyses, data from the HCs were not analyzed

further.

Data Visualization

Heatmaps visualizing either the log2fc or calibrated Ct values were created using the

pheatmap package in R [19]. Log2fc and Ct values were scaled row-wise to z-scores

using the *scale* function while ignoring missing values. After confirming normality of

the data via assessing QQ-Plots, an unpaired t-test using the log2fc values was

performed for each miRNA, comparing the sPD group with the LRRK2$_{MC}$. The p-value

threshold to determine significance was corrected for multiple testing by dividing by

the number of tests. The distribution of p-values was examined and visualized via

histograms. For miRNAs that the t-test revealed to be significantly differentially

expressed between the groups, ROC analysis was performed. PCA was performed on

calibrated Ct values using the built-in *prcomp* function of R. As the function does not

accept missing values, they were imputed by setting them to the mean Ct value from

all individuals. Two-dimensional plots were generated by graphing PC1 and PC2

values.

Least Absolute Shrinkage and Selection Operator regression models

The *caret* package [20] was used for building the LASSO regression models. For

missing values, the mean of the Ct values of all patients was used. We selected

sensitivity and specificity in differentiating between the sPD and LRRK2$_{MC}$ group,

determined by Leave-One-Out-Cross-Validation (LOOCV) [21], as our experimental

read-outs.

Random forest Prediction Model

For building a RF model that predicts group affiliation to either the sPD or the LRRK2$_{MC}$ group, the *randomForest* package [22] was used. To obtain mean values of sensitivity and specificity along with 95% confidence intervals, 100 models were constructed with each model containing 200 trees. For the RF model we did not use LOOCV, as sensitivity and specificity of the classification is already tested on out-of-bag samples reducing the risk of over-fitting. Along with this, we generated proximity heatmaps describing similarity of patient pairs based on the number of trees that classify those two patients in the same terminal node. Missing values were computed using the *rfImpute* function. The Gini coefficient's mean decrease, which describes each decision node in terms of classification accuracy, was used to evaluate the importance of each variable. High purity of a node results in a low Gini coefficient. Consequently, variables that are more crucial for distinguishing between groups exhibit a greater mean decrease of the Gini coefficient compared to others.

Integration of CSF and Plasma data sets

To integrate the information from both the plasma and CSF datasets, combined variables were computed. First, Pearson's correlation analysis was performed between the Ct values of the eleven miRNAs reliably detectable in CSF (Ct$_{CSF}$) and the Ct values of the corresponding miRNAs detected in plasma (Ct$_{plasma}$). A two-tailed p-value cut-off was applied, and alpha was set to 0.05. All significant combinations were used to form new variables by multiplying the corresponding Ct values (Ct$_{multiplied}$). This generated 20 new variables, which were subsequently used to create a heatmap, perform t-tests and PCA and generate LASSO and RF models.

Identification of discriminatory miRNAs

In a final step, we wanted to select those miRNAs that most accurately discriminated between groups. This feature selection was performed for each of the analyses using the following criteria: 1) p-values resulting from the unpaired t-test were sorted in ascending order and the top five miRNAs with the smallest p-value were selected; 2) the loading scores of PC1 were assessed and the top five miRNAs were selected; 3) the five most influential miRNAs from the RF model were selected based on the mean decrease of the Gini coefficient; 4) the LASSO regression model automatically selects discriminatory features through shrinkage and elimination of less relevant variables by introducing penalties; in this case, three miRNAs were selected. As the CSF data set alone had proven to not efficiently discriminate between groups, these selection steps were only applied to the Ct_{plasma} and $Ct_{multiplied}$ data sets.

Target prediction enrichment analysis

We used the miRDB database [23] to predict gene targets of selected miRNAs. Next, for each miRNA we selected the top 100 genes sorted by Target score and performed Gene Ontology (GO) enrichment analysis in R using the ontology class "*biological processes*", the database org.HS.eg.db and the clusterProfiler package [24].

Results

Detection of miRNA Expression Patterns in Plasma and CSF

In plasma, a total of 58 miRNAs passed our selection criteria. 23 miRNAs showed stable expression in 29/30 individuals while the remaining 35 miRNAs were detected in all 30 individuals. When visualizing log2fc values using a heatmap, the group of LRRK2$_{MC}$, which included both symptomatic and asymptomatic mutation carriers, showed similar expression patterns and could be distinguished from the sPD group (Fig. 2A). When using the calibrated Ct$_{plasma}$ data, the expression patterns became less apparent but still noticeable (Fig. S5A). In contrast, in CSF only eleven miRNAs survived our selection process, with five miRNAs detected in all 31 individuals and six miRNAs in 30/31 individuals. Expression levels in CSF were generally lower compared to plasma. In CSF, no clear clustering was observable (Fig. 2B). Heatmaps based on calibrated Ct$_{CSF}$ also did not show any clear observable group-specific clusters (Fig. S5B).

Identification of single discriminatory miRNAs through t-test

After performing unpaired t-test on log2fc values in plasma, a total of 17 miRNAs were below a p-value threshold of 0.05 (Table S4). After correction for multiple comparisons by adjusting the p-value threshold to 0.0009, miR-29c-3p showed a significant difference (LRRK2$_{MC}$: -0.39 (SD: ±0.48); sPD: 0.62 (SD: ±0.54), t(56) = 4.21, p = 0.0007). ROC analysis of miR-29c-3p revealed a sensitivity of 90% and specificity of 90% with an AUC of 0.86 (Fig. S6). In CSF, only miR-223-3p showed a p-value below 0.05 (LRRK2$_{MC}$: -0.49 (SD: ±1.05); sPD: 0.62 (SD: ±1.20), t(9) = -2.2, p = 0.041) but due to correction for multiple testing did not pass the p-value threshold of 0.0045.

P-values in the plasma data set were not equally distributed but concentrated at the

lower end of the scale (Fig. 2C), while the p-values in the CSF data set appeared to be

randomly distributed (Fig. 2D). This indicates that the miRNA expression levels

measured in CSF alone seem to contain little information on group discrimination,

while in plasma a subset of miRNAs could be used to identify $LRRK2_{MC}$. As a

consequence, in the following analyses we primarily focused on the plasma dataset. We

sorted p-values from the plasma data set in ascending order and identified the top 5

miRNAs with the lowest p-value (Fig. 2E). These included miR-128-3p (log2fc in

$LRRK2_{MC}$: -1.01 (SD: ±0.66); sPD: 0.03 (SD: ±0.83), t(56) = 3.06, p = 0.0071), miR-

148a-3p ($LRRK2_{MC}$: -0.60 (SD: ±0.56); sPD: 0.38 (SD: ±0.73), t(56) = 3.31, p =

0.0042), miR-153-3p ($LRRK2_{MC}$: -0.97 (SD: ±1.19); sPD: 0.65 (SD: ±0.69), t(56) =

3.58, p = 0.0035), miR-29c-3p ($LRRK2_{MC}$: -0.39 (SD: ± SD: ±0.48); sPD: 0.62 (SD:

±0.54), t(56) = 4.21, p = 0.0007) and miR-425-5p ($LRRK2_{MC}$: -0.33 (SD: ±0.38); sPD:

0.54 (SD: ±0.55), t(56) = 4.01, p = 0.0010).

Detecting patient clusters using plasma miRNA expression levels and PCA

PCA explained 82% of the overall variance in the plasma dataset. PC1 (77.4% of

variance) explained the majority of the variance and appeared to discriminate groups,

while PC2 only explained 4.6% of the overall variance (Fig. 3A). Interestingly, all

$LRRK2_{MC}$ clustered together while the various *LRRK2* mutations could not be

distinguished (Fig. 3A). PC1 discriminated LRRK2MC from sPD with a sensitivity of

100% (9/9) and a specificity of 70% (7/10). PC1 and PC2 combined reached the same

sensitivity with a specificity of 80% (8/10) (Fig. S7A). Next, we performed PCAs

including the healthy controls (Fig. 4). While clustering was less clear when comparing

HC to sPD, for comparison of HC and $LRRK2_{MC}$ clear clustering was observable (Fig.

4A and 4B). The clearest clustering was achieved when comparing LRRK2$_{MC}$ with sPD

(Fig. 4C). When including all three groups in the PCA, clustering becomes less clear,

yet LRRK2$_{MC}$ remains notably separated from sPD and HCs (Fig. 4D). The PCA of the

CSF data explained 84.2% of the overall variance, with both PC1 (75.4% of variance)

and PC2 (8.8% of variance) not able to differentiate between the groups (Fig. 3B).

miRNA expression levels in CSF correlate to Plasma levels

A total of 20 miRNA combinations from Ct$_{plasma}$ and Ct$_{CSF}$ values showed significant

correlation (Fig. 3C) with R-values ranging from 0.36 to 0.51 (Table S5). Significantly

correlating combinations were used to obtain new variables by multiplication

(Ct$_{multiplied}$).

Multiplied Ct values discriminate groups in t-test and PCA

When creating a heatmap based on the Ct$_{multiplied}$ data, no clear group-specific clusters

were noticeable (Fig. S8A). In the t-test ten miRNA-combinations reached a p-value

below 0.05 (Table S6). The combination of miR-29c-3p from plasma and miR-27a-3p

from CSF passed the adjusted p-value threshold of 0.0025 (LRRK2$_{MC}$: 930.8 (SD:

±61.6); sPD: 813.6 (SD: ±55.9), t(18) = 4.2, p = 0.0007). PCA for Ct$_{multiplied}$

discriminated the groups while explaining 90.2% of the total variance (PC1: 83.2%,

PC2: 7%) (Fig. S8B). Groups were separated based on PC1 with a sensitivity of 100%

(9/9) and a specificity of 80% (8/10). Combining PC1 and PC2 did not improve

sensitivity or specificity (Figure S7B).

LASSO Regression and Random Forest Models predict group membership

LASSO regression and the RF model were performed using the Ct_{plasma} and the

$Ct_{multiplied}$ dataset (Fig. 5A). Sensitivity of the LASSO model was 88.8% (8/9) for both

Ct_{plasma} and $Ct_{multiplied}$. Specificity was 70.0% (7/10) (Ct_{plasma}) and 80.0% (8/10)

($Ct_{multiplied}$), respectively (Fig. 5B). Predictions from the RF models using Ct_{plasma} had a

mean sensitivity of 84.2% (95% CI: 83.1% – 85.3%) and a mean specificity of 70.1%

(95% CI: 69.9 – 70.2%). When using $Ct_{multiplied}$ mean sensitivity was 73.6% (95% CI:

72.5% – 74.6%) and mean specificity was 80 % (95% CI: 80% – 80%) (Figure 5B).

When comparing the two models using an unpaired t-test, sensitivity was significantly

better when using Ct_{plasma} (t(98) = -13.8, p<0.001), while specificity was higher when

using $Ct_{multiplied}$ (t(98) = 99, p<0.001). Classifications performed by the RF model were

further analyzed by looking into proximity scores of patient pairs. The resulting

heatmap for the Ct_{plasma} dataset indicated that patients within the same group exhibit

greater proximity to each other than patients from different groups (Fig. 5C). This

becomes even more apparent in the model using $Ct_{multiplied}$ (Fig. 5D). Next, mean

decrease of Gini scores were calculated for both of these RF models (Fig. 5E and 5F)

to assess variable influence on classification accuracy. Interestingly, in both RF models,

miR-223-3p had the highest score, indicating a great impact on the model performance.

This miRNA had already been observed in the plasma t-test, where it indicated a

potential for group discrimination with an uncorrected p-value of p=0.051.

Selection of discriminatory miRNAs identified by multiple tests or models

Finally, for each test or model, we identified those miRNAs, that showed the greatest

potential for group separation or prediction (Table 2). From the RF models, PCA and

the t-tests, we extracted five miRNAs each, while the LASSO model identified three

miRNAs. Interestingly, of the selected miRNAs, three were identified by more than one

method; 1) miR-29c-3p was identified in t-tests and LASSO, 2) miR-128-3p in t-test

and RF, and 3) miR-424-5p in LASSO and RF (Fig. 5G, Fig. S9). GO analysis was

performed on the four miRNAs and we identified associated biological functions and

processes (number of significant annotations identified: miR-223-3p: 25, miR-29c-3p:

60, miR-128-3p: 0, miR-424-5p: 3, see Figure S10). Interestingly, the annotations for

miR-223-3p included neuron death, regulation of neuron death and neuron apoptotic

process.

Discussion

In this proof-of-concept study, we examined the extracellular miRNA signatures in plasma and CSF derived from LRRK2$_{MC}$ and sPD patients for LRRK2-dependent patterns. We discovered that plasma miRNA expression levels can distinguish sPD from LRRK2$_{MC}$. MiRNAs have been extensively studied in PD, but to the best of our knowledge, this is the first study to use machine learning and a broad data set of miRNA expression levels to distinguish genetic from sporadic PD.

Our results show that PCA separated LRRK2$_{MC}$ from most individuals with sPD without differentiating between the various *LRRK2* mutations. Separation was also observable in PCA comparing LRRK2$_{MC}$ and HC as well as in an analysis comparing all three groups. This indicates that the LRRK2 mutation status has a measurable effect on the extracellular miRNA signature in LRRK2$_{MC}$. Some sPD individuals clustered in proximity to the LRRK2$_{MC}$ individuals, raising the interesting possibility that this method could be used to identify patients who would benefit from a LRRK2 targeted therapy. However, this question needs to be addressed using larger cohorts in the context of LRRK2-inhibitor trials. Regular assessment of the identified miRNA signatures could help monitor treatment efficacy in therapies targeting LRRK2, where a shift in miRNA expression levels could indicate target engagement.

We extracted discriminatory miRNAs from each analysis and found that some miRNAs were overlapping between models, namely miR-29c-3p, miR-128-3p, miR-424-5p and miR-223-3p.

miR-29c-3p has been reported to play a role in disease modulation by mediating neuroinflammation and has been found to be involved in apoptotic processes [25]. It has further been proposed as a PD specific marker in several studies, but contradicting findings on expression levels compared to controls exist: studies report either

downregulation [26,27] or upregulation [28] in serum of sPD patients compared to HCs.

One study found upregulation in serum of sPD patients, but no dysregulation in LRRK2

associated PD [29].

miR-223-3p has been reported to be upregulated in midbrain dopamine neurons [30]

and serum of PD patients [31]. It was shown to be involved in the modulation of

inflammasome activity [32]. LRRK2 has long been suspected to play a role in

inflammation and e.g., was shown to affect microglial activation and pro-inflammatory

cytokine production [33]. Further, the GO annotations identified for miR-223-3p

included processes specific to neurodegeneration underlining a possible influence in

neurodegenerative disorders such as PD.

miR-128-3p has previously been identified as a potential treatment target due to its

ability to protect neurons from apoptosis. The upregulation in the sPD group could

therefore be reflective of a compensatory mechanism, that might not be relevant in

individuals carrying a *LRRK2* mutation.

miR-424-5p was shown to be increased in the forebrain of PD patients, while being

associated with FOXO1 [34]. FOXO1 activity is induced by LRRK2 [35] and thereby

links altered miR-424-5p levels to LRRK2 activity.

The low RNA abundance in CSF compared to plasma made miRNA detection difficult,

resulting in the exclusion of many miRNAs from CSF analyses. This reduced the

complexity of miRNA signatures we could assess in CSF, which may explain the poor

performance of CSF-based analyses. An alternative explanation could be that LRRK2

protein expression levels are known to be low in the CNS [4,5] despite their apparent

relevance in the pathophysiology of PD. In contrast, peripheral organs such as the lung

and the kidney display higher expression levels, which could also explain the increased

accuracy of models based on Ct_{plasma} miRNA profiles in identifying $LRRK2_{MC}$. We

have further analyzed the relation of miRNA expression levels in CSF and plasma and found that for most miRNAs, these two biofluids seem to display very different signatures. However, through correlation analysis we have identified a subset of miRNAs whose expression levels in the CNS and the periphery seem related. We convoluted correlated miRNA data sets in order to test whether $Ct_{multiplied}$ would provide RF models with significantly higher sensitivity or specificity, which was not the case. While RF models based on $Ct_{multiplied}$ still performed well and slightly different than RF models based on Ct_{plasma}, when facing the discussed classification problem there seems to be no clear benefit from adding CSF data to the model.

In summary, this proof-of-concept study showed promising results and deepens our understanding of LRRK2-associated PD, but it has limitations we want to address. First, the number of individuals included in the present study is relatively small. When utilizing group sorting algorithms for classification, larger sample sizes improve robustness and replicability. Ideally, the data set should be divided into a training and a testing data set to avoid overfitting. Additionally, a completely independent data set should then be used to replicate the predictions. While we did perform cross-validation using LOOCV or out-of-bag samples, this cannot fully replace validation in a testing or replication cohort. The results therefore have to be considered with caution until replicated in larger cohorts. Further, we selected miRNAs to be included in our qPCR panel based on the literature and therefore only considered miRNAs that have already been reported as dysregulated in PD or other neurodegenerative diseases. This may have introduced a bias and preventing discovery of novel contributing miRNAs. Finally, we found no evident advantage to employing multi-layered or composited readouts over standard t-tests. While we still believe that in diseases as complex as PD

basing classifications on multiple variables is an interesting approach, this concept has

still to be improved and repeated in larger and more complex cohorts.

Conclusion

In conclusion, in this proof-of-concept study we showed that LRRK2 mutation status

impacts the extracellular miRNA signature measured in plasma and shows promise to

separate LRRK2$_{MC}$ from sPD. Monitoring changes of the extracellular miRNA

signatures upon e.g. LRRK2 inhibition could be used to study drug efficacy or target

engagement. The potential of multi-layered approaches trying to identify sporadic PD

patients with a relevant role of LRRK2-related pathways are another potential

application which needs to be thoroughly assessed in larger cohorts.

List of Abbreviations

CSF: Cerebrospinal fluid

Ct_{CSF}: Ct values from CSF dataset

$Ct_{multiplied}$: multiplied Ct values from CSF and plasma dataset

Ct_{plasma}: Ct values from plasma dataset

Fc: Fold change

GO: Gene Ontology

HC: Healthy control

LASSO: least absolute shrinkage and selection operator

LEDD: levodopa equivalent daily dose

Log2fc: $\log_2$(Fold change)

LOOCV: Leave-One-Out Cross-Validation

LRRK2: Leucine-Rich Repeat Kinase 2

$LRRK2_{MC}$: *LRRK2* mutation carrier

miRNA: microRNA

MoCA: Montreal Cognitive Assessment

PCA: Principal component analysis

PD: Parkinson's disease

qPCR: real-time quantitative PCR

RF: Random Forest

RT: Reverse transcription

sPD: sporadic PD

UPDRS: Unified Parkinson Disease Rating Scale

Declarations

Ethics approval and consent to participate

The study was approved by the Hertie Institute for Clinical Brain Research Biobank and the ethics committee of the medical faculty of the University of Tübingen and University Clinic Tübingen (project ID: 199/2011B01). All participants gave written, informed consent.

Availability of data and materials

The generated and analyzed data are available from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The project was supported and funded by the Johannes-Dichgans-scholarship, the Deutsche Forschungsgemeinschaft (Projectnumber D27.15350), the Hertie Stiftung and the Eva Theers Stiftung.

Authors' Contributions

FK conceptualized the study. FK and TG supervised the study. LJB performed the experiments and the statistical analyses. LJB generated all figures. LJB and FK drafted a first version of the manuscript. All authors read, edited and approved the final manuscript.

Acknowledgements

Samples were obtained from the Neuro-Biobank of the University of Tuebingen, Germany (https://www.hih-tuebingen.de/en/about-us/core-facilities/biobank/). This biobank is supported by the local University, the Hertie Institute and the DZNE.

Supplementary Material

Figure S1: Identification of an outlier via PCA

Figure S2: Processing of Raw data

Figure S3: Group-wise Expression Distribution of miRNAs

Figure S4: Calculation of $\Delta\Delta$Ct values

Figure S5: Heatmaps based on Ct values

Figure S6: ROC curve for miR-29c-3p

Figure S7: PCA graphs including for evaluation of sensitivity and specificity

Figure S8: Heatmap and PCA resulting from multiplied variables

Figure S9: Scatterplots of log2fc values for the four discussed miRNAs

Figure S10: GO analysis of identified miRNAs

Table S1: included miRNAs with literature references

Table S2: Melting points of miRNAs detected in plasma

Table S3: Melting points of miRNAs detected in CSF

Table S4: Statistic details from plasma t-tests

Table S5: statistics of correlation matrix

Table S6: t-test with $Ct_{multiplied}$ data

539   **References**

540   [1]   Zimprich A, Biskup S, Leitner P, Lichtner P, Farrer M, Lincoln S, Kachergus J,

541   Hulihan M, Uitti RJ, Calne DB, Stoessl AJ, Pfeiffer RF, Patenge N, Carbajal IC,

542   Vieregge P, Asmus F, Müller-Myhsok B, Dickson DW, Meitinger T, Strom TM,

543   Wszolek ZK, Gasser T (2004) Mutations in LRRK2 Cause Autosomal-Dominant

544   Parkinsonism with Pleomorphic Pathology. *Neuron* **44**, 601–607.

545   [2]   Paisán-Ruíz C, Jain S, Evans EW, Gilks WP, Simón J, van der Brug M, de

546   Munain AL, Aparicio S, Gil AM, Khan N, Johnson J, Martinez JR, Nicholl D,

547   Carrera IM, Peña AS, de Silva R, Lees A, Martí-Massó JF, Pérez-Tur J, Wood

548   NW, Singleton AB (2004) Cloning of the Gene Containing Mutations that Cause

549   PARK8-Linked Parkinson's Disease. *Neuron* **44**, 595–600.

550   [3]   Correia Guedes L, Ferreira JJ, Rosa MM, Coelho M, Bonifati V, Sampaio C

551   (2010) Worldwide frequency of G2019S LRRK2 mutation in Parkinson's

552   disease: A systematic review. *Parkinsonism Relat Disord* **16**, 237–242.

553   [4]   Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A,

554   Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, Olsson I, Edlund K, Lundberg

555   E, Navani S, Szigyarto CA-K, Odeberg J, Djureinovic D, Takanen JO, Hober S,

556   Alm T, Edqvist P-H, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P,

557   Schwenk JM, Hamsten M, von Feilitzen K, Forsberg M, Persson L, Johansson

558   F, Zwahlen M, von Heijne G, Nielsen J, Pontén F (2015) Tissue-based map of

559   the human proteome. *Science (80-)* **347**,.

560   [5]   The Human Protein Atlas.

561   [6]   Jaleel M, Nichols RJ, Deak M, Campbell DG, Gillardon F, Knebel A, Alessi DR

562   (2007) LRRK2 phosphorylates moesin at threonine-558: characterization of how

563   Parkinson's disease mutants affect kinase activity. *Biochem J* **405**, 307–317.

[7] Simón-Sánchez J, Schulte C, Bras JM, Sharma M, Gibbs JR, Berg D, Paisan-

Ruiz C, Lichtner P, Scholz SW, Hernandez DG, Krüger R, Federoff M, Klein C,

Goate A, Perlmutter J, Bonin M, Nalls MA, Illig T, Gieger C, Houlden H,

Steffens M, Okun MS, Racette BA, Cookson MR, Foote KD, Fernandez HH,

Traynor BJ, Schreiber S, Arepalli S, Zonozi R, Gwinn K, van der Brug M, Lopez

G, Chanock SJ, Schatzkin A, Park Y, Hollenbeck A, Gao J, Huang X, Wood

NW, Lorenz D, Deuschl G, Chen H, Riess O, Hardy JA, Singleton AB, Gasser

T (2009) Genome-wide association study reveals genetic risk underlying

Parkinson's disease. *Nat Genet* **41**, 1308–1312.

[8] Schiesling C, Kieper N, Seidel K, Krüger R (2008) Review: Familial Parkinson's

disease – genetics, clinical phenotype and neuropathology in relation to the

common sporadic form of the disease. *Neuropathol Appl Neurobiol* **34**, 255–271.

[9] Kalia L V, Lang AE (2015) Parkinson's disease. *Lancet (London, England)* **386**,

896–912.

[10] Azeggagh S, Berwick DC (2022) The development of inhibitors of leucine-rich

repeat kinase 2 (LRRK2) as a therapeutic strategy for Parkinson's disease: the

current state of play. *Br J Pharmacol* **179**, 1478–1495.

[11] Jennings D, Huntwork-Rodriguez S, Henry AG, Sasaki JC, Meisner R, Diaz D,

Solanoy H, Wang X, Negrou E, Bondar V V., Ghosh R, Maloney MT, Propson

NE, Zhu Y, Maciuca RD, Harris L, Kay A, LeWitt P, King TA, Kern D,

Ellenbogen A, Goodman I, Siderowf A, Aldred J, Omidvar O, Masoud ST, Davis

SS, Arguello A, Estrada AA, de Vicente J, Sweeney ZK, Astarita G, Borin MT,

Wong BK, Wong H, Nguyen H, Scearce-Levie K, Ho C, Troyer MD (2022)

Preclinical and clinical evaluation of the LRRK2 inhibitor DNL201 for

Parkinson's disease. *Sci Transl Med* **14**,.

[12]   Bartel DP (2004) MicroRNAs: genomics, biogenesis, mechanism, and function.
590   *Cell* **116**, 281–297.

591   [13]   Burgos K, Malenica I, Metpally R, Courtright A, Rakela B, Beach T, Shill H,
592   Adler C, Sabbagh M, Villa S, Tembe W, Craig D, Van Keuren-Jensen K (2014)
593   Profiles of Extracellular miRNA in Cerebrospinal Fluid and Serum from Patients
594   with Alzheimer's and Parkinson's Diseases Correlate with Disease Status and
595   Features of Pathology. *PLoS One* **9**, e94839.

596   [14]   Gui Y, Liu H, Zhang L, Lv W, Hu X (2015) Altered microRNA profiles in
597   cerebrospinal fluid exosome in Parkinson disease and Alzheimer disease.
598   *Oncotarget* **6**, 37043–37053.

599   [15]   Gehrke S, Imai Y, Sokol N, Lu B (2010) Pathogenic LRRK2 negatively regulates
600   microRNA-mediated translational repression. *Nature* **466**, 637–641.

601   [16]   Ramakers C, Ruijter JM, Deprez RHL, Moorman AF. (2003) Assumption-free
602   analysis of quantitative real-time polymerase chain reaction (PCR) data.
603   *Neurosci Lett* **339**, 62–66.

604   [17]   R Core Team (2022) A language and environment for statistical computing.

605   [18]   Livak KJ, Schmittgen TD (2001) Analysis of Relative Gene Expression Data
606   Using Real-Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods* **25**, 402–
607   408.

608   [19]   Kolde R (2019) pheatmap: Pretty Heatmaps.

609   [20]   Kuhn M (2008) Building Predictive Models in R Using the caret Package. *J Stat*
610   *Softw* **28**,.

611   [21]   Kohavi R (2001) A Study of Cross-Validation and Bootstrap for Accuracy
612   Estimation and Model Selection. **14**,.

[22]   Liaw A, Wiener M (2002) Classification and Regression by randomForest. *R*

News **2**, 18–22.

[23] Chen Y, Wang X (2020) miRDB: an online database for prediction of functional

microRNA targets. *Nucleic Acids Res* **48**, D127–D131.

[24] Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L,

Fu X, Liu S, Bo X, Yu G (2021) clusterProfiler 4.0: A universal enrichment tool

for interpreting omics data. *Innov* **2**, 100141.

[25] Wang R, Yang Y, Wang H, He Y, Li C (2020) MiR-29c protects against

inflammation and apoptosis in Parkinson's disease model in vivo and in vitro by

targeting SP1. *Clin Exp Pharmacol Physiol* **47**, 372–382.

[26] Botta-Orfila T, Morató X, Compta Y, Lozano JJ, Falgàs N, Valldeoriola F, Pont-

Sunyer C, Vilas D, Mengual L, Fernández M, Molinuevo JL, Antonell A, Martí

MJ, Fernández-Santiago R, Ezquerra M (2014) Identification of blood serum

micro-RNAs associated with idiopathic and LRRK2 Parkinson's disease. *J*

Neurosci Res **92**, 1071–1077.

[27] Bai X, Tang Y, Yu M, Wu L, Liu F, Ni J, Wang Z, Wang J, Fei J, Wang W,

Huang F, Wang J (2017) Downregulation of blood serum microRNA 29 family

in patients with Parkinson's disease. *Sci Rep* **7**, 5411.

[28] Ozdilek B, Demircan B (2021) Serum microRNA expression levels in Turkish

patients with Parkinson's disease. *Int J Neurosci* **131**, 1181–1189.

[29] Soto M, Fernández M, Bravo P, Lahoz S, Garrido A, Sánchez-Rodríguez A,

Rivera-Sánchez M, Sierra M, Melón P, Roig-García A, Naito A, Casey B, Camps

J, Tolosa E, Martí M-J, Infante J, Ezquerra M, Fernández-Santiago R (2023)

Differential serum microRNAs in premotor LRRK2 G2019S carriers from

Parkinson's disease. *npj Park Dis* **9**, 15.

[30] Briggs CE, Wang Y, Kong B, Woo T-UW, Iyer LK, Sonntag KC (2015)

Midbrain dopamine neurons in Parkinson′s disease exhibit a dysregulated miRNA and target-gene network. *Brain Res* **1618**, 111–121.

[31] Mancuso R, Agostini S, Hernis A, Zanzottera M, Bianchi A, Clerici M (2019) Circulatory miR-223-3p Discriminates Between Parkinson's and Alzheimer's Patients. *Sci Rep* **9**, 9393.

[32] Bauernfeind F, Rieger A, Schildberg FA, Knolle PA, Schmid-Burgk JL, Hornung V (2012) NLRP3 Inflammasome Activity Is Negatively Controlled by miR-223. *J Immunol* **189**, 4175–4181.

[33] Moehle MS, Webber PJ, Tse T, Sukar N, Standaert DG, DeSilva TM, Cowell RM, West AB (2012) LRRK2 Inhibition Attenuates Microglial Inflammatory Responses. *J Neurosci* **32**, 1602–1611.

[34] Thomas RR, Keeney PM, Bennett JP (2012) Impaired Complex-I Mitochondrial Biogenesis in Parkinson Disease Frontal Cortex. *J Parkinsons Dis* **2**, 67–76.

[35] Kanao T, Venderova K, Park DS, Unterman T, Lu B, Imai Y (2010) Activation of FoxO by LRRK2 induces expression of proapoptotic proteins and alters survival of postmitotic dopaminergic neuron in Drosophila. *Hum Mol Genet* **19**, 3747–3758.

[36] Dorval V, Mandemakers W, Jolivette F, Coudert L, Mazroui R, Strooper B, Hébert SS (2014) Gene and MicroRNA transcriptome analysis of Parkinson's related LRRK2 mouse models. *PLoS One* **9**, e85510.

[37] Heman-Ackah SM, Hallegger M, Rao MS, Wood MJA (2013) RISC in PD: the impact of microRNAs in Parkinson's disease cellular and molecular pathogenesis. *Front Mol Neurosci* **6**,.

[38] Wang J, Chen C, Zhang Y (2020) An investigation of microRNA-103 and microRNA-107 as potential blood-based biomarkers for disease risk and

progression of Alzheimer's disease. *J Clin Lab Anal* **34**,.

[39] Li N, Pan X, Zhang J, Ma A, Yang S, Ma J, Xie A (2017) Plasma levels of miR-
137 and miR-124 are associated with Parkinson's disease but not with
Parkinson's disease with depression. *Neurol Sci* **38**, 761–767.

[40] Alexandrov PN, Dua P, Hill JM, Bhattacharjee S, Zhao Y, Lukiw WJ (2012)
microRNA (miRNA) speciation in Alzheimer's disease (AD) cerebrospinal fluid
(CSF) and extracellular fluid (ECF). *Int J Biochem Mol Biol* **3**, 365–73.

[41] Zhou L, Yang L, Li Y, Mei R, Yu H, Gong Y, Du M, Wang F (2018) MicroRNA-
128 Protects Dopamine Neurons from Apoptosis and Upregulates the Expression
of Excitatory Amino Acid Transporter 4 in Parkinson's Disease by Binding to
AXIN1. *Cell Physiol Biochem* **51**, 2275–2289.

[42] Chen Q, Deng N, Lu K, Liao Q, Long X, Gou D, Bi F, Zhou J (2021) Elevated
plasma miR-133b and miR-221-3p as biomarkers for early Parkinson's disease.
Sci Rep **11**, 15268.

[43] Cardo LF, Coto E, Ribacoba R, Menéndez M, Moris G, Suárez E, Alvarez V
(2014) MiRNA Profile in the Substantia Nigra of Parkinson's Disease and
Healthy Subjects. *J Mol Neurosci* **54**, 830–836.

[44] Mo M, Xiao Y, Huang S, Cen L, Chen X, Zhang L, Luo Q, Li S, Yang X, Lin
X, Xu P (2017) MicroRNA expressing profiles in A53T mutant alpha-synuclein
transgenic mice and Parkinsonian. *Oncotarget* **8**, 15–28.

[45] Oliveira SR, Dionísio PA, Correia Guedes L, Gonçalves N, Coelho M, Rosa
MM, Amaral JD, Ferreira JJ, Rodrigues CMP (2020) Circulating Inflammatory
miRNAs Associated with Parkinson's Disease Pathophysiology. *Biomolecules*
10,.

[46] Ding H, Huang Z, Chen M, Wang C, Chen X, Chen J, Zhang J (2016)

Identification of a panel of five serum miRNAs as a biomarker for Parkinson's

disease. *Parkinsonism Relat Disord* **22**, 68–73.

[47] Tatura R, Kraus T, Giese A, Arzberger T, Buchholz M, Höglinger G, Müller U

(2016) Parkinson's disease: SNCA-, PARK2-, and LRRK2- targeting

microRNAs elevated in cingulate gyrus. *Parkinsonism Relat Disord* **33**, 115–

121.

[48] Nair VD, Ge Y (2016) Alterations of miRNAs reveal a dysregulated molecular

regulatory network in Parkinson's disease striatum. *Neurosci Lett* **629**, 99–104.

[49] Marques TM, Kuiperij HB, Bruinsma IB, van Rumund A, Aerts MB, Esselink

RAJ, Bloem BR, Verbeek MM (2017) MicroRNAs in Cerebrospinal Fluid as

Potential Biomarkers for Parkinson's Disease and Multiple System Atrophy. *Mol*

Neurobiol **54**, 7736–7745.

[50] Li L, Ren J, Pan C, Li Y, Xu J, Dong H, Chen Y, Liu W (2021) Serum miR-214

Serves as a Biomarker for Prodromal Parkinson's Disease. *Front Aging Neurosci*

13,.

[51] Khoo SK, Petillo D, Kang UJ, Resau JH, Berryhill B, Linder J, Forsgren L,

Neuman LA, Tan AC (2012) Plasma-Based Circulating MicroRNA Biomarkers

for Parkinson's Disease. *J Parkinsons Dis* **2**, 321–331.

[52] dos Santos MCT, Barreto-Sanz MA, Correia BRS, Bell R, Widnall C, Perez LT,

Berteau C, Schulte C, Scheller D, Berg D, Maetzler W, Galante PAF, Nogueira

da Costa A (2018) miRNA-based signatures in cerebrospinal fluid as potential

diagnostic tools for early stage Parkinson's disease. *Oncotarget* **9**, 17455–17465.

[53] Han K (2017) The diagnostic efficacy of serum miR-103a, miR-30b, miR-29a

relative expression levels for Parkinson's disease. *Shadong Med J* **57**, 72–74.

[54] Yılmaz ŞG, Geyik S, Neyal AM, Soko ND, Bozkurt H, Dandara C (2016)

Hypothesis: Do miRNAs Targeting the Leucine-Rich Repeat Kinase 2 Gene (

LRRK2) Influence Parkinson's Disease Susceptibility? *Omi A J Integr Biol* **20**,

224–228.

[55]   Yao Y-F, Qu M-W, Li G-C, Zhang F-B, Rui H-C (2018) Circulating exosomal

miRNAs as diagnostic biomarkers in Parkinson's disease. *Eur Rev Med*

Pharmacol Sci **22**, 5278–5283.

[56]   Miñones-Moyano E, Porta S, Escaramís G, Rabionet R, Iraola S, Kagerbauer B,

Espinosa-Parrilla Y, Ferrer I, Estivill X, Martí E (2011) MicroRNA profiling of

Parkinson's disease brains identifies early downregulation of miR-34b/c which

modulate mitochondrial function. *Hum Mol Genet* **20**, 3067–3078.

[57]   Chen Y, Gao C, Sun Q, Pan H, Huang P, Ding J, Chen S (2017) MicroRNA-

4639 Is a Regulator of DJ-1 Expression and a Potential Early Diagnostic Marker

for Parkinson's Disease. *Front Aging Neurosci* **9**,.

[58]   Lin X, Wang R, Li R, Tao T, Zhang D, Qi Y (2022) Diagnostic Performance of

miR-485-3p in Patients with Parkinson's Disease and its Relationship with

Neuroinflammation. *NeuroMolecular Med* **24**, 195–201.

[59]   Li J, Yin Y, Wang R, Zhang P, Zhao J, Liu R, Chen Y (2021) Expressions and

clinical significance of miR-124 and miR-494 in elderly patients with Parkinson

disease. *Chinese J Behav Med Brain Sci* **12**, 294–298.

[60]   Zhou Y, Lu M, Du R-H, Qiao C, Jiang C-Y, Zhang K-Z, Ding J-H, Hu G (2016)

MicroRNA-7 targets Nod-like receptor protein 3 inflammasome to modulate

neuroinflammation in the pathogenesis of Parkinson's disease. *Mol*

Neurodegener **11**, 28.

[61]   Wu L, Zhao W, Kong F, Yang F, Zheng J (2020) SERUM MIR-9A AND MIR-

133B, DIAGNOSTIC MARKERS FOR PARKINSON'S DISEASE, ARE

UPREGULATED AFTER LEVODOPA TREATMENT. *Acta Medica Mediterr*

36,.

Table 1: Overview of clinical data.

Disease status	LRRK2 mutated		Sporadic PD (n = 10)	Healthy Control (n = 11)
	PD (n = 6)	Asymptomatic (n = 4)		
Age (years)	66.8 ± 8.7	52.3 ± 19.6	64.5 ± 10.9	60.1 ± 14.5
Gender (m:f)	1:5	0:4	2:8	2:9
DD (years)	8.8 ± 3.6	-	7 ± 4.1	-
MoCA	25 ± 4.1 (5/6)	28.5 ± 1.5	25.3 ± 3.7 (9/10)	28.3 ± 1.5 (4/11)
UPDRS-III	21.7 ± 8.2	1.0 ± 0.8 (3/4)	17.4 ± 10.1 (8/10)	0.7 ± 0.9 (3/11)
LEDD (mg)	612.8 ± 295.6	0.0 ± 0.0	357.5 ± 380.0	0.0 ± 0.0

745 DD: Disease duration, MoCA: Montreal-Cognitive-Assessment, UPDRS-III: Unified

746 Parkinson's Disease Rating Scale, LEDD: L-Dopa equivalent daily dose. Age, Disease

747 duration, MoCA, UPDRS-III and LEDD stated as mean value with standard deviation.

748 MoCA and UPDRS-III were not accessible for some patients and numbers in brackets

749 give fraction of accessible data.

Table 2: miRNAs selected as most influential by the different models.

t-test selected	PCA selected	LASSO selected	Random forest selected
miR-29c-3p	miR-19b-3p	**miR-29c-3p**	miR-26a-5p
miR-128-3p	miR-27a-3p	**miR-424-5p**	miR-106a-5p
miR-148a-3p	miR-28-5p	miR-532-5p	**miR-128-3p**
miR-153-3p	miR-30b-5p		*miR-223-3p*
miR-425-5p	miR-185-5p		**miR-424-5p**

The table displays miRNAs that were identified as relevant from each model or test. miRNAs highlighted in bold were identified in more than one analysis. Of note, miR-223-3p (italic) was extracted from both the Ct_{plasma} and the $Ct_{multiplied}$ RF models.

Figures

Figure 1: Study Design Overview. The experimental cohort compromised 31 patients categorized into three groups: individuals carrying a *LRRK2* mutation, sporadic PD patients and healthy controls. A total of 91 miRNAs, selected by reviewing relevant literature, were quantified in CSF and plasma samples using reverse transcription and qPCR. Subsequently, the obtained raw Ct values were calibrated and fc and log2fc values were calculated. miRNAs that were expressed only in subgroups were excluded. Group differences were then analyzed using t-tests, PCA, Random Forest and LASSO regression. Created with BioRender.com.

Figure 2: Visualizing miRNA signatures and group differences. A: Heatmap based on log2fc values from the plasma dataset, normalized per row. Rows represent different miRNAs while columns represent patients. Grey cells represent missing values. Column sorting function was inactivated to not interfere with group arrangement. The similarity indicated by the column dendrograms is not proportional between groups. **B:** Heatmap based on log2fc values from CSF dataset, normalized per row. This heatmap includes a smaller amount of miRNAs (represented in rows) as they were excluded from the CSF dataset due to missing values. **C:** Histogram showing the distribution of uncorrected p-values obtained from the plasma and the **D:** CSF datasets after performing unpaired t-tests using log2fc values. **E:** Scatterplots display log2fc values from sPD patients and LRRK2$_{MC}$. Five miRNAs with the lowest p-value in the t-tests were selected. Thick line indicates mean while the box indicates the standard deviation.

Figure 3: PCA graphs and correlation of plasma and CSF data. A: PCA graph from plasma and **B:** CSF Ct values. Ellipses indicate 95% confidence interval. Numbers indicate *LRRK2* mutation (1: G2019S, 1*: G2019S + G1819, 2: R1441C, 3: I2020T). In the plasma PCA graph, group ellipses overlap while the blue LRRK2$_{MC}$ ellipse trends to only include individuals of the LRRK2$_{MC}$ group. Based on PCA, the LRRK2$_{MC}$ group could be interpreted as a subpopulation of all PD patients. **C:** Correlation matrix presenting statistically significant ($p < 0.05$) correlations between Ct$_{plasma}$ and Ct$_{CSF}$ values of the eleven miRNAs included in the CSF dataset.

Figure 4: Plasma PCA plots comparing all groups. A: sPD and HC. **B:** LRRK2$_{MC}$ and HC. **C:** LRRK2$_{MC}$ and sPD. **D:** All groups. Arrows point towards the respective mean of PC1 and PC2. In **A**, **B** and **C** clear separation can be observed as indicated my arrows pointing towards opposing directions. When including all groups in **D**, clustering is less apparent but still observable.

Figure 5: Group prediction with LASSO and Random Forest. A: Overview of performed analyses and respective read-outs. **B:** Sensitivity and specificity values acquired by LASSO regression and Random Forest, using Ct_{plasma} and $Ct_{multiplied}$. **C:** Heatmaps display the proximity scores calculated in RF for all sPD patients (red) and $LRRK2_{MC}$ (blue) using Ct_{plasma} or **D:** $Ct_{multiplied}$. Groups clearly separate along mutation status. **E:** Mean Decrease Gini scores were calculated after building the respective RF models for both the Ct_{plasma} and **F:** the $Ct_{multiplied}$ data set. Values were sorted in descending order and the top ten miRNAs are displayed. High decrease of the Gini coefficient translates to a high impact on the performance of the respective RF model. **G:** From each analysis, most influential or discriminatory miRNAs were extracted.

Chord diagram displays the overlap of miRNAs selected from the different analyses.

Numbers in brackets display set size and grey connecting lines indicate miRNA

overlaps.

Supplementary Material

Figure S1: Identification of an outlier via PCA. A: The PCA graph on the left
displays data points from all sPD patients and LRRK2$_{MC}$. One of the asymptomatic
LRRK2$_{MC}$ clearly behaved as an outlier and was therefore excluded from further
analysis. **B:** PCA was repeated with the outlier excluded.

Figure S2: Processing of Raw data. Overview of data processing and selection
process. Created with BioRender.com.

826 **Figure S3: Group-wise Expression Distribution of miRNAs. A:** The histograms

827 show the number of individuals any given miRNA (x-axis) was found to be expressed

828 in, either in plasma or **B:** CSF. Colors indicate group affiliation.

830
831 **Figure S4: Calculation of ΔΔCt values.** Overview of calculation of ΔΔCt values.

832 Created with BioRender.com.

834
835 **Figure S5: Heatmaps based on Ct values. A:** Heatmaps for Ct_{plasma} and **B:** Ct_{CSF},

836 normalized per row. Rows represent miRNAs, columns represent individuals. Grey

837 cells represent missing values.

Figure S6: ROC curve for miR-29c-3p. ROC curve displaying the sensitivity and
specificity of a classification in LRRK2$_{MC}$ and sPD based on plasma log2fc values of
miR-29c-3p.

844
845

Figure S7: PCA graphs including lines for evaluation of discrimination based on
sensitivity and specificity. A: Ct_{plasma}. PC1 (solid line) achieves a sensitivity of 100%
(9/9) and a specificity of 70% (7/10). PC1 and PC2 combined (dotted line) reach the
same sensitivity with a specificity of 80% (8/10). B: $Ct_{multiplied}$. PC1 achieves separation
with sensitivity of 100% (9/9) and specificity of 80% (8/10).

Figure S8: Heatmap and PCA resulting from multiplied variables. A: $Ct_{multiplied}$ are

plotted as a heatmap for all groups. Again, rows display miRNAs, while columns

display individuals. **B:** When performing PCA with $Ct_{multiplied}$, group separation was

observable.

Figure S9: Scatterplots of log2fc values for the four discussed miRNAs. Thick line indicates

mean while the box indicates the standard deviation.

Figure S10: GO analysis of identified miRNAs. A: miR-223-3p. 25 significant annotations were

identified and sorted by adjusted p-value. Top 15 annotations are displayed. **B**: miR-29c-3p. 60

significant annotations were identified, top 15 annotations are displayed. **C**: miR-128-3p. There

were no annotations with an adjusted p-value < 0.05. 6 annotations with an adjusted p-value < 0.01

are displayed. **D**: miR-424-5p. 3 Annotations with an adjusted p-value < 0.05 were identified.

Table S1: included miRNAs with literature references.

miRNA	Context	Reference
miR-16-5p	Striatal tissue of LRRK2 KO mice	[36]
miR-1-3p	Exosomal miRNAs in CSF of PD patients	[14]
miR-103a-3p	Striatal tissue of LRRK2 KO mice	[36]
miR-106a-5p	Brain tissue of PD patients	[30]
miR-107	Algorithm based predicted interaction with LRRK2	[37]
	Plasma of AD and PD patients	[38]
miR-10a-5p	Exosomal miRNAs in CSF of PD patients	[14]
miR-122-5p	Striatal tissue of LRRK2 KO mice	[36]
miR-124-3p	Plasma of PD patients	[39]
miR-125b-5p	CSF of AD patients	[40]
miR-128-3p	Brain tissue of mouse models of PD	[41]
miR-129-5p	Algorithm based predicted interaction with LRRK2	[37]
miR-132-3p	Brain tissue of PD patients	[30]
miR-133b	Plasma of PD patients	[42]
miR-135a-5p	Brain tissue of PD patients	[30]
miR-135b-5p	Brain tissue of PD patients	[43]
miR-136-5p	Algorithm based predicted interaction with LRRK2	[37]
miR-137-3p	Plasma of PD patients	[39]
miR-141-5p	Algorithm based predicted interaction with LRRK2	[37]
miR-144-5p	CSF of PD patients	[44]
	Algorithm based predicted interaction with LRRK2	[37]
miR-145-5p	Brain tissue of PD patients	[30]
miR-146a-5p	Serum of LRRK2 PD patients	[45]
miR-148a-3p	Brain tissue of PD patients	[30]
miR-153-3p	Exosomal miRNAs in CSF of PD patients	[14]
miR-155-5p	Serum of LRRK2 PD patients	[45]
miR-15b-5p	Serum of PD patients	[46]
miR-181a-5p	Serum of PD patients	[46]
	Algorithm based predicted interaction with LRRK2	[37]
miR-185-5p	Serum of PD patients	[46]
	Algorithm based predicted interaction with LRRK2	[37]
miR-190a-5p	Brain tissue of PD patients	[34]
miR-195-5p	Serum of PD patients	[46]
miR-198	Brain tissue of PD patients	[43]
miR-199a-3p	Brain tissue of PD patients	[34]
miR-199b-5p	Brain tissue of PD patients	[47]
miR-19a-5p	Serum of LRRK2 PD patients	[26]
	Algorithm based predicted interaction with LRRK2	[37]
miR-19b-3p	Serum of PD patients	[26]
	Algorithm based predicted interaction with LRRK2	[37]
	Exosomal miRNAs in CSF of PD patients	[14]
miR-200a-3p	CSF of PD patients	[44]

	Algorithm based predicted interaction with LRRK2	[37]
miR-200b-3p	Algorithm based predicted interaction with LRRK2	[37]
miR-204-5p	Brain tissue of PD patients	[48]
miR-205-5p	CSF of PD patients	[49]
	Algorithm based predicted interaction with LRRK2	[37]
miR-214-5p	Serum of PD patients	[50]
miR-219a-2-3p	Brain tissue of PD patients	[48]
miR-221-3p	Plasma of PD patients	[42]
miR-222-3p	Plasma of PD patients	[51]
miR-223-3p	Brain tissue of PD patients	[30]
miR-24-3p	CSF of PD patients	[49]
miR-25-3p	Algorithm based predicted interaction with LRRK2	[37]
miR-26a-5p	Brain tissue of PD patients	[30]
miR-27a-3p	CSF of PD patients	[52]
miR-28-5p	Brain tissue of PD patients	[30]
miR-29a-3p	Serum of LRRK2 PD patients	[26]
miR-29c-3p	Serum of LRRK2 PD patients	[26]
miR-301b-5p	Algorithm based predicted interaction with LRRK2	[37]
miR-30b-5p	Serum of PD patients	[53]
miR-31-5p	Serum of PD advanced PD patients	[50]
miR-3143	Whole blood of PD patients	[54]
miR-3200-3p	Brain tissue of PD patients	[48]
miR-328-5p	Algorithm based predicted interaction with LRRK2	[37]
miR-331-5p	Exosomal miRNAs in Plasma of PD patients	[55]
miR-335-3p	Serum of LRRK2 PD patients	[45]
miR-335-5p	Serum of LRRK2 PD patients	[45]
miR-34b-5p	Brain tissue of PD patients	[56]
miR-34c-5p	Brain tissue of PD patients	[56]
miR-374a-5p	Brain tissue of PD patients	[30]
miR-382-5p	Brain tissue of PD patients	[48]
miR-384	Algorithm based predicted interaction with LRRK2	[37]
miR-409-3p	Exosomal miRNAs in CSF of PD patients	[14]
miR-410-5p	Algorithm based predicted interaction with LRRK2	[37]
miR-421	Brain tissue of PD patients	[48]
miR-423-5p	CSF of PD patients	[52]
miR-424-5p	Brain tissue of PD patients	[34]
miR-425-5p	Brain tissue of PD patients	[48]
miR-429	Brain tissue of PD patients	[34]
	Algorithm based predicted interaction with LRRK2	[37]
miR-4421	Brain tissue of PD patients	[48]
miR-451a	Brain tissue of PD patients	[34]
miR-454-5p	Algorithm based predicted interaction with LRRK2	[37]
miR-4639-5p	Plasma of PD patients	[57]
miR-4671-3p	Whole blood of PD patients	[54]

miR-485-5p	Serum of PD patients	[58]
miR-488-3p	Brain tissue of PD patients	[47]
miR-494-5p	Serum of PD patients	[59]
miR-505-5p	Exosomal miRNAs in Plasma of PD patients	[55]
miR-532-5p	Brain tissue of PD patients	[30]
miR-542-3p	CSF of PD patients	[44]
miR-544a	Brain tissue of PD patients	[47]
miR-561-3p	Whole blood of PD patients	[54]
miR-579-3p	Whole blood of PD patients	[54]
miR-626	Plasma of PD patients	[51]
miR-7-5p	Serum of PD patients	[60]
miR-744-5p	Brain tissue of PD patients	[30]
miR-92a-3p	Brain tissue of PD patients	[30]
miR-95-3p	Brain tissue of PD patients	[48]
miR-9-5p	Serum of PD patients	[61]

869   AD = Alzheimer's disease, PD = Parkinson's disease

Table S2: Melting points of miRNAs detected in plasma. Only miRNAs which were selected for further analysis steps are displayed. Low standard deviations indicate constant amplification of the target.

Target	Mean Melting Point [°C]	Standard deviation [°C]
hsa-miR-103a-3p	69.71	0.31
hsa-miR-106a-5p	71.71	0.34
hsa-miR-107	69.73	0.34
hsa-miR-10a-5p	71.77	0.66
hsa-miR-122-5p	70.59	0.31
hsa-miR-125b-5p	71.37	0.38
hsa-miR-128-3p	69.75	0.11
hsa-miR-132-3p	71.20	0.16
hsa-miR-133b	71.54	0.21
hsa-miR-135a-5p	70.62	0.13
hsa-miR-135b-5p	69.36	0.32
hsa-miR-136-5p	70.30	0.13
hsa-miR-1-3p	70.65	0.13
hsa-miR-144-5p	71.39	0.18
hsa-miR-145-5p	69.09	0.32
hsa-miR-146a-5p	70.36	0.17
hsa-miR-148a-3p	69.78	0.27
hsa-miR-153-3p	69.92	0.22
hsa-miR-155-5p	69.63	0.19
hsa-miR-15b-5p	69.34	0.25
hsa-miR-16-5p	70.60	0.18
hsa-miR-181a-5p	70.26	0.17
hsa-miR-185-5p	70.65	0.16
hsa-miR-190a-5p	69.29	0.26
hsa-miR-195-5p	69.31	0.28
hsa-miR-199a-3p	69.94	0.13
hsa-miR-19a-5p	70.44	0.15
hsa-miR-19b-3p	69.88	0.15
hsa-miR-200b-3p	71.37	0.13
hsa-miR-204-5p	69.22	0.15
hsa-miR-221-3p	69.93	0.51
hsa-miR-222-3p	69.91	0.40
hsa-miR-223-3p	70.84	0.20
hsa-miR-24-3p	74.47	0.18
hsa-miR-25-3p	69.82	0.26
hsa-miR-26a-5p	70.88	0.12
hsa-miR-27a-3p	70.86	0.20
hsa-miR-28-5p	70.30	0.15
hsa-miR-29a-3p	70.19	0.23

hsa-miR-29c-3p	70.40	0.13
hsa-miR-30b-5p	69.20	0.10
hsa-miR-31-5p	69.22	1.82
hsa-miR-335-3p	70.89	0.13
hsa-miR-335-5p	71.43	0.25
hsa-miR-374a-5p	70.21	0.16
hsa-miR-382-5p	71.78	0.26
hsa-miR-409-3p	70.32	0.26
hsa-miR-421	73.12	0.29
hsa-miR-423-5p	70.77	0.29
hsa-miR-424-5p	69.30	0.12
hsa-miR-425-5p	70.09	0.36
hsa-miR-451a	70.11	0.27
hsa-miR-505-5p	70.23	0.23
hsa-miR-532-5p	69.43	0.45
hsa-miR-542-3p	70.47	0.13
hsa-miR-744-5p	73.34	0.43
hsa-miR-7-5p	70.12	0.36
hsa-miR-95-3p	69.48	0.38
hsa-miR-9-5p	73.70	0.49

875 **Table S3: Melting points of miRNAs detected in CSF.** Only miRNAs which were selected for
876 further analysis steps are displayed. Low standard deviations indicate constant amplification of the
877 target.
878

Target	Mean Melting Point [°C]	Standard deviation [°C]
hsa-miR-16-5p	70.84	0.47
hsa-miR-125b-5p	71.32	0.43
hsa-miR-145-5p	69.12	0.33
hsa-miR-19b-3p	69.75	0.27
hsa-miR-204-5p	69.25	0.15
hsa-miR-223-3p	70.66	0.38
hsa-miR-24-3p	74.19	1.05
hsa-miR-26a-5p	71.05	0.84
hsa-miR-27a-3p	70.83	0.29
hsa-miR-29c-3p	70.55	0.25
hsa-miR-424-5p	69.42	0.22

880   **Table S4: Statistic details from plasma t-tests.** Alpha was corrected to 0.0009 by dividing by the

881   number of tests.

t-statistic	p-value	miRNA
-1.74	0.1019	hsa-miR-1-3p
-2.10	0.0524	hsa-miR-103a-3p
-2.72	0.0146	hsa-miR-106a-5p
-2.50	0.0239	hsa-miR-107
-0.25	0.8073	hsa-miR-10a-5p
2.62	0.0218	hsa-miR-122-5p
0.72	0.4803	hsa-miR-125b-5p
-3.06	0.0071	hsa-miR-128-3p
-1.98	0.0658	hsa-miR-132-3p
-1.39	0.1841	hsa-miR-133b
-2.74	0.0161	hsa-miR-135a-5p
-1.25	0.2296	hsa-miR-135b-5p
-1.89	0.0764	hsa-miR-136-5p
1.37	0.1998	hsa-miR-144-5p
-1.56	0.1384	hsa-miR-145-5p
-1.90	0.0783	hsa-miR-146a-5p
-3.31	0.0042	hsa-miR-148a-3p
-3.58	0.0035	hsa-miR-153-3p
-2.02	0.0597	hsa-miR-155-5p
-1.57	0.1357	hsa-miR-15b-5p
0.08	0.9378	hsa-miR-16-5p
-1.84	0.0851	hsa-miR-181a-5p
-3.00	0.0081	hsa-miR-185-5p
-2.23	0.0431	hsa-miR-190a-5p
-2.60	0.0202	hsa-miR-195-5p
-2.04	0.0588	hsa-miR-199a-3p
-1.02	0.3222	hsa-miR-19a-5p
-1.96	0.0734	hsa-miR-19b-3p
-1.19	0.2502	hsa-miR-200b-3p
-0.94	0.3590	hsa-miR-204-5p
-1.37	0.1908	hsa-miR-221-3p
-2.06	0.0574	hsa-miR-222-3p
-2.10	0.0514	hsa-miR-223-3p
-1.15	0.2666	hsa-miR-24-3p
-0.82	0.4257	hsa-miR-25-3p
-2.02	0.0658	hsa-miR-26a-5p
-1.86	0.0822	hsa-miR-27a-3p
-2.33	0.0358	hsa-miR-28-5p
-1.73	0.1064	hsa-miR-29a-3p
-4.21	**0.0007***	hsa-miR-29c-3p

-2.07	0.0535	hsa-miR-30b-5p
-1.55	0.1505	hsa-miR-31-5p
-1.51	0.1489	hsa-miR-335-3p
-2.24	0.0397	hsa-miR-335-5p
-2.61	0.0186	hsa-miR-374a-5p
-1.21	0.2421	hsa-miR-382-5p
-1.27	0.2233	hsa-miR-409-3p
-1.31	0.2068	hsa-miR-421
-2.35	0.0360	hsa-miR-424-5p
-4.01	0.0010	hsa-miR-425-5p
-0.62	0.5441	hsa-miR-451a
-1.36	0.1945	hsa-miR-505-5p
-2.08	0.0528	hsa-miR-532-5p
0.25	0.8022	hsa-miR-542-3p
-1.26	0.2261	hsa-miR-7-5p
-1.10	0.2890	hsa-miR-744-5p
-2.75	0.0148	hsa-miR-9-5p
1.68	0.1106	hsa-miR-95-3p

883   **Table S5: statistics of correlation matrix.** Correlations calculated after Pearson. Only significant

884   miRNA combinations ($p < 0.05$) are displayed.

miRNA Plasma	miRNA CSF	R	R^2	p-value
miR-16-5p	miR-125-5p	0.39	0.15	0.037
miR-16-5p	miR-145-5p	0.39	0.15	0.042
miR-16-5p	miR-19b-3p	0.40	0.16	0.034
miR-16-5p	miR-223-3p	0.43	0.19	0.021
miR-16-5p	miR-27a-3p	0.46	0.21	0.013
miR-19b-3p	miR-125-5p	0.38	0.15	0.041
miR-19b-3p	miR-19b-3p	0.44	0.20	0.018
miR-19b-3p	miR-223-3p	0.48	0.23	0.010
miR-19b-3p	miR-27a-3p	0.45	0.20	0.014
miR-204-5p	miR-125b-5p	0.37	0.14	0.043
miR-204-5p	miR-19b-3p	0.44	0.19	0.018
miR-204-5p	miR-223-3p	0.40	0.16	0.032
miR-204-5p	miR-27a-3p	0.48	0.23	0.007
miR-204-5p	miR-29c-3p	0.41	0.17	0.023
miR-223-3p	miR-223-3p	0.51	0.26	0.005
miR-26a-5p	miR-223-3p	0.49	0.24	0.007
miR-26a-5p	miR-27a-3p	0.36	0.13	0.032
miR-27a-3p	miR-223-3p	0.50	0.25	0.006
miR-27a-3p	miR-27a-3p	0.39	0.15	0.032
miR-29c-3p	miR-27a-3p	0.38	0.14	0.043

886 **Table S6: t-test with Ct$_{multiplied}$ data.** p-value threshold was corrected for multiple testing by
887 dividing by the number of tests and set to 0.0025.

miRNA Plasma	miRNA CSF	t-statistic	p-value
miR-16-5p	miR-125-5p	0.8	0.4483
miR-16-5p	miR-145-5p	1.1	0.3086
miR-16-5p	miR-19b-3p	2.0	0.0599
miR-16-5p	miR-223-3p	2.1	0.0500
miR-16-5p	miR-27a-3p	1.4	0.1898
miR-19b-3p	miR-125-5p	1.5	0.1602
miR-19b-3p	miR-19b-3p	2.8	0.0132
miR-19b-3p	miR-223-3p	2.8	0.0119
miR-19b-3p	miR-27a-3p	2.1	0.0544
miR-204-5p	miR-125b-5p	1.0	0.3343
miR-204-5p	miR-19b-3p	2.1	0.0517
miR-204-5p	miR-223-3p	2.3	0.0324
miR-204-5p	miR-27a-3p	1.5	0.1457
miR-204-5p	miR-29c-3p	1.3	0.2131
miR-223-3p	miR-223-3p	2.9	0.0113
miR-26a-5p	miR-223-3p	2.6	0.0223
miR-26a-5p	miR-27a-3p	2.2	0.0467
miR-27a-3p	miR-223-3p	2.6	0.0204
miR-27a-3p	miR-27a-3p	2.1	0.0532
miR-29c-3p	miR-27a-3p	4.2	**0.0007***