Conservation of molecular responses upon viral infection in the non-vascular plant *Marchantia polymorpha*

Eric Moner-Ros[1*], Tamara Jiménez-Góngora[1*], Luis Villar-Martín[1], Víctor M. González-Miguel[2], Denis Kutnjak[3] and Ignacio Rubio-Somoza[1,4,]

[1]Molecular Reprogramming and Evolution (MoRE) Laboratory, Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus UAB-Edifci CRAG. 08193 Cerdanyola del Vallés, Spain.

[2]Data Analysis area, Bioinformatics Core Unit, Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus UAB-Edifci CRAG. 08193 Cerdanyola del Vallés, Spain.

[3]National Institute of Biology, Department of Biotechnology and Systems Biology, Ljubljana, Slovenia.

[4]Consejo Superior de Investigaciones Científicas (CSIC), Barcelona 08001, Spain.

*Those authors contributed equally to this work

To whom correspondence should be addressed: Ignacio.rubio@cragenomica.es

Abstract

After their transition from water to land around 450 million years ago, plants colonized new habitats facing unprecedented pathogenic microbes. That expansion was mostly supported by a growing anatomical complexity based on the acquisition of developmental innovations, such as roots, stomata and vascular tissue. Despite several of those innovations became central for the interaction between plants and their associated microbes, little is known about their impact on plant immune programs and on the diversification of infection strategies of their pathogens. A paradigmatic case is the close relationship between plant vasculature and viruses. Vascular tissues provide a unique cellular environment for viral replication and existence, besides constituting a fast track for viral systemic spread throughout the plant. Since most of our knowledge about plant-virus interactions come from studies in vascular plants, we here present a comparative study to contribute to the understanding of the evolution of plant-virus interactions by molecularly characterizing the interplay between the bryophyte Marchantia polymorpha and viruses. Virome analysis of Marchantia plants shows that they are primarily associated with RNA viruses in natural settings. Additional molecular characterization of the interaction between Marchantia and tobacco mosaic virus (TMV) show conserved basic

processes with vascular plants and divergent features. Viral infection triggers an extensive

transcriptional reprogramming in Marchantia encompassing broad range defence responses,

inhibition of cell cycle and photosynthesis and a sustained wound response that prevents further

viral movement. Additionally, infected plants show premature aging and organ maturation.

Notably, we found that some core responses that occur in infected areas in Marchantia, were

described to be restricted to vascular tissues in Nicotiana, suggesting that evolutionary

appearance of developmental innovations that became central in plants-virus interactions

resulted in re-routing of defence responses. Finally, we uncover the conserved role of a

transcription factor interacting with the TMV silencing suppressor p126 in specifically abrogating

TMV infection.

Introduction

It is estimated that plants developed a functional vascular system through repurposing of

preexisting genes around 420 million years ago, which facilitated the growth of larger plants by

providing mechanical support and enabling long distant transport of water, nutrients, and

signalling molecules (Bowles et al., 2022, Lucas et al., 2013). Viruses are obligate parasites that

can infect virtually any living organism hijacking central host cellular functions to replicate and

spread within and intra hosts. Coincidentally with land colonization, the repertoire of viruses

(virome) associated to plants enlarged and diversified both in nature and infection strategies.

Although our current knowledge lacks information about virome composition at intermediately

divergent plant lineages, it appears that viruses with large DNA genomes majorly infecting

unicellular chlorophyte algae were displaced by an increasing population of viruses with RNA

and small DNA genomes found in vascular plants (Mushegian et al., 2016). Plant vasculature has

become central for the success of viral infections in tracheophytes, not only serving as fast track

for viral systemic spread, but also providing a unique cellular environment for some viruses

which replication is restricted to that tissue (Kappagantu et al., 2020). Accordingly, vascular cells

are thought to present specific responses to viral infection when compared to other leaf cell-

types (Collum and Culver, 2017).

Our current understanding of the molecular events governing plant-virus interactions is vastly

shaped by studies focusing on vascular plants, mainly crops. Plants respond to viral infections via

at least three different but interconnected defence mechanisms (Mandadi and Scholthof, 2013).

Firstly, plants have an extracellular surveillance system that monitors the presence of pathogens

through the recognition of highly conserved microbe patterns dubbed as Pattern-associated

molecular patterns (PAMPs), such as viral derived double-strand RNAs (dsRNAS), the bacterial

flagellin or the fungal chitin. Detection of those molecules by specialized surface-localized

receptor kinases (Pattern recognition receptors, PRRs), triggers a first layer of defence called

PAMP-triggered immunity (PTI). PRRs expression is often restricted to focal points for microbial

interaction and invasion (Beck et al., 2014, Yotsui et al., 2023). Two additional surveillance

mechanisms monitor the intracellular presence of microbe-derived molecules. Highly stable

dsRNAs produced by viruses during their life cycle, independently of the DNA or RNA nature of

their genomes, are detected by the RNA silencing machinery which constitutes a conserved

antiviral defence mechanism both in plants and animals (Ding et al., 2018, Ding and Voinnet,

2007). Additionally, RNA silencing machinery also recognizes dsRNA generated from the

conversion of exogenous and endogenous single strand RNAs (ssRNA) by RNA-dependent RNA

polymerases (RDRs). RDR1 and RDR6 have been involved in antiviral defence in *Arabidopsis*

thaliana and *Nicotiana tabacum* (Garcia-Ruiz et al., 2010, Rakhshandehroo et al., 2017, Ying et

al., 2010), even though *Nicotiana benthamiana* has only a functional RDR6 (Bally et al., 2015).

Upon recognition, viral dsRNAs are trimmed in small RNAs (visRNAs) of 20 to 24 nucleotides in

length by members of the DICER-like family, mainly DCL2, DCL3 and DCL4 in Arabidopsis and

Nicotiana (Garcia-Ruiz et al., 2010, Lewsey and Carr, 2009). The resulting visRNA duplexes are

subsequently loaded into proteins from the ARGONAUTE family, which are the core elements of

the RNA-induced silencing complex (RiSC), and used to scan the intracellular space for the source

of dsRNA precursors in a sequence complementary manner abrogating viral infection (Lopez-

Gomollon and Baulcombe, 2022). In turn, viruses produce countermeasures, known as silencing

suppressors, targeted to interfere with different steps of the host RNA silencing machinery

(Lopez-Gomollon and Baulcombe, 2022). In response to these silencing suppressors, host micro

RNAs (miRNA, one of the two main classes of endogenous sRNAs in plants) are no longer

functional, allowing the translation of their targets and licensing a new layer of defence. Among

the miRNA targets involved in such counter-counter measure, members of the Nucleotide-

binding domain leucine-rich repeat (NLR) family are central. Thus, it has been suggested that

host miRNAs might work as sensors for the presence of pathogen-derived silencing suppressors

enabling a rapid host defence response based on the activation of otherwise silenced miRNA

targets (Vasseur et al., 2022). NLRs are core elements of the third intracellular surveillance

system for directly monitoring the intracellular presence of pathogen-derived molecules, such

as the prototypical case of the tobacco mosaic virus (TMV) replicase by the gene *N* in tobacco

(Erickson et al., 1999), or their impact on host cellular homeostasis. NLR-activation triggers the

so-called Effector-triggered immunity (ETI), which normally triggers a hypersensitive response

(HR) culminating in cell death. Those three defence systems, PTI, RNA silencing and ETI largely

rely on the action of the plant hormone Salicylic acid (SA). PTI and ETI responses encompass

activation of SA signalling (Pruitt et al., 2021) while SA activates RDR1 in Arabidopsis and

Nicotiana (Lee et al., 2016, Ying et al., 2010, Yu et al., 2003). Additionally, RDR1 expression is also

induced by treatment with jasmonates (Liu et al., 2009, Xu et al., 2013), another central hormone

in plant defence.

A Central question in plant-microbe interactions is how the evolution of different plant

developmental trajectories, including the acquisition of major innovations, impacts host defence

and pathogen infection strategies. In animals, comparative studies have proved very informative

about the origin and diversification of their innate immune responses against virus (Iwama and

Moran, 2023). Likewise, comparative studies including non-vascular plants, like the liverwort

Marchantia polymorpha, have been instrumental in understanding the evolution of the

interactions between plants and pathogens other than viruses (Carella et al., 2018, Gimenez-

Ibanez et al., 2019, Redkar et al., 2022). Incorporating bryophytes into those comparative studies

not only provides important information about how plant immune programs may have evolved,

but it also bears the potential to determine the contribution of developmental innovations, such

as roots, stomata or vasculature, to their diversification.

To enable comparative studies towards understanding the evolution of plant-virus interactions,

we here present the characterization of the molecular interplay between the non-vascular

liverwort *Marchantia polymorpha* and viruses. We firstly defined the composition of the virome

of Marchantia plants living in the wild and used that information to select a virus with similar

genomic features to those found in nature for further molecular characterization. Additionally,

the selected virus should be able to infect a broad range of hosts. Following those criteria, we

focused on the first ever isolated virus, the tobacco mosaic virus (TMV). TMV is a ssRNA virus

from the Tobamovirus genus that has been extensively used for studying plant-virus interactions

in a wealth of plant species, including several Solanaceae species, like tomato and tobacco, and

the model plant *Arabidopsis thaliana*, allowing for comparative studies. TMV infects through

wound entry points from where it moves cell-to-cell via plasmodesmata, until it reaches the

vascular system for systemic infection of distal parts of the plant following the stream of
photoassimilates (Liu and Nelson, 2013).

Our results show that *Marchantia polymorpha* is associated with RNA viruses in natural
populations that can infect lab grown Marchantia plants. Additionally, TMV can replicate in
Marchantia plants triggering a conserved transcriptional reprogramming encompassing
increased wounding and PTI responses in detriment of photosynthesis and cell cycle progression
and leading to aging and senescence. Induction of a wealth of NLR proteins and core elements
from the RNA silencing machinery were also observed. Noteworthy, the general response of the
RNA silencing machinery observed in Marchantia upon TMV infection is equivalent to that found
specifically in vascular tissues of *Nicotiana benthamiana*. We additionally describe a conserved
host interactor with the TMV silencing suppressor p126 that triggers cross-species defence
against TMV.

Results

***Marchantia polymorpha* is infected by RNA viruses in nature.**

To identify viruses infecting *Marchantia polymorpha* in nature, we performed a virome
analysis of a composite plant sample. Marchantia individuals were obtained from three open air
botanical gardens throughout Europe: Tübingen (Germany), Bern (Switzerland) and Cambridge
(UK). In addition to the expected background of bacteriophage sequences, we detected several
known viruses, including some capable of infecting or being associated with plants, fungi or
invertebrates (Supplementary table 1). Putatively new virus-like sequences were identified in the
dataset, based on i) their relatively high average sequencing depths and ii) similarity to viruses
previously associated with plants (Supplementary Table 2). Two previously described and three
newly identified viruses were hence selected for further RT-qPCR confirmation of their
association with the different *M. polymorpha* individuals, along with plants from the Tak-1
reference accession grown in sterile conditions. Specific primers to detect Ligustrum necrotic
ringspot virus (LNRV, known to infect plants), Hubei picorna-like virus 51, (detected in a
metagenomics study and putatively associated with dragonfly) a newly identified picorna-like
virus (which we named Tübingen picorna-like virus 1, with average mapping depth of 2230x and
contig length of 7801 nts), a new tombus-like virus (Tübingen tombus-like virus 1, with average
mapping depth of 218x and contig length of 4521 nts), and a new sobemo-like virus (Bern
sobemo-like virus 1, with average mapping depth of 88x and contig length of 2569 nts) were

used for these analysis. While none of those viruses were detected in negative control Tak-1

plants, individuals from Tübingen were associated with all tested viruses but the Bern sobemo-

like virus 1, which in turn was present in individuals from Bern along with Tübingen picorna-like

virus 1. Individuals from the Cambridge population were associated with Hubei picorna-like virus

51 and Tübingen picorna-like virus 1 (Supplementary Table 3).

Given that Tübingen picorna-like virus 1 was present in all tested populations, we decided to

further characterize its genomic features and interaction with *Marchantia polymorpha*. Indeed,

Tübingen picorna-like virus 1 had the highest mapping depth of all contigs in our virome analysis

and showed a similar genome structure as other unclassified picorna-like viruses (Shi et al.,

2016), consisting of two predicted genome open reading frames (ORFs) containing a RNA-

dependent RNA polymerase (RdRp) and a coat protein motifs (Figure 1A). Phylogenetic analysis

(Figure 1B) of the RdRp protein sequence and its closest NCBI GenBank homologues showed

high similarity (80% identity of aligned sequences) of this virus isolate with a recently reported

plant associated picorna-like virus (Trichosanthes kirilowii picorna-like virus, QKK82970.1). Much

lower identities were observed for Tübingen picorna-like virus 1 (29-42%) and other viruses

associated with diverse sources (plants, gastropods, water, crayfish, artropods and bird feces).

However, given that they were discovered in metagenomics studies their association with these

hosts is only speculative.

Next, we tested whether the Tübingen picorna-like virus 1 was able to infect *M. polymorpha* by

incubating two-weeks old Tak-1 individuals with extracts isolated from the Tübingen population,

which contained the highest levels of Tübingen picorna-like virus 1, as determined by RT-qPCR

analysis, or from Tak-1 plants grown in sterile conditions. Individual Marchantia plants were

collected one and two weeks after incubation and the levels of viral accumulation were

measured by RT-qPCR. Individuals exposed to extracts from infected plants showed a trend of

increased viral levels over time, while no amplification was observed in plants incubated with

extracts from Tak-1 plants (Figure 1C). This indicated that Tübingen picorna-like virus 1 can

replicate in Marchantia.

Collectively, those results show that *Marchantia polymorpha* can be infected by RNA viruses in

nature.

Tobacco mosaic virus infects *Marchantia polymorpha*

Since its identification in the 19th century, tobacco mosaic virus has been extensively used for

studying plant-virus interactions in several vascular plant species like Arabidopsis and Nicotiana

(Scholthof, 2004). As a first step for comparative studies on plant-virus interactions between

vascular and non-vascular plants, we aimed to infect *Marchantia polymorpha* plants with a GFP

labelled clone of TMV (Peart et al., 2002). Sap from leaves of healthy and infected *Nicotiana*

benthamiana plants were isolated and used to incubate two-weeks old Tak-1 plants grown in

sterile conditions. After incubation, Marchantia plants were transferred to soil and part of the

remaining sap was inoculated mechanically with an abrasive in basal leaves from 1 month old

Nicotiana plants, as control for its infectivity. After two weeks, Nicotiana plants showed systemic

GFP signal under UV light along with symptoms of infection (Supplementary Figure 1), whereas

Tak-1 plants displayed GFP signal in discrete infection foci that increased their size over time

(Figure 2A), suggestive of viral replication. This was further corroborated by RT-qPCR

amplification of pooled samples of four GFP positive individuals at the first (T1) and second

weeks after incubation (T2), in comparison with mock and freshly incubated (T0) plants (Figure

2B). Histological sections of the infected area showed presence of the virus in all cell types, as

inferred from GFP fluorescence, suggesting no preference for a specific cell-type during viral

infection (Figure 2C).

Those results confirmed that TMV can infect *M. polymoprha* plants pointing to Marchantia-TMV

as a suitable pathosystem for comparative studies.

TMV infection triggers profound molecular reprogramming in *M. polymoprha*

To characterize the molecular reprogramming caused by TMV infection, we performed

transcriptomic assays. TMV-GFP infected tissue was collected in four pools including 5 plants

each, along with the very same area from mock treated plants as control (Supplementary Figure

2A). RNA was isolated and the presence of TMV was confirmed by RT-qPCR in each pool

(Supplementary Figure 2B). Principal component analysis (PCA, Supplementary Figure 2C) of the

transcriptome supported the RT-qPCR results showing a clear separation of infected and mock

samples. Since separation among infected samples correlated with viral RT-qPCR levels, we

selected the three replicates that had the highest levels of TMV and clustered together for

further analysis. A total of 1879 genes were found to be up regulated while 1645 were reduced

(padjust<0.05, Log2(fold-change) =1) in response to TMV infection (Supplementary Table 4).

Consistently with an ongoing pathogenic interaction, we found several defence related marker

genes among the differentially expressed genes (DEG). Members of the pathogenesis-related

(PR) family are prime markers of immune responses in several plant species, including

Marchantia (Carella et al., 2019, Gimenez-Ibanez et al., 2019) and we found that the expression

of several homologues showed opposite expression patterns when compared to mock treated

plants (Supplementary Table 5). Moreover, we observed that a conserved set of PTI markers,

shown to be activated in Marchantia upon interaction with bacteria (Gimenez-Ibanez et al.,

2019), oomycetes (Carella et al., 2019) and fungi (Redkar et al., 2022), were also induced in

response to TMV infection (Figure 3A). Additionally, a significant number of intracellular

receptors from the NLR family displayed upregulated expression upon TMV infection ($p < 6.245e-$

05 with those described by Bowman et al., (2017); $p < 1.308e-05$ with those described by Chia

et al., (2022); Supplementary Table 6). Nevertheless, none of them showed significant sequence

similarity to the NLR N gene triggering ETI responses in TMV infected tobacco plants (Whitham

et al., 1994). Likewise, several components from the RNA silencing machinery upregulated in

tobacco plants upon TMV infection (Sheng et al., 2019), were also differentially expressed in

response to TMV in Marchantia plants. The non-canonical *MpDCL1b (Mp6g09830)*, *MpDCL4*

(Mp7g11720), and *MpRDR6 (Mp8g14970)* were upregulated, whereas *MpDCL3 (Mp1g02840)*,

MpAGO9 (Mp8g08610) and *MpNRPE1b (Mp6g00230)* were suppressed.

Gene Ontology (GO) analysis of the upregulated genes revealed an enrichment (padjust<0.05)

of biological processes related to reactive oxygen species, wounding, light intensity, heat

responses and responses to fungus and the presence of their constituent chitin (Supplementary

Table 7, Supplementary Figure 3A). Photosynthesis, DNA packaging and cell cycle, microtubule

organization or regulation of salicylic acid metabolism were among the downregulated processes

(padjust<0.05; Supplementary Table 8, Supplementary Figure 3B).

Notably, TMV is mainly transmitted through wounds and wounding response was one of the

upregulated processes in the GO analysis, which prompted us analysing this response. ERF

transcription factors are central to trigger wounding responses in vascular plants such as *A.*

thaliana (Heyman et al., 2018) with a conserved set of ERF transcription factors being also

recently described to respond to wounding in *Marchantia polymorpha* plants (Liang et al., 2022).

A subset of these wounding responsive ERF transcription factors showed the same expression

pattern upon TMV infection as that observed upon intensive wounding (Figure 3B, Liang et al.,

2022). To determine the whole transcriptional overlap between wounding and TMV responses,

we reanalysed published tissue incision transcriptomic responses (Liang et al., 2022) using the

same pipeline we used for the TMV samples. Comparison of both datasets showed a significant

overlap both in up-regulated (593 genes; p<6.631e-235) and down-regulated genes (162 genes;

p<2.137e-70; Figure 3C), suggesting that TMV infection contributes to a sustained wound

response in Marchantia.

Chitin is one of the most abundant polysaccharides in nature and it is a primary component of

cell walls in fungi and exoskeletons of arthropods such as insects, which in turn can work as viral

vectors in a range of plants species (Campbell, 1996). According to the GO enrichment in

response to chitin and fungi, and besides the fungal responsive genes described above as part

of a conserved PTI response, we also observed upregulation of a set of fungal responsive WRKY

transcription factors (Figure 3D; Redkar et al., 2022, Yotsui et al., 2023).

Photosynthetic deficiency is a hallmark for several stress responses, including those involving

viral infections. Accordingly, downregulation of transcripts involved in photosynthesis was one

of the main processes affected in our GO study as consequence of TMV infection. We determined

the photosynthetic efficiency in TMV infected areas from 10 Tak-1 plants after 2-weeks of

infection along with the very same area in mock treated plants. TMV infected areas showed a

significant reduction in photosynthetic competency when compared to similar areas from mock

treated plants (Figure 3E). Notably, parts of those very same plants distant to the infection foci,

showed also reduced photosynthetic capability, although to a lesser extent (Figure 3F). Those

results suggest that TMV responses might be systemic in *Marchantia polymorpha* plants.

As previously observed in *Nicotiana tabacum*, Marchantia plants infected with TMV are

compromised in cell cycle progression and DNA packaging (Jada et al., 2014). To further

corroborate the impact of TMV infection in the cell cycle, we infected Marchantia plants carrying

the G2-M reporter proMpCYCB;1:D-box-GUS (Hernandez-Garcia et al., 2021). TMV infected

plants showed reduced levels of the CYCB1 reporter as compared to mock treated plants (Figure

3G). That result supports the transcriptomic data pointing to reduced cell division upon TMV

infection.

Overall, those results show that TMV engages in a compatible interaction with *Marchantia*

polymorpha.

Salicylic and OPDA-dependent signaling pathways are active at TMV infected areas

The plant hormones Salicylic acid (SA) and Jasmonic acid (JA) are central in defence responses

against most pathogens of vascular plants, acting both in a synergistic and antagonistic manner.

Whereas the SA biosynthesis and signalling pathway are mostly conserved in *Marchantia*
polymoprha (Jia et al., 2023), the active molecules binding to the COI1 receptor and triggering
jasmonate signaling are a group of long chain polyunsaturated fatty acids, including dinor-OPDA,
instead of JA-Ile (Kneeshaw et al., 2022, Monte et al., 2018). Therefore, we compared the SA and
jasmonate-related transcriptomic responses with those found during TMV infection. To that end,
we used the time course datasets obtained on incubation of Marchantia with 1mM SA during 2,
6 and 12 hours (Jeon et al., 2022), SA concentration shown to promote defence against bacterial
and to increase susceptibility to fungal pathogens (Jeon et al., 2022, Matsui et al., 2020).
Comparison of the DEGs identified on SA treatment showed a significant overlap with those
responding to TMV infection (Figure 4A; Supplementary Figure 4). A total of 225 genes were
commonly up-regulated by 2 hours SA treatment or TMV (p<1.799e-100), 306 (p < 1.382e-172)
were shared after 6 hours treatment, and 92 (p < 3.580e-47) by 12 hours of SA incubation.
Likewise, TMV infection shared 47 down regulated genes (p<1.618e-14) with 2 hours, 38 (p <
2.897e-13) with the 6 hours, and 17 genes with 12 hours incubation samples (p < 1.720e-05),
showing that TMV infection activates SA-dependent transcriptional responses.

To ascertain a possible contribution from the jasmonate signalling pathway, we compared the
TMV transcriptional response with that of plants treated with dnr-OPDA (Monte et al., 2018) or
having a constitutively activated jasmonate pathway as a consequence of the mutation of the
JAZ1 signalling repressor (jaz1 KO plants; Monte et al., 2019). A total of 143 and 81 genes were
commonly up-regulated upon TMV infection and *jaz1* deficiency or dnr-OPDA treatment, while
238 were downregulated in TMV infected and jaz1 KO plants (Figure 4B)., suggesting that TMV
infection also activates the jasmonate signalling pathway in Marchantia.

To further understand which were the processes under control of SA and JA signalling pathways
on TMV infection, we carried out a GO analysis of the commonly regulated genes. This showed
that the JA pathway was responsible for activation in Marchantia of wounding, reactive oxygen
species and virus responses, in addition to the catabolism of several compounds, including lipids,
phenylpropanoids and salicylic acid. Interestingly, some of these processes were also
upregulated by SA treatment, including wounding, SA and lignin catabolism among others. The
SA response pathway was on the other side also responsible for the observed transcriptional up-
regulation of defence genes against bacteria and fungi, as well as responsive to chitin (Figure 4C,
Supplementary Table 9)

Furthermore, amongst the downregulated processes by JAZ1 deficiency and TMV infection were
translational elongation, mucilage and cell wall biosynthesis, response to fructose,
photosynthesis, microtubule-based process, growth, and response to abscisic acid (ABA; Figure
4D, Supplementary Table 9)

Collectively, those results show that SA and JA signalling pathways are activated in Marchantia
during TMV infection.

TMV infection triggers aging in *Marchantia polymorpha*

Wound response, mitotic cell arrest, and reduced photosynthetic efficiency are hallmarks for
plant cell senescence (Espinoza et al., 2007, Gan, 2003). Senescence is the final stage of the life
cycle of a cell, tissue or organism and it is gradually triggered throughout aging. Additionally, SA
and JA, along with ET, contribute to senescence in *Arabidopsis thaliana* (Schommer et al., 2008,
Yu et al., 2021). Senescence can be also triggered by the interaction with different pathogens,
including the crucifer infecting TMV-Cg strain capable of compatible interactions with the
Arabidopsis UK-4 ecotype (Espinoza et al., 2007, Lim and Nam, 2005). Since all those senescence
hallmarks were present in our molecular characterization of the Marchantia-TMV interaction,
we first checked the expression levels of several senescence marker genes in our transcriptomic
data (Kulshrestha et al., 2022). We found that many of those senescence markers, such as
MpMYB14 (Mp5g19050), *MpORE1* (Mp2g07720), *MpNYE1* (Mp1g17090) and *MpWRKY7*
(Mp3g17660), were induced upon TMV infection, hence suggesting that TMV infection could
trigger aging and tissue maturation in Marchantia. Transcriptional trajectories orchestrating
maturation and aging in Marchantia, both at organism and tissue level, have been recently
reported through transcriptional studies over developmental time or by analysing sections along
the thalli from top to base (Wang et al., 2023). Clustering the transcriptional profile of mock and
TMV infected plants with those datasets, based on Pearson's correlation coefficient and
regression model, showed that TMV infection promotes maturation and aging. Thus, while the
transcriptome from mock treated plants clustered with juvenile state samples (slide 2 and Day
1), that from TMV infected plants clustered with slide 4 and day 31 of development (Figure 5A,
5B).

Those results indicated that TMV infection triggers senescence in Marchantia, a characteristic
trait from compatible plant - virus interactions, and aging.

MpNAC7 is a conserved interactor of the TMV silencing suppressor p126 and enhances defence against TMV in tobacco plants.

The TMV replicase p126 is essential for viral replication and movement, but it also works as a silencing suppressor in vascular plants (Ding et al., 2004). To gain insights into its possible role during TMV infection in Marchantia plants, we first established its subcellular location. We found that p126 fused to GFP localized in cytoplasmatic bodies in mesophyll cells when transiently expressed in *Nicotiana benthamiana* leaves (Figure 6A) consistently with earlier studies in onion epidermal cells, where the full length p126 protein fused to GFP was found associated to the endoplasmic reticulum (dos Reis Figueira et al., 2002). Similarly, Marchantia plants expressing this very same construct, showed discrete cytoplasmic spot-like inclusions in oil body cells (Figure 6A), showing that p126 subcellular location is conserved between Nicotiana and Marchantia.

Constitutive expression of viral-derived silencing suppressors often leads to plant developmental defects, like growth stunting (Kasschau et al., 2003). Accordingly, two-weeks old Marchantia plants constitutively expressing p126 were smaller in size when compared to control plants (Figure 6B).

The TMV replicase p126 is known to associate with several host factors in Arabidopsis, including TOM1 (Yamanaka et al., 2000), GDI2 (Kramer et al., 2011) and the transcription factor ATAF2 (Wang et al., 2009). To assess whether those interactions were conserved between p126 and the corresponding orthologs, we performed Y2H assays. Orthologs were identified by blasting the Arabidopsis protein sequences at MarpolBase (https//Marchantia.info) and PlantRegMap (http://plantregmap.gao-lab.org/). The P126 full length and helicase domain (nt 2442 to 3636) were fused to the GAL4 binding domain and used as baits, while the putative interactors fused to the GAL4-activation domain, were used as preys. Notably, the only conserved interaction was that of the ortholog of the NAC transcription factor ATAF2, *MpNAC7* (Mp6g02620) (Figure 6C). As for p126-ATAF2 binding (Wang et al., 2009), p126-MpNAC7 interaction could be confirmed *in planta* using bimolecular luminescence complementation (Figure 6D). Moreover, interaction with p126 was found to trigger proteasome dependent ATAF2 degradation in the context of TMV infection (Wang et al., 2009), and in our assays luminescence signal from p126-MpNAC7 interaction was always lower when TMV was co-infiltrated together with MpNAC7 and p126. A similar reduction was not observed in the presence of TuMV, indicating that specific MpNAC7 degradation during TMV infection is a conserved feature in Marchantia (Figure 6D).

In Arabidopsis, ATAF2 expression is triggered by TMV infection (Wang et al., 2009). Likewise, we

found that MpNAC7 expression is also increased on TMV infection (Log2(fold-change)= 0.991),

wounding (Log2(fold-change)= 2.67), dnr-OPDA and after 6 hours of SA treatment (Log2(fold-

change)= 1,23) in Marchantia. Likewise, ATAF2 is induced by wounding, SA and JA treatments

(Delessert et al., 2005) and contributes to defence against TMV in Arabidopsis (Wang et al.,

2009). Here, to ascertain the role of MpNAC7 in defence against TMV, and rule out a possible

connection with RNA silencing- mediated responses, we infected 1 month old Nicotiana plants

deficient in RDR6 (rdr6i; (Schwach et al., 2005)) with MpNAC7, or GUS as control, in the presence

of TMV or TuMV. TMV and TuMV strains were labelled with the ROS1 gene which triggers

anthocyanin production, and hence we followed presence of the viruses by the purple colour

(Bedoya et al., 2012). We observed that co-expression of MpNAC7 abrogated local and systemic

TMV infection with respect to plants co-expressing the GUS control. By contrast, MpNAC7 co-

expression was of no effect on TuMV infected plants, that displayed a similar local and systemic

presence of the virus as the GUS co-infiltrated controls (Figure 6E, Supplementary Figure 5).

Collectively, those results show that molecular interaction between the TMV silencing

suppressor p126 and the host NAC transcription factor is conserved in Marchantia, as well as the

role of this transcription factor in defence.

Discussion

In this work, we have characterized the interaction of the non-vascular liverwort *Marchantia*

polymoprha with viruses. First, we have defined known and unknown viruses that associate with

Marchantia in natural settings. These studies show that Marchantia is primarily associated with

RNA viruses in nature, validating previous predictions based on the presence of viral gene

hallmarks in the transcriptomes from the 1000 Plants Genome Project (Mushegian et al., 2016).

We also showed that at least one of these newly discovered viruses is able to replicate in

laboratory grown Marchantia Tak-1 plants. Second, we have found that TMV engages into a

compatible interaction with Marchantia plants. That interaction encompasses an extensive

transcriptional reprogramming reminiscent of that found in vascular plants such as Arabidopsis

and Nicotiana, albeit involving also previously uncovered differences like the activation of a

sustained wounding response. Third, we have found that the TMV silencing suppressor p126

displays similar subcellular location in Marchantia cells, as in Nicotiana, and that its interaction

with a NAC transcription factor, MpNAC7, also specifically abrogates TMV infection in Nicotiana

plants. All those commonalities show that core molecular events of plant-virus interactions
found in vascular plants are also present in non-vascular plants.

Marchantia plants infected with TMV undergo a transcriptional reprogramming consistent with
a pathogenic process and characterized by the activation of the defence related PR proteins and
a conserved PTI response. Perception of dsRNAs have been found to trigger PTI in response to
TMV infection in Arabidopsis, which in turn contributes to antiviral defence (Korner et al., 2013).
Indeed, dsRNA-triggered PTI is dependent on the Arabidopsis PRR co-receptor SERK1 (Niehl et
al., 2016), which has a corresponding ortholog in the Marchantia genome (Mp7g09160), and
which was shown to be involved in PTI against *Pseudomonas syringae,* and to be targeted by the
PTI suppressors AvrPto (Gimenez-Ibanez et al., 2019). Thus, it will be interesting in future studies
to analyse the role of MpSERK1 in viral PTI responses. Additionally, TMV infection triggers the
expression of a significant number of NLR genes that can potentially contribute to an ETI
response. In the absence of a DCL2 gene in Marchantia, we also observed changes on the
expression of core genes of the RNA silencing machinery in response to TMV infection. In
contrast to the repression of elements from the RNA-Directed DNA methylation pathway, such
as *MpDCL3*, *MpAGO9* and *MpNRPE1b*, the central elements within the RNA antiviral machinery
MpDCL4 and *MpRDR6* were activated. Such general switch on RNA silencing pathways in
Marchantia is reminiscent of that found specifically restricted to the vasculature of *Nicotiana*
benthamiana in response to TMV (Collum and Culver, 2017). Those results suggest an
evolutionary re-routing of defence mechanisms after acquisition of novel tissues or cell-types
that become new focal points for plant-virus interactions.

Interestingly, our results also uncover a previously unknown potential role of wound response
as an antiviral defence mechanism. Plant cells undergoing wound responses experience
inhibition of their photosynthetic ability becoming sink tissues on the transport of
photoassimilates (Quilliam et al., 2006), attracting TMV movement (Brants, 1961). Thus, it is
tempting to speculate that sustained wound response in Marchantia plants, along with local and
systemic decreased on photosynthetic capability and senescence confine TMV infection
preventing its movement throughout the plant.

Additionally, salicylic and jasmonate pathways are central for defence against several pathogens.
A spatially complementary synergistic interaction of both was observed to be necessary for
defence against *Pseudomonas syringae* in Arabidopsis (Betsuyaku et al., 2018, Salguero-Linares
et al., 2022). Likewise, their synergistic interaction is responsible for local and systemic resistance

against TMV in *Nicotiana benthamiana* (Zhu et al., 2014). In line with what was found in
Nicotiana, our results show that both pathways are activated in Marchantia plants upon TMV
infection, whilst initial results, show that both pathways are antagonistically involved in
Marchantia in the defence against fungal and bacterial pathogens (Gimenez-Ibanez et al., 2019,
Matsui et al., 2020).

Our work also highlights the conservation of central players, such as MpNAC7, in the specific
defence against TMV. Future research might focus on comparative studies to ascertain the
conservation of the genetic networks under MpNAC7 and ATAF2 regulation both in Marchantia
and Arabidopsis.

Collectively, results from this work suggest the presence of evolutionary conserved and divergent
plant strategies to cope with viral infections. Those might include gain and loss of defence
strategies, such as those involving wound responses. Evolution of plant-virus interactions might
also encompass the gradual acquisition of new elements within the underlying gene networks,
as those physically interacting with pathogen-derived proteins. Finally, defence mechanisms
might undergo re-location to newly pathogen-targeted cell-types and tissues over evolutionary
times.

Material and methods

Plant growth conditions

Male *Marchantia polymorpha* plants Takaragaike-1 (Tak-1) were grown on plates with half
strength Gamborg´s B5 medium (Duchefa) containing 1% agar and 1% sucrose under continuous
light and 22ºC.

Nicotiana benthamiana plants were grown on soil under long day conditions (16h light/ 8h dark)
and 25ºC.

Infection assays

Frozen tissue from Marchantia plants from the Tübingen population or lab grown Tak-1 plants
was homogenized in liquid nitrogen cold mortars, resuspended in phosphate buffer pH7.2,
filtered using 70 µM filters and kept at 4ºC. Two-week old Tak-1 plants grown in sterile conditions
in plates were vacuum incubated with each of both extracts and transferred to plates containing

wet soil. Eight Tak-1 individuals exposed to each sap were collected either one or two weeks

after incubation to assess viral replication.

Two weeks-old *Nicotiana benthamiana* plants were infiltrated with an agrobacterium solution

(0.5 O.D) carrying the TMV-GFP clone (Peart et al., 2002). Infected Nicotiana plants were grown

for 3 additional weeks along with non-infected plants and upper leaves displaying infection

symptoms were collected, along with equivalent ones from non-infected plants and frozen in

liquid nitrogen. Subsequently, leaf tissue was grinded in a liquid nitrogen cold mortar. Sap was

isolated from grinded leaf tissue using phosphate buffer pH 7.2, filtered with 70 µM filters and

maintained at 4ºC. Two-weeks old Tak-1 plants were incubated either in TMV-GFP containing sap

or sap from non-infected plants (mock) for 5 minutes in vacuum. After incubation, Tak-1 plants

were transferred to plates containing wet soil and brought back to the growth chamber. GFP

signal was monitored 1 and 2 weeks after incubation using a portable UV lamp or with a

magnifying stereoscope couple to a UV light source.

RNA isolation and expression analysis

Total RNA was extracted from Marchantia plants using the protocol described in Box et al.,

(2011). Isolated RNA was further cleaned using the RNeasy Kit (Qiagen). 500 ngs of total RNA

were DNase I treated (Fisher Scientific) and converted into cDNA using the RevertAid first strand

kit (Fisher Scientific) following the manufacturer instructions. RT-qPCR experiments were

performed in a lightcycler (ROCHE) device. Relative expression changes were calculated using

the $2^{-\Delta\Delta Ct}$ method using *MpACTIN7* (Mp6g11010) gene as housekeeping. Primers utilized can be

found in supplementary table 8.

RNA-seq analysis

For virome studies, 1 µg of total RNA from each of the 14 individuals included in the study was

pooled and 4 µg from the mixture were used for preparing a single library using total ribo-

depleted RNA following the manufacturer instructions (Illumina TruSeq Stranded total RNA LT

sample prep Kit, plant) and sequenced on a Illumina NovaSeq 6000 device in 2x150 bp mode. A

total of 306,895,552 reads were obtained. The reads were trimmed and further analysed using

QIAGEN CLC Genomics Workbench 21 (https://digitalinsights.qiagen.com/). Dataset was

analysed for the presence of known and new viruses using an established virus detection

pipeline (Pecman et al., 2017), which includes: reads trimming, *de novo* assembly of trimmed

reads, mapping of reads and contigs to NCBI viral RefSeq database (search for known viruses)

and pFam domain search for contig sequences (search for known and unknown viruses). To

detect known viruses, results of reads mapping to viral RefSeq database were inspected and

filtered for viruses for which at least 1000 nts of the reference sequence were covered by

mapped reads (Supplementary table 1). In parallel, to detect known viruses and search for new

ones, results of the pFam domain search of contig sequences were filtered for hits to viral

proteins. Then, this candidate list of viral contigs was additionally queried for similarity against

complete NCBI *nr* database (October 2020) using blastx (Gish and States, 1993) and the similarity

search results were visualized and taxonomically classified using MEGAN 6 (Huson et al., 2016).

Contigs corresponding to cellular organisms in this analysis were excluded from the list to obtain

a final list of putative viral-like sequences (Supplementary table 2). Trimmed sequencing reads

were mapped to the list of putative viral-like sequences, to estimate the abundance of

corresponding nucleic acids in the sequenced sample and the results were reported as average

mapping depth (number of times a position is covered by sequencing reads, averaged over a

complete contig length; Supplementary table 2).

For a newly discovered picorna-like virus, named Tübingen picorna-like virus 1, additional

analyses were performed, aiming to ascertain its association with *M. polymorpha*. Amino acid

sequence of ORF 1 of the virus (containing RNA-dependent polymerase (RdRp) domain) was

queried for similarity against complete NCBI *nr* database (January 2022) using blastp. Fifteen

best hits according to bit score values (which have at least 33% identity with the conserved part

of the RdRp domain of the queried virus) were included into phylogenetic analysis. Alignment of

the sequences of the conserved part of RdRp was performed using MAFFT 7.48 (Katoh and

Standley, 2013), followed by alignment trimming using TrimAI 1.3 in Automated 1 mode (Capella-

Gutiérrez et al., 2009). Then, best protein substitution model was selected using Model Finder

(Kalyaanamoorthy et al., 2017) and maximum likelihood phylogenetic tree inference was

performed based on the trimmed alignment using IQ-Tree 1.6.12 (Nguyen et al., 2015), using

1000 ultrafast bootstrap replicates (Hoang et al., 2018). The resulting phylogenetic tree was

visualized using iTOL 6.5 (Letunic and Bork, 2021) and viral taxa on the tree were coloured

according to the associated sample source, as designated in NCBI GenBank.

For analysing the transcriptional response in Marchantia plants to TMV infection, 1 ug of total

RNA was used for preparing each stranded library by mRNA enrichment following manufacturer

indications (TrueSeq kit from Illumina). Libraries were sequenced on a Novaseq 6000 device. A

total of 20-21 million pairs were obtained per sample. Adapter removal and trimming of low

quality bases was done using Trim Galore! v0.6.1
(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), removing reads shorter
than 20 bps and non-paired reads after the trimming step. Ribosomal RNA reads were removed
using SortMeRNA v2.1b (Kopylova et al., 2012), removing also reads that become unpaired after
this filtering step.

Cleaned reads together with the transcriptome of *Marchantia polymorpha* were used to quantify
gene expression at transcript level using Salmon v1.1.0 (Patro et al., 2017).

For this, the file MpTak1v5.1_r1.mrna.fasta containing the Marchantia transcriptome (v5.1) was
downloaded from https://marchantia.info/download/tak1v5.1/ (as of 19-Jan-2021). This file was
indexed using *Salmon index* and then used as input for *Salmon quant*, which was ran with
parameters -l A, –validateMappings, --recoverOrphans, --rangeFactorizationBins 4, --seqBias,
and –gcBias. Mapping rate was found to be in the 89.5%-92.58% range.

The R progam tximport v1.14.2 (Soneson et al., 2015) was used to aggregate Salmon's transcript
expression estimates at gene level and the *DESeqDataSetFromTximport* from the R package
DESeq2 v1.26.0 (Love et al., 2014) was used to create an R object with count data and
experimental information for all samples. A low-expression filter was then applied to remove any
gene with less than 10 counts across all 8 analyzed samples, resulting in a final set of 14,980
genes (from 19,217 annotated genes). Then, the *DESeq* function, which, among other things,
computes size factors for sample normalization, was applied to the filtered object. DESeq2's *rlog*
function was used to transform count data to the log2 scale while minimizing for differences
between samples for low-expressed genes. The transformed data was fed to DESeq2's *plotPCA*
function to visualize a PCA plot of the samples.

The analysis of differential expression was done using DESeq2's *results* function with the filtered
and normalized object as input and parameter *alpha = 0.05.*

All datasets used in this study for comparative analysis (wound response and salicylic acid time
course) were analysed in the very same way as described above.

All Gene ontology analysis (GO) were performed using PLAZA
(https://bioinformatics.psb.ugent.be/plaza/). Redundancy on representative categories was
reduced using Revigo (http://revigo.irb.hr/).

For correlation analysis of transcriptome responses between developmental trajectories and
TMV response, raw reads were processed as described above, except that rRNA was removed
using ribodetector_cpu v0.2.7 (Deng et al., 2022) and Salmon version was v1.9.0. Salmon's
counts were converted into a DESeq2 object and a low-expression filter was applied as described
above. Next, the existing batch effect between time course samples and sliced tissue ones was
removed using the Combat_seq function from the sva package (Leek et al., 2023), v3.48.0. Then,
values for biological replicates were averaged and Pearson correlation was computed for the
resulting count matrix and displayed using the Heatmap function from the ComplexHeatmap
package (Gu, 2022). The batch correction, correlation and heatmap plotting steps were based
on the R script used in (Wang et al., 2023) and kindly provided by the authors.

Constructs and plant transformation

cDNA from Marchantia mock treated plants was used to amplify the coding regions of *MpTOM1*
(Mp3g00660), *MpGDI2* (Mp6g05010) and *MpNAC7* (Mp6g02620) by PCR using Phusion
polymerase (NEB). Arabidopsis *Bobber1* (At5g53400) was amplified using cDNA synthesized from
RNA from Arabidopsis infolorescences. DNA was isolated from four-weeks old plants using
extraction buffer (200mM Tris pH 7.5, 25 mM EDTA, 250 mM Na Cl, 0.5% SDS) followed by
isopropanol precipitation, washed with Ethanol 70%, and resuspended in MiliQ sterile water.
Isolated DNA was used as template to amplify the genomic region covering the promoter and
degron to generate the MpCYCD1;1B:GUS reporter by amplification in 5 overlapping fragments
that were included in a final PCR to reconstruct the final region. P126 was amplified using DNA
from the TMV-GFP containing plasmid. All obtained fragments were further incubated with dATP
and Taq DNA polymerase (NEB) for 1 hour before cloned into the gateway compatible vector
PCR8/GW/TOPO TA (Invitrogen). An entry vector containing the Arabidopsis AGO1 was kindly
provided by Pablo Manavella. Inserts were cloned into their final vectors through LR reaction
using LR clonase II (Invitrogen). All plasmids were submitted to Sanger sequencing to corroborate
their integrity before used in the corresponding experiments. A list of used primers and
constructs can be found in supplementary table 10 and supplementary table 11.

Plant transformation was performed as described in Ishizaki et al., (2008)and selected in
Gamborg´s B5 media supplemented with Kanamycin 100mg/ml and Cefotaxime 100mg/ml.

Histology and imaging

Marchantia plants were embedded in 3% agarose and cross-sections of around 200 μM were obtained using a vibratome. A total of 12 MpCYCD1;1B:GUS reporter plants, either TMV infected or mock, were fixed in 90 % acetone, and GUS activity was assayed as described in Blazquez et al., (1997). Confocal imaging was performed using Leica SP5. Images were processed using Fiji (Schindelin et al., 2012).

Protein-protein interaction assays

Interactions in the yeast two-hybrid system was assayed on selective medium (L$^-$, W$^-$, H$^-$). The assay was repeated 3 times. For the luciferase complementation assay, one-month old rdr6i Nicotiana plants were used. *Agrobacterium tumefaciens* cultures with the different constructs were mixed in the same amounts at a final OD$_{600}$=0.6. Leaves were imaged three days after inoculation using a LAS4000 device.

Photosynthetic measurements

10 infected and 10 mock treated plants were subjected to 10 measures each using a IMAGING-PAM Maxi device (Heinz Walz GmbH). Quantification was done using the ImagingWing integrated software.

Acknowledgments

We would like to Juan Jose López-Moya, Miguel A. Blázquez and Salomé Prat for critical reading of the manuscript. Sebastian Schornack, Phil Carella, Alexander Summers, the CUBG curation team (Samuel Brockington and Mar Millan), Klaus Schlaeppi, Katja Rembold, Alba Gonzalez, Detlef Weigel, Alexandra Kehl and Brigitte Fiebig for their help in obtaining Marchantia individuals for the virome assays. We would like to thank Eduardo Bejarano for providing the plasmid containing TMV-GFP and Antonio Darós for providing TuMV-ROS1 and TMV-ROS1 containing plasmids. Pablo Manavella for the AtAGO1-pEntr plasmid. Phil Carella for an updated annotation of NLR genes in Marchantia. We would also thank to Jia-Wei Wang, Long Wang and Jie Xu for sharing the code used for correlation analysis between TMV transcriptional responses and the different samples from Marchantia developmental trajectories.

The work in the MoRE lab is funded by RYC-2015-19154 (funded by MCIN/AEI/ 10.13039/501100011033 and by "ESF Investing in your future"), RTI2018-097262-B-I00 (funded by MCIN/AEI/ 10.13039/501100011033 and by "ERDF A way of making Europe") and through the "Severo Ochoa Programme for Centres of Excellence in R&D" 2016-2019 (SEV-2015-0533)

and 2020-2023 (CEX2019-000902-S) funded by MCIN/AEI/ 10.13039/501100011033 and the
CERCA programme from the Generalitat de Catalunya to I-R-S. T J-G was recipient of a
Postdoctoral Fellowship from CRAG through the "Severo Ochoa Programme for Centres of
Excellence in R&D" 2016-2019 (SEV-2015- 0533). L. V-M was supported by BES-2016-076986
(funded by MCIN/AEI/ 10.13039/501100011033 and by "ESF Investing in your future"). The work
at NIB was funded by Slovenian Research and Innovation Agency core and project financing P4-
0407, P4-0165 and J4-4553.

Data access: Raw data from the virome has been deposited in SRA and GenBank under Bioproject
(PRJNA1022866) and BioSample (SAMN37631837). Raw data from TMV infected vs. Mock
treated plants have been deposit GEO (GSE241827). Reconstructed genomic sequences of three
novel viruses uncovered by virome analysis were deposited in NCBI GenBank (OR640980-82).

Figure legends

Figure 1. *Marchantia polymorpha is infected by RNA viruses.*

(A) Genome structure of Tübingen picorna-like virus 1, newly discovered in composite *M.*
polymorpha sample with two predicted open reading frames (ORFs) and corresponding
average reads mapping depth in sequenced composite sample.

(B) Maximum likelihood phylogenetic tree obtained based on the alignment of the
conserved part of RdRd of Tübingen picorna-like virus 1 and fifteen other most similar
viruses. The tree was midpoint rooted. Coloured virus names correspond to the sources
associated with a particular virus, according to the legend on the right side of the panel.
Numbers next to the branches represent statistical support (%) of the clades according to
the ultrafast bootstrap analysis. Codes next to the taxon names in round brackets are the
corresponding NCBI GenBank protein accession numbers. Next to them, fractions in square
brackets represent % of identity of Tübingen picorna-like virus 1 to other individual taxons
for the aligned part of the RdRp domain.

(C) Results of the viral concentration measurements for *M. polmorpha* plants inoculated
with Tübingen picorna-like virus 1 positive plant extracts. On the right, relative
concentrations of Tübingen picorna-like virus 1 in inoculated plants one- and two-weeks
post inoculation are shown; measurements of concentrations in individual plants are shown
as dots and the box-whisker plots represent the distribution of values for the two time

points (at 1 WPI, x represent an outlier measurement). On the top of the plot, the horizontal

connecting line designates statistically significant difference in the means of the two

distributions, with the corresponding p-value, according to the nonparametric Wilcoxon

rank sum test.

Figure 2. *Tobacco mosaic virus replicates in Marchantia plants*.

(A) GFP fluorescent signal increases in plants after 1 week, (left panel) compared to 2-weeks

infected plants (right panel). Both images show the same part or an infected plant under

a magnification device under UV light. Red shows chlorophyll autofluorescence. White

bar scale shows 100 µms.

(B) TMV-GFP transcripts increase over time in infected plants. Pools of three plants were

used for each time point. T0 corresponds to samples collected immediately after sap

incubation, while T1 and T2 correspond to samples collected 1 and 2 weeks after

incubation.

(C) TMV-GFP infects all cell types. Transversal section of and infected area under confocal

image. The picture shows the merge between green (virus) and red (autofluorescence)

channels. Merge was performed using Fiji. P stands for Photosynthetic region, S for

storage region and E stands for lower epidermis. White bar scale shows 1 µm.

Figure 3. *TMV triggers stress-related molecular reprogramming in Marchantia.*

(A) Set of wound responsive ERF transcription factors.

(B) Venn diagram showing that transcriptional reprogramming upon TMV infection largely

overlaps with that found upon wounding.

(C) Set of chitin and TMV responsive WRKY transcription factors in Marchantia.

(D) Set of common PTI marker genes responsive to fungi, bacteria and TMV.

(E) Photosynthetic efficiency (Fv/Fm) in infected areas compared to equivalent areas from

mock treated plants. Statistical assessment was performed using a parametric T-test.

(F) Photosynthetic efficiency (Fv/Fm) in non-infected areas, from infected plants, compared

to equivalent areas from mock treated plants. Statistical assessment was performed

using a parametric T-test.

(G) TMV infected plants show lower cell cycle activity as inferred from MpCYCD1;1B:GUS

reporter. Left, staining from reporter plants incubated in sap from mock plants. Right,

staining from reporter plants infected with TMV.

Figure 4. *Salicylic and jasmonate signalling pathways are activated during Marchantia-TMV*

interaction.

(A) Venn diagrams showing the overlap between the transcriptional responses under TMV

infection ad a salicylic treatment time course (2 and 6 hours).

(B) Overlap between the trancriptome responses to TMV and jaz1ko mutants with

constitutively active jasmonate signalling, left, and treatment with the active dnr-OPDA

jasmonate.

(C) Biological processes commonly upregulated in response to TMV infection, SA treatment

(2,6 and 12h) and activation of jasmonate signalling.

(D) Biological processes commonly downregulated in response to TMV infection, SA

treatment (12h) and activation of jasmonate signalling.

Figure 5. *TMV infection triggers maturation and aging in Marchantia*.

(A) Pairwise comparison between mock and TMV samples along with sliced thallus samples.

(B) Pairwise comparison between mock and TMV samples along with samples from

Marchantia developmental series (D, states the days after gemma germination

Marchantia samples were collected)

Figure 6. *Conservation of the interaction between Marchantia and the TMV silencing suppressor*

p126.

(A) P126-GFP shows accumulation in cytoplasmic bodies both in *Nicotiana benthamiana*

and *Marchantia polymorpha* under confocal microscopy. Scale bar is 100 µMs.

(B) Overexpression of p126 inhibits growth (area) when compared to Tak-1 control plants.

Analysis was done on 6 individual plants per genotype and using a two tailed-t-test.

(C) MpNAC7 interacts with the helicase domain of p126 in yeast.

(D) MpNAC7 interacts both with the helicase and the full length p126 in planta.

(E) Transient expression of MpNAC7 in Nicotiana plants abrogates local and systemic TMV

infection. TMV presence is shown as purple.

Supplementary figure 1. *Nicotiana benthamiana plants inoculated with sap from Marchantia*

infection experiments.

(A) Nicotiana plants inoculated with sap from mock treated plants. Upper panel, top view

of plant with white light. Down panel, same Nicotiana plant under UV light.

(B) Nicotiana plants inoculated with sap from TMV-GFP treated plants. Upper panel, top
view of plant with white light showing crinkled and yellowish leaves as result of TMV
infection. Down panel, same Nicotiana plant under UV light showing GFP signal from
TMV infection in leaves showing symptoms.

(C) Inset from a TMV-GFP infected leaf showing symptoms of infection (reduced chlorophyll
content) in the upper panel, coinciding with TMV-derived GFP signal under UV light.

Supplementary figure 2. *Sampling and validation of samples for transcriptomic assays.*

(A) Example of areas collected from mock, upper panel, and TMV-GFP infected plants for
transcriptomic assays.

(B) RT-qPCR validation of TMV-GFP levels in the 4 pools used for RNA-seq.

(C) Principal component analysis of the different replicates from RNA-seq experiment.

Supplementary figure 3. *Gene ontology analysis of biological processes to which DEGs in respond*
to TMV infection belong.

(A) GO categories from upregulated genes in response to TMV infection.

(B) Selected GO categories from downregulated genes in response to TMV infection.

Supplementary figure 4. *Overlap between DEGs responding to 12 hours of incubation in salicylic*
acid and those responsive TMV infection.

Supplementary figure 5. *MpNAC7 specifically abrogates TMV infection in Nicotiana plants.*

(A) Picture of inoculated leaves showing that MpNAC7 abrogates local TMV but not TuMV
infection (purple colour).

(B) Effect in systemic leaves.

(C) Upper view of infected plants.

Supplementary table 1. *Detection of viral sequences of known viruses*.

Supplementary table 2. *Detection of known and putative novel virus-like sequences*.

Supplementary table 3. *Demographic study for the presence of a set of known and newly*
identified virus in individuals from the different sampling sites.

Supplementary table 4. *List of DEG in response to TMV infection in Marchantia.*

Supplementary table 5. *List of Pathogenesis-related (PR) genes differentially expressed upon TMV infection.*

Supplementary table 6. *List of NLR genes differentially expressed upon TMV infection*

Supplementary table 7. *List of GO categories of the upregulated biological processes in response to TMV infection.*

Supplementary table 8. *List of GO categories of the downregulated biological processes in response to TMV infection.*

Supplementary table 9. *List of GO categories of the biological processes commonly responsive to TMV infection and salicylic acid time course treatment, jaz1ko mutants and dnr-OPDA treatment.*

Supplementary table 10. *Table of construct used in this study.*

Supplementary table 11. *Table of primers used in this study.*

Bibliography:

BALLY, J., NAKASUGI, K., JIA, F., JUNG, H., HO, S. Y., WONG, M., PAUL, C. M., NAIM, F., WOOD, C. C., CROWHURST, R. N., HELLENS, R. P., DALE, J. L. & WATERHOUSE, P. M. 2015. The extremophile Nicotiana benthamiana has traded viral defence for early vigour. *Nat Plants,* 1**,** 15165.

BECK, M., WYRSCH, I., STRUTT, J., WIMALASEKERA, R., WEBB, A., BOLLER, T. & ROBATZEK, S. 2014. Expression patterns of flagellin sensing 2 map to bacterial entry sites in plant shoots and roots. *J Exp Bot,* 65**,** 6487-98.

BEDOYA, L. C., MARTINEZ, F., ORZAEZ, D. & DAROS, J. A. 2012. Visual tracking of plant virus infection and movement using a reporter MYB transcription factor that activates anthocyanin biosynthesis. *Plant Physiol,* 158**,** 1130-8.

BETSUYAKU, S., KATOU, S., TAKEBAYASHI, Y., SAKAKIBARA, H., NOMURA, N. & FUKUDA, H. 2018. Salicylic Acid and Jasmonic Acid Pathways are Activated in Spatially Different Domains Around the Infection Site During Effector-Triggered Immunity in Arabidopsis thaliana. *Plant Cell Physiol,* 59**,** 439.

BLAZQUEZ, M. A., SOOWAL, L. N., LEE, I. & WEIGEL, D. 1997. LEAFY expression and flower initiation in Arabidopsis. *Development,* 124**,** 3835-44.

BOWLES, A. M. C., PAPS, J. & BECHTOLD, U. 2022. Water-related innovations in land plants evolved by different patterns of gene cooption and novelty. *New Phytol,* 235**,** 732-742.

BOWMAN, J. L., KOHCHI, T., YAMATO, K. T., JENKINS, J., SHU, S., ISHIZAKI, K., YAMAOKA, S., NISHIHAMA, R., NAKAMURA, Y., BERGER, F., ADAM, C., AKI, S. S., ALTHOFF, F., ARAKI, T., ARTEAGA-VAZQUEZ, M. A., BALASUBRMANIAN, S., BARRY, K., BAUER, D., BOEHM, C. R., BRIGINSHAW, L., CABALLERO-PEREZ, J., CATARINO, B., CHEN, F., CHIYODA, S., CHOVATIA, M., DAVIES, K. M., DELMANS, M., DEMURA, T., DIERSCHKE, T., DOLAN, L., DORANTES-

ACOSTA, A. E., EKLUND, D. M., FLORENT, S. N., FLORES-SANDOVAL, E., FUJIYAMA, A., FUKUZAWA, H., GALIK, B., GRIMANELLI, D., GRIMWOOD, J., GROSSNIKLAUS, U., HAMADA, T., HASELOFF, J., HETHERINGTON, A. J., HIGO, A., HIRAKAWA, Y., HUNDLEY, H. N., IKEDA, Y., INOUE, K., INOUE, S. I., ISHIDA, S., JIA, Q., KAKITA, M., KANAZAWA, T., KAWAI, Y., KAWASHIMA, T., KENNEDY, M., KINOSE, K., KINOSHITA, T., KOHARA, Y., KOIDE, E., KOMATSU, K., KOPISCHKE, S., KUBO, M., KYOZUKA, J., LAGERCRANTZ, U., LIN, S. S., LINDQUIST, E., LIPZEN, A. M., LU, C. W., DE LUNA, E., MARTIENSSEN, R. A., MINAMINO, N., MIZUTANI, M., MIZUTANI, M., MOCHIZUKI, N., MONTE, I., MOSHER, R., NAGASAKI, H., NAKAGAMI, H., NARAMOTO, S., NISHITANI, K., OHTANI, M., OKAMOTO, T., OKUMURA, M., PHILLIPS, J., POLLAK, B., REINDERS, A., ROVEKAMP, M., SANO, R., SAWA, S., SCHMID, M. W., SHIRAKAWA, M., SOLANO, R., SPUNDE, A., SUETSUGU, N., SUGANO, S., SUGIYAMA, A., SUN, R., SUZUKI, Y., TAKENAKA, M., et al. 2017. Insights into Land Plant Evolution Garnered from the Marchantia polymorpha Genome. *Cell,* 171**,** 287-304 e15.

BOX, M. S., COUSTHAM, V., DEAN, C. & MYLNE, J. S. 2011. Protocol: A simple phenol-based method for 96-well extraction of high quality RNA from Arabidopsis. *Plant Methods,* 7**,** 7.

BRANTS, D. H. 1961. The Influence of Meristematic Tissue and Injuries on the Transport of Tobacco Mosaic Virus in Nicotiana tabacum L. cultivar. Samsun. *Acta Botanica Neerlandica,* 10**,** 113-163.

CAMPBELL, R. N. 1996. Fungal transmission of plant viruses. *Annu Rev Phytopathol,* 34**,** 87-108.

CAPELLA-GUTIÉRREZ, S., SILLA-MARTÍNEZ, J. M. & GABALDÓN, T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics,* 25**,** 1972-1973.

CARELLA, P., GOGLEVA, A., HOEY, D. J., BRIDGEN, A. J., STOLZE, S. C., NAKAGAMI, H. & SCHORNACK, S. 2019. Conserved Biochemical Defenses Underpin Host Responses to Oomycete Infection in an Early-Divergent Land Plant Lineage. *Curr Biol,* 29**,** 2282-2294 e5.

CARELLA, P., GOGLEVA, A., TOMASELLI, M., ALFS, C. & SCHORNACK, S. 2018. Phytophthora palmivora establishes tissue-specific intracellular infection structures in the earliest divergent land plant lineage. *Proc Natl Acad Sci U S A,* 115**,** E3846-E3855.

CHIA, K.-S., KOURELIS, J., VICKERS, M., SAKA, I. T., KAMOUN, S. & CARELLA, P. 2022. The N-terminal executioner domains of NLR immune receptors are functionally conserved across major plant lineages. *bioRxiv*.

COLLUM, T. D. & CULVER, J. N. 2017. Tobacco mosaic virus infection disproportionately impacts phloem associated translatomes in Arabidopsis thaliana and Nicotiana benthamiana. *Virology,* 510**,** 76-89.

DELESSERT, C., KAZAN, K., WILSON, I. W., VAN DER STRAETEN, D., MANNERS, J., DENNIS, E. S. & DOLFERUS, R. 2005. The transcription factor ATAF2 represses the expression of pathogenesis-related genes in Arabidopsis. *Plant J,* 43**,** 745-57.

DENG, Z. L., MUNCH, P. C., MRECHES, R. & MCHARDY, A. C. 2022. Rapid and accurate identification of ribosomal RNA sequences via deep learning. *Nucleic Acids Res,* 50**,** e60.

DING, S. W., HAN, Q., WANG, J. & LI, W. X. 2018. Antiviral RNA interference in mammals. *Curr Opin Immunol,* 54**,** 109-114.

DING, S. W. & VOINNET, O. 2007. Antiviral immunity directed by small RNAs. *Cell,* 130**,** 413-26.

DING, X. S., LIU, J., CHENG, N. H., FOLIMONOV, A., HOU, Y. M., BAO, Y., KATAGI, C., CARTER, S. A. & NELSON, R. S. 2004. The Tobacco mosaic virus 126-kDa protein associated with virus replication and movement suppresses RNA silencing. *Mol Plant Microbe Interact,* 17**,** 583-92.

DOS REIS FIGUEIRA, A., GOLEM, S., GOREGAOKER, S. P. & CULVER, J. N. 2002. A nuclear
localization signal and a membrane association domain contribute to the cellular
localization of the Tobacco mosaic virus 126-kDa replicase protein. *Virology,* 301, 81-9.
ERICKSON, F. L., HOLZBERG, S., CALDERON-URREA, A., HANDLEY, V., AXTELL, M., CORR, C. &
BAKER, B. 1999. The helicase domain of the TMV replicase proteins induces the N-
mediated defence response in tobacco. *Plant J,* 18, 67-75.
ESPINOZA, C., MEDINA, C., SOMERVILLE, S. & ARCE-JOHNSON, P. 2007. Senescence-associated
genes induced during compatible viral interactions with grapevine and Arabidopsis. *J Exp*
Bot, 58, 3197-212.
GAN, S. 2003. Mitotic and postmitotic senescence in plants. *Sci Aging Knowledge Environ,* 2003,
RE7.
GARCIA-RUIZ, H., TAKEDA, A., CHAPMAN, E. J., SULLIVAN, C. M., FAHLGREN, N., BREMPELIS, K. J.
& CARRINGTON, J. C. 2010. Arabidopsis RNA-dependent RNA polymerases and dicer-like
proteins in antiviral defense and small interfering RNA biogenesis during Turnip Mosaic
Virus infection. *Plant Cell,* 22, 481-96.
GIMENEZ-IBANEZ, S., ZAMARRENO, A. M., GARCIA-MINA, J. M. & SOLANO, R. 2019. An
Evolutionarily Ancient Immune System Governs the Interactions between Pseudomonas
syringae and an Early-Diverging Land Plant Lineage. *Curr Biol,* 29, 2270-2281 e4.
GISH, W. & STATES, D. J. 1993. Identification of protein coding regions by database similarity
search. *Nat Genet,* 3, 266-72.
GU, Z. 2022. Complex Heatmap Visualization. *iMeta*.
HERNANDEZ-GARCIA, J., BRIONES-MORENO, A. & BLAZQUEZ, M. A. 2021. Origin and evolution
of gibberellin signaling and metabolism in plants. *Semin Cell Dev Biol,* 109, 46-54.
HEYMAN, J., CANHER, B., BISHT, A., CHRISTIAENS, F. & DE VEYLDER, L. 2018. Emerging role of the
plant ERF transcription factors in coordinating wound defense responses and repair. *J*
Cell Sci, 131.
HOANG, D. T., CHERNOMOR, O., VON HAESELER, A., MINH, B. Q. & VINH, L. S. 2018. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution,* 35,
518-522.
HUSON, D. H., BEIER, S., FLADE, I., GÓRSKA, A., EL-HADIDI, M., MITRA, S., RUSCHEWEYH, H. J. &
TAPPU, R. 2016. MEGAN Community Edition - Interactive Exploration and Analysis of
Large-Scale Microbiome Sequencing Data. *Plos Computational Biology,* 12.
ISHIZAKI, K., CHIYODA, S., YAMATO, K. T. & KOHCHI, T. 2008. Agrobacterium-mediated
transformation of the haploid liverwort Marchantia polymorpha L., an emerging model
for plant biology. *Plant Cell Physiol,* 49, 1084-91.
IWAMA, R. E. & MORAN, Y. 2023. Origins and diversification of animal innate immune responses
against viral infections. *Nat Ecol Evol,* 7, 182-193.
JADA, B., SOITAMO, A. J., SIDDIQUI, S. A., MURUKESAN, G., ARO, E. M., SALAKOSKI, T. & LEHTO,
K. 2014. Multiple different defense mechanisms are activated in the young transgenic
tobacco plants which express the full length genome of the Tobacco mosaic virus, and
are resistant against this virus. *PLoS One,* 9, e107778.
JEON, H.-W., IWAKAWA, H., MARAMOTO, S., HERRFURTH, C., GUSTCHE, N., SCHLÜTER, T.,
KYOZUKA, J., MIYAUCHI, S., FEUSSNER, I., ZACHGO, S. & NAKAGAMI, H. 2022. Contrasting
and conserved roles of NPR pathways in diverged land plant lineages. *bioRxiv*.
JIA, X., WANG, L., ZHAO, H., ZHANG, Y., CHEN, Z., XU, L. & YI, K. 2023. The origin and evolution of
salicylic acid signaling and biosynthesis in plants. *Mol Plant,* 16, 245-259.
KALYAANAMOORTHY, S., MINH, B. Q., WONG, T. K. F., VON HAESELER, A. & JERMIIN, L. S. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature*
Methods, 14, 587-+.

KAPPAGANTU, M., COLLUM, T. D., DARDICK, C. & CULVER, J. N. 2020. Viral Hacks of the Plant
Vasculature: The Role of Phloem Alterations in Systemic Virus Infection. *Annu Rev Virol,*
7**,** 351-370.
KASSCHAU, K. D., XIE, Z., ALLEN, E., LLAVE, C., CHAPMAN, E. J., KRIZAN, K. A. & CARRINGTON, J.
C. 2003. P1/HC-Pro, a viral suppressor of RNA silencing, interferes with Arabidopsis
development and miRNA unction. *Dev Cell,* 4**,** 205-17.
KATOH, K. & STANDLEY, D. M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Molecular Biology and Evolution,* 30**,** 772-
780.
KNEESHAW, S., SORIANO, G., MONTE, I., HAMBERG, M., ZAMARRENO, A. M., GARCIA-MINA, J.
M., FRANCO-ZORRILLA, J. M., KATO, N., UEDA, M., REY-STOLLE, M. F., BARBAS, C.,
MICHAVILA, S., GIMENEZ-IBANEZ, S., JIMENEZ-ALEMAN, G. H. & SOLANO, R. 2022.
Ligand diversity contributes to the full activation of the jasmonate pathway in
Marchantia polymorpha. *Proc Natl Acad Sci U S A,* 119**,** e2202930119.
KOPYLOVA, E., NOÉ, L. & TOUZET, H. 2012. SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data. *Bioinformatics,* 28**,** 3211-3217.
KORNER, C. J., KLAUSER, D., NIEHL, A., DOMINGUEZ-FERRERAS, A., CHINCHILLA, D., BOLLER, T.,
HEINLEIN, M. & HANN, D. R. 2013. The immunity regulator BAK1 contributes to
resistance against diverse RNA viruses. *Mol Plant Microbe Interact,* 26**,** 1271-80.
KRAMER, S. R., GOREGAOKER, S. P. & CULVER, J. N. 2011. Association of the Tobacco mosaic virus
126kDa replication protein with a GDI protein affects host susceptibility. *Virology,* 414**,**
110-8.
KULSHRESTHA, S., JIBRAN, R., VAN KLINK, J. W., ZHOU, Y., BRUMMELL, D. A., ALBERT, N. W.,
SCHWINN, K. E., CHAGNE, D., LANDI, M., BOWMAN, J. L. & DAVIES, K. M. 2022. Stress,
senescence, and specialized metabolites in bryophytes. *J Exp Bot,* 73**,** 4396-4411.
LEE, W. S., FU, S. F., LI, Z., MURPHY, A. M., DOBSON, E. A., GARLAND, L., CHALUVADI, S. R., LEWSEY,
M. G., NELSON, R. S. & CARR, J. P. 2016. Salicylic acid treatment and expression of an
RNA-dependent RNA polymerase 1 transgene inhibit lethal symptoms and meristem
invasion during tobacco mosaic virus infection in Nicotiana benthamiana. *BMC Plant*
Biol, 16**,** 15.
LEEK, J. T., JOHNSON, W. E., PARKER, H. S., FERTIG, E. J., JAFFE, A.E., ZHANG, Y., STOREY, J. D. &
TORRES, L. C. 2023. sva: Surrogate Variable Analysis.
LETUNIC, I. & BORK, P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. *Nucleic Acids Research,* 49**,** W293-W296.
LEWSEY, M. G. & CARR, J. P. 2009. Effects of DICER-like proteins 2, 3 and 4 on cucumber mosaic
virus and tobacco mosaic virus infections in salicylic acid-treated plants. *J Gen Virol,* 90**,**
3010-3014.
LIANG, Y., HEYMAN, J., XIANG, Y., VANDENDRIESSCHE, W., CANHER, B., GOEMINNE, G. & DE
VEYLDER, L. 2022. The wound-activated ERF15 transcription factor drives Marchantia
polymorpha regeneration by activating an oxylipin biosynthesis feedback loop. *Sci Adv,*
8**,** eabo7737.
LIM, P. O. & NAM, H. G. 2005. The molecular and genetic control of leaf senescence and longevity
in Arabidopsis. *Curr Top Dev Biol,* 67**,** 49-83.
LIU, C. & NELSON, R. S. 2013. The cell biology of Tobacco mosaic virus replication and movement.
Front Plant Sci, 4**,** 12.
LIU, Y., GAO, Q., WU, B., AI, T. & GUO, X. 2009. NgRDR1, an RNA-dependent RNA polymerase
isolated from Nicotiana glutinosa, was involved in biotic and abiotic stresses. *Plant*
Physiol Biochem, 47**,** 359-68.

LOPEZ-GOMOLLON, S. & BAULCOMBE, D. C. 2022. Roles of RNA silencing in viral and non-viral
plant immunity and in the crosstalk between disease resistance systems. *Nat Rev Mol*
Cell Biol, 23**,** 645-662.
LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol,* 15**,** 550.
LUCAS, W. J., GROOVER, A., LICHTENBERGER, R., FURUTA, K., YADAV, S. R., HELARIUTTA, Y., HE, X.
Q., FUKUDA, H., KANG, J., BRADY, S. M., PATRICK, J. W., SPERRY, J., YOSHIDA, A., LOPEZ-
MILLAN, A. F., GRUSAK, M. A. & KACHROO, P. 2013. The plant vascular system: evolution,
development and functions. *J Integr Plant Biol,* 55**,** 294-388.
MANDADI, K. K. & SCHOLTHOF, K. B. 2013. Plant immune responses against viruses: how does a
virus cause disease? *Plant Cell,* 25**,** 1489-505.
MATSUI, H., IWAKAWA, H., HYON, G. S., YOTSUI, I., KATOU, S., MONTE, I., NISHIHAMA, R.,
FRANZEN, R., SOLANO, R. & NAKAGAMI, H. 2020. Isolation of Natural Fungal Pathogens
from Marchantia polymorpha Reveals Antagonism between Salicylic Acid and Jasmonate
during Liverwort-Fungus Interactions. *Plant Cell Physiol,* 61**,** 265-275.
MONTE, I., FRANCO-ZORRILLA, J. M., GARCIA-CASADO, G., ZAMARRENO, A. M., GARCIA-MINA, J.
M., NISHIHAMA, R., KOHCHI, T. & SOLANO, R. 2019. A Single JAZ Repressor Controls the
Jasmonate Pathway in Marchantia polymorpha. *Mol Plant,* 12**,** 185-198.
MONTE, I., ISHIDA, S., ZAMARRENO, A. M., HAMBERG, M., FRANCO-ZORRILLA, J. M., GARCIA-
CASADO, G., GOUHIER-DARIMONT, C., REYMOND, P., TAKAHASHI, K., GARCIA-MINA, J.
M., NISHIHAMA, R., KOHCHI, T. & SOLANO, R. 2018. Ligand-receptor co-evolution shaped
the jasmonate pathway in land plants. *Nat Chem Biol,* 14**,** 480-488.
MUSHEGIAN, A., SHIPUNOV, A. & ELENA, S. F. 2016. Changes in the composition of the RNA
virome mark evolutionary transitions in green plants. *BMC Biol,* 14**,** 68.
NGUYEN, L. T., SCHMIDT, H. A., VON HAESELER, A. & MINH, B. Q. 2015. IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.
Molecular Biology and Evolution, 32**,** 268-274.
NIEHL, A., WYRSCH, I., BOLLER, T. & HEINLEIN, M. 2016. Double-stranded RNAs induce a pattern-
triggered immune signaling pathway in plants. *New Phytol,* 211**,** 1008-19.
PATRO, R., DUGGAL, G., LOVE, M. I., IRIZARRY, R. A. & KINGSFORD, C. 2017. Salmon provides fast
and bias-aware quantification of transcript expression. *Nature Methods,* 14**,** 417-+.
PEART, J. R., COOK, G., FEYS, B. J., PARKER, J. E. & BAULCOMBE, D. C. 2002. An EDS1 orthologue
is required for N-mediated resistance against tobacco mosaic virus. *Plant J,* 29**,** 569-79.
PRUITT, R. N., LOCCI, F., WANKE, F., ZHANG, L., SAILE, S. C., JOE, A., KARELINA, D., HUA, C.,
FROHLICH, K., WAN, W. L., HU, M., RAO, S., STOLZE, S. C., HARZEN, A., GUST, A. A.,
HARTER, K., JOOSTEN, M., THOMMA, B., ZHOU, J. M., DANGL, J. L., WEIGEL, D.,
NAKAGAMI, H., OECKING, C., KASMI, F. E., PARKER, J. E. & NURNBERGER, T. 2021. The
EDS1-PAD4-ADR1 node mediates Arabidopsis pattern-triggered immunity. *Nature,* 598**,**
495-499.
QUILLIAM, R. S., SWARBRICK, P. J., SCHOLES, J. D. & ROLFE, S. A. 2006. Imaging photosynthesis in
wounded leaves of Arabidopsis thaliana. *J Exp Bot,* 57**,** 55-69.
RAKHSHANDEHROO, F., REZAEE, S. & PALUKAITIS, P. 2017. Silencing the tobacco gene for RNA-
dependent RNA polymerase 1 and infection by potato virus Y cause remodeling of
cellular organelles. *Virology,* 510**,** 127-136.
REDKAR, A., GIMENEZ IBANEZ, S., SABALE, M., ZECHMANN, B., SOLANO, R. & DI PIETRO, A. 2022.
Marchantia polymorpha model reveals conserved infection mechanisms in the vascular
wilt fungal pathogen Fusarium oxysporum. *New Phytol,* 234**,** 227-241.
SALGUERO-LINARES, J., SERRANO, I., RUIZ-SOLANI, N., SALAS-GOMEZ, M., PHUKAN, U. J.,
GONZALEZ, V. M., BERNARDO-FAURA, M., VALLS, M., RENGEL, D. & COLL, N. S. 2022.

Robust transcriptional indicators of immune cell death revealed by spatiotemporal
transcriptome analyses. *Mol Plant,* 15**,** 1059-1075.
SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M., PIETZSCH, T.,
PREIBISCH, S., RUEDEN, C., SAALFELD, S., SCHMID, B., TINEVEZ, J. Y., WHITE, D. J.,
HARTENSTEIN, V., ELICEIRI, K., TOMANCAK, P. & CARDONA, A. 2012. Fiji: an open-source
platform for biological-image analysis. *Nat Methods,* 9**,** 676-82.
SCHOLTHOF, K. B. 2004. Tobacco mosaic virus: a model system for plant biology. *Annu Rev*
Phytopathol, 42**,** 13-34.
SCHOMMER, C., PALATNIK, J. F., AGGARWAL, P., CHETELAT, A., CUBAS, P., FARMER, E. E., NATH,
U. & WEIGEL, D. 2008. Control of jasmonate biosynthesis and senescence by miR319
targets. *PLoS Biol,* 6**,** e230.
SCHWACH, F., VAISTIJ, F. E., JONES, L. & BAULCOMBE, D. C. 2005. An RNA-dependent RNA
polymerase prevents meristem invasion by potato virus X and is required for the activity
but not the production of a systemic silencing signal. *Plant Physiol,* 138**,** 1842-52.
SHENG, Y., YANG, L., LI, C., WANG, Y. & GUO, H. 2019. Transcriptomic changes in Nicotiana
tabacum leaves during mosaic virus infection. *3 Biotech,* 9**,** 220.
SONESON, C., LOVE, M. I. & ROBINSON, M. D. 2015. Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Res,* 4**,** 1521.
VASSEUR, F., BALDRICH, P., JIMENEZ-GONGORA, T., VILLAR-MARTIN, L., WEIGEL, D. & RUBIO-
SOMOZA, I. 2022. miRNA472 deficiency enhances Arabidopsis thaliana defence without
reducing seed production. *bioRxiv,* doi.org/10.1101/2022.12.13.520224
WANG, L., WAN, M. C., LIAO, R. Y., XU, J., XU, Z. G., XUE, H. C., MAI, Y. X. & WANG, J. W. 2023. The
maturation and aging trajectory of Marchantia polymorpha at single-cell resolution. *Dev*
Cell, 58**,** 1429-1444 e6.
WANG, X., GOREGAOKER, S. P. & CULVER, J. N. 2009. Interaction of the Tobacco mosaic virus
replicase protein with a NAC domain transcription factor is associated with the
suppression of systemic host defenses. *J Virol,* 83**,** 9720-30.
WHITHAM, S., DINESH-KUMAR, S. P., CHOI, D., HEHL, R., CORR, C. & BAKER, B. 1994. The product
of the tobacco mosaic virus resistance gene N: similarity to toll and the interleukin-1
receptor. *Cell,* 78**,** 1101-15.
XU, T., ZHANG, L., ZHEN, J., FAN, Y., ZHANG, C. & WANG, L. 2013. Expressional and regulatory
characterization of Arabidopsis RNA-dependent RNA polymerase 1. *Planta,* 237**,** 1561-9.
YAMANAKA, T., OHTA, T., TAKAHASHI, M., MESHI, T., SCHMIDT, R., DEAN, C., NAITO, S. &
ISHIKAWA, M. 2000. TOM1, an Arabidopsis gene required for efficient multiplication of
a tobamovirus, encodes a putative transmembrane protein. *Proc Natl Acad Sci U S A,* 97**,**
10107-12.
YING, X. B., DONG, L., ZHU, H., DUAN, C. G., DU, Q. S., LV, D. Q., FANG, Y. Y., GARCIA, J. A., FANG,
R. X. & GUO, H. S. 2010. RNA-dependent RNA polymerase 1 from Nicotiana tabacum
suppresses RNA silencing and enhances viral infection in Nicotiana benthamiana. *Plant*
Cell, 22**,** 1358-72.
YOTSUI, I., MATSUI, H., MIYAUCHI, S., IWAKAWA, H., MELKONIAN, K., SCHLUTER, T., MICHAVILA,
S., KANAZAWA, T., NOMURA, Y., STOLZE, S. C., JEON, H. W., YAN, Y., HARZEN, A., SUGANO,
S. S., SHIRAKAWA, M., NISHIHAMA, R., ICHIHASHI, Y., IBANEZ, S. G., SHIRASU, K., UEDA,
T., KOHCHI, T. & NAKAGAMI, H. 2023. LysM-mediated signaling in Marchantia
polymorpha highlights the conservation of pattern-triggered immunity in land plants.
Curr Biol, 33**,** 3732-3746 e8.
YU, D., FAN, B., MACFARLANE, S. A. & CHEN, Z. 2003. Analysis of the involvement of an inducible
Arabidopsis RNA-dependent RNA polymerase in antiviral defense. *Mol Plant Microbe*
Interact, 16**,** 206-16.

YU, X., XU, Y. & YAN, S. 2021. Salicylic acid and ethylene coordinately promote leaf senescence. *J*
Integr Plant Biol, 63**,** 823-827.
ZHU, F., XI, D. H., YUAN, S., XU, F., ZHANG, D. W. & LIN, H. H. 2014. Salicylic acid and jasmonic
acid are essential for systemic resistance against tobacco mosaic virus in Nicotiana
benthamiana. *Mol Plant Microbe Interact,* 27**,** 567-77.

1021

a

b

c

A

Name	ID	Relative expression level
MpPR9	Mp521510	4.04
MpSYP13B	Mp2g21910	2.34
MpCML42	Mp6g18000	1.79
MpWRKY22	Mp7g17200	1.74
MpACRE132	Mp1g27170	1.22

B

Name	ID	Relative expression level
MpERF1	Mp7g09350	4.24
MpERF16	Mp4g22290	3.11
MpERF10	Mp7g00860	2.83
MpERF21	Mp5g22160	1.68
MpERF8	Mp6g18650	1.48
MpERF18	Mp5g12480	1.14
MpERF15	Mp7g09350	1.06
MpAP2L1	Mp8g11450	-1.62
MpERF17	Mp4g22280	-2.38
MpERF12	Mp6g06870	-2.95
MpERF13	Mp6g08690	-3.25

C

D

Name	ID	Relative expression level
MpWRKY7	Mp3g17660	3.56
MpWRKY6	Mp1g08960	2.52
MpWRKY12	Mp4g00200	1.89
MpWRKY10	Mp7g06550	1.79
MpWRKY9	Mp7g17200	1.74
MpWRKY13	Mp4g00180	1.64
MpWRKY1	Mp7g00010	1.28
MpWRKY14	Mp6g16800	1.16

E

F

G

A

Nicotiana benthamiana   *Marchantia polymorpha*

B

independent t-test, p = 1.07e-04

Area (cm^2)

P35S:p126-citrine   Tak 1

C

-AD/-BD	L⁻W⁻H⁻	L⁻W⁻
-/p126		
MpGDI2/p126		
MpTOM1/p126		
MpNAC7/p126		
-/Δp126		
MpGDI2/Δp126		
MpNAC7/Δp126		
MpTOM1/Δp126		

D

-N-Luc/C-Luc-	TMV	TuMV
MpNAC7/AtBOB1		
AtAGO1/p126		
AtAGO1/Δp126		
MpNAC7/p126		
MpNAC7/Δp126		

High
Low

E

TMV-ROS1/GUS

TMV-ROS1/*MpNac7*

Inoculated leaves   Systemic leaves