Conservation of molecular responses upon viral infection in the non-vascular plant *Marchantia polymorpha*

Eric Ros-Moner [1*], Tamara Jiménez-Góngora[1*], Luis Villar-Martín[1], Lana Vogrinec[2,3], Víctor M. González-Miguel[4], Denis Kutnjak[2] and Ignacio Rubio-Somoza[1,5]

[1]Molecular Reprogramming and Evolution (MoRE) Laboratory, Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus UAB-Edifci CRAG. 08193 Cerdanyola del Vallés, Spain.

[2]National Institute of Biology, Department of Biotechnology and Systems Biology, Ljubljana, Slovenia.

[3]Jožef Stefan International Postgraduate School, Ljubljana, Slovenia.

[4]Data Analysis area, Bioinformatics Core Unit, Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus UAB-Edifci CRAG. 08193 Cerdanyola del Vallés, Spain.

[5]Consejo Superior de Investigaciones Científicas (CSIC), Barcelona 08001, Spain.

*Those authors contributed equally to this work

To whom correspondence should be addressed: Ignacio.rubio@cragenomica.es

Abstract

After plants transitioned from water to land around 450 million years ago, they faced novel pathogenic microbes. Their colonization of diverse habitats was driven by anatomical innovations like roots, stomata, and vascular tissue, which became central to plant-microbe interactions. However, the impact of these innovations on plant immunity and pathogen infection strategies remains poorly understood. Here, we explore plant-virus interactions in the bryophyte *Marchantia polymorpha* to gain insights into the evolution of these relationships. Virome analysis reveals that *Marchantia* is predominantly associated with RNA viruses. Comparative studies with tobacco mosaic virus (TMV) show that *Marchantia* shares core defense responses with vascular plants but also exhibits unique features, such as a sustained wound response preventing viral spread. Additionally, general defense responses in Marchantia are equivalent to those restricted to vascular tissues in Nicotiana, suggesting that evolutionary acquisition of developmental innovations results in re-routing of defense responses in vascular plants.

Introduction

It is estimated that plants developed a functional vascular system through repurposing of pre-existing genes around 420 million years ago, which facilitated the growth of larger plants by providing mechanical support and enabling long distant transport of water, nutrients, and signalling molecules (Bowles et al., 2022, Lucas et al., 2013). Viruses are obligate parasites that can infect virtually any living organism hijacking central host cellular functions to replicate and spread within and intra hosts. Coincidentally with land colonization, the repertoire of viruses (virome) associated to plants enlarged and diversified both in nature and infection strategies. Although our current knowledge lacks information on virome composition at intermediately divergent plant lineages, it appears that viruses with large DNA genomes majorly infecting unicellular chlorophyte algae were displaced by an increasing population of viruses with RNA and small DNA genomes found in vascular plants (Mushegian et al., 2016). Plant vasculature has become central for the success of viral infections in tracheophytes, not only serving as fast track for viral systemic spread, but also providing a unique cellular environment for some viruses which replication is restricted to that tissue (Kappagantu et al., 2020). Accordingly, vascular cells are thought to present specific responses to viral infection when compared to other leaf cell-types (Collum and Culver, 2017).

Our current understanding of the molecular events governing plant-virus interactions is vastly shaped by studies focusing on vascular plants, mainly crops. Plants respond to viral infections via at least three different but interconnected defense mechanisms (Mandadi and Scholthof, 2013). Firstly, plants have an extracellular surveillance system that monitors the presence of pathogens through the recognition of highly conserved microbe patterns dubbed as Pattern-associated molecular patterns (PAMPs), such as viral derived double-strand RNAs (dsRNAS), the bacterial flagellin or the fungal chitin. Detection of those molecules by specialized surface-localized receptor kinases (Pattern recognition receptors, PRRs), triggers a first layer of defense called PAMP-triggered immunity (PTI). PRRs expression is often restricted to focal points for microbial interaction and invasion (Beck et al., 2014, Yotsui et al., 2023). Two additional surveillance mechanisms monitor the intracellular presence of microbe-derived molecules. Highly stable dsRNAs produced by viruses during their life cycle, independently of the DNA or RNA nature of their genomes, are detected by the RNA silencing machinery which constitutes a conserved antiviral defense mechanism both in plants and animals (Ding et al., 2018, Ding and Voinnet, 2007). Additionally, RNA silencing machinery also recognizes dsRNA generated from the

conversion of exogenous and endogenous single strand RNAs (ssRNA) by RNA-dependent RNA polymerases (RDRs). RDR1 and RDR6 have been involved in antiviral defense in *Arabidopsis thaliana* and *Nicotiana tabacum* (Garcia-Ruiz et al., 2010, Rakhshandehroo et al., 2017, Ying et al., 2010), even though *Nicotiana benthamiana* has only a functional RDR6 (Bally et al., 2015). Upon recognition, viral dsRNAs are trimmed in small RNAs (visRNAs) of 21 to 24 nucleotides in length by members of the DICER-like family, mainly DCL2, DCL3 and DCL4 in Arabidopsis and Nicotiana (Garcia-Ruiz et al., 2010, Lewsey and Carr, 2009). The resulting visRNA duplexes are subsequently loaded into proteins from the ARGONAUTE family, which are the core elements of the RNA-induced silencing complex (RiSC), and used to scan the intracellular space for the source of dsRNA precursors in a sequence complementary manner abrogating viral infection (Lopez-Gomollon and Baulcombe, 2022). In turn, viruses produce countermeasures, known as silencing suppressors, targeted to interfere with different steps of the host RNA silencing machinery (Lopez-Gomollon and Baulcombe, 2022). In response to these silencing suppressors, host micro RNAs (miRNA, one of the two main classes of endogenous sRNAs in plants) are no longer functional, allowing the translation of their targets and licensing a new layer of defense. Among the miRNA targets involved in such counter-counter measure, members of the Nucleotide-binding domain leucine-rich repeat (NLR) family are central. Thus, it has been suggested that host miRNAs might work as sensors for the presence of pathogen-derived silencing suppressors enabling a rapid host defense response based on the activation of otherwise silenced miRNA targets (Vasseur et al., 2022, Silvestri et al., 2024). NLRs are core elements of the third intracellular surveillance system for directly monitoring the intracellular presence of pathogen-derived molecules, such as the prototypical case of the tobacco mosaic virus (TMV) replicase by the gene *N* in tobacco (Erickson et al., 1999), or their impact on host cellular homeostasis. NLR-activation triggers the so-called Effector-triggered immunity (ETI), which normally leads to a hypersensitive response (HR) culminating in cell death. Those three defense systems, PTI, RNA silencing and ETI largely rely on the action of the plant hormone Salicylic acid (SA). PTI and ETI responses encompass activation of SA signalling (Pruitt et al., 2021) while SA activates RDR1 in Arabidopsis and Nicotiana (Lee et al., 2016, Ying et al., 2010, Yu et al., 2003). Additionally, RDR1 expression is also induced by treatment with jasmonates (Liu et al., 2009, Xu et al., 2013), another central hormone in plant defense. Many of those immunity components and networks were already present in the last common land plant ancestor, while others have been acquired alongside the increasing plant cellular and anatomical complexity (de Vries et al., 2018, Belanger et al., 2023, Chia et al., 2024b).

A central question in plant-microbe interactions is how the evolution of different plant

developmental trajectories, including the acquisition of major innovations, impacts host defense

and pathogen infection strategies. In animals, comparative studies have proved very informative

about the origin and diversification of their innate immune responses against virus (Iwama and

Moran, 2023). Likewise, comparative studies including non-vascular plants, like the liverwort

Marchantia polymorpha, have been instrumental in understanding the evolution of the

interactions between plants and pathogens other than viruses (Carella et al., 2018, Gimenez-

Ibanez et al., 2019, Redkar et al., 2022). Comparative studies in plants have traditionally relied

on both model plant species and microbes, despite the lack of prior experimental evidence for

those interactions to occur in nature. Incorporating bryophytes into those comparative studies

not only provides important information on how plant immune programs may have evolved, but

it also bears the potential to determine the contribution of developmental innovations, such as

roots, stomata or vasculature, to their diversification.

To enable comparative studies towards understanding the evolution of plant-virus interactions,

we here characterized the molecular interplay between the non-vascular liverwort *Marchantia*

polymorpha and viruses. We firstly defined the composition of the virome of Marchantia plants

living in the wild and used that information to select a virus with similar genomic features to

those found in nature for further molecular characterization, and which could be tagged with a

fluorescent protein that enables monitoring of infection. Additionally, the selected virus should

be able to infect a broad range of hosts, in which their molecular interactions have been studied.

Following those criteria, we focused on the first ever isolated virus, the tobacco mosaic virus

(TMV). TMV is a ssRNA virus from the Tobamovirus genus showing a broad host range in nature

(Zamfir et al., 2023). Additionally, TMV has been extensively used for studying plant-virus

interactions in a wealth of plant species, including several Solanaceae species, like tomato and

tobacco, and the model plant *Arabidopsis thaliana*, which provide strong foundation for

comparative studies. TMV infects through wound entry points from where it moves cell-to-cell

via plasmodesmata, until it reaches the vascular system for systemic infection of distal parts of

the plant following the stream of photoassimilates (Liu and Nelson, 2013).

Our results show that *Marchantia polymorpha* is associated with RNA viruses in natural

populations that can replicate in lab grown Marchantia plants. Additionally, TMV can replicate in

Marchantia plants triggering a conserved transcriptional reprogramming encompassing

increased wounding and PTI responses in detriment of photosynthesis and cell cycle progression

and leading to changes in life-history traits, such as aging and senescence. Induction of a wealth

of NLR proteins and core elements from the RNA silencing machinery were also observed,

including DCL4 that, in absence of a DCL2 gene in the Marchantia genome (Belanger et al., 2023),

seems to be the main antiviral DCL protein according to the predominant 21 nuleotides in length

presented by TMV-derived vsRNAs. Noteworthy, the general response of the RNA silencing

machinery observed in Marchantia upon TMV infection is equivalent to that found specifically in

vascular tissues of *Nicotiana benthamiana*, suggesting an evolutionary anatomical rerouting of

antiviral defense towards newly acquired focal points for viral infection.

We additionally identified a conserved host interactor with the TMV silencing suppressor p126

that triggers cross-species defense against TMV.

Results

***Marchantia polymorpha* is associated to RNA viruses in nature.**

To identify viruses associated with *Marchantia polymorpha* in nature, we performed a

virome analysis of two composite plant samples including 21 individuals growing naturally, not

cultivated, in 5 different locations spanning 4 different countries and accessible to possible viral

vectors. First included Marchantia individuals obtained from three open air botanical gardens

throughout Europe: Tübingen (Germany), Bern (Switzerland) and Cambridge (UK). The second

encompassed individuals growing open air at the botanical garden in Osnabrück (Germany) and

individuals growing in urban sites in Ljubljana (Slovenia). In addition to the expected background

of bacteriophage sequences, we detected several known viruses, including some capable of

infecting or being associated with plants, fungi or invertebrates (Supplementary Data 1).

Putatively new virus-like sequences, possibly associated with *M. polymorpha*, were selected for

further analyses from the first dataset, based on i) their relatively high average sequencing

depths and ii) similarity to viruses previously associated with eukaryotic organisms, including

plants and invertebrates (Supplementary Data 2). Two previously described and three newly

identified viruses were hence selected for further RT-qPCR confirmation of their association with

the different *M. polymorpha* individuals, along with plants from the Tak-1 reference accession

grown in sterile conditions. Specific primers to detect Ligustrum necrotic ringspot virus (LNRV,

known to infect plants), Hubei picorna-like virus 51, (detected in a metagenomics study and

putatively associated with diverse sample types by datamining – see results below) a newly

identified picorna-like virus (which we named Tübingen picorna-like virus 1, with average

mapping depth of 2230x and contig length of 7801 nts), a new tombus-like virus (Tübingen

tombus-like virus 1, with average mapping depth of 218x and contig length of 4521 nts), and a

new sobemo-like virus (Bern sobemo-like virus 1, with average mapping depth of 88x and contig

length of 2569 nts) were used for these analysis. While none of those viruses were detected in

negative control Tak-1 plants, individuals from Tübingen were associated with all tested viruses

but the Bern sobemo-like virus 1, which in turn was present in individuals from Bern along with

Tübingen picorna-like virus 1. Individuals from the Cambridge population were associated with

Hubei picorna-like virus 51 and Tübingen picorna-like virus 1 (Supplementary Data 3). Further,

we used a datamining approach to search for the presence of conserved RdRp domain

fingerprints (palmprints) of the selected viruses in global short reads sequence datasets using

Serratus infrastructure (Edgar et al., 2022). The results of the analysis suggested that palmprint

RdRp sequences with high similarity to those of queried viruses, can be associated with diverse

sample types, mostly including environmental samples, such as wastewater, water, rhizosphere,

soil and plants (Supplementary Data 4).

Given that Tübingen picorna-like virus 1 was present in all tested populations from the first

sampling, we decided to further characterize its genomic features and interaction with

Marchantia polymorpha. Indeed, Tübingen picorna-like virus 1 had the highest mapping depth

of all contigs in our virome analysis and showed a similar genome structure as other unclassified

picorna-like viruses (Shi et al., 2016), consisting of two predicted genome open reading frames

(ORFs), first containing domains with homology to superfamily 3 helicase, peptidase C3 and RNA-

dependent RNA polymerase (RdRp) and second three domains with homology to a coat protein

with 8-stranded jelly roll β-barrel motif (Figure 1A). Phylogenetic analysis (Figure 1B) of the RdRp

protein sequence and its closest NCBI GenBank homologues showed high similarity (80% identity

of aligned sequences) of this virus isolate with a recently reported plant associated picorna-like

virus (Trichosanthes kirilowii picorna-like virus, QKK82970.1). Much lower identities were

observed for Tübingen picorna-like virus 1 (29-42%) and other viruses associated with diverse

sources (plants, gastropods, water, crayfish, artropods and bird feces). However, given that they

were discovered in metagenomics studies their association with these hosts is only speculative.

Thus, we performed and additional phylogenetic analysis, comparing the RdRp sequence of

Tübingen picorna-like virus 1 with most similar RdRp sequences from RNA Viruses in

Metatranscriptomes (RVMT) database (https://riboviria.org/) (Neri et al., 2022). The analysis

revealed high similarity of the viral RdRp to several sequences included in family-level cluster

f.0032 with order *Picornavirales* (Supplementary Fig. 1A). Even if this cluster contains diverse

representatives, none of those has confirmed host associations to the date (Supplementary Data

5). Similarly, as found in Serratus analysis, sequences most similar to investigated virus were

associated, e.g., with wastewater, wetlands, and peat soil samples (Supplementary Fig. 1b,

Supplementary Data 5).

Next, we tested whether the Tübingen picorna-like virus 1 was able to replicate in *M. polymorpha*

by incubating two-weeks old Tak-1 individuals with extracts isolated from the Tübingen

population, which contained the highest levels of Tübingen picorna-like virus 1, as determined

by RT-qPCR analysis, or from Tak-1 plants grown in sterile conditions. Individual Marchantia

plants were collected one and two weeks after incubation and the levels of viral accumulation

were measured by RT-qPCR. Individuals exposed to extracts from virus-positive plants showed a

trend of increased viral levels over time, while no amplification was observed in plants incubated

with extracts from Tak-1 plants (Figure 1C). This indicated that Tübingen picorna-like virus 1 can

replicate in Marchantia.

Collectively, those results show that *Marchantia polymorpha* can be associated with RNA viruses

in nature.

Additionally, we performed data mining to search for possible DNA viruses associated with

Marchantia polymorpha. Sequencing data from 74 biological samples (obtained from natural

environments; (Beaulieu et al., 2023)) was analyzed for the presence of viral contigs. Analysis

revealed that most of the examined samples contained virus-like sequences, with the majority

exhibiting similarity to bacteriophages (Supplementary Data 6). Within the dataset we also found

virus-like sequences with similarity to members of *Phycodnaviridae,* which can infect unicellular

green algae. Since only few relatively short contig sequences for this group of viruses were

identified in different datasets, and the data was primarily not produced with microbiome study

in mind (e.g., no surface washing) the association with *M. polymorpha* with viruses from this

taxon is dubious and shall be further explored in future studies.

Tobacco mosaic virus replicates in *Marchantia polymorpha*

Since its identification in the 19[th] century, tobacco mosaic virus has been extensively used for

studying plant-virus interactions in several vascular plant species like Arabidopsis and Nicotiana

(Scholthof, 2004). As a first step for comparative studies on plant-virus interactions between

vascular and non-vascular plants, we aimed to infect *Marchantia polymorpha* plants with a GFP

labelled clone of TMV (Peart et al., 2002). Sap from leaves of healthy and infected *Nicotiana benthamiana* plants were isolated and used to incubate two-week-old Tak-1 plants grown in sterile conditions. After incubation, Marchantia plants were transferred to soil and part of the remaining sap was mechanically inoculated with an abrasive (carborundrum) in basal leaves from 1 month old Nicotiana plants, as control for its infectivity. After two weeks, Nicotiana plants showed systemic GFP signal under UV light along with symptoms of infection (Supplementary Fig. 2), whereas Tak-1 plants displayed GFP signal in discrete infection foci that increased their size over time (Figure 2A, Supplementary Fig. 3), suggestive of viral replication. This was further corroborated by RT-qPCR amplification of pooled samples including four GFP positive individuals at the first (T1) and second weeks after incubation (T2), in comparison with mock and freshly incubated (T0) plants (Figure 2B, Supplementary Fig. 4). sRNA profiling in GFP positive areas from Marchantia thalli identified TMV-derived vsRNAs enriched in 21 nucleotides long fragments, consistent with DCL4-mediated processing, and mapping to the TMV-GFP sequence (Figure 2C, Supplementary Fig. 5).

Histological sections of the infected area showed presence of the virus in all cell types, as inferred from GFP fluorescence and when compared to sections from mock treated plants, suggesting no preference for a specific cell-type during viral infection (Figure 2D).

Those results confirmed that TMV can replicate in *M. polymoprha* plants pointing to Marchantia-TMV as a suitable pathosystem for comparative studies.

TMV infection triggers profound molecular reprogramming in *M. polymoprha*

To characterize the molecular reprogramming caused by TMV infection, we performed transcriptomic assays. TMV-GFP infected tissue from 4-week-old Marchantia plants (2 weeks after inoculation) was collected in four pools including samples from 5 plants each, along with the very same area from mock treated plants as control (Supplementary Fig. 6A). RNA was isolated and the presence of TMV was confirmed by RT-qPCR in each pool (Supplementary Fig. 6B). Principal component analysis (PCA, Supplementary Fig. 6C) of the transcriptome supported the RT-qPCR results showing a clear separation of infected and mock samples. Since separation among infected samples correlated with viral RT-qPCR levels, we selected the three replicates that had the highest levels of TMV and clustered together for further analysis. A total of 1879 genes were found to be up regulated while 1645 were reduced (padjust<0.05, Log2(fold-change) =>1 or <-1) in response to TMV infection (Supplementary Data 7). Consistently with an ongoing

pathogenic interaction, we found several defense related marker genes among the differentially
expressed genes (DEG). Members of the pathogenesis-related (PR) family are prime markers of
immune responses in several plant species, including Marchantia (Carella et al., 2019, Gimenez-
Ibanez et al., 2019) and we found that the expression of several homologues showed opposite
expression profiles when compared to mock treated plants (Supplementary Table 1). Several PTI
marker genes have been identified in Marchantia based on their orthology to those established
in Arabidopsis (Gimenez-Ibanez et al., 2019) and their activation in the presence of bacterial and
fungal extracts (Gimenez-Ibanez et al., 2019, Redkar et al., 2022). We observed that a set of those
conserved PTI markers were also induced in response to TMV infection (Figure 3A). Additionally,
a significant number of intracellular receptors from the NLR family displayed upregulated
expression upon TMV infection ($p < 6.245e-05$ with those described by (Bowman et al., 2017); p
$< 1.308e-05$ with those described by (Chia et al., 2024; Supplementary Table 2). Nevertheless,
in line with the absence of TNL genes in Marchantia (Chia et al., 2024), none of those showed
significant sequence similarity to the NLR N gene triggering ETI responses in TMV infected
tobacco plants (Whitham et al., 1994). Likewise, several components from the RNA silencing
machinery upregulated in tobacco plants upon TMV infection (Sheng et al., 2019), were also
differentially expressed in response to TMV in Marchantia plants. The non-canonical MpDCL1b
(Mp6g09830), MpDCL4 (Mp7g11720), and MpRDR6 (Mp8g14970) were upregulated, whereas
MpDCL3 (Mp1g02840), MpAGO9 (Mp8g08610) and MpNRPE1b (Mp6g00230) were suppressed.

Gene Ontology (GO) analysis of the upregulated genes revealed an enrichment (padjust<0.05)
of biological processes related to reactive oxygen species, wounding, light intensity, heat
responses and responses to fungus and the presence of their constituent chitin (Supplementary
Data 8, Supplementary Fig. 7A). Photosynthesis, DNA packaging and cell cycle, microtubule
organization or regulation of salicylic acid metabolism were among the downregulated processes
(padjust<0.05; Supplementary Data 9, Supplementary Fig. 7B).

Notably, TMV is mainly transmitted through wounds and wounding response was one of the
processes to which upregulated genes belonged in the GO analysis, which prompted us analysing
this response. ERF transcription factors are central to trigger wounding responses in vascular
plants such as *A. thaliana* (Heyman et al., 2018) with a conserved set of ERF transcription factors
being also recently described to respond to wounding in *Marchantia polymorpha* plants (Liang
et al., 2022). A subset of these wounding responsive ERF transcription factors showed the same
expression pattern upon TMV infection as that observed upon intensive wounding (Figure 3B,

(Liang et al., 2022). To determine the whole transcriptional overlap between wounding and TMV

responses, we reanalysed published tissue incision transcriptomic responses (Liang et al., 2022)

using the same pipeline we used for the TMV samples. Comparison of both datasets showed a

significant overlap both in up-regulated (593 genes; $p<6.631e-235$) and down-regulated genes

(162 genes; $p<2.137e-70$; Figure 3C), suggesting that TMV infection might contribute to a

sustained wound response in Marchantia.

Chitin is one of the most abundant polysaccharides in nature and it is a primary component of

cell walls in fungi and exoskeletons of arthropods such as insects, which in turn can work as viral

vectors in a range of plants species (Campbell, 1996). According to the GO enrichment in

response to chitin and fungi, and besides the fungal responsive genes described above as part

of a conserved PTI response, we also observed upregulation of a set of fungal responsive WRKY

transcription factors (Figure 3D; (Redkar et al., 2022, Yotsui et al., 2023).

Photosynthetic deficiency is a hallmark for several stress responses, including those involving

viral infections. Accordingly, downregulation of transcripts involved in photosynthesis was one

of the main processes affected in our GO study as consequence of TMV infection. We determined

the photosynthetic efficiency in TMV infected areas from 4-weeks old Tak-1 plants after 2-weeks

of infection along with the very same area in mock treated plants. TMV infected areas showed a

significant reduction in photosynthetic competency when compared to similar areas from mock

treated plants (Figure 3E). Notably, parts of those very same plants distant to the infection foci,

showed also reduced photosynthetic capability, although to a lesser extent (Figure 3F). Those

results suggest that TMV responses might be systemic in *Marchantia polymorpha* plants.

Overall, those results show that TMV engages in a compatible interaction with *Marchantia*

polymorpha.

Salicylic and OPDA-dependent signaling pathways are active at TMV infected areas.

The plant hormones Salicylic acid (SA) and Jasmonic acid (JA) are central in defense responses

against most pathogens of vascular plants, acting both in a synergistic and antagonistic manner.

Whereas the SA biosynthesis and signalling pathway are mostly conserved in *Marchantia*

polymoprha (Jia et al., 2023), the active molecules binding to the COI1 receptor and triggering

jasmonate signaling are a group of long chain polyunsaturated fatty acids, including dinor-OPDA,

instead of JA-Ile (Kneeshaw et al., 2022, Monte et al., 2018). Therefore, we compared the SA and

jasmonate-related transcriptomic responses with those found during TMV infection. To that end,

we used the time course datasets obtained on incubation of Marchantia with 1mM SA during 2,

6 and 12 hours (Jeon et al., 2022), SA concentration shown to promote defense against bacterial

and to increase susceptibility to fungal pathogens (Jeon et al., 2022, Matsui et al., 2020).

Comparison of the DEGs identified on SA treatment showed a significant overlap with those

responding to TMV infection (Figure 4A; Supplementary Fig. 8). A total of 225 genes were

commonly up-regulated by 2 hours SA treatment or TMV (p<1.799e-100), 306 (p < 1.382e-172)

were shared after 6 hours treatment, and 92 (p < 3.580e-47) by 12 hours of SA incubation.

Likewise, TMV infection shared 47 down regulated genes (p<1.618e-14) with 2 hours, 38 (p <

2.897e-13) with the 6 hours, and 17 genes with 12 hours incubation samples (p < 1.720e-05),

showing that TMV infection activates SA-dependent transcriptional responses.

To ascertain a possible contribution of the jasmonate signalling pathway, we compared the TMV

transcriptional response with that of plants treated with dn-OPDA (Monte et al., 2018) or having

a constitutively active jasmonate pathway as a consequence of the mutation of the JAZ signalling

repressor (Mpjaz-1 KO plants; (Monte et al., 2019). A total of 143 and 81 genes were commonly

up-regulated upon TMV infection and *MpJAZ* deficiency or dn-OPDA treatment, while 238 were

downregulated in TMV infected and Mpjaz-1 KO plants (Figure 4B), suggesting that TMV infection

also activates the jasmonate signalling pathway in Marchantia. Finally, we found that 64 genes

were commonly up-regulated by TMV infection, SA 6 h treatment and absence of a functional

MpJAZ protein, while 54 genes were up-regulated by TMV infection, SA 6h and dn-OPDA

treatments (Figure 4C).

Next, we performed a GO analysis in the different sets of genes that were found to be commonly

regulated by TMV infections and each of the different hormone-related transcriptomes (i.e. Up

regulated genes by TMV and SA 6 h treatment from Figure 4A). That approach showed that the

JA pathway might contribute to the activation of wounding, reactive oxygen species and virus

responses, in addition to the catabolism of several compounds, including lipids,

phenylpropanoids and salicylic acid, in TMV infected Marchantia plants. Interestingly, some of

these processes were also commonly upregulated by TMV infection and SA treatment, including

wounding, SA and lignin catabolism among others. The SA response pathway might on the other

side also contribute to the observed transcriptional up-regulation of defense genes against

bacteria and fungi, as well as response to chitin in TMV infected plants (Figure 4D,

Supplementary Data 10). Amongst the downregulated processes by MpJAZ deficiency and TMV

infection were translational elongation, mucilage and cell wall biosynthesis, response to

fructose, photosynthesis, microtubule-based process, growth, and response to abscisic acid

(ABA; Figure 4E, Supplementary Data 10)

Furthermore, genes found to be commonly upregulated by TMV, SA and dn-OPDA treatment

were related to oxylipin and phenylalanine metabolism, wounding and drought recovery, among

other Gene ontology groups (Figure 4F; Supplementary Data 11). Meanwhile, those genes found

to be commonly upregulated by TMV infection, SA 6 h treatment and enhanced JA signalling in

Mpjaz-1 plants were related to antiviral defense and lipid catabolism, which has been also

recently found to contribute to SA activation and antiviral defense (Xu et al., 2022); Figure 4G,

Supplementary Data 11).

Collectively, those results suggest that SA and JA signalling pathways are activated in Marchantia

during TMV infection.

TMV infection triggers aging in *Marchantia polymorpha.*

Interactions with pathogens, including viruses, can result in the modification of life-history traits

and developmental trajectories both in plants and animal (Pagan et al., 2008, Sievert, 1978).

Senescence is the final stage of the life cycle of a cell, tissue or organism and it is gradually

triggered throughout aging. This process can be accelerated by the interaction with different

pathogens, including the crucifer infecting TMV-Cg strain capable of compatible interactions with

the Arabidopsis UK-4 ecotype (Espinoza et al., 2007, Lim and Nam, 2005). Wound response,

mitotic cell arrest, and reduced photosynthetic efficiency are hallmarks for plant cell senescence

(Espinoza et al., 2007, Gan, 2003). Additionally, SA and JA, along with ET, contribute to

senescence in *Arabidopsis thaliana* (Schommer et al., 2008, Yu et al., 2021). Since all those

senescence hallmarks were present in our molecular characterization of the Marchantia-TMV

interaction, we first checked the expression levels of several senescence marker genes in our

transcriptomic data (Kulshrestha et al., 2022). We found that many senescence markers, such as

MpMYB14 (Mp5g19050), MpORE1 (Mp2g07720), MpNYE1 (Mp1g17090) and MpWRKY7

(Mp3g17660), were induced upon TMV infection, hence suggesting that TMV infection could

trigger aging and tissue maturation in Marchantia. Transcriptional trajectories orchestrating

maturation and aging in Marchantia, both at organism and tissue level, have been recently

reported to be very similar through transcriptional studies over developmental time, including

individuals from different ages (spanning from 1 day after germination - D1- to 31 days old

Marchantia plants -D31) , or by analysing sections along the thalli from top to base, being slice 1

(S1) at the top, considered the youngest part of the organ, and being slice 6 (S6) at the base and

considered the oldest (Wang et al., 2023). Thus, while S1 samples were more similar to D1, those

from S5 showed more commonalities to those from D31 plants. To assay whether TMV infection

could modify the life-history of Marchantia plants we compared the transcriptional profile

obtained from GFP positive areas from 4-week-old TMV-GFP infected Marchantia plants and that

from equivalent zones from mock treated individuals with those from the maturation and

developmental trajectories. Clustering of the transcriptional profile of mock and TMV infected

plants with those datasets, based on Pearson's correlation coefficient and regression model,

showed that TMV infection promotes maturation and aging. Thus, while the transcriptome from

mock treated plants clustered with juvenile state samples (slice 2 and Day 1), that from TMV

infected plants clustered with slice 4 and day 31 of development (Figure 5A, 5B).

Those results indicated that TMV infection triggers premature tissue maturation and senescence

in Marchantia, a characteristic trait from compatible plant - virus interactions.

392 MpNAC7 is a conserved interactor of the TMV silencing suppressor p126 and enhances defense

393 against TMV in tobacco plants.

The TMV replicase p126 is essential for viral replication and movement, but it also works as a

silencing suppressor in vascular plants (Ding et al., 2004). To gain insights into its possible role

during TMV infection in Marchantia plants, we first established its subcellular localization. We

found that p126 fused to GFP accumulated within the cytoplasm (Figure 6A) consistently with

earlier studies in onion epidermal cells, where the full length p126 protein fused to GFP was

found to be cytoplasmic (dos Reis Figueira et al., 2002).

Constitutive expression of viral-derived silencing suppressors often leads to plant developmental

defects, like growth stunting (Kasschau et al., 2003). Accordingly, four-weeks old Marchantia

plants constitutively expressing p126 were smaller in size when compared to control plants

(Figure 6B, Supplementary Fig. 9).

The TMV replicase p126 is known to associate with several host factors in Arabidopsis, including

TOM1 (Yamanaka et al., 2000), GDI2 (Kramer et al., 2011) and the transcription factor ATAF2

(Wang et al., 2009). To assess whether those interactions were conserved between p126 and the

corresponding orthologs in Marchantia, we performed Y2H assays. Orthologs were identified by

blasting the Arabidopsis protein sequences at MarpolBase (https//Marchantia.info) and

PlantRegMap (http://plantregmap.gao-lab.org/). The p126 full length and helicase domain

(Δp126, nt 2442 to 3636) were fused to the GAL4 binding domain and used as baits, while the

putative interactors fused to the GAL4-activation domain, were used as preys. Notably, the only

conserved interaction was that of the ortholog of the NAC transcription factor ATAF2, MpNAC7

(Mp6g02620), with Δp126 (Figure 6C). As for Δp126-ATAF2 binding (Wang et al., 2009) Δp126-

MpNAC7 interaction could be confirmed *in planta* using bimolecular luminescence

complementation (Figure 6D). Additionally, both the helicase domain alone and the full-length

p126 were found to interact *in planta* with either NAC transcription factors. Such discrepancy

between the results obtained by the two different protein-protein assays can be potentially

attributed to the presence of additional protein partners in Nicotiana or variable conditions ,

such as intracellular pH or growth temperature, that modify protein folding allowing the

interaction with the full length p126.

In Arabidopsis, ATAF2 expression is triggered by TMV infection (Wang et al., 2009). Likewise, we

found that MpNAC7 expression is also increased on TMV infection (Log2(fold-change)= 0.991),

wounding (Log2(fold-change)= 2.67), dn-OPDA and after 6 hours of SA treatment (Log2(fold-

change)= 1,23) in Marchantia. Likewise, ATAF2 is induced by wounding, SA and JA treatments

(Delessert et al., 2005) and contributes to defense against TMV in Arabidopsis (Wang et al.,

2009). Here, to ascertain the role of MpNAC7 in defense against TMV, and rule out a possible

link with RNA silencing- mediated responses, we infected 1 month old Nicotiana plants deficient

in RDR6 (rdr6i; (Schwach et al., 2005)) with MpNAC7, or GUS as control, in the presence of TMV

or an unrelated ssRNA virus, such as turnip mosaic virus (TuMV). TMV and TuMV strains were

labelled with the ROS1 gene which triggers anthocyanin production, and hence we followed

presence of the viruses by the purple colour (Bedoya et al., 2012). We observed that co-

expression of MpNAC7 abrogated local and systemic TMV infection with respect to plants co-

expressing the GUS control. By contrast, MpNAC7 co-expression was of no effect on TuMV

infected plants, that displayed a similar local and systemic presence of the virus as the GUS co-

infiltrated controls (Figure 6E, Supplementary Fig. 10).

Collectively, those results show that molecular interaction between the TMV silencing

suppressor p126 and the host NAC transcription factor is conserved in Marchantia, as well as the

role of this transcription factor in defense.

Discussion

In this work, we have characterized the interaction of the non-vascular liverwort *Marchantia*
polymorpha with viruses. First, we have defined known and unknown viruses that associate with
Marchantia. This showed that Marchantia is primarily associated with RNA viruses in nature,
validating previous predictions based on the presence of viral gene hallmarks in the
transcriptomes from the 1000 Plants Genome Project (Mushegian et al., 2016). We also showed
that at least one of these newly discovered viruses (Tübingen picorna-like virus 1) can replicate
in laboratory grown Marchantia Tak-1 plants. In-depth phylogenetic analyses placed this virus
within a larger, poorly characterised cluster composed of viruses discovered with metagenomics
approaches, mostly associated with environmental samples, within the *Picornavirales* order.
Viruses from this order infect mostly animals, but also protists and plants (Koonin et al., 2020)
Tübingen picorna-like virus 1 codes for the hallmark protein domains characteristic of
picornaviruses, but seems to lack movement protein (MP) that is present in most of the plant-
infecting viruses, with the exception of some vertically transmitted viruses, e.g., from families
Paritiviridae and *Amalgavirida (Dolja et al., 2020)*.Although kinetics of viral replication is not
linear, in our infection experiment we observed only modest virus titre increase over time, which
might be connected to limited ability of the virus to move from cell-to-cell due to the lack of MP.
Few viruses have been identified by metagenomic approaches in non-vascular plants, such as *M.*
polymorpha, some with (Debat et al., 2023, Mifsud et al., 2022) and other without MP
(Mushegian et al., 2016, Vendrell-Mir et al., 2021). On the other hand, it is predicted that
plasmodesmata have evolved along with the acquisition of vasculature to assist in the transport
of nutrients throughout the plant by most likely changing the way they control their aperture
(Benitez et al., 2018, Hernandez-Hernandez et al., 2020). Therefore, it is plausible that the way
in which viruses interact with host plasmodesmata to move from cell-to-cell has also differed
over time. Additionally, it has also been recently proposed that viruses lacking a MP can
systemically infect vascular plants exploiting host proteins that are also present in Marchantia,
such as those belonging to the Phloem Protein 2 (PP2) gene family (Ying et al., 2024). Since we
only started to uncover the *M. polymorpha* virome, and, little is known about the structure and
function of *M. polymorpha* plasmodesmata (Zeinsmaster D.D and Carothers Z.B, 1974), involved
also in interaction with MP in vascular plants, further research building on the here developed
M. polymorpha inoculation procedures, will contribute to shed a light on the repertoire of viruses
infecting non-vascular plants. A recent study proposed a horizontal transfer of MP-encoding
viruses infecting vascular plants to non-vascular plants (Butkovic et al., 2023), whereas discovery
of Tübingen picorna-like virus 1 suggests another possible route of virus emergence in non-

vascular plants. This may be linked with horizontal virus transfer from invertebrates or protists

and could, very speculatively, represent a missing link in a recently suggested evolution of single

jelly roll type CP to MP (Butkovic et al., 2023).

Second, we have found that TMV engages into a compatible interaction with Marchantia plants.

That interaction encompasses an extensive transcriptional reprogramming reminiscent of that

found in vascular plants such as Arabidopsis and Nicotiana, albeit involving also previously

undescribed processes like the activation of a sustained wounding response here found in the

Marchantia-TMV interaction. Third, we have found that the TMV p126 silencing suppressor

displays similar subcellular localization in Marchantia cells, as in onion epidermal cells (dos Reis

Figueira et al., 2002), and that it interacts with MpNAC7, a NAC transcription factor ortholog of

the Arabidopsis ATAF2 factor involved in defense against TMV (Wang et al., 2009). Additionally,

transient expression of MpNAC7 specifically abrogates TMV infection in Nicotiana plants. All

those commonalities show that core molecular events of plant-virus interactions found in

vascular plants might be also present in non-vascular plants.

TMV infected Marchantia plants undergo a transcriptional reprogramming consistent with a

pathogenic process and characterized by the activation of defense related PR proteins and a

conserved PTI response. Perception of dsRNAs has been found to trigger PTI in response to TMV

infection in Arabidopsis, which in turn contributes to antiviral defense (Korner et al., 2013).

Indeed, dsRNA-triggered PTI is dependent on the Arabidopsis PRR co-receptor SERK1 (Niehl et

al., 2016), which has a corresponding ortholog in the Marchantia genome (Mp7g09160). This

very same Marchantia gene has been suggested as the ortholog of the Arabidopsis BAK1 gene,

which is central for PTI responses against *Pseudomonas syringae* (Gimenez-Ibanez, 2019).

Mp7g09160 has been shown to interact with the *P. syringae* effector AvrPto, which abrogates

PTI responses in Arabidopsis through interaction with AtBAK1 (Gimenez-Ibanez et al., 2019)..

Thus, it will be interesting in future studies to analyse the role of Mp7g09160 in viral PTI

responses. Additionally, TMV infection triggers the expression of a significant number of NLR

genes that can potentially contribute to an ETI response. While in *Nicotiana benthamiana*, both

DCL2 and DCL4 are processing TMV RNAs into 22 and 21 nucleotides long vsRNAs (Guo and

Wong, 2017), the solely presence of DCL4 in Marchantia results in a predominant processing in

21 nucleotides long vsRNAs. We also observed changes on the expression of core genes of the

RNA silencing machinery in response to TMV infection. In contrast to the repression of elements

from the RNA-Directed DNA methylation pathway, such as MpDCL3, MpAGO9 and MpNRPE1b,

the central elements within the RNA antiviral machinery MpDCL4 and MpRDR6 were activated.

Such general switch on RNA silencing pathways is reminiscent of that found in *Nicotiana*

benthamiana in response to TMV, although in *benthamiana* it is specifically restricted to the

vasculature (Collum and Culver, 2017). These results suggest an evolutionary re-routing of

defense mechanisms after acquisition of novel tissues or cell-types that become new focal points

for plant-virus interactions.

Interestingly, our results also uncover a previously unknown role of the wound response as an

antiviral defense mechanism. Plant cells undergoing wound responses experience inhibition of

their photosynthetic ability becoming sink tissues for photoassimilates (Quilliam et al., 2006),

attracting TMV movement (Brants, 1961). This wound response might be, at least partially,

sustained by the activation of the salicylic and jasmonate pathways. Thus, it is tempting to

speculate that sustained wound response in Marchantia plants, along with local and systemic

decreased on photosynthetic capability, confine TMV infection limiting its movement throughout

the plant. Additionally, premature senescence of infected tissues might further abrogate viral

infection. Therefore, modification of host´s life-history traits, such as senescence, can constitute

a defensive mechanism in Marchantia.

Salicylic and jasmonate pathways are central for defense against several pathogens. Although

the interaction between both hormones has been traditionally considered antagonistic, recent

studies suggest that such antagonism might have undergone evolutionary fine tuning, and hence

depend on each plant-microbe interaction (de Vries et al., 2018). Antagonism of the salicylic and

jasmonate pathways has been described in defense against bacterial and fungal pathogens in

Marchantia (Gimenez-Ibanez, 2019; Matsui, 2020). Our comparative analysis of the

transcriptional changes induced in response to TMV infection and those found in response to

treatments with Salicylic and jasmonates (dn-OPDA), or in Mpjaz-1 KO plants, show that both SA

and jasmonate pathways might contribute to the activation of genes involved in

phenylpropanoid metabolism. Those findings are in line with Salicylic and jasmonates triggering

this very same pathway in the moss *Physcomitrium patens,* which in turn is responsible for the

production of defense-related compounds. (de Vries et al., 2018). Likewise, in line with their

synergistic interaction being responsible for local and systemic resistance against TMV in

Nicotiana benthamiana (Zhu et al., 2014), both salicylic and jasmonates were observed to

activate genes involved in antiviral defense responses to TMV infection in Marchantia. Finally, a

set of SA/dn-OPDA commonly upregulated genes might contribute to the above mentioned
sustained wound response, and therefore, to contain virus spread.

Our work also highlights the conservation of central players, such as MpNAC7, in the specific
defense against TMV. Future comparative research studies will help to ascertain the conservation
of the genetic networks under MpNAC7 and ATAF2 regulation both in Marchantia and
Arabidopsis.

Collectively, results from this work suggest the presence of evolutionary conserved and divergent
plant strategies to cope with viral infections. Those might include gain and loss of defense
strategies, such as those involving wound responses. Evolution of plant-virus interactions might
also encompass the gradual acquisition of new elements within the underlying gene networks,
as those physically interacting with pathogen-derived proteins. Finally, defense mechanisms
might undergo anatomical re-location to newly pathogen-targeted cell-types and tissues over
evolutionary times.

Methods

Plant growth conditions

Male *Marchantia polymorpha* plants Takaragaike-1 (Tak-1) were grown on plates with half
strength Gamborg´s B5 medium (Duchefa) containing 1% agar and 1% sucrose under continuous
light and 22ºC.

Nicotiana benthamiana plants were grown on soil under long day conditions (16h light/ 8h dark)
and 25ºC.

Arabidopsis thaliana plants were grown on soil under long day conditions and 23 ºC.

Infection assays

Frozen tissue from Marchantia plants from the Tübingen population or lab grown Tak-1 plants
was homogenized in liquid nitrogen cold mortars, resuspended in phosphate buffer pH7.2,
filtered using 70 µM filters and kept at 4ºC. Two-week old Tak-1 plants grown in sterile conditions
in plates were vacuum incubated with each of both extracts and transferred to plates containing

wet soil. Eight Tak-1 individuals exposed to each sap were collected either one or two weeks

after incubation to assess viral replication.

Two week-old *Nicotiana benthamiana* plants were infiltrated with an agrobacterium solution

(0.5 O.D) carrying the TMV-GFP clone (Peart et al., 2002). Infected Nicotiana plants were grown

for 3 additional weeks along with non-infected plants and upper leaves displaying infection

symptoms were collected, along with equivalent ones from non-infected plants and frozen in

liquid nitrogen. Subsequently, leaf tissue was grinded in a liquid nitrogen cold mortar. Sap was

isolated from grinded leaf tissue using phosphate buffer pH 7.2, filtered with 70 µM filters and

maintained at 4ºC. Two-weeks old Tak-1 plants were incubated either in TMV-GFP containing sap

or sap from non-infected plants (mock) for 5 minutes in vacuum. After incubation, Tak-1 plants

were transferred to plates containing wet soil and brought back to the growth chamber. GFP

signal was monitored 1 and 2 weeks after incubation using a portable UV lamp or with a

magnifying stereoscope couple to a UV light source.

RNA isolation and expression analysis

Total RNA was extracted from Marchantia plants and Arabidopsis inflorescences using the

protocol described in (Box et al., 2011). Marchantia isolated RNA was further cleaned using the

RNeasy Kit (Qiagen). 500 ngs of total RNA were DNase I treated (Fisher Scientific) and converted

into cDNA using the RevertAid first strand kit (Fisher Scientific) following the manufacturer

instructions. RT-qPCR experiments were performed in a lightcycler (ROCHE) device. Relative

expression changes were calculated using the $2^{-\Delta\Delta Ct}$ method using *MpACTIN7* (Mp6g11010) gene

as housekeeping. Primers utilized can be found in supplementary table 3.

Samples for transcriptome assays were obtained from GFP positive areas collected from 5 4-

week-old Marchantia plants per replicate (4 replicates total) after two weeks incubation with

extracts from Nicotiana plants containing TMV-GFP. Equivalent areas were collected from mock

treated 4-week-old Marchantia plants as control. RNA was

RNA-seq analysis

For virome studies, 1 µg of total RNA from each of the 21 individuals included in the study was

pooled and 4 µg from the mixture were used for preparing two independent libraries using total

ribo-depleted RNA following the manufacturer instructions (Illumina TruSeq Stranded total RNA

LT sample prep Kit, plant). The first library containing samples from 14 individuals was sequenced

on an Illumina NovaSeq 6000 device in 2x150 bp mode, while the second one containing RNA

from 7 individuals, was sequenced in an Illumina NovaseqX device. A total of 306,895,552 and

506,670,250 reads were respectively obtained. The reads were trimmed and further analysed

using QIAGEN CLC Genomics Workbench 21 (https://digitalinsights.qiagen.com/). Dataset was

analysed for the presence of known and new viruses using an established virus detection

pipeline (Pecman et al., 2017), which includes: reads trimming, *de novo* assembly of trimmed

reads, mapping of reads and contigs to NCBI viral RefSeq database (search for known viruses)

and pFam domain search for contig sequences (search for known and unknown viruses). To

detect known viruses, results of reads mapping to viral RefSeq database were inspected and

filtered for viruses for which at least 1000 nts of the reference sequence were covered by

mapped reads (Supplementary Data 1). In parallel, to detect known viruses and search for new

ones, results of the pFam domain search of contig sequences were filtered for hits to viral

proteins. Then, this candidate list of viral contigs was additionally queried for similarity against

complete NCBI *nr* database (October 2020) using blastx (Gish and States, 1993) and the similarity

search results were visualized and taxonomically classified using MEGAN 6 (Huson et al., 2016).

Contigs corresponding to cellular organisms in this analysis were excluded from the list to obtain

a final list of putative viral-like sequences (Supplementary Data 2). Trimmed sequencing reads

were mapped to the list of putative viral-like sequences, to estimate the abundance of

corresponding nucleic acids in the sequenced sample and the results were reported as average

mapping depth (number of times a position is covered by sequencing reads, averaged over a

complete contig length; Supplementary Data 2).

The presence of selected viruses (three new and two known) was tested by RT-qPCR in RNA

extracts of collected samples as described above and datamining approach was used to search

for association of those viruses with different sample types linked with publicly available short

read datasets. This was performed by querying Serratus infrastructure palmprint database

(Edgar et al., 2022) by RdRp sequences of selected viruses using palmID tool

(https://serratus.io/palmid).

For a newly discovered picorna-like virus, named Tübingen picorna-like virus 1, additional

analyses were performed, aiming to ascertain its association with *M. polymorpha*. Putative

protein domains were annotated on the viral genome by InterProScan search

(https://www.ebi.ac.uk/interpro/search/sequence/, May 2024) of the two discovered open

reading frames against all included databases. Amino acid sequence of ORF 1 of the virus

(containing RNA-dependent polymerase (RdRp) domain) was queried for similarity against

complete NCBI *nr* database (January 2022) using blastp. Fifteen best hits according to bit score

values (which have at least 33% identity with the conserved part of the RdRp domain of the

queried virus) were included into phylogenetic analysis. Alignment of the sequences of the

conserved part of RdRp was performed using MAFFT 7.48 (Katoh and Standley, 2013), followed

by alignment trimming using TrimAI 1.3 in Automated 1 mode (Capella-Gutiérrez et al., 2009).

Then, best protein substitution model was selected using Model Finder (Kalyaanamoorthy et al.,

2017) and maximum likelihood phylogenetic tree inference was performed based on the

trimmed alignment using IQ-Tree 1.6.12 (Nguyen et al., 2015) using 1000 ultrafast bootstrap

replicates (Hoang et al., 2018). The resulting phylogenetic tree was visualized using iTOL 6.5

(Letunic and Bork, 2021) and viral taxa on the tree were coloured according to the associated

sample source, as designated in NCBI GenBank. To obtain even more in-depth information about

the phylogenetic position of the newly discovered virus, its sequence was aligned also against

members of the most related family-level cluster of RdRp sequences (f.0032) from RNA Viruses

in Metatranscriptomes (RVMT) database (https://riboviria.org/) (Neri et al., 2022). Alignment

and phylogenetic analyses were performed as described above, excluding TrimAI step.

For analysing the transcriptional response in Marchantia plants to TMV infection, 1 ug of total

RNA was used for preparing each stranded library by mRNA enrichment following manufacturer

indications (TrueSeq kit from Illumina). Libraries were sequenced on a Novaseq 6000 device. A

total of 20-21 million pairs were obtained per sample. Adapter removal and trimming of low

quality bases was done using Trim Galore! v0.6.1

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), removing reads shorter

than 20 bps and non-paired reads after the trimming step. Ribosomal RNA reads were removed

using SortMeRNA v2.1b (Kopylova et al., 2012), removing also reads that become unpaired after

this filtering step.

Cleaned reads together with the transcriptome of *Marchantia polymorpha* were used to quantify

gene expression at transcript level using Salmon v1.1.0 (Patro et al., 2017).

For this, the file MpTak1v5.1_r1.mrna.fasta containing the Marchantia transcriptome (v5.1) was

downloaded from https://marchantia.info/download/tak1v5.1/ (as of 19-Jan-2021). This file was

indexed using *Salmon index* and then used as input for *Salmon quant*, which was ran with

parameters -l A, –validateMappings, --recoverOrphans, --rangeFactorizationBins 4, --seqBias,

and –gcBias. Mapping rate was found to be in the 89.5%-92.58% range.

The R progam tximport v1.14.2 (Soneson et al., 2015) was used to aggregate Salmon's transcript
expression estimates at gene level and the *DESeqDataSetFromTximport* from the R package
DESeq2 v1.26.0 (Love et al., 2014) was used to create an R object with count data and
experimental information for all samples. A low-expression filter was then applied to remove any
gene with less than 10 counts across all 8 analyzed samples, resulting in a final set of 14,980
genes (from 19,217 annotated genes). Then, the *DESeq* function, which, among other things,
computes size factors for sample normalization, was applied to the filtered object. DESeq2's *rlog*
function was used to transform count data to the log2 scale while minimizing for differences
between samples for low-expressed genes. The transformed data was fed to DESeq2's *plotPCA*
function to visualize a PCA plot of the samples.

The analysis of differential expression was done using DESeq2's *results* function with the filtered
and normalized object as input and parameter *alpha = 0.05.*

All datasets used in this study for comparative analysis (wound response and salicylic acid time
course) were analysed in the very same way as described above.

All Gene ontology analysis (GO) were performed using PLAZA
(https://bioinformatics.psb.ugent.be/plaza/). Redundancy on representative categories was
reduced using Revigo (http://revigo.irb.hr/).

For correlation analysis of transcriptome responses between developmental trajectories and
TMV response, raw reads were processed as described above, except that rRNA was removed
using ribodetector_cpu v0.2.7 (Deng et al., 2022) and Salmon version was v1.9.0. Salmon's
counts were converted into a DESeq2 object and a low-expression filter was applied as described
above. Next, the existing batch effect between time course samples and sliced tissue ones was
removed using the Combat seq function from the sva package (Leek et al., 2023), v3.48.0. Then,
values for biological replicates were averaged and Pearson correlation was computed for the
resulting count matrix and displayed using the Heatmap function from the ComplexHeatmap
package (Gu, 2022). The batch correction, correlation and heatmap plotting steps were based
on the R script used in (Wang et al., 2023) and kindly provided by the authors.

DNA-seq data mining

Data mining was utilized to search for possible DNA viruses associated with *M. polymorpha.*
DNA-seq data from a recent study of *M. polymorpha* from natural populations (Beaulieu et al.,
2023) was retrieved from the SRA database (BioProject PRJNA931118). 74 biosample accessions

were selected to encompass diverse geographical locations and various *M. polymorpha*

subspecies. Datasets were analyzed using a customized pipeline for virus detection, designed

with the Snakemake workflow management system (v.7.32.4) (Köster et al., 2021). Some of the

chosen accessions contained sequencing data from multiple runs, which were merged before

analysis. The downloaded reads were processed using a custom script: (available at:

https://github.com/LanaVogrinec/snakemake-marchantia). Briefly, reads were trimmed with

trimmomatic (v.0.39) (Bolger et al., 2014) and assembled into contigs with MEGAHIT (v.1.2.9) (Li

et al., 2015). A similarity search was performed against the NCBI *nr* database (February 2024)

using DIAMOND (v.0.9.14) blastx (Buchfink et al., 2015) and the hits were taxonomically classified

with MEGAN6 (v.6.25.9) (Huson et al., 2016). Contig sequences classified as viral (group 'Viruses'

in MEGAN analysis) were extracted and are available, together with their classifications, in

Supplementary Data 6.

sRNA library construction and data processing

TMV-GFP positive areas were collected from 2 Marchantia plants infected for 2 weeks in two

biological replicates. RNA was isolated following the protocol described in (Box et al., 2011). 10

ngs of total RNA were used to build two independent libraries following manufacturer indications

(RealSeq-AC, Somagenics). Libraries were sequenced in an Illumina NovaSeq device. A total of

26-35 million single-end reads (101 bp) were obtained per sample. Adapter removal and

trimming of low quality bases was done using Trim Galore! v0.6.1

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) with parameters *-q 15 --*

clip_R1 1 --length 16.

Cleaned reads were then mapped to the TMV-GFP sequence using hisat2 (v2.1.0, (Kim et al.,

2019)) with default parameters. The resulting sam files were sorted and indexed using Samtools

(v1.15.1, (Danecek et al., 2021)) and the length distribution of reads mapping to the TMV-GFP

sequence was computed using samtools stats.

Constructs and plant transformation

cDNA from Marchantia mock treated plants was used to amplify the coding regions of *MpTOM1*

(Mp3g00660), *MpGDI2* (Mp6g05010) and *MpNAC7* (Mp6g02620) by PCR using Phusion

polymerase (NEB). Arabidopsis *Bobber1* (At5g53400) and *ATAF2* (At5g08790) were amplified

using cDNA synthesized from RNA from Arabidopsis inflorescences. p126 and Δp126 were

amplified using DNA from the TMV-GFP containing plasmid. All obtained fragments were further

incubated with dATP and Taq DNA polymerase (NEB) for 1 hour before cloned into the gateway
compatible vector PCR8/GW/TOPO TA (Invitrogen) or directly cloned to pENTR D-TOPO
(Invitrogen) in the case of ATAF2. An entry vector containing the Arabidopsis AGO1 was kindly
provided by Pablo Manavella. Inserts were cloned into their final vectors through LR reaction
using LR clonase II (Invitrogen). All plasmids were submitted to Sanger sequencing to corroborate
their integrity before used in the corresponding experiments. A list of primers used, and
constructs can be found in supplementary table 3 and supplementary table 4.

Plant transformation was performed as described in (Ishizaki et al., 2008) and selected in
Gamborg´s B5 media supplemented with Hygromycin 100mg/ml and Cefotaxime 100mg/ml.

Histology and imaging

Marchantia plants were embedded in 8% agarose and cross-sections of around 200 μM were
obtained using a vibratome. Confocal imaging was performed using Leica SP5. Images were
processed using Fiji (Schindelin et al., 2012).

Protein-protein interaction assays

Interactions in the yeast two-hybrid system were assayed on selective medium (L$^-$, W$^-$, H$^-$). The
assay was repeated 3 times. For the luciferase complementation assay, one-month old rdr6i
Nicotiana plants were used. *Agrobacterium tumefaciens* cultures with the different constructs
were mixed in the same amounts at a final OD$_{600}$=0.6. Leaves were imaged three days after
inoculation using a LAS4000 device. The assay was repeated 3 times with equivalent results.

Photosynthetic measurements

10 measurements per plant were performed in 5 infected and 5 mock treated individuals using
an IMAGING-PAM Maxi device (Heinz Walz GmbH). Quantification was done using the
ImagingWing integrated software.

Reporting summary

Further information on research design is available in the reporting summary linked to this
article.

Data availability

Raw data from the virome has been deposited in SRA and GenBank under Bioproject

PRJNA1022866 (https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1022866) and

BioSamples (SAMN37631837 and SAMN40978195). Raw data from TMV infected vs. Mock

treated plants RNA-seq and TMV sRNA-seq have been deposit in SRA (Bioproject PRJNA1010197,

https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1010197). Reconstructed genomic

sequences of three novel viruses uncovered by virome analysis were deposited in NCBI GenBank

(OR640980 https://www.ncbi.nlm.nih.gov/nuccore/OR640980, OR640981

https://www.ncbi.nlm.nih.gov/nuccore/OR640981, OR640982

https://www.ncbi.nlm.nih.gov/nuccore/OR640982). Raw data for other analysis included in this

work can be found in the accompanying Source data file.

Code availability

Original code for DNA virome analysis can be accessed at

https://github.com/LanaVogrinec/snakemake-marchantia, which has deposited at ZENODO

under the ID 10.5281/zenodo.13359141 (https://zenodo.org/records/13359141).

References

BALLY, J., NAKASUGI, K., JIA, F., JUNG, H., HO, S. Y., WONG, M., PAUL, C. M., NAIM, F., WOOD, C.
C., CROWHURST, R. N., HELLENS, R. P., DALE, J. L. & WATERHOUSE, P. M. 2015. The
extremophile Nicotiana benthamiana has traded viral defence for early vigour. *Nat*
Plants, 1**,** 15165.
BEAULIEU, C., LIBOUREL, C., MBADINGA ZAMAR, D. L., MAHBOUBI, K. E., HOEY, D. J., KELLER, J.,
GIROU, C., CLEMENTE, H. S., DIOP, I., AMBLARD, E., THÉRON, A., CAUET, S., RODDE, N.,
ZACHGO, S., HALPAPE, W., MEIERHENRICH, A., LAKER, B., BRAUTIGAM, A., GREIFF, G. R.,
SZOVENYI, P., CHENG, S., TANIZAWA, Y., LEEBENS-MACK, J. H., SCHMUTZ, J., WEBBER, J.,
GRIMWOOD, J., JACQUET, C., DUNAND, C., NELSON, J. M., ROUX, F., PHILIPPE, H.,
SCHORNACK, S., BONHOMME, M. & DELAUX, P.-M. 2023. The Marchantia
pangenome reveals ancient mechanisms of plant adaptation to the environment.
*bioRxiv***,** 2023.10.27.564390.
BECK, M., WYRSCH, I., STRUTT, J., WIMALASEKERA, R., WEBB, A., BOLLER, T. & ROBATZEK, S. 2014.
Expression patterns of flagellin sensing 2 map to bacterial entry sites in plant shoots and
roots. *J Exp Bot,* 65**,** 6487-98.
BEDOYA, L. C., MARTINEZ, F., ORZAEZ, D. & DAROS, J. A. 2012. Visual tracking of plant virus
infection and movement using a reporter MYB transcription factor that activates
anthocyanin biosynthesis. *Plant Physiol,* 158**,** 1130-8.
BELANGER, S., ZHAN, J. & MEYERS, B. C. 2023. Phylogenetic analyses of seven protein families
refine the evolution of small RNA pathways in green plants. *Plant Physiol,* 192**,** 1183-
1203.

BENITEZ, M., HERNANDEZ-HERNANDEZ, V., NEWMAN, S. A. & NIKLAS, K. J. 2018. Dynamical
784   Patterning Modules, Biogeneric Materials, and the Evolution of Multicellular Plants.
785   *Front Plant Sci,* 9**,** 871.
786   BOWLES, A. M. C., PAPS, J. & BECHTOLD, U. 2022. Water-related innovations in land plants
787   evolved by different patterns of gene cooption and novelty. *New Phytol,* 235**,** 732-742.
788   BOWMAN, J. L., KOHCHI, T., YAMATO, K. T., JENKINS, J., SHU, S., ISHIZAKI, K., YAMAOKA, S.,
789   NISHIHAMA, R., NAKAMURA, Y., BERGER, F., ADAM, C., AKI, S. S., ALTHOFF, F., ARAKI, T.,
790   ARTEAGA-VAZQUEZ, M. A., BALASUBRMANIAN, S., BARRY, K., BAUER, D., BOEHM, C. R.,
791   BRIGINSHAW, L., CABALLERO-PEREZ, J., CATARINO, B., CHEN, F., CHIYODA, S., CHOVATIA,
792   M., DAVIES, K. M., DELMANS, M., DEMURA, T., DIERSCHKE, T., DOLAN, L., DORANTES-
793   ACOSTA, A. E., EKLUND, D. M., FLORENT, S. N., FLORES-SANDOVAL, E., FUJIYAMA, A.,
794   FUKUZAWA, H., GALIK, B., GRIMANELLI, D., GRIMWOOD, J., GROSSNIKLAUS, U.,
795   HAMADA, T., HASELOFF, J., HETHERINGTON, A. J., HIGO, A., HIRAKAWA, Y., HUNDLEY, H.
796   N., IKEDA, Y., INOUE, K., INOUE, S. I., ISHIDA, S., JIA, Q., KAKITA, M., KANAZAWA, T.,
797   KAWAI, Y., KAWASHIMA, T., KENNEDY, M., KINOSE, K., KINOSHITA, T., KOHARA, Y., KOIDE,
798   E., KOMATSU, K., KOPISCHKE, S., KUBO, M., KYOZUKA, J., LAGERCRANTZ, U., LIN, S. S.,
799   LINDQUIST, E., LIPZEN, A. M., LU, C. W., DE LUNA, E., MARTIENSSEN, R. A., MINAMINO,
800   N., MIZUTANI, M., MIZUTANI, M., MOCHIZUKI, N., MONTE, I., MOSHER, R., NAGASAKI,
801   H., NAKAGAMI, H., NARAMOTO, S., NISHITANI, K., OHTANI, M., OKAMOTO, T.,
802   OKUMURA, M., PHILLIPS, J., POLLAK, B., REINDERS, A., ROVEKAMP, M., SANO, R., SAWA,
803   S., SCHMID, M. W., SHIRAKAWA, M., SOLANO, R., SPUNDE, A., SUETSUGU, N., SUGANO,
804   S., SUGIYAMA, A., SUN, R., SUZUKI, Y., TAKENAKA, M., et al. 2017. Insights into Land Plant
805   Evolution Garnered from the Marchantia polymorpha Genome. *Cell,* 171**,** 287-304 e15.
806   BOX, M. S., COUSTHAM, V., DEAN, C. & MYLNE, J. S. 2011. Protocol: A simple phenol-based
807   method for 96-well extraction of high quality RNA from Arabidopsis. *Plant Methods,* 7**,**
808   7.
809   BRANTS, D. H. 1961. The Influence of Meristematic Tissue and Injuries on the Transport of
810   Tobacco Mosaic Virus in Nicotiana tabacum L. cultivar. Samsun. *Acta Botanica*
811   *Neerlandica,* 10**,** 113-163.
812   BUTKOVIC, A., DOLJA, V. V., KOONIN, E. V. & KRUPOVIC, M. 2023. Plant virus movement proteins
813   originated from jelly-roll capsid proteins. *PLoS Biol,* 21**,** e3002157.
814   CAMPBELL, R. N. 1996. Fungal transmission of plant viruses. *Annu Rev Phytopathol,* 34**,** 87-108.
815   CAPELLA-GUTIÉRREZ, S., SILLA-MARTÍNEZ, J. M. & GABALDÓN, T. 2009. trimAl: a tool for
816   automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics,* 25**,**
817   1972-1973.
818   CARELLA, P., GOGLEVA, A., HOEY, D. J., BRIDGEN, A. J., STOLZE, S. C., NAKAGAMI, H. &
819   SCHORNACK, S. 2019. Conserved Biochemical Defenses Underpin Host Responses to
820   Oomycete Infection in an Early-Divergent Land Plant Lineage. *Curr Biol,* 29**,** 2282-2294
821   e5.
822   CARELLA, P., GOGLEVA, A., TOMASELLI, M., ALFS, C. & SCHORNACK, S. 2018. Phytophthora
823   palmivora establishes tissue-specific intracellular infection structures in the earliest
824   divergent land plant lineage. *Proc Natl Acad Sci U S A,* 115**,** E3846-E3855.
825   COLLUM, T. D. & CULVER, J. N. 2017. Tobacco mosaic virus infection disproportionately impacts
826   phloem associated translatomes in Arabidopsis thaliana and Nicotiana benthamiana.
827   *Virology,* 510**,** 76-89.
828   CHIA, K. S., KOURELIS, J., TEULET, A., VICKERS, M., SAKAI, T., WALKER, J. F., SCHORNACK, S.,
829   KAMOUN, S. & CARELLA, P. 2024a. The N-terminal domains of NLR immune receptors
830   exhibit structural and functional similarities across divergent plant lineages. *Plant Cell,*
831   36**,** 2491-2511.

CHIA, K. S., KOURELIS, J., TEULET, A., VICKERS, M., SAKAI, T., WALKER, J. F., SCHORNACK, S.,
KAMOUN, S. & CARELLA, P. 2024b. The N-terminal domains of NLR immune receptors
exhibit structural and functional similarities across divergent plant lineages. *Plant Cell*.
DANECEK, P., BONFIELD, J. K., LIDDLE, J., MARSHALL, J., OHAN, V., POLLARD, M. O., WHITWHAM,
A., KEANE, T., MCCARTHY, S. A., DAVIES, R. M. & LI, H. 2021. Twelve years of SAMtools
and BCFtools. *Gigascience,* 10.
DE VRIES, S., DE VRIES, J., VON DAHLEN, J. K., GOULD, S. B., ARCHIBALD, J. M., ROSE, L. E. &
SLAMOVITS, C. H. 2018. On plant defense signaling networks and early land plant
evolution. *Commun Integr Biol,* 11**,** 1-14.
DEBAT, H., GARCIA, M. L. & BEJERMAN, N. 2023. Expanding the Repertoire of the Plant-Infecting
Ophioviruses through Metatranscriptomics Data. *Viruses,* 15.
DELESSERT, C., KAZAN, K., WILSON, I. W., VAN DER STRAETEN, D., MANNERS, J., DENNIS, E. S. &
DOLFERUS, R. 2005. The transcription factor ATAF2 represses the expression of
pathogenesis-related genes in Arabidopsis. *Plant J,* 43**,** 745-57.
DENG, Z. L., MUNCH, P. C., MRECHES, R. & MCHARDY, A. C. 2022. Rapid and accurate
identification of ribosomal RNA sequences via deep learning. *Nucleic Acids Res,* 50**,** e60.
DING, S. W., HAN, Q., WANG, J. & LI, W. X. 2018. Antiviral RNA interference in mammals. *Curr
Opin Immunol,* 54**,** 109-114.
DING, S. W. & VOINNET, O. 2007. Antiviral immunity directed by small RNAs. *Cell,* 130**,** 413-26.
DING, X. S., LIU, J., CHENG, N. H., FOLIMONOV, A., HOU, Y. M., BAO, Y., KATAGI, C., CARTER, S. A.
& NELSON, R. S. 2004. The Tobacco mosaic virus 126-kDa protein associated with virus
replication and movement suppresses RNA silencing. *Mol Plant Microbe Interact,* 17**,**
583-92.
DOLJA, V. V., KRUPOVIC, M. & KOONIN, E. V. 2020. Deep Roots and Splendid Boughs of the Global
Plant Virome. *Annu Rev Phytopathol,* 58**,** 23-53.
DOS REIS FIGUEIRA, A., GOLEM, S., GOREGAOKER, S. P. & CULVER, J. N. 2002. A nuclear
localization signal and a membrane association domain contribute to the cellular
localization of the Tobacco mosaic virus 126-kDa replicase protein. *Virology,* 301**,** 81-9.
EDGAR, R. C., TAYLOR, B., LIN, V., ALTMAN, T., BARBERA, P., MELESHKO, D., LOHR, D.,
NOVAKOVSKY, G., BUCHFINK, B., AL-SHAYEB, B., BANFIELD, J. F., DE LA PENA, M.,
KOROBEYNIKOV, A., CHIKHI, R. & BABAIAN, A. 2022. Petabase-scale sequence alignment
catalyses viral discovery. *Nature,* 602**,** 142-147.
ERICKSON, F. L., HOLZBERG, S., CALDERON-URREA, A., HANDLEY, V., AXTELL, M., CORR, C. &
BAKER, B. 1999. The helicase domain of the TMV replicase proteins induces the N-
mediated defence response in tobacco. *Plant J,* 18**,** 67-75.
ESPINOZA, C., MEDINA, C., SOMERVILLE, S. & ARCE-JOHNSON, P. 2007. Senescence-associated
genes induced during compatible viral interactions with grapevine and Arabidopsis. *J Exp
Bot,* 58**,** 3197-212.
GAN, S. 2003. Mitotic and postmitotic senescence in plants. *Sci Aging Knowledge Environ,* 2003**,**
RE7.
GARCIA-RUIZ, H., TAKEDA, A., CHAPMAN, E. J., SULLIVAN, C. M., FAHLGREN, N., BREMPELIS, K. J.
& CARRINGTON, J. C. 2010. Arabidopsis RNA-dependent RNA polymerases and dicer-like
proteins in antiviral defense and small interfering RNA biogenesis during Turnip Mosaic
Virus infection. *Plant Cell,* 22**,** 481-96.
GIMENEZ-IBANEZ, S., ZAMARRENO, A. M., GARCIA-MINA, J. M. & SOLANO, R. 2019. An
Evolutionarily Ancient Immune System Governs the Interactions between Pseudomonas
syringae and an Early-Diverging Land Plant Lineage. *Curr Biol,* 29**,** 2270-2281 e4.
GISH, W. & STATES, D. J. 1993. Identification of protein coding regions by database similarity
search. *Nat Genet,* 3**,** 266-72.

GU, Z. 2022. Complex Heatmap Visualization. *iMeta*.

GUO, S. & WONG, S. M. 2017. Deep sequencing analysis reveals a TMV mutant with a poly(A)
tract reduces host defense responses in Nicotiana benthamiana. *Virus Res,* 239, 126-135.

HERNANDEZ-HERNANDEZ, V., BENITEZ, M. & BOUDAOUD, A. 2020. Interplay between turgor
pressure and plasmodesmata during plant development. *J Exp Bot,* 71, 768-777.

HEYMAN, J., CANHER, B., BISHT, A., CHRISTIAENS, F. & DE VEYLDER, L. 2018. Emerging role of the
plant ERF transcription factors in coordinating wound defense responses and repair. *J
Cell Sci,* 131.

HOANG, D. T., CHERNOMOR, O., VON HAESELER, A., MINH, B. Q. & VINH, L. S. 2018. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution,* 35,
518-522.

HUSON, D. H., BEIER, S., FLADE, I., GÓRSKA, A., EL-HADIDI, M., MITRA, S., RUSCHEWEYH, H. J. &
TAPPU, R. 2016. MEGAN Community Edition - Interactive Exploration and Analysis of
Large-Scale Microbiome Sequencing Data. *Plos Computational Biology,* 12.

ISHIZAKI, K., CHIYODA, S., YAMATO, K. T. & KOHCHI, T. 2008. Agrobacterium-mediated
transformation of the haploid liverwort Marchantia polymorpha L., an emerging model
for plant biology. *Plant Cell Physiol,* 49, 1084-91.

IWAMA, R. E. & MORAN, Y. 2023. Origins and diversification of animal innate immune responses
against viral infections. *Nat Ecol Evol,* 7, 182-193.

JEON, H.-W., IWAKAWA, H., MARAMOTO, S., HERRFURTH, C., GUSTCHE, N., SCHLÜTER, T.,
KYOZUKA, J., MIYAUCHI, S., FEUSSNER, I., ZACHGO, S. & NAKAGAMI, H. 2022. Contrasting
and conserved roles of NPR pathways in diverged land plant lineages. *bioRxiv*.

JIA, X., WANG, L., ZHAO, H., ZHANG, Y., CHEN, Z., XU, L. & YI, K. 2023. The origin and evolution of
salicylic acid signaling and biosynthesis in plants. *Mol Plant,* 16, 245-259.

KALYAANAMOORTHY, S., MINH, B. Q., WONG, T. K. F., VON HAESELER, A. & JERMIIN, L. S. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature
Methods,* 14, 587-+.

KAPPAGANTU, M., COLLUM, T. D., DARDICK, C. & CULVER, J. N. 2020. Viral Hacks of the Plant
Vasculature: The Role of Phloem Alterations in Systemic Virus Infection. *Annu Rev Virol,*
7, 351-370.

KASSCHAU, K. D., XIE, Z., ALLEN, E., LLAVE, C., CHAPMAN, E. J., KRIZAN, K. A. & CARRINGTON, J.
C. 2003. P1/HC-Pro, a viral suppressor of RNA silencing, interferes with Arabidopsis
development and miRNA unction. *Dev Cell,* 4, 205-17.

KATOH, K. & STANDLEY, D. M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Molecular Biology and Evolution,* 30, 772-
780.

KIM, D., PAGGI, J. M., PARK, C., BENNETT, C. & SALZBERG, S. L. 2019. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol,* 37, 907-
915.

KNEESHAW, S., SORIANO, G., MONTE, I., HAMBERG, M., ZAMARRENO, A. M., GARCIA-MINA, J.
M., FRANCO-ZORRILLA, J. M., KATO, N., UEDA, M., REY-STOLLE, M. F., BARBAS, C.,
MICHAVILA, S., GIMENEZ-IBANEZ, S., JIMENEZ-ALEMAN, G. H. & SOLANO, R. 2022.
Ligand diversity contributes to the full activation of the jasmonate pathway in
Marchantia polymorpha. *Proc Natl Acad Sci U S A,* 119, e2202930119.

KOONIN, E. V., DOLJA, V. V., KRUPOVIC, M., VARSANI, A., WOLF, Y. I., YUTIN, N., ZERBINI, F. M. &
KUHN, J. H. 2020. Global Organization and Proposed Megataxonomy of the Virus World.
Microbiol Mol Biol Rev, 84.

KOPYLOVA, E., NOÉ, L. & TOUZET, H. 2012. SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data. *Bioinformatics,* 28, 3211-3217.

KORNER, C. J., KLAUSER, D., NIEHL, A., DOMINGUEZ-FERRERAS, A., CHINCHILLA, D., BOLLER, T.,
HEINLEIN, M. & HANN, D. R. 2013. The immunity regulator BAK1 contributes to
resistance against diverse RNA viruses. *Mol Plant Microbe Interact,* 26**,** 1271-80.
KRAMER, S. R., GOREGAOKER, S. P. & CULVER, J. N. 2011. Association of the Tobacco mosaic virus
126kDa replication protein with a GDI protein affects host susceptibility. *Virology,* 414**,**
110-8.
KULSHRESTHA, S., JIBRAN, R., VAN KLINK, J. W., ZHOU, Y., BRUMMELL, D. A., ALBERT, N. W.,
SCHWINN, K. E., CHAGNE, D., LANDI, M., BOWMAN, J. L. & DAVIES, K. M. 2022. Stress,
senescence, and specialized metabolites in bryophytes. *J Exp Bot,* 73**,** 4396-4411.
LEE, W. S., FU, S. F., LI, Z., MURPHY, A. M., DOBSON, E. A., GARLAND, L., CHALUVADI, S. R., LEWSEY,
M. G., NELSON, R. S. & CARR, J. P. 2016. Salicylic acid treatment and expression of an
RNA-dependent RNA polymerase 1 transgene inhibit lethal symptoms and meristem
invasion during tobacco mosaic virus infection in Nicotiana benthamiana. *BMC Plant*
Biol, 16**,** 15.
LEEK, J. T., JOHNSON, W. E., PARKER, H. S., FERTIG, E. J., JAFFE, A.E., ZHANG, Y., STOREY, J. D. &
TORRES, L. C. 2023. sva: Surrogate Variable Analysis.
LETUNIC, I. & BORK, P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. *Nucleic Acids Research,* 49**,** W293-W296.
LEWSEY, M. G. & CARR, J. P. 2009. Effects of DICER-like proteins 2, 3 and 4 on cucumber mosaic
virus and tobacco mosaic virus infections in salicylic acid-treated plants. *J Gen Virol,* 90**,**
3010-3014.
LIANG, Y., HEYMAN, J., XIANG, Y., VANDENDRIESSCHE, W., CANHER, B., GOEMINNE, G. & DE
VEYLDER, L. 2022. The wound-activated ERF15 transcription factor drives Marchantia
polymorpha regeneration by activating an oxylipin biosynthesis feedback loop. *Sci Adv,*
8**,** eabo7737.
LIM, P. O. & NAM, H. G. 2005. The molecular and genetic control of leaf senescence and longevity
in Arabidopsis. *Curr Top Dev Biol,* 67**,** 49-83.
LIU, C. & NELSON, R. S. 2013. The cell biology of Tobacco mosaic virus replication and movement.
Front Plant Sci, 4**,** 12.
LIU, Y., GAO, Q., WU, B., AI, T. & GUO, X. 2009. NgRDR1, an RNA-dependent RNA polymerase
isolated from Nicotiana glutinosa, was involved in biotic and abiotic stresses. *Plant*
Physiol Biochem, 47**,** 359-68.
LOPEZ-GOMOLLON, S. & BAULCOMBE, D. C. 2022. Roles of RNA silencing in viral and non-viral
plant immunity and in the crosstalk between disease resistance systems. *Nat Rev Mol*
Cell Biol, 23**,** 645-662.
LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol,* 15**,** 550.
LUCAS, W. J., GROOVER, A., LICHTENBERGER, R., FURUTA, K., YADAV, S. R., HELARIUTTA, Y., HE, X.
Q., FUKUDA, H., KANG, J., BRADY, S. M., PATRICK, J. W., SPERRY, J., YOSHIDA, A., LOPEZ-
MILLAN, A. F., GRUSAK, M. A. & KACHROO, P. 2013. The plant vascular system: evolution,
development and functions. *J Integr Plant Biol,* 55**,** 294-388.
MANDADI, K. K. & SCHOLTHOF, K. B. 2013. Plant immune responses against viruses: how does a
virus cause disease? *Plant Cell,* 25**,** 1489-505.
MATSUI, H., IWAKAWA, H., HYON, G. S., YOTSUI, I., KATOU, S., MONTE, I., NISHIHAMA, R.,
FRANZEN, R., SOLANO, R. & NAKAGAMI, H. 2020. Isolation of Natural Fungal Pathogens
from Marchantia polymorpha Reveals Antagonism between Salicylic Acid and Jasmonate
during Liverwort-Fungus Interactions. *Plant Cell Physiol,* 61**,** 265-275.

MIFSUD, J. C. O., GALLAGHER, R. V., HOLMES, E. C. & GEOGHEGAN, J. L. 2022. Transcriptome
Mining Expands Knowledge of RNA Viruses across the Plant Kingdom. *J Virol,* 96**,**
e0026022.
MONTE, I., FRANCO-ZORRILLA, J. M., GARCIA-CASADO, G., ZAMARRENO, A. M., GARCIA-MINA, J.
M., NISHIHAMA, R., KOHCHI, T. & SOLANO, R. 2019. A Single JAZ Repressor Controls the
Jasmonate Pathway in Marchantia polymorpha. *Mol Plant,* 12**,** 185-198.
MONTE, I., ISHIDA, S., ZAMARRENO, A. M., HAMBERG, M., FRANCO-ZORRILLA, J. M., GARCIA-
CASADO, G., GOUHIER-DARIMONT, C., REYMOND, P., TAKAHASHI, K., GARCIA-MINA, J.
M., NISHIHAMA, R., KOHCHI, T. & SOLANO, R. 2018. Ligand-receptor co-evolution shaped
the jasmonate pathway in land plants. *Nat Chem Biol,* 14**,** 480-488.
MUSHEGIAN, A., SHIPUNOV, A. & ELENA, S. F. 2016. Changes in the composition of the RNA
virome mark evolutionary transitions in green plants. *BMC Biol,* 14**,** 68.
NERI, U., WOLF, Y. I., ROUX, S., CAMARGO, A. P., LEE, B., KAZLAUSKAS, D., CHEN, I. M., IVANOVA,
N., ZEIGLER ALLEN, L., PAEZ-ESPINO, D., BRYANT, D. A., BHAYA, D., CONSORTIUM, R. N. A.
V. D., KRUPOVIC, M., DOLJA, V. V., KYRPIDES, N. C., KOONIN, E. V. & GOPHNA, U. 2022.
Expansion of the global RNA virome reveals diverse clades of bacteriophages. *Cell,* 185**,**
4023-4037 e18.
NGUYEN, L. T., SCHMIDT, H. A., VON HAESELER, A. & MINH, B. Q. 2015. IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.
Molecular Biology and Evolution, 32**,** 268-274.
NIEHL, A., WYRSCH, I., BOLLER, T. & HEINLEIN, M. 2016. Double-stranded RNAs induce a pattern-
triggered immune signaling pathway in plants. *New Phytol,* 211**,** 1008-19.
PAGAN, I., ALONSO-BLANCO, C. & GARCIA-ARENAL, F. 2008. Host responses in life-history traits
and tolerance to virus infection in Arabidopsis thaliana. *PLoS Pathog,* 4**,** e1000124.
PATRO, R., DUGGAL, G., LOVE, M. I., IRIZARRY, R. A. & KINGSFORD, C. 2017. Salmon provides fast
and bias-aware quantification of transcript expression. *Nature Methods,* 14**,** 417-+.
PEART, J. R., COOK, G., FEYS, B. J., PARKER, J. E. & BAULCOMBE, D. C. 2002. An EDS1 orthologue
is required for N-mediated resistance against tobacco mosaic virus. *Plant J,* 29**,** 569-79.
PRUITT, R. N., LOCCI, F., WANKE, F., ZHANG, L., SAILE, S. C., JOE, A., KARELINA, D., HUA, C.,
FROHLICH, K., WAN, W. L., HU, M., RAO, S., STOLZE, S. C., HARZEN, A., GUST, A. A.,
HARTER, K., JOOSTEN, M., THOMMA, B., ZHOU, J. M., DANGL, J. L., WEIGEL, D.,
NAKAGAMI, H., OECKING, C., KASMI, F. E., PARKER, J. E. & NURNBERGER, T. 2021. The
EDS1-PAD4-ADR1 node mediates Arabidopsis pattern-triggered immunity. *Nature,* 598**,**
495-499.
QUILLIAM, R. S., SWARBRICK, P. J., SCHOLES, J. D. & ROLFE, S. A. 2006. Imaging photosynthesis in
wounded leaves of Arabidopsis thaliana. *J Exp Bot,* 57**,** 55-69.
RAKHSHANDEHROO, F., REZAEE, S. & PALUKAITIS, P. 2017. Silencing the tobacco gene for RNA-
dependent RNA polymerase 1 and infection by potato virus Y cause remodeling of
cellular organelles. *Virology,* 510**,** 127-136.
REDKAR, A., GIMENEZ IBANEZ, S., SABALE, M., ZECHMANN, B., SOLANO, R. & DI PIETRO, A. 2022.
Marchantia polymorpha model reveals conserved infection mechanisms in the vascular
wilt fungal pathogen Fusarium oxysporum. *New Phytol,* 234**,** 227-241.
SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M., PIETZSCH, T.,
PREIBISCH, S., RUEDEN, C., SAALFELD, S., SCHMID, B., TINEVEZ, J. Y., WHITE, D. J.,
HARTENSTEIN, V., ELICEIRI, K., TOMANCAK, P. & CARDONA, A. 2012. Fiji: an open-source
platform for biological-image analysis. *Nat Methods,* 9**,** 676-82.
SCHOLTHOF, K. B. 2004. Tobacco mosaic virus: a model system for plant biology. *Annu Rev
Phytopathol,* 42**,** 13-34.

SCHOMMER, C., PALATNIK, J. F., AGGARWAL, P., CHETELAT, A., CUBAS, P., FARMER, E. E., NATH,
U. & WEIGEL, D. 2008. Control of jasmonate biosynthesis and senescence by miR319
targets. *PLoS Biol,* 6**,** e230.
SCHWACH, F., VAISTIJ, F. E., JONES, L. & BAULCOMBE, D. C. 2005. An RNA-dependent RNA
polymerase prevents meristem invasion by potato virus X and is required for the activity
but not the production of a systemic silencing signal. *Plant Physiol,* 138**,** 1842-52.
SHENG, Y., YANG, L., LI, C., WANG, Y. & GUO, H. 2019. Transcriptomic changes in Nicotiana
tabacum leaves during mosaic virus infection. *3 Biotech,* 9**,** 220.
SIEVERT, R.C.. 1978. Effect of Potato Virus Y and Tobacco Mosaic Virus on Field-Grown Burley
Tobacco. *PHYTOPATHOLOGY,* 68**,** 823-825.
SILVESTRI, A., BANSAL, C. & RUBIO-SOMOZA, I. 2024. After silencing suppression: miRNA targets
strike back. *Trends Plant Sci.*
SONESON, C., LOVE, M. I. & ROBINSON, M. D. 2015. Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Res,* 4**,** 1521.
VASSEUR, F., BALDRICH, P., JIMENEZ-GONGORA, T., VILLAR-MARTIN, L., WEIGEL, D. & RUBIO-
SOMOZA, I. 2022. miRNA472 deficiency enhances Arabidopsis thaliana defence without
reducing seed production. *bioRxiv,* doi.org/10.1101/2022.12.13.520224
VENDRELL-MIR, P., PERROUD, P. F., HAAS, F. B., MEYBERG, R., CHARLOT, F., RENSING, S. A.,
NOGUE, F. & CASACUBERTA, J. M. 2021. A vertically transmitted amalgavirus is present
in certain accessions of the bryophyte Physcomitrium patens. *Plant J,* 108**,** 1786-1797.
WANG, L., WAN, M. C., LIAO, R. Y., XU, J., XU, Z. G., XUE, H. C., MAI, Y. X. & WANG, J. W. 2023. The
maturation and aging trajectory of Marchantia polymorpha at single-cell resolution. *Dev
Cell,* 58**,** 1429-1444 e6.
WANG, X., GOREGAOKER, S. P. & CULVER, J. N. 2009. Interaction of the Tobacco mosaic virus
replicase protein with a NAC domain transcription factor is associated with the
suppression of systemic host defenses. *J Virol,* 83**,** 9720-30.
WHITHAM, S., DINESH-KUMAR, S. P., CHOI, D., HEHL, R., CORR, C. & BAKER, B. 1994. The product
of the tobacco mosaic virus resistance gene N: similarity to toll and the interleukin-1
receptor. *Cell,* 78**,** 1101-15.
XU, T., ZHANG, L., ZHEN, J., FAN, Y., ZHANG, C. & WANG, L. 2013. Expressional and regulatory
characterization of Arabidopsis RNA-dependent RNA polymerase 1. *Planta,* 237**,** 1561-9.
XU, X. J., GENG, C., JIANG, S. Y., ZHU, Q., YAN, Z. Y., TIAN, Y. P. & LI, X. D. 2022. A maize
triacylglycerol lipase inhibits sugarcane mosaic virus infection. *Plant Physiol,* 189**,** 754-
771.
YAMANAKA, T., OHTA, T., TAKAHASHI, M., MESHI, T., SCHMIDT, R., DEAN, C., NAITO, S. &
ISHIKAWA, M. 2000. TOM1, an Arabidopsis gene required for efficient multiplication of
a tobamovirus, encodes a putative transmembrane protein. *Proc Natl Acad Sci U S A,* 97**,**
10107-12.
YING, X. B., DONG, L., ZHU, H., DUAN, C. G., DU, Q. S., LV, D. Q., FANG, Y. Y., GARCIA, J. A., FANG,
R. X. & GUO, H. S. 2010. RNA-dependent RNA polymerase 1 from Nicotiana tabacum
suppresses RNA silencing and enhances viral infection in Nicotiana benthamiana. *Plant
Cell,* 22**,** 1358-72.
YOTSUI, I., MATSUI, H., MIYAUCHI, S., IWAKAWA, H., MELKONIAN, K., SCHLUTER, T., MICHAVILA,
S., KANAZAWA, T., NOMURA, Y., STOLZE, S. C., JEON, H. W., YAN, Y., HARZEN, A., SUGANO,
S. S., SHIRAKAWA, M., NISHIHAMA, R., ICHIHASHI, Y., IBANEZ, S. G., SHIRASU, K., UEDA,
T., KOHCHI, T. & NAKAGAMI, H. 2023. LysM-mediated signaling in Marchantia
polymorpha highlights the conservation of pattern-triggered immunity in land plants.
Curr Biol, 33**,** 3732-3746 e8.

YU, D., FAN, B., MACFARLANE, S. A. & CHEN, Z. 2003. Analysis of the involvement of an inducible
Arabidopsis RNA-dependent RNA polymerase in antiviral defense. *Mol Plant Microbe*
Interact, 16**,** 206-16.
YU, X., XU, Y. & YAN, S. 2021. Salicylic acid and ethylene coordinately promote leaf senescence. *J*
Integr Plant Biol, 63**,** 823-827.
ZAMFIR, A. D., BABALOLA, B. M., FRAILE, A., MCLEISH, M. J. & GARCIA-ARENAL, F. 2023.
Tobamoviruses Show Broad Host Ranges and Little Genetic Diversity Among Four Habitat
Types of a Heterogeneous Ecosystem. *Phytopathology,* 113**,** 1697-1707.
ZEINSMASTER D.D & CAROTHERS Z.B 1974. The Fine Structure of Oogenesis in Marchantia
polymorpha. *American Journal of Botany,* 61**,** 499-512.
ZHU, F., XI, D. H., YUAN, S., XU, F., ZHANG, D. W. & LIN, H. H. 2014. Salicylic acid and jasmonic
acid are essential for systemic resistance against tobacco mosaic virus in Nicotiana
benthamiana. *Mol Plant Microbe Interact,* 27**,** 567-77.

Acknowledgments

We would like to thank Juan Jose López-Moya, Miguel A. Blázquez and Salomé Prat for critical

reading of the manuscript. Sebastian Schornack, Phil Carella, Alexander Summers, the CUBG

curation team (Samuel Brockington and Mar Millan), Klaus Schlaeppi, Katja Rembold, Alba

Gonzalez, Detlef Weigel, Alexandra Kehl, Brigitte Fiebig, Luisa Lanfranco, Mara Novero, Isabel

Monte, Adrián Vojtassák, Sabine Zachgo and Claudia Gieshoidt for their help in obtaining

Marchantia individuals for the virome assays. Neža Pajek Arambašić for help with pipeline

automation. We would like to thank Eduardo Bejarano for providing the plasmid containing TMV-

GFP and Antonio Darós for providing TuMV-ROS1 and TMV-ROS1 containing plasmids. Pablo

Manavella for the AtAGO1-pEntr plasmid. Phil Carella for an updated annotation of NLR genes in

Marchantia. We would also thank to Jia-Wei Wang, Long Wang and Jie Xu for sharing the code

used for correlation analysis between TMV transcriptional responses and the different samples

from Marchantia developmental trajectories.

The work in the MoRE lab is funded by RYC-2015-19154 (funded by MCIN/AEI/

10.13039/501100011033 and by "ESF Investing in your future"), RTI2018-097262-B-I00 (funded

by MCIN/AEI/ 10.13039/501100011033 and by "ERDF A way of making Europe") and through

the "Severo Ochoa Programme for Centres of Excellence in R&D" 2016-2019 (SEV-2015-0533)

and 2020-2023 (CEX2019-000902-S) funded by MCIN/AEI/ 10.13039/501100011033,CSIC-MCIN

(Grant 202240I201) and the CERCA and SGR (2021-SGR-00875) programmes from the

Generalitat de Catalunya to I-R-S. T J-G was recipient of a Postdoctoral Fellowship from CRAG

through the "Severo Ochoa Programme for Centres of Excellence in R&D" 2016-2019 (SEV-2015-

0533). L. V-M was supported by BES-2016-076986 (funded by MCIN/AEI/

10.13039/501100011033 and by "ESF Investing in your future"). The work at NIB was funded by

Slovenian Research and Innovation Agency core and project financing P4-0407, P4-0165 and J4-

4553.

Authors contributions

I.R-S. and T.J-G. conceptualized the study and designed experiments. I.R-S. supervised the study

and acquired funding. T.J-G. and I.R-S. isolated RNAs for virome analysis. D.K. and L.V. performed

virome analysis. E.R-M. and L.V-M. performed expression analysis of presence of viruses in

natural Marchantia populations and Marchantia infection with natural viruses. T.J-G. and I.R-S.

performed histological and expression analysis of viral infections. I.R-S. and V.M.G-M. performed

RNA isolation, sRNA libraries and analysis for transcriptomic assays. E.R-M. carried out all

phenotypic and p126 histological studies. I.R-S. carried out the protein-protein interaction assays

and performed infection assays in Nicotiana. I.R-S. and D.K. wrote the manuscript with input

from all authors.

1121

1122   Corresponding author

1123   Correspondence to ignacio.rubio@cragenomica.es

1124

Ethics declarations

Competing interests

The authors declare no competing interests

Figure legends

Figure 1. Marchantia polymorpha is infected by RNA viruses.

(A) Genome structure of Tübingen picorna-like virus 1, newly discovered in composite *M.*

polymorpha sample with two predicted open reading frames (ORFs) and corresponding

average reads mapping depth in sequenced composite sample. Black empty boxes

represent predicted protein domains detected by InterProScan: S3H – superfamily 3

helicase, 3C-Pro – peptidase C3, RdRp – RNA-dependent RNA polymerase and CP-SJR – coat

protein with 8-stranded jelly roll β-barrel motif.

(B) Maximum likelihood phylogenetic tree obtained based on the alignment of the

conserved part of RdRd of Tübingen picorna-like virus 1 and fifteen other most similar

viruses. The tree was midpoint rooted. Coloured virus names correspond to the sources

associated with a particular virus, according to the legend on the right side of the panel.
Numbers next to the branches represent statistical support (%) of the clades according to
the ultrafast bootstrap analysis. Branch length represents the average number of amino acid
substitutions per site. Codes next to the taxon names in round brackets are the
corresponding NCBI GenBank protein accession numbers. Next to them, fractions in square
brackets represent % of identity of Tübingen picorna-like virus 1 to other individual taxons
for the aligned part of the RdRp domain.

(C) Results of the viral concentration measurements for *M. polymorpha* plants inoculated
with Tübingen picorna-like virus 1 positive plant extracts. On the right, relative
concentrations of Tübingen picorna-like virus 1 in inoculated plants one- and two-weeks
post inoculation are shown; measurements of concentrations in individual plants (from 3
technical replicates each) are shown as dots (n=8 for both time points) and the box-whisker
plots represent the distribution of values (showing median as a thick line, first and third
quartile as boxes and minimum and maximum as whiskers) for the two time points (at 1
WPI, x represent an outlier measurement). On the top of the plot, the horizontal connecting
line designates statistically significant difference in the means of the two distributions, with
the corresponding p-value, according to the nonparametric Wilcoxon rank sum test (two-
sided). Source data are provided as a Source data File. BIORENDER was used to prepare this
figure.

Figure 2. Tobacco mosaic virus replicates in Marchantia plants.

(A) GFP fluorescent signal (in yellow) increases in thalli from 3-week-old plants after 1 week,
(left panel) compared to thalli from 4 weeks old and 2-weeks infected plants (right
panel). Both images show the same part or an infected plant under a magnification
device under UV light. Magenta shows chlorophyll autofluorescence. White bar scale
shows 100 µms. This experiment was repeated 4 times, and one additional replicate is
shown in Supplementary Fig. 3.
(B) TMV-GFP transcripts increase over time in infected plants. Pools of three infected plants
were used for each time point. The result from 3 technical replicates is shown, with two
complementary replicates showing the same trend in Supplementary Fig. 4. Mock, 4
weeks old plants incubated with sap from untreated plants. T0, 2 weeks old plants
collected right after incubation with sap from infected plants. T1, 3 weeks old plants
collected after 1 week from incubation. T2, 4 weeks old plants collected after 3 weeks

from incubation. Expression of TMV is relative to the expression of Marchantia actine.

Source data are provided as a Source data File.

(C) Length distribution of TMV-derived vsRNAs populations found in thalli from 4 weeks old

plants after 2 weeks of infection. Nts stans for nucleotides.

(D) TMV-GFP infects all cell types. Transversal sections of an Marchantia thalli from a mock

treated plant that was 4 weeks old (left) and equivalent area in a plant 2 weeks after

infection (right) under confocal image. The pictures show the merge between green (in

yellow, GFP-labelled virus) and red (in magenta, autofluorescence) channels. Merge was

performed using Fiji. P stands for Photosynthetic region, S for storage region and E stands

for lower epidermis. White bar scale shows 100 µm. This experiment was repeated 3

times showing similar results.

Figure 3. TMV triggers stress-related molecular reprogramming in Marchantia.

(A) Set of common PTI marker genes responsive to bacterial and fungal extracts and TMV in

infected areas in thalli from 4 weeks old plants after 2 weeks of infection compared to

equivalent areas from mock treated plants. DESeq2 results using the default parameters

(Wald test and Benjamini-Hochberg p adjustment method)

(B) Set of wound and TMV responsive ERF transcription factors. DESeq2 results using the

default parameters (Wald test and Benjamini-Hochberg p adjustment method)

(C) Venn diagram showing that transcriptional reprogramming upon TMV infection largely

overlaps with that found upon wounding.

(D) Set of chitin and TMV responsive WRKY transcription factors in Marchantia. DESeq2

results using the default parameters (Wald test and Benjamini-Hochberg p adjustment

method)

(E) Photosynthetic efficiency (Fv/Fm) in infected areas in thalli from 4 weeks old plants after

2 weeks of infection compared to equivalent areas from mock treated plants. 5 biological

and 10 technical replicates for each were assayed in both infected and mock treated

plants. Statistical assessment was performed using a two-sided parametric T-test. Length

of the whiskers: 1.5 x IQR. Central line: median. Box width: IQR (i.e. Q3-Q1). Source data

are provided as a Source data File. Marchantia plants and cameras depicting measured

areas come from Biorender. Light green area indicated TMV-GFP positive areas.

(F) Photosynthetic efficiency (Fv/Fm) in non-infected areas, in thalli from 4 weeks old plants

after 2 weeks of infection, compared to equivalent areas from mock treated plants. 5

biological and 10 technical replicates for each were assayed in both infected and mock

treated plants. Statistical assessment was performed using a two-sided parametric T-

test. Length of the whiskers: 1.5 x IQR. Central line: median. Box width: IQR (i.e. Q3-Q1).

Source data are provided as a Source data File. Marchantia plants and cameras depicting

measured areas come from Biorender. Light green area indicated TMV-GFP positive

areas.

Figure 4. Salicylic and jasmonate signalling pathways are activated during Marchantia-TMV

interaction.

(A)   Venn diagrams showing the overlap between the transcriptional responses under TMV

infection ad a salicylic treatment time course (2 and 6 hours).

(B)   Overlap between the trancriptome responses to TMV and Mpjaz1ko mutants with

constitutively active jasmonate signalling (left), and treatment with the active dn-OPDA

jasmonate (right).

(C)   Overlap between the trancriptome responses commonly found in TMV infection and the

1217   different hormone-related transcriptomes.

(D)   Biological processes to which genes commonly upregulated in response to TMV

infection, SA treatment (2,6 and 12h) and activation of jasmonate signalling belong

(E)   Biological processes to which genes commonly downregulated in response to TMV

infection, SA treatment (12h) and activation of jasmonate signalling belong.

(F)   Biological processes to which commonly upregulated in response to TMV infection, SA

treatment (6 h) and activation of jasmonate signalling in Mpjaz1KO plants belong.

(G) Biological processes to which commonly upregulated in response to TMV infection, SA (6

h) and dn-OPDA treatments belong.

Results in panels 4D-G come from a hypergeometric method with BY (BenjaminiYekutieli) p

adjustment with minimal size of genes annotated for testing of 5 and p-value cutoff of 0.05.

Figure 5. TMV infection triggers maturation and aging in Marchantia.

(A)   Pairwise comparison between transcriptomes from mock and TMV samples along with

those from sliced thallus samples. Slice 1(S1) corresponds to the top of thalli (youngest

part), while slice 6 (S6), corresponds to the base of thalli encompassing the oldest part.

(B)   Pairwise comparison between transcriptomes from mock and TMV samples along with

those from Marchantia developmental series Day 1 (D1) samples correspond to 1 day

after germination (youngest), while day 31 (31D) corresponds to fully mature Marchantia

thalli 31 days after germination.

Figure 6. Conservation of the interaction between Marchantia and the TMV silencing suppressor

p126.

(A) p126-GFP shows accumulation in the cytoplasm of Marchantia cells in cross-sections of

thalli from 4 weeks old plants (right) compared to thalli from untransformed plants of

the same age plants (left). The pictures show the merge between green (GFP) and red

(autofluorescence) channels. P stands for Photosynthetic region, S for storage region and

E stands for lower epidermis. Merge was performed using Fiji Scale bars are 100 µMs.

(B) Overexpression of p126 inhibits growth (area) when compared to Tak-1 control plants.

Analysis was done in 4 weeks old plants in 3 biological replicates (n=12). Statistical

significance was assessed using a two-sided Wilcoxon-test with Holm p adjustment.

Length of the whiskers: 1.5 x IQR. Central line: median. Box width: IQR (i.e. Q3-Q1).

(C) MpNAC7 interacts with the helicase domain of p126 in yeast.

(D) MpNAC7 and ATAF2 interact both with the helicase and the full length p126 in planta.

(E) Transient expression of MpNAC7 in Nicotiana plants abrogates local and systemic TMV

infection. TMV presence is shown as purple.

1251

A Tübingen picorna-like virus 1

ORF 1: S3H, 3C-Pro, RdRp
ORF 2: CP-SJR, CP-SJR, CP-SJR

Reads mapping depth

B

Beihai picorna-like virus 93 (APG76720) [29%]
Wenzhou channeled applesnail virus 2 (APG78498) [30%]
Riboviria sp. (QRQ90142.1) [30%]
Riboviria sp. (QRQ90158.1) [30%]
Trichosanthes kirilowii picorna-like virus (QKK82968.1) [37%]
Trichosanthes kirilowii picorna-like virus (QKK82970.1) [80%]
Tübingen picorna-like virus 1
Wenzhou picorna-like virus 42 (APG78519) [33%]
Shahe picorna-like virus 12 (APG77372) [42%]
Cragig virus 7 (QBP32757.1) [37%]
Ginkgo biloba picorna-like virus (QKK82974.1) [38%]
Wenzhou picorna-like virus 41 (APG78514) [38%]
Marnaviridae sp. (UGO57158.1) [35%]
Beihai picorna-like virus 102 (APG78882) [35%]
Shahe heteroptera virus 4 (APG77358) [35%]
Wenzhou picorna-like virus 40 (APG78510) [35%]

Branch length: 0.5

Source:
crayfish
mollusc
fresh water
plant
freshwater arthropod
bird feces

C

Extracts from Tübingen picorna-like virus 1 positive individuals + controls (Tak-1)

Inoculation of Tübingen picorna-like virus 1 negative individuals (Tak-1)

Measurement of Tübingen picorna-like virus 1 concentrations one and two weeks post inoculation

p = 0.02

Relative concentration of Tübingen picorna-like virus 1

Weeks post inoculation

A

B

C

Length (nts)
- 20
- 21
- 22
- 23
- 24
- 25

D

Mock

TMV-GFP

A *PTI Marker genes*

Name	ID	Relative expression level	padj
MpPR9	Mp5g21510	4.04	1.18E-21
MpSYP13B	Mp2g21910	2.34	0.00089
MpCML42	Mp6g18000	1.79	3.01E-17
MpWRKY22	Mp7g17200	1.74	1.82E-32
MpACRE132	Mp1g27170	1.22	1.30E-10

B *Wound and TMV responsive ERFs*

Name	ID	Relative expression level	padj
MpERF1	Mp7g09350	4.24	0.28
MpERF21	Mp5g22160	1.68	5.91E-05
MpERF8	Mp6g18650	1.48	1.33E-41
MpERF18	Mp5g12480	1.14	0.0015
MpERF15	Mp7g09350	1.06	0.028
MpAP2L1	Mp8g11450	-1.62	4.18E-13
MpERF10	Mp7g00860	-2.83	9.05E-11
MpERF17	Mp4g22280	-2.38	0.00058
MpERF12	Mp6g06870	-2.95	0.014
MpERF16	Mp4g22290	-3.11	0.0004
MpERF13	Mp6g08690	-3.25	4.72E-05

C

TMV Down — 1419
Wounding Up — 900
59
0 — 0
1250
0 — 0 — 0
248
616 — 0 — 167
0 — 0
13
TMV Up
Wounding Down

D *Chitin and TMV responsive WRKYs*

Name	ID	Relative expression level	padj
MpWRKY7	Mp3g17660	3.56	1.68E-34
MpWRKY6	Mp1g08960	2.52	3.71E-17
MpWRKY12	Mp4g00200	1.89	1.90E-11
MpWRKY10	Mp7g06550	1.79	7.31E-06
MpWRKY9	Mp7g17200	1.74	1.82E-32
MpWRKY13	Mp4g00180	1.64	7.72E-11
MpWRKY1	Mp3g00010	1.28	3.79E-25
MpWRKY14	Mp6g16800	1.16	0.013

E *Infected area*

Independent t-test, p=6.92e-05

Photosynthetic efficiency (y-axis: 0.0 to 0.9)

Mock | TMV-GFP

F *Non-infected area*

Independent t-test, p=0.0012

Photosynthetic efficiency (y-axis: 0.0 to 0.9)

Mock | TMV-GFP

A

Tak-1

P35S:p126-citrine

B

wilcoxon-test, p = 4.43e-05

Area (cm²)

Tak-1   *P35S:p126-citrine*

C

-AD/-BD | L⁻W⁻H⁻ | L⁻W⁻

-/p126
MpGDI2/p126
MpTOM1/p126
MpNAC7/p126
-/Δp126
MpGDI2/Δp126
MpNAC7/Δp126
MpTOM1/Δp126

D

-N-Luc/C-Luc- | GUS | TMV

MpNAC7/AtBOB1
AtATAF2/AtBOB1
AtAGO1/Δp126
AtAGO1/p126
MpNAC7/Δp126
MpNAC7/p126
AtATAF2/Δp126
AtATAF2/p126

High

Low

E

TMV-ROS1/GUS

TMV-ROS1/MpNAC7

Inoculated leaves   Systemic leaves