Dynamics of the viral community on the cheese surface during maturation and persistence

across production years

Thomas Paillet,[a] Quentin Lamy-Besnier,[b] Clarisse Figueroa,[a]* Marie-Agnès Petit,[b] Eric

Dugat-Bony[a]#

[a]Université Paris-Saclay, INRAE, AgroParisTech, UMR SayFood, 91120 Palaiseau, France

[b]Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78352 Jouy-en-Josas,

France

Running Head: Viral dynamics and persistence on the cheese surface

#Address correspondence to Eric Dugat-Bony, eric.dugat-bony@inrae.fr.

*Present address: Université de Paris Cité, INSERM, IAME, UMR 1137, 75018 Paris, France

ABSTRACT

The surface of smear-ripened cheeses constitutes a dynamic microbial ecosystem resulting from the

successive development of different microbial groups. Recent studies indicate that a viral community,

mainly composed of bacteriophages, coexists with cellular microorganisms in this ecosystem, but its

ecological significance remains to be elucidated. In this work, we studied a French smear-ripened

cheese by both viral metagenomics and 16S metabarcoding approaches to assess both the succession

of phages and bacterial communities on the cheese surface on the short-term during the ripening

period, and their temporal variations on the long-term in ready-to-eat cheeses over the years of

production. We observed a clear transition of the phage community structure during ripening with a

decreased relative abundance of viral species (vOTUs) associated with *Lactococcus* phages, which

were replaced by vOTUs associated with phages infecting ripening bacteria such as *Brevibacterium,*

Glutamicibacter, Pseudoalteromonas and *Vibrio*. The dynamics of the phage community was strongly

associated with bacterial successions observed on the cheese surface. Finally, while some variations in

the distribution of phages were observed in ready-to-eat cheeses produced at different dates spanning

more than 4 years of production, a core of abundant vOTUs was systematically detected. This result

revealed the long-term persistence of the main phages in the cheese production environment. Together,

these findings offer novel perspectives on the ecology of bacteriophages in smear-ripened cheese and

emphasize the significance of incorporating bacteriophages in the microbial ecology studies of

fermented foods.

IMPORTANCE

Smear-ripened cheeses are microbial ecosystems made up of various microorganisms including

bacteria, yeasts and also viruses such as bacteriophages, which infect and regulate bacterial

populations. In this work, a French smear-ripened cheese was used to study how these viruses and

bacteria interact over time in both the short-term during cheese production, and the long-term across

several years of production. It revealed that the composition of the bacteriophage community shifts

during the ripening process, aligning with the bacterial successions observed on the cheese surface

between lactic acid bacteria and ripening bacteria. Additionally, the main bacteriophages were found

consistently in cheese products made over a 4-years period, showing that they represent a persistent

component of the cheese-making environment. This research highlights the importance of considering

these bacteriophages when studying the microbial life of fermented foods like cheese.

INTRODUCTION

Smear-ripened cheeses are surface-ripened cheeses developing a typical viscous, red-orange smear on

their surface during ripening (1). In France, 14 out of the 46 PDO (Protected Designation of Origin)

cheeses belong to this category, *i.e.* Abondance, Beaufort, Chevrotin, Comté, Epoisses, Langres,

Livarot, Maroilles, Mont d'Or, Morbier, Munster, Ossau-Iraty, Pont l'Evêque and Reblochon. These

cheeses collectively accounted for 61% of the total annual volume of French PDO cheeses marketed,

with 128 kilotons in 2021 (https://www.inao.gouv.fr/eng/Publications/Donnees-et-cartes/Informations-

economiques). Due to their unique ripening process, involving frequent washes with saline and/or

alcoholic solutions, smear-ripened cheeses host a peculiar and diverse microbiota composed of lactic

acid bacteria (LAB, mainly starter cultures added at the beginning of the process for milk

acidification), yeasts and salt-tolerant bacteria belonging to the Actinomycetota, Bacillota and

Pseudomonadota phyla (1). The surface microbiota of smear-ripened cheeses is considered to be

responsible for the typical flavour and organoleptic properties of this type of cheese (2). From the past

two decades, numerous studies have been conducted to describe the composition of this microbiota

using culture-dependent methods (3, 4), molecular fingerprinting (5–8), and more recently amplicon-

based metagenomics also commonly referred to as metabarcoding (9–12).

Time series studies also reveals microbial successions occurring on the surface of smear-ripened

cheeses during the maturation process (8, 13, 14). Lactic acid bacteria (LAB), usually originating from

starter cultures, grow first in the milk and represent the dominant microorganisms in the curd. Yeasts,

e.g. Debaryomyces hansenii and *Geotrichum candidum*, which exhibit acid tolerance and metabolize

lactate, subsequently colonize the cheese surface, resulting in its deacidification. With the pH increase,

the establishment of a diverse bacterial community is progressively observed. The most common

bacterial taxa detected at the end of ripening on smear-ripened cheese belong to coryneform bacteria

(*e.g.* species of the *Glutamicibacter*, *Brevibacterium*, *Corynebacterium* or *Brachybacterium* genera),

Staphylococcus species and halophilic or halotolerant gram-negative bacteria (*e.g.* species of the

Psychrobacter, *Halomonas*, *Pseudoalteromonas, Hafnia, Vibrio*, *Pseudomonas* or *Proteus* genera)

(15).

Microbial successions in cheese during the ripening process result from the interplay between abiotic

factors, microbial interactions and complex ecological processes such as dispersion, diversification,

environmental selection, and ecological drift (16). While abiotic factors' impact on cheese microbial

community assembly has been extensively studied (17, 18), and some important microbial interactions

have been revealed as well (19–21), the effects of bacteriophages (or phages), *i.e.* viruses that infect

and replicate within bacteria, remain largely understudied. Their role in community assembly was

evidenced in many natural ecosystems (22) and recent work suggests the same applies to fermented

foods (23, 24). In the dairy industry, the impact of LAB phages is well documented because their lytic

activity can disturb the milk acidification step, causing delay in the production and even total loss of

production (25, 26). Consequently, the most studied dairy phages are *Lactococcus*, *Streptococcus*,

Lactobacillus and *Leuconostoc* phages, infecting the main starter cultures (27). Viral metagenomics

has been recently employed to characterize the bacteriophage communities in dairy samples, including

whey (28) and cheeses (29, 30). These investigations have demonstrated that such viral communities

are not restricted to LAB phages, but encompass a diverse array of phages that could potentially also

infect non-inoculated and ripening bacteria during cheese production. The recent application of

shotgun metagenomics combined with metaHiC on three washed-rind cheeses also enabled to

associate hundreds putative phage contigs with metagenome assembled genomes (MAGs) of their

bacterial hosts, particularly those belonging to Actinomycetota and Pseudomonadota phyla (14).

Beyond metagenomics, only a few studies have documented the isolation of such virulent phages

infecting ripening bacteria in cheese, *i.e.* targeting *Propionibacterium freudeunreichii* and

Brevibacterium aurantiacum (31–33). We recently isolated five additional virulent phages, targeting

Glutamicibacter arilaitensis (phages Montesquieu and Voltaire), *Brevibacterium aurantiacum* (phage

Rousseau), *Psychrobacter aquimaris* (phage d'Alembert) and *Leuconostoc falkenbergense* (phage

Diderot)*, from the surface of a French smear-ripened cheese suggesting that viral infection is likely to

occur on such ecosystems for most of the dominant bacteria (34). However, little is known about their

ecology.

In this study, we extended our investigations on the same smear-ripened cheese in order to evaluate

the impact of phage infections on community assembly. More specifically, our objective was to

enhance our understanding of the temporal distribution of bacteriophages and their bacterial hosts on

the cheese surface across two distinct timescales (Figure 1). On the short-term, cheeses were obtained

directly from the production facility at five distinct ripening stages over 28 days (three biological

replicates per stage), to analyze phage succession throughout the ripening process. On the long-term,

ready-to-eat cheeses, *i.e.* after storage and transport to the retail shop, of the same brand and variety

were sampled in 2017, 2019, and 2022 (three biological replicates per production year), to assess the

temporal variation of the phage community in this ecosystem over the years of production.

RESULTS

Overview of the cheese surface virome dataset.

The viral fraction was extracted from the surface of twenty-four smear-ripened cheese samples,

comprising 15 samples obtained during the ripening process to investigate the ecological succession of

phages (short-term study), and 9 samples collected post-transport and storage to examine the temporal

variation in the phage community across production years (long-term study) (Fig. 1). Subsequently,

the extracted viral fraction was subjected to viral metagenomics techniques. The experimental

procedures used for nucleic acids extraction and sequencing library preparation limit this study to the

description of double-stranded DNA viruses, excluding single-stranded DNA and RNA viruses.

The cheese viromes had an average sequencing depth of 11,585,587 sequencing reads and were

assembled into 5,753 contigs. Viral contigs were selected by considering 2 Kb as the minimum size

for viral sequence fragment and by using a combination of viral sequence detection tools (see material

and methods section). After clustering based on sequence homology, 331 species-level viral

operational taxonomic units (vOTUs) were obtained (Table 1). The vast majority of the vOTUs (*i.e.*

284) were detected in samples from both the short-term and long-term studies. The percentage of reads

mapping to these vOTUs reached 91.8% in average, revealing the low level of non-viral DNA

contamination in the studied viromes. Detailed information about the vOTUs including their size and

annotations are available in Table S1. The vOTUs sizes ranged from 2,002 to 144,530 bp, 79 contigs

(24%) being >10 Kb (Fig. 2A). This included 25 vOTUs (7.6%) flagged as complete genomes and 12

(3.6%) as high-quality genomes according to CheckV (35). Twenty-seven vOTUs (8.2%) were

identified as temperate phages by Vibrant (36) if they were excised from a larger contig or if they

encode an integrase. A bacterial host was successfully predicted for most vOTUs (272/331, i.e.

82.2%), spanning three bacterial phyla, namely Actinomycetota, Bacillota and Pseudomonadota (Fig.

2B). *Lactococcus* was the predicted genus containing the highest phage diversity (182 vOTUs),

followed by *Leuconostoc* (23 vOTUs). *Pseudoalteromonas* (18 vOTUs), *Psychrobacter* (17 vOTUs)

and *Brevibacterium* (6 vOTUs). Only 40.4% vOTUs (134/331) shared sequence homology (>30%

identity × coverage based on BLAT score) to a known dairy phage (Table S2). However, those vOTUs

represented 97% of the relative abundance observed in the dataset (Table 2) showing that most of the

dominant phages present in this ecosystem have already close-relatives that have been isolated and

characterized. Indeed, the most abundant detected phages matched either with the well characterized

Lactococcus phages from the 949 (*Audreyjarvisvirus* genus), 936 (*Skunavirus* genus) and P335 groups

or to the five virulent phages isolated from the same type of cheese (same variety, same brand) in our

previous work (34), namely *Glutamicibacter* phages Montesquieu and Voltaire, *Brevibacterium* phage

Rousseau, *Psychrobacter* phage D'Alembert and *Leuconostoc* phage Diderot (*Limdunavirus* genus).

Successional dynamics of the cheese virome throughout the ripening process.

While we have identified many phages in the cheese viromes, our next objective was to determine how

comparable the phage communities were across samples collected throughout the ripening process,

with a particular focus on each washing operation (short-term study). In the manufacturing process of

the studied cheese, the initial three washing operations (named W1, W2 and W3) utilize a NaCl

solution, while the last two operations (W4 and W5) utilize a NaCl solution containing increasing

concentrations of alcoholic liquor (Fig. 1). The dataset consisted in 15 samples (triplicate samples for

each washing operation) and 318 vOTUs (Table 1).

The median Shannon index across samples was 2.969 (ranging from 2.631 to 3.442). A slight decrease

in the viral diversity, as estimated by this alpha-diversity metric, was noted from W3 onwards.

However, the decrease was not statistically significant, suggesting that there was no major alteration of

the viral diversity related to the washing operations ($p > 0.05$, Kruskal-Wallis test; Fig. 3A).

An overall sparsity of 29% was observed on the abundance table reflecting that all the studied cheese

viromes shared most of the detected vOTUs. However, structural variation in the virome composition

was assessed by computing Bray-Curtis dissimilarity (Fig. 3B) and the PERMANOVA test indicated a

significant effect of the washing operation ($p = 0.001$, $r^2 = 0.658$).

To further investigate this result, we selected vOTUs with a normalized relative abundance above

0.005 % in average (168 contigs) and represented their distribution across samples through a heatmap

(Fig. 3C). We observed a clear structure in the vOTUs distribution across our samples. Based on the

total within sum of square (Fig. S1), we estimated to five the optimal number of vOTUs clusters

present in the dataset (labelled I to V in Fig. 3C). Clusters I and II were characterized by vOTUs

whose abundance did not vary with the ripening and that corresponded mainly to virulent *Lactococcus*

phages belonging to the 936 group (*Skunavirus* genus). Cluster V contained vOTUs whose abundance

decreased with ripening and that corresponded to other *Lactococcus* phages, close to the P335 group

containing both temperate and virulent members. Finally, clusters III and IV were essentially

composed of a few vOTUs that increased in relative abundance with ripening. These mainly

corresponded to phages targeting ripening bacteria such as *Brevibacterium*, *Glutamicibacter*,

Pseudoalteromonas and *Vibrio*, and non-starter lactic acid bacteria (NSLAB) such as *Leuconostoc*.

These few vOTUs corresponded to uncharacterized phages, except *Glutamicibacter* phage

Montesquieu, *Brevibacterium* phage Rousseau and *Leuconostoc* phage Diderot which we previously

isolated from the same type of cheese (34).

Both the principal coordinate analysis of the Bray-Curtis dissimilarity (Fig. 3B) and the heatmap

visualization (Fig. 3C) revealed a comparable virome composition between samples collected from

W1 and W2, as well as between those from W4 and W5. This observation suggests the succession of

two predominant viral communities on the cheese surface during the ripening process, corresponding

to these respective stages. Samples from W3 displayed a unique composition, wherein Cluster III and

IV were found in higher relative abundance compared to W1-W2 but not as abundant as in W4-W5.

Cluster V was also detected in lower abundance than in W1-W2 but at a similar level as in W4-W5.

The virome composition captured by those samples is therefore likely to reflect a transition stage.

We finally applied the DESeq2 method (37, 38) in order to statistically identify which vOTUs

differentiate the two predominant viral communities represented by W1-2 and W4-5 samples (Fig.

S2). In total, 22 vOTUs were differentially abundant between the two communities (adjusted $p < 0.05$,

log2 fold change >3 or <-3). Interestingly, two groups of vOTUs emerged, with readable outcomes:

vOTUs corresponding to phages infecting starter cultures, such as *Lactococcus* and *Streptococcus*

phages, had a negative log2 fold change meaning their relative abundances significantly decreased

during the ripening process. Conversely, vOTUs of virulent phages targeting NSLAB and ripening

bacteria (e.g. *Brevibacterium, Glutamicibacter, Pseudoalteromonas* and *Vibrio*) as well as unknown

phages were significantly more abundant in W4-5 samples compared to W1-2 samples (positive log2

fold change) reflecting a higher population level for such phages on the cheese surface in later

ripening stages. Among them were two vOTUs with high similarity to the genomes of

Glutamicibacter phage Montesquieu (2017-3_NODE2; 47,703 bp; 100% identity $\times$ coverage by

BLAT), and *Brevibacterium* phage Rousseau (2022-5_NODE4; 41,077 bp; 54% identity $\times$ coverage

by BLAT but 98% identity at the nucleotidic level over a portion of >9 kb). In this group, we also

detected a vOTU partially related to *Brevibacterium* phage AGM1 (SA2-0_NODE10; 37148 bp; 3.5%

identity $\times$ coverage by BLAT but 88% identity at the nucleotidic level over a portion of 1324 nt),

which was isolated from a Canadian washed-rind cheese (32).

Relationships between cheese virome and bacteriome throughout the ripening process.

Having observed a succession of viral communities on the cheese surface during the ripening process,

we sought to investigate whether it was associated with bacterial community changes. We therefore

profiled bacterial communities on the same samples using 16S metabarcoding. A total of 773,059

reads were obtained and clustered into 195 operational taxonomic units (OTUs) assigned to 14

different bacterial genera (Fig. 4A). As observed for viruses, the composition of the bacteriome of the

cheese surface varied through the ripening process and a clear transition in the bacterial community

structure was detected from W3 onwards. *Lactococcus* was the dominant genus in samples from W1

and W2, and was progressively replaced by typical surface aerobic bacteria such as members of

Psychrobacter, *Vibrio*, *Glutamicibacter* and *Pseudoalteromonas* genera.

The metabarcoding data, providing relative proportions, was complemented by bacterial counts to

introduce a quantitative aspect into the bacterial dynamics (Fig. 4 B and C). We observed an

approximately 2-log increase in aerobic bacteria between W1 and W5, rising from $\sim 10^{8}$ to $\sim 10^{10}$

CFU/g. In contrast, lactic acid bacteria remained stable over time ($\sim 10^{8}$ CFU/g) throughout the entire

experiment. Based on these results, the observed change in the relative abundance of *Lactococcus*

sequences through metabarcoding was primarily attributed to the growth of aerobic bacteria.

Then, a Mantel test, comparing two distance matrices, demonstrated the statistical relationships

between the virome and bacteriome datasets ($p = 0.001$; $r = 0.3151$; 999 permutations). This result

indicates that changes observed on the phage community structure is associated with bacterial

successions on the cheese surface. We further calculated spearman correlations between the relative

abundances of the main phage groups identified in the metavirome and of bacterial genera in the

bacteriome in order to identify co-abundance or mutual-exclusion patterns (Fig. 4D). Phages targeting

ripening bacteria, such as *Psychrobacter* phage d'Alembert, *Glutamicibacter* phage Montesquieu and

Brevibacterium phage Rousseau exhibited positive correlations with bacterial genera characteristics of

the surface flora of smear-ripened cheese such as *Glutamicibacter*, *Mesonia*, *Psychrobacter*,

Pseudoalteromonas and *Marinomonas*. <u>Conversely, these phages were negatively associated with</u>

<u>associa</u>ted with *Lactococcus*, *Leuconostoc* and *Ruminococcus*. Notably, an opposite pattern emerged

for LAB phages belonging to KSY1, P335 and 987 groups, <u>which are recognized for infecting</u>

<u>acidifying starter cultures such as *Lactococcus lactis* and/or *Streptococcus thermophilus*.</u>

Temporal variation of the cheese virome across production years.

As phage succession was evidenced in the short-term, specifically during the ripening process, we next

wanted to test whether the composition of the virome remains comparable on the long-term

perspective. To address this, we analysed the long-term dataset comprising 297 vOTUs detected on

nine cheese samples (Table 1). Theses samples were collected after ripening, transport and storage

across three different productions years (Fig. 1). We first focused on determining the proportion of

vOTUs shared across production years. Utilizing the full dataset without relative abundance filtering,

we observed that 56.6% of the vOTUs (168) were consistently detected in all production years, with

75.4% (224) observed in at least two different production years (Fig. 5A). Notably, samples from the

year 2022 had a higher number of unique vOTUs (48, 32.1% of the total) compared to samples from

the years 2017 and 2019 (3 and 11, respectively). We also conducted the same analysis focusing on

the most abundant vOTUs, defined by an average normalized relative abundance > 0.005% (99

vOTUs in total) (Table 1, Fig. 5B). A substantial majority of the vOTUs, *i.e.* 89.9%, were shared

across all three production years and this value rose to 97% when considering only two different years.

This result underscores the stability of the cheese surface virome in terms of composition

(presence/absence) and emphasizes the long-term persistence of dominant phages in the studied

ecosystem.

The effect of the production year (2017, 2019 and 2022) on the structure of the cheese virome was

further investigated by computing Bray-Curtis dissimilarity on the vOTU abundance table without

relative abundance filtering (Fig. 5C). The principal coordinate analysis revealed a distinct grouping of

samples consistent with the production years. The first axis clearly discriminates samples from year

2019 and samples from years 2017 and 2022. The second axis discriminates samples from years 2017

and 2022. The PERMANOVA test confirmed a significant effect of the production year ($p = 0.002$, R^2

$= 0.814$) on the distance metrics indicating structural variations of the phage community across

production years. Given the consistent presence of the most abundant vOTUs across all production

years (Fig. 5B), such variations could be attributed to the fluctuations in the presence/absence of

phages with lower abundance, as well as to the distribution of dominant phages in terms of relative

abundances.

Heatmap visualization was then used to identify phages explaining such structural variations (Fig. 5E)

in combination with differential abundance analysis (Fig. S3). Whereas the general profiles of all the

viromes were very similar, a few vOTUs were more specifically detected in one or two production

years. For example, the second most abundant vOTU in 2017 and 2019 (SS2-4_NODE1) which is

highly similar to *Lactococcus* phage 949 (*Audreyjarvisvirus* genus) was statistically less abundant in

2022 (adjusted p <0.05, log2 fold change <-3) contrary to a group of other distinct vOTUs also

predicted to infect *Lactococcus* but either not identified or assigned to the 936 (*Skunavirus* genus) and

P335 groups. Regarding *Psychrobacter* phages, the vOTU closely related to phage d'Alembert (2017-

0_NODE1) was more detected in 2017 than on the two other years (adjusted p <0.05, log2 fold change

>3). However, two vOTUs without sequence homology with known dairy phages but predicted to

infect also *Psychrobacter* (2022-0_NODE3 and 2022-5_NODE3) were detected later in 2022 in higher

abundance (adjusted p <0.05, log2 fold change >3). Notably, *Glutamicibacter* phage Voltaire was

present in low abundance in 2017 and 2019 (0.036% and 0.004% relative abundance in average,

respectively) but no more detected in 2022.

We finally performed 16S metabarcoding analysis in order to determine whether structural variations

were observed in the composition of the bacterial community as well. The surface of the studied

smear-ripened cheese was characterized by ten dominant bacterial genera. Similar to what we

observed for phages, these genera were consistently detected in all samples (Fig. 5D). However,

variations in their relative abundance were detected across the three production years.

Together, our results indicate the presence of a core microbiome consisting of dominant

microorganisms and their associated viruses that define the surface of the cheese under study. The

composition of this core microbiome seems to persist in the long-term timescale, although changes in

the microbial balance can be observed between production batches,

DISCUSSION

Recent studies of cheese samples using viral metagenomics (29, 30) or exploration of cheese microbial

metagenomes (14, 39) have revealed that the cheese environment harbours a diverse bacteriophage

community whose targets go beyond lactic acid starter cultures. The previous isolation of a few

representatives of these non-starter phages (32–34) suggests that phage infection occurs in this

ecosystem during the ripening process. Recently, the deliberate addition of one of those phages,

Brevibacterium phage AGM9, was proven to slow down the development of the orange rind color in a

model system mimicking a smear-ripened cheese (40). In the present study, we described the temporal

variation of the viral community of a French smear-ripened cheese at two timescales: short-term (28

days corresponding to duration the ripening process) and long-term (spanning four years of

production). The virome was predominantly composed of vOTUs associated with a variety of

Lactococcus phages as well as the *Glutamicibacter* phage Montesquieu, a virulent phage we had

previously isolated from the same cheese variety (34). This phage targets the ripening bacterium

Glutamicibacter arilaitensis. Several other known dairy phages, such as *Brevibacterium* phage

Rousseau, *Leuconostoc* phage Diderot, *Psychrobacter* phage d'Alembert, *Glutamicibacter* phage

Voltaire, and a novel vOTU displaying minimal sequence homology to *Brevibacterium* phage AGM1

were also detected, but at lower abundance. Notably, while known dairy phages represented 97% of

the relative abundance in our dataset, 197 vOTUs (59.6% of the total) remained unidentified and 49

(17.8%) had no predicted host. The diversity of these low-abundant phages is therefore not yet

completely sampled and, in particular, our results underscores the necessity to isolate a more

comprehensive collection of phages from cheese and other fermented foods in general. For example, a

few vOTUs partially aligned with genomic sequences from phages infecting halotolerant bacteria but

isolated from natural environments (e.g., *Pseudoalteromonas*, *Halomonas*, *Vibrio*, and *Proteus*

species). Even though these bacteria are not intentionally inoculated into smear-ripened cheese, they

are frequently detected in such products and tend to dominate by the end of the ripening process (15).

Their phages should therefore be looked for in future isolation initiatives.

The dynamics of bacterial and fungal communities has been extensively studied in a wide diversity of

cheese products during ripening (13, 41–45), enabling us to accurately describe successions of cellular

organisms occurring in cheese production processes (16). In this study, we present the first analysis of

viral dynamics throughout cheese ripening. Similar to the observed transitions in bacteria and fungi,

there was a distinct shift in viral composition over the course of the ripening process. Some vOTUs

associated with LAB starter phages, primarily targeting *Lactococcus lactis*, were progressively

replaced by vOTUs specific to phages that infect ripening bacteria. When comparing phage to

bacterial dynamics, there was a significant relationship between phage trajectories and bacterial

successions on the cheese surface. Specifically, the relative abundance of several phages, such as

Glutamicibacter phage Montesquieu, *Psychrobacter* phage d'Alembert and *Lactococcus* phages

belonging to the 936 (*Skunavirus* genus), KSY1 and P335 groups was positively correlated to the

relative abundance of their predicted bacterial hosts. This co-occurrence pattern raises questions about

the overall impact of phages on bacterial successions in cheese during ripening. However, it is

important to note that phage susceptibility is generally strain-dependent, as demonstrated in previous

studies on LAB strains (summarized in (46)) and in undefined starter cultures where multiple strains

of the same species stably co-exist with distinct phages (47–49). The narrow host range observed for

phages infecting ripening bacteria recently isolated from cheese rind also corroborates this result (32,

34). Unfortunately, in the present work the composition of the bacterial community was assessed at

the genus level, which prevent capturing such strain-level diversity. Future investigations at such level

would therefore be highly relevant to better understand the ecological role of phages in cheese rind

microbial communities.

Surprisingly, despite the changes in bacterial and viral communities' composition during the

production process, the relative abundance of some vOTUs remained very stable during ripening.

They mainly corresponded to virulent *Lactococcus* phages, belonging to the *Skunavirus* genus

(formerly 936 group). Among *Lactococcus* phages, this genus is by far the most frequently detected in

the dairy industry (50). Given that *Lactococcus lactis* predominantly proliferates during milk

acidification and maintains consistent concentration levels throughout the ripening period in this type

of cheese (13), we theorize that the maintenance of *Skunavirus* is indicative of the stability of phage

particles produced at the onset of cheese maturation. The remarkable stability of *Skunavirus* particles

in comparison to other *Lactococcus* phage groups, especially the P335 group, has been reported earlier

(51, 52). In contrast, we suggest that the relative decline observed in our experiment for certain

Lactococcus phages, specifically those affiliated to the P335 group, denotes their temporal instability.

Nevertheless, a more comprehensive examination of this phenomenon warrants dedicated

investigations.

Microbial communities of washed-rind cheese are generally dominated by environmental

microorganisms detected in processing environments, the so-called "house" microbiota (53), which is

specific to each production facility and provides a microbial signature distinguishing cheeses

belonging to the same variety but manufactured by distinct producers (10). Recently, the analysis of

several Quebec's terroir cheeses revealed that the dominant microorganisms remain stable from year

to year, which could be linked to typical manufacturing practices and consistency in the use of starter

and ripening cultures by cheesemakers (54). Here, we describe a similar observation for

bacteriophages. We indeed identified a core-virome composed of the most abundant vOTUs,

consistently detected in three different years of production. Previous work on an undefined starter

culture used for the production of a Swiss-type cheese, propagated for decades in the same dairy

environment, revealed that phages and bacteria stably coexist over time in this system and suggests

that this may contribute to the stable maintenance of the cheese starter culture over years (47). Phages

and ripening bacteria present on the surface of natural smear-ripened cheese are contaminating the

cheese production environment, and are therefore likely to repeatedly contaminate cheese from one

production cycle to another (34, 53). Nevertheless, further research is needed to elucidate the

contributions of phages to both the long-term maintenance of the strain-level diversity of ripening

bacteria, and the occurrence of punctual microbial disequilibrium leading to organoleptic defects.

These findings may be specific to the type of cheese under study, the strain-level diversity of the main

bacteria, and the operational conditions used for cheese production.

In conclusion, the observed dynamics of the cheese virome throughout the ripening process, coupled

with the persistence of the most dominant phages across production years, support an important role of

bacteriophages in the cheese microbial ecosystem. The acquisition of new knowledge about phage

ecology is now essential for a comprehensive understanding of microbial successions during milk

fermentation. Moreover, this knowledge may offer cheesemakers novel avenues to refine and control

their production processes.

MATERIAL AND METHODS

Cheese samples.

(i) Short-term study design: French smear-ripened cheeses, all from the same production batch, were

collected directly from a cheese plant at five different stages during the ripening process (Figure 1).

Three different cheeses were sampled per stage, and considered as biological replicates. The stages,

labeled W1 to W5, correspond to distinct washing steps. The initial three washes utilized a NaCl

solution, while the final two employed a NaCl solution supplemented with increasing concentrations

of alcoholic liquor (standard alcohol concentration of 40%). At each stage, three distinct cheeses were

sampled and designated as biological replicates A, B, and C. Cheeses were immediately stored at 4°C

after sampling and processed within 48h.

(ii) Long-term study design: ready-to-consume cheeses of the same type and same brand as the

previously described cheeses were purchased in a local supermarket in December 2017, November

2019 and February 2022, spanning >4 years of production (Figure 1). Three different cheeses, with the

same production date, were sampled each year and used as biological replicates. Cheeses were

immediately stored at 4°C after sampling and processed within 48h.

For both short-term and long-term studies, cheese samples were analysed immediately after reception

at the lab. Using sterile knives, the rind, approximately 2-3 mm thick, was carefully separated from the

core. It was then blended and processed for microbial counts, viral DNA extraction for metavirome

analysis, and microbial DNA extraction for extensive amplicon sequencing targeting the 16S rRNA

gene, which will be subsequently referred to as 16S metabarcoding.

Microbiological analysis.

Bacteria were enumerated by plating serial dilutions (10^{-1} to 10^{-7}) of one gram of cheese rind mixed in

9 mL of physiological water (9 g/L NaCl) on two different culture media as described in our previous

study (29). Brain Heart Infusion Agar (BHI, Biokar Diagnostics) supplemented with 50 mg/L

amphotericin (Sigma Aldrich, Saint-Louis, MO, USA) was used to count total aerobic bacteria after 48

h of incubation at 28°C. Man, Rogosa and Sharpe Agar (MRS, Biokar Diagnostics, Allonne, France)

supplemented with 50 mg/L amphotericin was used to count lactic acid bacteria after 48 h of

incubation at 30°C under anaerobic conditions.

Viral DNA extraction and metavirome analysis.

Extraction of the viral fraction from cheese rind was performed according to protocol P4 detailed in

(29) including a filtration step and a chloroform treatment. DNA was extracted from the viral particles

according to the protocol described in the same study and sent to Eurofins Genomics for high

throughput sequencing using the Illumina NovaSeq platform (2 × 150 bp paired-end reads,

approximately 10 million reads per sample).

All the details about the tools, versions and parameters used in the following pipeline are available in

scripts deposited in the GitLab repository (https://forgemia.inra.fr/eric.dugat-bony/cheese_virome).

Briefly, raw reads were quality filtered using Trimmomatic v0.39 (55). Then a single assembly was

computed for the collection of triplicate reads from each sample with Spades v3.15.3 (56), using either

the complete dataset of trimmed reads available or after subsampling the dataset to 1.5 million,

150,000 or 15,000 trimmed reads per sample. We noted that some abundant contigs were assembled

into longer, nearly complete contigs, after subsampling. Among the 5,753 contigs, those >2kb were

selected and clustered following an approach adapted from (57). Succinctly, a pairwise alignment was

first performed for all contigs using BLAT (58). Then, contigs with a self-alignment > 110% of contig

length, corresponding to chimeras, were removed. Remaining contigs were clustered at the species

level (90% identity × coverage) and the longest contig within each cluster was selected as the

representative sequence. The final contig dataset consisted in 3,122 dereplicated contigs.

Among them, 332 were identified of viral origin (vOTUs) by using a combination of three detection

tools: VIBRANT v1.2.1 (36), VirSorter2 v2.2.4 (59) and CheckV v0.8.1 (35). To be declared as viral,

a contig had to meet at least one of the following criteria: declared "complete", "high" or "medium"

quality by either VIBRANT or CheckV, declared "full" by VirSorter2. The bacterial host of the

vOTUs was predicted using iPHoP (60). Finally, all sequences were compared by BLAT to an in-

house database consisting of genome sequences from 32 common dairy phages (listed in Table S1) in

order to identify potential related phages with known taxonomy, verified host and lifestyle (30%

identity × coverage minimal cutoff). When appropriate, the host genus predicted by iPHoP was

replaced by the genus of the bacterial host of the closest relative phage identified by the BLAT search.

One vOTU, corresponding to the genome of the phage PhiX174 which is routinely used as control in

Illumina sequencing runs to monitor sequencing quality, was discarded resulting in a final dataset of

331 vOTUs.

In order to evaluate the relative abundance of each vOTU in each metavirome sample, trimmed reads

were mapped against the vOTUs using bwa-mem2 v2.2.1 (61) and counted with Msamtools v1.0.0

profile with the options --multi=equal --unit=ab –nolen (https://github.com/arumugamlab/msamtools).

The output files from all samples were joined into an abundance table and processed using the R

package phyloseq v1.38.0 (62). For each experimental dataset (short-term and long-term studies), read

counts were rarefied to the minimum depth observed in one individual sample. To facilitate

comparison of abundance levels among vOTUs, read counts were divided by contig length resulting in

an abundance table containing the mean coverage per vOTU in each sample. Subsequently,

normalization of the table was conducted using the total sum scaling method (TSS) (63), which simply

transforms the abundance table into relative abundance table through scaling by each sample's library

size, thus enabling comparisons between samples. Making the assumption that one vOTU corresponds

to one species (which is wrong each time several contigs belonging to the same phage genome are

present), Bray-Curtis dissimilatory index, as computed by the distance function from the R package

phyloseq, was used to compare viral communities between samples and the effect of different

variables on their structure was assessed using permutational analysis of variance as computed by the

adonis2 function from the R package vegan v2.6-2.

For the dynamic dataset, differential analysis was performed using the DESeq function implemented

in the R package DESeq2 v1.38.0 (37) and for which an official extension exists within the R package

phyloseq (https://joey711.github.io/phyloseq-extensions/DESeq2.html). This method was shown to

perform well for the detection of differentially abundant taxa in microbiome data (38). Because the

DESeq function already includes a normalization step (by the median of ratios method), this analysis

started from raw counts. The Wald test was used for comparing the abundance of all vOTUs between

two sample groups and vOTUs were considered differentially abundant if the p value was > 0.01 after

correction for multiple testing using the Benjamini and Hochberg method, log2 fold change was > 3 or

☐ -3 and average raw counts was > 1500.

Microbial DNA extraction and 16S metabarcoding profiles.

Total DNA extraction from the cheese surface was performed as previously described (10). PCR

amplification of the V3-V4 regions of the 16S rRNA gene was performed with the primers V3F (5'-

ACGGRAGGCWGCAG-3')   and   V4R   (5'-TACCAGGGTATCTAATCCT-3')

carrying   the   Illumina   5'-CTTTCCCTACACGACGCTCTTCCGATCT-3'   and   the   5'-

GGAGTTCAGACGTGTGCTCTTCCGATCT-3' tails, respectively. The reaction was performed

using 10 ng of extracted DNA, 0.5 μM primer, 0.2 mM dNTP, and 2.5 U of the MTP Taq DNA

polymerase (Sigma-Aldrich, USA). The amplification was carried out using the following program:

94°C for 60 s, 30 cycles at 94°C for 60 s, 65°C for 60 s, 72°C for 60 s, and a final elongation step at

72°C for 10 min. The resulting PCR products were sent to the @BRIDGe platform (INRAE, Jouy-en-

Josas, France) for library preparation and sequencing. Briefly, amplicons were purified using a

magnetic beads CleanPCR (Clean NA, GC biotech B.V., The Nederlands), the concentration was

measured using a Nanodrop spectrophotometer (Thermo Scientific, USA) and the amplicon quality

was assessed on a Fragment Analyzer (AATI, USA) with the reagent kit ADNdb 910 (35-1,500 bp).

Sample multiplexing was performed by adding tailor-made 6 bp unique indexes during the second

PCR step which was performed on 50–200 ng of purified amplicons using the following program:

94°C for 10 min, 12 cycles at 94°C for 60 s, 65°C for 60 s, 72°C for 60 s, and a final elongation step at

72°C for 10 min. After purification and quantification (as described above), all libraries were pooled

with equal amounts in order to generate equivalent number of raw reads for each library. The DNA

concentration of the pool was quantified on a Qubit Fluorometer (Thermofisher Scientific, USA) and

adjusted to a final concentration between 5 and 20 nM for sequencing. The pool was denatured (NaOH

0.1N) and diluted to 7 pM. The PhiX Control v3 (Illumina, USA) was added to the pool at 15% of the

final concentration, as described in the Illumina procedure, and the mixture was loaded onto the

Illumina MiSeq cartridge according to the manufacturer's instructions using MiSeq Reagent Kit v3 (2

× 250 bp paired-end reads).

Paired-end reads were analysed using FROGS v3.2 (64), according to the standard operating

procedure. Briefly, operational taxonomic units (OTUs) were built using Swarm with an aggregation

distance of 1 and the --fastidious option (65), and each OTU that accounted for <0.005% of the total

dataset of sequences was discarded, as previously recommended (66). Lastly, OTU affiliation was

checked using the EzBiocloud database v52018 (67). The abundance table was processed using the R

package phyloseq v1.38.0 (62).

Data availability.

Raw sequencing data for cheese virome and bacterial microbiome were deposited at the sequence read

archive (SRA) of the NCBI (https://www.ncbi.nlm.nih.gov/sra/) as part of bioprojects PRJNA984302

(short-term study) and PRJNA984735 (long-term study).

ACKNOWLEDGMENTS

T.P. is the recipient of a doctoral fellowship from the French Ministry of Higher Education, Research

and Innovation (MESRI) and the MICA department of the French National Research Institute for

Agriculture, Food and Environment (INRAE). For the metabarcoding analysis, this work has benefited

from the facilities and expertise of @BRIDGe (Université Paris-Saclay, INRAE, AgroParisTech,

GABI, 78350 Jouy-en-Josas, France). We are also grateful to the INRAE MIGALE bioinformatics

facility (MIGALE, INRAE, 2020. Migale bioinformatics Facility, doi:

10.15454/1.5572390655343293E12) for providing computing and storage resources.

TABLES

Table 1: summary statistics about the metavirome dataset.

	Complete dataset (24 samples)	Short-term study (15 samples)	Long-term study (9 samples)
Total number of vOTUs > 2 Kb	331	318	297
Number of vOTUs with relative abundance > 0.005%	157	168	99
Percentage of reads mapping against vOTUs (± SD)	91.8 ± 12.1	94.3 ± 7.0	87.6 ± 17.4

499 **Table 2: Most abundant dairy phages detected on the cheese metavirome.** Phages that have been

500 previously isolated from the same cheese variety are highlighted in bold. When classification is

501 available at the International Committee on Taxonomy of Viruses (ICTV), the viral genus is denoted

502 in brackets.

Closest relatives	Relative abundance*
Lactococcus 949 group *(Audreyjarvisvirus)*	0.56
***Glutamicibacter* Montesquieu**	**0.23**
Lactococcus 936 group *(Skunavirus)*	0.18
Lactococcus P335 group	1.54E-03
***Brevibacterium* Rousseau**	**1.08E-03**
***Psychrobacter* D'Alembert**	**1.74E-04**
Leuconostoc* Diderot *(Limdunavirus)	**1.14E-04**
***Glutamicibacter* Voltaire**	**1.70E-05**
Streptococcus 987 group	7.87E-06
Lactococcus KSY1 group *(Chopinvirus)*	5.94E-07
Non dairy phages**	0.03

*The relative abundance value used here is not normalized by genome size, it corresponds to the

relative proportion of reads mapping to all vOTUs belonging to a given phage group present in our in-

house dairy phage database (Table S1).

**vOTUs without any match to our in-house dairy phage database (30% identity × coverage minimal

cutoff).

FIGURE LEGENDS

Figure 1: Schematic representation of the experimental plan followed for the study of temporal

variation in the cheese virome composition. For the short-term study, cheese samples were collected

in triplicate after each washing operation during the ripening process (blue arrows indicate washes

performed with a NaCl solution, while red arrows indicate washes performed with a NaCl solution

containing alcoholic liquor). For the long-term study, cheese samples were collected in triplicate after

storage and transport at three different production years. All samples were analysed through viral

metagenomics, 16S metabarcoding and microbial plate counts.

Figure 2: Characteristics of the cheese virome dataset. A) Size distribution of the vOTUs. B)

Distribution of the vOTUs per bacterial host genus.

Figure 3: Viral succession on the cheese surface throughout the ripening process. A) Comparison

of the viral diversity, estimated by the Shannon index, at the different ripening steps. B) Principal

coordinate analysis of the Bray-Curtis dissimilarity. Samples are coloured according the washing

operation. C) Heatmap representing the normalized relative abundance of the most abundant vOTUs in

the different samples. The colours on the heatmap represent the log-transformed relative abundance,

and range from blue (low relative abundance) to red (high relative abundance). Host genus, phage

group and lifestyle of the different vOTUs are indicated when available. Hierarchical clustering using

the complete linkage method was used to define the order of rows (vOTUs). The five vOTUs clusters

detected in the dataset are boxed in red and labelled I to V.

Figure 4: Relationship between phage succession and the composition of cheese bacterial

community during the ripening process. A) Composition of the bacterial community assessed by a

metabarcoding approach targeting the V3-V4 regions of the 16S rRNA gene. Three different cheeses

from the same batch were analysed at each sampling point. Data were aggregated at the genus level.

B) Aerobic bacteria counts expressed in log(CFU/g). C) Lactic acid bacteria counts expressed in

log(CFU/g). D) Heatmap of Spearman correlations (r values, ranging from -1 to +1) describing the

relationship between the main phage groups detected in the virome dataset and bacterial genus-level

relative abundances. Hierarchical clustering using the complete linkage method was used to define the

order of rows (bacterial genera) and columns (phage groups).

Figure 5: Temporal variation of the cheese surface virome across production years. A) Venn

diagram constructed from the dataset without filtering, comprising 297 vOTUs. B) Venn diagram

constructed from the filtered dataset comprising 99 vOTUs with an averaged normalized relative

abundance value > 0.005%. C) Principal coordinate analysis of the Bray-Curtis dissimilarity. Samples

are coloured according the production year (2017, 2019 and 2022). D) Composition of the bacterial

community assessed by a metabarcoding approach targeting the V3-V4 regions of the 16S rRNA gene.

Three different cheeses from the same production year were analysed. Data were aggregated at the

genus level. E) Heatmap representing the normalized relative abundance of the most abundant vOTUs

in each sample. The colours on the heatmap represent the log-transformed relative abundance, and

range from blue (low relative abundance) to red (high relative abundance). Host genus, phage group

and lifestyle of the different vOTUs are indicated when available. Hierarchical clustering using the

complete linkage method was used to define the order of rows (vOTUs).

SUPPLEMENTARY MATERIAL

Table S1. Detailed annotation of the viral operational taxonomic units.

Table S2. Common dairy phages and associated information.

Figure S1: Estimation of the optimal number of vOTUs clusters in the short-term dataset. The

total within-cluster-sum-of-square was calculated for each number of clusters and the elbow point was

used to determine the optimal number of clusters.

Figure S2: Differential abundance analysis of the virome data between the two stable stages

represented by W1-2 and W4-5 samples (short-term study). Log2 fold change values obtained for

differential abundant vOTUs (dots) are represented and the dot colour indicate the predicted host

genus.

Figure S3: Differential abundance analysis of the virome data between the three years of

production (long-term study). A) Comparison between years 2017 and 2019. B) Comparison

between years 2017 and 2022. C) Comparison between years 2019 and 2022. Log2 fold change values

obtained for differential abundant vOTUs (dots) are represented and the dot colour indicate the

predicted host genus.

REFERENCES

1. Mounier J, Coton M, Irlinger F, Landaud S, Bonnarme P. 2017. Chapter 38 - Smear-Ripened

Cheeses, p. 955–996. *In* McSweeney, PLH, Fox, PF, Cotter, PD, Everett, DW (eds.), Cheese

(Fourth Edition). Academic Press, San Diego.

2. Corsetti A, Rossi J, Gobbetti M. 2001. Interactions between yeasts and bacteria in the smear

surface-ripened cheeses. International Journal of Food Microbiology 69:1–10.

3. Larpin-Laborde S, Imran M, Bonaïti C, Bora N, Gelsomino R, Goerges S, Irlinger F, Goodfellow M,

Ward AC, Vancanneyt M, Swings J, Scherer S, Guéguen M, Desmasures N. 2011. Surface

microbial consortia from Livarot, a French smear-ripened cheese. Canadian Journal of

Microbiology 57:651–660.

4. Lavoie K, Touchette M, St-Gelais D, Labrie S. 2012. Characterization of the fungal microflora in

raw milk and specialty cheeses of the province of Quebec. Dairy Sci & Technol 92:455–468.

5. Feurer C, Irlinger F, Spinnler H e., Glaser P, Vallaeys T. 2004. Assessment of the rind microbial

diversity in a farmhouse-produced vs a pasteurized industrially produced soft red-smear cheese

using both cultivation and rDNA-based methods. Journal of Applied Microbiology 97:546–556.

6. Mounier J, Monnet C, Jacques N, Antoinette A, Irlinger F. 2009. Assessment of the microbial

diversity at the surface of Livarot cheese using culture-dependent and independent

approaches. International Journal of Food Microbiology 133:31–37.

7. Mounier J, Gelsomino R, Goerges S, Vancanneyt M, Vandemeulebroecke K, Hoste B, Scherer S,

Swings J, Fitzgerald GF, Cogan TM. 2005. Surface Microflora of Four Smear-Ripened Cheeses.

Appl Environ Microbiol 71:6489–6500.

8.   Rea MC, Görges S, Gelsomino R, Brennan NM, Mounier J, Vancanneyt M, Scherer S, Swings J,

Cogan TM. 2007. Stability of the Biodiversity of the Surface Consortia of Gubbeen, a Red-Smear

Cheese. Journal of Dairy Science 90:2200–2210.

9.   Delcenserie V, Taminiau B, Delhalle L, Nezer C, Doyen P, Crevecoeur S, Roussey D, Korsak N,

Daube G. 2014. Microbiota characterization of a Belgian protected designation of origin cheese,

Herve cheese, using metagenomic analysis. Journal of Dairy Science 97:6046–6056.

10. Dugat-Bony E, Garnier L, Denonfoux J, Ferreira S, Sarthou A-S, Bonnarme P, Irlinger F. 2016.

Highlighting the microbial diversity of 12 French cheese varieties. International Journal of Food

Microbiology 238:265–273.

11. Quigley L, O'Sullivan O, Beresford TP, Ross RP, Fitzgerald GF, Cotter PD. 2012. High-throughput

sequencing for detection of subpopulations of bacteria not previously associated with artisanal

cheeses. Applied and Environmental Microbiology 78:5717–5723.

12. Wolfe BE, Button JE, Santarelli M, Dutton RJ. 2014. Cheese rind communities provide tractable

systems for in situ and in vitro studies of microbial diversity. Cell 158:422–433.

13. Irlinger F, Monnet C. 2021. Temporal differences in microbial composition of Époisses cheese

rinds during ripening and storage. Journal of Dairy Science 104:7500–7508.

14. Saak CC, Pierce EC, Dinh CB, Portik D, Hall R, Ashby M, Dutton RJ. 2023. Longitudinal, Multi-

Platform Metagenomics Yields a High-Quality Genomic Catalog and Guides an In Vitro Model

for Cheese Communities. mSystems 8:e00701-22.

15. Kothe CI, Bolotin A, Kraïem B-F, Dridi B, Renault P. 2021. Unraveling the world of halophilic and

halotolerant bacteria in cheese by combining cultural, genomic and metagenomic approaches.

International Journal of Food Microbiology 358:109312.

16.   Mayo B, Rodríguez J, Vázquez L, Flórez AB. 2021. Microbial Interactions within the Cheese

618   Ecosystem and Their Application to Improve Quality and Safety. 3. Foods 10:602.

619   17.   De Filippis F, Genovese A, Ferranti P, Gilbert JA, Ercolini D. 2016. Metatranscriptomics reveals

620   temperature-driven functional changes in microbiome impacting cheese maturation rate. Sci

621   Rep 6.

622   18.   Dugat-Bony E, Bonnarme P, Fraud S, Catellote J, Sarthou A-S, Loux V, Rué O, Bel N, Chuzeville S,

623   Helinck S. 2019. Effect of sodium chloride reduction or partial substitution with potassium

624   chloride on the microbiological, biochemical and sensory characteristics of semi-hard and soft

625   cheeses. Food Research International 125:108643.

626   19.   Irlinger F, Mounier J. 2009. Microbial interactions in cheese: implications for cheese quality and

627   safety. Current Opinion in Biotechnology 20:142–148.

628   20.   Kastman EK, Kamelamela N, Norville JW, Cosetta CM, Dutton RJ, Wolfe BE. 2016. Biotic

629   interactions shape the ecological distributions of Staphylococcus species. mBio 7:e01157-16.

630   21.   Pham N-P, Landaud S, Lieben P, Bonnarme P, Monnet C. 2019. Transcription Profiling Reveals

631   Cooperative Metabolic Interactions in a Microbial Cheese-Ripening Community Composed of

632   Debaryomyces hansenii, Brevibacterium aurantiacum, and Hafnia alvei. Front Microbiol 10.

633   22.   Brown TL, Charity OJ, Adriaenssens EM. 2022. Ecological and functional roles of bacteriophages

634   in contrasting environments: marine, terrestrial and human gut. Current Opinion in

635   Microbiology 70:102229.

636   23.   Ledormand P, Desmasures N, Dalmasso M. 2020. Phage community involvement in fermented

637   beverages: an open door to technological advances? Critical Reviews in Food Science and

638   Nutrition 1–10.

639   24.   Paillet T, Dugat-Bony E. 2021. Bacteriophage ecology of fermented foods: anything new under

640   the sun? Current Opinion in Food Science 40:102–111.

641   25.   Garneau JE, Moineau S. 2011. Bacteriophages of lactic acid bacteria and their impact on milk

642   fermentations. Microb Cell Fact 10 Suppl 1:S20.

643   26.   Pujato S a., Quiberoni A, Mercanti D j. 2019. Bacteriophages on dairy foods. Journal of Applied

644   Microbiology 126:14–30.

645   27.   Marcó MB, Moineau S, Quiberoni A. 2012. Bacteriophages and dairy fermentations.

646   Bacteriophage 2:149–158.

647   28.   Muhammed MK, Kot W, Neve H, Mahony J, Castro-Mejía JL, Krych L, Hansen LH, Nielsen DS,

648   Sørensen SJ, Heller KJ, Sinderen D van, Vogensen FK. 2017. Metagenomic analysis of dairy

649   bacteriophages: extraction method and pilot study on whey samples derived from using

650   undefined and defined mesophilic starter cultures. Appl Environ Microbiol 83:e00888-17.

651   29.   Dugat-Bony E, Lossouarn J, De Paepe M, Sarthou A-S, Fedala Y, Petit M-A, Chaillou S. 2020. Viral

652   metagenomic analysis of the cheese surface: A comparative study of rapid procedures for

653   extracting viral particles. Food Microbiology 85:103278.

654   30.   Queiroz LL, Lacorte GA, Isidorio WR, Landgraf M, de Melo Franco BDG, Pinto UM, Hoffmann C.

655   2022. High Level of Interaction between Phages and Bacteria in an Artisanal Raw Milk Cheese

656   Microbial Community. mSystems 0:e00564-22.

657   31.   Cheng L, Marinelli LJ, Grosset N, Fitz-Gibbon ST, Bowman CA, Dang BQ, Russell DA, Jacobs-Sera

658   D, Shi B, Pellegrini M, Miller JF, Gautier M, Hatfull GF, Modlin RL. 2018. Complete genomic

659   sequences of Propionibacterium freudenreichii phages from Swiss cheese reveal greater

660   diversity than Cutibacterium (formerly Propionibacterium) acnes phages. BMC Microbiology

661   18:19.

662 32. de Melo AG, Rousseau GM, Tremblay DM, Labrie SJ, Moineau S. 2020. DNA tandem repeats

663   contribute to the genetic diversity of Brevibacterium aurantiacum phages. Environmental

664   Microbiology 22:3413–3428.

665 33. Gautier M, Rouault A, Sommer P, Briandet R. 1995. Occurrence of Propionibacterium

666   freudenreichii bacteriophages in swiss cheese. Appl Environ Microbiol 61:2572–2576.

667 34. Paillet T, Lossouarn J, Figueroa C, Midoux C, Rué O, Petit M-A, Dugat-Bony E. 2022. Virulent

668   Phages Isolated from a Smear-Ripened Cheese Are Also Detected in Reservoirs of the Cheese

669   Factory. 8. Viruses 14:1620.

670 35. Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, Kyrpides NC. 2021. CheckV assesses

671   the quality and completeness of metagenome-assembled viral genomes. 5. Nat Biotechnol

672   39:578–585.

673 36. Kieft K, Zhou Z, Anantharaman K. 2020. VIBRANT: automated recovery, annotation and curation

674   of microbial viruses, and evaluation of viral community function from genomic sequences.

675   Microbiome 8:90.

676 37. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

677   RNA-seq data with DESeq2. Genome Biology 15:550.

678 38. McMurdie PJ, Holmes S. 2014. Waste Not, Want Not: Why Rarefying Microbiome Data Is

679   Inadmissible. PLOS Computational Biology 10:e1003531.

680 39. Walsh AM, Macori G, Kilcawley KN, Cotter PD. 2020. Meta-analysis of cheese microbiomes

681   highlights contributions to multiple aspects of quality. Nat Food 1:500–510.

40.   de Melo AG, Lemay M-L, Moineau S. 2023. The impact of Brevibacterium aurantiacum virulent

phages on the production of smear surface-ripened cheeses. International Journal of Food

Microbiology 400:110252.

41.   De Pasquale I, Di Cagno R, Buchin S, De Angelis M, Gobbetti M. 2014. Microbial Ecology

Dynamics Reveal a Succession in the Core Microbiota Involved in the Ripening of Pasta Filata

Caciocavallo Pugliese Cheese. Applied and Environmental Microbiology 80:6243–6255.

42.   Flórez AB, Mayo B. 2006. Microbial diversity and succession during the manufacture and

ripening of traditional, Spanish, blue-veined Cabrales cheese, as determined by PCR-DGGE.

International Journal of Food Microbiology 110:165–171.

43.   Fuka MM, Wallisch S, Engel M, Welzl G, Havranek J, Schloter M. 2013. Dynamics of Bacterial

Communities during the Ripening Process of Different Croatian Cheese Types Derived from Raw

Ewe's Milk Cheeses. PLoS ONE 8:e80734.

44.   Larpin S, Mondoloni C, Goerges S, Vernoux J-P, Guéguen M, Desmasures N. 2006. *Geotrichum*

candidum dominates in yeast population dynamics in Livarot, a French red-smear cheese. FEMS

Yeast Research 6:1243–1253.

45.   Pangallo D, Šaková N, Koreňová J, Puškárová A, Kraková L, Valík L, Kuchta T. 2014. Microbial

diversity and dynamics during the production of May bryndza cheese. International Journal of

Food Microbiology 170:38–43.

46.   Ortiz Charneco G, De Waal PP, Van Rijswijck IMH, Van Peij NNME, Van Sinderen D, Mahony J.

2023. Bacteriophages in the Dairy Industry: A Problem Solved? Annu Rev Food Sci Technol

14:367–385.

47. Somerville V, Berthoud H, Schmidt RS, Bachmann H-P, Meng YH, Fuchsmann P, von Ah U, Engel

P. 2021. Functional strain redundancy and persistent phage infection in Swiss hard cheese

starter cultures. ISME J 1–12.

48. Erkus O, de Jager VC, Spus M, van Alen-Boerrigter IJ, van Rijswijck IM, Hazelwood L, Janssen

PW, van Hijum SA, Kleerebezem M, Smid EJ. 2013. Multifactorial diversity sustains microbial

community stability. The ISME journal 7:2126–2136.

49. Spus M, Wardhana YR, Wolkers-Rooijackers JCM, Abee T, Smid EJ. 2023. Lytic bacteriophages

affect the population dynamics of multi-strain microbial communities. Microbiome Res Rep 2.

50. Jolicoeur AP, Lemay M-L, Beaubien E, Bélanger J, Bergeron C, Bourque-Leblanc F, Doré L,

Dupuis M-È, Fleury A, Garneau JE, Labrie SJ, Labrie S, Lacasse G, Lamontagne-Drolet M, Lessard-

Hurtubise R, Martel B, Menasria R, Morin-Pelchat R, Pageau G, Samson JE, Rousseau GM,

Tremblay DM, Duquenne M, Lamoureux M, Moineau S. 2023. Longitudinal Study of Lactococcus

Phages in a Canadian Cheese Factory. Applied and Environmental Microbiology 0:e00421-23.

51. Madera C, Monjardín C, Suárez JE. 2004. Milk contamination and resistance to processing

conditions determine the fate of Lactococcus lactis bacteriophages in dairies. Applied and

Environmental Microbiology 70:7365–7371.

52. Wagner N, Brinks E, Samtlebe M, Hinrichs J, Atamer Z, Kot W, Franz CMAP, Neve H, Heller KJ.

2017. Whey powders are a rich source and excellent storage matrix for dairy bacteriophages.

International Journal of Food Microbiology 241:308–317.

53. Bokulich NA, Mills DA. 2013. Facility-Specific "House" Microbiome Drives Microbial Landscapes

of Artisan Cheesemaking Plants. Applied and Environmental Microbiology 79:5214–5223.

54.   Raymond-Fleury A, Lessard M-H, Chamberland J, Pouliot Y, Dugat-Bony E, Turgeon SL, St-Gelais

725   D, Labrie S. 2022. Analysis of Microbiota Persistence in Quebec's Terroir Cheese Using a

726   Metabarcoding Approach. 7. Microorganisms 10:1381.

727   55.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

728   data. Bioinformatics 30:2114–2120.

729   56.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,

730   Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA.

731   2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing.

732   J Comput Biol 19:455–477.

733   57.   Shah SA, Deng L, Thorsen J, Pedersen AG, Dion MB, Castro-Mejía JL, Silins R, Romme FO,

734   Sausset R, Jessen LE, Ndela EO, Hjelmsø M, Rasmussen MA, Redgwell TA, Leal Rodríguez C,

735   Vestergaard G, Zhang Y, Chawes B, Bønnelykke K, Sørensen SJ, Bisgaard H, Enault F, Stokholm J,

736   Moineau S, Petit M-A, Nielsen DS. 2023. Expanding known viral diversity in the healthy infant

737   gut. 5. Nat Microbiol 8:986–998.

738   58.   Kent WJ. 2002. BLAT—The BLAST-Like Alignment Tool. Genome Res 12:656–664.

739   59.   Guo J, Bolduc B, Zayed AA, Varsani A, Dominguez-Huerta G, Delmont TO, Pratama AA, Gazitúa

740   MC, Vik D, Sullivan MB, Roux S. 2021. VirSorter2: a multi-classifier, expert-guided approach to

741   detect diverse DNA and RNA viruses. Microbiome 9:37.

742   60.   Roux S, Camargo AP, Coutinho FH, Dabdoub SM, Dutilh BE, Nayfach S, Tritt A. 2022. iPHoP: an

743   integrated machine-learning framework to maximize host prediction for metagenome-

744   assembled virus genomes. preprint. Microbiology.

745   61.   Vasimuddin Md, Misra S, Li H, Aluru S. 2019. Efficient Architecture-Aware Acceleration of BWA-

746   MEM for Multicore Systems, p. 314–324. *In* 2019 IEEE International Parallel and Distributed

747   Processing Symposium (IPDPS).

748   62.   McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive analysis and

749   graphics of microbiome census data. PLOS ONE 8:e61217.

750   63.   Lin H, Peddada SD. 2020. Analysis of microbial compositions: a review of normalization and

751   differential abundance analysis. 1. npj Biofilms Microbiomes 6:1–13.

752   64.   Escudié F, Auer L, Bernard M, Mariadassou M, Cauquil L, Vidal K, Maman S, Hernandez-Raquet

753   G, Combes S, Pascal G. 2017. FROGS: Find, Rapidly, OTUs with Galaxy Solution. Bioinformatics

754   https://doi.org/10.1093/bioinformatics/btx791.

755   65.   Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M. 2014. Swarm: robust and fast clustering

756   method for amplicon-based studies. PeerJ 2:e593.

757   66.   Bokulich NA, Subramanian S, Faith JJ, Gevers D, Gordon JI, Knight R, Mills DA, Caporaso JG.

758   2013. Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing.

759   Nat Methods 10:57–59.

760   67.   Yoon S-H, Ha S-M, Kwon S, Lim J, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: a

761   taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. Int

762   J Syst Evol Microbiol 67:1613–1617.

Study of phage succession during ripening
(short-term)

Study of temporal variations
(long-term)

Wash 1 (W1)   Wash 2 (W2)   Wash 3 (W3)   Wash 4 (W4)   Wash 5 (W5)

Curd production

Day 1   Storage 4°C and transport   Day 28

2017

2019

2022

Viral metagenomics   16S metabarcoding   Plate counts

A SHANNON INDEX

Kruskal-Wallis test, Chi2 = 6.767 , p = 0.149

B BRAY-CURTIS

Permanova, R2 = 0.658 , p = 0.001 ***

Kinetic
- W1
- W2
- W3
- W4
- W5

PCOA2 [22.5%]

PCOA1 [57.4%]

C

I
II
III
IV
V

Host_genus
Group
Lifestyle
W1-A W1-B W1-C W2-A W2-B W2-C W3-A W3-B W3-C W4-A W4-B W4-C W5-A W5-B W5-C

log10
-1
-3
-5

Host_genus
- Lactococcus
- Streptococcus
- Leuconostoc
- Brevibacterium
- Glutamicibacter
- Pseudoalteromonas
- Vibrio
- Enterococcus
- No prediction

Group
- Lactococcus 936 group
- Lactococcus P335 group
- Lactococcus 949 group
- Streptococcus 987 group
- Leuconostoc Diderot
- Brevibacterium Rousseau
- Glutamicibacter Montesquieu
- No hit

Lifestyle
- Temperate
- Virulent
- No prediction

A Whole dataset (297 contigs)

B Filtered dataset (99 contigs)

C BRAY-CURTIS
Permanova, R2 = 0.814 , p = 0.002 **

D Bacterial community composition