Phylogenetic diversity of flowering plants declines across the growing season in Rocky

Mountain wildflower communities

Key words: co-flowering, network analysis, Rocky Mountain Biological Laboratory,

phylogenetic diversity, elevational gradient

Manuscript type: major article

Word count: 6764

Abstract

Phylogenetic diversity is an important axis of biodiversity associated with many

ecosystem functions, but its variation over time in temperate communities has rarely been

explored. Plants' flowering phenology is key for reproduction: it determines synchrony among

plants, mutualists, antagonists, and favorable environmental conditions. Individual species'

flowering phenologies combine to determine which species' flowers co-occur in time and space

to form co-flowering communities. We recorded plants' flowering phenology across elevations

in the Colorado Rocky Mountains and predicted: 1) flowering species will form subsets of co-

flowering species, 2) groups of co-flowering species will be more distantly related than the

broader community, 3) marginal early and late season conditions will constrain flowering to

species groups that are less phylogenetically diverse and 4) the entire community will have

higher phylogenetic diversity at higher elevations. We identified co-flowering groups using a

network analysis and quantified their phylogenetic diversity. Co-flowering communities had

higher phylogenetic diversity early in the season and at higher elevations, but became less

diverse by all metrics at the end of the season across elevations. Our results indicate that

phylogenetic diversity varies consistently across the season and that conditions in the late season

and at low elevations might constrain diversity.

Words: 197/200

Key words: co-flowering, network analysis, Rocky Mountain Biological Laboratory,

phylogenetic diversity, elevational gradient

Introduction

How biological diversity is distributed through space and time has been a central question in ecology and evolution for centuries (Linnaeus 1799; Darwin 1859; MacArthur 1969; Hillebrand 2004; Schluter and Pennell 2017). Most recently, the distribution of evolutionary (phylogenetic) diversity across contemporary communities has received a great deal of attention (Webb et al. 2002; Mayfield and Levine 2010; Molina-Venegas et al. 2021; Tietje et al. 2023). Higher phylogenetic diversity within a community indicates greater evolutionary divergence of community members, and it is thought that phylogenetic diversity may capture ecologically relevant variation among species that is missed when quantifying diversity by species richness or variation in one or a few traits (Cadotte et al. 2009; Owen et al. 2019; Huang et al. 2020; E-Vojtkó et al. 2023). Phylogenetic diversity is often observed to correlate poorly with variation in individual traits (CaraDonna and Inouye 2015; Gerhold et al. 2015; Maitner et al. 2022; E-Vojtkó et al. 2023), yet it does appear to be associated with community-level functions (Mazzochini et al. 2019; Molina-Venegas et al. 2021). For example, phylogenetic diversity correlates more positively with biomass than do other diversity metrics (Cadotte et al. 2008), and is associated with greater ecosystem stability and productivity (Cadotte et al. 2012; Cadotte 2013; Davies et al. 2016; Mazzochini et al. 2019) and increased resistance to species invasion (Galland et al. 2019). Extinction risk has also been observed to be phylogenetically clustered, indicating phylogenetic variation in the ecological vulnerability of species (Willis et al. 2008; Eiserhardt et al. 2015; González-Orozco et al. 2016). Here, we investigate patterns of phylogenetic diversity in plants that flower together in the same plant communities, and how their diversity changes across the growing season.

The timing of flowering (flowering phenology) is a key component of plant reproduction,

as it determines the synchrony (or lack thereof) between plants and their reproductive mutualists

and antagonists (Elzinga et al. 2007; Forrest and Miller-Rushing 2010; Munguía-Rosas et al.

2011; Freimuth et al. 2022), and with the environmental conditions favorable for reproduction

(Jentsch et al. 2009). At the community level, individual species' flowering phenologies combine

to determine which species' flowers co-occur in time and space (co-flower), shaping the floral

resources available to pollinators (Blaauw and Isaacs 2014), and affecting whether plant species

might compete with or facilitate one another to attract pollinators (Levin and Anderson 1970;

Brown and Kodric-Brown 1979; Moeller 2004; Moeller and Geber 2005a; Ghazoul 2006;

Kharouba et al. 2018). Importantly, nearly 90% of flowering plants rely on animal pollinators for

reproduction (Ollerton et al. 2011), but pollinator populations are declining worldwide

(Biesmeijer et al. 2006; Koh et al. 2016), disrupting plant-pollinator networks and driving

declines in plant populations (Burkle et al. 2013; Potts et al. 2016). Plants are also changing their

phenology in response to climate change and biodiversity loss (Menzel et al. 2006; Cleland et al.

2007; CaraDonna et al. 2014; Wolf et al. 2017), and such shifts in phenology are expected to

alter interspecific interactions (Kharouba et al. 2018; Pareja-Bonilla et al. 2023; Austin et al.

2024). An understanding of how the phylogenetic composition of co-flowering plant

communities changes across the growing season could provide a foundation for increasing our

understanding of the relationship between species composition and functioning of these

communities, and the potential consequences of environmental change and biodiversity loss for

flowering.

Of particular importance is when in the season phylogenetic diversity peaks and when it

is lowest, given the positive associations between phylogenetic diversity and ecosystem

productivity and stability (Cadotte et al. 2012; Cadotte 2013; Davies et al. 2016; Mazzochini et

al. 2019), and the potential sensitivity of low diversity communities to phylogenetically clustered

declines or extinctions (Willis et al. 2008; Eiserhardt et al. 2015; González-Orozco et al. 2016).

A central prediction of community phylogenetics is that 'environmental filtering' that favors

only the species adapted to a particular set of conditions will result in a lower diversity of species

establishing in a community, due to phylogenetic conservation of multi-trait strategies for the

necessary environmental adaptations (Rejmánek 1996; Webb 2000; Webb et al. 2002). Evidence

for this pattern has been found more often at broad spatial or environmental scales and

environmental extremes, though not universally (Ma et al. 2016; Park et al. 2020; Maitner et al.

2022). For flowering communities, this would predict that groups of species co-flowering at one

time of the season would be less diverse (more closely related) than the community as a whole,

particularly at environmental extremes that might occur early and late in the season at the

margins of suitable flowering conditions.

In contrast, closely-related species are predicted to compete more strongly at the local

scale, leading to communities of more distantly related species (higher phylogenetic diversity),

now known as the 'competition-relatedness hypothesis' (Darwin 1859; Ricciardi and Mottiar

2006). This hypothesis would predict that groups of species co-flowering and interacting would

be more diverse than expected by chance than the community as a whole. Considering both

hypotheses of environmental filtering and competition-relatedness, we might expect local-scale

co-flowering communities to be overall more diverse than expected, but to become more closely-

related and less diverse at the environmental extremes of the season, due to increased

environmental filtering. Nevertheless, studies of community phylogenetic assembly have found

mixed support for environmental filtering and competition-relatedness hypotheses (Daehler

2001; Proche\u015f et al. 2007; Thuiller et al. 2010; Jones et al. 2013; Ma et al. 2016; Cadotte et al.

2018; Park et al. 2020). A persistent challenge has been addressing how the niche and fitness

differences of potentially competing species relate to phylogeny, given that phylogeny is often

not aligned with specific traits (Mayfield and Levine 2010; Maitner et al. 2022). Indeed, for

flowering communities, floral traits are known to be highly variable phylogenetically (Vamosi

and Vamosi 2010; CaraDonna and Inouye 2015; O'Meara et al. 2016), and so any

phylogenetically structured environmental filtering or competitive interactions may be shaped by

other phylogenetically conserved non-floral traits of species (but see (Martín-Hernanz et al.

2023; Xiang et al. 2023).

Despite the importance of understanding the diversity of co-flowering communities and

testing predictions for their structure, little work has quantified the phylogenetic relationships of

co-flowering species. Studies on floral networks by Albor et al. (2020, 2022) found low

phylogenetic signal in the overlap of co-flowering species pairs but did not examine variation in

phylogenetic diversity of co-flowering communities. Studies by Wolowski et al. (2017) of

hummingbird-pollinated plants in the Brazilian Atlantic rainforest, and by Shrestha et al. (2019)

of floral color communities in Australian herbs, each found some evidence for non-random

phylogenetic structure (either less or more diverse than expected by chance), depending on the

environment and subset of the community considered. Notably, Wolowski and colleagues found

lower diversity and potential evidence of environmental filtering in lowland relative to montane

sites, but they did not quantify phylogenetic diversity over time during the season (Wolowski et

al. 2017). Shrestha and colleagues found variable temporal patterns in their data, depending on

the pollinator type of the floral community analyzed (Shrestha et al. 2019). To our knowledge,

no other studies have examined phylogenetic composition of flowering communities across a

season.

Within a flowering community at a particular site, groups of species that flower at the

same time within a season can be identified using the framework of a network analysis (Proulx et

al. 2005; Arceo-Gómez et al. 2018). A network analysis quantifies the connections ('edges')

between entities ('nodes') and identifies whether certain groups of nodes form clusters of strong

connections to one another (a 'module'), with weaker connections outside of the group. As

proposed by Arceo-Gómez and colleagues (2018), the duration and intensity of flowering

overlap can be used to define the strength of connections (edges) between pairs of flowering

species (nodes), and these pairwise connections can be used to form a network of flowering

interactions across the entire community. The extent to which the network has groups (modules)

of species with more interactions with one another than expected by chance can be quantified as

the network's modularity. Modules then represent subsets of species within communities that co-

flower more strongly with each other than with other species (Arceo-Gómez et al. 2018). Such

network analyses have been used to investigate trait similarity within co-flowering modules and

how plant-pollinator interactions may impact community assembly processes (Albor et al. 2020,

2022; Wei et al. 2021; Suárez-Mariño et al. 2022).

Here we use network analyses to identify the presence of co-flowering modules across

seasons and quantify the phylogenetic diversity within each module. To do this, we tracked

flowering phenology for plant communities at the Rocky Mountain Biological Laboratory

(RMBL) in the central Colorado Rocky Mountains. The flowering species in this region are

some of the best studied globally for flowering phenology (Dunne et al. 2003; Inouye 2008;

Forrest et al. 2010; Aldridge et al. 2011; CaraDonna et al. 2014; Powers et al. 2022; Rivest et al.

2023) and plant-pollinator interactions (Bosch and Waser 1999; Forrest and Thomson 2010;

Brosi and Briggs 2013; Waser and Price 2016; Gallagher and Campbell 2017; Ogilvie et al.

2017; Bain et al. 2022; Faust and Iler 2022; Schiffer et al. 2023). Different combinations of co-

flowering species within these communities have been observed to either facilitate or interfere

with pollination of one another (Forrest et al. 2010; Bain et al. 2022; Faust and Iler 2022;

Arrowsmith et al. 2023; Schiffer et al. 2023), though the phylogenetic context of these

interactions has not been explored.

Two previous studies have explored aspects of phylogenetic relationships in RMBL plant

communities, to our knowledge. CaraDonna and Inouye (2015) tested for phylogenetic signal in

multiple phenological traits of 60 species at RMBL, asking whether phenological traits tended to

be more similar for more closely related species. They found significant phylogenetic signal in

peak flowering and first flowering dates, but not in end date, flowering duration, or the

sensitivities of any phenological trait to environmental variation (CaraDonna and Inouye 2015).

Thus, the traits that will shape flowering overlap between these species include a mix of traits

that are likely to follow patterns of phylogenetic relationships and those that will not. In a study

of community phylogenetic diversity across elevations in the RMBL area, Bryant and colleagues

(2008) found that plant communities tended to be increasingly overdispersed (more diverse than

expected at random) as elevation increased up to observations at 3800 m above sea level, with a

peak in absolute phylogenetic diversity circa 3000 m elevation, indicating that elevational

gradients are likely important for shaping plant community diversity in this system.

Our study of the phenology of phylogenetic diversity in co-flowering communities

includes wildflower communities in three meadows that range from a montane site at 2815 m

above sea level to a subalpine site just below treeline at 3380 m. We quantify the distribution of

flowering abundance of species in these communities and we use a network analysis to test

whether flowering phenology is clustered into groups (modules) of co-flowering species across

the season. We then quantify phylogenetic diversity across the season, and test whether it differs

across modules of species flowering together. We test the general prediction that phylogenetic

diversity will be higher than expected at these local scales and within co-flowering modules

(overdispersed) but relatively lower (more underdispersed) early and late in the season when

environmental conditions are most marginal. Finally, we ask whether overdispersion increases

with elevation in co-flowering communities, as has been previously observed for co-occurring

plants along the same gradient (Bryant et al. 2008).

Materials & Methods

<u>Study Sites</u>

We collected data at three sites in Washington Gulch near the Rocky Mountain

Biological Laboratory (RMBL, Gothic, Colorado, USA) from June to August in 2021 and 2022.

RMBL is in the East River valley of the West Elk mountains, approximately 10 kilometers from

Crested Butte, Colorado. The relatively short growing season lasts 3-5 months, and snowmelt

date is variable, but has been arriving earlier on average with climate change (Aldridge et al.

2011; CaraDonna et al. 2014). Study sites were located at 2815 m (38°53'50"N, 106°58'43"W),

3165 m (38°57'38"N, 107°01'53"W), and 3380 m (38°58'10"N, 107°01'53"W) in elevation.

These sites are part of a larger, long-term study of plant functional traits over an elevational

gradient at RMBL established in 2003 (Bryant et al. 2008; Sloat et al. 2015). All sites are

approximately 50 m^2 and have similar slopes and aspects (Sloat et al. 2015). Using quadrats, we

defined five 1.2 m x 1.2 m plots spaced 5 m apart from each other along a subtransect running

downslope within each site. We did not control for density or other factors when placing the

plots so we could capture an unbiased sample of the sites. We then marked the plots with flags at

the corners and strings around the edges. We removed these markings at the end of each summer,

so exact plot locations shifted slightly each year. The lowest and middle sites are in the montane

vegetation zone, while the highest is subalpine (Ackerfield 2015). Across our sites, both soil pH

and average July soil temperature decrease with elevation (Bryant et al. 2008). We identified

forbs in the plots using the *Flora of Colorado* (Ackerfield 2015) and reference specimens at the

Rocky Mountain Biological Laboratory Herbarium (herbarium code RMBL).

Field Data Collection

Within the plots, we recorded all newly open flowers or inflorescences once per week

from early June to mid-August for ten weeks of observation over each of the two summers. We

counted flowers for all forbs in all plots, which resulted in counts for 56 species (see also

Results). We counted flowers as 'flowering units', which we defined as either a single flower

(most species), a capitula (Asteraceae), or an umbel (Apiaceae). We considered flowers open

once the petals were separated and able to be counted. Each week, we tagged all new flowering

units with a unique color of yarn and recorded the number of flowering units per species. We did

not have predefined sample sizes and included all flowering units, which resulted in 3,660

flowering units in 2021, 1,973 in 2022, and a total of 5,633 flowering units across all sites and

both years for our sample size (see also Results; Fig. 1). We only counted flowering units once

as they opened, so counts each week were not cumulative. Marked flowering units were very

rarely still open by the time of data collection for the next week, so recounting of previously-

marked flowering units was not required to capture the flowering community each week (first

author initials, pers. obs.). We never observed wilted or dead flowering units that we hadn't

previously tagged. All flowering units across all individuals of a species in all plots were

combined to produce a single count of flowering units per species, per site, each week. Counts

across weeks produced a distribution of flowering abundance across the 10 weeks of observation.

This type of weekly abundance data has previously been shown to capture variation in co-

flowering for co-flowering network analyses in similar-length growing seasons (Arceo-Gómez et

al. 2018).

For species that fruited during the study period, we also quantified successful fruit and

seed development at the end of each species' flowering period. These data are not used for

further analyses in this study, and data and associated Methods are available through the

Environmental Data Initiative portal under package ID edi.1478.1.

<u>Identifying Modules in Co-flowering Networks</u>

To test whether co-flowering communities were significantly modular, we used a

network analysis with flowering species as the nodes and flowering overlap as the connections

between them (edges). To quantify flowering overlap between each species pair in a given site

and year, we calculated Schoener's Index of niche overlap (Schoener 1970). We used the number

of newly open flowers per week for each species to quantify its flowering phenology distribution

and calculated Schoener's Index between the distributions of a species pair as:

$SI = 1 - 1/2\sum |Pik - Pjk|$

where Pik and Pjk are the proportion of species i and j flowering in week k (Forrest et al. 2010).

Values range from 0 to 1, and fall at or near 0 when there is no flowering overlap and close to or

at 1 for near or complete overlap in flowering distributions. We calculated Schoener's Index in

the R package spaa with the command niche.overlap() and method "schoener" (Zhang 2016).

We constructed co-flowering networks for each site for 2021 and 2022 (six total) with

Schoener's Index values as edge weights (Arceo-Gómez et al. 2018; Albor et al. 2020, 2022). An

edge between two species will have more weight if they have a greater flowering overlap (i.e. a

larger Schoener's Index value). We constructed networks using the igraph R package (Csardi and

Nepusz 2006). We calculated modularity using (Blondel et al. 2008)'s algorithm via the

cluster.louvain() command in igraph (Table 1). Within the cluster.louvain() command, we

specified weighted edges (according to their Schoener's Index) and a resolution of 1.0.

Resolution tells the algorithm whether to group the nodes into many, small modules or few,

larger modules, and 1 is a standard value to not bias the algorithm towards module size

(Newman and Girvan 2004; Csardi and Nepusz 2006).

The modularity algorithm has two outputs: modularity values (Q), which range from -1 to

1, and which nodes (species) belong in which module. A modularity of Q = 1 represents a

perfectly modular network in which modules have no connections (edges) outside their module.

A Q = 0 indicates that the structure is no different than random, and Q = -1 results when there is

no clustering in the network. To calculate the statistical significance of modularity (Q), we

generated 1000 null networks for each of the six site and year combinations using the

nullmodel() command and method "r2d" in the bipartite and vegan R packages (Dormann and

Strauss 2014). We then calculated modularity (Q) of the null networks and a z-score for the

observed modularity relative to the null distribution (Table 1). The z-score reflects how many

standard deviations the actual network's modularity is away from the mean modularity of all

1000 null networks, and z-scores above 2 are considered significant (Dormann and Strauss

2014).

We also used Gephi (Bastian et al. 2009) to calculate betweenness centrality, which is a

network metric that estimates how many times a species connects otherwise unconnected species

(ranging from zero to positive numbers of these connections), and can be used to identify

keystone species with relatively high values that contribute uniquely to community overlap in

flowering (Martín González et al. 2010; Arceo-Gómez et al. 2018). We constructed network

visualizations in Gephi (Bastian et al. 2009).

<u>Phylogenetic Diversity of Co-Flowering Modules</u>

To quantify and test predictions about phylogenetic diversity of co-flowering species

across the season, we calculated phylogenetic diversity metrics for each site, and then for the

communities of species flowering in each week and in each co-flowering module identified by

the network analyses. We used the 'ALLMB.tre' angiosperm phylogeny from Smith and Brown

(2018) to calculate all phylogenetic metrics. This phylogeny includes all genera in our dataset

and 49 of 56 species observed in our plots. If a species was missing from the phylogeny but was

the only member of its genus in the plots, we replaced it with a closely related congener in the

Smith and Brown phylogeny (Supplementary material Table S1). Only the genus *Senecio* had

multiple species in the plots missing from the phylogeny. We replaced *Senecio integerrimus*

Nutt. with sister species *S. triangularis* Hook. (Pelser et al. 2007). We removed *Senecio*

crassulus A. Gray from all phylogenetic analyses because it has no available genetic or

phylogenetic information.

To quantify phylogenetic diversity, we calculated three metrics: Faith's Phylogenetic

Diversity (Faith 1992), Mean Phylogenetic Distance (MPD), and Mean Nearest Taxon Distance

(MNTD) in the R package geiger (Pennell et al. 2014). Faith's Phylogenetic Diversity is the total

evolutionary distance (branch length) across a phylogenetic tree describing the relationships

among the taxa in a dataset. This metric will increase if relationships are more distant across the

tree (species are more evolutionarily divergent). Mean Phylogenetic Distance is the mean

distance between pairs of species, and so it also increases as evolutionary relationships are more

distant across the tree, but will be less sensitive to rare cases of very distantly-related species that

would otherwise elevate Faith's Phylogenetic Diversity. Mean Nearest Taxon Distance is the

mean of the distance to the closest relative of each taxon, and so it reflects the typical distance to

any species' closest relative in the dataset and increases as closest relatives are more distant.

Therefore, MNTD is more reflective of the tendency of species to have closer relatives than it is

of the overall evolutionary diversity across the entire tree (community of species) that is captured

by the former metrics (for further discussion, see review by (Tucker et al. 2017)).

To test whether a plot had higher phylogenetic diversity than expected (overdispersion)

or less than expected (underdispersion), we calculated standard effect sizes (SES) for all three

metrics in the picante R package (Kembel et al. 2010) using the ses.metric() function. Positive

SES values indicate that the community in question is more phylogenetically diverse than a

random draw of the same number of species from the regional species pool (overdispersion), and

a *P* value above 0.975 indicates statistical significance of the SES. Negative SES values indicate

that the species group in question is less phylogenetically diverse than a random draw of the

same number of species from the regional species pool (underdispersion), and a *P* value below

0.025 indicates statistical significance of the SES. *P* values are significant if the SES absolute

value is greater than 1.96. We used each metric's "sample.pool" null model to draw a random

sample of the same number of species as the focal community from the pool of all species

observed across all RMBL sites with 5000 iterations.

We analyzed all data in R version 4.2.3 (R Core Team 2023).

Results

<u>Flowering Observations</u>

We recorded data for 56 species and 5,633 flowering units across all three sites and years

(Fig. 1). We counted 3,660 flowering units in 2021 and 1,973 in 2022. Across both years, the

lowest site had 33 species and 1,994 flowering units, the middle site had 29 species and 2,108

flowering units, and the highest site had 26 species and 1,531 flowering units. Five species

occurred at all three sites: *Potentilla gracilis* Douglas ex Hook., *Viola nuttallii* Pursh,

Ranunculus inamoenus Greene, *Mertensia brevistyla* S. Watson, and *Erigeron speciosus* DC

(Fig. 1).

Our dataset of pairwise observations of flowering overlap within sites and years yielded a

total of 1,262 pairwise observations, with site-year combinations ranging from a low of 136 pairs

at the High site in 2022 to a high of 300 pairs at the Low site in 2022 (Supplemental File

Schoeners_all.csv). Over half (57.2%) of the 1,262 species pair observations across sites and

years had low or no overlap in their flowering phenology (Schoener's Index at or near 0; Figs. S1

& S2). Over three quarters (76.4%) of these species pairs fell below a Schoener's Index of 0.25

(lower quarter of the index, which ranges from 0 to 1). A small subset of species pairs had high

flowering overlap: 58 of the 1,262 pairs (<5%) had Schoener's Index values in the upper quartile

of the index (at or above 0.75).

<u>Co-Flowering Network Modularity</u>

We constructed networks for flowering communities at each site in each year. Network

size ranged from 19 to 26 nodes (species) per network and from 153 to 325 edges (Schoener's

Index of overlapping flowering between species) each (Table 1; Fig. 2). All sites in both years

were significantly modular ($Q > 0.35$ for all; Z-scores > 2 for all; Table 1), indicating that certain

groups of species overlapped in their flowering more than with the other species outside of their

module (Fig. 2). The High site had the lowest modularity (2021: $Q= 0.42$; 2022: $Q=0.36$) and the

Low site had higher modularity (2021: $Q=0.44$; 2022: $Q=0.56$) in both years (Table 1),

indicating tighter clustering of the modules (weaker overlap with other modules) in the Low site.

All three sites had three modules each in 2021 (Fig. 2). In 2022, the High and Middle sites again

had three modules, while the Low site had four (Fig. 2). The duration of flowering of species

within each module varied, but ranged from 2-4 weeks, depending on the site, year, and time of

the summer (Fig. 1).

For all sites and years, modules corresponded with flowering at the early, mid, and late

growing season, and approximate dates of the modules were relatively consistent for sites across

both years (Fig. 1). A species' module depended on when the largest proportion of its flowers

were open. For example, if 90% of a species' flowers were open in weeks 1 and 2, but 10% of its

flowers opened in weeks 4 and 5, the modularity algorithm would place it in the early season

module. Thus, it is possible for species flowering in the same week to be in different modules,

depending on the other weeks when they had open flowers (Fig. 1). For the low site with an

additional module in 2022, there were two middle-season modules spanning the middle of the

growing season in July, but one had more overlapping flowering with the early season module

and the other was less connected to modules before or after it (Figs. 1 & 2).

367   Betweenness centrality values indicated that a small set of species had large numbers of

connections to otherwise unconnected species. The highest value of this metric was 197.03 for

Viola nuttallii at the low elevation site in 2022 (Table S2). The average betweenness centrality

value across our dataset was 8.6. In 2021, the average betweenness centrality was 5.6, and

highest values were >30, seen for *Achillea millefolium, Dasiphora fruticosa, Heliomeris*

multiflora and *Noccaea fendleri*. In 2022, the average betweenness centrality was 11.44, and

highest values were >50, seen for *Tragopogon dubius, Artemisia tridentata, Lupinus bakeri,*

Erigeron speciosus, Ericameria nauseosa and *Viola nuttallii.*

<u>Phylogenetic Diversity of Co-Flowering Modules</u>

377   The 56 species in our plots spanned 24 families: 12 at the lowest site, 14 at the middle,

and 16 at the highest. Five families (Asteraceae, Ranunculaceae, Rosaceae, Violaceae,

Boraginaceae) occurred at all three sites (Fig. 3). At the site level, across the entire season, the

Low site had the highest PD and lowest MPD in both years (Table 1). This indicates that the

Low site tended to have lower diversity (per MPD) but had some divergent taxa elevating PD

(Table 1). The only significant deviation from expected occurred for the High site in 2021, when

it had significantly higher diversity than expected (overdispersion; Table 1). MNTD values were

also the highest at the High site in both years, consistent with higher divergence between each

flowering species and its closest relatives at this site (Table 1).

386   Across weeks, phylogenetic diversity metrics generally indicated weak overdispersion

(higher diversity than expected, but not significantly) before declining to underdispersion (lower

diversity than expected) by the end of the season (Fig. 4). In 2021, underdispersion was

significant for all metrics by Week 10 for both Middle and High sites, and for MPD at the Low

site (Fig. 4). In 2022, underdispersion was weaker for the Middle and High sites in the later

weeks, but highly significant for the Low site for all three metrics for the last three weeks of the

season (Fig. 4).

For co-flowering modules of species flowering together, early-season modules were more

overdispersed than late-season modules for all sites, years, and phylogenetic metrics (Fig. 5).

Notably, all three metrics indicated significant underdispersion (SES < -1.96) for the late season

module in 2021, and MPD was consistently lowest and strongly significant with an SES less than

-4. This indicates that species co-flowering in the late season module are consistently more

closely related to one another. In 2022, as for the weekly data, the tendency for overdispersion in

the late season was present but weaker for the Middle and High sites, but continued to be highly

significant for the Low site.

Discussion

Despite the importance of flowering phenology for plants' reproductive success (Elzinga

et al. 2007; Forrest and Miller-Rushing 2010) and persistence during environmental change

(Cleland et al. 2007; Anderson 2016), the distribution of phylogenetic diversity across flowering

seasons has been largely undescribed. Here, we find that Rocky Mountain wildflower

communities are structured into modules of species that flower together, and that phylogenetic

diversity varies among modules. Co-flowering modules tend towards phylogenetically

overdispersion early in the season and at higher elevation, but become strongly underdispersed

and depauperate in diversity at the end of the flowering season at all elevations.

Co-flowering was significantly modular at all three sites and in both years. The total

richness of flowering taxa was approximately 30 species at each site (ranging from 33 at low

elevation to 26 at high elevation), but for nearly all site-year combinations this diversity was

divided into three modules of co-flowering species across the season. At the low elevation site in

2022, co-flowering species were further divided into an additional mid-season module. These

seasonal modules ranged from 15% to 55% of all species that were observed to flower across the

season at a site. Co-flowering communities were consistent across years: only 15 species

switched modules from 2021 to 2022 across all sites, despite 2021 having nearly twice the

abundance of flowers, with earlier snowmelt and less rain than 2022 (Prather et al. 2023). Thus,

communities of co-flowering species contained a fraction of the total flowering diversity at each

site, and co-flowered as largely the same communities year to year.

We predicted that phylogenetic diversity would be higher than expected within co-

flowering modules (overdispersed) but relatively lower (more underdispersed) early and late in

the season when environmental conditions are most marginal. We did not find significant

overdispersion within any module in either year, and instead we found that modules tended

towards overdispersion only earlier in the season (though not significantly). Phylogenetic

diversity did decline towards the end of the growing season by all phylogenetic metrics, resulting

in significant underdispersion by the end of the season. This seasonal pattern was consistent

whether species were grouped by flowering week or by module, with declines at the end of the

season beginning at roughly the same time as the transition to the flowering of the late season

module. This variation in phylogenetic composition across co-flowering modules was associated

with clear differences in the representation of regional taxonomic diversity. In particular, species

in the Asteraceae and, to a lesser extent, the Rosaceae dominated the modules at the end of the

flowering season. The prevalence of these taxa at the end of the season may be familiar to field

biologists (first author initials, pers. obs), but to our knowledge, lower phylogenetic diversity of

later flowering modules has not been previously reported. One previous study at RMBL found

phylogenetic signal (clustering) in first and peak flowering dates, but not in last flowering date or

flowering duration of species (CaraDonna and Inouye 2015). Those previously reported patterns

are consistent with our finding that the late flowering community is phylogenetically clustered

into relatively fewer clades, and that it occurs as a distinct module of species that begin their

flowering late in the season.

Our observation of lower diversity at the end of the season might reflect more marginal

conditions that are suitable for only a subset of species. Soil organic carbon, soil respiration, and

net ecosystem productivity all decline throughout the season in this region (Moyes and Bowling

2013; Sloat et al. 2015). Declines in each of these variables have been associated with decreasing

plant diversity in other systems (Tilman et al. 1997; Johnson et al. 2008; Lehmann et al. 2020).

Air temperature is also known to increase from June to August at this site, and has been shown to

impact flowering phenology, soil temperature, and pollinator behavior (Harte et al. 1995;

Aldridge et al. 2011; Ogilvie et al. 2017; Cordes et al. 2020). In contrast, while low soil moisture

has been observed to cause declines in species richness and phylogenetic diversity in other

systems (Deng et al. 2016; Wagg et al. 2017; Serafini et al. 2019; Miao et al. 2022), it is unlikely

to be a factor here. Soil moisture in our system is affected by snowmelt timing and precipitation

from summer thunderstorms, and has significant interannual variability (Pederson et al. 2011;

Waser and Price 2016; Campbell et al. 2019; Mooney et al. 2021), which would not explain our

consistent results.

Elevational differences could also be informative regarding the relationship between

abiotic factors and phylogenetic diversity. Our high-elevation site as a whole was more

overdispersed across both years by two of three diversity metrics (MPD and PD), and

consistently more overdispersed than our low-elevation site. The low elevation site deviated little

from null expectations and varied from weakly over- to underdispersed across metrics and years.

A previous study by Bryant et al. (2008) across this same elevational gradient also found that

phylogenetic overdispersion of plants increased with elevation, and that this relationship was

predicted by lower soil temperature, lower pH, and higher total nitrogen as elevation increased.

Similarly, Wolowski and colleagues found lower diversity and potential evidence of

environmental filtering in lowland relative to montane sites for hummingbird-pollinated plants in

the Brazilian Atlantic rainforest (Wolowski et al. 2017). These findings and ours suggest a

pattern of lower phylogenetic diversity (underdispersion) of the flowering communities that

occur in the warmer conditions at lower elevations and at the end of the season, and direct tests

of this relationship would be valuable.

Studies in other systems have supported the opposite of our finding: underdispersion at

high elevations, associated with abiotic filtering, wherein only a subset of evolutionary lineages

share adaptations to the high elevation environment (Webb et al. 2002; Graham and Fine 2008;

Dehling et al. 2014). These contrasting results highlight the uncertainty around which abiotic

conditions are likely to impose phylogenetic filtering in different systems. We find evidence of

filtering of flowering species only at the end of the flowering season, and thus the cooler/wetter

conditions that occur early in the season and at higher elevations might be more favorable (less

of a filter) for the species flowering across our gradient. It is noteworthy that the overdispersion

at our highest site was weaker in 2022, when this region experienced a later snowmelt and wetter

and cooler conditions (Prather et al. 2023), which may have imposed a stronger phylogenetic

filter on our highest elevation site.

481   In addition to the effects of abiotic conditions, biotic interactions among co-flowering

plants and their pollinators are likely to be shaping the diversity of flowering species across the

season. Co-flowering has been studied as a mechanism of both and competition among plants,

often specifically in the context of pollination (Sargent and Ackerly 2008; Carvalheiro et al.

2014; Briscoe Runquist et al. 2016). Pollinators can structure plant communities and co-

flowering networks (Sargent and Ackerly 2008; Wei et al. 2021), and may impact co-flowering

modules directly or indirectly through their interactions with abiotic conditions (Gallagher and

Campbell 2017). Previous work at RMBL has found that pollen and pollinator limitation are

often species-specific and/or context-dependent (Forrest and Thomson 2010; Gezon et al. 2016;

Schiffer et al. 2023), which suggests that the mechanisms shaping co-flowering community

assembly will depend on the particular species and context.

492   On the one hand, co-flowering is a potential mechanism of facilitation by which plants

might overcome pollinator limitation and attract more pollinators to a multi-species patch

(Ghazoul 2006). Indeed, some evidence suggests that co-flowering can increase pollinator

visitation rates and the fitness of participating species by increasing the rewards for pollinators

(Brown and Kodric-Brown 1979; Moeller 2004; Ghazoul 2006; Ogilvie and Thomson 2016;

Bain et al. 2022). For example, in a recent study at RMBL, the experimental removal of the

dominant plant species, *Helianthella quinquenervis,* immediately lowered pollinator visitation

rates and network robustness for remaining species (Bain et al. 2022). In our network analysis,

H. quinquenervis (Asteraceae) was observed as a middle or late season module species at the

middle elevation site. It had betweenness centrality values that were lower than the average

betweenness centrality of other species in both years, indicating low overlap with species outside

of its co-flowering module and the potential for close interactions within its module, which

would be dominated by related species in the Asteraceae in the late season. Similarly positive co-

flowering interactions among related species have been seen in other systems. *Clarkia* congeners

on the west coast of North America share pollinators and experience increased fitness and bee

visitation through overlapping flowering (Moeller 2004; Moeller and Geber 2005*b*). Community

members with similar floral shapes and sizes increased the seedset of *Delphinium barbeyi* at

RMBL (Arrowsmith et al. 2023). Notably, facilitative interactions have been predicted and

observed to be more common among distantly related species, which could result in

overdispersion (Valiente-Banuet and Verdú 2007; Cavender-Bares et al. 2009), but the evidence

above for pollinator facilitation among related species would predict that co-flowering modules

would be more underdispersed.

On the other hand, sharing pollinators in time and space can lead to lower fitness through

increased resource competition among plants (due to niche overlap) and/or reproductive

interference (Levin and Anderson 1970; Johnson et al. 2022). These negative interactions could

be stronger for closely related species if niches are phylogenetically conserved, though that need

not be the case (Mayfield and Levine 2010). For example, work in the North American Great

Lakes region showed that invasive *Lythrum salicaria* reduced bee visits to a native congener

when they co-flowered (Brown et al. 2002). In a recent study at RMBL, Faust and Iler (2022)

found that co-flowering decreased the seedset of *Linum lewisii*, but only when the plants were

relatively less drought-stressed. Their results suggest that abiotic factors can mediate biotic co-

flowering interactions, and the interactions may become more facilitative as environmental

stresses increase (Faust and Iler 2022). That multiple interacting abiotic and biotic mechanisms

might be shaping these co-flowering communities is suggested by the significant modularity in

their networks. Arceo-Gomez et al. (2018) suggested that modularity in a network indicates

multiple mechanisms affecting community assembly, because random or single mechanism-

driven assembly would not result in clustering of groups of interacting species in time. Thus,

future work should ideally combine tests of multiple abiotic and biotic interactions across

phylogenetic divergence of community members to understand their assembly.

Finally, we note that flowering timing in these communities is already being affected by

climate change. Globally, plants are flowering earlier with the advancing onset of spring (Menzel

et al. 2006; Cleland et al. 2007; CaraDonna et al. 2014; Wolf et al. 2017), and this trend has been

documented extensively at RMBL (Forrest et al. 2010; CaraDonna et al. 2014; Powers et al.

2022; Rivest et al. 2023). The patterns of co-flowering that we observe in this study are likely to

be changing over time, and may not be optimized for current abiotic and biotic conditions. In

particular, if climate change causes warm late season conditions to become more common earlier

in the season (Inouye 2008; Pederson et al. 2011), then we might expect that the highest diversity

early season modules are experiencing the most adverse effects. Studies of co-flowering network

change over time would reveal whether co-flowering module communities shift their

phenologies together, whether species diversity tends to be gained or lost in the process, and the

potential roles of phylogenetic diversity in shaping response to environmental change.

In summary, we find that co-flowering communities are significantly modular, containing

a subset of flowering species diversity at a site, and forming phenologically distinct communities

that are largely consistent across years. We also find that late season modules have lower

phylogenetic diversity according to all metrics (PD, MPD & MNTD) than early season modules,

and that this decline in diversity occurred at all three elevations. These patterns are consistent

with environmental filtering of phylogenetic diversity in the late season, but more work is

needed to understand the abiotic and biotic conditions that might impose filtering on flowering of

species at these sites. Across sites, our high elevation site contained more phylogenetic diversity

than the low elevation site in both years, consistent with previous observations of overdispersion

of the entire plant community at higher elevations, but contrasting with the prediction that high

elevations should impose intense filtering and contain lower phylogenetic diversity. Our results

clarify how phylogenetic diversity is distributed across sets of co-flowering species and across

elevations in a subalpine plant community, but also highlight open questions about the

mechanisms impacting the composition of co-flowering groups, and the evolutionary history of

traits that allow for early or late season flowering.

Figures

Fig. 1: Density distributions for the number of flowering units (flowers or inflorescences) opened each

week, for all species flowering at each site. Solid lines represent 2021 data and dashed lines represent

2022 data. Week 1 start dates were June 8, 2021 and June 6, 2022, and Week 10 start dates were August

9, 2021 and August 8, 2022. Lines are color coded by flowering module assigned by the network analysis

(magenta for Early, orange for Mid, turquoise for a Mid2 module present only in the 2022 Low elevation

site, and green for Late).

Fig. 2: Network analyses of flowering time for species at each site and year, with nodes (species) color-coded according to the co-flowering module assigned by the analysis. Modules shown in purple flower early in the summer, followed by orange modules in the middle of the summer and green modules late in the summer. Node size represents betweenness centrality from the network analysis, with larger nodes showing species that co-flower with more species. Lines connecting species show edges weighted according to their Schoener's Index of flowering overlap, where thicker lines have higher values (more flowering overlap) and thinner lines have lower values (less flowering overlap).

589   Fig. 3: Histograms of the number of flowering units (flowers or inflorescences) color coded by family.

590   Shown are panels for the Low elevation site (a, d), the Middle elevation site (b, e) and the High elevation

591   site (c, f) for 2021 (a, b, c) and 2022 (d, e, f) flowering seasons. Each season is partitioned into co-

592   flowering modules from the network analysis (Early, Mid, and Late, with an additional mid-season

593   module in the 2022 Low elevation site).

Fig. 4: Standard effect sizes for metrics of phylogenetic diversity of the flowering community by week,

for all sites and years. Metrics include mean nearest taxon distance (pink), mean phylogenetic distance

(green), and Faith's phylogenetic diversity (blue). Shown are panels for the Low elevation site (a, d), the

Middle elevation site (b, e), and the High elevation site (c, f) for 2021 (a-c) and 2022 (d-f) flowering

seasons. SES values of 1.96 or greater indicate significant overdispersion (greater diversity than

expected), while values of -1.96 and below indicate significant underdispersion (lower diversity than

expected).

Fig. 5: Standard effect sizes for metrics of phylogenetic diversity of species in co-flowering modules

identified in the network analyses, for all sites and years. Metrics include mean nearest taxon distance

(pink), mean phylogenetic distance (green), and phylogenetic diversity (blue). Shown are panels for the

Low elevation site (a, d), the Middle elevation site (b, e), and the High elevation site (c, f) for 2021 (a, b,

c) and 2022 (d, e, f) flowering seasons. Within sites and years, panels show phylogenetic diversity for

each co-flowering module, including those flowering in early, mid, and late season. SES values of 1.96 or

greater indicate significant overdispersion (greater diversity than expected), while values of -1.96 and

below indicate significant underdispersion (lower diversity than expected).

Table 1: Metrics from the network and phylogenetic analyses for flowering communities at each site

(High, Medium, Low) and year (2021, 2022). High site is at 3380 m, Medium is at 3165 m, and Low is at

2815 m. From the network analyses of co-flowering, shown are Modularity (Q scores), Z-score for

modularity significance, and the number of Edges (species pairs) in each network. From analyses of

phylogenetic diversity, shown are diversity metrics (Value), standard effect sizes (SES) for those values,

and *P*-values for significance (for each phylogenetic diversity metric, including Faith's Phylogenetic

Diversity (PD), Mean Phylogenetic Diversity (MPD), and Mean Nearest Taxon Distance (MNTD).

Significant P-values (those over above 0.975 or below 0.025) are indicated with an asterisk.

	Co-Flowering Networks			Faith's PD			MPD			MNTD		
Site	Modularity (Q)	Z-score	Edges	Value	SES	*P*-value	Value	SES	*P*-value	Value	SES	*P*-value
2021 Flowering Season												
High	0.42	19.75	171	1637.1	1.45	0.93	242.4	1.86	0.99*	131.8	0.37	0.64
Medium	0.48	28.22	231	1503.5	-0.80	0.21	221.8	0.15	0.52	97.5	-1.40	0.08
Low	0.44	24.27	231	1720.7	0.33	0.63	220.3	0.03	0.48	123.6	0.42	0.65
2022 Flowering Season												
High	0.36	14.56	153	1428.8	0.27	0.60	228.4	0.32	0.58	126.6	0.02	0.50
Medium	0.43	26.86	276	1677.2	-0.09	0.47	232.6	0.94	0.83	106.2	-0.59	0.28
Low	0.56	43.12	325	1841.5	-0.10	0.46	216.9	-1.16	0.13	114.4	0.25	0.59

References

Ackerfield, J. 2015. Flora of Colorado. Botanical Research Institute of Texas.

Albor, C., G. Arceo-Gómez, and V. Parra-Tabla. 2020. Integrating floral trait and flowering time

distribution patterns help reveal a more dynamic nature of co-flowering community assembly

processes. The Journal of ecology 108:2221–2231.

Albor, C., T.-L. Ashman, A. Stanley, C. Martel, and G. Arceo-Gómez. 2022. Flower colour and flowering

phenology mediate plant–pollinator interaction assembly in a diverse co-flowering community.

Functional ecology n/a.

Aldridge, G., D. W. Inouye, J. R. K. Forrest, W. A. Barr, and A. J. Miller-Rushing. 2011. Emergence of a

mid-season period of low floral resources in a montane meadow ecosystem associated with climate

change. The Journal of ecology 99:905–913.

Anderson, J. T. 2016. Plant fitness in a rapidly changing world. The New phytologist 210:81–87.

Arceo-Gómez, G., R. L. Kaczorowski, and T.-L. Ashman. 2018. A Network Approach to Understanding

Patterns of Coflowering in Diverse Communities. International journal of plant sciences 179:569–

582.

Arrowsmith, K. C., V. A. Reynolds, H. M. Briggs, and B. J. Brosi. 2023. Community context mediates

effects of pollinator loss on seed production. Ecosphere 14:e4569.

Austin, M. W., A. B. Smith, K. M. Olsen, P. C. Hoch, K. N. Krakos, S. P. Schmocker, and N. E. Miller-

Struttmann. 2024. Climate change increases flowering duration, driving phenological reassembly and

elevated co-flowering richness. The New phytologist.

Bain, J. A., R. G. Dickson, A. M. Gruver, and P. J. CaraDonna. 2022. Removing flowers of a generalist

plant changes pollinator visitation, composition, and interaction network structure. Ecosphere

13:e4154.

Bastian, M., S. Heymann, and M. Jacomy. 2009. Gephi: An Open Source Software for Exploring and

Manipulating Networks. Proceedings of the International AAAI Conference on Web and Social

Media 3:361–362.

Biesmeijer, J. C., S. P. M. Roberts, M. Reemer, R. Ohlemüller, M. Edwards, T. Peeters, A. P. Schaffers,

et al. 2006. Parallel Declines in Pollinators and Insect-Pollinated Plants in Britain and the

Netherlands. Science 313:351–354.

Blaauw, B. R., and R. Isaacs. 2014. Larger patches of diverse floral resources increase insect pollinator

density, diversity, and their pollination of native wildflowers. Basic and applied ecology 15:701–711.

Blondel, V. D., J.-L. Guillaume, R. Lambiotte, and E. Lefebvre. 2008. Fast unfolding of communities in

large networks. Journal of statistical mechanics 2008:P10008.

Bosch, M., and N. M. Waser. 1999. Effects of local density on pollination and reproduction in

Delphinium nuttallianum and Aconitum columbianum (Ranunculaceae). American journal of botany

86:871–879.

Briscoe Runquist, R., D. Grossenbacher, S. Porter, K. Kay, and J. Smith. 2016. Pollinator-mediated

assemblage processes in California wildflowers. Journal of evolutionary biology 29:1045–1058.

Brosi, B. J., and H. M. Briggs. 2013. Single pollinator species losses reduce floral fidelity and plant

reproductive function. Proceedings of the National Academy of Sciences of the United States of

America 110:13044–13048.

Brown, B. J., R. J. Mitchell, and S. J. Graham. 2002. Competition for pollination between an invasive

species (Purple loosestrife) and a native congener. Ecology 83:2328–2336.

Brown, J. H., and A. Kodric-Brown. 1979. Convergence, Competition, and Mimicry in a Temperate

Community of Hummingbird-Pollinated Flowers. Ecology 60:1022–1035.

Bryant, J. A., C. Lamanna, H. Morlon, A. J. Kerkhoff, B. J. Enquist, and J. L. Green. 2008. Microbes on

mountainsides: Contrasting elevational patterns of bacterial and plant diversity. Proceedings of the

National Academy of Sciences 105:11505–11511.

Burkle, L. A., J. C. Marlin, and T. M. Knight. 2013. Plant-pollinator interactions over 120 years: loss of

species, co-occurrence, and function. Science 339:1611–1615.

Cadotte, M. W. 2013. Experimental evidence that evolutionarily diverse assemblages result in higher

productivity. Proceedings of the National Academy of Sciences of the United States of America

110:8996–9000.

Cadotte, M. W., S. E. Campbell, S.-P. Li, D. S. Sodhi, and N. E. Mandrak. 2018. Preadaptation and

Naturalization of Nonnative Species: Darwin's Two Fundamental Insights into Species Invasion.

Annual review of plant biology 69:661–684.

Cadotte, M. W., B. J. Cardinale, and T. H. Oakley. 2008. Evolutionary history and the effect of

biodiversity on plant productivity. Proceedings of the National Academy of Sciences of the United

States of America 105:17012–17017.

Cadotte, M. W., J. Cavender-Bares, D. Tilman, and T. H. Oakley. 2009. Using phylogenetic, functional

and trait diversity to understand patterns of plant community productivity. PloS one 4:e5695.

Cadotte, M. W., R. Dinnage, and D. Tilman. 2012. Phylogenetic diversity promotes ecosystem stability.

Ecology 93:S223–S233.

Campbell, D. R., P. Sosenski, and R. A. Raguso. 2019. Phenotypic plasticity of floral volatiles in

response to increasing drought stress. Annals of botany 123:601–610.

CaraDonna, P. J., A. M. Iler, and D. W. Inouye. 2014. Shifts in flowering phenology reshape a subalpine

plant community. Proceedings of the National Academy of Sciences of the United States of America

111:4916–4921.

CaraDonna, P. J., and D. W. Inouye. 2015. Phenological responses to climate change do not exhibit

phylogenetic signal in a subalpine plant community. Ecology 96:355–361.

Carvalheiro, L. G., J. C. Biesmeijer, G. Benadi, J. Fründ, M. Stang, I. Bartomeus, C. N. Kaiser-Bunbury,

et al. 2014. The potential for indirect effects between co-flowering plants via shared pollinators

depends on resource abundance, accessibility and relatedness. Ecology letters 17:1389–1399.

Cavender-Bares, J., K. H. Kozak, P. V. A. Fine, and S. W. Kembel. 2009. The merging of community

ecology and phylogenetic biology. Ecology letters 12:693–715.

Cleland, E. E., I. Chuine, A. Menzel, H. A. Mooney, and M. D. Schwartz. 2007. Shifting plant phenology

in response to global change. Trends in ecology & evolution 22:357–365.

Cordes, L. S., D. T. Blumstein, K. B. Armitage, P. J. CaraDonna, D. Z. Childs, B. D. Gerber, J. G. A.

Martin, et al. 2020. Contrasting effects of climate change on seasonal survival of a hibernating

mammal. Proceedings of the National Academy of Sciences of the United States of America

117:18119–18126.

Csardi, G., and T. Nepusz. 2006. The igraph software package for complex network research. Computer

Science.

Daehler, C. C. 2001. Darwin's naturalization hypothesis revisited. The American naturalist 158:324–330.

Darwin, C. 1859. On the Origin of Species. Murray, London.

Davies, T. J., M. C. Urban, B. Rayfield, M. W. Cadotte, and P. R. Peres-Neto. 2016. Deconstructing the

relationships between phylogenetic diversity and ecology: a case study on ecosystem functioning.

Ecology 97:2212–2222.

Dehling, D. M., S. A. Fritz, T. Töpfer, M. Päckert, P. Estler, K. Böhning-Gaese, and M. Schleuning.

2014. Functional and phylogenetic diversity and assemblage structure of frugivorous birds along an

elevational gradient in the tropical Andes. Ecography no–no.

Deng, L., K. Wang, J. Li, G. Zhao, and Z. Shangguan. 2016. Effect of soil moisture and atmospheric

humidity on both plant productivity and diversity of native grasslands across the Loess Plateau,

China. Ecological engineering 94:525–531.

Dormann, C. F., and R. Strauss. 2014. A method for detecting modules in quantitative bipartite networks.

Methods in ecology and evolution / British Ecological Society 5:90–98.

Dunne, J. A., J. Harte, and K. J. Taylor. 2003. Subalpine meadow flowering phenology responses to

climate change: Integrating experimental and gradient methods. Ecological monographs 73:69–86.

Eiserhardt, W. L., F. Borchsenius, C. M. Plum, A. Ordonez, and J.-C. Svenning. 2015. Climate-driven

extinctions shape the phylogenetic structure of temperate tree floras. Ecology letters 18:263–272.

Elzinga, J. A., A. Atlan, A. Biere, L. Gigord, A. E. Weis, and G. Bernasconi. 2007. Time after time:

flowering phenology and biotic interactions. Trends in ecology & evolution 22:432–439.

E-Vojtkó, A., F. de Bello, Z. Lososová, and L. Götzenberger. 2023. Phylogenetic diversity is a weak

proxy for functional diversity but they are complementary in explaining community assembly

patterns in temperate vegetation. The Journal of ecology.

Faith, D. P. 1992. Conservation evaluation and phylogenetic diversity. Biological conservation 61:1–10.

Faust, M. N., and A. M. Iler. 2022. Pollinator-mediated reproductive consequences of altered co-

flowering under climate change conditions depend on abiotic context. Climate Change Ecology

3:100043.

Forrest, J., D. W. Inouye, and J. D. Thomson. 2010. Flowering phenology in subalpine meadows: Does

climate variation influence community co-flowering patterns? Ecology 91:431–440.

Forrest, J., and A. J. Miller-Rushing. 2010. Toward a synthetic understanding of the role of phenology in

ecology and evolution. Philosophical transactions of the Royal Society of London. Series B,

Biological sciences 365:3101–3112.

Forrest, J., and J. D. Thomson. 2010. Consequences of variation in flowering time within and among

individuals of Mertensia fusiformis (Boraginaceae), an early spring wildflower. American journal of

botany 97:38–48.

Freimuth, J., O. Bossdorf, J. F. Scheepens, and F. M. Willems. 2022. Climate warming changes

synchrony of plants and pollinators. Proceedings. Biological sciences / The Royal Society

289:20212142.

Gallagher, M. K., and D. R. Campbell. 2017. Shifts in water availability mediate plant-pollinator

interactions. The New phytologist 215:792–802.

Galland, T., G. Adeux, H. Dvořáková, A. E-Vojtkó, I. Orbán, M. Lussu, J. Puy, et al. 2019. Colonization

resistance and establishment success along gradients of functional and phylogenetic diversity in

experimental plant communities. The Journal of ecology 107:2090–2104.

Gerhold, P., J. F. Cahill, M. Winter, I. V. Bartish, and A. Prinzing. 2015. Phylogenetic patterns are not

proxies of community assembly mechanisms (they are far better). Functional ecology 29:600–614.

Gezon, Z. J., D. W. Inouye, and R. E. Irwin. 2016. Phenological change in a spring ephemeral:

implications for pollination and plant reproduction. Global change biology 22:1779–1793.

Ghazoul, J. 2006. Floral diversity and the facilitation of pollination. The Journal of ecology 94:295–304.

González-Orozco, C. E., L. J. Pollock, A. H. Thornhill, B. D. Mishler, N. Knerr, S. W. Laffan, J. T.

Miller, et al. 2016. Phylogenetic approaches reveal biodiversity threats under climate change. Nature

climate change 6:1110–1114.

Graham, C. H., and P. V. A. Fine. 2008. Phylogenetic beta diversity: linking ecological and evolutionary

processes across space in time. Ecology letters 11:1265–1277.

Harte, J., M. S. Torn, F.-R. Chang, B. Feifarek, A. P. Kinzig, R. Shaw, and K. Shen. 1995. Global

Warming and Soil Microclimate: Results from a Meadow-Warming Experiment. Ecological

applications: a publication of the Ecological Society of America 5:132–150.

Hillebrand, H. 2004. On the generality of the latitudinal diversity gradient. The American naturalist

163:192–211.

Huang, M., X. Liu, M. W. Cadotte, and S. Zhou. 2020. Functional and phylogenetic diversity explain

different components of diversity effects on biomass production. Oikos 129:1185–1195.

Inouye, D. W. 2008. Effects of Climate Change on Phenology, Frost Damage, and Floral Abundance of

Montane Wildflowers. Ecology 89:353–362.

Jentsch, A., J. Kreyling, J. Boettcher-treschkow, and C. Beierkuhnlein. 2009. Beyond gradual warming:

extreme weather events alter flower phenology of European grassland and heath species. Global

change biology 15:837–849.

Johnson, C. A., P. Dutt, and J. M. Levine. 2022. Competition for pollinators destabilizes plant

coexistence. Nature 607:721–725.

Johnson, D., G. K. Phoenix, and J. P. Grime. 2008. Plant community composition, not diversity, regulates

soil respiration in grasslands. Biology letters 4:345–348.

Jones, E. I., S. L. Nuismer, and R. Gomulkiewicz. 2013. Revisiting Darwin's conundrum reveals a twist

on the relationship between phylogenetic distance and invasibility. Proceedings of the National

Academy of Sciences of the United States of America 110:20627–20632.

Kembel, S. W., P. D. Cowan, M. R. Helmus, W. K. Cornwell, H. Morlon, D. D. Ackerly, S. P. Blomberg,

et al. 2010. Picante: R tools for integrating phylogenies and ecology. Bioinformatics 26:1463–1464.

Kharouba, H. M., J. Ehrlén, A. Gelman, K. Bolmgren, J. M. Allen, S. E. Travers, and E. M. Wolkovich.

2018. Global shifts in the phenological synchrony of species interactions over recent decades.

Proceedings of the National Academy of Sciences of the United States of America 115:5211–5216.

Koh, I., E. V. Lonsdorf, N. M. Williams, C. Brittain, R. Isaacs, J. Gibbs, and T. H. Ricketts. 2016.

Modeling the status, trends, and impacts of wild bee abundance in the United States. Proceedings of

the National Academy of Sciences of the United States of America 113:140–145.

Lehmann, J., C. M. Hansel, C. Kaiser, M. Kleber, K. Maher, S. Manzoni, N. Nunan, et al. 2020.

Persistence of soil organic carbon caused by functional complexity. Nature geoscience 13:529–534.

Levin, D. A., and W. W. Anderson. 1970. Competition for Pollinators between Simultaneously Flowering

Species. The American naturalist 104:455–467.

Linnaeus, C. 1799. Species plantarum. Impensis G. C. Nauk.

MacArthur, R. H. 1969. Patterns of communities in the tropics. Biological journal of the Linnean Society.

Linnean Society of London 1:19–30.

Ma, C., S.-P. Li, Z. Pu, J. Tan, M. Liu, J. Zhou, H. Li, et al. 2016. Different effects of invader-native

phylogenetic relatedness on invasion success and impact: a meta-analysis of Darwin's naturalization

hypothesis. Proceedings. Biological sciences / The Royal Society 283.

Maitner, B. S., D. S. Park, B. J. Enquist, and K. M. Dlugosch. 2022. Where we've been and where we're

going: the importance of source communities in predicting establishment success from phylogenetic

relationships. Ecography 2022.

Martín González, A. M., B. Dalsgaard, and J. M. Olesen. 2010. Centrality measures and the importance of

generalist species in pollination networks. Ecological Complexity 7:36–43.

Martín-Hernanz, S., R. G. Albaladejo, S. Lavergne, E. Rubio, M. Marín-Rodulfo, J. Arroyo, and A.

Aparicio. 2023. Strong conservatism of floral morphology during the rapid diversification of the

genus Helianthemum. American journal of botany 110:e16155.

Mayfield, M. M., and J. M. Levine. 2010. Opposing effects of competitive exclusion on the phylogenetic

structure of communities. Ecology letters 13:1085–1093.

Mazzochini, G. G., C. R. Fonseca, G. C. Costa, R. M. Santos, A. T. Oliveira-Filho, and G. Ganade. 2019.

Plant phylogenetic diversity stabilizes large-scale ecosystem productivity. Global ecology and

biogeography: a journal of macroecology 28:1430–1439.

Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aasa, R. Ahas, K. Alm-Kübler, et al. 2006. European

phenological response to climate change matches the warming pattern. Global change biology

12:1969–1976.

Miao, R., Y. Miao, Y. Liu, M. Guo, G. Fu, R. Xiao, Z. Yang, et al. 2022. Late-season drought exerts more

negative effects on plant diversity and cover than early-season drought through changing soil

moisture in a semi-arid grassland. Journal of vegetation science: official organ of the International

Association for Vegetation Science 33.

Moeller, D. a., and M. a. Geber. 2005*a*. Ecological context of the evolution of self-pollination in Clarkia

xantiana: population size, plant communities, and reproductive assurance. Evolution; international

journal of organic evolution 59:786–799.

Moeller, D. A. 2004. Facilitative Interactions Among Plants Via Shared Pollinators. Ecology 85:3289–

3301.

Moeller, D. A., and M. A. Geber. 2005*b*. ECOLOGICAL CONTEXT OF THE EVOLUTION OF SELF-

POLLINATION IN CLARKIA XANTLANA: POULATION SIZE, PLANT COMMUNITIES, AND

REPRODUCTTIVE ASSURANCE. Evolution 59:786–799.

Molina-Venegas, R., M. Á. Rodríguez, M. Pardo-de-Santayana, C. Ronquillo, and D. J. Mabberley. 2021.

Maximum levels of global phylogenetic diversity efficiently capture plant services for humankind.

Nature ecology & evolution 5:583–588.

Mooney, E., M. Mullins, J. Den Uyl, S. Trail, P. Nguyen, J. Owens, E. Godtfredsen, et al. 2021. Early

snowmelt reduces aphid abundance (Aphis asclepiadis) by creating water-stressed host plants

(Ligusticum porteri) and altering interactions with ants. Arthropod-plant interactions 15:33–46.

Moyes, A. B., and D. R. Bowling. 2013. Interannual variation in seasonal drivers of soil respiration in a

semi-arid Rocky Mountain meadow. Biogeochemistry 113:683–697.

Munguía-Rosas, M. A., J. Ollerton, V. Parra-Tabla, and J. A. De-Nova. 2011. Meta-analysis of

phenotypic selection on flowering phenology suggests that early flowering plants are favoured.

Ecology letters 14:511–521.

Newman, M. E. J., and M. Girvan. 2004. Finding and evaluating community structure in networks.

Physical review. E, Statistical, nonlinear, and soft matter physics 69:026113.

Ogilvie, J. E., S. R. Griffin, Z. J. Gezon, B. D. Inouye, N. Underwood, D. W. Inouye, and R. E. Irwin.

2017. Interannual bumble bee abundance is driven by indirect climate effects on floral resource

phenology. Ecology letters 20:1507–1515.

Ogilvie, J. E., and J. D. Thomson. 2016. Site fidelity by bees drives pollination facilitation in sequentially

blooming plant species. Ecology 97:1442–1451.

Ollerton, J., R. Winfree, and S. Tarrant. 2011. How many flowering plants are pollinated by animals?

Oikos 120:321–326.

O'Meara, B. C., S. D. Smith, W. S. Armbruster, L. D. Harder, C. R. Hardy, L. C. Hileman, L. Hufford, et

al. 2016. Non-equilibrium dynamics and floral trait interactions shape extant angiosperm diversity.

Proceedings. Biological sciences / The Royal Society 283.

Owen, N. R., R. Gumbs, C. L. Gray, and D. P. Faith. 2019. Global conservation of phylogenetic diversity

captures more than just functional diversity. Nature communications 10:859.

Pareja-Bonilla, D., M. Arista, L. P. C. Morellato, and P. L. Ortiz. 2023. Better soon than never: climate

change induces strong phenological reassembly in the flowering of a Mediterranean shrub

community. Annals of botany.

Park, D. S., X. Feng, B. S. Maitner, K. C. Ernst, and B. J. Enquist. 2020. Darwin's naturalization

conundrum can be explained by spatial scale. Proceedings of the National Academy of Sciences

117:10904–10910.

Pederson, G. T., S. T. Gray, T. Ault, W. Marsh, D. B. Fagre, A. G. Bunn, C. A. Woodhouse, et al. 2011.

Climatic Controls on the Snowmelt Hydrology of the Northern Rocky Mountains. Journal of climate

24:1666–1687.

Pelser, P. B., B. Nordenstam, J. W. Kadereit, and L. E. Watson. 2007. An ITS phylogeny of tribe

Senecioneae (Asteraceae) and a new delimitation of Senecio L. Taxon 56:1077–1104.

Pennell, M. W., J. M. Eastman, G. J. Slater, J. W. Brown, J. C. Uyeda, R. G. FitzJohn, M. E. Alfaro, et al.

2014. geiger v2.0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic

trees. Bioinformatics 30:2216–2218.

Potts, S. G., V. Imperatriz-Fonseca, H. T. Ngo, M. A. Aizen, J. C. Biesmeijer, T. D. Breeze, L. V. Dicks,

et al. 2016. Safeguarding pollinators and their values to human well-being. Nature 540:220–229.

Powers, J. M., H. M. Briggs, R. G. Dickson, X. Li, and D. R. Campbell. 2022. Earlier snow melt and

reduced summer precipitation alter floral traits important to pollination. Global change biology

28:323–339.

Prather, R. M., N. Underwood, R. M. Dalton, B. Barr, and B. D. Inouye. 2023. Climate data from the

Rocky Mountain Biological Laboratory (1975-2022). Ecology e4153.

Proches, Ş., J. R. U. Wilson, D. M. Richardson, and M. Rejmánek. 2007. Searching for phylogenetic

pattern in biological invasions. Global ecology and biogeography: a journal of macroecology 17:5–

10.

Proulx, S. R., D. E. L. Promislow, and P. C. Phillips. 2005. Network thinking in ecology and evolution.

Trends in ecology & evolution 20:345–353.

R Core Team. 2023. R: A language and environment for statistical computing.

Rejmánek, M. 1996. A theory of seed plant invasiveness: The first sketch. Biological conservation

78:171–181.

Ricciardi, A., and M. Mottiar. 2006. Does Darwin's naturalization hypothesis explain fish invasions?

Biological invasions 8:1403–1407.

Rivest, S., B. D. Inouye, and J. R. K. Forrest. 2023. Warmer springs increase potential for temporal

reproductive isolation among habitat patches in subalpine flowering plants. The Journal of ecology

111:2257–2268.

Sargent, R. D., and D. D. Ackerly. 2008. Plant–pollinator interactions and the assembly of plant

communities. Trends in ecology & evolution 23:123–130.

Schiffer, A., X. Loy, C. Morozumi, and B. J. Brosi. 2023. Differences in individual flowering time

change pollen limitation and seed set in three montane wildflowers. American journal of botany

110:1–14.

Schluter, D., and M. W. Pennell. 2017. Speciation gradients and the distribution of biodiversity. Nature

546:48–55.

Schoener, T. W. 1970. Nonsynchronous Spatial Overlap of Lizards in Patchy Habitats. Ecology 51:408–

418.

Serafini, J., P. Grogan, and L. Aarssen. 2019. Summer precipitation limits plant species richness but not

overall productivity in a temperate mesic old-field meadow. Journal of vegetation science: official

organ of the International Association for Vegetation Science 30:832–844.

Shrestha, M., A. G. Dyer, J. E. Garcia, and M. Burd. 2019. Floral colour structure in two Australian

herbaceous communities: it depends on who is looking. Annals of botany 124:221–232.

Sloat, L. L., A. N. Henderson, C. Lamanna, and B. J. Enquist. 2015. The Effect of the Foresummer

Drought on Carbon Exchange in Subalpine Meadows. Ecosystems 18:533–545.

Smith, S. A., and J. W. Brown. 2018. Constructing a broadly inclusive seed plant phylogeny. American

journal of botany 105:302–314.

Suárez-Mariño, A., G. Arceo-Gómez, C. Albor, and V. Parra-Tabla. 2022. Co-flowering modularity and

floral trait similarity help explain temporal changes in plant–pollinator network structure. Plant

Ecology 223:1289–1304.

Thuiller, W., L. Gallien, I. Boulangeat, F. De Bello, T. Münkemüller, C. Roquet, and S. Lavergne. 2010.

Resolving Darwin's naturalization conundrum: a quest for evidence. Diversity and Distributions

16:461–475.

Tietje, M., A. Antonelli, F. Forest, R. Govaerts, S. A. Smith, M. Sun, W. J. Baker, et al. 2023. Global

hotspots of plant phylogenetic diversity. The New phytologist 240:1636–1646.

Tilman, D., C. L. Lehman, and K. T. Thomson. 1997. Plant diversity and ecosystem productivity:

theoretical considerations. Proceedings of the National Academy of Sciences of the United States of

America 94:1857–1861.

Tucker, C. M., M. W. Cadotte, S. B. Carvalho, T. J. Davies, S. Ferrier, S. A. Fritz, R. Grenyer, et al.

2017. A guide to phylogenetic metrics for conservation, community ecology and macroecology.

Biological reviews of the Cambridge Philosophical Society 92:698–715.

Valiente-Banuet, A., and M. Verdú. 2007. Facilitation can increase the phylogenetic diversity of plant

communities. Ecology letters 10:1029–1036.

Vamosi, J. C., and S. M. Vamosi. 2010. Key innovations within a geographical context in flowering

plants: towards resolving Darwin's abominable mystery. Ecology letters 13:1270–1279.

Veldhuisen, L. 2023. Community-level flowering & fitness data across an elevational gradient, Rocky

Mountain Biological Lab, 2021-2022.

Wagg, C., M. J. O'Brien, A. Vogel, M. Scherer-Lorenzen, N. Eisenhauer, B. Schmid, and A. Weigelt.

2017. Plant diversity maintains long-term ecosystem productivity under frequent drought by

increasing short-term variation. Ecology 98:2952–2961.

Waser, N. M., and M. V. Price. 2016. Drought, pollen and nectar availability, and pollination success.

Ecology 97:1400–1409.

Webb, C. O. 2000. Exploring the Phylogenetic Structure of Ecological Communities: An Example for

Rain Forest Trees. The American naturalist 156:145–155.

Webb, C. O., D. D. Ackerly, M. A. McPeek, and M. J. Donoghue. 2002. Phylogenies and Community

Ecology. Annual review of ecology and systematics 33:475–505.

Wei, N., R. L. Kaczorowski, G. Arceo-Gómez, E. M. O'Neill, R. A. Hayes, and T.-L. Ashman. 2021.

Pollinators contribute to the maintenance of flowering plant diversity. Nature 1–5.

Willis, C. G., B. Ruhfel, R. B. Primack, A. J. Miller-Rushing, and C. C. Davis. 2008. Phylogenetic

patterns of species loss in Thoreau's woods are driven by climate change. Proceedings of the National

Academy of Sciences 105:17029–17033.

Wolf, A. A., E. S. Zavaleta, and P. C. Selmants. 2017. Flowering phenology shifts in response to

biodiversity loss. Proceedings of the National Academy of Sciences of the United States of America

114:3463–3468.

Wolowski, M., L. G. Carvalheiro, and L. Freitas. 2017. Influence of plant-pollinator interactions on the

assembly of plant and hummingbird communities. The Journal of ecology 105:332–344.

Xiang, G., Y. Jiang, J. Lan, L. Huang, L. Hao, Z. Liu, and J. Xia. 2023. Different influences of

phylogenetically conserved and independent floral traits on plant functional specialization and

pollination network structure. Frontiers in plant science 14:1084995.

Zhang, J. 2016. *Package "spaa"* (0.2.2 ed.).

| | Co-Flowering Networks | | | Faith's PD | | | MPD | | | MNTD | | |
Site	Modularity (Q)	Z-score	Edges	Value	SES	*P*-value	Value	SES	*P*-value	Value	SES	*P*-value
2021 Flowering Season												
High	0.42	19.75	171	1637.1	1.45	0.93	242.4	1.86	0.99*	131.8	0.37	0.64
Medium	0.48	28.22	231	1503.5	-0.80	0.21	221.8	0.15	0.52	97.5	-1.40	0.08
Low	0.44	24.27	231	1720.7	0.33	0.63	220.3	0.03	0.48	123.6	0.42	0.65
2022 Flowering Season												
High	0.36	14.56	153	1428.8	0.27	0.60	228.4	0.32	0.58	126.6	0.02	0.50
Medium	0.43	26.86	276	1677.2	-0.09	0.47	232.6	0.94	0.83	106.2	-0.59	0.28
Low	0.56	43.12	325	1841.5	-0.10	0.46	216.9	-1.16	0.13	114.4	0.25	0.59

Low site (2815 m)   Middle site (3165 m)   High site (3380 m)

2021

2022

Mid2

Module Color Key
Early
Mid
Late