**Hyperactivation of MEK1 in cortical glutamatergic neurons results in projection axon**

**deficits and aberrant motor learning**

*George R. Bjorklund[1,2], *Katherina P. Rees[1], Kavya Balasubramanian[1], Lauren T. Hewitt[1,3], Kenji

Nishimura[1,4], Jason M. Newbern[1#]

[1]School of Life Sciences and [2]School of Biological and Health Systems Engineering, Arizona

State University, Tempe, AZ, 85287

[3]Present address: Porter Neuroscience Center, Eunice Kennedy-Shriver National Institute of

Child Health and Human Development, National Institutes of Health, Bethesda, Maryland, 20892

[4]Present address: Institute for Neuroscience, The University of Texas-Austin, Austin, Texas,

78712

*These authors contributed equally

[#]Corresponding author & ORCID ID# 0000-0003-2865-0112

**_Keywords_**

connectivity, cortex, development, RASopathy, kinase, axon

**_Summary statement:_**

MEK-ERK1/2 hyperactivation in developing cortical excitatory neurons is sufficient to decrease

long-range axonal outgrowth, which coincides with reduced Arc expression and deficits in aspects

of skilled motor learning by adulthood.

***Abstract***

Abnormal Extracellular Regulated Kinase 1/2 (ERK1/2) signaling is linked to multiple

neurodevelopmental diseases, especially the RASopathies, which typically exhibit ERK1/2

hyperactivation in neurons and non-neuronal cells. To better understand how excitatory neuron-

autonomous ERK1/2 activity regulates the development of the mouse motor cortex, we

conditionally expressed a hyperactive MEK1$^{S217/221E}$ variant using *Nex/NeuroD6:Cre*. Our results

show that MEK1$^{S217/221E}$ expression led to persistent hyperactivation of ERK1/2 in neocortical

axons, but not excitatory neuron somas or nuclei. We noted reduced axonal arborization of

multiple subcortical target domains in mutants and reduced cortical expression of the activity

dependent gene, ARC. These changes did not coincide with significant differences in voluntary

locomotor activity or motor performance in the accelerating rotarod task. However, motor learning

in a single-pellet retrieval task was significantly diminished in *Nex/NeuroD6:Cre*; *MEK1$^{S217/221E}$*

mutants. Restriction of MEK1$^{S217/221E}$ expression to layer V cortical neurons recapitulated axonal

outgrowth deficits, however, had no effect on motor learning. Collectively, these results indicate

that within the cortex, glutamatergic neuron-autonomous hyperactivation of MEK1 is sufficient to

drive deficits in axon outgrowth, activity dependent gene expression, and skilled motor learning.

## *Introduction*

Many extracellular cues important for forebrain development, such as fibroblast growth factors (FGFs) and neurotrophins, activate receptor tyrosine kinases (RTKs) to modulate a wide range of cellular behaviors. Ligand-bound RTKs initiate intracellular signaling via numerous kinase cascades, including the PI3K, PLC, PKC, and Extracellular Regulated Kinase 1/2 (ERK1/2) pathways (Lemmon and Schlessinger, 2010). Activation of ERK1/2 downstream of RTK signaling has been well-studied, which requires RAS-dependent activation of RAF isoforms and subsequent phosphorylation of MEK1 and MEK2 at two conserved residues in the kinase domain, $Ser^{218/222}$ and $Ser^{222/226}$, respectively (Klomp et al., 2021; Lavoie et al., 2020). Activated MEK1/2 then phosphorylates ERK1 and ERK2, which ultimately act upon hundreds of cytosolic and nuclear substrates (Ünal et al., 2017). ERK1/2 signaling has frequently been linked to the control of cell proliferation, differentiation, survival, synaptic plasticity, and learning (Atkins et al., 2001; Gómez-Palacio-Schjetnan and Escobar, 2008; Jeanneteau et al., 2010; Kalil and Dent, 2014; Mebratu and Tesfaigzi, 2009; Nowaczyk et al., 2014; Polleux and Snider, 2010; Thomas and Huganir, 2004). However, this broad repertoire obscures the intricate, context-dependent functions of ERK1/2 signaling that are highly contingent on the precise developmental stage, cell type, and/or extracellular cue (Santini et al., 2019; Shaul and Seger, 2007). Developing neurons in particular exhibit complex, subtype specific responses to modifications in ERK1/2 activity that are not fully understood (Ryu and Lee, 2016).

Key regulators in the RTK-ERK1/2 signaling network are frequently mutated in a family of developmental syndromes known collectively as the "RASopathies" (Jindal et al., 2015; Kim and Baek, 2019; Rauen, 2013; Rauen, 2022). Well-defined mutations have been identified in individual syndromes within the RASopathy superfamily, such as Noonan syndrome (NS), Neurofibromatosis type 1 (NF1), Costello syndrome (CS), and Cardio-facio-cutaneous syndrome (CFC). NS and NF1 are common RASopathies with a combined frequency of ~1:2000 births (Rauen, 2013; Tidyman and Rauen, 2009). However, >50 distinct RASopathy mutations in over 10 different genes within the RTK-ERK1/2 network have been reported (Brown et al., 2012;

Gutmann et al., 2012; Jindal et al., 2015; Kaul et al., 2015; Rauen, 2022). Upstream RASopathy

mutations in NF1, SHP2, and RAS have been shown to modulate PI3K/Akt, PKC, and/or

Rho/ROCK activity, in addition to the ERK1/2 pathway (Anastasaki and Gutmann, 2014; Brown

et al., 2012; Castellano and Downward, 2011; Kaul et al., 2015). ERK1/2 hyperactivation is,

however, thought to be a core, common feature of nearly all RASopathies. Thus, pharmacological

inhibitors of ERK1/2 regulators have been an important focus for therapeutic development

(Guilding et al., 2007; López-Juárez et al., 2017; Nakamura et al., 2009; Wang et al., 2012; Wu

et al., 2011). ERK1/2 also represents a convergent target downstream of multiple genetic risk

factors for autism and is abnormally activated in models of Fragile X and 16p11.2 deletion

syndrome (Chung et al., 2021; Osterweil et al., 2013; Wen et al., 2016). Therefore, understanding

the cell-specific effects of ERK1/2 activity in the developing forebrain may shed light on potential

neuropathological processes in multiple developmental disorders (Bernardino et al., 2022; Li et

al., 2005; Roskoski, 2019).

MRI-based studies suggest that differences in both cortical structural and functional

connectivity are a critical aspect of intellectual disability and cognitive changes in RASopathies

and autism (Aydin et al., 2016; Chang et al., 2016; Fattah et al., 2021; Filippi et al., 2013; Ibrahim

et al., 2017; Owen et al., 2018; Tomson et al., 2015; Zamboni et al., 2007). Changes in motor

cortex function have been detected in RASopathies and individuals with Noonan Syndrome note

that treatments for cognitive and musculoskeletal abnormalities are a high priority (Bernardino et

al., 2022; Doherty et al., 2023; Tiemens et al., 2023). Yet, it is difficult to discern whether functional

changes are a result of aberrant signaling in neurons, myelinating oligodendrocytes, or complex

interactions during development. Studies have shown a specific role for ERK1/2 signaling in

regulating rodent myelinating glial development *in vivo* (Ishii et al., 2012; Ishii et al., 2013;

Newbern et al., 2011), and pharmacological MEK1/2 inhibition rescues myelination deficits

caused by RASopathic signaling in mature oligodendrocytes (López-Juárez et al., 2017; Mayes

et al., 2013). Effects of upstream RASopathy mutations in SynGAP1, NF1, and RAS that activate

multiple downstream cascades, selectively in neurons, have been explored (Clement et al., 2012;

Cui et al., 2008; Kim et al., 2023; Rumbaugh et al., 2006). However, the neuron-autonomous

effects of downstream ERK1/2 hyperactivation in developing cortical excitatory circuits are less

clear. We previously reported subtype-specific functions for ERK1/2 in the development of cortical

inhibitory and excitatory neurons with a series of Cre-dependent conditional mouse lines. Deletion

of *Mek1/2* (*Map2k1/2*) in postmitotic excitatory neurons resulted in reduced corticospinal axon

extension into the lumbar spinal cord followed by death of a subset of layer V CTIP2$^+$ neurons in

the neonatal motor cortex (Xing et al., 2016). In contrast, hyperactivation of ERK1/2 signaling in

excitatory neurons via overexpression of a Cre-dependent, phosphomimic *MEK1$^{S217/221E}$* construct

did not alter CTIP2$^+$ density, but led to reductions in corticospinal elongation by P3 (Xing et al.,

2016).

Given the marked excitatory neuron subtype-specific outcomes associated with modifying

ERK1/2 activity, we sought to understand whether MEK1 hyperactivation in excitatory neurons

during development is sufficient to alter cortical axon outgrowth to multiple targets and ultimately

affect cognition. We utilized a *Nex/Neurod6:Cre* mouse to conditionally express a hyperactive

*MEK1$^{S217/221E}$* construct in newborn, postmitotic cortical glutamatergic neurons, but not

oligodendrocytes or cortical inhibitory neurons (Goebbels et al., 2006; Krenz et al., 2008).

*MEK1$^{S217/221E}$* selectively drives downstream hyperactivation of ERK1/2, independent of directly

manipulating parallel RTK-linked cascades. Moreover, *Nex:Cre*-directed expression minimizes

cellular contributions from glia and inhibitory neurons known to be modified by changes in ERK1/2

signaling (Angara et al., 2020; Holter et al., 2021; Kang and Lee, 2019; Knowles et al., 2023a;

Knowles et al., 2023b). This approach will help identify the excitatory neuron-autonomous

functions of selective hyperactivation of MEK-ERK1/2 signaling during long-range cortical circuit

formation.

Our results show that conditional *MEK1$^{S217/221E}$* overexpression leads to a persistent

increase in phosphorylated ERK1/2 (pERK1/2) levels in axonal projections that coincides with

reduced motor cortex derived axon arborization in intracortical and subcortical target domains.

Moreover, mutant cortices exhibited reduced expression of a canonical immediate early gene,

Arc, and deficits in learning a single-pellet retrieval task (Guo et al., 2015; Kawai et al., 2015;

Peters et al., 2017; Whishaw et al., 1993). To test whether these deficits were cortical layer V-

autonomous, we crossed the $MEK1^{217/221E}$ mouse line with a $Rbp4:Cre$ mouse. Viral tracing

analysis of contralateral corticocortical, corticostriatal, and corticospinal projections in $Rbp4:Cre$

$MEK1^{217/221E}$ mutants revealed a significant decrease in axonal innervation, but no deficit in skilled

motor learning. Collectively, these results indicate that excitatory neuron-specific hyperactivation

of MEK1 in the cortex is sufficient to disrupt the development of cortical axonal projections, the

expression of Arc, and select aspects of skilled motor learning.

**Materials and Methods**

*Transgenic Mice*

All animal experiments were performed in accordance with established procedures approved by the Institutional Animal Care and Use Committee of Arizona State University (Protocol# 22-1924R) and NIH guidelines for the use and care of laboratory animals. Mice used in this work were of a mixed genetic background, housed in standard conditions, and kept on a 12-hour light/dark cycle with *ad libitum* food and water. *NeuroD6/Nex-Cre (NeuroD6$^{tm1(cre)Kan}$)* mice were kindly provided by Dr. Klaus Nave and Dr. Sandra Goebbels (MGI:2668659) (Goebbels et al., 2006). *Rbp4:Cre* mice were obtained from the MMRRC (RRID:MMRRC_031125-UCD) (Gong et al., 2007). *Mek1$^{S217/221E}$ (Tg(CAG-cat,-Map2k1*)243Rbns)* mice were provided by Dr. Maike Krenz and Dr. Jeffrey Robbins (MGI:3822131) (Krenz et al., 2008). Cre-dependent *tdTomato/Ai9* (B6.-Gt(ROSA)26Sor$^{tom(CAG-tdTomato)Hze}$/J) mice were ordered from Jackson Laboratories (RRID:IMSR_JAX:007909) (Madisen et al., 2010). Each experiment was replicated a minimum of three times from independent litters.

Genomic DNA was extracted from toe or tail samples and amplified with standard polymerase chain reaction techniques. Primers for gene amplification are as follows 5'-3': Cre - GCTAAACATGCTTCATCGTCGG and GATCTCCGGTATTGAAACTCCAGC amplify a 645 base pair allele. *Mek1$^{S217/221E}$* – GTACCAGCTCGGCGGAGACCAA and TTGATCACAGCAATGCTAACTTTC amplify a 600 base pair allele. *Ai9* – CTGTTCCTGTACGGCATGG and GGCATTAAAGCAGCGTATCC amplify a 196 base pair allele.

*Tissue Preparation*

Transcardial perfusions were performed on mice using a 4% paraformaldehyde/PBS solution and dissected brains were post-fixed overnight at 4°C. Brains were then cryopreserved via serial incubations in 15% and 30% sucrose/PBS solutions prior to embedding in Tissue-Tek O.C.T. (Tissue-Tek 4583) and rapid freezing. Cryostat sections were collected in cold PBS for free-floating staining or mounted on Fisherbrand Superfrost Plus slides (Fisher Scientific 12-550-

162  15) and air-dried prior to staining. For vibratome sectioned material, fixed adult brains were

163  mounted in agarose and sections were collected in PBS prior to free-floating

164  immunohistochemistry.

### *Viral Injections and Image Quantification*

P1 pups were removed from their home cage as a group and individually cryo-

anesthetized on ice for 4-6 minutes. Viral injections were then performed immediately with 50nl

of solution using a 32-gauge beveled needle fitted to a Hamilton 5µl neuros syringe mounted on

a manipulator arm. After injections, pups were placed on a 37°C heated surface for recovery and

returned to their home cage as a group. Adeno-Associated Viral vectors containing a Cre-

dependent tdTomato construct under the control of a strong CAG promoter (AAV-CAG-FLEX-

tdTomato, UNC Viral Vector Core) were used for viral tracing experiments. For quantification of

virally labeled RFP$^+$ axonal arborization in the forebrain, we first manually assessed the number

of RFP$^+$ neurons in M1 from confocal images of at least three sections near the center of the

injection site. Confocal images of axonal target domains from at least three sections were

automatically thresholded in ImageJ using the Otsu method (Otsu, 1979). The total number of

RFP$^+$ pixels within the target domain was normalized to the number of labeled M1 neurons within

the injection site to provide a relative estimate of axonal arborization. To quantify axon elongation

along the spinal cord, RFP$^+$ immunolabeling in the cervical dorsal funiculus was normalized to

immunolabeling in hindbrain corticobulbar tract (CBT). Representative images were cropped,

adjusted for brightness and contrast, and inverted in Photoshop for clear visualization of axonal

projections.

### *Immunohistochemistry*

Tissue sections were rinsed with PBS/0.1% Triton X-100 (PBST) and blocked with 5%

normal donkey serum in PBST at room temperature for ~1 hour. Primary antibodies were diluted

in PBST/5%NDS and incubated overnight at 4°C with gentle agitation. Primary antibodies used

were: rabbit anti-MEK1 (RRID:AB776273, Abcam ab32091), rabbit anti-phospho-p44/42 MAPK

(ERK1/2) (Thr202/Tyr204) (RRID:AB_2315112, Cell Signaling Technologies 4370), rabbit anti-

ERK2 (RRID:AB_732210, Abcam ab32081), rabbit anti-ARC (RRID:AB_887694, Synaptic

Systems 156-003), mouse anti-β3Tubulin (TUJ1, RRID:AB_10063408, Bioloegend 801202),

rabbit   anti-RFP   (RRID:AB_2209751,   Rockland   600-401-379),   chicken   anti-RFP

(RRID:AB_10704808, Rockland 600-901-379), and mouse anti-NeuN (RRID:AB_2298772,

Millipore Sigma MAB377). After rinsing in PBS 3X, secondary antibodies diluted in PBST/5%NDS

with DAPI (Sigma Aldrich 10236276001) were added and incubated overnight at 4°C with gentle

agitation. Secondary antibodies include Alexa Fluor 488, 546, 568, and 647 conjugated to anti-

rabbit IgG, anti-mouse IgG, or anti-chicken IgY (Invitrogen). Imaging was performed using a Zeiss

LSM 800 or Leica SP5 laser scanning confocal microscope.

***Confocal Image Analysis and Quantification***

The number of pERK1/2 labeled cells in M1 layer II/III were counted from 20x confocal

images in at least two hemi-sections per mouse and averaged. We quantified the level of pERK1/2

immunolabeling in high resolution confocal images of layer V neurons collected with a 63X 1.4NA

objective using the same acquisition settings. Individual NeuN[+] pyramidal shaped somas >100

$\mu m^2$ with a clear DAPI-labeled nucleus in cortical layer V of the motor cortex were randomly

selected by a blinded observer. The integrated density of pERK1/2 immunolabeling in the

cytoplasm and nucleus was then measured. A total of 68 control and 83 mutant neurons were

analyzed from at least three sections per mouse, four mice per genotype, representing a total

sampling area >14,000 $\mu m^2$ per genotype.

To estimate the levels of pERK1/2 immunoreactivity in TUJ1[+] callosal axons, standard

confocal images of the corpus collosum were collected and analyzed. Standard confocal images

of at least two tissue sections of the corpus collosum per mouse were collected and automatically

thresholded in ImageJ in an unbiased manner using the default settings of Moments

Autothreshold (Tsai, 1985). Colocalized pERK1/2[+]/TUJ1[+] pixels were determined using Pierre

Bourdoncle's Colocalization Plugin with default settings and normalized to the total number of

TUJ1$^+$ pixels. Representative images were subsequently acquired in Airyscan mode using

optimized settings to visualize the localization of pERK1/2 more clearly within TUJ1$^+$ axons.

To quantify CST elongation in *RBP4:Cre; Ai9$^{+/-}$* mice, confocal images of the dorsal

funiculus in at least three lumbar segments per control and *MEK1$^{S217/221E}$* expressing mice were

autothresholded using Moments and the relative change in the number of RFP$^+$ pixels was

determined. Representative images have been cropped and adjusted for brightness and contrast

in Photoshop for presentation.

***Western Blotting***

Dorsal cortices were lysed in RIPA buffer (0.05M Tris-HCl, pH 7.4, 0.5M NaCl, 0.25%

deoxycholic acid, 1% Triton X-100, and 1 mM EDTA, Millipore) and supplemented with 0.1% SDS,

protease inhibitor (Sigma-Aldrich P2714), and phosphatase inhibitor cocktail II and III (Sigma-

Aldrich P5726) via sonication. Lysates were then cleared by centrifugation and protein

concentrations determined by Bradford Assay using a Peirce Protein Assay Kit (Thermo Scientific

23200). Equal amounts of protein were denatured in reducing sample buffer, separated by SDS-

PAGE gels (BioRad 456-1035 and 456-8095), and transferred to PVDF membranes (BioRad 162-

0177). Blots were blocked with 5% low-fat milk TBS containing 0.1% Tween-20 (TBST) for 1hr at

room temperature, then incubated overnight at 4°C with primary antibodies in 4% BSA in TBST.

Primary antibodies used were: rabbit anti-MEK1 (RRID:AB_776273, Abcam ab32091), rabbit anti-

phospho-p44/42 MAPK (ERK1/2) (Thr202/Tyr204) (RRID:AB_2315112, Cell Signaling

Technologies 4370), rabbit anti-ERK2 (RRID:AB_732210, Abcam ab32081), rabbit anti-ARC

(RRID:AB_887694, Synaptic Systems 156-003), rabbit anti-GAPDH (RRID:AB_561053, Cell

Signaling Technologies 2118), and rabbit anti-β-actin (RRID:AB_10950489, Cell Signaling

Technologies 8457). After washing with TBST, membranes were incubated with donkey anti-

rabbit HRP-conjugated secondary antibodies (Jackson Immunoresearch 711-035-152) in 5%

milk/TBST for 2hr at room temperature. Membranes were washed with TBST and signal was

detected with SuperSignal West Pico chemiluminescent (Thermo Scientific 32106) and exposed

to radiographic film (Thermo Scientific 34089). Quantification of Western Blots was performed

using a high-resolution scanned image of the exposed radiographic film in a grayscale TIFF file

format. Bands were selected for quantification of integrated density in ImageJ following

background subtraction from an unlabeled region and normalized to either the loading control or

non-phosphorylated target.

***Open Field, Accelerating Rotarod, and Single-Pellet Retrieval***

Open field testing was performed in an open top opaque plexiglass box measuring

38x38x30cm. The area within the box is illuminated by a centered, overhead 40-watt spotlight.

Video recordings of open field behavior were scored by a blinded observer. Rotarod analysis was

performed on a San Diego Instruments accelerating rotarod. Animals were tested for three trials

per day for five consecutive days, accelerating the rotarod from 4 to 40 RPM with a 15min intertrial

interval. The latency to fall was measured for each of three trials during a day and averaged.

A single-pellet retrieval task was performed using only male mice that were acclimated to

sucrose pellets (TestDiet 1811555) in their home cages at least one week prior to testing. Mice

were habituated for 10-15min per day, for two days in the testing room and chamber, which

consists of a 15x10x20cm transparent acrylic box with a 5mm wide slit on the right and left side

of one end that opens to a 10mm high food platform. During the subsequent two training days,

mice were observed while single sucrose pellets were made regularly available during a 10-20min

session. Only mice that attempted at least 20 reaches were further analyzed. Daily testing was

performed for 10 min per day, for 5 days by placing a single pellet on the food platform. To

enhance motivation for obtaining sucrose pellets, *Nex:Cre;MEK1$^{S217/221E}$* mice were mildly food

restricted by limiting free feeding to 2-6 hours daily while maintaining at least 90% of their initial

weight. Due to unexpected negative health effects of mild food restriction in *RBP4:Cre;*

*MEK1$^{S217/221E}$* mice, these mutants and associated control mice were only restricted during a 2-4

hr period prior to a single 5 min session per day.

Attempts were scored as a "success" if the mouse was able to retrieve the sucrose pellet

with its preferred paw and place it in its mouth. If the mouse grasped the pellet but failed to return

it to its mouth, the attempt was scored as a "drop". If the mouse clearly reached for the pellet but

failed to grasp it, the attempt was categorized as a "miss". For each mouse, the proportion of

successful out of total attempts was calculated. Accelerating rotarod and single-pellet retrieval

results were analyzed using a two-way repeated measures ANOVA in SPSS.

**Results**

*Hyperactivation of MEK1 increases levels of pERK1/2 in cortical excitatory neuron axons*

To investigate the developmental effects of excitatory neuron-autonomous hyperactive MEK-ERK1/2 signaling within the cortex *in vivo*, we generated mice expressing *Nex/NeuroD6:Cre*, Cre-dependent hyperactive *Mek1/Map2k1$^{S217/221E}$*, and Cre-dependent tdTomato/RFP (*Ai9$^{+/-}$*) (Goebbels et al., 2006; Krenz et al., 2008; Madisen et al., 2010). As expected, E14.5 *Nex:Cre; Ai9$^{+/-}$* forebrains showed RFP expression in excitatory neurons and axons in the cortical plate (CP) and intermediate zone (IZ), but not in neural stem cells in the ventricular zone (VZ) (Fig. 1A-B) (Xing et al. 2016, Goebbels et al. 2006). Substantial MEK1 overexpression was apparent in recombined excitatory neuron somas and long-range axonal projections in *Nex:Cre; MEK1$^{S217/221E}$; Ai9$^{+/-}$* compared to controls (Fig. 1B-C, G-H). Western blots of dorsal cortical lysates from E14.5 *Nex:Cre; MEK1$^{S217/221E}$* embryos demonstrate a 5.29 ± 0.93 fold-increase in MEK1 expression relative to controls (Mean ± SEM, n=5, Student's t-test p=0.002) (Fig. 1K-L).

*MEK1$^{S217/221E}$* hyperphosphorylates ERK1/2 in cell free assays and in culture, but the extent of downstream effects *in vivo* varies in different tissue types (Alessi et al., 1994; Bueno, 2000; Cowley, 1994; Klesse et al., 1999; Krenz et al., 2008; Lajiness et al., 2014; Li et al., 2012). We directly measured whether excitatory neuronal *MEK1$^{S217/221E}$* overexpression coincides with increased phosphorylated ERK1/2 (pERK1/2) levels *in vivo*. E14.5 dorsal cortical lysates from mutant mice exhibited a significant 1.61 ± 0.19 fold-increase in pERK1/2 levels compared to controls (Mean ± SEM, n=10, Student's t-test p=0.005) (Fig. 1M-N). Immunolabeling for pERK1/2 in *Nex:Cre; MEK1$^{S217/221E}$; Ai9$^{+/-}$* mice revealed a relative qualitative enrichment of pERK1/2 in the intermediate zone and nascent RFP$^{+}$ cortical excitatory neuron axons (Fig. 1I-J, white arrows), while comparable levels of pERK1/2 were observed in the unrecombined VZ of control and mutant embryos.

We next evaluated MEK1 and pERK1/2 activation in adult brains. Adult *Nex:Cre; MEK1$^{S217/221E}$* cortical lysates revealed a statistically significant 1.61 ± 0.2 fold increase in MEK1

# E14.5 Cortex

**Figure 1. MEK1 hyperactivation selectively in immature glutamatergic neurons results in increased levels of pERK1/2 by E14.5**

**A-J.** Immunolabeling of E14.5 *Nex:Cre; MEK1$^{S217/221E}$; Ai9$^{+/-}$* mutant cortices shows a substantial increase of MEK1 expression in the intermediate zone (IZ) and cortical plate (CP) of mutants (F-H) when compared to *Nex:Cre; Ai9$^{+/-}$* controls (A-C) (n=4). RFP$^+$ axons within the IZ of mutants (I-J, white arrow heads) exhibited a qualitative enrichment in levels of phosphorylated ERK1/2 relative to controls (D-E) (n=4). We did not detect a notable difference in the levels of pERK1/2 in the ventricular zone (VZ), where recombination is essentially absent in these mice.

**K-N.** Western blots of E14.5 dorsal cortical lysates (K) reveal a 5-fold increase of MEK1 in mutants compared to controls (L) (Mean ± SEM, n=5, * = Student's t-test p=0.002). We also observed a significant increase in pERK1/2 levels in mutants compared to controls (M-N) (Mean ± SEM, n=10, * = Student's t-test p=0.005).

Scale bars: A, F = 100 µm, B-E, G-J = 60µm.

expression over controls (mean ± SEM, n=3, Student's t-test p=0.041) (Fig. 2A-B). The proportion of excitatory neurons in whole adult cortical lysates is relatively less than at E14.5 and may contribute to the reduced fold-increase in MEK1 in adult mutants when compared to E14.5 mutants. We did not detect a significant difference in pERK1/2 levels between *Nex:Cre; MEK1$^{S217/221E}$* and control whole cortical lysates (n=3, Student's t-test p=0.832) (Fig. 2C-D). MEK1 immunolabeling of the cortex revealed a substantial increase in mutant excitatory neurons somas and axons within the corpus callosum when compared to the controls (Fig. 2E-H).

Control forebrains immunolabeled for pERK1/2 showed the expected enrichment in a subset of neuronal somas in layer II/III, with relative low levels in the corpus callosum (Fig. 2I-J) (Hart and Balleine, 2016; Lee et al., 2021). Interestingly, in *Nex:Cre; MEK1$^{S217/221E}$* mutant mice, we observed a significant decrease in pERK1/2$^{+}$ layer II/III neurons in M1 when compared to controls (n=6, Student's t test p=0.002) (Fig. 2U). Moreover, semi-quantitative high-resolution analyses of pERK1/2 immunolabeling intensity in mutant layer V neurons revealed a reduction in the cytoplasm and the nucleus when compared to controls (n=68 control and 83 mutant neurons from four mice per genotype, Student's t-test p<0.01) (Fig. 2M-P, V). In contrast, we noted a qualitative increase in pERK1/2-immunolabeling in a number of mutant forebrain white matter tracts, such as the corpus callosum (Fig. 2L, yellow arrows), fimbria, and stria terminalis (Fig. S1), which are enriched in axons derived from excitatory populations in the cortex, hippocampus, and amygdala, respectively. High resolution analysis of pERK1/2 co-localization with TUJ1, a marker of axons, in the corpus callosum, demonstrated the presence of pERK1/2 in axons and a modest increase in co-localization in mutant axons compared to controls (n=4, Student's t test p = 0.054) (Fig. 2Q-T, W, yellow arrows). Collectively, these data suggest glutamatergic neuron-autonomous *MEK1$^{S217/221E}$* expression results in significant hyperactivation of ERK1/2 in the axonal compartment of multiple forebrain projection subtypes.

*MEK1$^{S217/221E}$ expression disrupts corticocortical and corticostriatal axon arborization*

We previously reported a reduction in corticospinal tract (CST) elongation in *Nex:Cre;*

# Adult Cortex

**A**

MEK1

GAPDH

**B** MEK1 / GAPDH

**C**

ERK2

P-ERK1/2

β-actin

**D** pERK1/2 / ERK2

**DAPI / MEK1**

**E** M1 *Control*

**F** MEK1

**G** M1 *Nex:Cre; MEK1^{S217/221E}*

**H** MEK1

**DAPI / pERK1/2**

**I** L2/3 L5 L6 cc st *Control*

**J** pERK1/2

**K** L2/3 L5 L6 cc st *Nex:Cre; MEK1^{S217/221E}*

**L** pERK1/2

## M1 Layer V

**DAPI / pERK1/2 / NeuN** | **pERK1/2**

**M** *Control* **N**

**O** *Nex:Cre; MEK1^{S217/221E}* **P**

## Corpus Callosum - Airyscan

**DAPI / pERK1/2 / TUJ1**

**Q** **R**

**S** **T**

**U** pERK1/2 labeled profiles in M1 Layer 2/3

**V** pERK1/2 Levels in M1 L5 Neurons

**W** pERK1/2 colabeling with TUJ1 in Corpus Callosum

p = 0.054

**Figure 2. Increased levels of pERK1/2 in cortical axons of adult Nex:Cre; MEK1$^{S217/221E}$ mice.**

**A-D.** Western Blots of adult cortical lysates revealed a 1.61 ± 0.20 fold increase in MEK1 in *Nex:Cre; MEK1$^{S217/221E}$* mutants when compared to control mice (Mean ± SEM, n=3, Student's t-test p=0.041) (A-B). No significant difference in pERK1/2 levels between mutant and control mice was detected in adult cortical lysates (Mean ± SEM, n=3, Student's t-test p=0.389) (C-D).

**E-W.** An increase of MEK1 immunolabeling was apparent in the cortex of adult *Nex:Cre; MEK1$^{S217/221E}$* mice (G-H) relative to controls (E-F). Phosphorylated ERK1/2 immunolabeling was qualitatively increased in the corpus callosum (dotted yellow outline) in adult mutant mice (K-L, yellow arrows), when compared to controls (I-J). Within the grey matter, we noted a significant reduction in the number pERK1/2 labeled neurons in layer II/III of mutants (L, U) (Mean ± SEM, n=6, * = Student's t test p=0.002). High resolution confocal imaging and analysis of NeuN$^+$ pyramidal neurons in M1 layer V (M-P, dotted yellow outlines) revealed a decrease in cytoplasmic and nuclear levels of pERK1/2 levels (O-P, V) when compared to controls (M-N) (Mean ± SEM, n=68 control and 83 mutant neurons from four mice per group, * = Student's t-test p<0.01). We detected a modest increase in pERK1/2/TUJ1 co-localization in the corpus callosum of mutants (S-T, yellow arrows) compared to controls (Q-R, quantified in W, Mean ± SEM, n=4, Student's t test p = 0.054).

Scale bars E,L = 500µm, M-P = 10µm, Q-T = 2 µm (M1 = primary motor cortex, cc = corpus callosum, st = striatum).

*MEK1$^{S217/221E}$* mice by postnatal day 3 (P3) that persisted to P30 (Xing et al., 2016). Here, we

asked whether MEK1$^{S217/221E}$ expression is sufficient to reduce axon arborization in two other

target domains, the contralateral cortex and striatum. To unilaterally label developing neurons

and their axons, a Cre-dependent tdTomato/RFP adeno-associated viral vector (AAV-*CAG-*

*FLEX-tdTomato*) was injected into the primary motor cortex (M1) at P1 (Fig. 3A). There was no

significant difference in the density of tdTomato-labeled cells per section between *Nex:Cre;*

*MEK1$^{S217/221E}$* (224.73 ± 26.73) and control (189.96 ± 14.20) motor cortices (mean ± SEM, n=5,

Student's t-test, p=0.284) (Fig. 3B-D). To estimate axonal arborization, tdTomato-labeled pixels

in contralateral M1 and striatum were quantified. There was a statistically significant reduction of

contralateral corticocortical arborization in *Nex:Cre; MEK1$^{S217/221E}$* mice (631.68 ± 27.51)

compared to controls (791.79 ± 42.88) when normalized to number of transduced cells in M1

(Mean ± SEM, n=5, Student's t-test, p=0.014) (Fig. 3E, F, H, I, K). Contralateral corticostriatal

projections into the dorsolateral striatum were also significantly reduced in the

*Nex:Cre;MEK1$^{S217/221E}$* mice (control: 255.67 ± 23.64, mutant: 155.44 ± 21.18, Mean ± SEM,

n=5, Student's t-test, p=0.013) (Fig. 3E, G, H, J, L). Together with our previous findings, cortical

glutamatergic neuron autonomous *MEK1$^{S217/221E}$* expression appears sufficient to hyperactivate

ERK1/2 during axonal development and reduce arborization into multiple long-range targets.

*ARC protein levels are reduced by MEK1 hyperactivation*

ERK1/2 is a known regulator of glutamatergic signaling and activity dependent plasticity

(Bramham et al., 2010; Thomas and Huganir, 2004). Whether *MEK1$^{S217/221E}$* expression alters

circuit-level cortical activity in adults is unclear. Activity-Regulated Cytoskeleton-associated gene

(ARC) is an immediate early gene (IEG) triggered by glutamate receptor signaling via ERK1/2

and other pathways (Cao et al., 2015; Chotiner et al., 2010; Correa et al., 2012; Ebert and

Greenberg, 2013; Pintchovski et al., 2009; Ying et al., 2002). Our data shows axonal arborization,

somal and nuclear levels of pERK1/2, are reduced in *Nex:Cre; MEK1$^{S217/221E}$* mutants, suggesting

## AAV:*CAG-FLEX-tdTomato* P1 Injection: Collect P30

**A**

*Control*

*Nex:Cre; MEK1^(S217/221E)*

**B** *Control* — M1 — tdTomato

**C** *Nex:Cre; MEK1^(S217/221E)* — M1

**D** AAV-tdTomato Labeled Neurons

Avg labeled neurons in M1 (per section)
CTRL / MUT

## Contralateral Corticocortical and Corticostriatal Projections

*Control* | *Nex:Cre; MEK1^(S217/221E)*

tdTomato

**E** M1, st
**F** M1
**G** st
**H** M1, st
**I** M1
**J**

**K** Contralateral axonal arborization - cortex

Axonal Arborization per Labeled Neuron (pixels$^2$/cell)
CTRL / MUT
*

**L** Contralateral axonal arborization - striatum

Axonal Arborization per Labeled Neuron (pixels$^2$/cell)
CTRL / MUT
*

**Figure 3. Nex:Cre; MEK1$^{S217/221E}$ mice exhibit disrupted contralateral corticocortical and corticostriatal arborization.**

**A-D.** An adeno-associated viral vector, AAV:*CAG-Flex-tdTomato* was injected into the left primary motor cortex at P1 and brains were collected at P30 as shown in the rostral to caudal serial sections surrounding the injection site (A) and representative images of the injected region in the primary motor cortex (M1) of the *Nex:Cre; MEK1$^{S217/221E}$* (C) and control (B) forebrains. No significant difference in the average density of RFP-labeled cells was detected between *Nex:Cre; MEK1$^{S217/221E}$* and control mice (Mean ± SEM, n=5, Student's t-test p=0.2839) (D).

**E-L.** Representative confocal images of contralateral corticocortical and corticostriatal (st = striatum) RFP-labeled axonal arborization in the *Nex:Cre; MEK1$^{S217/221E}$* (H-J) and control mice (E-G). *Nex:Cre; MEK1$^{S217/221E}$* mice showed a 19.57% ± 4.30 reduction of normalized intracortical axon arborization (I) compared to control mice (Mean ± SEM, n=5, Student's t-test p=0.014) (F, quantified in K). Cortical axon arborization in the mutant dorsolateral striatum was reduced 37.50% ± 9.14 (J) compared to control mice (Mean ± SEM, n=5, Student's t-test p=0.013) (G, quantified in L).

Scale bars: B, C, E, H = 500μm, F, G, I, J= 100μm.

that glutamatergic drive may be diminished. Indeed, western blots of adult whole cortical lysates

revealed a significant 37 +/- 11% decrease in ARC expression in *Nex:Cre;MEK1$^{S217/221E}$* mice

compared to controls (n=11, Student's t-test p<0.005) (Fig. 4A-B). Immunolabeling further

indicated a widespread decrease in cortical ARC levels *in vivo* (n=4) (Fig. 4C-F). Together, these

findings support that developmental expression of hyperactive MEK1 in glutamatergic neurons

reduces cortical ARC expression *in vivo.*

*Motor skill acquisition is disrupted in Nex:Cre; MEK1$^{S217/221E}$ mice*

Common phenotypes in RASopathic individuals include hypotonia, changes in motor

coordination and altered motor learning (Blüthgen et al., 2017; Davis et al., 2000; Monje et al.,

2005). We thus asked if hyperactivation of MEK1 in *Nex:Cre* expressing cells was sufficient to

drive changes in adult mouse motor behavior in open field, accelerating rotarod, and single-pellet

retrieval assays (Crawley, 1999). Open field testing revealed no significant difference in total

distance traveled between *Nex:Cre; MEK1$^{S217/221E}$* and control mice (Fig. 5A, n=12, Student's t-

test, p=0.075). Time spent in the center of the open field arena is a gauge of anxiety-related

behavior, however, we did not observe a significant difference (Fig. 5A, n=12, Student's t-test

p=0.21). We also did not observe a significant effect of genotype during accelerating rotarod

testing, which assesses motor coordination and balance (Fig. 5B, [$F_{(1, 27)}$=0.246, p=0.624], n=15

controls & 14 mutants). These data reveal that *Nex;Cre; MEK1$^{S217/221E}$* mice have relatively normal

global voluntary locomotor behavior, motor coordination, and balance.

We next utilized a single-pellet retrieval task to determine if there was a deficit in fine

forelimb motor function and learning, which relies heavily on the motor cortex (Chen et al., 2014;

Kawai et al., 2015; Li and Hollis, 2021). This task assays the ability of mice to reach through a

narrow slot with a single forelimb to retrieve a sucrose pellet. We measured the percentage of

successful retrievals over the total number of reaching attempts across 5 consecutive days of

testing. Importantly, we found that *Nex:Cre; MEK1$^{S217/221E}$* mutants exhibit a significant decrease

# Adult Cortex

**Figure 4. MEK1$^{S217/221E}$ reduces cortical ARC protein expression.**

**A-B.** Western Blots of adult cortical lysates showed a significant reduction in ARC in *Nex:Cre;*

*MEK1$^{S217/221E}$* mutants when compared to control mice (Mean ± SEM, n=11, Student's t-test,

p<0.005).

**C-F.** Immunolabeling revealed a qualitative reduction of ARC in *Nex:Cre; MEK1$^{S217/221E}$* cortices

(E-F) when compared to controls (C-D, n=5).

Scale bars = 200um.

in success rate relative to controls (Fig. 5C, main effect of genotype [$F_{(1,16)}$=11.04, p=0.004], n=11

controls & 8 mutants). The success rate in mutant mice was significantly decreased relative to

controls during the last three days of testing ( * = LSD post-hoc p < 0.05). However, no significant

difference in success rate was noted on the first day of testing (LSD post-hoc p=0.59) or in the

number of attempts [$F_{(1,16)}$=0.06, p=0.81]. Collectively, these results indicate that the *Nex:Cre;*

*MEK1$^{S217/221E}$* mice do not exhibit initial, basal differences in fine forelimb motor function, but

display a significant impairment in learning a skilled forelimb reaching and grasping task.

*Mouse model for studying layer V autonomous effects of ERK1/2 signaling*

Since *Nex:Cre* recombines in all cortical layers, it is unclear whether axonal deficits are

due to direct layer V specific effects or changes in upstream layers in the cortical circuit. To

determine the layer V autonomous functions of ERK1/2 signaling, we employed a *Rbp4:Cre*

mouse that drives recombination primarily in cortical layer V neurons in the cortex (Gong et al.,

2007; Kozorovitskiy et al., 2012) and outside of the nervous system in the liver (Steinhoff et al.,

2021; Thompson et al., 2017). Inducible deletion of *Erk1/2* in adult mice has been shown to trigger

hepatocyte death and increased mortality, while ERK1/2 hyperactivation induces hepatocyte

proliferation and carcinoma (Cingolani et al., 2022; Huynh et al., 2003; Ito et al., 1998). Attempts

to generate *Rbp4:Cre; Erk1$^{-/-}$; Erk2$^{fl/fl}$* loss-of-function mice did not result in viable births (0 of 85,

expected 1:4 births). However, *Rbp4:Cre; MEK1$^{S217/221E}$; Ai9$^{+/-}$* mice are viable into adulthood, but

exhibit hepatomegaly (Fig. S2M). Nonetheless, *Rbp4:Cre; MEK1$^{S217/221E}$; Ai9$^{+/-}$* mice express RFP

and elevated levels of MEK1 in cortical layer V somas and axon projections and provide a model

for evaluating layer V-autonomous effects (Fig. S2D-F, J-L).

*Layer V directed expression of MEK1$^{S217/221E}$ reduces cortical axon arborization*

We did not observe a decrease in RFP$^+$ cortical layer V neuron density in M1 of *Rbp4:Cre;*

*MEK1$^{S217/221E}$; Ai9$^{+/-}$* mutants relative to controls (Control = 372 ± 9, Mutant = 477 ± 38, Mean ±

# Locomotor and Motor Skill Assays - *Nex:Cre; MEK1*$^{S217/221E}$

**Figure 5. Motor skill acquisition is disrupted by MEK1 hyperactivation in cortical**

**glutamatergic neurons.**

**A.** Open field testing of *Nex:Cre; MEK1$^{S217/221E}$* (n=12) and control (n=12) mice revealed no

significant difference in distance traveled (Mean ± SEM, Student's t-test p=0.075) or time spent

in the center of the arena (Mean ± SEM, Student's t-test p=0.21) (A).

**B.** Accelerating rotarod testing revealed a significant effect of day [$F_{(4, 27)}$=6.144, p<0.001] over 5

days of testing, but no significant effect of genotype [$F_{(1, 27)}$=0.246, p=0.624] (Mean ± SEM, n=15

controls & 14 mutants) (B).

**C-D.** In the single-pellet retrieval task, the control success rate significantly improved over five

days from 12.39 ± 4.50 to 27.06 ± 5.11 percent success rate (Mean ± SEM, Students t-test D1 vs

D5 p=0.04, n=11) (C-D). The *Nex:Cre; MEK1$^{S217/221E}$* mutant mice exhibited a significantly lower

success rate relative to controls during the final three days of testing [$F_{(1,16)}$=11.04, p=0.004] (C-

D, Mean ± SEM, n=11 controls & 8 mutants, * = LSD post-hoc p < 0.05).

SEM, Student's t-test p=0.054), consistent with past findings assessing the proportion of layer V

CTIP2[+] neurons in *Nex:Cre; MEK1[S217/221E]* mice (Xing 2016) (Fig. S2B,E). We next asked if

hyperactive MEK1 expression in layer V is sufficient to reduce axon extension. All layer V derived

subcortical projections in the spinal cord dorsal funiculus were labeled by the Cre-dependent

tdTomato/RFP reporter, *Ai9[+/-]*, in *Rbp4:Cre* mice. We observed that *Rbp4:Cre; MEK1[S217/221E];*

*Ai9[+/-]* mice had a 28 +/- 8% reduction in RFP[+] pixel density in the lumbar dorsal funiculus

compared to controls (Fig. 6A-G) (Mean ± SEM, n=5, Student's t-test p < 0.05).

The widespread bilateral labeling of all cortical layer V neurons in *Rbp4:Cre; Ai9[+/-]* mice

made it challenging to determine the anatomical source of axonal labeling. Therefore, we utilized

focal neonatal injection of a Cre-dependent tdTomato/RFP AAV into M1 to unilaterally label

neurons at birth. Comparable numbers of neurons were labeled in M1 in control and *Rbp4:Cre;*

*MEK1[S217/221E]* mice at P30 (Mean ± SEM, n=5, Student's t-test p=0.400) (Fig. S3C). Although there

was no difference in RFP labeling in corticobulbar tract (CBT), mutants had reduced RFP labeling

in the cervical dorsal funiculus compared to controls, consistent with decreased CST elongation

(Fig. 7A-E, Fig. S3D-H). Intracortical arborization in contralateral M1 of *Rbp4:Cre; MEK1[S217/221E]*

mice was also significantly reduced when compared to control mice (Mean ± SEM, n=5, Student's

t-test p=0.019) (Fig. 7F-G, J-K, quantification in N). Contralateral dorsolateral striatal arborization

was also significantly reduced in *Rbp4:Cre;MEK1[S217/221E]* mice when compared to controls (Mean

± SEM, n=5, Student's t-test p=0.007) (Fig. 7H-I, L-M, quantification in O). Collectively, these

results indicate that hyperactive ERK1/2 signaling in cortical glutamatergic layer V neurons is

sufficient to reduce axonal elongation of multiple projection neurons.

*Reduction in cortical wiring of layer V is expendable for skilled motor learning*

Since *Rbp4:Cre;MEK1[S217/221E]* mice exhibit reduced axonal extension of cortical projection

neurons, we tested whether these mutants display deficits in motor behavior. Hepatomegaly was

observed in *Rbp4:Cre;MEK1[S217/221E]* mice and mouse models of chronic liver disease have been

shown to exhibit hypolocomotion in the open-field, but little change in motor learning or anxiety-

## P60: Lumbar Spinal Cord Cre dependent tdTomato Reporter

**G**

**Corticospinal Labeling**

**Figure 6. MEK1$^{S214/221E}$ expression in cortical layer V reduces corticospinal axon elongation in the spinal cord dorsal fasciculus.**

**A-G.** *RBP4:Cre; Ai9$^{+/-}$* control (A) and *RBP4:Cre; MEK1$^{S17/221E}$; Ai9$^{+/-}$* mutant (D) mice label descending cortical layer V derived corticospinal axons (dotted yellow box). Cross-sectional lumbar segments reveal a statistically significant 28% reduction of RFP$^+$ labeled pixels in the dorsal funiculus of *RBP4:Cre; MEK1$^{S217/221E}$; Ai9$^{+/-}$* mice (E-F, dotted yellow outline) compared to *RBP4:Cre; Ai9$^{+/-}$* controls (B-C, quantification in G, mean ± SEM, n=5, * = Student's t-test p < 0.05).

Scale bars A, D = 200um, B-C, E-F = 20um (dh = dorsal horn, cc = central canal, vh = ventral horn, df = dorsal funiculus).

**AAV:*CAG-FLEX-tdTomato* P1 Injection: Collect P30**

**Layer 5 Contralateral Corticocortical and Corticostriatal Projections**

**Figure 7. Corticospinal, corticocortical and corticostriatal layer V projections are reduced in *RBP4:Cre; MEK1^S217/221E* mutants.**

**A-E.** Unilateral AAV labeling of descending corticospinal axons derived from M1 in RBP4:Cre mice. Cross sections of cervical spinal cord segments reveal a reduction of descending RFP$^+$ corticospinal axons in the dorsal funiculus of *RBP4Cre; MEK1^S217/221E* mutants (D) compared to *RBP4:Cre* control mice (B) when normalized to corticobulbar tract labeling (E, Mean ± SEM, n=3, Student's t-test p=0.036).

**F-O.** Virally-labeled *RBP4Cre; MEK1^S217/221E* axons had significantly reduced arborization in the contralateral motor cortex (K) compared to *RBP4:Cre* controls (G) when normalized to the number of transduced cells in M1 (quantification in N, Mean ± SEM, n=5, * = Student's t-test p=0.019). Additionally, *RBP4Cre; MEK1^S217/221E* mice had significantly reduced arborization of the dorsolateral striatum (M) compared to *RBP4:Cre* controls (I) (quantification in O, Mean ± SEM, n=5, * Student's t-test p=0.007).

Scale bars: A,C, =200µm, B,D,= 20µm, F,H,J,L = 500µm, G,I,K,M = 150µm (dh = dorsal horn, df = dorsal funiculus, iz = intermediate zone, vh = ventral horn, M1 = primary motor cortex, st = striatum).

like phenotypes (Jover et al., 2006). When compared to control mice, *Rbp4:Cre;MEK1$^{S217/221E}$*

mutants showed a significant decrease in voluntary distance traveled in the open-field assay

(n=22 controls and 11 mutants, Student's t-test p=0.015) and a decrease in the proportion of time

spent in the center (Student's t-test p=0.001) (Fig. 8A). In contrast, no significant effect of

genotype was detected on the accelerating rotarod task ([$F_{(1,36)}$=0.007, p=0.933], n=26 controls &

12 mutants) (Fig. 8B).

In the single-pellet retrieval task, we found that mild food restriction in *Rbp4:Cre;*

*MEK1$^{S217/221E}$* mice resulted in health issues of unknown origin and spontaneous lethality. Thus,

we substantially reduced the period of food restriction relative to our *Nex:Cre* experiments and

only analyzed reaching and grasping behaviors during the first five minutes of each trial when

mice were highly motivated. In stark contrast to *Nex:Cre; MEK1$^{S217/221E}$* mutants, we found no

significant effect of genotype on the success rate between control and *Rbp4:Cre;MEK1$^{S217/221E}$*

mice (Fig. 8C-D, ([$F_{(1,24)}$=0.03, p=0.88], n=14 controls & 12 mutants). Taken together, these

results indicate that layer V specific hyperactivation of MEK1 in glutamatergic neurons is

insufficient to disrupt motor skilled learning.

## Locomotor and Motor Skill Assays - *RBP4:Cre; MEK1^{S217/221E}*

**Figure 8. MEK1$^{S217/221E}$ in cortical layer V glutamatergic neurons does not disrupt motor learning.**

**A.** Open field testing of RBP4*:Cre; MEK1$^{S217/221E}$* and control mice revealed that mutants exhibit a significant reduction in total distance traveled (Mean ± SEM, n= 22 controls & 11 mutants, Student's t-test p=0.01) and time spent in center of box (Mean ± SEM, n= 22 controls & 11 mutants, Student's t-test p=0.001).

**B.** Accelerating rotarod testing revealed no significant effect of genotype over 5 days of testing control and mutant mice ([$F_{(1,36)}$=0.007, p=0.933] (Mean ± SEM , n=26 controls & 12 mutants).

**C-D.** In the single-pellet retrieval task, there was no significant effect of genotype on daily success rate [$F_{(1,24)}$=0.03, p=0.88] (C, Mean ± SEM, n=14 controls & 12 mutants). Both mutants and controls improved their success rate over five days [$F_{(4,24)}$=5.81, p<0.01] (D, Mean ± SEM, n = 14 controls & 12 mutants).

**Discussion**

Resolving the cell-specific effects of pathological ERK1/2 activity may help inform the rational design of therapeutic approaches for multiple neurodevelopmental conditions. Here, we tested whether cortical excitatory neuron-directed expression of hyperactive MEK1$^{S217/221E}$ during early development modulates long-range axon outgrowth from sensorimotor cortices, expression of ARC, and motor-related behaviors. MEK1 hyperactivation selectively mimics a core downstream feature of multiple RASopathies without direct modulation of multiple RTK-linked signaling cascades associated with upstream mutations. We generated mice expressing Cre-dependent MEK1$^{S217/221E}$ and *Nex:Cre* to target cortical excitatory neurons, but not cortical GABAergic neurons or glia. Axonal levels of pERK1/2 are persistently increased in these mutants. Importantly, ERK1/2 hyperactivation coincides with reductions in corticocortical, corticostriatal, and corticospinal axon projections arising from motor cortices. We also observe lower expression of the activity-dependent gene, Arc, within the mature cortex of mutant mice. While many basic motor behaviors remain intact, skilled motor learning in a reaching task is inhibited in these mutants. Lastly, we demonstrate that a layer V-specific MEK1$^{S217/221E}$ mutant mouse recapitulates axonal outgrowth deficits, but motor learning is not altered. Overall, our data show that layer V autonomous expression of MEK1$^{S217/221E}$ disrupts the development of long-range cortical axon projections, while MEK1 hyperactivation in developing *Nex:Cre*-expressing neurons appears necessary for diminished skilled motor learning (Fig. S4).

**Compartment-specific ERK1/2 activation in *MEK1$^{S217/221E}$*-expressing neurons**

*In vitro* studies have been vital for quantitative studies of kinase signaling and pharmacological inhibitors, but the precise effect of RASopathy mutations on pERK1/2 levels and distribution are more challenging to measure *in vivo*. We generated mutants with increased hyperactive MEK1 levels in developing excitatory neurons and observed increased levels of pERK1/2 in axonal, but not somal and nuclear compartments. Recent studies have shown that callosal layer II/III projections have enriched ERK2/MAPK1 protein within axonal growth cones

(Engmann et al., 2022; Poulopoulos et al., 2019; Shigeoka et al., 2016). It will be important to

explore whether enrichment of ERK1/2 pathway components within the axon renders this

compartment relatively vulnerable to RASopathic mutations. High resolution measurement of

RASopathic kinase activation in distinct subcellular domains *in vivo* may further illuminate core

pathological mechanisms.

Paradoxically, the somal and nuclear levels of pERK1/2 were reduced in adult mutant

excitatory neurons. A similar phenomenon has been observed in Drosophila MEK1 mutants,

possibly due to early changes in the expression of morphogens during development (Goyal et al.,

2017). The compensatory mechanisms that appear to override the biochemical hyperactivity of

the MEK1$^{S217/221E}$ mutant in neuronal soma and nuclei may involve ERK1/2-directed

phosphatases enriched within somal compartments or developmental changes in upstream

ERK1/2 activators (Caunt et al., 2008; Jeffrey et al., 2007; Li et al., 2007; Lim et al., 2015; Parmar

et al., 2015; Poulopoulos et al., 2019). Nevertheless, pharmacological inhibitors of the ERK1/2

signaling cascade are actively being explored for clinical use in RASopathies (Guilding et al.,

2007; Hatzivassiliou et al., 2010; Morris et al., 2013; Nakamura et al., 2009; Zamorano et al.,

2018). For example, MEK1/2 inhibitors during postnatal stages reverses cardiac deficits and

tumorigenesis in RASopathy models and patients with Noonan Syndrome (Andelfinger et al.,

2019; Hilal et al., 2023; Wu et al., 2011). However, if developmental compensatory mechanisms

lead to paradoxically reduced levels of pERK1/2 in select neuronal compartments, it is unclear

whether pharmacological inhibition would be beneficial. Additional quantitative analyses of

compartment-specific changes in pERK1/2 levels in RASopathy models may help inform the use

of systemic pharmacological inhibitors.

**MEK-ERK1/2 hyperactivity reduces cortical long-range axon arborization**

MRI-based studies of individuals with NF1 and Noonan Syndrome have discovered

abnormalities in both functional and structural connectivity (Fattah et al., 2021; Filippi et al., 2013;

Rai et al., 2023). While oligodendrocyte mediated changes in the myelination of axons are likely

involved, the neuron-autonomous contributions to aberrant connectivity are less understood. We

show that neuron-specific MEK1[S217/221E] expressing mice did not exhibit significant loss of cortical

layer V neurons, but layer V-autonomous hyperactivation of ERK1/2 reduces the arborization of

corticocortical, corticostriatal, and corticospinal axons. *In vitro* studies have demonstrated ERK1/2

is necessary for aspects of trophic factor-induced axon outgrowth (Markus et al., 2002; Ozdinler

and Macklis, 2006), axon guidance (Bagnard et al., 2004; Forcet et al., 2002; Yun et al., 2005),

cytoskeletal organization (Atwal et al., 2003; Goold and Gordon-Weeks, 2005; Ray and Sturgill,

1987) and local protein translation in growth cones (Campbell and Holt, 2003). Whether

hyperactive ERK1/2 activity disrupts these molecular processes or modulates other intracellular

events associated with axon outgrowth *in vivo* is unclear. Axonal outgrowth in humans occurs

during gestational stages, which makes therapeutic intervention more difficult. RASopathies are

often diagnosed within the first years of life (Zenker et al., 2022). Therefore, it may be beneficial

to study postnatal stages of axonal refinement and pruning in RASopathies, which may be more

accessible to current treatment strategies.

Animal models of upstream RASopathic mutations in NF1, Ptpn11, and Ras have

observed changes in multiple RTK-linked downstream intracellular cascades, including PI3K/Akt,

JAK/STAT, PKC, mTOR and ERK1/2, and inhibitors of multiple cascades are capable of reversing

certain cellular deficits (Anastasaki and Gutmann, 2014; Brown et al., 2012; Langdon et al., 2012;

López-Juárez et al., 2017; Titus et al., 2017). Our work now shows that downstream MEK-ERK1/2

hyperactivity is sufficient to reduce long range axon arborization *in vivo*. Understanding whether

upstream mutations also drive neuron-autonomous deficits in long-range excitatory neuron

axonal outgrowth and may help inform the design of personalized, mutation-specific approaches

to RASopathy therapeutics.

**Nex:Cre; MEK1[S217/221E] mice exhibit decreased Arc and deficits in skilled motor learning**

Activity-regulated cytoskeletal gene *Arc* is an immediate early gene that rapidly responds

to neuronal activity and regulates synaptic plasticity (Steward and Worley, 2001; Steward et al.,

1998). *Arc* has been shown to be dysregulated in Fragile X and Angelman Syndrome (Greer et

al., 2010; Zalfa et al., 2003), however, whether RASopathic individuals exhibit aberrant *Arc*

expression is not clear. Several reports show that glutamate and BDNF drive Arc transcription

and translation via ERK1/2 (Chotiner et al., 2010; Correa et al., 2012; Nikolaienko et al., 2017;

Pintchovski et al., 2009; Rao et al., 2006; Ying et al., 2002). Notably, our results indicate that

MEK1 hyperactivation during early neuronal development unpredictably resulted in reduced

phosphorylated ERK1/2 levels in the soma and nucleus of adult neurons. It is perhaps not

surprising that ARC protein is also decreased in adult mutant cortices, in agreement with our

past report showing P21 *Nex:Cre; MEK1$^{S217/221E}$* mice have reduced levels of *Arc* mRNA (Xing

et al., 2016). Hippocampal excitatory neurons in a developing forebrain-directed RASopathy

mutant, *Emx1:Cre Ptpn11$^{D61Y}$*, exhibit analogous reductions in depolarization-induced pERK1/2

and BDNF promoter activity (Altmüller et al., 2017). We propose the reduced levels of pERK1/2-

ARC in cortical neuron soma and nuclei may be due to decreases in the development of cortical

glutamatergic synapses or excitatory activity as a consequence of diminished long-range axon

outgrowth.

In line with the reduced expression of *Arc*, our data show that *MEK1$^{S217/221E}$* expression in

*Nex:Cre* mice is sufficient to disrupt skilled motor learning. *Nex/Neurod6* predominantly drives

Cre expression in immature cortical excitatory neurons, however, we cannot exclude that other

*Nex/Neurod6* expressing neuronal populations could contribute to this behavioral phenotype

(Goebbels et al., 2006). Nonetheless, restricting MEK1$^{S217/221E}$ to layer V with *Rbp4:Cre* does not

result in decreased skilled motor learning even though we observed a decrease in layer V neuron

axon outgrowth. Motor learning is known to involve a complex, distributed set of neuronal

populations that are not targeted in *Rbp4:Cre* mice. For example, the activity of layer II/III neurons

in the motor cortex are recruited in a skilled reaching task and contain a sparse population of

pyramidal neurons with some of the highest levels of pERK1/2 in the normal adult mouse cortex

(Levy et al., 2020). We found a reduction in the number of layer II/III neurons with high levels of

pERK1/2 in the motor cortex of *Nex:Cre; MEK1$^{S217/221E}$* mice. Additional research is needed to

explore the normal functions of these high pERK1/2-expressing layer II/III neurons in cortical

processes and whether they are particularly vulnerable to RASopathic signaling.

***Funding***

This work was supported by the National Institutes of Health [K99/R00 NS076661 and R01

NS097537 to JMN].

***Acknowledgments***

We wish to thank Noah Fry, Becca Reinking-Herd, and Sarah Bjorklund, for their technical

assistance.

## References

Alessi, D. R., Saito, Y., Campbell, D. G., Cohen, P., Sithanandam, G., Rapp, U., Ashworth, A., Marshall, C. J. and Cowley, S. (1994). Identification of the sites in MAP kinase kinase-1 phosphorylated by p74raf-1. *The EMBO Journal* 13, 1610–1619.

Altmüller, F., Pothula, S., Annamneedi, A., Nakhaei-Rad, S., Montenegro-Venegas, C., Pina-Fernández, E., Marini, C., Santos, M., Schanze, D., Montag, D., et al. (2017). Aberrant neuronal activity-induced signaling and gene expression in a mouse model of RASopathy. *PLoS Genet* 13, e1006684.

Anastasaki, C. and Gutmann, D. H. (2014). Neuronal NF1/RAS regulation of cyclic AMP requires atypical PKC activation. *Human Molecular Genetics* 23, 6712–6721.

Andelfinger, G., Marquis, C., Raboisson, M.-J., Théoret, Y., Waldmüller, S., Wiegand, G., Gelb, B. D., Zenker, M., Delrue, M.-A. and Hofbeck, M. (2019). Hypertrophic Cardiomyopathy in Noonan Syndrome Treated by MEK-Inhibition. *J Am Coll Cardiol* 73, 2237–2239.

Angara, K., Pai, E. L.-L., Bilinovich, S. M., Stafford, A. M., Nguyen, J. T., Li, K. X., Paul, A., Rubenstein, J. L. and Vogt, D. (2020). *Nf1* deletion results in depletion of the *Lhx6* transcription factor and a specific loss of parvalbumin [+] cortical interneurons. *Proc. Natl. Acad. Sci. U.S.A.* 117, 6189–6195.

Atkins, C. M., Yon, M., Groome, N. P. and Sweatt, J. D. (2001). Regulation of Myelin Basic Protein Phosphorylation by Mitogen-Activated Protein Kinase During Increased Action Potential Firing in the Hippocampus. *Journal of Neurochemistry* 73, 1090–1097.

Atwal, J. K., Singh, K. K., Tessier-Lavigne, M., Miller, F. D. and Kaplan, D. R. (2003). Semaphorin 3F Antagonizes Neurotrophin-Induced Phosphatidylinositol 3-Kinase and Mitogen-Activated Protein Kinase Kinase Signaling: A Mechanism for Growth Cone Collapse. *The Journal of Neuroscience* 23, 7602–7609.

Aydin, S., Kurtcan, S., Alkan, A., Guler, S., Filiz, M., Yilmaz, T. F., Sahin, T. U. and Aralasmak, A. (2016). Relationship between the corpus callosum and neurocognitive disabilities in children with NF-1: diffusion tensor imaging features. *Clinical Imaging* 40, 1092–1095.

Bagnard, D., Sainturet, N., Meyronet, D., Perraut, M., Miehe, M., Roussel, G., Aunis, D., Belin, M. F. and Thomasset, N. (2004). Differential MAP kinases activation during semaphorin3A-induced repulsion or apoptosis of neural progenitor cells. *Mol Cell Neurosci* 25, 722–731.

Bernardino, I., Dionísio, A. and Castelo-Branco, M. (2022). Cortical inhibition in neurofibromatosis type 1 is modulated by lovastatin, as demonstrated by a randomized, triple-blind, placebo-controlled clinical trial. *Sci Rep* 12, 13814.

Blüthgen, N., van Bentum, M., Merz, B., Kuhl, D. and Hermey, G. (2017). Profiling the MAPK/ERK dependent and independent activity regulated transcriptional programs in the murine hippocampus in vivo. *Sci Rep* 7, 45101.

Bramham, C. R., Alme, M. N., Bittins, M., Kuipers, S. D., Nair, R. R., Pai, B., Panja, D., Schubert, M., Soule, J., Tiron, A., et al. (2010). The Arc of synaptic memory. *Exp Brain Res* 200, 125–140.

Brown, J. A., Diggs-Andrews, K. A., Gianino, S. M. and Gutmann, D. H. (2012). Neurofibromatosis-1 heterozygosity impairs CNS neuronal morphology in a cAMP/PKA/ROCK-dependent manner. *Molecular and Cellular Neuroscience* 49, 13–22.

Bueno, O. F. (2000). The MEK1-ERK1/2 signaling pathway promotes compensated cardiac hypertrophy in transgenic mice. *The EMBO Journal* 19, 6341–6350.

Campbell, D. S. and Holt, C. E. (2003). Apoptotic Pathway and MAPKs Differentially Regulate Chemotropic Responses of Retinal Growth Cones. *Neuron* 37, 939–952.

Cao, V. Y., Ye, Y., Mastwal, S., Ren, M., Coon, M., Liu, Q., Costa, R. M. and Wang, K. H. (2015). Motor Learning Consolidates Arc-Expressing Neuronal Ensembles in Secondary Motor Cortex. *Neuron* 86, 1385–1392.

Castellano, E. and Downward, J. (2011). RAS Interaction with PI3K: More Than Just Another Effector Pathway. *Genes Cancer* 2, 261–274.

Caunt, C. J., Armstrong, S. P., Rivers, C. A., Norman, M. R. and McArdle, C. A. (2008). Spatiotemporal Regulation of ERK2 by Dual Specificity Phosphatases. *Journal of Biological Chemistry* 283, 26612–26623.

Chang, Y. S., Owen, J. P., Pojman, N. J., Thieu, T., Bukshpun, P., Wakahiro, M. L. J., Marco, E. J., Berman, J. I., Spiro, J. E., Chung, W. K., et al. (2016). Reciprocal white matter alterations due to 16p11.2 chromosomal deletions versus duplications: White Matter Microstructure in 16p11.2 CNVs. *Hum. Brain Mapp.* 37, 2833–2848.

Chen, C.-C., Gilmore, A. and Zuo, Y. (2014). Study Motor Skill Learning by Single-pellet Reaching Tasks in Mice. *JoVE* 51238.

Chotiner, J. K., Nielson, J., Farris, S., Lewandowski, G., Huang, F., Banos, K., de Leon, R. and Steward, O. (2010). Assessment of the role of MAP kinase in mediating activity-dependent transcriptional activation of the immediate early gene *Arc/Arg3.1* in the dentate gyrus in vivo. *Learn. Mem.* 17, 117–129.

Chung, W. K., Roberts, T. P., Sherr, E. H., Snyder, L. G. and Spiro, J. E. (2021). 16p11.2 deletion syndrome. *Current Opinion in Genetics & Development* 68, 49–56.

Cingolani, F., Liu, Y., Shen, Y., Wen, J., Farris, A. B. and Czaja, M. J. (2022). Redundant Functions of ERK1 and ERK2 Maintain Mouse Liver Homeostasis Through Down-Regulation of Bile Acid Synthesis. *Hepatology Communications* 6, 980–994.

Clement, J. P., Aceti, M., Creson, T. K., Ozkan, E. D., Shi, Y., Reish, N. J., Almonte, A. G., Miller, B. H., Wiltgen, B. J., Miller, C. A., et al. (2012). Pathogenic SYNGAP1 mutations impair cognitive development by disrupting maturation of dendritic spine synapses. *Cell* 151, 709–723.

Correa, S. A. L., Hunter, C. J., Palygin, O., Wauters, S. C., Martin, K. J., McKenzie, C., McKelvey, K., Morris, R. G. M., Pankratov, Y., Arthur, J. S. C., et al. (2012). MSK1 Regulates Homeostatic and Experience-Dependent Synaptic Plasticity. *Journal of Neuroscience* 32, 13039–13051.

Cowley, S. (1994). Activation of MAP kinase kinase is necessary and sufficient for PC12 differentiation and for transformation of NIH 3T3 cells. *Cell* 77, 841–852.

Crawley, J. N. (1999). Behavioral phenotyping of transgenic and knockout mice: experimental design and evaluation of general health, sensory functions, motor abilities, and specific behavioral tests1Published on the World Wide Web on 2 December 1998.1. *Brain Research* 835, 18–26.

Cui, Y., Costa, R. M., Murphy, G. G., Elgersma, Y., Zhu, Y., Gutmann, D. H., Parada, L. F., Mody, I. and Silva, A. J. (2008). Neurofibromin regulation of ERK signaling modulates GABA release and learning. *Cell* 135, 549–560.

Davis, S., Vanhoutte, P., Pages, C., Caboche, J. and Laroche, S. (2000). The MAPK/ERK cascade targets both Elk-1 and cAMP response element-binding protein to control long-term potentiation-dependent gene expression in the dentate gyrus in vivo. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 20, 4563–4572.

Doherty, A. C., Huddleston, D. A., Horn, P. S., Ratner, N., Simpson, B. N., Schorry, E. K., Aschbacher-Smith, L., Prada, C. E. and Gilbert, D. L. (2023). Motor Function and Physiology in Youth With Neurofibromatosis Type 1. *Pediatr Neurol* 143, 34–43.

Ebert, D. H. and Greenberg, M. E. (2013). Activity-dependent neuronal signalling and autism spectrum disorder. *Nature* 493, 327–337.

Engmann, A. K., Hatch, J. J., Nanda, P., Veeraraghavan, P., Ozkan, A., Poulopoulos, A., Murphy, A. J. and Macklis, J. D. (2022). Neuronal subtype-specific growth cone and soma purification from mammalian CNS via fractionation and fluorescent sorting for subcellular analyses and spatial mapping of local transcriptomes and proteomes. *Nat Protoc* 17, 222–251.

Fattah, M., Raman, M. M., Reiss, A. L. and Green, T. (2021). PTPN11 Mutations in the Ras-MAPK Signaling Pathway Affect Human White Matter Microstructure. *Cerebral Cortex* 31, 1489–1499.

Filippi, C. G., Watts, R., Duy, L. A. N. and Cauley, K. A. (2013). Diffusion-Tensor Imaging

Derived Metrics of the Corpus Callosum in Children With Neurofibromatosis Type I.

*American Journal of Roentgenology* 200, 44–49.

Forcet, C., Stein, E., Pays, L., Corset, V., Llambi, F., Tessier-Lavigne, M. and Mehlen, P.

(2002). Netrin-1-mediated axon outgrowth requires deleted in colorectal cancer-

dependent MAPK activation. *Nature* 417, 443–447.

Goebbels, S., Bormuth, I., Bode, U., Hermanson, O., Schwab, M. H. and Nave, K.-A. (2006).

Genetic targeting of principal neurons in neocortex and hippocampus of NEX-Cre mice.

*genesis* 44, 611–621.

Gómez-Palacio-Schjetnan, A. and Escobar, M. L. (2008). In vivo BDNF modulation of adult

functional and morphological synaptic plasticity at hippocampal mossy fibers.

*Neuroscience Letters* 445, 62–67.

Gong, S., Doughty, M., Harbaugh, C. R., Cummins, A., Hatten, M. E., Heintz, N. and Gerfen, C.

R. (2007). Targeting Cre Recombinase to Specific Neuron Populations with Bacterial

Artificial Chromosome Constructs. *Journal of Neuroscience* 27, 9817–9823.

Goold, R. G. and Gordon-Weeks, P. R. (2005). The MAP kinase pathway is upstream of the

activation of GSK3β that enables it to phosphorylate MAP1B and contributes to the

stimulation of axon growth. *Molecular and Cellular Neuroscience* 28, 524–534.

Goyal, Y., Jindal, G. A., Pelliccia, J. L., Yamaya, K., Yeung, E., Futran, A. S., Burdine, R. D.,

Schüpbach, T. and Shvartsman, S. Y. (2017). Divergent effects of intrinsically active

MEK variants on developmental Ras signaling. *Nat Genet* 49, 465–469.

Greer, P. L., Hanayama, R., Bloodgood, B. L., Mardinly, A. R., Lipton, D. M., Flavell, S. W., Kim,

T.-K., Griffith, E. C., Waldon, Z., Maehr, R., et al. (2010). The Angelman Syndrome

Protein Ube3A Regulates Synapse Development by Ubiquitinating Arc. *Cell* 140, 704–

716.

Guilding, C., McNair, K., Stone, T. W. and Morris, B. J. (2007). Restored plasticity in a mouse

model of neurofibromatosis type 1 via inhibition of hyperactive ERK and CREB:

Restored plasticity in NF1 via ERK inhibition. *European Journal of Neuroscience* 25, 99–105.

Guo, J.-Z., Graves, A. R., Guo, W. W., Zheng, J., Lee, A., Rodríguez-González, J., Li, N., Macklin, J. J., Phillips, J. W., Mensh, B. D., et al. (2015). Cortex commands the performance of skilled movement. *eLife* 4, e10774.

Gutmann, D. H., Parada, L. F., Silva, A. J. and Ratner, N. (2012). Neurofibromatosis Type 1: Modeling CNS Dysfunction. *J. Neurosci.* 32, 14087–14093.

Hart, G. and Balleine, B. W. (2016). Consolidation of Goal-Directed Action Depends on MAPK/ERK Signaling in Rodent Prelimbic Cortex. *J. Neurosci.* 36, 11974–11986.

Hatzivassiliou, G., Song, K., Yen, I., Brandhuber, B. J., Anderson, D. J., Alvarado, R., Ludlam, M. J. C., Stokoe, D., Gloor, S. L., Vigers, G., et al. (2010). RAF inhibitors prime wild-type RAF to activate the MAPK pathway and enhance growth. *Nature* 464, 431–435.

Hilal, N., Chen, Z., Chen, M. H. and Choudhury, S. (2023). RASopathies and cardiac manifestations. *Front Cardiovasc Med* 10, 1176828.

Holter, M. C., Hewitt, L. T., Nishimura, K. J., Knowles, S. J., Bjorklund, G. R., Shah, S., Fry, N. R., Rees, K. P., Gupta, T. A., Daniels, C. W., et al. (2021). Hyperactive MEK1 Signaling in Cortical GABAergic Neurons Promotes Embryonic Parvalbumin Neuron Loss and Defects in Behavioral Inhibition. *Cerebral Cortex* 31, 3064–3081.

Huynh, H., Nguyen, T. T. T., Chow, K.-H. K.-P., Tan, P. H., Soo, K. C. and Tran, E. (2003). Over-expression of the mitogen-activated protein kinase (MAPK) kinase (MEK)-MAPK in hepatocellular carcinoma: Its role in tumor progression and apoptosis. *BMC Gastroenterol* 3, 19.

Ibrahim, A. F. A., Montojo, C. A., Haut, K. M., Karlsgodt, K. H., Hansen, L., Congdon, E., Rosser, T., Bilder, R. M., Silva, A. J. and Bearden, C. E. (2017). Spatial working memory in neurofibromatosis 1: Altered neural activity and functional connectivity. *NeuroImage: Clinical* 15, 801–811.

Ishii, A., Fyffe-Maricich, S. L., Furusho, M., Miller, R. H. and Bansal, R. (2012). ERK1/ERK2 MAPK Signaling is Required to Increase Myelin Thickness Independent of Oligodendrocyte Differentiation and Initiation of Myelination. *Journal of Neuroscience* 32, 8855–8864.

Ishii, A., Furusho, M. and Bansal, R. (2013). Sustained Activation of ERK1/2 MAPK in Oligodendrocytes and Schwann Cells Enhances Myelin Growth and Stimulates Oligodendrocyte Progenitor Expansion. *J. Neurosci.* 33, 175–186.

Ito, Y., Sasaki, Y., Horimoto, M., Wada, S., Tanaka, Y., Kasahara, A., Ueki, T., Hirano, T., Yamamoto, H., Fujimoto, J., et al. (1998). Activation of mitogen-activated protein kinases/extracellular signal-regulated kinases in human hepatocellular carcinoma. *Hepatology* 27, 951–958.

Jeanneteau, F., Deinhardt, K., Miyoshi, G., Bennett, A. M. and Chao, M. V. (2010). The MAP kinase phosphatase MKP-1 regulates BDNF-induced axon branching. *Nat Neurosci* 13, 1373–1379.

Jeffrey, K. L., Camps, M., Rommel, C. and Mackay, C. R. (2007). Targeting dual-specificity phosphatases: manipulating MAP kinase signalling and immune responses. *Nat Rev Drug Discov* 6, 391–403.

Jindal, G. A., Goyal, Y., Burdine, R. D., Rauen, K. A. and Shvartsman, S. Y. (2015). RASopathies: unraveling mechanisms with animal models. *Disease Models & Mechanisms* 8, 1167–1167.

Jover, R., Rodrigo, R., Felipo, V., Insausti, R., Sáez-Valero, J., García-Ayllón, M. S., Suárez, I., Candela, A., Compañ, A., Esteban, A., et al. (2006). Brain edema and inflammatory activation in bile duct ligated rats with diet-induced hyperammonemia: A model of hepatic encephalopathy in cirrhosis. *Hepatology* 43, 1257–1266.

Kalil, K. and Dent, E. W. (2014). Branch management: mechanisms of axon branching in the developing vertebrate CNS. *Nat Rev Neurosci* 15, 7–18.

Kang, M. and Lee, Y.-S. (2019). The impact of RASopathy-associated mutations on CNS development in mice and humans. *Mol Brain* 12, 96.

Kaul, A., Toonen, J. A., Cimino, P. J., Gianino, S. M. and Gutmann, D. H. (2015). Akt- or MEK-mediated mTOR inhibition suppresses Nf1 optic glioma growth. *Neuro-Oncology* 17, 843–853.

Kawai, R., Markman, T., Poddar, R., Ko, R., Fantana, A. L., Dhawale, A. K., Kampff, A. R. and Ölveczky, B. P. (2015). Motor Cortex Is Required for Learning but Not for Executing a Motor Skill. *Neuron* 86, 800–812.

Kim, Y. E. and Baek, S. T. (2019). Neurodevelopmental Aspects of RASopathies. *Mol.Cells* 42, 441–447.

Kim, Y. E., Kim, Y.-S., Lee, H.-E., So, K. H., Choe, Y., Suh, B.-C., Kim, J.-H., Park, S. K., Mathern, G. W., Gleeson, J. G., et al. (2023). Reversibility and developmental neuropathology of linear nevus sebaceous syndrome caused by dysregulation of the RAS pathway. *Cell Rep* 42, 112003.

Klesse, L. J., Meyers, K. A., Marshall, C. J. and Parada, L. F. (1999). Nerve growth factor induces survival and differentiation through two distinct signaling cascades in PC12 cells. *Oncogene* 18, 2055–2068.

Klomp, J. E., Klomp, J. A. and Der, C. J. (2021). The ERK mitogen-activated protein kinase signaling network: the final frontier in RAS signal transduction. *Biochemical Society Transactions* 49, 253–267.

Knowles, S. J., Stafford, A. M., Zaman, T., Angara, K., Williams, M. R., Newbern, J. M. and Vogt, D. (2023a). Distinct hyperactive RAS/MAPK alleles converge on common GABAergic interneuron core programs. *Development* 150, dev201371.

Knowles, S. J., Holter, M. C., Li, G., Bjorklund, G. R., Rees, K. P., Martinez-Fuentes, J. S., Nishimura, K. J., Afshari, A. E., Fry, N., Stafford, A. M., et al. (2023b). *Multifunctional requirements for ERK1/2 signaling in the development of ganglionic eminence derived glia and cortical inhibitory neurons*. elife.

Kozorovitskiy, Y., Saunders, A., Johnson, C. A., Lowell, B. B. and Sabatini, B. L. (2012).

    Recurrent network activity drives striatal synaptogenesis. *Nature* 485, 646–650.

Krenz, M., Gulick, J., Osinska, H. E., Colbert, M. C., Molkentin, J. D. and Robbins, J. (2008).

    Role of ERK1/2 signaling in congenital valve malformations in Noonan syndrome. *Proc.*

    *Natl. Acad. Sci. U.S.A.* 105, 18930–18935.

Lajiness, J. D., Snider, P., Wang, J., Feng, G.-S., Krenz, M. and Conway, S. J. (2014). SHP-2

    deletion in postmigratory neural crest cells results in impaired cardiac sympathetic

    innervation. *Proc. Natl. Acad. Sci. U.S.A.* 111,.

Langdon, Y., Tandon, P., Paden, E., Duddy, J., Taylor, J. M. and Conlon, F. L. (2012). SHP-2

    acts via ROCK to regulate the cardiac actin cytoskeleton. *Development* 139, 948–957.

Lavoie, H., Gagnon, J. and Therrien, M. (2020). ERK signalling: a master regulator of cell

    behaviour, life and fate. *Nat Rev Mol Cell Biol* 21, 607–632.

Lee, J., Urban-Ciecko, J., Park, E., Zhu, M., Myal, S. E., Margolis, D. J. and Barth, A. L. (2021).

    FosGFP expression does not capture a sensory learning-related engram in superficial

    layers of mouse barrel cortex. *Proc. Natl. Acad. Sci. U.S.A.* 118, e2112212118.

Lemmon, M. A. and Schlessinger, J. (2010). Cell Signaling by Receptor Tyrosine Kinases. *Cell*

    141, 1117–1134.

Levy, S., Lavzin, M., Benisty, H., Ghanayim, A., Dubin, U., Achvat, S., Brosh, Z., Aeed, F.,

    Mensh, B. D., Schiller, Y., et al. (2020). Cell-Type-Specific Outcome Representation in

    the Primary Motor Cortex. *Neuron* 107, 954-971.e9.

Li, Y. and Hollis, E. (2021). Basal Forebrain Cholinergic Neurons Selectively Drive Coordinated

    Motor Learning in Mice. *J. Neurosci.* 41, 10148–10160.

Li, W., Cui, Y., Kushner, S. A., Brown, R. A. M., Jentsch, J. D., Frankland, P. W., Cannon, T. D.

    and Silva, A. J. (2005). The HMG-CoA Reductase Inhibitor Lovastatin Reverses the

    Learning and Attention Deficits in a Mouse Model of Neurofibromatosis Type 1. *Current*

    *Biology* 15, 1961–1967.

Li, C., Scott, D. A., Hatch, E., Tian, X. and Mansour, S. L. (2007). *Dusp6* ( *Mkp3* ) is a negative

    feedback regulator of FGF-stimulated ERK signaling during mouse development.

    *Development* 134, 167–176.

Li, X., Newbern, J. M., Wu, Y., Morgan-Smith, M., Zhong, J., Charron, J. and Snider, W. D.

    (2012). MEK Is a Key Regulator of Gliogenesis in the Developing Brain. *Neuron* 75,

    1035–1050.

Lim, B., Dsilva, C. J., Levario, T. J., Lu, H., Schüpbach, T., Kevrekidis, I. G. and Shvartsman, S.

    Y. (2015). Dynamics of Inductive ERK Signaling in the Drosophila Embryo. *Current*

    *Biology* 25, 1784–1790.

López-Juárez, A., Titus, H. E., Silbak, S. H., Pressler, J. W., Rizvi, T. A., Bogard, M., Bennett,

    M. R., Ciraolo, G., Williams, M. T., Vorhees, C. V., et al. (2017). Oligodendrocyte Nf1

    Controls Aberrant Notch Activation and Regulates Myelin Structure and Behavior. *Cell*

    *Reports* 19, 545–557.

Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L.,

    Palmiter, R. D., Hawrylycz, M. J., Jones, A. R., et al. (2010). A robust and high-

    throughput Cre reporting and characterization system for the whole mouse brain. *Nat*

    *Neurosci* 13, 133–140.

Markus, A., Zhong, J. and Snider, W. D. (2002). Raf and Akt Mediate Distinct Aspects of

    Sensory Axon Growth. *Neuron* 35, 65–76.

Mayes, D. A., Rizvi, T. A., Titus-Mitchell, H., Oberst, R., Ciraolo, G. M., Vorhees, C. V.,

    Robinson, A. P., Miller, S. D., Cancelas, J. A., Stemmer-Rachamimov, A. O., et al.

    (2013). Nf1 Loss and Ras Hyperactivation in Oligodendrocytes Induce NOS-Driven

    Defects in Myelin and Vasculature. *Cell Reports* 4, 1197–1212.

Mebratu, Y. and Tesfaigzi, Y. (2009). How ERK1/2 activation controls cell proliferation and cell

    death: Is subcellular localization the answer? *Cell Cycle* 8, 1168–1175.

Monje, P., Hernández-Losa, J., Lyons, R. J., Castellone, M. D. and Gutkind, J. S. (2005). Regulation of the transcriptional activity of c-Fos by ERK: A novel role for the prolyl isomerase Pin1. *Journal of Biological Chemistry* 280, 35081–35084.

Morris, E. J., Jha, S., Restaino, C. R., Dayananth, P., Zhu, H., Cooper, A., Carr, D., Deng, Y., Jin, W., Black, S., et al. (2013). Discovery of a Novel ERK Inhibitor with Activity in Models of Acquired Resistance to BRAF and MEK Inhibitors. *Cancer Discovery* 3, 742–750.

Nakamura, T., Gulick, J., Pratt, R. and Robbins, J. (2009). Noonan syndrome is associated with enhanced pERK activity, the repression of which can prevent craniofacial malformations. *Proc. Natl. Acad. Sci. U.S.A.* 106, 15436–15441.

Newbern, J. M., Li, X., Shoemaker, S. E., Zhou, J., Zhong, J., Wu, Y., Bonder, D., Hollenback, S., Coppola, G., Geschwind, D. H., et al. (2011). Specific Functions for ERK/MAPK Signaling during PNS Development. *Neuron* 69, 91–105.

Nikolaienko, O., Eriksen, M. S., Patil, S., Bito, H. and Bramham, C. R. (2017). Stimulus-evoked ERK-dependent phosphorylation of activity-regulated cytoskeleton-associated protein (Arc) regulates its neuronal subcellular localization. *Neuroscience* 360, 68–80.

Nowaczyk, M. J. M., Thompson, B. A., Zeesman, S., Moog, U., Sanchez-Lara, P. A., Magoulas, P. L., Falk, R. E., Hoover-Fong, J. E., Batista, D. A. S., Amudhavalli, S. M., et al. (2014). Deletion of *MAP2K2/MEK2*: a novel mechanism for a RASopathy?: *MAP2K2/MEK2* deletion. *Clin Genet* 85, 138–146.

Osterweil, E. K., Chuang, S.-C., Chubykin, A. A., Sidorov, M., Bianchi, R., Wong, R. K. S. and Bear, M. F. (2013). Lovastatin Corrects Excess Protein Synthesis and Prevents Epileptogenesis in a Mouse Model of Fragile X Syndrome. *Neuron* 77, 243–250.

Otsu, N. (1979). A Threshold Selection Method from Gray-Level Histograms. *IEEE Trans. Syst., Man, Cybern.* 9, 62–66.

Owen, J. P., Bukshpun, P., Pojman, N., Thieu, T., Chen, Q., Lee, J., D'Angelo, D., Glenn, O. A., Hunter, J. V., Berman, J. I., et al. (2018). Brain MR Imaging Findings and Associated

Outcomes in Carriers of the Reciprocal Copy Number Variation at 16p11.2. *Radiology* 286, 217–226.

Ozdinler, P. H. and Macklis, J. D. (2006). IGF-I specifically enhances axon outgrowth of corticospinal motor neurons. *Nat Neurosci* 9, 1371–1381.

Parmar, M. S., Jaumotte, J. D., Zigmond, M. J. and Cavanaugh, J. E. (2015). ERK1, 2, and 5 expression and activation in dopaminergic brain regions during postnatal development. *Int. j. dev. neurosci.* 46, 44–50.

Peters, A. J., Lee, J., Hedrick, N. G., O'Neil, K. and Komiyama, T. (2017). Reorganization of corticospinal output during motor learning. *Nat Neurosci* 20, 1133–1141.

Pintchovski, S. A., Peebles, C. L., Joo Kim, H., Verdin, E. and Finkbeiner, S. (2009). The Serum Response Factor and a Putative Novel Transcription Factor Regulate Expression of the Immediate-Early Gene Arc/Arg3.1 in Neurons. *Journal of Neuroscience* 29, 1525–1537.

Polleux, F. and Snider, W. (2010). Initiating and Growing an Axon. *Cold Spring Harbor Perspectives in Biology* 2, a001925–a001925.

Poulopoulos, A., Murphy, A. J., Ozkan, A., Davis, P., Hatch, J., Kirchner, R. and Macklis, J. D. (2019). Subcellular transcriptomes and proteomes of developing axon projections in the cerebral cortex. *Nature* 565, 356–360.

Rai, B., Naylor, P. E., Siqueiros-Sanchez, M., Wintermark, M., Raman, M. M., Jo, B., Reiss, A. L. and Green, T. (2023). Novel effects of Ras-MAPK pathogenic variants on the developing human brain and their link to gene expression and inhibition abilities. *Transl Psychiatry* 13, 245.

Rao, V. R., Pintchovski, S. A., Chin, J., Peebles, C. L., Mitra, S. and Finkbeiner, S. (2006). AMPA receptors regulate transcription of the plasticity-related immediate-early gene Arc. *Nat Neurosci* 9, 887–895.

Rauen, K. A. (2013). The RASopathies. *Annu Rev Genomics Hum Genet.* 14, 355–369.

Rauen, K. A. (2022). Defining RASopathy. *Disease Models & Mechanisms* 15, dmm049344.

Ray, L. B. and Sturgill, T. W. (1987). Rapid stimulation by insulin of a serine/threonine kinase in 3T3-L1 adipocytes that phosphorylates microtubule-associated protein 2 in vitro. *Proceedings of the National Academy of Sciences* 84, 1502–1506.

Roskoski, R. (2019). Targeting ERK1/2 protein-serine/threonine kinases in human cancers. *Pharmacological Research* 142, 151–168.

Rumbaugh, G., Adams, J. P., Kim, J. H. and Huganir, R. L. (2006). SynGAP regulates synaptic strength and mitogen-activated protein kinases in cultured neurons. *Proc Natl Acad Sci U S A* 103, 4344–4351.

Ryu, H.-H. and Lee, Y.-S. (2016). Cell type-specific roles of RAS-MAPK signaling in learning and memory: Implications in neurodevelopmental disorders. *Neurobiology of Learning and Memory* 135, 13–21.

Santini, C. C., Longden, J., Schoof, E. M., Simpson, C. D., Jeschke, G. R., Creixell, P., Kim, J., Wu, X., Turk, B. E., Rosen, N., et al. (2019). Global view of the RAF-MEK-ERK module and its immediate downstream effectors. *Sci Rep* 9, 10865.

Shaul, Y. D. and Seger, R. (2007). The MEK/ERK cascade: From signaling specificity to diverse functions. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1773, 1213–1226.

Shigeoka, T., Jung, H., Jung, J., Turner-Bridger, B., Ohk, J., Lin, J. Q., Amieux, P. S. and Holt, C. E. (2016). Dynamic Axonal Translation in Developing and Mature Visual Circuits. *Cell* 166, 181–192.

Steinhoff, J. S., Lass, A. and Schupp, M. (2021). Biological Functions of RBP4 and Its Relevance for Human Diseases. *Front. Physiol.* 12, 659977.

Steward, O. and Worley, P. F. (2001). Selective targeting of newly synthesized Arc mRNA to active synapses requires NMDA receptor activation. *Neuron* 30, 227–240.

Steward, O., Wallace, C. S., Lyford, G. L. and Worley, P. F. (1998). Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites. *Neuron* 21, 741–751.

Thomas, G. M. and Huganir, R. L. (2004). MAPK cascade signalling and synaptic plasticity. *Nat Rev Neurosci* 5, 173–183.

Thompson, S. J., Sargsyan, A., Lee, S.-A., Yuen, J. J., Cai, J., Smalling, R., Ghyselinck, N., Mark, M., Blaner, W. S. and Graham, T. E. (2017). Hepatocytes Are the Principal Source of Circulating RBP4 in Mice. *Diabetes* 66, 58–63.

Tidyman, W. E. and Rauen, K. A. (2009). The RASopathies: developmental syndromes of Ras/MAPK pathway dysregulation. *Current Opinion in Genetics & Development* 19, 230–236.

Tiemens, D. K., Kleimeier, L., Leenders, E., Wingbermühle, E., Roelofs, R. L., Sibbles, B., Oostwegel, F. S. M., Vroonland, E., van Leeuwen, C., Niessen, H., et al. (2023). The most important problems and needs of rasopathy patients with a noonan syndrome spectrum disorder. *Orphanet J Rare Dis* 18, 198.

Titus, H. E., López-Juárez, A., Silbak, S. H., Rizvi, T. A., Bogard, M. and Ratner, N. (2017). Oligodendrocyte RasG12V expressed in its endogenous locus disrupts myelin structure through increased MAPK, nitric oxide, and notch signaling. *Glia* 65, 1990–2002.

Tomson, S. N., Schreiner, M. J., Narayan, M., Rosser, T., Enrique, N., Silva, A. J., Allen, G. I., Bookheimer, S. Y. and Bearden, C. E. (2015). Resting state functional MRI reveals abnormal network connectivity in neurofibromatosis 1. *Human Brain Mapping* 36, 4566–4581.

Tsai, W.-H. (1985). Moment-preserving thresolding: A new approach. *Computer Vision, Graphics, and Image Processing* 29, 377–393.

Ünal, E. B., Uhlitz, F. and Blüthgen, N. (2017). A compendium of ERK targets. *FEBS Lett* 591, 2607–2615.

Wang, Y., Kim, E., Wang, X., Novitch, B. G., Yoshikawa, K., Chang, L.-S. and Zhu, Y. (2012). ERK Inhibition Rescues Defects in Fate Specification of Nf1-Deficient Neural Progenitors and Brain Abnormalities. *Cell* 150, 816–830.

Wen, Y., Alshikho, M. J. and Herbert, M. R. (2016). Pathway Network Analyses for Autism Reveal Multisystem Involvement, Major Overlaps with Other Diseases and Convergence upon MAPK and Calcium Signaling. *PLoS ONE* 11, e0153329.

Whishaw, I. Q., Pellis, S. M., Gorny, B., Kolb, B. and Tetzlaff, W. (1993). Proximal and distal impairments in rat forelimb use in reaching follow unilateral pyramidal tract lesions. *Behavioural Brain Research* 56, 59–76.

Wu, X., Simpson, J., Hong, J. H., Kim, K.-H., Thavarajah, N. K., Backx, P. H., Neel, B. G. and Araki, T. (2011). MEK-ERK pathway modulation ameliorates disease phenotypes in a mouse model of Noonan syndrome associated with the Raf1(L613V) mutation. *J Clin Invest* 121, 1009–1025.

Xing, L., Larsen, R. S., Bjorklund, G. R., Li, X., Wu, Y., Philpot, B. D., Snider, W. D. and Newbern, J. M. (2016). Layer specific and general requirements for ERK/MAPK signaling in the developing neocortex. *eLife* 5, e11123.

Ying, S.-W., Futter, M., Rosenblum, K., Webber, M. J., Hunt, S. P., Bliss, T. V. P. and Bramham, C. R. (2002). Brain-Derived Neurotrophic Factor Induces Long-Term Potentiation in Intact Adult Hippocampus: Requirement for ERK Activation Coupled to CREB and Upregulation of *Arc* Synthesis. *J. Neurosci.* 22, 1532–1540.

Yun, M.-S., Kim, S.-E., Jeon, S. H., Lee, J.-S. and Choi, K.-Y. (2005). Both ERK and Wnt/β-catenin pathways are involved in Wnt3a-induced proliferation. *Journal of Cell Science* 118, 313–322.

Zalfa, F., Giorgi, M., Primerano, B., Moro, A., Di Penta, A., Reis, S., Oostra, B. and Bagni, C. (2003). The Fragile X Syndrome Protein FMRP Associates with BC1 RNA and Regulates the Translation of Specific mRNAs at Synapses. *Cell* 112, 317–327.

Zamboni, S. L., Loenneker, T., Boltshauser, E., Martin, E. and Il'yasov, K. A. (2007). Contribution of diffusion tensor MR imaging in detecting cerebral microstructural changes in adults with neurofibromatosis type 1. *AJNR Am J Neuroradiol* 28, 773–776.

Zamorano, C., Fernández-Albert, J., Storm, D. R., Carné, X. and Sindreu, C. (2018). Memory
Retrieval Re-Activates Erk1/2 Signaling in the Same Set of CA1 Neurons Recruited
During Conditioning. *Neuroscience* 370, 101–111.

Zenker, M., Edouard, T., Blair, J. C. and Cappa, M. (2022). Noonan syndrome: improving
recognition and diagnosis. *Arch Dis Child* 107, 1073–1078.

## Adult Forebrain

**Supplemental Figure 1. Increased levels of pERK1/2 in hippocampal axons of adult Nex:Cre; MEK1$^{S217/221E}$ mice.**

**A-D.** Immunolabeling of forebrain sections at the level of the adult dorsal hippocampus showed a prominent increase of pERK1/2 within the mutant fimbria and stria terminalis (C-D) relative to controls (A-B), which are enriched in axons derived from excitatory pyramidal neurons in the hippocampal CA1-3 and the amygdala, respectively (n=4, fi = fimbria, st = stria terminalis). Scale bars = 200μm.

# Adult Forebrain

# Adult Spinal Cord

# Adult Liver

**Supplemental Figure 2. Conditional expression of MEK1$^{S217/221E}$ in RBP4:Cre mice.**

**A-F.** Representative confocal images of *RBP4:Cre; Ai9$^{+/-}$* cortices. Both *RBP4:Cre; Ai9$^{+/-}$* mice control and *RBP4:Cre; MEK1$^{S217/221E}$; Ai9$^{+/-}$* mutants express RFP/tdTomato in cortical layer V (A-B, D-E). Immunolabeling of MEK1 showed a robust increase of expression restricted to cortical layer V in mutants (F) compared to controls (C).

**G-L.** Representative images of cross-sectional lumbar segments showing that RFP is detected in descending projections in the dorsal funiculus of controls (G-H) and mutants (J-K). High resolution images reveal increased MEK1 expression in RFP-labeled axons of mutants (K-L) compared to controls (H-I).

**M.** Gross dissection of the mutant liver was suggestive of hepatomegaly when compared to littermate controls (M). (M1 = primary motor cortex, cc = corpus collosum, st = striatum, S = septum, dh = dorsal horn, vh = ventral horn, cc = central canal, df = dorsal funiculus).

Scale bars = Forebrain: A-F = 500 µm, Spinal Cord: G, K = 500µm, H-I, K-L = 50µm, Liver: M = 1cm.

## AAV:*CAG-FLEX-tdTomato* P1 Injection: Collect P30

## Corticobulbar Tract Labeling

**Supplemental Figure 3. Corticobulbar tract axonal extension is not affected in *RBP4:Cre; MEK1$^{S217/2121E}$* mice.**

**A-C.** Representative AAV injection sites in M1. AAV-tdTomato labeled cell counts in the motor cortex showed no significant differences between *RBP4:Cre;MEK1$^{S217/221E}$* (B) and *RBP4:Cre* (A) mice as quantified in C (RFP$^{+}$ cells per section, mean ± SEM, n=5, Student's t-test p=0.4). **D-H.** Representative images of corticobulbar tract labeling. Quantification of tdTomato labeled axons in the corticobulbar tract revealed no significant differences between the *RBP4:Cre; MEK1$^{S217/221E}$* (F-G) and *RBP4:Cre* (D-E) control mice (normalized to labeled neurons, mean ± SEM, n=4, Student's t-test, p=0.818) (H).

Scale bars: A,B,D,F = 500µm, E,G= 20µm.

**Supplemental Figure 4. Model of aberrant circuit development in excitatory neuron-specific *MEK1$^{217/221E}$* mice.**

**A-C.** *MEK1$^{S217/221E}$* expressing layer V neurons (red) exhibit reductions in corticospinal axon elongation along the dorsal funiculus (A), reduced arborization in contralateral cortex and striatum (B), and persistent increases in axonal pERK1/2 (C - green dots) when compared to control neurons (black). These axonal deficits in *Nex:Cre; MEK1$^{S217/221E}$* mutants coincide with decreased levels of pERK1/2 in the soma and nucleus and reduced expression of activity regulated cytoskeleton-associated protein, ARC (blue), which may underly select deficits in learning a skilled forelimb task (C).