Cortical high gamma power relates to movement speed and is disproportionately reduced after stroke

Benjamin Haverland[1,2], Lena S. Timmsen[1,2], Silke Wolf[1], Charlotte J. Stagg[3,4], Lukas Frontzkowski[1], Jan Feldheim[1], Focko L. Higgen[1], Christian Gerloff[1], Robert Schulz[1], Till R. Schneider[2], Bettina C. Schwab[5,2,*,#], Fanny Quandt[1,*,#]

1 Department of Neurology, University Medical Center Hamburg-Eppendorf, 20246 Hamburg, Germany
2 Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, 20246 Hamburg, Germany
3 Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, OX3 9DU Oxford, UK
4 Research Council Brain Network Dynamics Unit, Nuffield Department of Clinical Neurosciences, University of Oxford, OX3 9DU Oxford, UK
5 Biomedical Signals and Systems, Technical Medical Centre, University of Twente, 7522 NB Enschede, The Netherlands
* These authors contributed equally to this work

Corresponding authors: Fanny Quandt, University Medical Center Hamburg-Eppendorf, Martinistraße 52, 20246 Hamburg, Germany, Email: f.quandt@uke.de; Bettina C. Schwab, University of Twente, Drienerlolaan 5, 7522 NB Enschede, The Netherlands, Email: b.c.schwab@utwente.nl

Abstract

Motor cortical high gamma oscillations (60 to 90 Hz) occur at the onset of movement and are spatially focused over the contralateral primary motor cortex. They have been shown to be prokinetic in nature and contain information on various movement parameters. Although high gamma oscillations are widely recognized for their significance in human motor control, their precise function on a cortical level remains elusive. Further, their relevance in human stroke pathophysiology is unknown. Here, we aim to investigate how cortical high gamma oscillations relate to motor performance and motor skill acquisition, and test for possible alterations after ischemic stroke.

We performed an observational study investigating the relationship between cortical high gamma oscillations and motor function. We recorded magnetoencephalography data during a thumb movement speed task in 14 chronic stroke patients (age = 65.4 ± 9.3 years, 7 females), 15 age-matched control participants (age = 64.5 ± 8.4 years, 7 females) and 29 healthy young participants (age = 25.4 ± 4.6 years, 13 females).

Motor cortical high gamma oscillations showed a strong relation with *movement speed* as trials with higher *movement speed* were associated with greater high gamma power. Stroke patients showed reduced cortical high gamma power, surpassing the effect attributable to decreased *movement speed* in these patients. The decrease of high gamma power, however, did not relate to residual motor function or structural disconnection. In slower movements, we observed distinct high gamma peaks for each movement, whereas in faster movements high gamma activation smeared to fewer distinct peaks. Even though motor skill acquisition was evident in all groups, it was not linked to high gamma power.

Our study is the first to quantify high gamma oscillations after stroke, revealing a reduction in movement-related high gamma power. Moreover, we provide strong evidence for a pivotal role of motor cortical high gamma oscillations in encoding *movement speed*. Our findings underscore the potential for exploring non-invasive brain stimulation within the high gamma frequency range to enhance motor rehabilitation after stroke.

Introduction

Gamma oscillations (25 to 140 Hz) have been found in various areas of the human brain[1] and have been linked to neural communication[2,3] as well as to a wide range of cognitive processes.[4] At the onset of motor actions, spatially focused movement-related high-gamma oscillations (~ 60 to 90 Hz) occur over the contralateral motor cortex[5], which have been shown to be prokinetic.[6–10] High gamma oscillations have been recorded in motor cortical areas from extracellular recordings (single- or multi-unit, local field potentials (LFPs))[11,12], electrocorticographic (ECoG) recordings[7,13], as well as non-invasive magnetoencephalography (MEG)[6,14] and electroencephalography (EEG).[15] In particular, invasive cortical recordings showed that high gamma oscillations contain decoding information on position, velocity and predominantly movement speed.[13,16,17]

Although high gamma oscillations seem to play a crucial role in human motor control, they have so far never been quantified in stroke patients. As over 50% of all stroke patients suffer from persistent hand motor deficits[18], understanding the neurophysiological processes of motor coding after stroke is key for the development of interventions to improve recovery and motor performance. Moreover, the study of human populations with impaired motor function may shed light on the general role of movement-related high gamma oscillations. For example, if high gamma oscillations purely reflect movement kinematics, they should be proportionally scaled down with impaired motor function. If high gamma oscillations rather reflect the intent to move or the cognitive effort, they would be expected to be preserved or even enhanced with restricted motor function.

In the current study, we characterize the role of movement-related high gamma oscillations in motor control and motor skill acquisition by analysing MEG data from stroke patients, healthy age-matched control participants and healthy young participants. We hypothesize that high gamma oscillations relate to *movement speed* and explore their role in motor control in the chronic phase after stroke.

Materials and methods

Participants and setting

Three different groups were recruited within the study: (i) 16 well-recovered stroke patients, (ii) 18 healthy control participants age-matched to the stroke patients and (iii) 30 healthy young participants (age 18 to 35 years). Stroke patients, with a first-ever clinical stroke at least 6 months prior, who experienced a deficit of the upper extremity for at least 24 hours and showed a structural lesion on a clinical magnetic resonance (MR) or computer tomography image were included in the study.

Psychotropic medication was an exclusion criterion in all three groups. All participants were right-handed according to the Edinburgh Handedness Inventory. One healthy young participant and three age-matched older controls had to be excluded because they did not fulfil all inclusion criteria retrospectively (neurological conditions, handedness). Two stroke patients were excluded because of extensive artifacts in the MEG data due to metal implants. The number of recruited stroke patients was based on prior comparable observational studies.[19,20]

Multimodal imaging was obtained, consisting of MEG measurements during the execution of a motor task as well as structural MR imaging. Standardized clinical testing was acquired in stroke patients and age-matched control participants by means of the Fugl-Meyer Assessment of the Upper Extremity (UEFM), Action Research Arm Test (ARAT), Nine Hole Peg Test (NHPT), Box and Block Test (BBT), whole hand grip force, key grip force, National Institutes of Health Stroke Scale (NIHSS), modified Rankin Scale (mRS) and Mini Mental State Examination (MMSE).

Participants gave written informed consent and the study was performed according to the Declaration of Helsinki. The study was approved by the local ethics committee of the Medical Association of Hamburg (2021-10410-BO-ff).

Motor Task

Participants were seated comfortably in a CTF-MEG system (CTF Systems, Coquitlam, Canada) in front of a screen with their performing arm fixated in a splint, next to a two-button response box (Current Designs, Philadelphia, PA) that was placed in a position where the thumb of the participant was able to easily reach both buttons. The task consisted of alternately pressing two buttons for a total of four button presses in reaction to a 'ready – steady – go' cue using the thumb while keeping the arm as relaxed as possible (Fig. 2A).

First, participants performed 20 trials to familiarize themselves with the task. Afterwards, they were instructed to perform the task as fast as possible using standardized instructions. Each participant performed six blocks with 40 trials each. After each trial, participants received feedback on their absolute task performance and the improvement relative to their performance in the current and the previous block. After each block, additional feedback was given in form of a bar indicating the mean performance for each completed block. If participants pressed the button before the go-cue, took more than one second before initiating the first button press, or pressed an incorrect order of buttons, an error message was displayed. There were differences in the experimental set-up between groups of young and control/stroke participants to account for behavioural differences and needs between these

Table 1 Clinical and demographic data of stroke patients

Patient	Sex	Age	Lesion side	TAS (months)	Stroke location	MRS	NIHSS	UEFM	ARAT	Grip force	KG force	NHPT	BBT
1	F	70	L	52	TC, CR	1	0	65	1.00	1.11	1.16	1.09	0.93
2	F	79	R	8	TC, CR, CI	3	2	46	0.77	0.09	0.60	0.20	0.53
3	M	74	R	153	BG, CI, CR	2	1	50	0.95	0.51	0.60	0.64	0.66
4	F	71	R	52	BG	1	1	63	1.00	0.97	1.07	0.93	0.96
5	F	74	R	43	PLCI	1	0	60	0.98	0.89	0.80	0.83	0.82
6	F	43	L	38	PLCI	0	0	66	1.00	1.59	1.26	1.22	1.11
7	F	73	L	39	MED	1	1	58	0.98	0.92	0.88	1.11	0.94
8	M	65	L	9	BG	1	0	66	1.00	0.93	1.28	1.16	1.15
9	M	61	R	9	PLCI	1	0	66	0.98	0.89	1.12	0.88	0.91
10	M	58	L	6	CR	1	0	64	1.00	1.05	1.00	1.05	1.00
11	M	57	L	51	MED	1	2	63	0.93	0.98	1.14	0.77	0.9
12	F	67	R	129	PO	0	0	64	1.00	0.97	1.40	0.89	1.07
13	M	62	R	9	CR	1	1	60	0.95	0.68	0.90	0.52	0.64
14	M	61	R	16	PRE	1	0	64	0.95	0.69	0.75	0.81	0.91
Mean stroke	F: 7	65.4		45.6		1.1	0.6	61.1***	0.96**	0.88	1.00	0.86	0.89*
SD stroke		9.3		45.3		0.7	0.8	6.1	0.06	0.33	0.25	0.28	0.18
Mean control	F: 7	64.5				0	0	65.7	1	1.18	0.97	1.02	1
SD control		8.4				0	0	0.7	0.01	0.7	0.09	0.14	0.07

TAS: time after stroke, KG: key grip, SD: standard deviation, TC: thalamocapsular, CR: corona radiata, CI: capsula interna, BG: basal ganglia, PLCI: posterior limb of the capsula interna, MED: media infarct, PO: pons, PRE: precentral gyrus; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, statistical test of stroke against control group. Values of ARAT, grip force, key grip force, NHPT and BBT are ratios between the affected and not affected hand in stroke patients and performing and not performing hand in control participants.

groups. Young participants performed the task with their left (non-dominant) hand, while stroke patients used the hand contralateral to the lesioned hemisphere. The performing hand of age-matched control participants was chosen to match stroke patients. Moreover, while in the young group lights in the measuring chamber were turned off, the light was kept on for stroke patients and older participants. Further, the size of the feedback bar (linear versus sigmoidal increase) was adapted in the age-matched control group and in stroke participants to account for differences in *movement speed*.

Data acquisition

MEG and MRI data acquisition

MEG data were recorded with 275 axial gradiometers of a CTF-MEG system with a sampling rate of 1200 Hz. The head position was continuously measured, and participants were given instructions to help them return to their initial head position after each block of the task. Electrooculogram (horizontal and vertical) and electrocardiogram were recorded via bipolar channels. Time points of button presses and visual stimuli were sent to the acquisition program via LPT triggers.

We acquired MRI data from 21 young participants, all 15 control participants and 11 stroke patients. The remaining participants did not consent to MR-imaging. We acquired imaging data using a 3 T Prisma MRI scanner (Siemens Healthineers, Erlangen, Germany) equipped with a 64-channel head coil. For anatomical imaging in stroke patients and control participants, a 3-dimensional magnetization-prepared rapid gradient echo sequence (repetition time (TR) = 2500 ms, echo time (TE) = 2.15 ms, flip angle 8°, 288 coronal slices with a voxel size of 0.8 × 0.8 × 0.8 mm³) was used. In young participants, the parameters of the sequence were slightly different (TR = 2300 ms, TE = 2.98 ms, flip angle 9°, 256 coronal slices with a voxel size of 1 × 1 × 1 mm³). T2-weighted images were acquired by using a fluid-attenuated inversion recovery sequence (TR = 9210 ms, TE = 92 ms, TI = 2500 ms, flip angle 140°, 70 axial slices with a voxel size of 0.9 × 0.9 × 2.0 mm³) for stroke lesion delineation. In addition, multi-shell diffusion-weighted images (DWI) were obtained covering the whole brain with gradients (b = 500, 1000 and 2000 s/mm²) applied along 96 non-collinear directions with the following sequence parameters: TR = 5000 ms, TE = 76 ms, slice thickness = 2 mm, in-plane resolution = 1 × 1 mm. For patient 14 the sequence parameters varied slightly: TR = 5000 ms, TE = 78 ms, slice thickness = 2 mm, in-plane resolution = 1 x 1 mm.

Data analysis

Data analysis was conducted using the software package Fieldtrip[21] in MATLAB (R2021b, The MathWorks, Inc.). Statistical analysis was performed in R version 4.1.3.[22]

Clinical and behavioural data

ARAT, NHPT, BBT, whole hand grip force and key grip force scores of the performing hand were divided by scores of the non-performing hand to obtain a ratio score for each measure. For UEFM and MMSE we used the absolute test scores (UEFM: range 0 to 66, MMSE: range 0 to 30). Within the motor task, *movement speed* was defined as the reciprocal of the time between the first and the fourth button press in each trial. *Improvement* was defined in two different ways:

(i) on a trial level as the percentage increase in *movement speed* from the current to the next trial:

$$\textit{trial-wise improvement} = \frac{\textit{movement speed} \text{ in the next trial - } \textit{movement speed} \text{ in the current trial}}{\textit{movement speed} \text{ in the current trial}} \times 100$$

(ii) across blocks as the change from first block to best block:

$$\textit{block-wise improvement} = \frac{\textit{movement speed} \text{ in the best block - } \textit{movement speed} \text{ in the first block}}{\textit{movement speed} \text{ in the first block}} \times 100$$

MEG Data

MEG data were bandpass filtered from 1 to 120 Hz and bandstop filtered at 49 to 51 Hz and 99 to 101 Hz (to eliminate the line noise artefact and its first harmonic) using a fourth-order butterworth filter. Data were segmented into trials with the length of -3.2 s to +3.5 s with respect to the time point of the first button press (0 s). Artifacts were carefully removed by a combination of visual artifact rejection as well as an independent component analysis, in which components representing eyeblinks, cardiac activity, or muscle artifacts were removed from the data. Error trials, trials in which participants took more than 3 s to complete all four button presses as well as trials in which the movement duration (time from first to fourth button press) deviated more than three scaled absolute deviations from the individual block median were excluded, leaving 205.5 ± 16 (mean ± standard deviation) trials per participant in the young group (5960 trials in total), 191.5 ± 25.8 trials per participant in the stroke group (2681 trials in total) and 205.6 ± 14.6 trials per participant in the control group (3084 trials in total).

Source space activity was estimated using the dynamic imaging of coherent sources (DICS) beamforming method.[23] Head models, giving a geometrical description of the head, were computed based on individual T1-weighted MR images (21 young participants, 11 stroke patients, 15 control participants) or the Colin27 Average Brain[24] (8 young participants, 3 stroke patients) using the single-shell method.[25] Source models, stating the location of the source in the brain volume, consisted of a regularly spaced 8 mm 3D grid warped on an individual or standard head model respectively. At each location, a projection matrix describing signal propagation to the electrodes (the leadfield matrix) was computed. Time-frequency averaged cross-spectral density matrices for the movement period (0 to 0.6 s) and baseline period (-1.6 to -1 s) at 75 Hz with ± 15 Hz frequency smoothing (17 slepian tapers) were computed and source space activity was estimated at each grid point. To noise-normalize source power, the difference of power in the movement period and baseline period was divided by the baseline period for each trial in each participant. Brain regions were identified by masking of the source representation with the Brainnetome atlas.[26] For each participant, the voxel with maximal power in the primary motor cortex (M1, regions A4hf, A4ul, A4t, A4tl), premotor cortex (PMC, regions A6vl, A6cvl, A6dl, A6cdl) and supplementary motor area (SMA, region A6m) contralateral to their performing hand was detected. The power values of these voxels were used for further analyses.

To obtain a time-frequency representation, we computed real-valued DICS filters at 75 Hz ± 15 Hz frequency smoothing for the periods 0 to 1.5 s (task period) and -1.6 to -0.1 s (baseline period) and multiplied the filters of the voxels with maximal power in each region with the time-series data. In the resulting source level time-series data, we estimated power in the range 60 to 90 Hz (steps of 2 Hz and 0.05s, 0.25 s moving time window, 5 slepian tapers, resulting in ± 12 Hz frequency smoothing). We identified the high gamma peak frequency for each participant by averaging the data within the time interval from 0 to 0.6 s over trials and determined the frequency with the highest power value. To identify the number of peaks in participant-level frequency-averaged time courses, we used the MATLAB function findpeaks to detect local maxima within a time interval starting at the first button press until 0.2 s after the last button press. Peaks were required to have a minimum width of 0.05 s and had to be separated by at least 21% of the movement duration.

MRI-Data

Quantification of structural disconnection

Stroke lesions were delineated with ITK-SNAP[27] and registered to a Montreal National Institute (MNI) 1 mm³ template. Hemispheres of all patients with left-sided strokes were flipped to the other side. In one patient, the stroke lesion was not detectable on the MRI anymore (Patient 1). To calculate the structural disconnection for each patient, lesion masks were analysed using the

network modification NEMO2-toolbox.[28] In brief, Change of Connectivity (ChaCo) values were calculated by overlaying the MNI-registered lesion masks onto a normative structural connectome derived from 420 healthy participants from the Human Connectome Project. The ChaCo scores resemble the number of disrupted streamlines divided by the total number of streamlines connecting a brain region to the rest of the network. We applied the Brainnetome 246 atlas[26] to compute ChaCo values for ipsilesional M1, ventral premotor cortex (PMv, regions A6vl, A6cvl), dorsal premotor cortex (PMd, A6dl, A6cdl) and SMA. The mean of ChaCo values of PMv and PMd was used as ChaCo of the PMC. In agreement with pre-established studies, we removed all values below 0.02 due to noise artifacts.[28]

Fractional anisotropy of the corticospinal tract

Preprocessing and reconstruction of MR-images were performed using QSIPrep 0.16.1[29], which is based on Nipype 1.8.5[30] (RRID:SCR_002502). Tract segmentation of the corticospinal tract (CST) was performed using TractSeg.[31,32] Fractional Anisotropy (FA) values of corresponding DWI voxels were extracted using MRtrix3.[33] Mean CST-FA values of the hemisphere contralateral to the performing hand (affected hemisphere in stroke patients) were divided by the mean CST-FA values of the hemisphere ipsilateral to the performing hand to obtain ratio scores. See Supplementary material for more details.

Statistical analysis

Group differences

Differences in patient characteristics, *movement speed*, block-wise improvement, high gamma power and high gamma peak frequency were assessed using two-tailed *t*-tests and Wilcoxon rank-sum tests as appropriate.

High gamma power compared to baseline

Paired two-tailed *t*-tests were calculated in each source voxel between movement period and baseline and corrected for multiple comparisons via cluster-based nonparametric statistical tests[34] with 1000 randomizations and a cluster threshold of $p = 0.025$. Data from participants performing the task with the right hand were flipped across hemispheres prior to the analysis.

High gamma power matched for movement speed

To evaluate whether group differences in high gamma power are solely due to differences in *movement speed* or also due to an effect of group, we matched each trial of each stroke patient to a trial with the same *movement speed* (± 1%) of a control participant performing the task with the same hand as well as each trial of each young participant to a trial of a control participant. Trials were then averaged within the participants and the group

difference was assessed with a one-tailed *t*-tests with the alternative hypothesis being that stroke patients have less high gamma power than control participants and that control participants have less high gamma power than young participants. Given the random selection of trials in the matching process, we performed the analysis 1000 times and used the median *p*-value as the indicator of significance.

Modelling the association of behavioural data and high gamma oscillations

Linear mixed-effects models were calculated for each group and motor region (M1, PMC) to assess the association between (i) high gamma power and *movement speed* as well as (ii) high gamma power and *trial-wise improvement,* controlled for *movement speed*. Models included participants as a random intercept. Further, linear regression models were fitted to (iii) evaluate the relation of high gamma peak frequency and *movement speed*, (iv) high gamma power and *block-wise improvement*, controlled for *movement speed* and (v) high gamma power and clinical scores, ChaCo or CST-FA. In all models (age-matched controls, stroke patients), the performing hand was added as a fixed effect. To ensure homoscedasticity and linearity of the residuals, high gamma power was transformed to log(100+high gamma power) in models (i) and (ii).

Number of high gamma peaks

We modelled the dependent variable number of high gamma peaks with independent variables group, *movement speed* and performing hand using cumulative link models (CLM, ordinal package) with a flexible logit link function for the regions M1 and PMC.

Visual inspection of residual plots did not reveal any obvious deviations from homoscedasticity or normality, proportional odds assumption held true in the CLMs. In linear mixed-effects models and CLMs, significance was obtained by likelihood ratio tests of the full model against a reduced model without the effect in question. In the case of multiple comparisons, *p*-values were corrected accordingly using the false discovery rate. Full statistical test and model results are provided in a tabular form in the Supplementary material.

Results

Participants

Fourteen stroke patients (age = 65.4 ± 9.3 (mean ± standard deviation) years, 7 females, 8 with lesion of the right hemisphere, performing hand: 8 x left hand, 6 x right hand), 15 control participants (age = 64.5 ± 8.4 years, 7 females, performing hand: 10 x left hand, 5 x right hand),

Figure 1 Stroke patient characteristics. (**A**) Stroke lesions of the 10 patients with available MRI lesion data, overlayed on a T1-weighted image in MNI standard space. Colour indicates the number of stroke patients with a lesion at the voxel. Left-hemispheric lesions were flipped to the right hemisphere. (**B**) Structural disconnection of the 10 stroke patients. Colour indicates number of stroke patients with impaired connection between regions of M1 and the respective region. (**C**) Distribution of CST-FA ratios. There was no significant group difference. (**D**) Distribution of UEFM-scores. Asterisks indicate the significant group difference. Significance markers: ***$p < 0.001$

and 29 young participants (age = 25.4 ± 4.6 years, 13 females) passed inclusion criteria. Stroke patients and control participants showed no signs of cognitive impairment according to the MMSE (stroke: mean 28.9 points, range 27 to 30 points; control: mean 29.7 points, range 28 to 30 points).

Clinical scores and structural imaging

Stroke patients mostly showed only mild impairment of the upper extremity (Table 1, mean UEFM = 61.1), with a significant difference between stroke patients and control participants in UEFM (Fig. 1D), ARAT and BBT (UEFM: $p < 0.001$, ARAT: $p = 0.004$, BBT: $p = 0.047$) but no significant difference in NHPT ($p = 0.17$), whole hand ($p = 0.08$) and key grip force ratio ($p = 0.69$). CST-FA values were not significantly different between stroke patients and control participants (Fig. 1C, $p = 0.29$). Stroke patients had a mean ChaCo of 0.19 (range 0 to 0.59) in M1 and 0.14 (range 0 to 0.55) in PMC. Most frequently disrupted streamlines from M1 were the ones targeting ipsilateral motor areas (Fig. 1B).

Improvement in movement speed

On average, stroke patients had a significantly lower *movement speed* than control participants (Fig. 2C, $p_{cor} = 0.001$; corrected for two comparisons), and control participants had a significantly lower *movement speed* than young participants ($p_{cor} < 0.001$; corrected for two comparisons). All three groups improved their average

movement speed from block 1 to block 6 (Fig. 2B, stroke patients: $p_{cor} = 0.02$, control participants: $p_{cor} = 0.01$, young participants: $p_{cor} = 0.001$; corrected for three comparisons). Comparing the initial and the individual best block, stroke patients increased their average *movement speed* by 22.6 ± 12.2 % (mean ± standard deviation), control participants by 15.3 ± 7.1 % and young participants by 18.0 ± 10.9 %. *Block-wise improvement* showed no significant differences between the three groups, even without correction for multiple comparisons (Fig. 2D, stroke/control: $p = 0.07$, control/young: $p = 0.33$).

Movement-related high gamma power

We observed an increase in high gamma power during the movement period (0 to 0.6 s, 60 to 90 Hz) relative to baseline with a maximum over the contralateral motor cortices (Fig. 3A). The increase of high gamma power during movement was significant compared to baseline in each group (Supplementary Fig. 1, stroke patients: $p_{cor} = 0.02$, control participants: $p_{cor} = 0.001$, young participants: $p_{cor} = 0.001$; corrected for three comparisons). The increase in high gamma power occurred with movement onset and was most pronounced in M1 and PMC (Fig. 6C, Supplementary Fig. 2), prompting us to focus further analysis on these regions. Stroke patients had significantly lower high gamma power than control participants in the voxel with maximal power in regions M1 and PMC (Fig. 3B, M1: $p_{cor} = 0.04$, PMC: $p_{cor} = 0.04$; corrected for four comparisons) and control participants had significantly lower high gamma power than young participants (M1: p_{cor}

Figure 2 Motor task and behavioural data. (**A**) The motor task contained one training block with 20 trials and six measurement blocks with 40 trials each. In each trial, participants had to perform four alternating button presses as fast as possible in response to a visual cue. They received feedback on their *movement speed* in each trial. Additional feedback showing the block mean of *movement speed* was given after each block. There was a 1 minute rest period between blocks. (**B**) *Movement speed* for each trial averaged in each group. An increase in *movement speed* over the course of the trials was evident in all groups. Shaded areas indicate ± standard deviation. (**C**) Distribution of participant mean of *movement speed*. Asterisks indicate significant group differences. (**D**) Distribution of participant mean of *block-wise improvement*. *Block-wise improvement* represents the percentage *improvement* in *movement speed* from the first to the individual best block. There were no significant group differences. Significance markers: **$p < 0.01$, ***$p < 0.001$

= 0.04, PMC: p_{cor} = 0.03; corrected for four comparisons). When matching performance (*movement speed*) and handedness in stroke patients and age-matched controls (trials in each group 1206.1 ± 2.7 (mean ± standard deviation)), stroke patients still had a significantly lower high gamma power than control participants in both motor regions (Fig. 3C, M1: p_{cor} = 0.04, PMC: p_{cor} = 0.04; corrected for two comparisons). In contrast, the group difference in high gamma power between young and control participants was not significantly different anymore after matching for *movement speed*, even without correction for multiple comparisons (trials in each group: 1470.8 ± 2.1, M1: p = 0.25, PMC: p = 0.15).

Relation of high gamma power, structure and clinical scores

High gamma power in stroke patients from either motor region was not significantly related to the structural disconnection of M1 or PMC (ChaCo) (M1: p = 0.47, PMC: p = 0.79) or the structural integrity of the CST (CST-FA) (M1: p = 0.93, PMC: p = 0.86). Furthermore, there was no

significant relation between high gamma power in either motor region and residual motor function assessed by clinical scores (UEFM: M1: p = 0.74, PMC: p = 0.47; ARAT: M1: p = 0.09, PMC: p = 0.32; BBT: M1: p = 0.48, PMC: p = 0.75; NHPT: M1: p = 0.25, PMC: p = 0.64; grip force: M1: p = 0.81, PMC: p = 0.98; key grip force: M1: p = 0.82, PMC: p = 0.84).

Relation of high gamma power and movement speed

Movement speed significantly related to high gamma power in linear mixed-effects models in the young and control participant groups in both motor regions, as trials with higher *movement speed* were associated with greater high gamma power (Fig. 4, Supplementary Fig. 3, young participants: M1: p_{cor} < 0.001, PMC: p_{cor} < 0.001; control participants: M1: p_{cor} < 0.001, PMC: p_{cor} = 0.002; corrected for six comparisons). In stroke patients, the association of *movement speed* and high gamma power showed a trend in the same direction, however, did not reach significance (M1: p_{cor} = 0.14, PMC: p_{cor} = 0.14; corrected for six comparisons).

Figure 3 Spatial distribution and group differences of high gamma power. (**A**) Group mean high gamma power (0 to 0.6 s, 60 to 90 Hz) overlayed on a standard brain. Colour indicates the percentage increase in high gamma power compared to baseline. Activity of participants that performed the task with the right hand was flipped across hemispheres. High gamma activity is evident mainly in motor regions contralateral to movement. (**B**) Distribution of participant mean high gamma power in M1 and PMC. Asterisks indicate significant group differences. (**C**) Group difference (control participants vs. stroke patients) in high gamma power when matched for *movement speed* and performing hand. Boxplots and dots show mean high gamma power of each participant over 1000 repetitions of the matching process. Significance marker: *$p < 0.05$

Figure 4 Relation between high gamma power in M1 and movement speed. Effect plots of the fixed effect *movement speed* in trial-level linear mixed-effects models for the relation between *movement speed* and high gamma power. Model results showed that *movement speed* significantly related to high gamma power in young and control participant groups. Asterisks indicate significance of the effect *movement speed* compared to a reduced model. HGP: high gamma power. Significance markers: ***$p < 0.001$

High gamma peak frequency

High gamma peak frequency significantly differed between young (M1: $\bar{f}_{peak}$ = 79.4 Hz) and control participants (M1: $\bar{f}_{peak}$ = 70.7 Hz), with significantly lower high gamma peak frequency in the older controls (Fig. 5A, M1: $p_{cor} < 0.001$, PMC: $p_{cor} < 0.001$; corrected for four comparisons). High gamma peak frequency was not significantly different between stroke patients (M1: $\bar{f}_{peak}$ = 71.9 Hz) and control participants (M1: $p_{cor} = 0.95$, PMC: $p_{cor} = 0.95$; corrected for four comparisons). Linear regression models showed

no significant relation between *movement speed* and high gamma peak frequency in any group after correction for multiple comparisons (stroke patients: M1: $p = 0.37$, PMC: $p = 0.048$, $p_{cor} = 0.22$; control participants: M1: $p = 0.76$, PMC: $p = 0.60$; young participants: M1: $p = 0.33$, PMC: $p = 0.07$; p_{cor} corrected for six comparisons).

Number of high gamma peaks

High gamma power time courses revealed differences of high gamma power evolution over time between the three groups (Fig. 6A). Young participants exhibited fewer high

Figure 5 Characteristics of high gamma peak frequency. Distribution of participant's mean high gamma peak frequency. Asterisks indicate significant group differences. We did not test for a difference between young participants and stroke patients. Significance markers: ***$p <$ 0.001

gamma peaks over the time course of movement compared to control participants, the difference between control participants and stroke patients was not significant (Fig. 6B, Supplementary Fig. 4A, control/young: M1: $p_{cor} =$ 0.02, PMC: $p_{cor} <$ 0.001; stroke/control: M1: $p_{cor} =$ 0.40, PMC: $p_{cor} =$ 0.58; corrected for four comparisons). The difference in number of high gamma peaks across groups, however, turned out to be a result of differences in *movement speed*. When modelling the number of high gamma peaks as a function of group and *movement speed*, we observed a significant effect of *movement speed* (Fig. 6D, Supplementary Fig. 4B, M1: $p_{cor} <$ 0.001, PMC: $p_{cor} =$ 0.006; corrected for two comparisons), but not group (M1: $p_{cor} =$ 0.98, PMC: $p_{cor} =$ 0.58; corrected for two comparisons). Hence, participants with faster *movement speed* had a significantly higher probability of having fewer high gamma peaks. Specifically, in M1, if *movement speed* was greater than 1.88 /s (corresponding to ~ 7.5 presses/s), the probability was highest that the participants would exhibit only one high gamma peak. In contrast, if *movement speed* was below 0.62 /s (corresponding to ~ 2.5 presses/s), the probability was highest that the participant would exhibit one individual high gamma peak for each button press.

Relation of high gamma power and improvement

Neither *improvement* across trials, defined as the percentage increase in *movement speed* from the current to the next trial (*trial-wise improvement*) nor *improvement* across blocks defined as the change from first block to best block (*block-wise improvement*) showed a significant association with high gamma power in any group or region after correction for multiple comparisons (*trial-wise*: stroke patients: M1: $p =$ 0.37, PMC: $p =$ 0.93; control participants: M1: $p =$ 0.08, PMC: $p =$ 0.10; young participants: M1: $p =$ 0.52, PMC: $p =$ 0.93; *block-wise*: stroke patients: M1: $p =$ 0.10, PMC: $p =$ 0.26; control participants: M1: $p =$ 0.28, PMC: $p =$ 0.23; young participants: M1: $p =$ 0.03, $p_{cor} =$ 0.17, PMC: $p =$ 0.12; p_{cor} corrected for six comparisons).

Discussion

Our study is the first to the best of our knowledge to characterize cortical high gamma oscillations during dexterous thumb movements with differing performance levels across age groups and in stroke patients. We demonstrated that faster *movement speed* strongly related to increased cortical high gamma power over motor areas during movement onset. In contrast, *improvement* of performance did not significantly relate to high gamma power. Moreover, with lower *movement speeds*, we observed distinct high gamma peaks for each movement, while in faster movements, high gamma activation became smeared over time. Even though *movement speed* showed the strongest association with high gamma power, stroke pathology resulted in an additional decrease of cortical motor high gamma power.

Decreased cortical high gamma power in stroke patients

In humans, studies on movement-related high gamma oscillations are restricted. Most reports on high gamma oscillations have utilized invasive ECoG recordings, which are not available in stroke patients. EEG, which can be measured easily in stroke patients, can smear the focal high gamma oscillations due to spatial blurring at the skull. MEG recordings, however, offer the opportunity to measure motor cortical high gamma oscillations non-invasively and with high spatial precision in patients with stroke pathology. In line with the literature in healthy cohorts, we observed high gamma oscillations in chronic stroke patients at the onset of contralateral movements, located over primary motor cortex and premotor cortices. We observed that chronic stroke patients, even though exhibiting largely very mild hand motor impairment, showed less cortical high gamma power compared to age-matched controls. This difference even remained significant after matching for *movement speed*. These results are paralleled by an existing study in rodents, in which the peri-infarct cortex of mice after stroke showed a deficit in low-gamma power under anesthesia.[35] Moreover, the disproportional decrease of high gamma power in stroke patients suggests that high gamma power may not purely reflect movement kinematics. This suggestion is further supported by findings of significant high gamma power in motor imagery.[36,37] The decrease of high gamma power was not related to clinical motor function scores or structural integrity of the corticospinal tract. Moreover, the structural disconnection of the primary motor cortex to other brain areas was not significantly related to the decrease of high gamma power.

Topological organization of high gamma oscillations

High gamma oscillations are more spatially focused compared to lower frequencies[7] and better aligned with

Figure 6 High gamma power time course. Data shown are from the voxel with maximal power in M1, results are similar in PMC (Supplementary Fig. 4). (**A**) Group-level high gamma power time courses. Dotted lines indicate average time points of button presses. (**B**) Distribution of number of individual high gamma peaks. Number of peaks was counted in participant mean of high gamma power time courses. Asterisk indicates the significant group difference. (**C**) Group-level time-frequency spectrograms. The time point of the first button press corresponds to 0 s. Dotted lines indicate average time points of button presses. Note the different scaling of the colour bar in the different groups. (**D**) Effect plot of the independent variable *movement speed* in a cumulative link model that models the relationship between number of high gamma peaks, *movement speed* and group with participant-level data from all three groups combined. The effect of *movement speed* was significant compared to a reduced model, the effect of group (not shown) was not significant. Significance markers: *p < 0.05, ***p < 0.001

somatotopy.[5,14] In comparison, modulation of beta activity (13 to 30 Hz) may reflect a more general, widespread activation signal during and after movement with only limited specificity for the limb component that was moved.[9] In simple motor tasks, high gamma oscillations have been described to be localized to M1.[5–7] With complex tasks, a more widespread cortical area is recruited[38], consisting ofPMd and PMv as part of the network for hand motor control.[39,40] PMC is connected with both the arm area of the primary motor cortex and the spinal cord[41] and is thought to be involved in connecting external cues with forthcoming movement.[39] A reason for the observed elevated high gamma power levels in PMC might be that our task required to constantly maximize motor performance. It may have thus involved a more complex network, including PMC, compared to simpler tasks like finger abductions or fist clenching.

The role of high gamma oscillations in motor control

High gamma oscillations are thought to be prokinetic in nature and have been shown to strongly synchronize during active but not passive movements.[8,42,43] At the onset of movement, in addition to the motor cortex, also basal ganglia and thalamus exhibit gamma band oscillations.[44–46] Invasive recordings from the basal ganglia in patients with deep brain stimulation[36,42,47–49] revealed that movement-related high gamma power from the contralateral subthalamic nucleus and globus pallidus interna increased with movement amplitude, movement velocity and force production, stressing its potential significance in movement coding. Also, links to behaviour have been established in movement disorders, in which less high gamma power in the contralateral subthalamic nucleus is associated with higher symptom severity in Parkinson's disease (PD).[49] Similarly, PD patients in the dopamine-depleted state show pathologically impaired movements and less high gamma power.[47,48] In line, dopaminergic medication, as a prokinetic drug, leads to increased high gamma power in PD patients.[50] Furthermore, patients suffering from hyperkinetic movements, such as medication-induced dyskinesias, present with pronounced high gamma power.[51] Recent studies have examined the subcortical-cortical communication in the gamma band and found that fast

reaction times were preceded by enhanced subcortical spike to cortical gamma phase coupling[52] underlining the role of gamma activity within the subcortical-cortical network for motor performance. In cortical invasive recordings, high gamma oscillations up to very high frequencies (600 to 1000 Hz) contained decoding information on position, velocity and predominantly movement speed.[13] Whereas single-unit activity in M1 is thought to mainly tune direction, larger neuronal populations as reflected by LFPs seem to represent movement speed.[16,17,53] Here, for the first time using non-invasive recording in humans, we demonstrate a clear association of cortical high gamma power (60 to 90 Hz) and *movement speed* in healthy individuals with different age ranges. In stroke patients, this association may potentially hold true in a similar way, but did not reach significance in our study.

Time course of high gamma oscillations

High gamma oscillations have been shown to be most pronounced around movement onset, subsiding over the course of movement even when force production is maintained.[8] Hence, high gamma oscillations seem to encode, at least to some extent, kinematic variables or cognitive processes rather than purely active contraction of muscles. With repetitive movements, a distinct burst of high gamma power has been described, with a tendency for individual gamma bursts to blur at a movement frequency of 3 Hz.[8] We showed that the observed inter-subject difference in high gamma peaks was driven by differences in *movement speed*, where with faster movements individual peaks blurred and the probability to exhibit only one high gamma peak increased. This contrasts with observations for power at lower frequencies: beta-band power modulations have still been observed at a movement frequency of 4 Hz.[54] Hence, for faster repeating movements, high gamma oscillations may code the overall motor action rather than every single movement separately.

Effects of high gamma peak frequency

The range of motor high gamma oscillations has been reported to be variable across studies, ranging from 75 to 100 Hz[7] over 60 to 90 Hz[8] to 55 to 375 Hz[47]. In the hippocampal CA1 region, studies have described a slow-, mid- and fast-frequency gamma band. It remains unclear whether distinct high gamma bands encode separate forms of motor processing. Over the lifespan, the high gamma frequency range has been reported to vary depending on age with an increase from the very young child to adulthood.[55,56] Here, we observe a decrease of the high gamma peak frequency with age with significantly lower peak frequency in the older controls (79.4 Hz versus 70.7 Hz). These findings are in contrast to a previous study, which found no association between high gamma

peak frequency and age in a large cohort with a simple motor task.[57] The reason for the differences might be the different analysis methods or discrepancies in motor tasks. Intra-participant gamma peak frequency has been reported to be consistent over recording sessions, over hemispheres and for movement of the same body part with differing movement vigour.[42,49,58] Also in our study, high gamma peak frequency did not significantly change with varying *movement speed*.

The role of high gamma power in motor skill acquisition

Recent work suggested that motor-related high gamma oscillations may also play a role in motor learning[59,60] and some studies have shown an increase in motor performance (faster reaction times, learning of movement parameters) after high gamma transcranial alternating current stimulation (tACS).[61–66] Further, a study analysing transcranial magnetic stimulation-induced intracortical inhibition reported a decrease in intracortical inhibition after high gamma tACS, correlating with improved motor learning scores.[67] In our study, the level of motor skill acquisition was similar in stroke patients compared to young and age-matched control participants, consistent with the notion that there is no significant difference in implicit motor learning in the affected hand after stroke compared to healthy controls.[68] Hence, there seems to be potential for further rehabilitation to improve motor skills even in a cohort of mildly impaired stroke patients. We investigated the relationship between high gamma power and *improvement* in *movement speed* over time and did not find a significant association. Our findings suggest that high gamma power does at least not strongly relate to motor skill acquisition in this dexterous finger movement task.

Limitations

Several limitations should be considered for this study. First, the sample size of stroke patients was limited. It is therefore possible that we missed findings because of restricted statistical power. Further, most stroke patients exhibited only mild impairment, limiting generalizability to more severe stroke patients. Nevertheless, we were able to show a highly significant relationship of movement speed and high gamma power within the two healthy cohorts and significant group differences between stroke patients and control participants. Second, as we specifically investigated thumb movements, the generalisability to movement of other parts of the upper extremity is unknown. Still, high gamma oscillations are elicited by movements of different body parts in a similar manner[5,7,8], which suggests that our findings may potentially generalize to other movement paradigms. Finally, we found no association between high gamma power and motor skill acquisition, while it is important to

note that our study misses any evaluation of long-term effects neglecting the learning-performance distinction.[69] We therefore are only able to draw conclusions on short-term motor skill acquisition.

In conclusion, characterizing cortical movement-related high gamma oscillations, we found a strong positive relationship between *movement speed* and high gamma oscillations within the motor network in younger and older healthy participants. For the first time, we quantified high gamma oscillations in stroke patients with motor impairment and observed a decrease in cortical motor high gamma power going beyond the association to *movement speed*. We suggest that enhancing cortical high gamma activity through tACS could potentially serve as a therapeutic approach to improve motor performance in stroke patients.

Acknowledgements

We thank Karin Reimann and Christiane Reißmann for assistance in data recording and participant recruitment and Sophie Grigutsch for valuable discussions.

Funding

This work was supported by the Medical Faculty of the University Medical Center Hamburg-Eppendorf ("Tandemförderung" to B.C.S. & F.Q.), the German Research Foundation (DFG; SFB 936 - 178316478, project Z2 to B.C.S. & F.Q.; SCHW 2023/2-1 to B.C.S.), the Gemeinnützige Hertie-Stiftung (Hertie Network of Excellence in Clinical Neuroscience, to F.Q.) and the Else Kröner-Fresenius-Stiftung (2020_EKES.16 to R.S.).

Competing interests

The authors report no competing interests.

Supplementary material

Supplementary material is available at bioRχiv online.

Data availability

The data that support the findings of this study are available from the corresponding author, upon reasonable request.

References

1. Buzsáki G, Wang XJ. Mechanisms of gamma oscillations. *Annu Rev Neurosci*. 2012;35:203-225. doi:10.1146/annurev-neuro-062111-150444
2. Ni J, Wunderle T, Lewis CM, Desimone R, Diester I, Fries P. Gamma-Rhythmic Gain Modulation. *Neuron*. 2016;92(1):240-251. doi:10.1016/j.neuron.2016.09.003
3. Misselhorn J, Schwab BC, Schneider TR, Engel AK. Synchronization of Sensory Gamma Oscillations Promotes Multisensory Communication. *eNeuro*. 2019;6(5):ENEURO.0101-19.2019. doi:10.1523/ENEURO.0101-19.2019
4. Herrmann CS, Munk MHJ, Engel AK. Cognitive functions of gamma-band activity: memory match and utilization. *Trends Cogn Sci*. 2004;8(8):347-355. doi:10.1016/j.tics.2004.06.006
5. Miller KJ, Leuthardt EC, Schalk G, et al. Spectral changes in cortical surface potentials during motor movement. *J Neurosci*. 2007;27(9):2424-2432. doi:10.1523/JNEUROSCI.3886-06.2007
6. Cheyne D, Bells S, Ferrari P, Gaetz W, Bostan AC. Self-paced movements induce high-frequency gamma oscillations in primary motor cortex. *NeuroImage*. 2008;42(1):332-342. doi:10.1016/j.neuroimage.2008.04.178
7. Crone NE, Miglioretti DL, Gordon B, Lesser RP. Functional mapping of human sensorimotor cortex with electrocorticographic spectral analysis. II. Event-related synchronization in the gamma band. *Brain*. 1998;121 (Pt 1:2301-2315. doi:10.1093/brain/121.12.2301
8. Muthukumaraswamy SD. Functional properties of human primary motor cortex gamma oscillations. *J Neurophysiol*. 2010;104(5):2873-2885. doi:10.1152/JN.00607.2010
9. Pfurtscheller G, Graimann B, Huggins JE, Levine SP, Schuh LA. Spatiotemporal patterns of beta desynchronization and gamma synchronization in corticographic data during self-paced movement. *Clin Neurophysiol*. 2003;114(7):1226-1236. doi:10.1016/s1388-2457(03)00067-1
10. Ulloa JL. The Control of Movements via Motor Gamma Oscillations. *Front Hum Neurosci*. 2021;15:787157. doi:10.3389/fnhum.2021.787157
11. Donoghue JP, Sanes JN, Hatsopoulos NG, Gaál G. Neural discharge and local field potential oscillations in primate motor cortex during voluntary movements. *J Neurophysiol*. 1998;79(1):159-173. doi:10.1152/jn.1998.79.1.159
12. Khorasani A, Heydari Beni N, Shalchyan V, Daliri MR. Continuous Force Decoding from Local Field Potentials of the Primary Motor Cortex in Freely Moving Rats. *Sci Rep*. 2016;6:35238. doi:10.1038/srep35238

13. Hammer J, Pistohl T, Fischer J, et al. Predominance of Movement Speed Over Direction in Neuronal Population Signals of Motor Cortex: Intracranial EEG Data and A Simple Explanatory Model. *Cereb Cortex*. 2016;26(6):2863-2881. doi:10.1093/cercor/bhw033

14. Quandt F, Reichert C, Hinrichs H, Heinze HJ, Knight RT, Rieger JW. Single trial discrimination of individual finger movements on one hand: A combined MEG and EEG study. *NeuroImage*. 2012;59(4):3316-3324. doi:10.1016/j.neuroimage.2011.11.053

15. Ball T, Demandt E, Mutschler I, et al. Movement related activity in the high gamma range of the human EEG. *NeuroImage*. 2008;41(2):302-310. doi:10.1016/j.neuroimage.2008.02.032

16. Anderson NR, Blakely T, Schalk G, Leuthardt EC, Moran DW. Electrocorticographic (ECoG) correlates of human arm movements. *Exp Brain Res*. 2012;223(1):1-10. doi:10.1007/s00221-012-3226-1

17. Wang PT, McCrimmon CM, King CE, et al. Characterization of electrocorticogram high-gamma signal in response to varying upper extremity movement velocity. *Brain Struct Funct*. 2017;222(8):3705-3748. doi:10.1007/s00429-017-1429-8

18. Pennati GV, Plantin J, Carment L, et al. Recovery and prediction of dynamic precision grip force control after stroke. *Stroke*. Published online 2020:944-951. doi:10.1161/STROKEAHA.119.026205

19. Gerloff C, Bushara K, Sailer A, et al. Multimodal imaging of brain reorganization in motor areas of the contralesional hemisphere of well recovered patients after capsular stroke. *Brain*. 2006;129(Pt 3):791-808. doi:10.1093/brain/awh713

20. Quandt F, Bönstrup M, Schulz R, et al. The functional role of beta-oscillations in the supplementary motor area during reaching and grasping after stroke: A question of structural damage to the corticospinal tract. *Hum Brain Mapp*. 2019;40(10):3091-3101. doi:10.1002/hbm.24582

21. Oostenveld R, Fries P, Maris E, Schoffelen JM. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput Intell Neurosci*. 2011;2011. doi:10.1155/2011/156869

22. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing; 2022. https://www.R-project.org/

23. Gross J, Kujala J, Hamalainen M, Timmermann L, Schnitzler A, Salmelin R. Dynamic imaging of coherent sources: Studying neural interactions in the human brain. *Proc Natl Acad Sci U S A*. 2001;98(2):694-699. doi:10.1073/pnas.98.2.694

24. Holmes CJ, Hoge R, Collins L, Woods R, Toga AW, Evans AC. Enhancement of MR Images Using Registration for Signal Averaging: *J Comput Assist Tomogr*. 1998;22(2):324-333. doi:10.1097/00004728-199803000-00032

25. Nolte G. The magnetic lead field theorem in the quasi-static approximation and its use for magnetoencephalography forward calculation in realistic volume conductors. *Phys Med Biol*. 2003;48(22):3637-3652. doi:10.1088/0031-9155/48/22/002

26. Fan L, Li H, Zhuo J, et al. The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cereb Cortex*. 2016;26(8):3508-3526. doi:10.1093/cercor/bhw157

27. Yushkevich PA, Piven J, Hazlett HC, et al. User-guided 3D active contour segmentation of anatomical structures: significantly improved efficiency and reliability. *NeuroImage*. 2006;31(3):1116-1128. doi:10.1016/j.neuroimage.2006.01.015

28. Kuceyeski A, Maruta J, Relkin N, Raj A. The Network Modification (NeMo) Tool: elucidating the effect of white matter integrity changes on cortical and subcortical structural connectivity. *Brain Connect*. 2013;3(5):451-463. doi:10.1089/brain.2013.0147

29. Cieslak M, Cook PA, He X, et al. QSIPrep: an integrative platform for preprocessing and reconstructing diffusion MRI data. *Nat Methods*. 2021;18(7):775-778. doi:10.1038/s41592-021-01185-5

30. Gorgolewski K, Burns CD, Madison C, et al. Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in python. *Front Neuroinformatics*. 2011;5:13. doi:10.3389/fninf.2011.00013

31. TractSeg: Automatic White Matter Bundle Segmentation - Git Hub. Accessed September 29, 2023. https://github.com/MIC-DKFZ/TractSeg

32. Wasserthal J, Neher P, Maier-Hein KH. TractSeg - Fast and accurate white matter tract segmentation. *NeuroImage*. 2018;183:239-253. doi:10.1016/j.neuroimage.2018.07.070

33. MRtrix3 - GitHub. Accessed September 29, 2023. https://github.com/MRtrix3/mrtrix3/releases/tag/3.0.3

34. Maris E. Statistical testing in electrophysiological studies. *Psychophysiology*. 2012;49(4):549-565. doi:10.1111/j.1469-8986.2011.01320.x

35. Hazime M, Alasoadura M, Lamtahri R, et al. Prolonged deficit of low gamma oscillations in the peri-infarct cortex of mice after stroke. *Exp Neurol*. 2021;341:113696. doi:10.1016/j.expneurol.2021.113696

36. Fischer P, Pogosyan A, Cheeran B, et al. Subthalamic nucleus beta and gamma activity is modulated depending on the level of imagined grip

force. *Exp Neurol*. 2017;293:53-61.
doi:10.1016/j.expneurol.2017.03.015

37. Miller KJ, Schalk G, Fetz EE, den Nijs M, Ojemann JG, Rao RPN. Cortical activity during motor execution, motor imagery, and imagery-based online feedback. *Proc Natl Acad Sci U S A*. 2010;107(9):4430-4435. doi:10.1073/pnas.0913697107

38. Donoghue JP, Sanes JN. Motor areas of the cerebral cortex. *J Clin Neurophysiol*. 1994;11(4):382-396.

39. Chouinard PA, Paus T. The primary motor and premotor areas of the human cerebral cortex. *Neuroscientist*. 2006;12(2):143-152. doi:10.1177/1073858405284255

40. Rizzolatti G, Fogassi L, Gallese V. Motor and cognitive functions of the ventral premotor cortex. *Curr Opin Neurobiol*. 2002;12(2):149-154. doi:10.1016/s0959-4388(02)00308-2

41. Dum RP, Strick PL. The origin of corticospinal projections from the premotor areas in the frontal lobe. *J Neurosci*. 1991;11(3):667-689. doi:10.1523/JNEUROSCI.11-03-00667.1991

42. Brücke C, Huebl J, Schönecker T, et al. Scaling of movement is related to pallidal γ oscillations in patients with dystonia. *J Neurosci*. 2012;32(3):1008-1019. doi:10.1523/JNEUROSCI.3860-11.2012

43. Liu X, Wang S, Yianni J, et al. The sensory and motor representation of synchronized oscillations in the globus pallidus in patients with primary dystonia. *Brain*. 2008;131(Pt 6):1562-1573. doi:10.1093/brain/awn083

44. Androulidakis AG, Kühn AA, Chen CC, et al. Dopaminergic therapy promotes lateralized motor activity in the subthalamic area in Parkinson's disease. *Brain*. 2007;130(2):457-468. doi:10.1093/BRAIN/AWL358

45. Kempf F, Brücke C, Salih F, et al. Gamma activity and reactivity in human thalamic local field potentials. *Eur J Neurosci*. 2009;29(5):943-953. doi:10.1111/j.1460-9568.2009.06655.x

46. Muthukumaraswamy SD. Temporal dynamics of primary motor cortex γ oscillation amplitude and piper corticomuscular coherence changes during motor control. *Exp Brain Res*. 2011;212(4):623-633. doi:10.1007/s00221-011-2775-z

47. Anzak A, Tan H, Pogosyan A, et al. Subthalamic nucleus activity optimizes maximal effort motor responses in Parkinson's disease. *Brain*. 2012;135(Pt 9):2766-2778. doi:10.1093/brain/aws183

48. Joundi RA, Brittain JS, Green AL, Aziz TZ, Brown P, Jenkinson N. Oscillatory activity in the subthalamic nucleus during arm reaching in Parkinson's disease. *Exp Neurol*. 2012;236(2):319-326. doi:10.1016/J.EXPNEUROL.2012.05.013

49. Lofredi R, Neumann WJ, Bock A, et al. Dopamine-dependent scaling of subthalamic gamma bursts with movement velocity in patients with Parkinson's disease. *eLife*. 2018;7:e31895. doi:10.7554/eLife.31895

50. Litvak V, Eusebio A, Jha A, et al. Movement-related changes in local and long-range synchronization in parkinson's disease revealed by simultaneous magnetoencephalography and intracranial recordings. *J Neurosci*. 2012;32(31):10541-10553. doi:10.1523/JNEUROSCI.0767-12.2012

51. Swann NC, de Hemptinne C, Miocinovic S, et al. Gamma Oscillations in the Hyperkinetic State Detected with Chronic Human Brain Recordings in Parkinson's Disease. *J Neurosci*. 2016;36(24):6445-6458. doi:10.1523/JNEUROSCI.1128-16.2016

52. Fischer P, Lipski WJ, Neumann WJ, et al. Movement-related coupling of human subthalamic nucleus spikes to cortical gamma. *eLife*. 2020;9:e51956. doi:10.7554/eLife.51956

53. Mehring C, Rickert J, Vaadia E, Cardosa de Oliveira S, Aertsen A, Rotter S. Inference of hand movements from local field potentials in monkey motor cortex. *Nat Neurosci*. 2003;6(12):1253-1254. doi:10.1038/nn1158

54. Toma K, Mima T, Matsuoka T, et al. Movement Rate Effect on Activation and Functional Coupling of Motor Cortical Areas. *J Neurophysiol*. 2002;88(6):3377-3385. doi:10.1152/jn.00281.2002

55. Cheyne D, Jobst C, Tesan G, Crain S, Johnson B. Movement-related neuromagnetic fields in preschool age children. *Hum Brain Mapp*. 2014;35(9):4858-4875. doi:10.1002/hbm.22518

56. Johnson B, Jobst C, Al-Loos R, He W, Cheyne D. Individual differences in motor development during early childhood: An MEG study. *Dev Sci*. 2020;23(5):e12935. doi:10.1111/desc.12935

57. Bardouille T, Bailey L. Evidence for age-related changes in sensorimotor neuromagnetic responses during cued button pressing in a large open-access dataset. *NeuroImage*. 2019;193:25-34. doi:10.1016/J.NEUROIMAGE.2019.02.065

58. Cheyne D, Ferrari P. MEG studies of motor cortex gamma oscillations: evidence for a gamma "fingerprint" in the brain? *Front Hum Neurosci*. 2013;7:575. doi:10.3389/fnhum.2013.00575

59. Akkad H, Dupont-Hadwen J, Kane E, et al. Increasing human motor skill acquisition by driving theta-gamma coupling. *eLife*. 2021;10:e67355. doi:10.7554/eLife.67355

60. Nowak M, Zich C, Stagg CJ. Motor Cortical Gamma Oscillations: What Have We Learnt and Where Are We Headed? *Curr Behav Neurosci Rep*. 2018;5(2):136-142. doi:10.1007/s40473-018-0151-z

61. Bologna M, Guerra A, Paparella G, et al. Transcranial Alternating Current Stimulation Has Frequency-Dependent Effects on Motor Learning in Healthy Humans. *Neuroscience*. 2019;411:130-139. doi:10.1016/j.neuroscience.2019.05.041

62. Guerra A, Bologna M, Paparella G, et al. Effects of Transcranial Alternating Current Stimulation on Repetitive Finger Movements in Healthy Humans. *Neural Plast*. 2018;2018:4593095. doi:10.1155/2018/4593095

63. Joundi RA, Jenkinson N, Brittain JS, Aziz TZ, Brown P. Driving oscillatory activity in the human cortex enhances motor performance. *Curr Biol*. 2012;22(5):403-407. doi:10.1016/j.cub.2012.01.024

64. Santarnecchi E, Biasella A, Tatti E, Rossi A, Prattichizzo D, Rossi S. High-gamma oscillations in the motor cortex during visuo-motor coordination: A tACS interferential study. *Brain Res Bull*. 2017;131:47-54. doi:10.1016/j.brainresbull.2017.03.006

65. Sugata H, Yagi K, Yazawa S, et al. Modulation of Motor Learning Capacity by Transcranial Alternating Current Stimulation. *Neuroscience*. 2018;391:131-139. doi:10.1016/j.neuroscience.2018.09.013

66. Spooner RK, Wilson TW. Spectral specificity of gamma-frequency transcranial alternating current stimulation over motor cortex during sequential movements. *Cereb Cortex*. 2023;33(9):5347-5360. doi:10.1093/cercor/bhac423

67. Nowak M, Hinson E, van Ede F, et al. Driving Human Motor Cortical Oscillations Leads to Behaviorally Relevant Changes in Local GABAA Inhibition: A tACS-TMS Study. *J Neurosci*. 2017;37(17):4481-4492. doi:10.1523/JNEUROSCI.0098-17.2017

68. Kal E, Winters M, van der Kamp J, et al. Is Implicit Motor Learning Preserved after Stroke? A Systematic Review with Meta-Analysis. *PloS One*. 2016;11(12):e0166376. doi:10.1371/journal.pone.0166376

69. Kantak SS, Winstein CJ. Learning-performance distinction and memory processes for motor skills: a focused review and perspective. *Behav Brain Res*. 2012;228(1):219-231. doi:10.1016/j.bbr.2011.11.028