The five homologous CiaR-controlled Ccn sRNAs of *Streptococcus pneumoniae*

modulate Zn-resistance.

Nicholas R. De Lay[1,4#], Nidhi Verma[1], Dhriti Sinha[1], Abigail Garrett[2], Maximillian Osterberg[3],

Spencer Reiling[1], Daisy Porter[1], David P. Giedroc[3], and Malcolm E. Winkler[2]

[1]Department of Microbiology and Molecular Genetics, McGovern Medical School, University of

Texas Health Science Center, Houston, TX 77030, USA

[2]Department of Biology, Indiana University Bloomington, Bloomington, Indiana 47405

[3]Department of Chemistry, Indiana University, Bloomington, Bloomington, Indiana 47405

[4]MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences, University of

Texas Health Science Center, Houston, TX 77030, USA

Running Title: Loss of the Ccn sRNAs cause Zn hypersensitivity

Keywords: small RNA; post-transcriptional regulation; Zn homeostasis; Mn homeostasis;

CiaRH

Correspondence should be sent to Nicholas R. De Lay (Tel: +1 (713) 500-6293; Email:

nicholas.r.delay@uth.tmc.edu)

ABSTRACT

Zinc is a vital transition metal for *Streptococcus pneumoniae*, but is deadly at high concentrations. Zn intoxication of *S. pneumoniae* results from a deficiency in Mn, which is required for key metabolic enzymes and defense against oxidative stress. Here, we report our identification and characterization of the function of the five homologous, CiaRH-regulated Ccn sRNAs in controlling *S. pneumoniae* virulence and metal homeostasis. We show that deletion of all five *ccn* genes (*ccnA*, *ccnB*, *ccnC*, *ccnD*, and *ccnE*) from *S. pneumoniae* strains D39 (serotype 2) and TIGR4 (serotype 4) causes Zn hypersensitivity and an attenuation of virulence in a murine invasive pneumonia model. We provide evidence that addition of Zn disproportionately impairs Mn uptake by the $\Delta ccnABCDE$ mutants. Consistent with a response to Mn starvation, expression of genes encoding the CzcD Zn exporter and the Mn-independent ribonucleotide reductase, NrdD-NrdG, were increased in the $\Delta ccnABCDE$ mutant relative to its isogenic *ccn*$^+$ parent strain. The growth inhibition by Zn that occurs as the result of loss of the *ccn* genes is rescued by supplementation with Mn or Oxyrase™, a reagent that removes dissolved oxygen. Lastly, we found that the Zn-dependent growth inhibition of the $\Delta ccnABCDE$ strain was not altered by deletion of *sodA*, whereas the *ccn*$^+$ $\Delta sodA$ strain phenocopied the $\Delta ccnABCDE$ strain. Overall, our results indicate that the Ccn sRNAs have a crucial role in preventing oxidative stress in *S. pneumoniae* during exposure to excess Zn by modulating Mn uptake.

AUTHOR SUMMARY

Zn and Mn are essential micronutrients for many bacteria, including *Streptococcus pneumoniae*. While Zn performs vital structural or catalytic roles in certain proteins, in excess, Zn can inhibit Mn uptake by *S. pneumoniae* and displace, but not functionally replace Mn from key enzymes including superoxide dismutase A (SodA). Here, we show that the Ccn small regulatory RNAs

promote *S. pneumoniae* Mn uptake and resistance to the oxidative stress. Furthermore, we

demonstrate that these small regulatory RNAs modulate the ability of *S. pneumoniae* to cause

invasive pneumonia. Altogether, these findings reveal a new layer of regulation of *S. pneumoniae*

Zn and Mn homeostasis and suggest that there are factors in addition to known transporters that

modulate intracellular Mn levels.

INTRODUCTION

Small regulatory RNAs have been established as fundamental regulators of gene expression in

bacteria and are involved in controlling nearly every aspect of bacterial physiology, metabolism,

and behavior (*1-3*). Two basic classes of small regulatory RNAs have been identified and

characterized, those that control gene expression by directly interacting with transcripts via

hydrogen bonding between complementary or wobble base-pairs and others that indirectly

effect transcript abundance by titrating an RNA or DNA-binding protein (*4, 5*). Interactions

between the former class of riboregulators, henceforth referred to as sRNAs, and their cognate

target transcripts can result in changes in their transcription, translation, and/or stability

depending on many factors include the sequence, accessibility, structure, and location of the

sRNA binding site. One of the most facile modes of regulation discovered involves the sRNA

binding within or adjacent to the translation initiation region blocking the 16S rRNA within the

30S ribosomal subunit from base-pairing with the complementary Shine-Delgarno sequence, or

ribosome binding site, within the mRNA. Many other elegant mechanisms of sRNA-based gene

regulation have been uncovered (*6-8*). While a large amount of progress has been made

towards understanding the contribution of sRNAs to the response of Gram-negative bacteria

such as *Escherichia coli* to internally and externally derived stresses, environmental cues, and

host interactions, much less headway has been achieved in understanding the functions of

sRNAs in Gram-positive bacteria, particularly, *Streptococcus pneumoniae.*

The Gram-positive, ovoid diplococcus *S. pneumoniae* is a leading cause of lower respiratory

infection morbidity and mortality worldwide resulting in nearly 2 million deaths per year (*9*). We

and others have discovered 100s of putative sRNAs in *S. pneumoniae* (*10-15*), but the functions

of almost all of them remains a mystery. Among the first sRNAs identified in *S. pneumoniae*

were the five homologous Ccn sRNAs (CcnA, CcnB, CcnC, CcnD, and CcnE), which were

shown to be transcribed in response to activation of the CiaRH two-component system (*15, 16*);

expression of the CiaRH two-component systems is induced by penicillin and sialic acid (*17,*

18). Shortly after the discovery of the five Ccn sRNAs, Tsui, Mukerjee (Sinha), et al

demonstrated that CcnA negatively regulates competence and the *comCDE* mRNA encoding

the precursor of the competence stimulating peptide and the two-component system that

responds to this signal and activates competence (*15*). Schnorpfeil et al then formally

demonstrated that the five Ccn sRNAs negatively regulate competence by base-pairing with the

comCDE mRNA (*19*). Other likely targets post-transcriptionally regulated by the Ccn sRNAs

were identified in that study including mRNAs encoding components of a galactose transporter

(*spd_0090*), a formate-nitrate transporter (*nirC*), branched-chain amino acid transporter (*brnQ*)

and a toxin *(shetA)*, but direct regulation of these targets by the Ccn sRNAs has not yet been

shown (*19*). One of these five homologous sRNAs, CcnE, has also been implicated in *S.*

pneumoniae virulence in a murine invasive pneumonia model (*12*).

Here, we report our discovery of a role for the five Ccn sRNAs in controlling *S. pneumoniae*

virulence and Zn resistance. Specifically, we show that deletion of the genes encoding the five

Ccn sRNAs attenuates the virulence of *S. pneumoniae* strains D39 and TIGR4 in a murine

invasive pneumonia model. Additionally, we show that loss of the Ccn sRNAs leads *S.*

pneumoniae D39 and TIGR4 to become hypersensitive to Zn toxicity. Altogether, our results

indicate that the Ccn sRNAs promote Mn uptake leading into increased Zn resistance due to an

increase in the amount of active superoxide dismutase A (SodA).

RESULTS

The Ccn sRNAs are important for *S. pneumoniae* pathogenesis. Work from a prior study (*12*)

indicated that deletion of one of the five Ccn sRNA genes (*ccnE*) reduced *S. pneumoniae* serotype

4 strain TIGR4 virulence in a murine invasive pneumoniae model. In that study, the authors also

discovered by Tn-seq that transposon insertions in *ccnE* reduced *S. pneumoniae* strain TIGR4

fitness in murine lungs, whereas transposon insertions in *ccnA* had no significant impact on its

fitness in the murine lung, nasopharynx, or blood. To determine the contribution of the Ccn sRNAs

to the virulence of the *S. pneumoniae* serotype 2 strain D39 in a murine invasive pneumonia

model, we initially made single deletions of *ccnA*, *ccnB*, *ccnC, ccnD*, or *ccnE* and a quintuple

deletion of all five *ccn* genes in a strain background harboring the *rpsL1* allele (IU1781). Next, we

evaluated the growth of these strains in BHI broth at 37°C in an atmosphere of 5% CO_2. Neither

deletion of any single Ccn sRNA gene or all five of them had any significant effect on growth rate,

although the growth yield of the Δ*ccnE* strain was slightly reduced (Figs. 1A, S1A, and S1C). We

then determined the consequence of these deletions on *S. pneumoniae* pathogenicity in a murine

invasive pneumonia model (see Materials and Methods). While removal of any single *ccn* gene

had no significant impact on its virulence in mice (Fig. S2), deletion of all five *ccn* genes attenuated

S. pneumoniae strain D39 pathogenicity increasing median survival time from 43 h to 67 h (Fig.

2A).

To confirm that the *ccn* genes are generally important for *S. pneumoniae* virulence and is not

an attribute specific to strain D39, we also deleted all five *ccn* genes from a serotype 4 TIGR4

strain harboring the *rpsL1* allele (NRD10220) and measured the impact of these deletions on its

growth and virulence using the same murine invasive pneumonia model. As shown in Fig. 1D,

deletion of all five *ccn* genes did have an effect on growth rate of *S. pneumoniae* TIGR4 in BHI

broth incubated at 37°C with an atmosphere of 5% CO_2. To validate the effect of the *ccnABCDE*

deletion on *S. pneumoniae* TIGR4 growth, we reconstructed the quintuplet deletion strain from

scratch, but this time we subsequently replaced the mutant *rpsL* allele conferring streptomycin

resistance that aided in construction of the unmarked deletions of the *ccn* genes back to the wild

type *rpsL* allele. Next, we tested the growth of that strain and its parental *ccn*[+] TIGR4 strain in BHI

broth at 37°C under an atmosphere of 5% CO_2. Again, we saw a substantial reduction in growth

rate when all five *ccn* genes were deleted (Fig. S3A). The *ccnABCDE* deletion also resulted in a

marked attenuation of *S. pneumoniae* TIGR4 virulence increasing the survival rate of ICR outbred

mice from 0% to 50% (Fig. 2B). Two of the mice that survived infection with the TIGR4

Δ*ccnABCDE* strain had no detectable bacteria in the blood and the other two mice had 1000 and

2750 CFUs per mL of blood, respectively, which was far below 10^7 bacteria found in moribund

mice that were infected with the *ccn*[+] parent strain. Our results show that the *ccn* genes are

important for *S. pneumoniae* pathogenesis.

The Ccn sRNAs impact expression of Zn and Mn-related genes. To discover a basis for the

defect in *S. pneumoniae* virulence caused by the deletion of the five *ccn* genes, we compared

global gene expression by high throughput RNA-sequencing (RNA-seq) between *S. pneumoniae*

strain D39 or TIGR4 and the derived Δ*ccnABCDE* mutant strains grown to exponential phase

(OD_{620} between 0.15 and 0.2) in BHI broth at 37°C in an atmosphere of 5% CO_2. In the *S.*

pneumoniae D39 strain background, the *ccnABCDE* deletion resulted in down-regulation of 3

genes and up-regulation of 113 genes by 2-fold or more ($P_{adj} < 0.05$) (Table S3). In contrast,

deletion of the *ccn* genes from the TIGR4 strain resulted in down-regulation of 25 genes and up-

regulation of 97 genes by 2-fold or greater ($P_{adj} < 0.05$) (Table S4). 37 genes were up-regulated

by 2-fold ($P_{adj}<0.05$) in the *ccnABCDE* deletion strain in both the D39 and TIGR4 backgrounds

(Table 1); among these differentially expressed genes were iron uptake system genes (*piuB*,

piuC, *piuD*, and *piuA*), a Zn-responsive ECF (energy-coupling factor) transport gene

SPD_1267/SP_1438, and *czcD* encoding a Zn/Cd exporter that provides Zn and Cd resistance.

To validate our RNA-seq data, we first measured abundance of *piuB*, *spd_1267*, and *czcD*

transcripts in RNA samples isolated for the RNA-seq experiment from *S. pneumoniae* strain D39

and derived Δ*ccnABCDE* strain by reverse transcriptase droplet digital PCR (RT-ddPCR).

Consistent with our RNA-seq data the *piuB, spd_1267,* and *czcD* transcripts were up-regulated

by 3.47, 10.5, and 1.93-fold respectively in the Δ*ccnABCDE* strain compared to its parental D39

strain grown in BHI broth (Fig. 3A, B, and C). Using RT-ddPCR analysis of the RNA samples

isolated from exponential cultures of *S. pneumoniae* TIGR4 and derived Δ*ccnABCDE* mutant

strain grown in BHI broth at 37°C under an atmosphere of 5% CO_2, we also observed a 2.45

increase in the abundance of the *czcD* mRNA in the *ccn* mutant as compared to its parental strain

(Fig. 3E). Altogether, these data suggested to us that removal of the *ccn* genes from *S.*

pneumoniae was leading to an increase in the intracellular abundance of Zn relative to Mn, and

to cope with this stress, the *ccn* mutant strain was increasing expression of the CzcD Zn exporter.

Table 1: Genes significantly, differentially expressed between a Δ*ccnABCDE* and *ccn*[+]

strain in both the *S. pneumoniae* D39 and TIGR4 background during exponential growth

in BHI broth[a]

D39 locus tag	Gene	Known or predicted function	D39 fold change	TIGR4 fold change
SPD_0025		tRNA-specific adenosine-34 deaminase	84.3	144
SPD_0027	*dut*	deoxyuridine 5'-triphosphate nucleotidohydrolase	3.52	4.39
SPD_0028		hypothetical protein	3.80	3.40
SPD_0029	*radA*	DNA repair protein	3.55	2.80
SPD_0090		galactose ABC transport protein	2.09	2.00
SPD_0104		aggregation-promoting factor	2.69	2.27
SPD_0222	*gpmB1*	phosphoglycerate mutase family protein	25.0	22.9
SPD_0243	*uppS*	undecaprenyl diphosphate synthase	5.97	4.42
SPD_0244	*cdsA*	phosphatidate cytidylyltransferase	5.50	4.49
SPD_0245	*eep*	intramembrane protease	5.57	4.19
SPD_0246	*proS*	prolyl-tRNA synthetase	5.91	4.68
SPD_0247	*bglA*	6-phospho-β-glucosidase	3.57	3.35
SPD_0308	*clpL*	ATP-dependent protease subunit	13.7	2.71
SPD_0460	*dnaK*	protein chaperone	3.97	2.01
SPD_0501	*licT*	β-glucoside operon antiterminator	2.91	4.77
SPD_0502	*bglF*	β-glucoside PTS transporter subunit	3.07	5.65

SPD_0503	bglA-2	6-phospho- β-glucosidase	2.57	4.79
SPD_0615	glnH3	degenerate glutamine ABC transporter subunit	11.6	4.05
SPD_0616	glnQ3	glutamine ABC transporter subunit	8.90	3.07
SPD_0617	glnP3b	glutamine ABC transporter subunit	11.1	3.63
SPD_0618	glnP3a	glutamine ABC transporter subunit	11.8	2.98
SPD_0775		acetyltransferase	3.29	2.71
SPD_1045		degenerate DUF3884 domain protein	4.73	3.16
SPD_1046	lacG-2	6-phospho-β-galactosidase	3.56	2.92
SPD_1267		ECF transporter subunit	11.1	2.14
SPD_1638	czcD	Cd/Zn exporter	2.69	2.66
SPD_1649	piuB	Fe uptake transporter subunit	5.13	2.43
SPD_1650	piuC	Fe uptake transporter subunit	4.45	2.01
SPD_1651	piuD	Fe uptake transporter subunit	4.22	2.15
SPD_1652	piuA	Fe uptake transporter subunit	4.38	2.25
SPD_1748	pneA2	lantibiotic peptide	2.19	2.16
SPD_1749	lanM	lanthionine biosynthesis protein	2.49	2.29
SPD_1750	wrbA	FAD-dependent flavoprotein	3.00	2.69
SPD_1751		hypothetical protein	2.56	3.16
SPD_1752	clyB	toxin secretion ABC transporter	3.58	3.22
SPD_1753		epidermin leader peptide processing serine protease	2.44	2.87
SPD_1932	malP	malodextrin phosphorylase	2.58	2.58

aRNA extraction and mRNA-seq analyses were performed as described in *Materials and Methods*.

RNA was prepared from cultures of strains IU1781 (D39 *rpsL1*), NRD10176 (D39 *rpsL1*

Δ*ccnABCDE*), NRD10220 (TIGR4 *rpsL1*), and NRD10266 (TIGR4 *rpsL1* Δ*ccnABCDE*) (Table S1

and S2). Fold changes (2.0-fold cut-off) and adjusted P-values (Pval <0.05) are based on three

independent biological replicates.

Absence of the *ccn* genes causes S. *pneumoniae* to become hypersensitive to Zn. If the

absence of the *ccn* genes from *S. pneumoniae* leads to an imbalance of transition metals with

higher levels of Zn relative to Mn, then we would expect that increasing the concentration of Zn

present in the medium would disproportionately impair the growth of the Δ*ccnABCDE* mutant

relative to the isogenic *ccn*+ strain. Previous studies have indicated that Becton-Dickinson (BD)

BHI broth typically contains ~20 µM Zn and 200 nM Mn (*20, 21*). We first compared growth of

strain D39 and derived Δ*ccnA*, Δ*ccnB*, Δ*ccnC*, Δ*ccnD*, Δ*ccnE*, and Δ*ccnABCDE* strains in BHI

broth alone or supplemented with 0.2 mM Zn at 37°C under an atmosphere of 5% CO_2. No

significant difference was observed in growth rate between strain D39 and derived Δ*ccnA*, Δ*ccnB*,

Δ*ccnC*, and Δ*ccnD* mutant strains in BHI in the presence or absence of 0.2 mM added Zn (Fig.

S1), although the growth yield for the *ccnE* mutant was lower in BHI in the presence and absence

of Zn. As noted above, growth the strain D39 and derived Δ*ccnABCDE* mutant strain was similar

in BHI broth alone (Fig. 1A.) In contrast, the absence of all five *ccn* genes led to an obvious

impairment in growth rate in BHI supplemented with 0.2 mM Zn (Fig. 1B) This growth deficiency

relative to the *ccn*⁺ parental strain was also observed for the Δ*ccnABCDE* strain when Zn was

increased in BHI broth to 0.4 mM (Fig. 1C). Consistently, addition of Zn at 0.4 mM severely

reduced the growth rate of the *ccn*⁺ D39 strain. Reintroduction of *ccnC* and *ccnD* expressed from

their native promoters at ectopic loci partially ameliorated the growth defect of the Δ*ccnABCDE*

mutant strain in BHI supplemented with 0.2 mM or 0.4 mM Zn, whereas insertion of copies of

ccnA, ccnB, and *ccnD* with their native promoter at the *bgaA* and CEP loci completely corrected

the growth deficiency of this quintuple *ccn* mutant (Figs. 1B and 1C) under these growth

conditions. To verify that the Zn hypersensitivity caused by the deletion of all five *ccn* genes was

not specific to the serotype 2 strain D39, we also tested the effect of the quintuple *ccn* deletion on

the growth of the serotype 4 TIGR4 strain in BHI broth supplemented with Zn. Consistent with our

results observed for strain D39, deletion of the Ccn sRNA genes from TIGR4 led to a growth

impairment in BHI when Zn was added at a final concentration of 0.2 or 0.4 mM (Figs. 1D-F and

S3). Curiously, Zn at the highest concentration used had less of an effect on TIGR4 growth than

it did on the D39 strain. Overall, these results indicate that Ccn sRNAs promote *S. pneumoniae*

Zn tolerance.

The Zn-hypersensitivity that occurs in the absence of the Ccn sRNA genes is due to a reduction in cellular Mn levels. *S. pneumoniae* is a Mn-centric bacteria encoding several Mn-requiring enzymes including superoxide dismutase (SodA), a capsule regulatory kinase (CpsB), phosphoglucomutase (Pgm), phosphopentomutase (DeoB), a cell division regulating phosphatase (PhpP), an aerobic ribonucleotide reductase (NrdEF), pyruvate kinase (PyK), and lactate dehydrogenase (Ldh). Mis-metalation of these Mn-dependent enzymes by Zn, which inhibits their enzymatic activity (*22, 23*), can occur when the internal ratio of Zn-to-Mn is high. Additionally, the substrate binding component of the PsaBCA Mn ATP binding cassette (ABC) type transporter, the only known Mn importer in *S. pneumoniae*, has been shown to bind Zn tightly, blocking Mn uptake (*20, 24*). Our RNA-seq and RT-ddPCR data above indicated that expression of the CzcD Zinc exporter, which is expressed in response to high levels of Zn relative to Mn (*25, 26*), is up-regulated in *S. pneumoniae* strains lacking the *ccn* genes (Fig. 3C and E). Based on these results and the published data mentioned above, we hypothesized that Zn-hypersensitivity caused by the removal of all five *ccn* genes from the *S. pneumoniae* genome is due to reduced Mn uptake. If this postulate is correct, then the Zn-dependent growth inhibition that occurs when the *S. pneumoniae* Δ*ccnABCDE* mutant strain is grown in BHI broth supplemented with 0.2 mM Zn should be rescued by inclusion of an equimolar amount of Mn into the medium. As shown in Fig. 4, the growth impairment of the Δ*ccnABCDE* mutant of *S. pneumoniae* D39 or TIGR4 strain in BHI broth with 0.2 mM Zn is cured by addition of 0.2 mM Mn consistent with our model.

To directly test whether or not the levels of transition metals are perturbed in strains lacking the *ccn* genes, we measured total cell associated transition metals in *S. pneumoniae* strain D39, derived Δ*ccnABCDE* mutant, and the Δ*ccnABCDE* strain complemented with *ccnA*, *ccnB*, and *ccnD* by inductively coupled plasma-mass spectrometry (ICP-MS). During exponential growth (OD_{620} of ~0.2) in BHI broth alone or supplemented with 0.2 mM Zn, there was no significant difference in Zn abundance among these strains (Fig 5A). Similarly, Mn levels were comparable

among all tested strains when grown to exponential phase in BHI broth alone (Fig. 5B). However,

there was a modest, but statistically significant (P = 0.039), 22% reduction in total cell associated

Mn in the ΔccnABCDE mutant compared to its parental ccn⁺ strain, which was partially

complemented by ccnA, ccnB, and ccnD. Thus, our evidence that Mn supplementation eliminated

the growth deficiency of the ccn⁻ strain caused by excess Zn and that the amount of Mn associated

with this mutant strain was reduced compared to the ccn⁺ strain when exposed to a high Zn

concentration suggest that the Ccn sRNAs are important for preserving Mn homeostasis, when

S. pneumoniae encounters a Zn-rich environment.

Oxidative stress due to reduced levels of active superoxide dismutase A contributes to the

Zn hypersensitivity of the *S. pneumoniae* strain lacking the Ccn sRNAs. To discover the

molecular basis for the Zn hypersensitivity caused by loss of the *ccn* genes, we turned to an RNA-

seq based approach. Briefly, we compared transcript abundance in RNA isolated from *S.*

pneumoniae strain D39 and derived ΔccnABCDE strain grown to exponential phase (OD$_{620}$ of

~0.2) at 37°C under an atmosphere of 5% CO_2 in BHI broth supplemented with 0.2 mM Zn. Similar

to our RNA-seq performed with these strains in the absence of Zn supplementation, we observed

a 2.33-fold increase in expression of the CzcD Zn exporter specifying mRNA and a 9.5-fold

increase in the Spd_1267 Zn-responsive ECF-type transporter producing mRNA in the

ΔccnABCDE mutant compared to its *ccn⁺* parent strain (Tables 2 and S5). Interestingly, we also

saw a 1.90-fold decrease (P_{adj} = 1.81x10^{-43}) in expression of the *sodA* mRNA, encoding

superoxide dismutase A, in the ΔccnABCDE mutant strain, which was just below our arbitrary

two-fold cutoff (Table S5). We subsequently measured the relative abundance of these transcripts

by RT-ddPCR and northern blot analysis (Figs. 3B, C, and D) and were able to confirm our RNA-

seq results indicating up-regulation of *spd_1267* and *czcD* mRNAs and down-regulation of the

sodA mRNA abundance when the *ccn* genes were deleted from *S. pneumoniae*. This result was

intriguing to us since a prior study found that Mn starvation of *S. pneumoniae* cells due to

exposure to high concentrations of Zn relative to Mn led to a reduction in the transcription of *sodA*

and a reduction in superoxide dismutase activity (*22*). Furthermore, Eijkelkamp *et* al discovered

that deletion of *sodA* had no significant impact on *S. pneumoniae* growth under Mn replete

conditions, but was vital for growth in media containing a high Zn-to-Mn ratio (*22*).

Table 2: Genes significantly, differentially expressed between a *S. pneumoniae* D39 and

derived Δ*ccnABCDE* strain in both BHI alone or supplemented with Zn[a]

D39 locus tag	Gene	Known or predicted function	Fold change (BHI)	Fold change (BHI+Zn)
SPD_0025		tRNA-specific adenosine-34 deaminase	84.3	49.0
SPD_0027	*dut*	deoxyuridine 5'-triphosphate nucleotidohydrolase	3.52	3.76
SPD_0028		hypothetical protein	3.80	3.02
SPD_0029	*radA*	DNA repair protein	3.55	2.95
SPD_0080	*pavB*	cell wall surface anchor family protein	6.69	6.32
SPD_0163		DNA binding protein	2.00	2.07
SPD_0222	*gpmB1*	phosphoglycerate mutase family protein	25.0	21.7
SPD_0243	*uppS*	undecaprenyl diphosphate synthase	5.97	7.61
SPD_0244	*cdsA*	phosphatidate cytidylyltransferase	5.50	7.77
SPD_0245	*eep*	intramembrane protease	5.57	8.59
SPD_0246	*proS*	prolyl-tRNA synthetase	5.91	9.65
SPD_0247	*bglA*	6-phospho-β-glucosidase	3.57	4.73
SPD_0277	*bglA-1*	6-phospho-β-glucosidase	3.53	2.73
SPD_0279	*celB*	cellobiose PTS transporter subunit	5.05	2.97
SPD_0308	*clpL*	ATP-dependent protease subunit	13.7	9.55
SPD_0350	*vraT*	cell wall-active antibiotic response protein	2.19	2.62
SPD_0351	*vraS*	two-component system histidine kinase	2.29	2.74
SPD_0352	*vraR*	two-component system response regulator	2.31	2.70
SPD_0353	*alkD*	degenerate DNA alkylation repair enzyme	2.11	2.67
SPD_0354	*alkD*	degenerate DNA alkylation repair enzyme	2.37	2.69
SPD_0458	*hrcA*	heat inducible transcription repressor	3.62	3.69
SPD_0459	*grpE*	heat shock protein	3.77	3.87
SPD_0460	*dnaK*	protein chaperone	3.97	3.82
SPD_0461	*dnaJ*	protein chaperone	3.50	3.52
SPD_0474	*blpZ*	immunity protein	2.40	2.05
SPD_0501	*licT*	β-glucoside operon antiterminator	2.91	5.26
SPD_0502	*bglF*	β-glucoside PTS transporter subunit	3.07	4.65
SPD_0503	*bglA-2*	6-phospho- β-glucosidase	2.57	3.75
SPD_0537		putative Zn-dependent protease	2.07	2.21

SPD_0615	glnH3	degenerate glutamine ABC transporter subunit	11.6	18.0
SPD_0616	glnQ3	glutamine ABC transporter subunit	8.90	16.8
SPD_0617	glnP3b	glutamine ABC transporter subunit	11.1	15.8
SPD_0618	glnP3a	glutamine ABC transporter subunit	11.8	15.1
SPD_0681		hypothetical protein	2.82	5.45
SPD_0701	ciaR	two-component response regulator	2.72	2.56
SPD_0702	ciaH	two-component histidine kinase	2.80	3.01
SPD_0775		acetyltransferase	3.29	3.61
SPD_0803		putative phage shock protein C		
SPD_0804		ABC transporter ATP-binding protein	2.28	3.01
SPD_0805		ABC transporter permease protein	2.43	3.15
SPD_0913		extracellular protein	3.39	3.31
SPD_0938		degenerate TN5252 relaxase	9.35	5.06
SPD_0940	rrfD	UDP-N-acetyl-D-mannosaminouronic acid dehydrogenase.	3.95	5.31
SPD_0942		hypothetical protein	2.25	2.41
SPD_0943		hypothetical proein	2.41	2.43
SPD_0944		nodulation protein L	2.24	2.38
SPD_0946		hypothetical protein	2.16	3.27
SPD_0947		hypothetical protein	2.69	3.97
SPD_0948	nikS	nikkomycin biosynthesis protein	3.73	4.29
SPD_0949		N-acetylneuraminate synthase	2.38	4.85
SPD_0950	mefE	macrolide ABCE transporter subunit	2.44	3.99
SPD_1045		degenerate DUF3884 domain protein	4.73	6.81
SPD_1046	lacG-2	6-phospho-b-galactosidase	3.56	7.28
SPD_1047	lacE-2	lactose PTS transporter subunit	4.21	6.33
SPD_1049	lacT	β-glucoside bgl operon antiterminator	3.23	3.48
SPD_1114		hypothetical protein	13.5	5.37
SPD_1267		ECF transporter subunit	11.1	9.53
SPD_1297	pdxS	pyridoxal 5'-phosphate synthase	2.02	2.04
SPD_1506	axe1	acetyl xylan esterase 1	3.62	2.68
SPD_1615		degenerate hypothetical protein	4.02	2.09
SPD_1638	czcD	Cd/Zn exporter	2.69	2.33
SPD_1709	groL	HSP60 family chaperone	2.58	2.38
SPD_1710	groES	HSP60 family chaperone	2.25	2.31
SPD_1716		hypothetical protein	2.56	5.62
SPD_1717		membrane protein	2.40	5.22
SPD_1718		LytR/AlgR family response regulator	2.44	4.58
SPD_1746		hypothetical protein	2.96	4.25
SPD_1747	pneA1	lantibiotic peptide	2.02	4.29
SPD_1748	pneA2	lantibiotic peptide	2.19	4.60
SPD_1749	lanM	lanthionine biosynthesis protein	2.49	2.37
SPD_1750	wrbA	FAD-dependent flavoprotein	3.00	2.85
SPD_1751		hypothetical protein	2.56	4.07
SPD_1752	clyB	toxin secretion ABC transporter	3.58	4.02
SPD_1753		epidermin leader peptide processing serine protease	2.44	3.00
SPD_1769		membrane protein	2.29	3.42

SPD_1932	malP	malodextrin phosphorylase	2.58	2.95
SPD_1933	malQ	4-α-glucanotransferase	2.76	2.77
SPD_1990		mannose PTS transporter subunit	2.01	13.8
SPD_1994	fucA	L-fuculose phosphate aldolase	2.35	8.69
SPD_2034	comFC	phosphororibosyltransferase domain protein	32.7	14.1
SPD_2035	comFA	DNA transporter ATPase	8.88	10.2
SPD_2068	htrA	serine protease	2.79	2.13
SPD_2069	parB	chromosome partitioning protein	3.04	2.88

aRNA extraction and mRNA-seq analyses were performed as described in *Materials and Methods*.

RNA was prepared from cultures of strains IU1781 (D39 *rpsL1*) and NRD10176 (D39 *rpsL1*

Δ*ccnABCDE*) grown to exponential phase in BHI alone or supplemented with 0.2 mM ZnSO$_4$

(Table S1 and S2). Fold changes (2.0-fold cut-off) and adjusted P-values (Pval <0.05) are based

on three independent biological replicates.

To initially examine whether the growth deficiency of the Δ*ccnABCDE* mutant relative to the

ccn$^+$ D39 strain was due in part to oxidative stress, we evaluated the impact of addition of

Oxyrase™, an enzyme mixture that removes molecular oxygen by reducing it to water, on the

growth of these strains in BHI broth alone or supplemented with 0.2 mM or 0.4 mM Zn (Fig 6A, B,

and C) under an atmosphere of 5% CO$_2$. Once again, we observed that deletion of *ccnA*, *ccnB*,

ccnC, *ccnD*, and *ccnE* from *S. pneumoniae* strain D39 had no significant impact on growth in BHI

alone. However, under these growth conditions, the addition of Oxyrase™ reduced the growth

rate of both the *ccn*$^+$ and ccn$^-$ strains to a similar extent (Fig. 6A). As we anticipated, addition of

Oxyrase™ to BHI supplemented with Zn (0.2 mM) improved the growth rate of the Δ*ccnABCDE*

strain to that observed for the *ccn*$^+$ D39 parent strain (Fig. 6B). Interestingly, addition of Oxyrase™

improved the growth rate of both strains in BHI with 0.4 mM Zn and eliminated any growth

differences between them (Fig. 6C). Finally, we examined the contribution of *sodA* to the growth

of *S. pneumoniae* D39 and derived Δ*ccnABCDE* mutant strain. In BHI broth alone or

supplemented with 0.2 mM Zn, deletion of *sodA* reduced the growth rate of the *ccn*$^+$ strain, but

had no significant reduction in growth rate of the ΔccnABCDE mutant strain (Fig. 7). Based on

these results, we concluded that the amount of functional SodA was negligible in the *S.*

pneumoniae strain lacking the Ccn sRNAs and thus, deleting *sodA* did not significantly impact its

growth, whereas this deletion does substantially impair growth of the isogenic *ccn*+ strain.

DISCUSSION

High density Tn-seq experiments performed more than a decade ago revealed that sRNAs

play a crucial role in regulating *S. pneumoniae* virulence including its ability to colonize the blood,

nasopharynx, and lungs of its host (*12*). While this discovery in itself may not be surprising, it is

astonishing that very little progress has been made towards understanding the functions of these

sRNAs given their importance in governing *S. pneumoniae* pathogenesis. Here, we investigated

the contribution of the five homologous Ccn sRNAs to *S. pneumoniae* pathogenesis and gene

regulation. Not only have we confirmed their crucial role in pneumococcal disease progression

(Fig. 2), but also discovered their extensive functions in regulating gene expression and Zn

resistance. Specifically, we found that exposure to relatively high, yet host-relevant, Zn

concentrations (0.2 mM) disproportionally inhibited growth (Fig 1, 4, 6, and 7) and modestly

reduced total cell associated Mn levels (Fig. 5) of *S. pneumoniae* strains lacking genes for the

five Ccn sRNAs. This Zn-dependent growth inhibition caused the *ccnABCDE* deletion was

completely alleviated by addition of Mn (Fig. 4) or Oxyrase™ (Fig. 6), which removes molecular

oxygen by reducing it to water. Furthermore, deletion of *sodA*, encoding the Mn-dependent

superoxide dismutase A, from *S. pneumoniae* resulted in a Zn-dependent growth inhibition

indistinguishable from that observed for the ΔccnABCDE strain; however, the same deletion had

no impact on the growth of the ΔccnABCDE strain (Fig. 7). Altogether, these results indicate that

the Ccn sRNAs promote Mn uptake and availability resulting in an increased abundance of active

SodA, which improves the growth of *S. pneumoniae* in a Zn-rich environment due to greater

protection from damaging reactive oxygen species.

How do the Ccn sRNAs promote an increased concentration of free intracellular Mn^{2+} cations?

Potential mechanisms through which the Ccn sRNAs could increase the concentration of

available Mn inside *S. pneumoniae* cells include that the Ccn sRNAs (1) negatively regulate

expression of a Mn exporter, (2) positively regulate expression of a Mn importer, (3) reduce

production of an intracellular protein or other factor that effectively chelates Mn, or (4) decrease

synthesis of a cellular component that would otherwise restrict access of Mn to the periplasmic

Mn binding component of the Mn importer. The main Mn exporter of *S. pneumoniae* is MntE (Fig.

8), as deletion of the encoding gene leads to accumulation of total cell associated Mn (*21, 27*).

MgtA, designated as a Ca efflux protein, appears to also export Mn, but has a very limited role in

this process (Fig. 8) (*28*). Neither MntE or MgtA were up-regulated in either *S. pneumoniae* strain

D39 or TIGR4 when the *ccn* genes were deleted (Tables S3 S4, and S5) making it unlikely that

the Ccn sRNAs increase intracellular Mn levels by down-regulating expression of these Mn

exporter genes. Additionally, we were unable to identify strong Ccn sRNA binding sites in the

translation initiation region of *mntE* or *mgtA*, which suggests that these sRNAs do not regulate

translation of these transcripts. Finally, if the Ccn sRNAs increase total cell-associated Mn levels

by down-regulating MntE expression, then we would expect that deletion of *mntE* would suppress

the Zn hypersensitivity of the *S. pneumoniae* Δ*ccnABCDE* mutant; however, this did not occur

(Fig S4A and C).

An alternative possibility is that the Ccn sRNAs promote Mn uptake by positively regulating

expression of the *psaBCA* operon encoding the only known Mn importer in *S. pneumoniae* (Fig.

8)(*29, 30*). Localized to the inner membrane, PsaB is the ATP binding component whereas PsaC

is the permease of this ABC-type transporter. PsaA, the substrate binding component, is located

in the periplasm, where it binds Mn. Once again, in our RNA-seq experiments, we did not observe

a decrease in expression of the *psaBCA* operon when the *ccn* genes were deleted from *S.*

pneumoniae strain D39 or TIGR4 (Tables S3, S4, and S5) indicating that the Ccn sRNAs do not

positively regulate expression of this Mn importer. Furthermore, if this was the case, then we

would expect that deletion of *psaR* encoding the repressor of the *psaBCA* operon (*31, 32*) might

suppress the Zn-dependent growth inhibition of the *S. pneumoniae* Δ*ccnABCDE* mutant;

however, we did not observe this (Fig. S4B and D).

While it remains possible that the Ccn sRNAs regulate production of a factor that chelates

intracellular Mn, we did not observe the up-regulation of any *known* Mn-binding proteins via RNA-

seq. A final possibility that we surmised could be occurring is that the Ccn sRNAs negatively

impact the production or architecture of a structure that could restrict Mn uptake by limiting the

diffusion of Mn into the periplasm, where it could be transported by the PsaBCA Mn importer into

the cytoplasm. For example, a physical structure that might serve as a permeability barrier to Mn

is the polysaccharide capsule that surrounds *S. pneumoniae* diplococci. To test this possibility,

we evaluated the impact of deleting the capsule biosynthetic genes on the Zn hypersensitivity of

the *S. pneumoniae* Δ*ccnABCDE* mutant. Interestingly, we observed that the *cps2ABCDETFGH*

deletion completely suppressed the Zn hypersensitivity of an *S. pneumoniae* strain lacking the

ccn genes (Fig. S4C and F). While the *ccnABCDE* deletion did not cause *S. pneumoniae* to

express the capsule biosynthetic genes at a higher level (Tables S3 S4, and S5), we did observe

increased expression of genes in the Leloir pathway (*galK* and *galT2*), which is involved in the

synthesis of a precursor for capsular polysaccharide synthesis (UDP-glucose), in the Δ*ccnABCDE*

mutant as compare to its isogenic *ccn*[+] parental D39 strain when grown in BHI broth

supplemented with 0.2 mM $ZnSO_4$ (Table S5 and Fig. 8). Notably, increased production of

capsular polysaccharide precursors has been shown to increase capsule production (*33, 34*). In

future studies, we plan on rigorously testing this model that the Ccn sRNAs decrease capsule

synthesis by reducing expression of genes involved in synthesizing capsular polysaccharide

precursors leading to an increase in Mn uptake (Fig. 8). Moreover, we intend to evaluate the

molecular mechanism by which the Ccn sRNAs regulate expression of these genes.

MATERIALS AND METHODS

Bacterial strains and growth conditions. Bacterial strains used in this study were derived

from encapsulated *S. pneumoniae* serotype 2 strain D39W (*14*) and TIGR4 and are listed in Table

S1. Strains were grown on plates containing trypticase soy agar II (modified; Becton-Dickinson

[BD]) and 5% (vol/vol) defribrinated sheep blood (TSAII BA) at 37°C in an atmosphere of 5% CO_2,

and liquid cultures were statically grown in BD brain heart infusion (BHI) broth at 37°C in an

atmosphere of 5% CO_2. Bacteria were inoculated into BHI broth from frozen cultures or single,

isolated colonies. For overnight cultures, strains were first inoculated into a 17-mm-diameter

polystyrene plastic tube containing 5 mL of BHI broth and then serially diluted by 100-fold into

four tubes; these cultures were then grown for 10 to 16 h. Cultures with an optical density at 620

nm (OD_{620}) of 0.1 to 0.4 were diluted to a starting OD_{620} between 0.002 and 0.005 in 5 mL of BHI

broth in 16-mm glass tubes. Growth was monitored by measuring OD_{620} using a Genesys 30

visible spectrophotometer (ThermoFisher Scientific). For antibiotic selections, TSAII BA plates

and BHI cultures were supplemented with 250 µg kanamycin per mL, 150 µg streptomycin per

mL, or 0.3 µg erythromycin per mL.

Construction and confirmation of mutants. Mutant strains were constructed by

transformation of competent *S. pneumoniae* D39 and TIGR4 derived strains with linear PCR

amplicons as described previously (*35, 36*). DNA amplicons containing antibiotic resistance

markers were synthesized by overlapping fusion PCR using the primers listed in Table S2.

Competence was induced in *S. pneumoniae* D39 or TIGR4 derived cells with CSP-1 or CSP-2,

respectively, synthetic competence stimulatory peptide. Unmarked deletions of the target genes

were constructed using the *kan*^{R}-*rpsL*$^+$ (Janus cassette) allele replacement method as described

previously (*37*). In the first step, the Janus cassette containing *rpsL+* allele and a kanamycin

resistance gene was used to disrupt target genes in an *rpsL1* or *rpsLK56T* (Str^R) strain

background, and transformants were selected for kanamycin resistance and screened for

streptomycin sensitivity. In the second step, the Janus cassette was eliminated by replacement

with a PCR amplicon lacking antibiotic markers and the resulting transformants were selected for

streptomycin resistance and screened for kanamycin sensitivity. Freezer stocks were made of

each strain from single colonies isolated twice on TSAII BA plates containing antibiotics listed in

Table S1. All strains were validated by PCR amplification and sequencing.

RNA extraction. To isolate RNA, strains were grown in 30 mL of BHI starting at an OD_{620} =

0.002 in 50 mL conical tubes. RNA was extracted from exponentially growing cultures of IU1781

(D39), NRD10220 (TIGR4), and their derived isogenic mutants lacking all five *ccn* genes,

NRD10176 (D39 Δccn) and NRD10266 (TIGR4 Δccn), at OD_{620} ≈ 0.2 using the FastRNA Pro Blue

Kit (MP Bio) according to the manufacturer's guidelines. Briefly, cells were collected by

centrifugation at 16,000 x g for 8 min at 4^{o}C. Cell pellets were resuspended in 1 mL of RNApro™

solution (MP Bio) and processed five-times for 40 sec at 400 rpm in a BeadBug™ homogenizer

(Benchmark Scientific). Cell debris was removed by centrifugation at 16,000 x g for 5 min at 4^{o}C.

After mixing 300 μL of chloroform with the supernatant, the aqueous and organic layers were

separated by centrifugation at 16,000 x g for 5 min at 4^{o}C. RNA was precipitated with 500 μL of

ethanol at -80^{o}C overnight. After collecting the precipitated RNA by centrifugation at 16,000 x g

for 15 min at 4^{o}C, the pellet was washed once with 75% ethanol and suspend in DEPC-treated

water. The amount and purity of all RNA samples isolated were assessed by NanoDrop

spectroscopy (Thermo Fisher).

Library preparation and mRNA-seq. cDNA libraries were prepared from total RNA Azenta Life
Sciences. Briefly, total RNA was subjected to rRNA-depletion using the FastSelect 5S/16S/23S
rRNA depletion kit for bacteria. Libraries were the generated with NEBNext Ultra™ II Directional
RNA Library Prep Kit. 150 bp paired-end read sequencing was performed using an Illumina
HiSeq4000 sequencer.

RNA-seq analysis. The raw sequencing reads were quality and adapter trimmed using Cutadapt
version 4.1 with a minimum length of 18 nucleotides. The trimmed reads were then mapped on
the *Streptococcus pneumoniae* D39 (Genbank CP000410) genome using Bowtie2 (*38*). HTseq
version 2.0.2 was used to generate read counts for the genes (*39*). Differential gene expression
was identified using the program DESeq2 with default parameters (*40*). Primary data from the
mRNA-seq analyses were submitted to the NCBI Gene Expression Omnibus (GEO) and have
the accession number GSE246655.

Reverse transcriptase-droplet digital PCR (RT-ddPCR) analysis. RT-ddPCR was performed
as described previously (*41*). Isolated RNA was treated with DNase (TurboDNase, Ambion) as
per manufacturer's instructions. Next, RNA (1 µg) was reverse transcribed using Superscript III
reverse transcriptase (Invitrogen) with random hexamers. RT and No RT control (NRT) sample
were utilized. These samples were diluted $1:10^1$, $1:10^2$, $1:10^3$, or $1:10^6$. Then, 2 µL of each diluted
RT and NRT sample was added to a 22 µL reaction mixture containing 11 µL of QX200™ ddPCR
™ Evagreen Supermix (Bio-Rad) and 1.1 µL of each 2 µM ddPCR primers (Table S6). A single
no template control (NTC) was included for each ddPCR primer pair used. Reactions were
performed using at least three independent biological replicates. Droplets were generated using
the QX200 Automated Droplet Generator (Bio-Rad), and end-point PCR was carried out using a
C1000 Touch™ thermal cycler (Bio-Rad) following the manufacturer's instructions. Quantification

of PCR-positive and PCR-negative droplets in each reaction, which provides absolute

quantification of the target transcript, was performed using the QX200 Droplet Reader (Bio-Rad).

This data was analyzed with QuantaSoft software (Bio-Rad) to determine the concentration of

each target expressed as copies per µL. Transcript copies were normalized to *tuf* mRNA (internal

control) and fold changes of transcripts corresponding to target genes in different mutants relative

to the WT parent were calculated. Statistical analysis was performed using Student's t-test with

GraphPad Prism version 10.0.0.

Northern blot analysis. Northern blotting was conducted as previously described (*13*). Briefly,

3 μg of isolated RNA was fractionated on 10% polyacrylamide gels containing 7% urea by

electrophoresis at 55 V and subsequently, transferred to a Zeta-probe membrane (Bio-Rad) using

a Trans-Blot SD semidry transfer apparatus (Bio-rad) at 4 mA per cm^2 with a maximum of 400

mA for 50 min. RNA was then UV-crosslinked to the membrane with a Spectroline UV crosslinker

with the "optimal crosslink" setting. 5'-Biotinylated probes were hybridized to the membrane

overnight at 42°C in ULTRAhyb (Ambion) hybridization buffer. Blots were developed according to

the BrightStar BioDetect kit protocol (Ambion), imaged with the ChemiDoc MP imager (Bio-Rad),

and individual band intensities were quantified using Image Lab software version 5.2.1 (Bio-Rad).

Signal intensities for each transcript were normalized to that of 5S rRNA, which served as a

loading control. Graphs of normalized abundance of each transcript for three biological replicates

were produced using GraphPad Prism version 10.0.0.

Inductively coupled plasma-mass spectrometry (ICP-MS) analysis. ICP-MS sample

preparation was based on a previous publication (*42*), with some modifications. Bacteria were

grown in BHI broth at 37°C with 5% CO_2 to OD_{620} = 0.2. Five mL of culture was centrifuged for 10

min in pre-chilled tubes at 12,400 x *g* at 4°C, and cell pellets were resuspended in 1.0 mL of

chilled BHI supplemented with 1 mM nitrilotriacetic acid (Sigma-Aldrich) (pH 7.2). Samples were

centrifuged for 7 min at 16,100 x g at 4°C, and supernatants were removed. Pellets were

centrifuged for an additional 3 min in the same way, and residual supernatant was removed. Cell

pellets were washed twice with centrifugation in the same way with 1.0 mL of chilled 10X PBS

(1.3M NaCl, 88mM Na_2HPO_4, 12mM NaH_2PO_4, pH 7.0) that had been treated with chelator.

Similar ICP-MS results were obtained when cells were washed with 1X PBS instead of 10X PBS.

Chelated PBS was prepared by mixing with 1% (wt/vol) Chelex-100 (BioRad), which was rotated

overnight at 4°C and passed through a 0.22 μm Steriflip (MilliporeSigma) filter. Before the last

centrifugation in PBS, samples were split into two 0.475 mL aliquots for ICP-MS analysis and

protein quantification. After removal of supernatants, pellets for ICP-MS were dried for 15 h at low

heat in an evaporative centrifuge and stored at -80°C until being processed for ICP-MS analysis.

Pellets for protein determination were suspended in in 100 μL of lysis buffer (1% (wt/vol) SDS

[Sigma], 0.1% w/v Triton X-100 [Mallinckrodt]) and stored at -80°C. Protein amount was

determined by using the DC protein assay (BioRad). For ICP-MS analysis, dried samples were

resuspended in 400 μL of 30% trace metal grade HNO_3 (Sigma). Samples and a 30% HNO_3 blank

were heated at 95° C for 10 min with shaking at 500 rpm. Samples were then diluted 100-fold to

a final volume of 3.0 mL with 2.5% HNO_3 containing the Pure Plus Internal Standard Mix (100

ppb, PerkinElmer). Samples were analyzed using an Agilent 8800 QQQ ICP-MS operating with

hydrogen (^{55}Mn detection) or helium (^{66}Zn detection) as collision gases to remove possible

interferences. ^{45}Sc or ^{72}Ge were used as internal references. Zn^{2+} and Mn^{2+} amounts were

calculated from standard curves made with Pure Plus Multi-Element Calibration Standard 3 (0.5-

100ppb, PerkinElmer). Metals amounts detected in the 30% HNO_3 blank were subtracted from all

samples. Metal amounts in samples were normalized relative to total protein amounts in the

matched samples.

Mouse models of infection: All procedures were approved in advance by UTHealth Animal

Welfare Committee and carried out as previously described (*41*). Male ICR mice (21-24 g; Envigo)

were anaesthetized by inhaling 4 to 5 % isoflurane. A total of 8 mice were intranasally inoculated

with 10^7 CFU of a specific *S. pneumoniae* strain suspended in 50 μL of 1 X PBS prepared from

cultures grown in BHI broth at 37°C in an atmosphere of 5% CO_2 to $OD_{620} \approx 0.1$. Mice were

monitored visually at 4 to 8 h intervals, and isoflurane-anesthetized moribund mice were

euthanized by cardiac puncture-induced exsanguination followed by cervical dislocation. Kaplan-

Meir survival curves and log-rank tests were generated using GraphPad Prism 10.0.0 software.

SUPPLEMENTAL MATERIALS

Supplemental Materials are available for this article.

ACKNOWLEDGEMENTS

This work was supported by NIGMS grant T32GM131994 (to M.O.), NIGMS grant R35GM118157

(to D.P.G.), NIGMS grant R35GM131767 (to M.E.W.), McGovern Medical Startup funds, and

NIAID grant R21AI171771 (to N.R.D.).

REFERENCES

1. D. G. Mediati, S. Wu, W. Wu, J. J. Tree, Networks of Resistance: Small RNA Control of Antibiotic Resistance. *Trends Genet* **37**, 35-45 (2021).
2. R. Ghandour, K. Papenfort, Small regulatory RNAs in *Vibrio cholerae*. *Microlife* **4**, uqad030 (2023).
3. K. Papenfort, S. Melamed, Small RNAs, Large Networks: Posttranscriptional Regulons in Gram-Negative Bacteria. *Annual Review of Microbiology*, (2023).
4. J. Hor, G. Matera, J. Vogel, S. Gottesman, G. Storz, Trans-Acting Small RNAs and Their Effects on Gene Expression in *Escherichia coli* and *Salmonella enterica*. *EcoSal Plus* **9**, (2020).
5. C. Pourciau, Y. J. Lai, M. Gorelik, P. Babitzke, T. Romeo, Diverse Mechanisms and Circuitry for Global Regulation by the RNA-Binding Protein CsrA. *Frontiers in Microbiology* **11**, 601352 (2020).
6. M. G. Jorgensen, J. S. Pettersen, B. H. Kallipolitis, sRNA-mediated control in bacteria: An increasing diversity of regulatory mechanisms. *Biochim Biophys Acta Gene Regul Mech* **1863**, 194504 (2020).

7. L. Bossi, N. Figueroa-Bossi, P. Bouloc, M. Boudvillain, Regulatory interplay between small
RNAs and transcription termination factor Rho. *Biochim Biophys Acta Gene Regul Mech*
1863, 194546 (2020).
8. F. Ponath, J. Hor, J. Vogel, An overview of gene regulation in bacteria by small RNAs
derived from mRNA 3' ends. *FEMS Microbiology Reviews* **46**, (2022).
9. G. B. D. L. R. I. Collaborators, Estimates of the global, regional, and national morbidity,
mortality, and aetiologies of lower respiratory infections in 195 countries, 1990-2016: a
systematic analysis for the Global Burden of Disease Study 2016. *Lancet Infect Dis* **18**,
1191-1210 (2018).
10. P. Acebo, A. J. Martin-Galiano, S. Navarro, A. Zaballos, M. Amblar, Identification of 88
regulatory small RNAs in the TIGR4 strain of the human pathogen *Streptococcus*
pneumoniae. *RNA* **18**, 530-546 (2012).
11. R. Kumar *et al.*, Identification of novel non-coding small RNAs from *Streptococcus*
pneumoniae TIGR4 using high-resolution genome tiling arrays. *BMC Genomics* **11**, 350
(2010).
12. B. Mann *et al.*, Control of virulence by small RNAs in *Streptococcus pneumoniae*. *PLoS*
Pathogens **8**, e1002788 (2012).
13. D. Sinha *et al.*, Redefining the sRNA Transcriptome in *Streptococcus pneumoniae* Serotype
2 Strain D39. *Journal of Bacteriology*, (2019).
14. J. Slager, R. Aprianto, J. W. Veening, Deep genome annotation of the opportunistic human
pathogen *Streptococcus pneumoniae* D39. *Nucleic Acids Research* **46**, 9971-9989
(2018).
15. H. C. Tsui *et al.*, Identification and characterization of noncoding small RNAs in
Streptococcus pneumoniae serotype 2 strain D39. *Journal of Bacteriology* **192**, 264-279
(2010).
16. A. Halfmann, M. Kovacs, R. Hakenbeck, R. Bruckner, Identification of the genes directly
controlled by the response regulator CiaR in *Streptococcus pneumoniae*: five out of 15
promoters drive expression of small non-coding RNAs. *Molecular Microbiology* **66**, 110-
126 (2007).
17. K. Hentrich *et al.*, *Streptococcus pneumoniae* Senses a Human-like Sialic Acid Profile via
the Response Regulator CiaR. *Cell Host Microbe* **20**, 307-317 (2016).
18. P. D. Rogers *et al.*, Gene expression profiling of the response of *Streptococcus pneumoniae*
to penicillin. *J Antimicrob Chemother* **59**, 616-626 (2007).
19. A. Schnorpfeil *et al.*, Target evaluation of the non-coding csRNAs reveals a link of the two-
component regulatory system CiaRH to competence control in *Streptococcus*
pneumoniae R6. *Molecular Microbiology* **89**, 334-349 (2013).
20. F. E. Jacobsen, K. M. Kazmierczak, J. P. Lisher, M. E. Winkler, D. P. Giedroc, Interplay
between manganese and zinc homeostasis in the human pathogen *Streptococcus*
pneumoniae. *Metallomics* **3**, 38-41 (2011).
21. J. E. Martin, J. P. Lisher, M. E. Winkler, D. P. Giedroc, Perturbation of manganese
metabolism disrupts cell division in *Streptococcus pneumoniae*. *Molecular Microbiology*
104, 334-348 (2017).
22. B. A. Eijkelkamp *et al.*, Extracellular zinc competitively inhibits manganese uptake and
compromises oxidative stress management in *Streptococcus pneumoniae*. *PloS One* **9**,
e89427 (2014).
23. A. L. McFarland, N. Bhattarai, M. Joseph, M. E. Winkler, J. E. Martin, Cellular Mn/Zn Ratio
Influences Phosphoglucomutase Activity and Capsule Production in *Streptococcus*
pneumoniae D39. *Journal of Bacteriology* **203**, e0060220 (2021).
24. C. A. McDevitt *et al.*, A molecular mechanism for bacterial susceptibility to zinc. *PLoS*
pathogens **7**, e1002357 (2011).

25. J. E. Martin *et al.*, The zinc efflux activator SczA protects *Streptococcus pneumoniae*
serotype 2 D39 from intracellular zinc toxicity. *Molecular Microbiology* **104**, 636-651
(2017).
26. A. D. Ogunniyi *et al.*, Central role of manganese in regulation of stress responses,
physiology, and metabolism in *Streptococcus pneumoniae*. *Journal of Bacteriology* **192**,
4489-4497 (2010).
27. J. W. Rosch, G. Gao, G. Ridout, Y. D. Wang, E. I. Tuomanen, Role of the manganese efflux
system *mntE* for signalling and pathogenesis in *Streptococcus pneumoniae*. *Molecular*
Microbiology **72**, 12-25 (2009).
28. J. E. Martin *et al.*, A Mn-sensing riboswitch activates expression of a Mn^{2+}/Ca^{2+} ATPase
transporter in Streptococcus. *Nucleic Acids Research* **47**, 6885-6899 (2019).
29. A. Dintilhac, G. Alloing, C. Granadel, J. P. Claverys, Competence and virulence of
Streptococcus pneumoniae: Adc and PsaA mutants exhibit a requirement for Zn and Mn
resulting from inactivation of putative ABC metal permeases. *Molecular Microbiology* **25**,
727-739 (1997).
30. R. Novak, J. S. Braun, E. Charpentier, E. Tuomanen, Penicillin tolerance genes of
Streptococcus pneumoniae: the ABC-type manganese permease complex Psa.
Molecular Microbiology **29**, 1285-1296 (1998).
31. N. S. Jakubovics, A. W. Smith, H. F. Jenkinson, Expression of the virulence-related Sca
(Mn^{2+}) permease in *Streptococcus gordonii* is regulated by a diphtheria toxin
metallorepressor-like protein ScaR. *Molecular Microbiology* **38**, 140-153 (2000).
32. J. W. Johnston, D. E. Briles, L. E. Myers, S. K. Hollingshead, Mn^{2+}-dependent regulation of
multiple genes in *Streptococcus pneumoniae* through PsaR and the resultant impact on
virulence. *Infection and Immunity* **74**, 1171-1180 (2006).
33. C. L. Ventura, R. T. Cartee, W. T. Forsee, J. Yother, Control of capsular polysaccharide
chain length by UDP-sugar substrate concentrations in *Streptococcus pneumoniae*.
Molecular Microbiology **61**, 723-733 (2006).
34. M. B. Ayoola *et al.*, Polyamine Synthesis Effects Capsule Expression by Reduction of
Precursors in *Streptococcus pneumoniae*. *Frontiers in Microbiology* **10**, 1996 (2019).
35. S. Ramos-Montanez *et al.*, Polymorphism and regulation of the *spxB* (pyruvate oxidase)
virulence factor gene by a CBS-HotDog domain protein (SpxR) in serotype 2
Streptococcus pneumoniae. *Molecular Microbiology* **67**, 729-746 (2008).
36. K. E. Bruce, B. E. Rued, H. T. Tsui, M. E. Winkler, The Opp (AmiACDEF) Oligopeptide
Transporter Mediates Resistance of Serotype 2 *Streptococcus pneumoniae* D39 to
Killing by Chemokine CXCL10 and Other Antimicrobial Peptides. *Journal of Bacteriology*
200, (2018).
37. K. M. Kazmierczak, K. J. Wayne, A. Rechtsteiner, M. E. Winkler, Roles of rel_{Spn} in stringent
response, global regulation and virulence of serotype 2 *Streptococcus pneumoniae* D39.
Molecular Microbiology **72**, 590-611 (2009).
38. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**,
357-359 (2012).
39. S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**, 166-169 (2015).
40. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
41. D. Sinha, J. P. Frick, K. Clemons, M. E. Winkler, N. R. De Lay, Pivotal Roles for
Ribonucleases in *Streptococcus pneumoniae* Pathogenesis. *mBio*, e0238521 (2021).
42. Y. Fu *et al.*, A new structural paradigm in copper resistance in *Streptococcus pneumoniae*.
Nat Chem Biol **9**, 177-183 (2013).
617

FIGURES

Figure 1: Growth phenotypes of *S. pneumoniae* strains harboring deletion of the *ccn* genes.

Growth characteristics at 37°C under an atmosphere of 5% CO_2 in BHI broth alone (A, D) or with

0.2 mM (B, E) or 0.4 mM (C,F) $ZnSO_4$ of following strains: (A, B, C) IU781 (D39), NRD10176

(Δccn), NRD10396 ($\Delta ccn//ccnA^+B^+D^+$), and NRD10397 ($\Delta ccn//ccnC^+D^+$); (D, E, F) NRD10220

(TIGR4) and NRD10266 (Δccn). Each point on the graph represents the mean OD_{620} value from

three independent cultures. Error bars, which in some cases are too small to observe in the graph,

represent the standard deviation (SD).

Figure 2: Virulence phenotypes of *S. pneumoniae* strains harboring deletion of the *ccn* genes.

Survival curve of ICR outbred mice after infection with ~10^7 CFU in a 50 μL inoculum of the

following *S. pneumoniae* strains: (A) IU1781 (D39) and NRD10176 (Δ*ccn*); (C) NRD10220

(TIGR4) and NRD10266 (Δ*ccn*). Eight mice were infected per strain. Disease progression of

animals was monitored, the time at which animals reached a moribund state was recorded, and

these mice were subsequently euthanized as described in Materials and Methods. A survival

curve was generated from this data and analyzed by Kaplan-Meier statistics and log rank test to

determine P-values.

Figure 3: Loss of the *ccn* genes perturbs the expression of Zn and Mn stress associated genes

in *S. pneumoniae*. Abundance of *piuB* (A), *spd_1267* (B), *czcD* (C, E), and *sodA* (D) mRNAs was

determined by RT-ddPCR (A, B, and C) or northern blot analyses (D) as described in Materials

and Methods for strain IU1781 (D39) and derived Δ*ccnABCDE* mutant strain (NRD10176; Δ*ccn*)

grown to exponential phase (OD_{620} of ~0.2) in BHI broth alone (BHI) or supplemented with 0.2

mM Zn (BHI+Zn) at 37°C under an atmosphere of 5% CO_2. (E) Expression of *czcD* was

determined by RT-ddPCR analyses for NRD10220 (TIGR4) and derived Δ*ccnABCDE* mutant

strain NRD10266 (Δ*ccn*) grown to exponential phase in BHI Broth at 37°C under an atmosphere

of 5% CO_2. Transcript levels were normalized to *tuf* (A, B, C, and E) or 5S rRNA (D). Values

represent the mean of three independent cultures and error bars indicate SEM.

Figure 4: Mn supplementation eliminates the Zn dependent growth inhibition of *S. pneumoniae*

Δ*ccnABCDE* mutant. Growth characteristics at 37°C under an atmosphere of 5% CO_2 in BHI broth

alone (BHI) or with 0.2 mM $ZnSO_4$ (BHI+Zn), 0.2 mM $MnCl_2$ (BHI+Mn), or 0.2 mM $ZnSO_4$ and

$MnCl_2$ (BHI+Zn+Mn) of strains (A) IU1781 (D39 *ccn*+), (B) NRD10176 (D39 Δ*ccn*), (C), NRD10220

(TIGR4 *ccn*+), and (D) NRD10266 (TIGR4 Δ*ccn*). Each point on the graph represents the mean

OD_{620} value from three independent cultures. Error bars, which in some cases are too small to

observe in the graph, represent the standard deviation (SD).

Figure 5: Deletion of the *ccn* genes reduces total cell associated Mn, but not Zn levels. Total cell associated Zn (A) and Mn (B) was measured from cells harvested from cultures of IU781 (D39), NRD10176 (Δccn), and NRD10396 ($\Delta ccn//ccnA^+B^+D^+$) grown to exponential growth phase (OD_{620} of ~0.2) in BHI broth alone (BHI) or with 0.2 mM ZnS04 (BHI+Zn) by ICP-MS and normalized to protein amounts. Results represent the mean of 5 to 9 replicates with values from each replicate shown as a point. Error bars indicate SEM.

Figure 6: Depletion of O_2 abolishes the Zn hypersensitivity of the *S. pneumoniae* ΔccnABCDE

mutant. Growth characteristics at 37°C under an atmosphere of 5% CO_2 in BHI broth alone (A)

or with 0.2 mM (B) or 0.4 mM (C) $ZnSO_4$ of IU781 (D39) and NRD10176 (Δccn) in the absence

or presence of 10% (volume/volume) Oxyrase™. Each point on the graph represents the mean

OD_{620} value from three independent cultures. Error bars, which in some cases are too small to

observe in the graph, represent the standard deviation (SD).

Figure 7: A *S. pneumoniae* ΔsodA mutant phenocopies the Zn hypersensitivity of a

ΔccnABCDE mutant strain. Growth characteristics at 37°C under an atmosphere of 5% CO_2 in

BHI broth alone (A) or with 0.2 mM $ZnSO_4$ (B) of IU781 (D39), NRD10176 (Δccn), NRD10533

(D39 ΔsodA), and NRD10534 (Δccn ΔsodA). Each point on the graph represents the mean

OD_{620} value from three independent cultures. Error bars, which in some cases are too small to

observe in the graph, represent the standard deviation (SD).

Figure 8: Model of the function of the Ccn sRNAs in regulating Mn homeostasis. The Ccn sRNAs down-regulate expression of Leloir pathway (*galK*, *galT2*, *galT1*, *galE2*) and galactose transport genes (*spd_0088*, *spd_0089*, *spd_0090*) resulting in reduced CPS synthesis and thus, a thinner capsule, which improves Mn uptake (pneumococcal cell on the left). In the absence of the Ccn sRNAs, increased expression of the Leloir pathway genes leads to increased production of capsule precursors and consequently, increased capsule thickness limiting Mn uptake (pneumococcal cell on the right). MgtA (Ca^{2+}/Mn^{2+} exporter); MntE (primary Mn^{2+} exporter); PsaBCA (only known Mn^{2+} importer); CzcD (Cd^{2+}/Zn^{2+} exporter); Pht (histidine triad proteins that bind Zn^{2+} and transfer to AdcAII); AdcBCAII (Zn^{2+} import system).