Genetic requirement of *dact1/2* to regulate noncanonical Wnt signaling and *calpain 8* during embryonic convergent extension and craniofacial morphogenesis

Shannon H. Carroll[1,2,3,], Sogand Schafer[1,2], Kenta Kawasaki[2,3], Casey Tsimbal[2,3],

Amélie M. Julé[4], Shawn A. Hallett[2,3], Edward Li[2] and Eric C. Liao[1,2,3]

[1]Center for Craniofacial Innovation, Children's Hospital of Philadelphia Research

Institute, Children's Hospital of Philadelphia, PA 19104, USA.

[2]Division of Plastic and Reconstructive Surgery, Department of Surgery, Children's

Hospital of Philadelphia, PA 19104, USA.

[3]Shriners Hospital for Children, Tampa, FL 33607, USA

[4] Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA

02115, USA

Abstract

Wnt signaling plays crucial roles in embryonic patterning including the regulation of convergent extension during gastrulation, the establishment of the dorsal axis, and later, craniofacial morphogenesis. Further, Wnt signaling is a crucial regulator of craniofacial morphogenesis. The adapter proteins Dact1 and Dact2 modulate the Wnt signaling pathway through binding to Disheveled. However, the distinct relative functions of Dact1 and Dact2 during embryogenesis remain unclear. We found that *dact1* and *dact2* genes have dynamic spatiotemporal expression domains that are reciprocal to one another suggesting distinct functions during zebrafish embryogenesis. Both *dact1* and *dact2* contribute to axis extension, with compound mutants exhibiting a similar convergent extension defect and craniofacial phenotype to the *wnt11f2* mutant. Utilizing single-cell RNAseq and an established noncanonical Wnt pathway mutant with a shortened axis (*gpc4*), we identified *dact1/2* specific roles during early development. Comparative whole transcriptome analysis between wildtype and *gpc4* and wildtype and *dact1/2* compound mutants revealed a novel role for *dact1/2* in regulating the mRNA expression of the classical calpain *capn8*. Over-expression of *capn8* phenocopies *dact1/2* craniofacial dysmorphology. These results identify a previously unappreciated role of *capn8* and calcium-dependent proteolysis during embryogenesis. Taken together, our findings highlight the distinct and overlapping roles of *dact1* and *dact2* in embryonic craniofacial development, providing new insights into the multifaceted regulation of Wnt signaling.

Introduction

Wnt signaling is a crucial regulator of embryogenesis through its regulation of body axis

patterning, cell fate determination, cell migration, and cell proliferation (Logan and

Nusse 2004, Steinhart and Angers 2018, Mehta, Hingole et al. 2021). Current

mechanistic understanding of Wnt signaling during embryogenesis includes an

extensive catalog of ligands, receptors, co-receptors, adaptors, and effector molecules

(Clevers and Nusse 2012, Niehrs 2012, Loh, van Amerongen et al. 2016, Mehta,

Hingole et al. 2021). The intricate spatiotemporal integration of Wnt signaling

combinations is an important focus of developmental biology and tissue morphogenesis

(Petersen and Reddien 2009, Clevers and Nusse 2012, Loh, van Amerongen et al.

2016, Wiese, Nusse et al. 2018). Disruptions of Wnt signaling-associated genes lead to

several congenital malformations which often affect multiple organ systems given their

pleotropic developmental functions (Hashimoto, Morita et al. 2014, Shi 2022).

Craniofacial anomalies are among the most common structural congenital

malformations and genes in the Wnt signaling pathway are frequently implicated (Ji,

Hao et al. 2019, Reynolds, Kumari et al. 2019, Huybrechts, Mortier et al. 2020).

Genetic approaches in zebrafish have identified a number of key Wnt regulators of early

development, with gastrulation and craniofacial phenotypes (Brand, Heisenberg et al.

1996, Hammerschmidt, Pelegri et al. 1996, Heisenberg, Brand et al. 1996, Piotrowski,

Schilling et al. 1996, Solnica-Krezel, Stemple et al. 1996, Schilling and Le Pabic 2009).

The *silberblick (slb)* mutant, later identified as a *wnt11f2* mutant allele, exhibits

gastrulation and midline craniofacial phenotypes that encompassed aspects of multiple

mutant classes. During early segmentation in the somite stage, the *wnt11f2* mutant

developed a shortened anterior-posterior axis and partially fused eyes (Heisenberg,

Brand et al. 1996). Subsequently, as the cranial prominences converge and the ethmoid

plate (EP) formed, instead of a fan-shaped structure observed in wildtype embryos, the

wnt11f2 mutant formed a rod-like EP with a significant deficiency of the medio-lateral

dimension (Heisenberg, Brand et al. 1996, Heisenberg and Nusslein-Volhard 1997).

Another mutant kny*pek (kny)*, identified as having a nonsense mutation in *gpc4*, an

extracellular Wnt co-receptor, was identified as a gastrulation mutant that also exhibited

a shortened body axis due to a defect in embryonic convergent extension (CE) (Solnica-

Krezel, Stemple et al. 1996). In contrast to the *slb/wnt11f2 mutant*, the *gpc4* mutant

formed an EP that is wider in the medio-lateral dimension than the wildtype, in the

opposite end of the EP phenotypic spectrum compared to *wnt11f2* (Topczewski, Sepich

et al. 2001, Rochard, Monica et al. 2016). These observations beg the question of how

defects in early patterning and CE of the embryo may be associated with later

craniofacial morphogenesis. The observation that *wnt11f2* and *gpc4* mutant share

similar CE dysfunction and axis extension phenotypes but contrasting craniofacial

morphologies (Heisenberg and Nusslein-Volhard 1997) supports a hypothesis that CE

mechanisms regulated by these Wnt pathway genes are specific to the temporal and

spatial contexts during embryogenesis.

Dact (aka Frodo, Dapper) are scaffolding proteins that regulate Dishevelled (Dvl)-

mediated Wnt signaling, both positively and negatively (Cheyette, Waxman et al. 2002,

Gloy, Hikasa et al. 2002, Waxman, Hocking et al. 2004, Gao, Wen et al. 2008, Wen,

Chiang et al. 2010, Ma, Liu et al. 2015, Lee, Cheong et al. 2018). Dact proteins bind

directly to Dvl (Gloy, Hikasa et al. 2002, Brott and Sokol 2005, Lee, Cheong et al. 2018)

and interact with and inhibit members of TGF-β and Nodal signaling pathways (Zhang,

Zhou et al. 2004, Su, Zhang et al. 2007, Meng, Cheng et al. 2008, Kivimae, Yang et al.

2011). In chick and Xenopus, *Dact2* and *Dact1* (respectively) are expressed in the

neural folds during neural crest delamination and are important in epithelial-

mesenchymal transition (EMT), Wnt signaling, and TGF-β signaling (Hikasa and Sokol

2004, Schubert, Sobreira et al. 2014, Rabadan, Herrera et al. 2016). In mouse embryos,

Dact1 is expressed predominantly in mesodermal tissues, as well as ectodermal-

derived tissues, (Hunter, Hikasa et al. 2006) and ablation of Dact1 results in defective

EMT and primitive streak morphogenesis, with subsequent posterior defects (Wen,

Chiang et al. 2010). Mouse embryonic Dact2 expression has been described in the oral

epithelium and ablation of *Dact2* causes increased cell proliferation (Li, Florez et al.

2013) and re-epithelialization in mice (Meng, Cheng et al. 2008) and zebrafish (Kim,

Kim et al. 2020).

Previous experiments using morpholinos to disrupt *dact1* and *dact2* in zebrafish found

dact1 morphants to be slightly smaller and to develop a normal body. In contrast, *dact2*

morphants were found to phenocopy described zebrafish gastrulation mutants, with

impaired CE, shortened body axis, and medially displaced eyes. Importantly, prior work

using morpholino-mediated gene disruption of *dact1* and *dact2* did not examine

craniofacial morphogenesis except to analyze *dact1* and *dact2* morphants head and eye

shapes under light microscopy (Waxman, Hocking et al. 2004). These experiments

were carried out at a time when morpholino was the accessible tool of gene disruption

(Nasevicius and Ekker 2000, Corey and Abrams 2001, Heasman 2002). Since

CRISPR/Cas9 targeted gene mutagenesis became popularized, many reports of

germline mutant phenotypes being discrepant from prior morpholino studies warranted

revisiting many of the prior work (Kok, Shin et al. 2015) and careful interpretation given

the caveats of each technology (Morcos, Vincent et al. 2015, Rossi, Kontarakis et al.

2015). More recently, a zebrafish CRISPR/Cas9 genetic *dact2* mutant has been

generated and studied, but unlike in the *dact2* morphant, no developmental phenotypes

were described (Kim, Kim et al. 2020).

Here we investigated the genetic requirement of *dact1* and *dact2* during embryogenesis

and craniofacial development using germline mutant alleles. We found an early

developmental role for *dact1* and *dact2* during gastrulation and body axis elongation.

We also characterized the abnormal craniofacial development of the *dact1/2* compound

mutants. We identified distinct transcriptomic profiles of wildtype, dact1/2, and gpc4

mutants during early development, including finding *calpain 8* (*capn8)* calcium-

dependent protease to be ectopically expressed in the *dact1/2* mutants. These results

elaborate on the cellular roles of dact1/2 and identify capn8 as a novel regulatory

candidate of embryogenesis.

Results

***dact1* and *dact2* have distinct expression patterns throughout embryogenesis**

To determine the spatiotemporal gene expression of *dact1* and *dact2* during

embryogenesis we performed wholemount RNA *in situ* hybridization (ISH) across key

time points (Fig. 1). Given that the described craniofacial phenotypes of the *dact2*

morphant and the *wnt11f2* mutant are similar (Heisenberg and Nusslein-Volhard 1997,

Waxman, Hocking et al. 2004), we also performed *wnt11f2* ISH to compare to *dact1* and

dact2 expression patterns.

During gastrulation at 8 hours post-fertilization (hpf; 75% epiboly), some regions

of *dact1* and *dact2* gene expression were shared and some areas are distinct to each

dact gene (Fig. 1A). Further, *dact* gene expression was distinct from *wnt11f2* in that

wnt11f2 expression was not detected in the presumptive dorsal mesoderm. Transcripts

of *dact1*, *dact2,* and *wnt11f2* were all detected in the blastoderm margin, as previously

described (Makita, Mizuno et al. 1998, Heisenberg, Tada et al. 2000, Gillhouse, Wagner

Nyholm et al. 2004). Transcripts of *dact2*, and to a lesser extent *dact1*, were also

detected in the prechordal plate and chordamesoderm (Fig. 1A). Additionally, *dact2*

gene expression was concentrated in the shield and presumptive organizer or

Nieuwkoop center along with *wnt11f2*. This finding is consistent with previously

described expression patterns in zebrafish and supports a role for *dact1 and dact2* in

mesoderm induction and *dact2* in embryo dorsalization (Thisse 2001, Gillhouse,

Wagner Nyholm et al. 2004, Muyskens and Kimmel 2007, Oteiza, Koppen et al. 2010).

At the end of gastrulation and during somitogenesis the differences in the domains of

dact1 and *dact2* gene expressions became more distinct (Fig. 1B,C). At tailbud stage,

dact1 transcripts were detected in the neuroectoderm and the posterior paraxial

mesoderm, whereas *dact2* transcripts were detected in the anterior neural plate,

notochord, and tailbud. Anterior notochord and tailbud expression overlapped with

wnt11f2 gene expression (Makita, Mizuno et al. 1998, Heisenberg, Tada et al. 2000).

The expression of *dact2* was unique in that its expression demarcated the anterior

border of the neural plate. As *dact2* morphants exhibited a craniofacial defect with

medially displaced eyes and midfacial hypoplasia (Waxman, Hocking et al. 2004), we

examined *dact1* and *dact2* expression in the orofacial tissues. At 24 hpf we found some

overlap but predominantly distinct expression patterns of *dact1* and *dact2* with *dact1*

being more highly expressed in the pharyngeal arches and *dact2* being expressed in

the midbrain/hindbrain boundary. Both *dact1* and *dact2* appeared to be expressed in the

developing oral cavity. At 48 hpf *dact1* expression is consistent with expression in the

developing craniofacial cartilage elements, while *dact2* expression appears within the

developing mouth. The distinct cellular expression profiles of *dact1* and *dact2* were

more clear in histological sections through the craniofacial region at 72 hpf. Utilizing

RNAscope *in situ* hybridization, we found that *dact2* and the epithelial gene *irf6* were co-

expressed in the surface and oral epithelium that surround the cartilaginous structures

(Fig. 1F). This is in contrast to *dact1* which was expressed in the developing cartilage of

the anterior neurocranium (ANC)/EP and palatoquadrate of the zebrafish larvae (Fig.

1F).

We examined the overall expression patterns of *dact1*, *dact2, gpc4,* and *wnt11f2*

using Daniocell single-cell sequencing data (Farrell, Wang et al. 2018). In general, we

found *dact1* spatiotemporal gene expression to be more similar to *gpc4* while *dact2*

gene expression was more similar to *wnt11f2* (Fig. S1). These results of shared but also

distinct domains of spatiotemporal gene expression of *dact1* and *dact2* suggest that the

dact paralogs may have some overlapping developmental functions while other roles

are paralog-specific.

dact1 and dact2 contribute to axis extension and dact1/2 compound mutants

exhibit a convergent extension defect

dact1 and *dact2* are known to interact with *disheveled* and regulate non-

canonical Wnt signaling (Gloy, Hikasa et al. 2002, Waxman, Hocking et al. 2004, Gao,

Wen et al. 2008, Wen, Chiang et al. 2010, Ma, Liu et al. 2015, Lee, Cheong et al. 2018)

and we have previously described the craniofacial anomalies of several zebrafish Wnt

mutants (Dougherty, Kamel et al. 2012, Kamel, Hoyos et al. 2013, Rochard, Monica et

al. 2016, Ling, Rochard et al. 2017, Alhazmi, Carroll et al. 2021). Previous work

investigated the effect of *dact1* and *dact2* disruption during zebrafish embryogenesis

using morpholinos and reported morphant phenotypes in embryonic axis extension and

eye fusion (Waxman, Hocking et al. 2004). However, the limitations associated with

morpholino-induced gene disruption (Kok, Shin et al. 2015, Morcos, Vincent et al. 2015,

Rossi, Kontarakis et al. 2015) and the fact that craniofacial morphogenesis was not

detailed for the *dact1* and *dact2* morphants, warranted the generation of mutant

germline alleles (Fig. S2). We created a *dact1* mutant allele (22 bp deletion, hereafter

dact1-/-) and a *dact2* mutant allele (7 bp deletion, hereafter *dact2-/-*), both resulting in a

premature stop codon and presumed protein truncation (Fig. S2A,B). Gene expression

of *dact1* and *dact2* was measured in pooled *dact1-/-*, *dact2-/-,* and *dact1/2-/-* embryos

(Fig. S2C). We found a decrease in *dact1* mRNA and an increase in *dact2* mRNA levels

in the respective CRISPR single mutants. We hypothesize that *dact2* mRNA levels are

maintained or elevated in the *dact2-/-* mutant due to the relative 3' position of the

deletion. In the *dact2-/-* embryos we found a slight increase in *dact1* mRNA levels,

suggesting a possible compensatory effect of dact2 disruption. The specificity of the

gene disruption was demonstrated by phenotypic rescue of the rod-like EP with the

injection of *dact1* or *dact2* mRNA. Injection of *dact1* mRNA or *dact2* mRNA or in

combination decreased the percentage of rod-like EP phenotype from near the

expected 25% (35% actual) to 2-7% (Fig. S2D,E).

Analysis of compound *dact1* and *dact2* heterozygote and homozygote alleles

during late gastrulation and early segmentation time points identified embryonic axis

extension anomalies (Fig. 2A,B). *dact1-/-* or *dact2-/-* homozygotes develop to be

phenotypically normal and viable. However, at 12 hpf, *dact2-/-* single mutants have a

significantly shorter body axis relative to wildtype. In contrast, body length shortening

phenotype was not observed in *dact1-/-* homozygotes. Compound heterozygotes of

dact1+/-; *dact2+/-* also developed normally but exhibited shorter body axis relative to

wildtype. The most significant axis shortening occurred in *dact1-/-; dact2-/-* double

homozygotes with a less severe truncation phenotype in the compound heterozygote

dact1+/-; dact2-/- (Fig. 2A,B). Interestingly, these changes in body axis extension do not

preclude the compound heterozygous larvae from reaching adulthood, except in the

dact1-/-; dact2-/- double homozygotes which did not survive from larval to juvenile

stages.

Body axis truncation has been attributed to impaired CE during gastrulation

(Tada and Heisenberg 2012). To delineate CE hallmarks in the *dact1-/-;dact2-/-*

mutants, we performed wholemount RNA ISH detecting genes that are expressed in

key domains along the body axis. At bud stage, *dact1-/-; dact2-/-* embryos demonstrate

bifurcated expression of *pax2a* and decreased anterior extension of *gsc* expression,

suggesting impaired midline convergence and anterior extension of the mesoderm (Fig.

2C). At the 1-2 somite stage, *zic1*, *pax2a* and *tbx6* are expressed in neural plate,

prospective midbrain and the tailbud, respectively, in both the wildtype and *dact1-/-;*

dact2-/- embryos. However, the spacing of these genes clearly revealed the shortening

of the anteroposterior body axis in the *dact1-/-; dact2-/-* embryos. Midline convergence

is decreased and the anterior border of the neural plate (marked by *zic1* expression)

was narrower in the *dact1-/-; dact2-/-* embryos (Fig. 2D). At the 10-somite stage (ss),

dact1-/-; dact2-/- embryos demonstrated decreased spacing between *ctslb1* and *pax2a*

gene expression, suggesting impaired lengthening of the anterior portion of the embryo.

Detection of muscle marker *myo1d* in the *dact1-/-; dact2-/-* embryos delineated impaired

posterior lengthening as well as reduced somitogenesis, evidenced by the decreased

number of somites (Fig. 2E). These data point to impaired CE of the mesoderm in

dact1/2-/- double mutants, which resulted in a shorter body axis. The aberrant CE and

axis extension in the *dact1/2-/-* phenotypes were similar to findings in other Wnt

mutants, such as *slb* and *kyn* (Heisenberg, Tada et al. 2000, Topczewski, Sepich et al.

2001) in that the body axis is truncated upon segmentation.

dact1/dact2 compound mutants exhibit axis shortening and craniofacial

dysmorphology

Given the defective converge phenotype and shortened axis in the *dact* mutants

during gastrulation, we examined the fish at 4 dpf for axis defects and for evidence of

defective morphogenesis in the craniofacial cartilages. Craniofacial morphology is an

excellent model for studying CE morphogenesis as many craniofacial cartilage elements

develop through this cellular mechanism (Kamel, Hoyos et al. 2013, Mork and Crump

2015, Sisson, Dale et al. 2015, Rochard, Monica et al. 2016). No craniofacial phenotype

was observed in *dact1* or *dact2* single mutants (data not shown). However, in-crossing

to generate *dact1/2-/-* compound homozygotes resulted in dramatic craniofacial

malformation (Fig. 3). Specificity of this phenotype to *dact1/2* was confirmed via rescue

with *dact1* or *dact2* mRNA injection (Fig. S2D,E). The *dact1/2-/-* mutant embryos

exhibited fully penetrant midfacial hypoplasia (Fig. 3A) however the degree of eye field

convergence in the midline varied between individuals. The forebrain protruded dorsally

and the mouth opening and ventral cartilage structures were displaced ventrally (Fig.

3A). Alcian blue staining of cartilage elements revealed severe narrowing of the EP into

a rod-like structure in 100 percent of double mutants, while the ventral cartilage

elements were largely unaffected (Fig. 3B). Notably the trabeculae extending posteriorly

from the EP and the rest of the posterior neurocranium exhibit wildtype morphology in

dact1/2. This *dact1/2-/-* double mutant phenotype is highly similar to that described for

wnt11f2 (*slb*) mutants, a key regulator of non-canonical Wnt signaling and CE (Kimmel,

Miller et al. 2001).

As axis lengthening was found to be affected by loss of *dact1* and *dact2* (Fig. 2A)

we measured body length in 5 dpf *dact1/2* compound mutants. Using the length of the

vertebral spine as a measure of body length we found a trend (p=0.06) towards an

effect of *dact1* and *dact2* on shortening of the body length. Similar to axis length during

gastrulation/segmentation, the shortening was most pronounced in *dact1/2-/-* double

homozygous mutants versus wildtype clutch-mates (Fig. 3C, Fig. S3A,B).

As *wnt11f2* signals via disheveled and since dact proteins are known to interact

with disheveled (Wong, Bourdelas et al. 2003, Zhang, Gao et al. 2006, Kivimae, Yang et

al. 2011), it is suspected that dact has a role in wnt11f2 signaling. Combinatorial gene

disruption with morpholinos showed that *dact2* morpholino exasperated the *wnt11*

morpholino midfacial/eye fusion defect (Waxman, Hocking et al. 2004). We

hypothesized that the shared phenotypes between *wnt11f2* and *dact1/2* mutants point

to these genes acting in the same signaling pathway. To test for genetic epistasis

between *wnt11f2*, *dact1,* and *dact2* genes we generated *wnt11f2/dact1/2-/-* triple

homozygous mutants. If *wnt11f2* and *dact1/2* had independent developmental

requirements, the *wnt11f2/dact1/2-/-* mutant may exhibit a phenotype distinct from

wnt11f2-/- or *dact1/2-/-* mutants. We found that the *wnt11f2/dact1/2-/-* triple

homozygous mutant phenotype of the linear rod-like EP was the same as the *wnt11f2-/-*

mutant or *dact1/2-/-* double mutant, without exhibiting additional or neo-phenotypes in

the craniofacial cartilages or body axis (Fig. 3D,E). This result supports *dact1 and dact2*

acting downstream of *wnt11f2* signaling during ANC morphogenesis, where loss of

dact1/2 function recapitulates a loss of *wnt11f2* signaling.

Lineage tracing of *dact1/2* mutant NCC movements reveals their ANC

composition

The EP forms from the convergence of a central frontal prominence-derived

structure with bilateral maxillary prominence-derived elements (Wada, Javidan et al.

2005, Swartz, Sheehan-Rooney et al. 2011, Dougherty, Kamel et al. 2012, Mork and

Crump 2015, Rochard, Monica et al. 2016).The stereotypic convergent migration of

cranial neural crest cells and their derivatives presents an excellent model to examine

CE movements and their effects on tissue morphogenesis. The zebrafish EP is formed

from the joining of a midline frontal prominence derived from the anteromost cranial

neural crest cell (NCC) population that migrate over the eyes and turn caudally, to join

paired lateral maxillary prominences derived from the second stream of cranial NCC

population that migrate rostrally (Kimmel, Miller et al. 2001, Wada, Javidan et al. 2005,

Schilling and Le Pabic 2009, Dougherty, Kamel et al. 2012, Mork and Crump 2015). The

EP that forms is a planar fan-shaped structure where we and others have shown that

the morphology is governed by Wnt signaling (Kimmel, Miller et al. 2001, Rochard,

Monica et al. 2016).

Given the rod-like EP we observed in the *dact1/2-/-* double mutants, we

hypothesized that the dysmorphology could be due to aberrant migration of the

anteromost midline stream of cranial NCCs resulting in fusion of the lateral maxillary

components. Conversely, an abrogated contribution from the second paired stream of

maxillary NCCs could lead to an EP composed entirely of the medial component. To

distinguish between these possibilities, we carried out lineage tracing

of the cranial NCC populations in wildtype and *dact1/2* mutants. The *dact1/2*

compound mutants were bred onto a *sox10:kaede* transgenic background, where we

and others have shown that the *sox10* reporter is a reliable driver of cranial neural crest

labeling (Wada, Javidan et al. 2005, Dutton, Antonellis et al. 2008, Swartz, Sheehan-

Rooney et al. 2011, Dougherty, Kamel et al. 2012, Kague, Gallagher et al. 2012, Mork

and Crump 2015). Cranial NCC populations in wildtype and *dact1/2* mutants were

targeted at 19 hpf to photoconvert Kaede reporter protein in either the anterior cranial

NCCs that contribute to the frontal prominence, or the second stream of NCCs that

contribute to the maxillary prominence, where the labeled cells were followed

longitudinally over 4.5 days of development (Fig. 4). We found that the anterior NCCs of

wildtype embryos migrated antero-dorsally to the eye and populated the medial EP. To

our surprise, the anterior cranial NCC also migrated to contribute to the median element

of the rod-like EP, suggesting the complex anterior then caudal migration of the anterior

NCC is not disrupted by *dact1/2* mutation (Fig. 4A, arrows). This finding is in contrast to

lineage tracing in another midline mutant with a similarly shaped rod-like EP, the *syu*

(*sonic hedgehog* null) mutant, where the anterior NCCs failed to populate the ANC

(Wada, Javidan et al. 2005).

Next, the second stream of NCC population that contribute to the maxillary

prominence was labeled, where they migrate and contribute to the lateral element of the

EP as expected in the wildtype (Fig. 4B). When the second stream of cranial NCCs

were labeled and followed in the *dact1/2* mutants, the cells were found to migrate

normally up to 36 hpf, but did not ultimately populate the EP in the mutant (arrows).

These results suggest that NCC migration itself is not regulated by *dact1/2* but that loss

of *dact1/2* hinders the second stream of NCCs' ability to populate the ANC by an

alternative means. Further, we have found that a rod-like EP can be formed from 2

different NCC origins, where in the *dact1/2* mutants the EP is contributed by the

anteromost frontonasal NCCs, in contrast to the similar rod-shaped EP of the *syu*

mutants that is formed from the more posterior stream of maxillary NCCs (Wada,

Javidan et al. 2005).

Genetic interaction of *dact1/2* with *gpc4* and *wls* to determine facial morphology

Given the role of Dact/dapper as modifiers of Wnt signaling, we hypothesized

that genetic interaction of *dact1/2* with *wls* and *gpc4* will modify facial morphology. Gpc4

is a glycoprotein that binds Wnt ligands and modulates Wnt signaling. *gpc4* zebrafish

mutants have impaired CE which leads to a shortened body axis (Topczewski, Sepich

et al. 2001). Wls is a posttranslational modifier of Wnt ligands which promotes their

secretion (Banziger, Soldini et al. 2006, Bartscherer, Pelte et al. 2006). We previously

described that these components of the Wnt/PCP pathway (*gpc4* receptor, *wls*

intracellular ligand chaperon, and Wnt ligands *wnt9a* and *wnt5b)* are required for

craniofacial morphogenesis, where each gene affects particular morphologic aspects of

chondrocytes arrangement in the cardinal axis of the ANC and Meckel's cartilage

(Rochard, Monica et al. 2016, Ling, Rochard et al. 2017). Using the EP as a

morphologic readout, we examined the genetic interaction of *dact1* and *dact2* with *wls*

and *gpc4*. Compound mutants of *dact1*, *dact2, gpc4* or *wls* were generated by breeding

the single alleles. Compared to wildtype ANC morphology, abrogation of *gpc4* led to

increased width in the transverse axis, but shorter in the antero-posterior axis (Rochard,

Monica et al. 2016). Disruption of *wls* leads to ANC morphology that is also wider in the

transverse dimension, but to a lesser degree than observed in *gpc4*. Additionally, in the

wls mutant, chondrocytes stack in greater layers in the sagittal axis (Rochard, Monica et

al. 2016).

Disruption of *gpc4* or *wls* in addition to *dact1/2* generated EP morphology that

contained phenotypic attributes from each single mutant, so that the resultant ANC

morphology represented a novel ANC form. The EP of a triple homozygous

gpc4/dact1/2-/- mutant was triangular, wider in the transverse axis and shorter in the

antero-posterior axis compared to the rod-like ANC observed in the *dact1/2-/-* double

mutant (Fig. 5A,B). Similarly, the ANC of a triple homozygous *wls/dact1/2-/-* mutant was

in the shape of a rod, shorter in the antero-posterior axis and thicker in the sagittal axis

compared to the *dact1/2-/-* double mutant, reflecting attributes of the *wls* mutant (Fig.

5C,D). In addition to the EP phenotypes, the triple homozygous *gpc4/dact1/2-/-* mutant

also had a short body axis and truncated tail similar but more severe than the *gpc4*

mutant (Fig. 5A). Since compound disruption of *dact1*, *dact2*, and *gpc4* or *wls* resulted

in a new phenotype we conclude that these genes function in different components of

Wnt signaling during craniofacial development.

As we analyzed the subsequent genotypes of our *dact1/dact2/gpc4* triple

heterozygote in-cross we gleaned more functional information about *dact1* and *dact2.*

We found that *dact1* heterozygosity in the context of *dact2-/-; gpc4-/-* was sufficient to

replicate the triple *dact1/dact2/gpc4* homozygous phenotype (Fig. 5E). In contrast,

dact2 heterozygosity in the context of *dact1-/-; gpc4-/-* double mutant produced ANC in

the opposite phenotypic spectrum of ANC morphology, appearing similar to the *gpc4-/-*

mutant phenotype (Fig. 5E). These results show that *dact1* and *dact2* do not have

redundant function during craniofacial morphogenesis, and that *dact2* function is more

indispensable than *dact1*. These results also suggest that *dact1* and *gpc4* may have

overlapping roles in craniofacial development.

dact1/2 and gpc4 regulate axis extension via overlapping and distinct cellular

pathways

Our analyses of axis extension and the hallmarks of a CE defect (namely

decreased length and increased width between early tissues) demonstrate that *dact1*

and/or *dact2* are required for CE and anterior-posterior axis lengthening during

gastrulation (Fig. 3). An axis lengthening and CE defect has also been described in

gpc4 (aka kny) mutants (Topczewski, Sepich et al. 2001). We also observe a defect in

axis lengthening in *gpc4-/-* in our hands (representative image Fig. 6A) that is grossly

similar to the *dact1/2-/-* mutants. Interestingly, the midfacial hypoplasia of the *wnt11f2*

(slb) mutant has been attributed to a defect in axis extension and anterior neural plate

patterning (Heisenberg and Nusslein-Volhard 1997), whereas defective axis extension

does not lead to midfacial hypoplasia in the *gpc4-/-* mutant. Therefore, we hypothesized

that by comparing and contrasting the gene expression changes in *dact1/2* versus *gpc4*

mutants during axis extension we could identify cell programs specifically responsible

for the anterior axis defect and subsequent midfacial hypoplasia. We performed single-

cell transcriptional analysis to compare *dact1/2* mutants, *gpc4* mutants, and wildtype

embryos during the segmentation stage. Single-cell encapsulation and barcoded cDNA

libraries were prepared from individual dissociated 4 ss wildtype, *dact1/2-/-* compound

mutant and *gpc4-/-* mutant embryos using the 10X Genomics Chromium platform and

Illumina next-generation sequencing. Genotyping of the embryos was not possible but

quality control analysis by considering the top 2000 most variable genes across the

dataset showed good clustering by genotype, indicating the reproducibility of individuals

in each group. Twenty clusters were identified using Louvain clustering and identity

was assigned by reviewing cluster-specific markers in light of published expression data

(Farrell, Wang et al. 2018, Farnsworth, Saunders et al. 2020, Bradford, Van Slyke et al.

2022) (Fig. 6B,C). Qualitatively, we did not observe any significant difference in cluster

abundance between genotype groups (Fig. S4). We found that *dact1*, *dact2*, and *gpc4*

were detected at various levels across clusters, though *dact1* expression was lower

than *dact2* (Fig. 6D), consistent with what we observed in RNA wholemount ISH

analysis (Fig. 1).

To assess the relative differences in gene expression between genotype groups,

we merged clusters into broader cell lineages: ectoderm, axial mesoderm, and paraxial

mesoderm (Fig. 7A). We focused on these cell types because they contribute

significantly to CE processes and axis establishment. For each of these cell lineages,

we performed independent pseudo-bulk differential expression analyses (DEA) of

wildtype vs. *dact1/2-/-* mutant and wildtype vs. *gpc4-/-* mutant (Fig. 7A). In all 3 cases,

we found differentially expressed genes (DEGs) that were commonly in *dact1-/-;dact2-/-*

and *gpc4-/-* mutant relative to wildtype (Fig. 7B). To address the hypothesis that *dact1*

and *dact2* regulate molecular pathways distinct from those regulated by *gpc4* we also

identified genes that were differentially expressed only in *dact1/2-/-* mutants or only in

gpc4-/- mutants (Fig. 7B). Functional analysis of these DEGs found unique enrichment

of intermediate filament genes in *gpc4-/-* whereas *dact1/2-/-* mutants had enrichment for

pathways associated with proteolysis (Fig. 7C, S5). Enrichment for pathways associated

with calcium-binding were found in both *gpc4-/-* and *dact1/2-/-*, although the specific

DEGs were distinct (Fig. S5). We performed functional analyses specifically for genes

that were differentially expressed in *dact1/2-/-* mutants, but not in *gpc4-/-* mutants, and

found enrichment in pathways associated with proteolysis (Fig. 7C) suggesting a novel

role for Dact in embryogenesis.

Interrogation of *dact1/2-/-* mutant-specific DEGs found that the calcium-

dependent cysteine protease *calpain 8* (*capn8*) was significantly overexpressed in

dact1/2-/- mutants in paraxial mesoderm (103 fold), axial mesoderm (33 fold), and in

ectoderm (3 fold; Fig. 7A). We also found that loss of *dact1/2* causes significant

changes to *capn8* expression pattern (Fig. 8A). Whereas *capn8* gene expression is

principally restricted to the epidermis of wildtype embryos, loss of *dact1/2* leads to

significant expansion of ectopic *capn8* gene expression in broader cell types such as in

mesodermal tissues (Fig. 8A). We corroborated this finding with wholemount RNA ISH

for *capn8* expression in wildtype versus *dact1/2-/-* 12 hpf embryos (Fig. 8B). The

expression of *smad1* was found to be decreased uniquely in the ectoderm of *dact1/2-/-*

embryos relative to wildtype (Fig. 7A), however this finding was not investigated further,

Capn8 is considered a "classical" calpain, with domain homology similar to

Capn1 and *Capn2* (Macqueen and Wilcox 2014). In adult human and mouse tissue,

Capn8 expression is largely restricted to the gastrointestinal tract (Sorimachi, Ishiura et

al. 1993, Macqueen and Wilcox 2014), however embryonic expression in mammals has

not been characterized. Proteolytic targets of Capn8 have not been identified, however,

other classical calpains have been implicated in Wnt and cell-cell/ECM signaling

(Konze, van Diepen et al. 2014), including in Wnt/Ca^{+2} regulation of CE in Xenopus

(Zanardelli, Christodoulou et al. 2013). To determine whether the *dact1/2-/-* mutant

craniofacial phenotype could be attributed to *capn8* overexpression, we performed

injection of *capn8* or *gfp* control mRNA into 1 cell-stage zebrafish embryos. In wildtype

zebrafish, exogenous *capn8* mRNA caused the distinct *dact1/2-/-* craniofacial

phenotype including a rod-like ANC at a very low frequency (1 in 142 injected embryos).

This craniofacial phenotype was not observed in wildtype larvae, or when wildtype

embryos were injected with an equal concentration of *gfp* mRNA (0 in 192 injected

embryos) (data not shown). When mRNA was injected into 1 cell-stage embryos

generated from *dact1/2+/-* interbreeding, *capn8* caused a significant increase in the

number of larvae with the mutant craniofacial phenotype when on a *dact1/2+/-* genetic

background (Fig. 8D, 0.0% vs. 7.5%). We did not find an effect of exogenous *capn8* on

any other genotype, including *dact1-/-,dact2+/-* which we suspect to be due to the

smaller number of those individuals in our experimental population. These findings

suggest a new contribution of capn8 to embryonic development as well as anterior

neural plate patterning and craniofacial development. Further, the regulation of *capn8*

by *dact* may be required for normal embryogenesis and craniofacial morphogenesis.

Discussion

In this study, we examined the genetic requirement of *dact1* and *dact2* during

early embryogenesis and craniofacial morphogenesis in zebrafish. Wnt signaling is

central to the orchestration of embryogenesis and numerous proteins have been

identified as modulators of Wnt signaling, including *Dact1* and *Dact2* (Cheyette,

Waxman et al. 2002). Several studies across *Xenopus*, zebrafish, and mouse have

ascribed roles to *dact1* and *dact2,* including both promoting and antagonizing Wnt

signaling, depending on the developmental context (Cheyette, Waxman et al. 2002,

Gloy, Hikasa et al. 2002, Waxman, Hocking et al. 2004, Gao, Wen et al. 2008, Wen,

Chiang et al. 2010, Ma, Liu et al. 2015, Lee, Cheong et al. 2018). Here, we show that

dact1 and *dact2* are required for axis extension during gastrulation and show an

example of CE defects during gastrulation associated with craniofacial defects. During

axis extension, we show that genetic disruption of *dact2,* but not *dact1,* resulted in a

significantly shortened axis relative to wildtype. This result is similar to what was

previously found using morpholinos to disrupt *dact1* and *dact2.* Interestingly, genetically

disrupted mutants of *dact1* or *dact2* developed to be phenotypically normal whereas

dact1/2 compound mutants displayed a severe dysmorphic craniofacial phenotype.

Again, this is largely similar to the previous morpholino study that found disruption of

each gene to cause only a slight and occasional dysmorphic cranial phenotype at 24 hpf

(Waxman, Hocking et al. 2004). Notably, embryos injected with a mixture of *dact1* and

dact2 morpholino were not characterized after 10 ss, and the singly injected embryos

were not characterized after 24 hpf (Waxman, Hocking et al. 2004). Therefore, by

analyzing genetic mutants of dact1 and dact2 our findings have largely validated the

previous morpholino literature as well as added new data on later developmental

outcomes.

The gene expression and genetic epistasis experiments carried out here support

that the *dact* paralogs are not redundant and have unique functions during different

stages of embryonic and larval development. We observed that *dact1* and *dact2* have

distinct spatiotemporal expression patterns throughout embryogenesis, suggesting

unique roles for each paralog in developmental processes. Differential expression of

Dact1 and *Dact2* was also described during odontogenesis in mice (Kettunen, Kivimae

et al. 2010). This aligns with previous findings of differential roles of *dact1* and *dact2* in

canonical vs. non-canonical Wnt signaling (Waxman, Hocking et al. 2004) and a specific

role for *dact2*, but not *dact1*, in TGF-β signaling (Su, Zhang et al. 2007, Schubert,

Sobreira et al. 2014). However, the lack of a resultant phenotype upon genetic ablation

of *dact1* or *dact2* individually suggests the capability of functional compensation. This is

puzzling given their distinct expression patterns and needs to be examined further.

We found that *dact1* and *dact2* contribute to axis extension, and their compound

mutants exhibit a shortened and widened body axis that is consistent with a CE defect

during gastrulation. This finding aligns with previous studies that have implicated *dact1*

and *dact2* in non-canonical Wnt signaling and regulation of embryonic axis extension

(Waxman, Hocking et al. 2004). Based on our gene expression and combinatorial

genetic analyses, we offer the hypothesis that *dact1* expression in the paraxial

mesoderm is required for dorsal CE during gastrulation through its role in noncanonical

Wnt/PCP signaling, similar to the defect observed upon *gpc4* disruption. Conversely, we

posit that *dact2* functions in the prechordal mesoderm to promote anterior migration

during gastrulation, a function which has also been ascribed to *wnt11f2* (Heisenberg

and Nusslein-Volhard 1997). It is only upon loss of both *dact1* and *dact2* functions that

the axis is significantly truncated and a craniofacial malformation results. Further

experiments with spatially restricted gene ablation or cell transplantation are required to

test this hypothesis.

Our results underscore the crucial roles of *dact1* and *dact2* in embryonic

development and suggest a connection between gastrulation movements and

subsequent craniofacial morphogenesis. Our finding that in *dact1-/-;dact2-/-* compound

mutants the first stream of cranial NCC migrate and contribute to the ANC, while the

second stream fails to contribute suggests the possibility of an anatomical barrier to

migration, rather than an autonomous defect of the cranial NCCs. Disruption of the

sonic hedgehog signaling pathway in zebrafish results in a similar phenotype to *dact1/2-*

/- and *wnt11f2-/-* mutants where the eyes converge medially and the EP narrows to a

rod shape. Interestingly, lineage tracing analysis in hedgehog-disrupted embryos found

the rod-like EP to consist solely of second stream-derived cranial NCCs (Wada, Javidan

et al. 2005). This is in contrast to the *dact1/2-/-* mutants, demonstrating two different

cellular mechanisms that result in a similar anatomical dysmorphology. It will be

important to test the generality of this phenomenon and determine if other mutants with

craniofacial abnormalities have early patterning differences. Further, a temporally

conditional genetic knockout is needed to definitively test the connection between early

and later development.

By comparing the transcriptome across different Wnt genetic contexts, i.e. *gpc4-*

/- with that of the *dact1/2-/-* compound mutant, we identified a novel role for *dact1/2* in

the regulation of proteolysis, with significant misexpression of *capn8* in the mesoderm of

dact1/2-/- mutants. Although at a very low frequency, ectopic expression of *capn8*

mRNA recapitulated the *dact1/2-/-* mutant craniofacial phenotype, suggesting that

inhibition of *capn8* expression in the mesoderm by dact is required for normal

morphogenesis. Genes involved in calcium ion binding were also differentially

expressed in the *dact1/2-/-* mutants and we predict that altering intracellular calcium

handling in conjunction with *capn8* overexpression would increase the frequency of the

recapitulated *dact1/2-/-* mutant phenotype.

Capn8 is described as a stomach-specific calpain and a role during

embryogenesis has not been previously described. Calpains are typically calcium-

activated proteases and it is feasible that Capn8 is active in response to Wnt/Ca^{2+}

signaling. A close family member, Capn2 has been found to modulate Wnt signaling by

degradation of beta-catenin (Zanardelli, Christodoulou et al. 2013, Konze, van Diepen et

al. 2014). Our findings suggest that dact-dependent suppression of *capn8* expression is

necessary for normal embryogenesis and craniofacial morphogenesis, further

expanding the functional repertoire of *dact1/2*. Continued research is required to test a

direct regulatory role of dacts on *capn8* expression. While our data suggests an

interaction between dact signaling and capn8 function, we did not find capn8

overexpression to be wholly sufficient to cause the rod-like EP phenotype, Further, we

did not test the necessity of *capn8* for craniofacial development in this study. This study

does however identify capn8 as a novel embryonic gene warranting further investigation

into its role during embryogenesis, with possible implications for known craniofacial or

other disorders. Recently, Capn8 has been implicated in EMT associated with cancer

metastasis (Zhong, Xu et al. 2022, Song, Cui et al. 2024) and Xenopus capn8 was

found to be required for cranial NCC migration (Cousin, Abbruzzese et al. 2011) which

further supports a role of *capn8* in cranial NCC migration and craniofacial

morphogenesis.

Another gene identified in our scRNA-seq data to be differentially expressed in

the *dact1/22-/-* but not the *gpc4-/-* embryos was *smad1*. Smad1 acts in the TGF-β

signaling pathway and dact2 has been described to inhibit TGF-β and Nodal signaling

by promoting the degradation of Nodal receptors (Zhang, Zhou et al. 2004, Su, Zhang et

al. 2007, Meng, Cheng et al. 2008, Kivimae, Yang et al. 2011). Zebrafish Nodal pathway

mutants (*cyc/ndr2*, *oep/tdgf1*, *sqt/ndr1*) exhibit medially displaced eyes (Hatta, Kimmel

et al. 1991, Brand, Heisenberg et al. 1996, Heisenberg and Nusslein-Volhard 1997,

Feldman, Gates et al. 1998, Zhang, Talbot et al. 1998) and it is robustly feasible that

dysregulation of TGF-β signaling in the *dact1/2-/-* mutant contributes to the craniofacial

phenotype. Future research will examine the role of *dact1* and *dact2* in the coordination

of Wnt and TGF-β signaling and the importance of this coordination in the context of

craniofacial development. Of note, Sonic Hedgehog (shh) signaling is a principal

regulator to the vertebrate midline (Chiang, Litingtung et al. 1996, Ribes, Balaskas et al.

2010), and important in the development of the zebrafish floorplate (Halpern, Hatta et al.

1997, Odenthal, van Eeden et al. 2000). Mutants with disrupted *shh* expression or

signaling (*cyc/ndr2*, *smo, oep/tdgf1*) exhibit medially displaced eyes similar to the

dact1/2 mutants (Brand, Heisenberg et al. 1996, Chen, Burgess et al. 2001). We did not

find any genes within the sonic hedgehog pathway to be differentially expressed in

dact1/2 mutants, though post-transcriptional interactions cannot be ruled out.

This study has uncovered the genetic requirement of *dact1* and *dact2* in

embryonic CE and craniofacial morphogenesis, delineated the genetic interaction with

Wnt genes and identified capn8 as a modifier of this process. Future work will delineate

the molecular differences across the different *dact1/2* and other Wnt mutants to further

identify determinants of craniofacial morphogenesis; and to connect these findings to

clinically important Wnt regulators of facial morphology and pathology.

Methods

Animals and CRISPR/Cas9 targeted mutagenesis

All animal husbandry and experiments were performed in accordance with approval

from Massachusetts General Hospital Institutional Animal Care and Usage Committee.

Zebrafish (*Danio rerio*) embryos and adults were maintained in accordance with

institutional protocols. Embryos were raised at 28.5 C in E3 medium (Carroll, Macias

Trevino et al. 2020) and staged visually and according to standardized developmental

timepoints (Westerfield 1993). All zebrafish lines used for experiments and gene editing

were generated from the Tubingen or AB strain. The *wnt11f2* mutant line and gpc4-/-

mutant line were obtained from Zebrafish International Resource Center (wnt11f2^tx226/+

and gpc4^hi1688Tg/+ respectively). The *wls*-/- mutant line was originally gifted to the lab and

independently generated, as previously described (Rochard, Monica et al. 2016). The

sox10:kaede transgenic line was previously generated and described by our lab

(Dougherty, Kamel et al. 2012).

CRISPR sgRNA guides were designed using computational programs ZiFiT Targeter

v4.2 (zifit.partners.org/ZiFit) (Sander, Zaback et al. 2007), crispr.mit.edu

(https://zlab.bio/guide-design-resouces) (Ran, Hsu et al. 2013), and ChopChop

(https://chopchop.cbu.uib.no) (Montague, Cruz et al. 2014) with traditional sequence

constraints. Guides were chosen that were predicted to give high efficiency and

specificity. Guides best meeting these parameters were selected in exon 2 of *dact1* and

exon 4 of *dact2*. No suitable gRNA with sufficient efficiency were identified for *dact2* 5'

of exon 4 and the resulting phenotype was reassuring compared to previous morpholino

published results. Guides for *dact1* and *dact2* and Cas9 protein were prepared and

microinjected into 1 cell-stage zebrafish embryos and founders were identified as

previously described (Carroll, Macias Trevino et al. 2020). Primers flanking the sgRNA

guide site were designed for genotyping and fragment analysis was performed on

genomic DNA to detect base pair insertion/deletion. Sanger sequencing was performed

to verify targeted gene mutation and confirm the inclusion of a premature stop codon.

dact1 forward primer: TACAGAAGCTGCTGAAGTACCG, *dact1* reverse primer:

CCCTCTCTCAAAGTGTTTTGGT, *dact2* forward primer:

TGAAGAGCTCCACTCCCCTGT, *dact2* reverse primer:

GCAGTTGAGGTCCATTCAGC.

RT qPCR analysis

Pooled wildtype, *dact1-/-*, *dact2-/-*, and *dact1/2-/-* fish were collected and RNA extractions

were performed using RNeasy Mini Kit (Qiagen). cDNA was generated using High

Capacity cDNA Reverse Transcription Kit (ThermoFisher). Quantitative PCR (qPCR) was

performed using *dact1* (Dr03152516_m1) and *dact2* (Dr03426298_s1) TaqMan Gene

Expression Assays. Expression was normalized to 18S rRNA expression (

Hs03003631_g1). TaQman Fast Advanced master mix (ThermoFisher) and a

StepOnePlus Real-Time PCR system (Applied Biosystems) were used to measure

relative mRNA levels, which were calculated using the ddCT method.

Microinjection of mRNA

Template DNA for *in vivo* mRNA transcription was generated by PCR amplification of

the gene of interest from a zebrafish embryo cDNA library and cloning into pCS2+8

destination plasmid. mRNA for injection was generated using an *in vitro* transcription kit

(Invitrogen mMessage mMachine). One-cell stage zebrafish embryos were injected with

2 nL of mRNA in solution. To test genetic knockout specificity, 150 or 300 pg or dact1,

dact2 or dact1 and dact2 mRNA was injected. For capn8 overexpression analysis 200

pg or GFP or capn8 mRNA was injected. Following phenotyping analysis, genotypes

were determined by fragment analysis of *dact1* and *dact2* genotyping PCR products.

Wholemount and RNAscope *in situ* hybridization

Wholemount *in situ* hybridization was performed as previously described (Carroll,

Macias Trevino et al. 2020). Zebrafish embryonic cDNA was used as a template to

generate riboprobes. Primers were designed to PCR amplify the specific riboprobe

sequence with a T7 promoter sequence linked to the reverse primer. *In vitro*

transcription was performed using a T7 polymerase and DIG-labeled nucleotides

(Roche). RNAscope was performed on sectioned zebrafish larvae as previously

described (Carroll, Macias Trevino et al. 2020). Probes were designed by and

purchased from ACD Bio. Hybridization and detection was performed according to the

manufacturer's protocol. Sections were imaged using a confocal microscope (Leica

SP8) and z-stack maximum projections were generated using Fiji software.

Alcian blue staining and imaging

Alcian blue staining and imaging were performed at 4 dpf as previously described

(Carroll, Macias Trevino et al. 2020). Briefly, larvae were fixed in 4% v/v formaldehyde

overnight at 4°C. Larvae were dehydrated in 50% v/v ethanol and stained with Alcian

blue as described (Walker and Kimmel 2007). Whole and dissected larvae were imaged

in 3% w/v methylcellulose using a Nikon Eclipse 80i compound microscope with a Nikon

DS Ri1 camera. Z-stacked images were taken and extended depth of field was

calculated using NIS Element BR 3.2 software. Images were processed using Fiji

software. After image capture embryos were genotyped by PCR and fragment analysis.

Axis measurements

Compound *dact1+/-*; *dact2+/-* zebrafish were in-crossed and progeny were collected

from 2 separate clutches and fixed in 4% formaldehyde at approximately 8ss. Embryos

were individually imaged using a Zeiss Axiozoom stereoscope and processed for DNA

extraction and genotyping (Westerfield 1993). Images were analyzed using Fiji. A circle

was drawn to overlay the yolk and the geometric center was determined using the

function on Fiji. Using the Fiji angle tool, lines were drawn from the center point to the

anterior-most point of the embryo and from the center to the posterior-most point of the

embryo. The resulting inner angle of these lines was determined. Each angle

measurement was then calculated as a ratio to the average angle of the wildtype

embryos. ANOVA was performed to determine statistical significance, $p < 0.05$.

Lineage analysis of cranial NCCs

Live embryos were mounted in 1% w/v low melt agarose and covered with E3 medium

containing 0.013% w/v tricaine. Wildtype control and dact1/2-/- compound mutants on a

Tg(*soxl0*:kaede) background were imaged on a Leica DMi8 confocal microscope and

photoconverted using the UV laser (404 nm) until the green kaede fluorescence

disappeared. For each embryo, one side was photoconverted and the contralateral side

served as an internal control. After photoconversion, embryos were removed from the

agarose and raised in E3 medium at 28.5 C until the required developmental timepoint,

at which time they were similarly re-mounted and re-imaged. Z-stacked images were

processed as maximum-intensity projections using Fiji software.

Single-cell RNA sequencing (scRNA-seq)

Single-cell transcriptomic analyses were performed on 10 zebrafish embryos, including

4 wildtype, 3 *dact1-/-;dact2-/-* compound mutant, and 3 *gpc4-/-* mutant embryos.

Embryos were collected at 4 ss with *dact1-/-,dact2-/-* and *gpc4-/-* mutants being

identified by their truncated body axis. Embryos were dechorionated with a short

(approximately 10 min) incubation in 1 mg/ml Pronase and then washed 3x in embryo

medium. Cell dissociation was performed with modifications as previously described

(Farrell, Wang et al. 2018). Each embryo was transferred to 50 μl DMEM/F12 media on

ice. To dissociate cells, media was replaced with 200 μl DPBS (without Ca^{2+} and Mg^{2+})

with 0.1% w/v BSA. Embryos were disrupted by pipetting 10x with a P200 pipette tip.

500 μl of DPBS + 0.1% BSA was added and cells were centrifuged at 300xg for 1 min.

Cell pellets were resuspended in 200 μl DPBS + 0.1% BSA and kept on ice. Just prior

to encapsulation, cells were passed through a 40 μm cell strainer, and cell counts and

viability were measured. After droplet encapsulation, barcoding, and library preparation

using the 10X Genomics Chromium Single Cell 3' kit (version 3), data were sequenced

on an Illumina NovaSeq 6000 sequencer.

FASTQ files were demultiplexed and aligned to the GRCz11 build of the

zebrafish genome using Cellranger (version 6.1.0) (Zheng, Terry et al. 2017). Raw

Cellranger count matrices were imported into R (version 4.1.2) using Seurat (version

4.1.0) (Hao, Hao et al. 2021). First, we reviewed data for quality and excluded any

droplet that did not meet all of the following criteria: (i) have at least 1,500 unique

molecular identifiers (UMIs), (ii) covering at least 750 distinct genes; (iii) have <5% of

genes mapping to the mitochondrial genome; and (iv) have a log10 of detected genes

per UMI >80%. After quality control, the dataset was also filtered to exclude genes with

a detection rate below 1 in 3,000 cells, leaving a total of 20,078 distinct genes

expressed across 19,457 cells for analysis.

The quality-controlled count data were normalized using Pearson's residuals

from the regularized negative binomial regression model, as implemented in

Seurat::SCTransfrom (Hafemeister and Satija 2019). When computing the SCT model,

the effect of the total number of UMIs and number of detected genes per cell were

regressed out. After normalization, the top 3,000 most variably expressed genes were

used to calculate principal components (PCs). Data were then integrated by source

sample using Harmony (version 0.1.0) (Korsunsky, Millard et al. 2019). A two-

dimensional uniform manifold approximation and projection (UMAP) (Becht, McInnes et

al. 2018) was then derived from the first 40 Harmony embeddings for visualization.

Using the integrated Harmony embeddings, clusters were defined with the Louvain

clustering method, as implemented within Seurat. A resolution of 0.3 was used for

cluster definition. Cluster identities were assigned by manually reviewing the results of

Seurat::FindAllMarkers, searching for genes associated to known developmental

lineages. Gene expression data for key markers that guided cluster identity assignment

were visualized using Seurat::DotPlot.

Following this detailed annotation, some clusters were grouped to focus

downstream analyses on 3 major lineages: ventral mesoderm (grouping cells from the

pronephros, vasculogenic/myeloid precursors, hematopoietic cells, heart primordium,

and cephalic mesoderm clusters), dorsal mesoderm (adaxial cells, segmental plate, and

paraxial mesoderm), ectoderm (CNS, mid/hindbrain boundary, spinal cord, and neural

crest). For those 3 lineages, single-cell level data were aggregated per sample and

cluster to perform pseudobulk differential expression analyses (DEA) contrasting

genotypes. Independent pairwise comparisons of *dact1-/-;dact2-/-* versus wildtype and

gpc4-/- vs wildtype were performed using DESeq2 (v1.34.0) (Love, Huber et al. 2014).

P-values were corrected for multiple testing using the default Benjamini-Hochberg

method; log2 fold change values were corrected using the apeglm shrinkage estimator

(Zhu, Ibrahim et al. 2019). Significance was defined as an adjusted p-value < 0.1 and

log2 fold change > 0.58 in absolute value. Heatmaps of the top most significant

differentially expressed genes were generated from the regularized log transformed

data using pheatmap (version 1.0.12). Overlap in significant genes across pairwise

comparisons were determined and visualized in Venn diagrams. Over-representation

analyses against the Gene Ontology (GO) database were ran using clusterProfiler

(version 4.2.2) (Wu, Hu et al. 2021), using as input the set of genes found to be

differentially expressed in the comparison of *dact1-/-;dact-/-* versus wildtype but not

gpc4-/- versus wildtype. Sequencing data have been deposited in GEO under accession

code GSE240264.

Statistical analysis

Analyses were performed using Prism Software (GraphPad) unless otherwise specified.

An unpaired Student's *t* test or one-way ANOVA with multiple comparisons was used as

indicated and a *P*-value <0.05 was considered significant. Graphs represent the mean

+/- the SEM and *n* represents biological replicates. For categorical data (normal vs.

mutant phenotype) a Fisher exact test was performed between *gfp* and *capn8* injected

embryos and the odds ratio was determined. The confidence interval was determined

by the Baptista-Pike method.

Acknowledgments

We thank Christoph Seiler, Adele Donohue and the Aqautics Facility team at Children's

Hospital of Philadelphia; and Jessica Bethoney at Massachusetts General Hospital

(MGH) for their excellent management of our zebrafish colonies and facilities. We thank

the MGH Next Generation Sequencing Core for cell encapsulation, cDNA library

preparation, and sequencing. Single-cell sequencing analysis was performed by the

Harvard Chan Bioinformatics Core. Work by A.J. was funded in part by the Harvard

Stem Cell Institute. We appreciate and acknowledge the generous funding support from

the Shriners Hospitals for Children. This work was supported by the National Institutes

of Health grant R01DE027983 to E.C.L.

References

Alhazmi, N., S. H. Carroll, K. Kawasaki, K. C. Woronowicz, S. A. Hallett, C. Macias Trevino, E. B. Li,
R. Baron, F. Gori, P. C. Yelick, M. P. Harris and E. C. Liao (2021). "Synergistic roles of Wnt
modulators R-spondin2 and R-spondin3 in craniofacial morphogenesis and dental
development." Sci Rep **11**(1): 5871.

Banziger, C., D. Soldini, C. Schutt, P. Zipperlen, G. Hausmann and K. Basler (2006). "Wntless, a
conserved membrane protein dedicated to the secretion of Wnt proteins from signaling cells."
Cell **125**(3): 509-522.

Bartscherer, K., N. Pelte, D. Ingelfinger and M. Boutros (2006). "Secretion of Wnt ligands
requires Evi, a conserved transmembrane protein." Cell **125**(3): 523-533.

Becht, E., L. McInnes, J. Healy, C. A. Dutertre, I. W. H. Kwok, L. G. Ng, F. Ginhoux and E. W.
Newell (2018). "Dimensionality reduction for visualizing single-cell data using UMAP." Nat
Biotechnol.

Bradford, Y. M., C. E. Van Slyke, L. Ruzicka, A. Singer, A. Eagle, D. Fashena, D. G. Howe, K. Frazer,
R. Martin, H. Paddock, C. Pich, S. Ramachandran and M. Westerfield (2022). "Zebrafish
information network, the knowledgebase for Danio rerio research." Genetics **220**(4).

Brand, M., C. P. Heisenberg, R. M. Warga, F. Pelegri, R. O. Karlstrom, D. Beuchle, A. Picker, Y. J.
Jiang, M. Furutani-Seiki, F. J. van Eeden, M. Granato, P. Haffter, M. Hammerschmidt, D. A. Kane,
R. N. Kelsh, M. C. Mullins, J. Odenthal and C. Nusslein-Volhard (1996). "Mutations affecting
development of the midline and general body shape during zebrafish embryogenesis."
Development **123**: 129-142.

Brott, B. K. and S. Y. Sokol (2005). "Frodo proteins: modulators of Wnt signaling in vertebrate
development." Differentiation **73**(7): 323-329.

Carroll, S. H., C. Macias Trevino, E. B. Li, K. Kawasaki, N. Myers, S. A. Hallett, N. Alhazmi, J.
Cotney, R. P. Carstens and E. C. Liao (2020). "An Irf6-Esrp1/2 regulatory axis controls midface
morphogenesis in vertebrates." Development **147**(24).

Chen, W., S. Burgess and N. Hopkins (2001). "Analysis of the zebrafish smoothened mutant
reveals conserved and divergent functions of hedgehog activity." Development **128**(12): 2385-
2396.

Cheyette, B. N., J. S. Waxman, J. R. Miller, K. Takemaru, L. C. Sheldahl, N. Khlebtsova, E. P. Fox,
T. Earnest and R. T. Moon (2002). "Dapper, a Dishevelled-associated antagonist of beta-catenin
and JNK signaling, is required for notochord formation." Dev Cell **2**(4): 449-461.

Chiang, C., Y. Litingtung, E. Lee, K. E. Young, J. L. Corden, H. Westphal and P. A. Beachy (1996).
"Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function." Nature
383(6599): 407-413.

Clevers, H. and R. Nusse (2012). "Wnt/beta-catenin signaling and disease." Cell **149**(6): 1192-
1205.

Corey, D. R. and J. M. Abrams (2001). "Morpholino antisense oligonucleotides: tools for
investigating vertebrate development." Genome Biol **2**(5): REVIEWS1015.

Cousin, H., G. Abbruzzese, E. Kerdavid, A. Gaultier and D. Alfandari (2011). "Translocation of the
cytoplasmic domain of ADAM13 to the nucleus is essential for Calpain8-a expression and cranial
neural crest cell migration." Dev Cell **20**(2): 256-263.

Dougherty, M., G. Kamel, V. Shubinets, G. Hickey, M. Grimaldi and E. C. Liao (2012). "Embryonic
fate map of first pharyngeal arch structures in the sox10: kaede zebrafish transgenic model." J
Craniofac Surg **23**(5): 1333-1337.

Dutton, J. R., A. Antonellis, T. J. Carney, F. S. Rodrigues, W. J. Pavan, A. Ward and R. N. Kelsh
(2008). "An evolutionarily conserved intronic region controls the spatiotemporal expression of
the transcription factor Sox10." BMC Dev Biol **8**: 105.

Farnsworth, D. R., L. M. Saunders and A. C. Miller (2020). "A single-cell transcriptome atlas for
zebrafish development." Dev Biol **459**(2): 100-108.

Farrell, J. A., Y. Wang, S. J. Riesenfeld, K. Shekhar, A. Regev and A. F. Schier (2018). "Single-cell
reconstruction of developmental trajectories during zebrafish embryogenesis." Science
360(6392).

Feldman, B., M. A. Gates, E. S. Egan, S. T. Dougan, G. Rennebeck, H. I. Sirotkin, A. F. Schier and
W. S. Talbot (1998). "Zebrafish organizer development and germ-layer formation require nodal-
related signals." Nature **395**(6698): 181-185.

Gao, X., J. Wen, L. Zhang, X. Li, Y. Ning, A. Meng and Y. G. Chen (2008). "Dapper1 is a
nucleocytoplasmic shuttling protein that negatively modulates Wnt signaling in the nucleus." J
Biol Chem **283**(51): 35679-35688.

Gillhouse, M., M. Wagner Nyholm, H. Hikasa, S. Y. Sokol and Y. Grinblat (2004). "Two
Frodo/Dapper homologs are expressed in the developing brain and mesoderm of zebrafish."
Dev Dyn **230**(3): 403-409.

Gloy, J., H. Hikasa and S. Y. Sokol (2002). "Frodo interacts with Dishevelled to transduce Wnt
signals." Nat Cell Biol **4**(5): 351-357.

Hafemeister, C. and R. Satija (2019). "Normalization and variance stabilization of single-cell
RNA-seq data using regularized negative binomial regression." Genome Biol **20**(1): 296.

Halpern, M. E., K. Hatta, S. L. Amacher, W. S. Talbot, Y. L. Yan, B. Thisse, C. Thisse, J. H.
Postlethwait and C. B. Kimmel (1997). "Genetic interactions in zebrafish midline development."
Dev Biol **187**(2): 154-170.

Hammerschmidt, M., F. Pelegri, M. C. Mullins, D. A. Kane, M. Brand, F. J. van Eeden, M.
Furutani-Seiki, M. Granato, P. Haffter, C. P. Heisenberg, Y. J. Jiang, R. N. Kelsh, J. Odenthal, R. M.
Warga and C. Nusslein-Volhard (1996). "Mutations affecting morphogenesis during gastrulation
and tail formation in the zebrafish, Danio rerio." Development **123**: 143-151.

Hao, Y., S. Hao, E. Andersen-Nissen, W. M. Mauck, 3rd, S. Zheng, A. Butler, M. J. Lee, A. J. Wilk,
C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J. Jain, A. Srivastava, T.
Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J. M. McElrath, C. A. Blish, R. Gottardo, P. Smibert
and R. Satija (2021). "Integrated analysis of multimodal single-cell data." Cell **184**(13): 3573-
3587 e3529.

Hashimoto, M., H. Morita and N. Ueno (2014). "Molecular and cellular mechanisms of
development underlying congenital diseases." Congenit Anom (Kyoto) **54**(1): 1-7.

Hatta, K., C. B. Kimmel, R. K. Ho and C. Walker (1991). "The cyclops mutation blocks
specification of the floor plate of the zebrafish central nervous system." Nature **350**(6316): 339-
341.

Heasman, J. (2002). "Morpholino oligos: making sense of antisense?" Dev Biol **243**(2): 209-214.

Heisenberg, C. P., M. Brand, Y. J. Jiang, R. M. Warga, D. Beuchle, F. J. van Eeden, M. Furutani-
Seiki, M. Granato, P. Haffter, M. Hammerschmidt, D. A. Kane, R. N. Kelsh, M. C. Mullins, J.
Odenthal and C. Nusslein-Volhard (1996). "Genes involved in forebrain development in the
zebrafish, Danio rerio." Development **123**: 191-203.

Heisenberg, C. P. and C. Nusslein-Volhard (1997). "The function of silberblick in the positioning
of the eye anlage in the zebrafish embryo." Dev Biol **184**(1): 85-94.

Heisenberg, C. P., M. Tada, G. J. Rauch, L. Saude, M. L. Concha, R. Geisler, D. L. Stemple, J. C.
Smith and S. W. Wilson (2000). "Silberblick/Wnt11 mediates convergent extension movements
during zebrafish gastrulation." Nature **405**(6782): 76-81.

Hikasa, H. and S. Y. Sokol (2004). "The involvement of Frodo in TCF-dependent signaling and
neural tissue development." Development **131**(19): 4725-4734.

Hunter, N. L., H. Hikasa, S. M. Dymecki and S. Y. Sokol (2006). "Vertebrate homologues of Frodo
are dynamically expressed during embryonic development in tissues undergoing extensive
morphogenetic movements." Dev Dyn **235**(1): 279-284.

Huybrechts, Y., G. Mortier, E. Boudin and W. Van Hul (2020). "WNT Signaling and Bone: Lessons
From Skeletal Dysplasias and Disorders." Front Endocrinol (Lausanne) **11**: 165.

Ji, Y., H. Hao, K. Reynolds, M. McMahon and C. J. Zhou (2019). "Wnt Signaling in Neural Crest
Ontogenesis and Oncogenesis." Cells **8**(10).

Kague, E., M. Gallagher, S. Burke, M. Parsons, T. Franz-Odendaal and S. Fisher (2012).
"Skeletogenic fate of zebrafish cranial and trunk neural crest." PLoS One **7**(11): e47394.

Kamel, G., T. Hoyos, L. Rochard, M. Dougherty, Y. Kong, W. Tse, V. Shubinets, M. Grimaldi and E.
C. Liao (2013). "Requirement for frzb and fzd7a in cranial neural crest convergence and
extension mechanisms during zebrafish palate and jaw morphogenesis." Dev Biol **381**(2): 423-
433.

Kettunen, P., S. Kivimae, P. Keshari, O. D. Klein, B. N. Cheyette and K. Luukko (2010). "Dact1-3
mRNAs exhibit distinct expression domains during tooth development." Gene Expr Patterns
10(2-3): 140-143.

Kim, D. H., E. J. Kim, D. H. Kim and S. W. Park (2020). "Dact2 is involved in the regulation of
epithelial-mesenchymal transition." Biochem Biophys Res Commun **524**(1): 190-197.

Kimmel, C. B., C. T. Miller and C. B. Moens (2001). "Specification and morphogenesis of the
zebrafish larval head skeleton." Dev Biol **233**(2): 239-257.

Kivimae, S., X. Y. Yang and B. N. Cheyette (2011). "All Dact (Dapper/Frodo) scaffold proteins
dimerize and exhibit conserved interactions with Vangl, Dvl, and serine/threonine kinases."
BMC Biochem **12**: 33.

Kok, F. O., M. Shin, C. W. Ni, A. Gupta, A. S. Grosse, A. van Impel, B. C. Kirchmaier, J. Peterson-
Maduro, G. Kourkoulis, I. Male, D. F. DeSantis, S. Sheppard-Tindell, L. Ebarasi, C. Betsholtz, S.
Schulte-Merker, S. A. Wolfe and N. D. Lawson (2015). "Reverse genetic screening reveals poor
correlation between morpholino-induced and mutant phenotypes in zebrafish." Dev Cell **32**(1):
97-108.

Konze, S. A., L. van Diepen, A. Schroder, R. Olmer, H. Moller, A. Pich, R. Weissmann, A. W. Kuss,
R. Zweigerdt and F. F. Buettner (2014). "Cleavage of E-cadherin and beta-catenin by calpain
affects Wnt signaling and spheroid formation in suspension cultures of human pluripotent stem
cells." Mol Cell Proteomics **13**(4): 990-1007.

Korsunsky, I., N. Millard, J. Fan, K. Slowikowski, F. Zhang, K. Wei, Y. Baglaenko, M. Brenner, P. R.
Loh and S. Raychaudhuri (2019). "Fast, sensitive and accurate integration of single-cell data
with Harmony." Nat Methods **16**(12): 1289-1296.

Lee, H., S. M. Cheong, W. Han, Y. Koo, S. B. Jo, G. S. Cho, J. S. Yang, S. Kim and J. K. Han (2018).
"Head formation requires Dishevelled degradation that is mediated by March2 in concert with
Dapper1." Development **145**(7).

Li, X., S. Florez, J. Wang, H. Cao and B. A. Amendt (2013). "Dact2 represses PITX2 transcriptional
activation and cell proliferation through Wnt/beta-catenin signaling during odontogenesis."
PLoS One **8**(1): e54868.

Ling, I. T., L. Rochard and E. C. Liao (2017). "Distinct requirements of wls, wnt9a, wnt5b and
gpc4 in regulating chondrocyte maturation and timing of endochondral ossification." <u>Dev Biol</u>
421(2): 219-232.

Logan, C. Y. and R. Nusse (2004). "The Wnt signaling pathway in development and disease."
<u>Annu Rev Cell Dev Biol</u> **20**: 781-810.

Loh, K. M., R. van Amerongen and R. Nusse (2016). "Generating Cellular Diversity and Spatial
Form: Wnt Signaling and the Evolution of Multicellular Animals." <u>Dev Cell</u> **38**(6): 643-655.

Love, M. I., W. Huber and S. Anders (2014). "Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2." <u>Genome Biol</u> **15**(12): 550.

Ma, B., B. Liu, W. Cao, C. Gao, Z. Qi, Y. Ning and Y. G. Chen (2015). "The Wnt Signaling
Antagonist Dapper1 Accelerates Dishevelled2 Degradation via Promoting Its Ubiquitination and
Aggregate-induced Autophagy." <u>J Biol Chem</u> **290**(19): 12346-12354.

Macqueen, D. J. and A. H. Wilcox (2014). "Characterization of the definitive classical calpain
family of vertebrates using phylogenetic, evolutionary and expression analyses." <u>Open Biol</u> **4**(4):
130219.

Makita, R., T. Mizuno, S. Koshida, A. Kuroiwa and H. Takeda (1998). "Zebrafish wnt11: pattern
and regulation of the expression by the yolk cell and No tail activity." <u>Mech Dev</u> **71**(1-2): 165-
176.

Mehta, S., S. Hingole and V. Chaudhary (2021). "The Emerging Mechanisms of Wnt Secretion
and Signaling in Development." <u>Front Cell Dev Biol</u> **9**: 714746.

Meng, F., X. Cheng, L. Yang, N. Hou, X. Yang and A. Meng (2008). "Accelerated re-
epithelialization in Dpr2-deficient mice is associated with enhanced response to TGFbeta
signaling." <u>J Cell Sci</u> **121**(Pt 17): 2904-2912.

Montague, T. G., J. M. Cruz, J. A. Gagnon, G. M. Church and E. Valen (2014). "CHOPCHOP: a
CRISPR/Cas9 and TALEN web tool for genome editing." <u>Nucleic Acids Res</u> **42**(Web Server issue):
W401-407.

Morcos, P. A., A. C. Vincent and J. D. Moulton (2015). "Gene Editing Versus Morphants."
<u>Zebrafish</u> **12**(5): 319.

Mork, L. and G. Crump (2015). "Zebrafish Craniofacial Development: A Window into Early
Patterning." <u>Curr Top Dev Biol</u> **115**: 235-269.

Muyskens, J. B. and C. B. Kimmel (2007). "Tbx16 cooperates with Wnt11 in assembling the
zebrafish organizer." <u>Mech Dev</u> **124**(1): 35-42.

Nasevicius, A. and S. C. Ekker (2000). "Effective targeted gene 'knockdown' in zebrafish." Nat Genet **26**(2): 216-220.

Niehrs, C. (2012). "The complex world of WNT receptor signalling." Nat Rev Mol Cell Biol **13**(12): 767-779.

Odenthal, J., F. J. van Eeden, P. Haffter, P. W. Ingham and C. Nusslein-Volhard (2000). "Two distinct cell populations in the floor plate of the zebrafish are induced by different pathways." Dev Biol **219**(2): 350-363.

Oteiza, P., M. Koppen, M. Krieg, E. Pulgar, C. Farias, C. Melo, S. Preibisch, D. Muller, M. Tada, S. Hartel, C. P. Heisenberg and M. L. Concha (2010). "Planar cell polarity signalling regulates cell adhesion properties in progenitors of the zebrafish laterality organ." Development **137**(20): 3459-3468.

Petersen, C. P. and P. W. Reddien (2009). "Wnt signaling and the polarity of the primary body axis." Cell **139**(6): 1056-1068.

Piotrowski, T., T. F. Schilling, M. Brand, Y. J. Jiang, C. P. Heisenberg, D. Beuchle, H. Grandel, F. J. van Eeden, M. Furutani-Seiki, M. Granato, P. Haffter, M. Hammerschmidt, D. A. Kane, R. N. Kelsh, M. C. Mullins, J. Odenthal, R. M. Warga and C. Nusslein-Volhard (1996). "Jaw and branchial arch mutants in zebrafish II: anterior arches and cartilage differentiation." Development **123**: 345-356.

Rabadan, M. A., A. Herrera, L. Fanlo, S. Usieto, C. Carmona-Fontaine, E. H. Barriga, R. Mayor, S. Pons and E. Marti (2016). "Delamination of neural crest cells requires transient and reversible Wnt inhibition mediated by Dact1/2." Development **143**(12): 2194-2205.

Ran, F. A., P. D. Hsu, J. Wright, V. Agarwala, D. A. Scott and F. Zhang (2013). "Genome engineering using the CRISPR-Cas9 system." Nat Protoc **8**(11): 2281-2308.

Reynolds, K., P. Kumari, L. Sepulveda Rincon, R. Gu, Y. Ji, S. Kumar and C. J. Zhou (2019). "Wnt signaling in orofacial clefts: crosstalk, pathogenesis and models." Dis Model Mech **12**(2).

Ribes, V., N. Balaskas, N. Sasai, C. Cruz, E. Dessaud, J. Cayuso, S. Tozer, L. L. Yang, B. Novitch, E. Marti and J. Briscoe (2010). "Distinct Sonic Hedgehog signaling dynamics specify floor plate and ventral neuronal progenitors in the vertebrate neural tube." Genes Dev **24**(11): 1186-1200.

Rochard, L., S. D. Monica, I. T. Ling, Y. Kong, S. Roberson, R. Harland, M. Halpern and E. C. Liao (2016). "Roles of Wnt pathway genes wls, wnt9a, wnt5b, frzb and gpc4 in regulating convergent-extension during zebrafish palate morphogenesis." Development **143**(14): 2541-2547.

Rossi, A., Z. Kontarakis, C. Gerri, H. Nolte, S. Holper, M. Kruger and D. Y. Stainier (2015). "Genetic compensation induced by deleterious mutations but not gene knockdowns." Nature **524**(7564): 230-233.

Sander, J. D., P. Zaback, J. K. Joung, D. F. Voytas and D. Dobbs (2007). "Zinc Finger Targeter
(ZiFiT): an engineered zinc finger/target site design tool." Nucleic Acids Res **35**(Web Server
issue): W599-605.

Schilling, T. F. and P. Le Pabic (2009). "Fishing for the signals that pattern the face." J Biol **8**(11):
101.

Schubert, F. R., D. R. Sobreira, R. G. Janousek, L. E. Alvares and S. Dietrich (2014). "Dact genes
are chordate specific regulators at the intersection of Wnt and Tgf-beta signaling pathways."
BMC Evol Biol **14**: 157.

Shi, D. L. (2022). "Wnt/planar cell polarity signaling controls morphogenetic movements of
gastrulation and neural tube closure." Cell Mol Life Sci **79**(12): 586.

Sisson, B. E., R. M. Dale, S. R. Mui, J. M. Topczewska and J. Topczewski (2015). "A role of
glypican4 and wnt5b in chondrocyte stacking underlying craniofacial cartilage morphogenesis."
Mech Dev **138 Pt 3**: 279-290.

Solnica-Krezel, L., D. L. Stemple, E. Mountcastle-Shah, Z. Rangini, S. C. Neuhauss, J. Malicki, A. F.
Schier, D. Y. Stainier, F. Zwartkruis, S. Abdelilah and W. Driever (1996). "Mutations affecting cell
fates and cellular rearrangements during gastrulation in zebrafish." Development **123**: 67-80.

Song, N., K. Cui, L. Zeng, Y. Fan, Z. Wang, P. Shi, W. Su and H. Wang (2024). "Calpain 8 as a
potential biomarker regulates the progression of pancreatic cancer via EMT and AKT/ERK
pathway." J Proteomics **301**: 105182.

Sorimachi, H., S. Ishiura and K. Suzuki (1993). "A novel tissue-specific calpain species expressed
predominantly in the stomach comprises two alternative splicing products with and without
Ca(2+)-binding domain." J Biol Chem **268**(26): 19476-19482.

Steinhart, Z. and S. Angers (2018). "Wnt signaling in development and tissue homeostasis."
Development **145**(11).

Su, Y., L. Zhang, X. Gao, F. Meng, J. Wen, H. Zhou, A. Meng and Y. G. Chen (2007). "The
evolutionarily conserved activity of Dapper2 in antagonizing TGF-beta signaling." FASEB J **21**(3):
682-690.

Swartz, M. E., K. Sheehan-Rooney, M. J. Dixon and J. K. Eberhart (2011). "Examination of a
palatogenic gene program in zebrafish." Dev Dyn **240**(9): 2204-2220.

Tada, M. and C. P. Heisenberg (2012). "Convergent extension: using collective cell migration
and cell intercalation to shape embryos." Development **139**(21): 3897-3904.

Thisse, B., Pflumio, S., Furthauer, M., Loppin, B., Heyer, V., Degrave, A., Woehl, R., Lux, A.,
Steffan, T., Charbonnier, X.Q., Thisse, C. (2001). "Expression of the zebrafish genome during
embryogenesis." from http://zfin.org.

Topczewski, J., D. S. Sepich, D. C. Myers, C. Walker, A. Amores, Z. Lele, M. Hammerschmidt, J.
Postlethwait and L. Solnica-Krezel (2001). "The zebrafish glypican knypek controls cell polarity
during gastrulation movements of convergent extension." Dev Cell **1**(2): 251-264.

Wada, N., Y. Javidan, S. Nelson, T. J. Carney, R. N. Kelsh and T. F. Schilling (2005). "Hedgehog
signaling is required for cranial neural crest morphogenesis and chondrogenesis at the midline
in the zebrafish skull." Development **132**(17): 3977-3988.

Walker, M. B. and C. B. Kimmel (2007). "A two-color acid-free cartilage and bone stain for
zebrafish larvae." Biotech Histochem **82**(1): 23-28.

Waxman, J. S., A. M. Hocking, C. L. Stoick and R. T. Moon (2004). "Zebrafish Dapper1 and
Dapper2 play distinct roles in Wnt-mediated developmental processes." Development **131**(23):
5909-5921.

Wen, J., Y. J. Chiang, C. Gao, H. Xue, J. Xu, Y. Ning, R. J. Hodes, X. Gao and Y. G. Chen (2010).
"Loss of Dact1 disrupts planar cell polarity signaling by altering dishevelled activity and leads to
posterior malformation in mice." J Biol Chem **285**(14): 11023-11030.

Westerfield, M. (1993). The zebrafish book : a guide for the laboratory use of zebrafish
(Brachydanio rerio). Eugene, OR, M. Westerfield.

Wiese, K. E., R. Nusse and R. van Amerongen (2018). "Wnt signalling: conquering complexity."
Development **145**(12).

Wong, H. C., A. Bourdelas, A. Krauss, H. J. Lee, Y. Shao, D. Wu, M. Mlodzik, D. L. Shi and J. Zheng
(2003). "Direct binding of the PDZ domain of Dishevelled to a conserved internal sequence in
the C-terminal region of Frizzled." Mol Cell **12**(5): 1251-1260.

Wu, T., E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan, X. Fu, S. Liu, X.
Bo and G. Yu (2021). "clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data." Innovation (Camb) **2**(3): 100141.

Zanardelli, S., N. Christodoulou and P. A. Skourides (2013). "Calpain2 protease: A new member
of the Wnt/Ca(2+) pathway modulating convergent extension movements in Xenopus." Dev Biol
384(1): 83-100.

Zhang, J., W. S. Talbot and A. F. Schier (1998). "Positional cloning identifies zebrafish one-eyed
pinhead as a permissive EGF-related ligand required during gastrulation." Cell **92**(2): 241-251.

Zhang, L., X. Gao, J. Wen, Y. Ning and Y. G. Chen (2006). "Dapper 1 antagonizes Wnt signaling by
promoting dishevelled degradation." J Biol Chem **281**(13): 8607-8612.

Zhang, L., H. Zhou, Y. Su, Z. Sun, H. Zhang, L. Zhang, Y. Zhang, Y. Ning, Y. G. Chen and A. Meng
(2004). "Zebrafish Dpr2 inhibits mesoderm induction by promoting degradation of nodal
receptors." Science **306**(5693): 114-117.

Zheng, G. X., J. M. Terry, P. Belgrader, P. Ryvkin, Z. W. Bent, R. Wilson, S. B. Ziraldo, T. D.
Wheeler, G. P. McDermott, J. Zhu, M. T. Gregory, J. Shuga, L. Montesclaros, J. G. Underwood, D.
A. Masquelier, S. Y. Nishimura, M. Schnall-Levin, P. W. Wyatt, C. M. Hindson, R. Bharadwaj, A.
Wong, K. D. Ness, L. W. Beppu, H. J. Deeg, C. McFarland, K. R. Loeb, W. J. Valente, N. G. Ericson,
E. A. Stevens, J. P. Radich, T. S. Mikkelsen, B. J. Hindson and J. H. Bielas (2017). "Massively
parallel digital transcriptional profiling of single cells." Nat Commun **8**: 14049.

Zhong, X., S. Xu, Q. Wang, L. Peng, F. Wang, T. He, C. Liu, S. Ni and Z. He (2022). "CAPN8 involves
with exhausted, inflamed, and desert immune microenvironment to influence the metastasis of
thyroid cancer." Front Immunol **13**: 1013049.

Zhu, A., J. G. Ibrahim and M. I. Love (2019). "Heavy-tailed prior distributions for sequence count
data: removing the noise and preserving large differences." Bioinformatics **35**(12): 2084-2092.

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

Figure Legends

Figure 1. Unique and shared *dact1* and *dact2* gene expression domains during zebrafish development. **(A-C)** Representative images of wholemount *in situ* hybridization showing *dact1*, *dact2,* and *wnt11f2* gene expression patterns. **A)** At 8 hpf, *dact2* and *wnt11f2* are highly expressed in the dorsal margin and presumptive Nieuwkoop center of the gastrulating embryo, with *dact1* being weakly detected (arrowhead). In contrast to *wnt11f2*, *dact1* and *dact2* are expressed in the presumptive dorsal mesoderm (asterisk). **B)** Lateral (anterior to the left of page) and anterior (dorsal side toward top of page) views of bud-stage embryos. *dact2* and *wnt11f2* transcripts are both detected in the tailbud (asterisk) while *dact2* is additionally expressed in the axial mesoderm (arrow). *dact1* gene expression is concentrated to the paraxial mesoderm and the neuroectoderm (open arrowheads). **C)** Lateral and flat-mount views of 4 ss embryos. *dact2* is expressed in the anterior neural plate and polster (P), notochord (N), paraxial and presomitic mesoderm (PM) and tailbud (TB). In contrast, *dact1* is expressed in the midbrain (MB) and the paraxial and presomitic mesoderm. **(D-E)** Representative lateral (anterior to left of page) images of wholemount *in situ* hybridization showing *dact1* and *dact2* expression patterns. **D)** At 24 hpf expression is detected in the developing head. **E)** At 48 hpf expression is detected in the developing craniofacial structures (arrow). **F)** Representative images of RNAscope *in situ* hybridization analysis of *dact1* (white) and *dact2* (yellow) and *irf6* (green) expression in transverse section of 72 hpf embryos. *dact1* is expressed in the ethmoid plate (ep) and palatoquadrate (pq) orofacial cartilage, while *dact2* is expressed in the oral epithelium

1096   (oe). The epithelial marker *irf6* is expressed in the oe and surface epithelium (se). Dapi

1097   (blue). Scale bar: 100 μm.

1098

Figure 2. Impaired convergent extension in *dact1/2* compound mutants. **A)** Inter-cross

of compound heterozygotes yield embryos with different degrees of axis extension that

correspond to the *dact1* and *dact2* genotypes. Representative lateral images of

embryos at 12 hpf. The yellow line indicates body axis angle measured from the anterior

point of the head, the center of the yolk, to the end of the tail. **B)** Quantification of body

axis angle. Numbers represent the difference in angle relative to the average wildtype

embryo. Asterisk indicates genotypes with angles significantly different from wildtype.

ANOVA p<0.5 n= 3-21 embryos. **C)** Representative bud stage wildtype and *dact1/2-/-*

mutant embryos stained for *gsc* (prechordal plate), *pax2a* (midbrain/hindbrain boundary)

and *krox20* (rhombomere 3). Asterisk indicates lack of *krox20* expression in *dact1/2-/-*

mutant. Scale bar = 200 μm **D)** Representative flat mounts of 1-2 ss wildtype and

dact1/2 mutant embryos stained for *zic1* (telencephalon), *pax2a* and *tbx6* (ventrolateral

mesoderm). **E)** Representative flat mounts of 10 ss wildtype and *dact1/2-/-* mutant

embryos stained for *ctsl1b* (hatching gland), *zic1*, *pax2a*, *krox20*, and *myo1d* (somites).

1113

Figure 3. Midface development requires *dact1* and *dact2*. **A)** Representative brightfield

images of wildtype and *dact1/2-/-* compound mutants at 4 dpf. 100 individuals were

analyzed from a *dact1/2+/-* double het cross. Lateral and ventral views show d*act1/2-/-*

compound mutants have a hypoplastic midface, medially displaced eyes, and a

displaced lower jaw. **B)** Representative flat-mount images of Alcian blue stained

ethmoid plate (EP) and visceral cartilage (VC) elements from 4 dpf wildtype and

d*act1/2-/-* compound mutants. d*act1/2-/-* mutants have a rod-shaped EP with no distinct

lateral and medial elements. No obvious differences were found in *dact1/2* mutant VC.

C) Representative brightfield image of 4 dpf wildtype and *dact1/2-/-* mutant. Bar

indicates vertebral spine length. Scale bar: 100 μm. **D)** Representative images of Alcian

blue stained *dact1/2-/-, wnt11f2-/-,* and *wnt11f2-/-,dact1/2-/-* compound mutants.

Embryos resulted from a *dact1+/-,dact2+/-,wnt11f2+/-* in-cross. Lateral and ventral

views show similar craniofacial phenotypes in each mutant. **E)** Representative flat-

mount images of Alcian blue stained EP show a similar phenotype between *dact1/2-/-,*

wnt11f2-/-, and *wnt11f2-/-,dact1/2-/-* compound mutants. Scale bar: 200 μm.

1129

Figure 4. Anterior neural crest cells of the *dact1/2-/-* mutant migrate to the midline and

populate the dysmorphic ethmoid plate. Lineage tracing of wildtype and *dact1/2-/-*

double mutant zebrafish embryos using Tg(*sox10*:kaede) line. *sox10*:kaede

fluorescence is shown in green and photo-converted kaede is shown in magenta and

highlighted with an arrow. Asterisks indicate that the cell population is absent. **A,B)**19

hpf embryo sagittal views showing photoconversion of anterior-most neural crest

population. At 36 hpf frontal images show the migration of photoconverted neural crest

cells to the frontal prominence in wildtype and *dact1/2-/-* double mutants. At 55 hpf,

frontal images show photoconverted neural crest cells populating the region of the

developing ANC in wildtype and *dact1/2-/-* mutants. At 4.5 dpf ventral images show

photoconverted neural crest cells populating the medial ethmoid plate in wildtype.

Similarly, neural crest cells in *dact1/2-/-* mutants populate the rod-shaped ethmoid plate.

Scale bar:100 μm. Representative images of 3 individual experiments.

1143

Figure 5. A nonoverlapping functional role for *dact1*, *dact2,* and *gpc4* and *wls*. **A)**

Representative Alcian blue stained wholemount images of wildtype, *dact1/2-/-* double

mutant, *gpc4-/-* mutant, and *gpc4/dact1/2-/-* triple mutants at 4 dpf. Low magnification

lateral images of embryos showing tail truncation in *dact1/2-/-* mutants, shortened and

kinked tail in *gpc4-/-* mutants, and a combinatorial effect in *gpc4/dact1/2-/-* triple

mutants. Higher magnification lateral images show a shortened midface and displaced

lower jaw in *dact1/2-/-* mutants, a shortened midface in *gpc4-/-* mutant, and a

combinatorial effect in *gpc4/dact1/2-/-* triple mutants. **B)** Representative flat-mount

images of dissected Alcian blue-stained cartilage elements. *dact1/2-/-* mutants have a

narrow rod-shaped ethmoid plate (EP) while *gpc4-/-* mutants have a broad and

shortened EP. *dact1/2/gpc4* triple mutants have a combinatorial effect with a short,

broad rod-shaped EP. In ventral cartilages (VC), *dact1/2-/-* mutants have a relatively

normal morphology while Meckel's cartilage in *gpc4-/-* mutants and *gpc4/dact1/2-/-* triple

mutants is truncated. **C,D)** Same as above except *wls-/-* mutant and *wls/dact1/2-/-* triple

mutant, with similar findings. **E)** Combinatorial genotypes of *dact1*, *dact2,* and *gpc4*.

dact2-/- contributed the *dact/gpc4* compound phenotype while *dact1-/-* did not. Scale

bar: 200 μm.

1161

Figure 6. Single-cell RNAseq of 4 ss wildtype, *dact1/2-/-* mutant and *gpc4-/-* mutants.

A) Summary schematic showing similar phenotypes in *dact1/2-/-* and *gpc4-/-* mutants at

12 hpf and divergent phenotypes at 4 dpf. Single-cell RNAseq was performed during

axis extension to compare and contrast *dact1/2-/-* and *gpc4-/-* transcriptional programs.

UMAP showing cluster identification. **B)** UMAP of cell clusters identified by Single-cell

RNAseq. **C)** Dot plot showing the most differentially expressed genes between clusters.

D) UMAP showing *dact1*, *dact2,* and *gpc4* expression in wildtype embryos.

1169

Figure 7. Pseudobulk differential expression analysis of single-cell RNAseq data. **A)**

Heatmaps showing the 50 most differentially expressed genes in 3 major cell types;

ectoderm (clusters 4,5,6,7), paraxial mesoderm (clusters 10,11,12), and lateral plate

mesoderm (clusters 15, 16, 17,18) between *dact1/2-/-* mutants and wildtype and *gpc4-/-*

mutants and wildtype. **B)** Venn diagrams showing unique and overlapping DEGs in

dact1/2-/- and *gpc4-/-* mutants. **C)** GO analysis of *dact1/2-/-* mutant-specific DEGs in

ectoderm showing enrichment for proteolytic processes.

1177

Figure 8. Expression of *capn8* is significantly dysregulated in *dact1/2-/-* mutants. **A)**

Single-cell RNAseq gene expression analysis of *capn8* in wildtype and *dact1/2-/-*

mutants. In wildtype embryos, *capn8* expression is restricted predominantly to the

epidermis whereas *capn8* is widely expressed throughout the embryo in *dact1/2-/-*

mutants, especially in the mesoderm. **B)** Wholemount *in situ* hybridization of *capn8*

expression in wildtype and *dact1/2-/-* mutant embryos at 2 ss. Staining corroborates the

single-cell RNAseq data, with expanded ectopic expression of *capn8* throughout the

embryo**.** Flat mounts are oriented anterior to the left. Scale bar: 100 μm. **C)** Brightfield

images and Alcian blue staining of the ethmoid plate show ectopic expression of *capn8*

mRNA (200 pg) at the 1-cell stage in *dact1+/-,dact2+/-* embryos recapitulates the

dact1/2-/- compound mutant craniofacial phenotype. The mutant craniofacial phenotype

did not manifest in gfp mRNA (200 pg) injected 1-cell stage *dact1+/-,dact2+/-* embryos.

Scale bar: 100 μm **D)** Quantification of mutant and normal craniofacial phenotype in 4

dpf larvae after mRNA injection at the 1-cell stage. Larvae were derived from *dact1/2+/-*

interbreeding. Larvae were uninjected or injected with 200 pg *gfp* control or *capn8*

mRNA. A Fisher exact test showed a significant effect of *capn8* mRNA injecting in the

dact1/2 double heterozygotes. p= 0.013.

1195

1196   **Supplemental Figures**

1197

1198   **Figure S1.** Daniocell single-cell RNAseq analysis with a display of *dact1*, *dact2*, *gpc4,*

1199   and *wnt11f2* in all cell clusters from 3-120 hpf of development (Farrell, Wang et al.

1200   2018). Cephalic mesoderm (ceph mes), mesenchyme, neural ectoderm, and muscle

1201   clusters are noted.

1202

Figure S2. Characterization of CRISPR/Cas9 generated *dact1-/-* and *dact2-/-* mutants.

A) Schematic representations of *dact1* and *dact2* exons, positions of guide RNA target

site, introduced premature stop codon (arrow), and sequences of mutations. **B)** DNA

fragment analysis of *dact1+/-* and *dact2+/-* animals showing wildtype (250 and 387 bp

respectively and mutant (228 and 380 bp respectively) **C)** Expression levels of *dact1*

and *dact2* mRNA by RT-qPCR in 12 hpf *dact1-/-* mutants, *dact2-/-* mutants, and

dact1/2-/- compound mutants. 8 embryos were pooled for mRNA isolation per sample.

D) Injection of *dact1* mRNA, *dact2* mRNA, or a combination of *dact1* and *dact2* mRNA

rescues the rod-shaped ethmoid plate phenotype in *dact1/2-/-* compound mutants.

Representative images of Alcian blue stained *dact1/2-/-* double mutant treated with 300

pg *dact1* mRNA and 300 pg *dact2* mRNA. Arrow highlights normal ethmoid plate (EP).

Visceral cartilage (VC) also appeared normal. **E)** Quantification of the mutant

craniofacial phenotype observed in a *dact1-/-,dact2+/-* breeding in-cross. Without mRNA

injection, the mutant phenotype was observed at approximately (35%) the expected

Mendelian ratio of 25%. Injection with *dact1* mRNA, *dact2* mRNA, or a combination of

dact1 and *dact2* mRNA decreased the frequency that the mutant craniofacial phenotype

was observed.

1220

1221   **Figure S3.** Loss of *dact1* and *dact2* tends to decrease total body length. **A)**

1222   Representative brightfield images of 4 dpf larvae of compound *dact1* and *dact2* mutant

1223   genotypes. Bar represents *dact1/2+/+* body length measurement. Scale bar: 100 μm. **B)**

1224   Scatter plot of body length measurement of compound *dact1* and *dact2* mutant

1225   genotypes at 4 dpf. ANOVA p=0.06. n=3-4.

1226

1227   **Figure S4.** Cluster abundance across genotype groups. Scatter box plots showing the

1228   frequency of each identified cell cluster between *dact1/2-/-*, *gpc4-/-*, and wildtype

1229   samples.

1230

1231   **Figure S5.** Loss of *gpc4* and loss of *dact1/2* leads to distinct changes in gene

1232   expression profiles but with some overlapping functions. GO analysis of DEGs identified

1233   between *gpc4-/-* and wildtype embryos and *dact1/2-/-* and wildtype embryos found

1234   changes in calcium ion binding and actin interaction in both mutants.

1235

Figure 1

Figure 2

Figure 3

A

	wildtype	*dact1/2 -/-*
lateral		
ventral		

B

	wildtype	*dact1/2 -/-*
EP		
VC		

C

wildtype *dact1/2 -/-*

D

dact1/2 -/- *wnt11f2 -/-* *wnt11f2-/-, dact1/2-/-*

lateral

ventral

E

EP n=6 n=3 n=3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

capn8 expression

wildtype   *dact1/2-/-*

1. floor plate
2. epidermis
3. periderm
4. central nervous system
5. mid/hindbrain boundary
6. spinal cord
7. neural crest
8. notochord
9. tail bud
10. adaxial cells
11. segmental plate
12. paraxial cells/somites
13. tail bud epidermis
14. pronephros
15. vasculogenic/myeloid cells
16. hematopoeitic cells
17. heart field
18. cephalic mesoderm
19. hatching gland/polster
20. otic placode
21. primordial germ cells

B

wildtype

dact1/2-/-

C

dact1+/-, dact2+/-   + *gfp*

dact1+/-, dact2+/-   + *capn8*

D

■ wildtype phenotype
□ *dact1/2-/-* phenotype

	uninj	gfp	capn8	uninj	gfp	capn8	uninj	gfp	capn8	uninj	gfp	capn8	uninj	gfp	capn8	uninj	gfp	capn8	uninj	gfp	capn8	uninj	gfp	capn8	uninj	gfp	capn8
dact1	+/+			+/+			+/+			+/-			+/-			+/-			-/-			-/-			-/-		
dact2	+/+			+/-			-/-			+/+			+/-			-/-			+/+			+/-			-/-		

Figure S1

Figure S2

A

dact1

Target site: TAACCAGCTCCAGGAATTAGACCGACAGA
Mutation: TAAC . . . . AGA
‑ 22bp

dact2

Target site: ACCGTTACCCCAGTCC
Mutation: ACCGTTA CC
‑ 7bp

B

228 250

380 387

C

relative expresssion

wt *dact1-/-*

wt *dact2-/-*

wt *dact1/2-/-*

D

Lateral

EP   VC

E

# of embryos

mutant
normal

uninjected 35%
dact1 300pg 2%
dact2 300pg 6%
dact1 + dact2 150pg 6%
dact1 + dact2 300pg 7%

Figure S3

Figure S4

Figure S5

gpc4-/- vs. wildtype

dact1/2-/- vs. wildtype