Distinct and interdependent functions of three RING proteins regulate recombination

during mammalian meiosis

Masaru Ito[1,2,5,*], Yan Yun[1,2], Dhananjaya S. Kulkarni[1,2], Sunkyung Lee[1,2], Sumit Sandhu[1,2],

Briana Nuñez[1,3], Linya Hu[2], Kevin Lee[2], Nelly Lim[2], Rachel M. Hirota[2], Rowan Prendergast[2],

Cynthia Huang[2], Ivy Huang[2] and Neil Hunter[1,2,4,6,*]

[1]Howard Hughes Medical Institute, University of California Davis, Davis, CA 95616, USA

[2]Department of Microbiology & Molecular Genetics, University of California Davis, Davis, CA

95616, USA

[3]Department of Biochemistry & Molecular Biology, Brown University, Providence, RI 02912, USA

[4]Department of Molecular & Cellular Biology, University of California Davis, Davis, CA 95616,

USA

[5]Current address: Institute for Protein Research, Osaka University, Osaka 565-0871, Japan

[6]Lead contact: nhunter@ucdavis.edu (N.H.)

[*]Correspondence: msrito2@protein.osaka-u.ac.jp (M.I.), nhunter@ucdavis.edu (N.H.)

Classification: CELL BIOLOGY

Keywords: fertility, meiosis, gamete, crossing over, recombination, aneuploidy, chromosome

segregation, DNA repair

SUMMARY

During meiosis, each pair of homologous chromosomes becomes connected by at least one

crossover, as required for accurate segregation at the first division, and adjacent crossovers are

widely separated thereby limiting total numbers. In coarsening models, crossover patterning

results from nascent recombination sites competing to accrue a limiting pro-crossover RING-

domain protein (COR) that diffuses between synapsed chromosomes. Here we delineate the

localization dynamics of three mammalian CORs in mouse and determine their

interdependencies. RNF212, HEI10 and a new member RNF212B show divergent

spatiotemporal dynamics along synapsed chromosomes, including profound differences in

spermatocytes and oocytes, that are not easily reconciled by elementary coarsening models.

Contrasting mutant phenotypes and genetic requirements indicate that RNF212B, RNF212, and

HEI10 play distinct but interdependent functions in regulating meiotic recombination and

coordinating the events of meiotic prophase-I by integrating signals from DNA breaks, homolog

synapsis, the cell-cycle, and incipient crossover sites.

SIGNIFICANCE

Meiosis produces haploid gametes by precisely halving the chromosome complement. Crossing

over between homologous chromosomes (homologs) is essential for their accurate segregation

and defects are associated with infertility, miscarriage, and congenital disease. Factors that

ensure crossing over between each pair of homologs include mammalian RING-domain

proteins RNF212, HEI10, and RNF212B, alleles of which are linked to infertility and heritable

variation in crossover rate. This study focuses on understanding the functions and relationships

between pro-crossover RING proteins (CORs) in mouse, providing important insights into their

roles in regulating recombination, the DNA repair process that gives rise to crossovers. Notably,

chromosomal localization dynamics of the three CORs are distinct and show striking sexual

dimorphism with important implications for models of crossover control.

INTRODUCTION

During sexual reproduction, parents contribute equally to their offspring by producing gametes

with precisely half the normal cellular ploidy. This is achieved via the two successive rounds of

chromosome segregation that occur during meiosis. Accurate segregation during the first

meiotic division (MI) requires crossing over between each pair of homologous chromosomes

(homologs). In combination with cohesion between sister-chromatids, crossovers create

connections called chiasmata that enable the stable biorientation of homolog pairs on the

meiosis-I spindle [1-3]. Defective crossing over can trigger cell death via the spindle checkpoint [4],

or cause aneuploidy due to chromosome missegregation [5,6], and is therefore associated with

infertility, miscarriage, and congenital syndromes such as Down, Turner and Klinefelter [7-10].

Crossing over is typically the minority outcome of meiotic recombination with most events being

repaired as noncrossovers without exchange of chromosome arms; for example, in mouse

spermatocytes only ~24 crossovers are selected from ~250 recombination events initiated by

programmed DNA double-strand breaks (DSBs) [11]. Crossover sites are selected in such a way

that each pair of chromosomes almost always obtains at least one crossover, termed crossover

assurance; adjacent events tend to be widely and evenly spaced via crossover interference

(which also limits total crossovers); and crossover numbers per nucleus show low variance

relative to DSBs, a buffering process known as crossover homeostasis [2, 3, 12]. The molecular

basis of crossover patterning remains unclear but manifests at the cytological level as the

selective retention/accumulation of certain pro-crossover factors at designated crossover sites

as prophase progresses [2, 3]. Among these are RING-domain proteins related to budding yeast

Zip3 [13], inferred to mediate protein modification by ubiquitin and the related small-protein

modifier SUMO [2, 14-21]. Species differ with respect to the number and subgroup (Zip3/RNF212

and HEI10) of these **C**ross**O**ver **R**ING proteins (CORs). For example, *Sordariales* and plant

genomes encode a single HEI10 homolog [13, 18, 22]; *Drosophila* encodes three Zip3/RNF212-like

proteins, two of which appear to be redundant [23] [21]; and *C. elegans* has four members with

homology to both subgroups [19, 20, 24].

Mammalian genomes encode both RNF212 and HEI10 (a.k.a. CCNBIP1) homologs with

interrelated functions that are essential for crossing over [15-17, 25]. RNF212 specifically localizes

between synapsed chromosomes, initially as numerous small foci all along the central region of

synaptonemal complexes (SCs), before undergoing HEI10-dependent repatterning to

accumulate at prospective crossover sites while being lost elsewhere [15-17]. RNF212 and other

CORs are inferred to stabilize pro-crossover factors that directly mediate the DNA events of

crossing over [15-19, 26, 27]. These include the MutSγ complex, a heterodimer of MSH4 and MSH5

that can bind D-loop and Holliday junction intermediates [28-30], and is inferred to stabilize and

protect them from dissociation by the conserved Bloom-helicase/decatenase complex, BLM-

TOPIIIα-RMI1-RMI2 (analogous to Sgs1-Top3-Rmi1 in budding yeast)[31-34]. In mouse

spermatocytes, MutSγ initially localizes to a majority of recombination sites during late zygotene

and early pachytene as homologs complete synapsis [11, 15]. Subsequently, MutSγ is lost from

most sites but retained at prospective crossover sites, which then accumulate crossover-specific

factors such as the MutLγ endonuclease [2, 29].

A key question for understanding meiotic crossover control is whether the dynamic focal

patterning of COR proteins along SCs is responsible for specifying crossover sites, as proposed

by recent models that invoke phase-transition/condensation, coarsening, and/or Ostwald

ripening processes [19, 35-37]. In these models, adjacent recombination sites compete to

accumulate a limiting amount of COR protein that diffuses along synaptonemal complexes, with

crossover designation ensuing at "winning" sites where COR has been accrued. Alternatively,

COR patterning may occur as a downstream consequence of crossover patterning by a distinct

process such as the mechanical stress envisioned in the beam-film model [38] [39]. The latter

possibility does not exclude COR coarsening, which could help reinforce initial crossover

designation or have other downstream functions.

Here we characterize mammalian CORs RNF212, HEI10, and a new member RNF212B [40-45].

Contrasting phenotypes, localization, and genetic dependencies imply that RNF212B, RNF212,

and HEI10 function as distinct modules that integrate signals from DSBs, synapsis, the cell-

cycle, and developing crossover complexes in order to effect the spatiotemporal coordination of

the major events of meiotic prophase-I. Moreover, the three CORs show diverse spatiotemporal

dynamics, including striking differences between males and females, features that are not

readily reconciled by coarsening models of crossover patterning.

RESULTS

***Rnf212b^{-/-}* mutant mice are sterile due to diminished crossing over**

Null mutations in mouse *Rnf212b* were created by pronuclear injection with CRISPR-Cas9,

targeting sequences immediately downstream of the first ATG (**Figures S1** and **S2**). Seven

mutated lines were obtained, the mutations and phenotypes of which are summarized in **Table**

S1. Two identical mutant lines, with a single nucleotide insertion that results in a frameshift after

the eighth codon and a premature stop codon after just 27 codons (**Figure S2B**), were chosen

for detailed phenotypic analysis. An independent *Rnf212b* mutant mouse line was recently

reported with similar phenotypes to those described below [45].

Heterozygous mutant animals were backcrossed four times and then bred to homozygosity.

Mature *Rnf212b*[-/-] mutant males had testes that were ~3-fold smaller than wild-type counterparts

(means ± SDs of 219.8 ± 16.8 mg per pair of testes from 8 wild-type mice versus 71.6 ± 7.8 mg

from 6 *Rnf212b*[-/-] mice; $p < 0.0001$, two-tailed *t* test; **Figure 1A**), the cauda epididymides were

devoid of sperm, and the animals were sterile. Testis sections from *Rnf212b*[-/-] mutant males

revealed that post-metaphase-I stages and spermatozoa were absent indicating loss of

spermatocytes at stage XII (**Figure 1B**). Sterility was also observed for female *Rnf212b*[-/-]

mutants, analysis of which is presented in **Figure 6**.

Rnf212[-/-] mutants are sterile because crossing over almost completely fails, resulting in

unconnected univalent chromosomes at metaphase I [15]. To determine whether this is also the

case for *Rnf212b*[-/-] mutants, chromosome spreads from diakinesis/metaphase-I spermatocytes

were analyzed (**Figure 1C**). In wild-type nuclei, all autosomes presented as bivalents with 24.1

± 2.3 chiasmata per nucleus (mean ± SD; n = 61 nuclei). Conversely, *Rnf212b*[-/-] nuclei

contained primarily univalents with a residual of 1-3 chiasmata (mean ± SD; 0.66 ± 0.75

chiasmata, n = 77 nuclei). Immunostaining of surface-spread pachytene-stage chromosomes

revealed that *Rnf212b*[-/-] mutant spermatocytes completely failed to form foci of the crossover-

specific marker MLH1 (**Figure 1D**), indicating defects in the designation and/or maturation of

crossover-specific recombination complexes. This inference was confirmed by analyzing two

other crossover markers, PRR19 and CDK2 (**Figure S3**) [16, 45, 46].

Irregular chromosome synapsis in *Rnf212b*[-/-] mutant spermatocytes

To begin to understand the defects leading to crossover failure in the *Rnf212b*[-/-] mutant,

progression through meiotic prophase-I was assessed by quantifying the fractions of

spermatocytes at each substage (leptonema, zygonema, pachynema and diplonema; **Figure**

1E). Stage distributions indicated a significant overrepresentation of zygotene-stage nuclei and

fewer pachytene nuclei in *Rnf212b*[-/-] mutant spermatocytes relative to wild type (p = 0.001, *G*

test) suggesting delayed or unstable synapsis (**Figure 1F**). Consistently, X-Y chromosomes

were frequently unsynapsed in mutant spermatocytes, although they were always in close

proximity (**Figure 1G** and **1H**). Unsynapsed X-Y chromosomes were over 5-fold more frequent

in *Rnf212b*[-/-] spermatocytes relative to wild type (28.6%, n = 57/199 cells from 3 mice and 5.7%,

n = 9/158 cells from 3 mice, respectively; p < 0.0001, Fisher's exact test) and were observed

specifically in mid/late pachytene cells (staining positive for the histone variant H1t) indicating

premature X-Y de-synapsis analogous to that seen in the *Rnf212*[-/-] mutants [15].

Intermediate steps of meiotic recombination are defective in *Rnf212b*[-/-] mutants

The modest synapsis defects of the *Rnf212b*[-/-] mutant suggest that early steps of recombination

may occur normally, with RNF212B promoting later steps of crossover designation and/or

maturation. To test these inferences, a variety of recombination markers were analyzed by

immunostaining spermatocyte chromosome spreads (**Figures 1I**, **S4** and **S5**). RAD51 and

DMC1 assemble onto resected DSB ends to form nucleoprotein filaments that catalyze

homologous pairing and DNA strand exchange [47]. The numbers of RAD51 and DMC1 foci

detected in zygotene spermatocytes were slightly elevated in the *Rnf212b*$^{-/-}$ mutant relative to

wild type (**Figure S4**), suggesting a mild perturbation of recombination as chromosomes

synapse that might reflect the delayed/unstable synapsis inferred above (**Figure 1G,H**).

Replication protein A (RPA; comprising RPA1, RPA2 and RPA3) binds single-stranded DNA

formed at DSB ends and at D-loops as strand exchange ensues [48]. Prominent immunostaining

foci of RPA2 emerge in zygonema, as DMC1 and RAD51 foci are diminishing, and then

disappear as DSB repair ensues during pachynema. In mid-zygonema, *Rnf212b*$^{-/-}$ and wild-type

spermatocytes formed similar numbers of RPA2 foci, but in subsequent stages levels were

significantly lower in *Rnf212b*$^{-/-}$ cells suggesting that DSBs are repaired faster in the mutant

(**Figure S5A,B**). More dramatic reductions were observed for foci of the ZMM proteins MSH4,

TEX11^{Zip4} and HFM1^{Mer3}, with ~3-fold fewer foci in the *Rnf212b*$^{-/-}$ mutant (**Figures 1I,J** and **S5C-**

F). The altered dynamics of RPA2 and ZMM factors in *Rnf212b*$^{-/-}$ mutant spermatocytes were

similar to those seen for *Rnf212*[-/-], although reduction of MSH4 foci appears to be more severe

in *Rnf212b*[-/-] cells (**Figures 1I,J** and **S5**) [15].

RNF212B dynamically localizes to synaptonemal complexes and recombination sites

To address the possibility that RNF212B functions locally to stabilize ZMMs at nascent

crossover sites, antibodies were raised against full-length RNF212B (isoform a; **Figure S1**) and

used to immunostain surface-spread spermatocyte chromosomes (**Figures 2A**,**B** and **S6**).

RNF212B was first detected in very early zygonema, specifically localizing to sites of synapsis

initiation marked by SYCP1 (73.1% of short SYCP1 stretches <2 μm colocalized with RNF212B

foci; n = 38/52 from 11 nuclei). As synapsis ensued, a punctate staining pattern emerged along

extending SCs, with focus numbers peaking in early pachynema, just after synapsis was

completed, at 175.9 ± 15.2 foci per nucleus (mean ± SD; n = 27 nuclei; **Figure 2B**). Focus

numbers then diminished throughout pachynema, but at the same time a small number of large,

bright "amplified" RNF212B foci emerged. By late pachynema, each SC had only one or two

large RNF212B foci with 25.0 ± 2.3 foci per nucleus (mean ± SD; n = 29 nuclei; **Figure 2B**). By

mid-diplonema, as homologs desynapsed, all RNF212B foci had disappeared.

Super-resolution structured illumination microscopy (SIM) showed that RNF212B, like RNF212,

localizes specifically to the central region of the SC (**Figure 2C**). The phenotypes of *Rnf212b$^{-/-}$*

mutants and the localization dynamics of the RNF212B protein indicate strong similarities with

RNF212. To begin to explore the relationship between RNF212B and RNF212, colocalization

was analyzed using SIM (**Figure 2D,E**). In late zygotene/early pachytene stage spermatocytes,

when focus numbers peaked, an average of 307.5 ± 50.0 RNF212B foci were detected per

nucleus compared to 374.6 ± 43.2 RNF212 foci, and 233.4 ± 38.7 RNF212B-RNF212 co-foci.

Thus, RNF212-RNF212B colocalization is high but incomplete with 75.9 ± 3.4% of RNF212B

foci colocalized with RNF212, whereas 62.1 ± 5.1% of RNF212 foci colocalized with RNF212B

(means ± SDs; *n* = 11 nuclei; **Figure 2D,E**; note that many more foci are resolved by SIM than

by the conventional fluorescence microscopy used in **Figure 2A,B**).

SIM was also used to analyze localization of RNF212B to ongoing recombination sites marked

by ZMM factor MSH4 (**Figure 2F,G**). In early pachynema, RNF212B were in 2-fold excess

relative to MSH4 (290.0 ± 38.3 RNF212B versus 149.4 ± 17.8 MSH4 foci per nucleus; means ±

SDs; n = 7 nuclei). Reflecting this excess, only 40.2 ± 6.2 % of RNF212B foci colocalized with

recombination sites marked by MSH4 foci; however, 77.2 ± 4.7% MSH4 foci colocalized with

RNF212B (means ± SDs; n = 7 nuclei; **Figure 2F,G**).

In late pachynema, numbers of amplified RNF212B foci per nucleus were indistinguishable from

those of the crossover marker MLH1 and colocalization was essentially absolute (**Figure 2H**;

99.7 ± 1.0 % of RNF212B foci colocalized with MLH1 foci and 99.0 ± 1.7 % of MLH1 foci

colocalized with RNF212B foci; means ± SDs; n = 15 nuclei). Thus, RNF212B foci in late-

pachytene spermatocytes exclusively localize to crossover sites.

Amplified RNF212B foci and HEI10 designate crossover sites by mid pachynema

We previously showed that immunostaining foci of the ubiquitin ligase HEI10 localize to

designated crossover sites marked by MLH1 foci in late pachynema [16, 17]. Unlike RNF212B and

RNF212, we did not detect numerous HEI10 foci in zygotene and early pachytene-stage

spermatocytes. Instead, crossover specific HEI10 foci materialized in pachytene, similar to other

crossover markers such as MLH1 (**Figure 3A,B**). By sub-staging pachytene spermatocytes, we

showed that initial emergence of MLH1 foci occurred only in nuclei with a full complement of

HEI10 foci, which was already present by mid pachynema (**Figure 3A,B**), i.e. crossover

designation must have occurred by this stage.

Having defined HEI10 foci as the earliest detectable marker of crossover designation, we set

out to determine the spatial-temporal relationship with amplification of RNF212B and RNF212

foci. The intensity of each RNF212B or RNF212 focus was measured from images with

matched exposures and normalized to the mean intensities of foci in early pachytene nuclei in

which HEI10 foci were not yet detected (n = 2,449 RNF212B foci and 2,513 RNF212 foci; 8

nuclei; **Figure 3C**,**D** and **S7A**). In mid-pachytene nuclei with at least one clear HEI10 crossover

focus on each autosomal SC, RNF212B foci at crossover sites (i.e. colocalizing with HEI10 foci)

were 11-fold brighter than other RNF212B foci in the same nuclei (i.e. the average of all other

foci from the same 8 nuclei; 6.29 ± 2.66 for 194 crossover foci versus 0.57 ± 0.59 for 2,004

other foci from 8 nuclei; means ± SDs; $p < 0.0001$, Mann-Whitney test) and 6.3-fold brighter

than the foci detected in early pachytene nuclei, consistent with the interpretation that crossover

foci grow at the expense other foci. The same analysis showed that RNF212 foci located at

crossover sites were only 3-fold brighter than other foci in the same nuclei (1.83 ± 1.05 for 194

crossover foci versus 0.60 ± 0.58 for 2,291 other foci form 8 nuclei; means ± SDs; $p < 0.0001$,

Mann-Whitney test) and 1.8-fold brighter than those in early pachytene. Thus, at designated

crossover sites in mid pachytene, RNF212B foci show stronger amplification than RNF212 foci.

In late pachytene spermatocytes, the only remaining RNF212B foci were crossover specific and

had intensities comparable to those of the crossover foci analyzed in mid-pachytene nuclei

(6.42 ± 4.84-fold brighter than early pachytene foci; 206 crossover foci from 8 nuclei; mean ±

SD) implying that RNF212B foci do not continue to enlarge between mid and late pachytene;

and that RNF212B is lost from other sites as opposed to being absorbed into crossover foci. By

contrast, RNF212 staining in late pachytene nuclei comprised a mixture of crossover foci and

other small residual foci. The crossover-associated RNF212 foci in these nuclei were 6-fold

brighter than other foci in the same nuclei (1.53 ± 1.13 for 206 crossover foci versus 0.24 ± 0.23

for 988 other foci from 8 nuclei; means ± SDs; $p < 0.0001$, Mann-Whitney test) but only 1.5-fold

brighter than those in early pachytene. Thus, crossover-associated RNF212 foci continue to

differentiate between mid and late pachynema but this may occur primarily via loss of RNF212

from noncrossover sites as opposed to continued growth of crossover-specific foci. Consistent

with these inferences, the total signal intensity of RNF212B per nucleus was unchanged from

early to mid pachynema, but then decreased from mid to late pachynema as RNF212B was lost

from noncrossover sites; by comparison, total signal intensity of RNF212 reduced during both

early to mid, and mid to late pachynema (1.00 ± 0.21, 0.97 ± 0.24, and 0.54 ± 0.33 for

RNF212B; and 1.00 ± 0.32, 0.69 ± 0.31, and 0.22 ± 0.10 for RNF212 in early, mid, and late

pachynema, respectively, n = 8 nuclei each; **Figure S7C**).

Per chromosome analysis for mid-pachytene autosomes highlighted additional distinctions

between RNF212B and RNF212 with respect to the amplification of brighter crossover-

correlated foci (**Figures 3E** and **S7B**). RNF212B foci at crossover sites were on average 5.1-

fold brighter relative to the brightest other focus along the same chromosome, whereas RNF212

foci were only 1.4-fold brighter (**Figure 3E**, left). When these relative intensities were plotted for

individual pairs of colocalizing RNF212-RNF212B crossover foci, almost no correlation was

detected (r^2 = 0.106), i.e. at mid pachynema, bright RNF212B foci are not coincident with bright

RNF212 foci, pointing to distinct amplification behaviors of the two proteins at designated

crossover sites (**Figure 3E**, right).

In mid-pachynema, 76.1% of designated crossover sites (n = 140/184; 8 nuclei) showed

amplification for RNF212B foci (≥2-fold brighter relative to the brightest other focus along the

same chromosome) but not for RNF212 foci. Oppositely only 0.5% of crossover sites (n =

1/184; 8 nuclei) showed amplification of only RNF212 but not RNF212B (**Figure 3E**, right).

Together, these data imply that crossover-specific amplification of RNF212B occurs earlier than

that of RNF212. The distinct behaviors of RNF212B and RNF212 were most pronounced in

nuclei transitioning between early and mid-pachynema, as HEI10 foci were emerging (i.e. nuclei

with <19 HEI10 foci per nucleus; **Figure 3F**). In these nuclei, RNF212B showed clear

amplification at designated crossover sites but RNF212 did not.

Together, our analysis of focus dynamics in spermatocytes shows that RNF212B foci amplify at

designated crossover sites during early-to-mid pachytene, coincident with the appearance of

HEI10 foci (**Figure S8**); and then RNF212B is lost from other sites without additional

amplification of crossover-associated foci. By contrast, RNF212 crossover foci differentiate later,

during mid-to-late pachytene; and loss of RNF212 from other sites, rather than growth, may be

the major cause of changes in relative intensity.

Interdependence between HEI10, RNF212B, and RNF212

Our previous studies evoked a model in which RNF212 establishes a precondition for

crossover/noncrossover differentiation by stabilizing nascent recombination intermediates and

rendering further progression dependent on HEI10 [16, 17]. Notably, in *Hei10*^{mei4/mei4} mutant

spermatocytes, abundant RNF212 foci persist along synapsed chromosomes and

recombination stalls at an intermediate step marked by the persistence of ZMM foci throughout

pachytene [16, 17]. Similarly, in early pachynema *Hei10*^{mei4/mei4} mutant spermatocytes, initial

RNF212B staining was indistinguishable from wild type, but this pattern of abundant foci along

SCs persisted until early diplonema (**Figure 4A,B**). Moreover, in sharp contrast to the

crossover-specific pattern of amplified RNF212B foci that developed in wild-type nuclei (24.8 ±

2.7 foci, mean ± SD from 16 nuclei), significantly amplified RNF212B foci were not detected

along the majority of chromosomes in *Hei10*^{mei4/mei4} mutant nuclei (**Figure 4C,D**). Specifically, for

54.4% of individual chromosomes from late pachytene *Hei10*^{mei4/mei4} spermatocyte nuclei

(marked by strong H1t staining), neither the first nor the second brightest focus was ≥1.5-fold

brighter than the third brightest focus along the same chromosome, and only 17.0% of individual

chromosomes had RNF212B foci that were ≥2-fold brighter than the third brightest focus along

the same chromosome (n = 93/171 and 29/171 chromosomes, respectively; 9 nuclei; **Figure**

4A,C). The most differentiated RNF212B foci were on average only 1.6-fold brighter (with a

maximum of 4.2-fold) relative to the third brightest focus along the same chromosome in late

pachytene $Hei10^{mei4/mei4}$ spermatocyte nuclei; whereas in wild-type mid-pachytene nuclei,

RNF212B foci at crossover sites were on average 6.7-fold brighter relative to the brightest other

focus along the same chromosome (**Figure 4D**; 33 wild-type chromosomes with two crossover

foci were used for comparison to $Hei10^{mei4/mei4}$ chromosomes; 8 nuclei). Thus, crossover-specific

patterning of RNF212B appears to be largely dependent on HEI10. Reciprocally,

immunostaining for HEI10 in $Rnf212b^{-/-}$ mutant spermatocytes revealed that formation of HEI10

foci is dependent on RNF212B (**Figure 4E**).

Our localization analysis is incompatible with RNF212B and RNF212 functioning as an obligate

heterocomplex of fixed stoichiometry. Even so, immunostaining analysis in $Rnf212b^{-/-}$ and

Rnf212$^{-/-}$ mutants indicated an interdependent relationship. RNF212 immunostaining was

diminished in *Rnf212b*$^{-/-}$ mutant meiocytes and, likewise, RNF212B immunostaining was

diminished in *Rnf212*$^{-/-}$ cells (**Figures 4F,G** and **S9**). Immunoblotting revealed that protein levels

of RNF212 and, to a greater extent, RNF212B were reduced in testis extracts from *Rnf212b*$^{-/-}$

and *Rnf212*$^{-/-}$ mutants, respectively, relative to wild-type controls (**Figure 4H**).

Interaction between RNF212B and RNF212 was detected by yeast two-hybrid assay (Y2H), and

self-interaction for both proteins was also inferred, although RNF212 self-interaction appeared

to be much weaker than that of RNF212B (**Figure 4I,J**). Thus, RNF212B and RNF212

physically interact and are mutually dependent for protein stability and thus chromosomal

localization. Additional analysis showed that an intact RING finger domain was essential for

RNF212B self-interaction by Y2H, for the stability of RNF212B and RNF212 *in vivo*, and thus for

the pro-crossover functions of RNF212B (**Figures S10** and **S11**). Interaction between HEI10

and RNF212B (but not HEI10 and RNF212) was also detected, which might reflect their

spatiotemporal colocalization at designated crossover sites (**Figure 4I,J**).

337   **Similar but distinct localization of RNF212B and RNF212 in mutants defective for**

338   **recombination and synapsis**

339   The relationship between RNF212B and RNF212 was further explored by analyzing their

340   localization in mutants defective for recombination or synapsis. Repair of SPO11-induced DSBs

341   is required for the alignment and synapsis of homologs. Thus, *Spo11*[-/-] mutant meiocytes are

342   generally defective for synapsis; however, extensive stretches of SC can form between non-

343   homologous axes [49, 50]. In *Spo11*[-/-] spermatocytes, RNF212B staining was still detected

344   specifically on synapsed regions marked by SYCP1, as previously seen for RNF212 (**Figure**

345   **5A**)[15]. However, intensities of RNF212B immunofluorescence measured in *Spo11*[-/-] nuclei with

346   late zygotene/early pachytene-like morphologies (well-developed SYCP3-staining axes and

347   extensive synapsis, albeit non-homologous) were ~50% lower than those in equivalent early-

348   pachytene *Spo11*[+/+] nuclei (**Figure 5B**). By contrast, intensities of SC-associated RNF212

349   immunostaining were comparable for *Spo11*[-/-] and *Spo11*[+/+] nuclei. Correspondingly, the

350   normalized ratio of RNF212B / RNF212 immunofluorescence intensity on synapsed regions was

351   reduced by ~50% in *Spo11*[-/-] spermatocytes (**Figure 5C**). Thus, SPO11-initiated recombination

352   differentially influences the association of RNF212B and RNF212 with synaptonemal

complexes, with only RNF212B showing a significant dependence.

SYCP1 is the major component of the SC central region. Although *Sycp1*[-/-] mutant meiocytes fail

to assemble SCs, recombination initiates normally and homolog axes become coaligned [51]. In

Sycp1[-/-] spermatocytes, RNF212B staining was barely detectable in early/mid zygotene-like

nuclei indicating that initial association with meiotic chromosomes is strongly dependent on the

SC central region (**Figure S12A**). However, relatively dim RNF212B foci could be detected

between aligned homologous axes in late zygotene/early pachytene-like nuclei that had

extensive homolog coalignment (**Figures 5D**). The majority of these RNF212B foci localized to

recombination sites, revealed by a high level of colocalization with RPA2 (71.0 ± 8.0%; mean ±

S.D., *n* = 8 nuclei), as seen previously for RNF212 localization in *Sycp1*[-/-] spermatocytes

(**Figures 5D** and **S12B**)[15]. Thus, normal localization of RNF212B to prophase-I chromosomes is

strongly dependent on synapsis, with respect to both timing and abundance, but RNF212B can

independently localize to recombination sites, albeit inefficiently.

RNF212B focus numbers in *Sycp1*[-/-] nuclei were reduced 2.4-fold relative to *Sycp1*[+/+] controls

(192.1 ± 18.3 foci in *Sycp1*[+/+] versus 81.4 ± 19.3 in *Sycp1*[-/-]; mean ± SDs; **Figure 5E**).

Correspondingly, the degree of RPA2 colocalization with RNF212B was reduced 2.4-fold (from

67.3 ± 3.7% in *Sycp1*[+/+] to 27.9 ± 6.0% in *Sycp1*[-/-]; means ± S.D.; n = 7 and 8 nuclei,

respectively; p = 0.0002, Mann-Whitney test) and RNF212B-RPA2 co-foci were reduced 1.9-

fold (109.4 ± 7.4 foci in *Sycp1*[+/+] versus 57.9 ± 14.7 in *Sycp1*[-/-]; mean ± SDs; **Figures 5F** and

S12C). In contrast, in the same nuclei, numbers of RNF212-RPA2 co-foci remained essentially

unchanged (122.4 ± 7.8 foci in *Sycp1*[+/+] versus 131.3 ± 17.4 in *Sycp1*[-/-]; mean ± SDs) despite a

1.5-fold reduction in total RNF212 focus numbers (220.3 ± 13.9 foci in *Sycp1*[+/+] versus 146.9 ±

21.4 in *Sycp1*[-/-]; mean ± SDs; **Figures 5E** and **S12C**). Moreover, the degree of RPA2-RNF212

colocalization was barely altered (75.2 ± 2.9% in *Sycp1*[+/+] versus 64.3 ± 11.6% in *Sycp1*[-/-]; mean

± SDs; p = 0.0578, Mann-Whitney test; **Figure 5F**).

We infer that the localization of RNF212B to recombination sites shows a greater dependence

on synapsis than does RNF212. Consequently, this differential dependency resulted in reduced

colocalization of RNF212B and RNF212 foci in *Sycp1*[-/-] spermatocytes (61.8 ± 9.0% of

RNF212B foci colocalized with RNF212, and 33.9 ± 6.3% of RNF212 foci colocalized with

RNF212B; versus 95.8 ± 2.4% and 83.8 ± 8.6% in *Sycp1*$^{+/+}$ nuclei; mean ± SDs; **Figure S12D**).

MLH3 and MLH1 constitute the MutLγ complex that facilitates the maturation and crossover-

specific resolution of double-Holliday junctions [29, 30]. Consistently, *Mlh3*$^{-/-}$ meiocytes are

proficient for synapsis and early steps of meiotic recombination, but defective for crossing over

[52]. We previously showed that MutLγ constrains HEI10 localization both temporally and spatially;

in *Mlh3* mutant spermatocytes HEI10 foci appear much earlier than in wild type (during

zygotene), are much more numerous (~90 per nucleus), and persist throughout pachytene [16]. In

contrast, RNF212B staining in *Mlh3*$^{-/-}$ spermatocytes was comparable to *Mlh3*$^{+/+}$ controls: focus

numbers peaked in early pachynema, then diminished as pachynema progressed with one or

two large RNF212B foci emerging on each SC (**Figures 5G**,**H** and **S12E,F**). Numbers of

amplified crossover-specific RNF212B foci in late pachytene *Mlh3*$^{-/-}$ nuclei were slightly lower

than in wild-type *Mlh3*$^{+/+}$ controls, possibly reflecting reduced stability or altered efficiency of

formation (**Figure S12F**). In addition, unlike wild type, some smaller foci remained on the SCs

until late pachynema in *Mlh3*$^{-/-}$ nuclei suggesting an additional signal following the maturation of

crossover sites may trigger the complete loss of RNF212B from noncrossover sites.

Nonetheless, this analysis indicates that the differentiation of RNF212B into crossover-specific

foci precedes crossing over and does not require crossover-specific factor, MutLγ. Importantly,

we can also infer that while crossover-specific patterning of RNF212B requires HEI10 (above,

Figure 4A-D), it does not require HEI10 *per se* to undergo crossover-specific patterning, i.e.

crossover-specific patterning of HEI10 appears to be downstream of initial crossover

designation and requires a maturation step that is dependent on MutLγ.

RNF212 staining in *Mlh3*$^{-/-}$ mutant spermatocytes was comparable to wild-type cells in early

pachynema, but the differentiation of crossover-specific foci was not as conspicuous; and

overall focus numbers remained relatively high throughout pachynema (**Figures 5G**,**I**, and

S12G). Moreover, numerous RNF212 foci persisted on synapsed regions in early diplotene

Mlh3$^{-/-}$ nuclei in which RNF212B staining was barely detectable. Together, localization studies in

mutant contexts extend our inference that the dynamics and regulation of RNF212B, RNF212,

and HEI10 are distinct.

RNF212B is essential for crossing over in oocytes

Rnf212b$^{-/-}$ mutation in females also caused sterility but unlike males, in which gametes were

absent (**Figure 1B**), the ovaries of 18 days postpartum (dpp) mice contained large numbers of

oocytes (**Figure 6A,B**). However, analysis of MLH1 foci in prophase-I nuclei (from fetal ovaries,

Figure 6C), and chiasmata in metaphase-I oocytes (from adult ovaries, **Figure 6D**) revealed

that a severe crossover defect is common to *Rnf212b$^{-/-}$* mutants of both sexes: MLH1 foci were

completely absent, and chiasmata were reduced over 20-fold (1.11 ± 0.92 chiasmata per

nucleus in *Rnf212b$^{-/-}$*, *n* = 38 oocytes; compared to 24.6 ± 2.4 in *Rnf212b$^{+/+}$*, *n* = 28 oocytes;

means ± SDs).

Sexually dimorphic chromosomal localization of RNF212B, RNF212, and HEI10

RNF212B: In fetal oocytes, the initial immunostaining pattern of RNF212B during

zygonema was similar to that observed in spermatocytes, with foci localized specifically to

regions of homolog synapsis (**Figure 6E,F**). Focus numbers also peaked in early pachynema

(defined as pachytene-stage oocytes with ≤4 MLH1 foci from embryonic day E16.5 ovaries) but

were 1.6-fold greater than in spermatocytes at the same stage (**Figure 6F**; 279.0 ± 50.4 foci per

oocyte nucleus, *n* = 11, versus 175.9 ± 15.2 foci per spermatocyte, *n* = 27; mean ± S.D.). Also

distinct from spermatocytes, high numbers of RNF212B foci persisted along SCs throughout

pachynema both during and after the emergence of amplified foci at designated crossover sites.

Indeed, even after a full complement of crossover sites had matured in mid/late pachytene

oocytes (with ≥20 MLH1 foci, **Figure 6F**), RNF212B foci averaged 229.5 ± 45.0 per nucleus

(from E18.5 ovaries; mean ± SD; n = 22; **Figure 6F**). Furthermore, RNF212B remained on

synapsed regions throughout diplonema, as homologs desynapsed, and disappeared only as

oocytes entered the dictyate stage (**Figure 6E,F**).

Given the distinct dynamics of RNF212B in oocytes, we also analyzed the timing and degree of

amplification of RNF212B and RNF212 at crossover sites (**Figure 6G,H**). In early/mid

pachytene oocytes (with ≤10 MLH1 foci from E18.5 ovaries), although crossover-associated

RNF212B foci were on average 1.8-fold brighter than other (noncrossover) RNF212B foci in the

same nuclei (1.76 ± 0.58 for 28 crossover foci versus 0.98 ± 0.40 for 1,375 other foci from 7

nuclei; means ± SDs; $p < 0.0001$, Mann-Whitney test; **Figure 6H**), their intensity distributions

overlapped; indeed, we readily observed individual SCs with crossover and non-crossover

RNF212B foci of similar intensities (**Figure 6G**, upper panels).

By mid/late-pachynema (oocytes with ≥20 MLH1 foci from E18.5 ovaries), crossover-associated

RNF212B foci were now 2.5-fold brighter than other RNF212B foci, suggesting continued

amplification after initial designation (2.63 ± 1.13 for 250 crossover foci versus 1.07 ± 0.43 for

1,468 other foci from 9 nuclei; means ± SDs; $p < 0.0001$, Mann-Whitney test; **Figure 6H**). Thus,

in oocytes, growth/amplification of RNF212B may be coincident with crossover-site maturation

marked by the emergence of MLH1 foci. This contrasts spermatocytes, in which amplified

RNF212B foci clearly precede the appearance of MLH1 (**Figure 3**). Again, while crossover

associated RNF212B foci were brighter than other foci at the population level, they were not

always the brightest foci along the same SC (**Figure 6G**, lower panels). Moreover, a consistent

pattern of RNF212B foci at and surrounding designated crossover sites was not observed:

some crossover foci were adjacent to other similarly bright foci, others were embedded in a

domain of fainter foci, and others by gaps of diminished RNF212B staining (**Figure 6G**).

Notably, the intensities of other (noncrossover) RNF212B foci and the total signal intensity of

RNF212B per oocyte nucleus were largely unchanged throughout pachynema (**Figure 6H** and

S13A), in sharp contrast to the loss of RNF212B foci from non-crossover sites in spermatocytes

(**Figures 2, 3** and **S7**).

RNF212: Abundant general staining of RNF212 was also retained throughout

pachynema in oocytes, but amplification of RNF212 at crossover sites appeared less

pronounced than that of RNF212B (**Figure 6G**). Crossover-associated RNF212 foci in early/mid

pachynema were only 1.3-fold brighter than other foci (1.32 ± 0.47 for 30 crossover foci versus

0.99 ± 0.45 for 1.596 other foci from 7 nuclei; means ± SDs; p = 0.0001, Mann-Whitney test). By

mid/late pachynema, crossover-associated RNF212 were now 1.6-fold brighter than other foci in

the same nuclei (1.87 ± 1.04 for 245 crossover foci versus 1.19 ± 0.75 for 1,580 other foci form

9 nuclei; means ± SDs; p < 0.0001, Mann-Whitney test) and 1.9-fold brighter than those in

early/mid pachytene consistent with continued amplification (**Figure 6H**). Intriguingly, the

intensity of RNF212 at noncrossover sites slightly increased in mid/late pachynema nuclei

suggesting that general loading of RNF212 along SCs may continue during pachytene (0.99 ±

0.45 to 1.19 ± 0.75 from early/mid to mid/late pachynema; means ± SDs; p < 0.0001, Mann-

Whitney test; **Figure 6H**). Moreover, total signal intensity of RNF212 per oocyte nucleus was

largely unchanged between early/mid to mid/late pachynema, as seen for RNF212B (**Figure**

S13A).

HEI10: Chromosomal localization of HEI10 was also sexually dimorphic. Unlike

spermatocytes, prominent HEI10 foci were already apparent in zygotene-stage oocytes,

specifically at regions of synapsis (**Figure S13B**). Focus numbers increased as synapsis

ensued, peaking in early pachynema (defined as pachytene-stage oocytes with ≤4 MLH1 foci)

with 123.9 ± 28.7 foci per nucleus (mean ± SD; n = 18 nuclei; **Figure 7A,D**). At this stage,

HEI10 foci were unevenly spaced and highly variable in size and intensity; with configurations

suggestive of growth, shrinkage, and/or fusion of foci as might be predicted by coarsening

models (**Figure 7A,F** and **S13B,D**). Focus numbers then reduced as pachynema progressed,

showing a largely crossover-specific pattern by mid pachynema (defined as pachytene-stage

oocytes with ≥20 MLH1 foci from E17.5 ovaries; **Figure 7 B–E**). At this stage, 86.2 ± 9.6 % of

HEI10 foci colocalized with MLH1 foci and 98.0 ± 2.7 % of MLH1 foci colocalized with HEI10

foci (means ± SDs; n = 14 nuclei).

Crossover-associated HEI10 foci in early-to-mid pachynema (pachytene-stage oocytes with 5-

15 MLH1 foci from E16.5 ovaries) were on average 2.3-fold brighter than other foci in the same

nuclei (1.80 ± 0.53 for 96 crossover foci versus 0.79 ± 0.48 for 1,089 other foci from 9 nuclei;

means ± SDs; *p* < 0.0001, Mann-Whitney test; **Figure 7B,F**). Even very faint emergent MLH1

foci in early pachynema (pachytene-stage oocytes with ≤4 MLH1 foci from E16.5 ovaries)

tended to be associated with slightly larger HEI10 foci suggesting that amplification

accompanies crossover designation (1.26 ± 0.39 for 33 crossover foci versus 1.00 ± 0.45 for

3,817 other foci from 26 nuclei; means ± SDs; *p* = 0.0003, Mann-Whitney test; **Figure 7A,F**).

However, MLH1 foci emerged when HEI10 foci still outnumbered crossovers by more than 3:1,

i.e. before HEI10 *per se* had attained a crossover-specific distribution. Moreover, while

crossover-associated HEI10 foci were on average 1.9-fold brighter relative to the brightest other

focus along the same chromosome in early/mid pachynema (**Figure 7G**), similarly bright HEI10

foci were often observed along the same SC (**Figure 7B**, magnified panels, and **Figure**

S13B,C). After mid pachynema (oocytes with ≥20 MLH1 foci from E16.5 ovaries), crossover-

associated HEI10 foci were 3.1-fold brighter than the foci detected in early pachynema and 1.7-

fold brighter than the crossover-specific foci detected in early/mid pachynema (3.06 ± 1.00 for

186 crossover foci from 6 nuclei ; mean ± SD; **Figure 7F**), suggesting continued amplification of

HEI10 after crossover designation. The intensities of other remaining (noncrossover) HEI10 foci

in mid pachynema were comparable to those of the foci detected in early pachynema, though

these foci were eventually lost (**Figure 7D-F**).

Together, these data reveal profound sexual dimorphism in the localization dynamics of

RNF212B, RNF212, and HEI10. Importantly, in oocytes, designation of crossover sites does not

involve general depletion of RNF212B and RNF212 from noncrossover sites; and amplification

may be a coincident or downstream event. In contrast, as crossover associated HEI10 foci

amplify, foci at other sites progressively diminish in number. While the timing of HEI10

amplification may be consistent with a role in crossover-site designation, crossover sites are

maturing well before a crossover-specific pattern of HEI10 is attained.

DISCUSSION

COR Patterning vis-a-vis Crossover Patterning

An elementary coarsening model, that can explain crossover patterning in Arabidopsis, posits

that a finite amount of HEI10 protein (fixed amount per unit length) diffuses along SCs and

becomes progressively absorbed into one or a few amplified foci which then designate which

recombination sites will mature into crossovers [35, 36]. Our data in mouse are hard to reconcile

with this model, revealing a more complex scenario in which pattering of a given COR protein

may precede, accompany, or follow the designation of crossover sites; can occur with or without

depletion of CORs from non-crossover sites; and shows striking sexual dimorphism

(summarized in **Figure 7H**).

Global and local regulation of recombination by RNF212B, RNF212, and HEI10

What might be the function(s) of the observed dynamics of RNF212, RNF212B and HEI10? Our

analysis emphasizes the interdependencies and distinctions between the three mammalian

CORs and their global and local functions that help coordinate key events of meiotic prophase I.

Most evident is regulating the progression of recombination occurring in the context of the SC.

Following initial DNA strand-exchange and homolog pairing, RNF212-RNF212B associates with

nascent SCs and acts to stabilize ZMM factors and pause the progression of recombination. In

this way, RNF212-RNF212B may act to protect nascent recombination intermediates from being

prematurely dissociated by the Bloom complex, which is inferred to mediate the default

noncrossover outcome of recombination via synthesis-dependent strand annealing [31-34, 53]. This

early function of RNF212-RNF212B appears to promote and stabilize synapsis, possibly acting

as a kind of proofreading mechanism that selectively reinforces SC assembled at recombination

sites, i.e. between homologous chromosomes. As prophase-I progresses, connection of

homologs by recombinational interactions is superseded by the SC. RNF212-RNF212B could

mediate this hand-off by ensuring that recombinational connections are not resolved until

mature SC has formed.

Pausing the progression of SC-associated recombination events may also be a prerequisite for

crossover/noncrossover differentiation and/or to allow time for crossover sites to mature.

Importantly, RNF212-RNF212B renders the progression of recombination dependent on HEI10

(**Figure 4**) and additional pro-crossover factors including kinase CDK2, and the CNTD1-PRR19

complex [16, 17, 45, 54, 55]. In the absence of HEI10, foci of RNF212-RNF212B and ZMM proteins

remain abundant and undifferentiated along SCs, and recombination continues to be stalled

throughout pachytene, with DSB repair being completed only as the SC central region

disassembles along with RNF212-RNF212B.

At designated crossover sites, local protection of intermediates via RNF212-RNF212B may

enable crossover-specific events including dHJ formation and recruitment of the crossover-

specific resolution machinery, organized around the MutLγ endonuclease. Local stabilization of

SC via RNF212-RNF212B may also explain why crossover sites are the last sites to desynapse

during diplotene [56]. We have suggested that patches of SC retained at crossover sites may help

coordinate the DNA events of crossing over with exchange of the underlying chromosome axes

to form chiasmata.

Distinct and interdependent functions of RNF212, RNF212B and HEI10

Our ability to discern potentially distinct functions of RNF212 and RNF212B is compromised by

their interdependence for protein stability, i.e. *Rnf212* and *Rnf212b* single mutant phenotypes

likely reflect diminished function of both proteins. However, localization dynamics of RNF212

and RNF212B reveal divergent behaviors of the two proteins that point to distinct function and

regulation.

Notably, in early pachytene spermatocytes, RNF212 foci outnumber RNF212B and up to 40%

don't detectably colocalize. Also, the amplification of crossover specific RNF212B foci was much

stronger than for RNF212 and occurred earlier, with the suggestion that differentiation may

occur via distinct processes, e.g. redistribution to crossover sites for RNF212B, versus the loss

from noncrossover sites for RNF212 (or via different proportions of redistribution and loss).

Thus, although RNF212 and RNF212B interact, these observations argue against the existence

of an obligate heterocomplex of fixed stoichiometry and suggest that the activity of RNF212B

may be modulated by switching binding partners as it accumulates at designated crossover

sites. For example, RNF212B-RNF212B self-interaction could become prominent, or RNF212B

could partner with HEI10 in order to locally accumulate and modulate E3 ligase activities at

designated crossover sites. Consistent with the latter possibility, HEI10-RNF212B interaction is

detected by Y2H, and differentiation of RNF212B foci is both dependent on HEI10 and

temporally indistinguishable from its appearance at prospective crossover sites (**Figures 4** and

S8).

Localization dynamics in mutant backgrounds also point to distinct functions and regulation of

the three CORs. For example, RNF212 localization along SCs remains robust in the absence of

SPO11-catalyzed DSBs while localization of RNF212B is much less efficient, suggesting that

DSB signaling may stabilize RNF212B. In this respect, RNF212B may be like budding yeast

Zip3, which requires phosphorylation at S/T-Q consensus sites for the DNA-damage response

(DDR) kinases Mec1^{ATR}/Tel1^{ATM} for normal localization [57]. Further distinctions are seen in the

absence of synapsis, with RNF212 showing a robust ability to localize to recombination sites,

while RNF212B localization was much weaker. Thus, inputs from both DSBs and synapsis

appear to stabilize RNF212B and thereby coordinate its activity in space and time, while

RNF212 might act more as an anchor for localization to SCs and recombination sites. We

previously showed that HEI10 foci in spermatocytes are largely dependent on both DSBs and

synapsis [16], consistent with our inference that HEI10 foci assemble only as crossover sites

differentiate, dependent on RNF212-RNF212B.

Defective crossover maturation in the *Mlh3* mutant impacts the dynamics of RNF212, RNF212B

and HEI10 to varying degrees. Although crossover-specific patterning of RNF212-RNF212B foci

still occurs in *Mlh3* spermatocytes, numerous small foci of RNF212, and to a lesser extent

RNF212B, also persist along SCs implying that a MutLγ-dependent signal associated with the

maturation of crossover sites enhances the general loss of RNF212-RNF212B from

noncrossover sites. We previously showed that HEI10 patterning is severely perturbed in *Mlh3*

mutant spermatocytes, with abnormally high numbers of foci (~90 per nucleus) forming much

earlier than normal (in zygonema) and then persisting at recombination sites until chromosomes

desynapse [16]. Thus, although HEI10 is required for crossover patterning of RNF212-RNF212B

foci, this function does not require crossover-specific patterning of HEI10 itself, i.e. crossover-

specific localization of HEI10 appears to be downstream of initial crossover designation.

Moreover, crossover-specific patterning of HEI10 requires an upstream maturation step that is

dependent on the MutLγ complex, likely the dHJ formation function described by Premkumar et

al. [58]. Consistently, in oocytes, MLH1 foci can be detected before HEI10 foci have attained a

crossover-specific pattern (**Figure 7A** and **S13BC**).

Collectively, these data suggest that RNF212, RNF212B and HEI10 may function as apical

effectors that coordinate meiotic prophase by integrating signals from synapsis, DSB repair, cell

cycle kinases, and maturating crossover sites (summarized in **Figure 7**).

Sexual dimorphism

Patterning of all three CORs along SCs shows striking sexual dimorphism. In spermatocytes,

complete differentiation of RNF212B results in only large crossover-specific foci; while in

oocytes, numerous foci persist throughout pachytene even after amplified crossover-specific

foci have formed and crossover sites have matured (marked by the appearance of MLH1 foci;

Figures 2A,H, and **6E,G**). RNF212 dynamics are similarly dimorphic. Thus, complete

differentiation of RNF212B or RNF212 is not a prerequisite for the patterning and maturation of

crossover sites. Possibly, accumulation of a threshold level of RNF212B at a given

recombination site is sufficient for it to mature into a crossover. However, weak amplification and

incomplete differentiation in females could render RNF212B crossover foci less stable and

potentially reversible, possibly accounting for the inefficiency of crossover maturation

characterized in human oocytes that can result in unconnected (achiasmate) homologs, or

homologs connected only by a single telomere-proximal chiasma, configurations that are prone

to segregation error [8, 9]. HEI10 shows striking temporal and spatial dimorphism with foci

emerging in mid pachytene spermatocytes already showing a crossover distribution; while in

oocytes, numerous foci first form during zygotene and then attain a crossover-specific pattern in

pachytene.

What could account for the sexually dimorphic behavior of mammalian CORs? A pertinent

difference may be the duration of pachytene, which lasts almost 7 days in mouse spermatocytes

[59] compared to just 1-2 days in oocytes [60]. Thus, if patterning dynamics are relatively slow,

differentiation may remain incomplete in oocytes. Also, oocyte SCs are longer than

spermatocytes and CORs appear to be generally more abundant. COR dimorphism might also

reflect dimorphism in the DNA-damage checkpoint and gamete quality control. We previously

showed that RNF212 is required for oocyte quality control as dictyate oocytes assemble into

primordial follicles [61], while the primary checkpoint in spermatocytes occurs much earlier, as the

sex-body assembles in mid pachytene [62].

FIGURES

Figure 1. RNF212B is required for crossing over via stabilization of ZMMs in mouse

spermatocytes.

(A) Reduced testis size in *Rnf212b$^{-/-}$* mutant males. Red bars indicate means. ****p ≤ 0.0001 for

a two-tailed *t* test. n, numbers of mice analyzed.

(B) Defective spermatogenesis in *Rnf212b$^{-/-}$* mutant males. Seminiferous tubule sections from

wild-type and *Rnf212b$^{-/-}$* testes stained with hematoxylin and eosin. rSt, round spermatid; eSt,

elongated spermatid. Arrowheads indicate metaphase cells.

(C) Chiasmata are diminished in *Rnf212b$^{-/-}$* spermatocytes. Diakinesis/metaphase-I stage

spermatocytes from wild-type and *Rnf212b$^{-/-}$* testes stained with DAPI.

(D) MLH1 foci are absent in *Rnf212b$^{-/-}$* spermatocytes. Mid-late pachytene-stage spermatocyte

nuclei from wild-type and *Rnf212b$^{-/-}$* testes immunostained for SYCP3 and MLH1.

(E) Representative images of successive prophase-I stages in chromosome spreads of wild-

type spermatocytes immunostained for SYCP3.

(F) Distributions of spermatocytes in successive prophase-I stages based on SYCP3 staining.

>250 unselected SYCP3-positive cells were analyzed per mouse. Bar graphs indicate means of

two mice with plots for each mouse. *p* = 0.001 for *G* test (total cells analyzed: 578 *Rnf212b$^{+/+}$*

cells and 549 *Rnf212b*$^{-/-}$ cells).

(G) Autosomal and X-Y synapsis in wild-type and *Rnf212b*$^{-/-}$ spermatocytes. Pachytene-stage

spermatocyte nuclei immunostained for SYCP3 and H1t are shown. The magnified images

show X-Y chromosomes.

(H) Premature desynapsis of X-Y chromosomes in *Rnf212b*$^{-/-}$ and *Rnf212*$^{-/-}$ spermatocytes. ns,

not significant ($p > 0.05$); ****$p \le 0.0001$ for Fisher's exact tests. Total numbers of cells analyzed

from 3 mice of each genotype are indicated below the X axis.

(I) Wild-type and *Rnf212b*$^{-/-}$ spermatocyte nuclei at the indicated stages immunostained for

SYCP3 and a ZMM factor MSH4. Representative chromosomes are magnified.

(J) MSH4 focus counts. Red bars indicate means ± SDs. ***$p \le 0.001$; ****$p \le 0.0001$ for two-

tailed Mann-Whitney tests. n, numbers of nuclei analyzed.

Scale bars, 10 μm for images of full nuclei and 2 μm for magnified panels.

Figure 2. RNF212B localization to synaptonemal complexes and recombination sites in

prophase I spermatocytes.

(A) RNF212B localization in spermatocyte nuclei at successive prophase I stages. Images show

surface-spread prophase-I spermatocyte nuclei immunostained for SYCP3, SYCP1, RNF212B

and H1t. Arrowheads highlight large RNF212B foci. The arrow indicate synapsed pseudo-

autosomal region between the X-Y chromosomes.

(B) Numbers of RNF212B foci per nucleus in spermatocytes at successive prophase I stages.

Red bars indicate means. Means of focus numbers are shown below the graph. Lept.,

leptonema; Zyg., zygonema; pach., pachynema; dip., diplonema. Numbers of nuclei analyzed

were 8, 12, 27, 29, 29, 19 and 11 for in leptonema, zygonema, early pachynema, mid

pachynema, late pachynema, early diplonema and mid diplonema, respectively.

(C) RNF212B localizes to the central region of the synaptonemal complex. SIM images of early

and late pachytene spermatocytes immunostained for SYCP3 and RNF212B. Arrowheads

highlight large RNF212B foci.

(D) RNF212B colocalization with RNF212 in late zygonema. SIM images of a late zygotene

spermatocyte immunostained for SYCP3, RNF212B and RNF212.

(E) Quantification of RNF212B colocalization with RNF212. Left, focus counts. Right, degree of

colocalization. Black and red bars indicate means. 3 late-zygotene and 8 early-pachytene nuclei

were analyzed by SIM imaging.

(F) RNF212B colocalization with MSH4 in early pachynema. SIM images of an early pachytene

spermatocyte immunostained for SYCP3, RNF212B and MSH4.

(G) Quantification of RNF212B colocalization with MSH4. Left, focus counts. Right, degree of

colocalization. Black and red bars indicate means. 7 early pachytene nuclei were analyzed by

SIM imaging.

(H) RNF212B colocalization with MLH1. A late pachytene spermatocyte immunostained for

SYCP3, RNF212B and MLH1. Arrowheads indicate crossover sites.

Magnified images in (A), (C), (D), (F), and (H) show representative chromosomes. Scale bars,

10 µm for full nuclei and 2 µm for magnified images.

Figure 3. Distinct timing and development of crossover-specific foci of HEI10, RNF212B

and RNF212

(A) HEI10 forms crossover foci by mid-pachynema. Pachytene spermatocytes nuclei

immunostained for SYCP3, HEI10, MLH1 and H1t. Top images show a representative H1t-

negative, mid-pachytene nucleus in which small MLH1 foci are emerging on just a few

chromosomes (indicated by arrows) whereas bright HEI10 foci are present on all autosomes.

Bottom images show a representative H1t-positive late-pachytene nucleus in which both MLH1

and HEI10 form clear, overlapping crossover-specific foci on all autosomes.

(B) Quantification of HEI10 and MLH1 foci in pachytene spermatocytes confirms that MLH1 foci

develop only in nuclei with a full complement of HEI10 foci. 25 pachytene nuclei were randomly

selected and analyzed.

(C) Differentiation of RNF212B at prospective crossover sites is stronger than RNF212.

Airyscan images of early, mid and late pachytene spermatocytes nuclei immunostained for

SYCP3, RNF212B, RNF212 and HEI10. In all cases, RNF212B and RNF212 image exposures

are matched. Arrowheads indicate crossover sites marked by HEI10.

(D) Quantification of focus intensities for RNF212B and RNF212. Black bars indicate means.

****$p \leq 0.0001$ for two-tailed Mann-Whitney tests. 8 nuclei for each stage were analyzed by

Airyscan imaging. Total numbers of foci analyzed: 2,449 RNF212B foci and 2,513 RNF212 foci

in early pachynema; 194 crossover foci, 2,004 other RNF212B foci and 2,291 other RNF212

foci in mid pachynema; 206 crossover foci and 988 other RNF212 foci in late pachynema. All, all

foci; COs, crossover foci colocalized with HEI10 foci; Others, other foci that don't colocalize with

HEI10 foci. ND, not detected. Quantification of focus intensities in representative individual

nuclei in (D) is shown in **Figure S7A**.

(E) Per chromosome analysis for autosomes in mid-pachytene spermatocyte nuclei. Intensity of

each crossover-associated RNF212B or RNF212 focus relative to that of the brightest other

(non-crossover) focus along the same chromosome plotted separately (left) and in two

dimensions (right). Red dashed lines indicate a relative focus intensity of 2. Detailed

representation of a per chromosome analysis is shown in **Figure S7B**. ****$p \leq 0.0001$ for two-

tailed Mann-Whitney tests. 186 crossover foci from 8 mid-pachytene nuclei were analyzed. Two

crossover-associated RNF212B foci with no other RNF212B foci along the same chromosome

were excluded from the plots.

(F) RNF212B differentiation without detectable RNF212 differentiation during early to mid

pachyenema. Airyscan images of an early- to mid-pachytene spermatocyte nucleus, with 8

HEI10 foci, immunostained for SYCP3, RNF212B, RNF212 and HEI10 are shown. Arrowheads

indicate crossover sites marked by HEI10.

Magnified images in (C) and (F) show representative chromosomes. Scale bars, 10 µm for full

nuclei and 2 µm for magnified images.

Figure 4. Interdependent relationship between HEI10, RNF212 and RNF212B

(A) Persistent RNF212B staining in *Hei10$^{mei4/mei4}$* spermatocytes. Images of a late-pachytene

nucleus immunostained for SYCP3, RNF212B, RNF212 and H1t. The magnified panels show

representative chromosomes with: (i) no apparent differentiation of RNF212B foci; or some

differentiation, indicated by arrowheads. Scale bars, 10 μm in images of full nuclei and 2 μm in

the magnified panels.

(B) Focus counts of RNF212B. Red bars indicate means. Total numbers of nuclei analyzed are

indicated below the X-axis. For late stages, only late pachytene nuclei were analyzed for

Hei10$^{+/+}$ whereas both late pachytene and early diplotene nuclei were analyzed for *Hei10$^{mei4/mei4}$*

because of the early desynapsis of *Hei10$^{mei4/mei4}$* cells [16]. ns, not significant ($p > 0.05$); ****$p \leq$

0.0001 for two-tailed Mann-Whitney tests.

(C) Differentiation of RNF212B foci. Autosomes in late-pachytene *Hei10$^{+/+}$* and *Hei10$^{mei4/mei4}$*

spermatocyte nuclei were categorized into four classes based on the degree of RNF212B

differentiation. Strong, weak or no differentiation represents autosomes where the brightest

RNF212B foci were ≥2-fold, 1.5-2-fold, or <1.5-fold brighter relative to the third brightest focus

along the same synaptonemal complex, respectively. Complete differentiation is autosomes that

have only one or two bright RNF212B foci. 8 late-pachytene $Hei10^{+/+}$ and 9 late-pachytene

$Hei10^{mei4/mei4}$ nuclei were analyzed by Airyscan imaging. $p < 0.0001$ for a G test.

(D) Intensity of the brightest RNF212B foci relative to the third brightest focus along the same

chromosome in late pachytene $Hei10^{mei4/mei4}$ and mid pachytene $Hei10^{+/+}$ spermatocyte nuclei.

Red bars indicate means. Nuclei analyzed were the same as in (C) and **Figure 3E**, respectively,

but chromosomes with two crossover foci were analyzed for $Hei10^{+/+}$ spermatocytes. Total

numbers of foci analyzed are indicated below the X-axis. ****$p \leq 0.0001$ for two-tailed Mann-

Whitney test.

(E) Crossover-specific immunostaining foci of HEI10 foci were absent from chromosome

spreads of $Rnf212b^{-/-}$ spermatocytes. Mid/late pachytene nuclei from wild-type and $Rnf212b^{-/-}$

spermatocytes were immunostained for SYCP3 and HEI10. Scale bars, 10 µm.

(F and G) RNF212 (F) and RNF212B (G) are interdependent for chromosomal localization.

Early pachytene spermatocyte nuclei from the indicated genotypes were immunostained for

SYCP3 and RNF212 (F) or RNF212B (G). Scale bars, 10 µm.

(H) Interdependent protein stability of RNF212B and RNF212. Whole-testis extract from

indicated genotypes were subjected to immunoblotting for RNF212 and RNF212B. Histone H3

is a loading control.

(I) Yeast two-hybrid assay showing interaction between RNF212B, RNF212 and HEI10, and

self-interactions for RNF212B and RNF212. Yeast cells expressing Gal4 activation (Gal4-AD)

and binding (Gal4-BD) domain fused to the indicated proteins, or empty vectors as controls,

were spotted with the indicated dilutions. Aureobasidin A selection was used to detect activation

of the *AUR1-C* reporter. * indicates less stringent selection for *ADE2* and *HIS3* reporters.

(J) A summary of interactions between RNF212B, RNF212 and HEI10 detected by yeast two-

hybrid assays in (I).

Figure 5. Chromosomal localization of RNF212B and RNF212 in mutants defective for

recombination and synapsis

(A) RNF212B and RNF212 localize to synapsed regions between non-homologous axes in

DSB-defective $Spo11^{-/-}$ spermatocytes. Images show a late zygotene/early pachytene-like

nucleus immunostained for SYCP3, SYCP1, RNF212B and RNF212.

(B and C) Staining intensities of RNF212B and RNF212 on synapsed regions. Signal intensities

per μm of SC per nucleus (B); and ratios of intensities per μm of SC for RNF212B relative to

RNF212 (C). Black and red bars indicate means. 14 early pachytene $Spo11^{+/+}$ nuclei and 10 late

zygotene/early pachytene-like $Spo11^{-/-}$ nuclei were analyzed.

(D) RNF212B and RNF212 localize to recombination sites between aligned homolog axes in

synapsis-defective $Sycp1^{-/-}$ spermatocytes. A late zygotene/early pachytene-like nucleus

immunostained for SYCP3, RNF212B, RNF212 and RPA2 is shown.

(E and F) Focus counts of RNF212B and RNF212 (E); and degree of RPA2 colocalization with

RNF212B and RNF212 (F). Black bars indicate means. 7 early pachytene $Sycp1^{+/+}$ nuclei and 8

late zygotene/early pachytene-like $Sycp1^{-/-}$ nuclei were analyzed.

(G) Differentiation of crossover-specific foci of RNF212B and RNF212 and persistence of

RNF212 staining in crossover-defective *Mlh3*$^{-/-}$ spermatocyte nuclei. Late pachytene and early

diplotene nuclei immunostained for SYCP3, RNF212B, and RNF212 are shown. Arrowheads

indicate large RNF212B foci.

(H and I) Numbers of RNF212B (H) and RNF212 (I) foci per nucleus in successive late

prophase I stages of spermatocytes. Black bars indicate means. Numbers of nuclei analyzed in

early pachynema, late pachynema, and early diplonema were 12, 20, and 13 for *Mlh3*$^{+/+}$; and

12, 18, and 13 for *Mlh3*$^{-/-}$.

ns, not significant ($p > 0.05$); ***$p \leq 0.001$; ****$p \leq 0.0001$ for two-tailed Mann-Whitney tests.

Magnified images in (D) and (G) show representative chromosomes. Scale bars, 10 µm for full

nuclei and 2 µm for magnified images.

Figure 6. RNF212B is required for crossing over and localizes to the synaptonemal

complexes and recombination sites in prophase I oocytes.

(A) Ovary sections from wild-type and *Rnf212b*$^{-/-}$ mutant females at 18 days postpartum (dpp)

immunostained for p63 and counterstained with hematoxylin.

(B) Oocyte counts at 18 dpp. Red bars indicate means. ns, not significant ($p > 0.05$, two-tailed *t*

test). n, numbers of mice analyzed.

(C) MLH1 foci are absent in *Rnf212b*$^{-/-}$ oocytes. Pachytene-stage oocyte nuclei from wild-type

and *Rnf212b*$^{-/-}$ ovaries at embryonic day 18.5 immunostained for SYCP3 and MLH1. Scale bars,

10 μm.

(D) Chiasmata are diminished in *Rnf212b*$^{-/-}$ oocytes. Metaphase-I oocytes from ovaries of ≥2-

month-old wild-type and *Rnf212b*$^{-/-}$ females stained with DAPI and immunostained for CREST

(centromeres). Scale bars, 10 μm.

(E) RNF212B localization at successive prophase-I stages of fetal oocyte nuclei from embryonic

days 15.5 (E15.5) and 17.5 (E17.5), and 0.5 day postpartum (0.5 dpp). Surface-spread

prophase-I oocyte nuclei were immunostained for SYCP3 and RNF212B. Arrowheads indicate

large, differentiated RNF212B foci. The magnified images show representative chromosomal

regions. Scale bars, 10 μm for images of full nuclei and 2 μm for magnified panels.

(F) Numbers of RNF212B foci in oocyte nuclei at successive prophase-I stages. Red bars

indicate means. Means of focus numbers are indicated below the graph. Numbers of nuclei

analyzed in leptonema (at E15.5), zygonema (at E15.5), early pachynema (pachytene nuclei

with ≤4 MLH1 foci at E16.5), mid/late pachynema (pachytene nuclei with ≥20 MLH1 foci at

E18.5), early diplonema (at 0.5 dpp), mid diplonema (at 0.5 dpp), late diplonema (at 0.5 dpp)

and dictyate stage (at 0.5 dpp) are 21, 22, 11, 22, 18, 11, 15 and 7, respectively.

(G) RNF212B colocalization with RNF212 and MLH1 in oocytes. Early pachytene (top) and

mid/late pachytene (bottom) oocytes from E18.5 ovaries immunostained for SYCP3, RNF212B,

RNF212 and MLH1. Arrowheads indicate crossover sites. The magnified images show

representative chromosomal regions. Scale bars, 10 μm for images of full nuclei and 2 μm for

magnified panels.

(H) Quantification of focus intensities for RNF212B and RNF212. Black bars indicate means. **p

≤ 0.01; ***p ≤ 0.001; ****p ≤ 0.0001 for two-tailed Mann-Whitney tests. 7 early/mid pachytene

(with ≤10 MLH1 foci) and 9 mid/late pachytene (with ≥20 MLH1 foci) nuclei from E18.5 ovaries

were analyzed. Total numbers of foci analyzed: 28 crossover RNF212B foci, 1,375 other

RNF212B foci, 30 crossover RNF212 foci and 1,596 other RNF212 foci in early/mid

pachynema; 250 crossover RNF212B foci, 1,468 other RNF212B foci, 245 crossover RNF212

foci and 1,580 other RNF212 foci. All, all foci; COs, crossover foci colocalized with MLH1 foci;

Others, other foci that don't colocalize with MLH1 foci.

Figure 7. Dynamic chromosomal localization of HEI10 in oocytes.

(A-C) HEI10 localization at successive prophase-I stages of fetal oocyte nuclei from embryonic

days 16.5 (E16.5) and 17.5 (E17.5). Surface-spread prophase-I oocyte nuclei in early

pachynema (A), early/mid pachynema (B), and mid pachynema (C) were immunostained for

SYCP3, HEI10 and MLH1. Arrowheads indicate crossover foci marked by MLH1. The asterisk

indicates an SC with an isolated bright HEI10 focus but without an MLH focus. The magnified

images show representative chromosomal regions. Scale bars, 10 µm for images of full nuclei

and 2 µm for magnified panels.

(D) Numbers of HEI10 and MLH1 foci in oocyte nuclei at successive prophase-I stages. Red

bars indicate means. Means of focus numbers are indicated below the graph. Numbers of nuclei

analyzed in leptonema (L; at E16.5), early and mid zygonema (E/MZ; at E16.5), late zygonema

(LZ; at E16.6), early pachynema (pachytene nuclei with ≤4 MLH1 foci at E16.5), early-to-mid

pachynema (E/MP; pachytene nuclei with 5-15 MLH1 foci at E16.5), and mid pachynema (MP;

pachytene nuclei with ≥20 MLH1 foci at E16.5 and E17.5), are 8, 8, 24, 18, 11 and 21,

respectively.

(E) Quantification of HEI10 and MLH1 foci in pachytene oocytes from E16.5 and E17.5 ovaries.

56 pachytene nuclei were randomly selected and analyzed.

(F) Quantification of HEI10 focus intensities. Red bars indicate means. ***$p \leq 0.001$; ****$p \leq$

0.0001 for two-tailed Mann-Whitney tests. 26 early pachytene (with ≤4 MLH1 foci), 9 early/mid

pachytene (with 5-15 MLH1 foci) and 6 mid pachytene (with ≥20 MLH1 foci) nuclei from E16.5

ovaries were analyzed. Total numbers of foci analyzed: 33 crossover foci and 3,817 other foci in

early pachynema; 96 crossover foci and 1,089 other foci in early/mid pachynema; 186 crossover

foci and 180 other foci. All, all foci; COs, crossover foci colocalized with MLH1 foci; Others,

other foci that don't colocalize with MLH1 foci.

(G) Per chromosome analysis for HEI10 focus intensity. Intensity of each crossover-associated

HEI10 focus relative to that of the brightest other (non-crossover) focus along the same

chromosome. Red bars indicate means. Mean intensities are indicated below the graph.

Detailed representation of per chromosome analysis is shown in **Figure S13B**. ****$p \leq 0.0001$

for two-tailed Mann-Whitney tests. 31, 92, and 111 crossover foci from 14 early pachytene, 9

early/mid pachytene, and 6 mid pachytene nuclei were analyzed, respectively.

(H) Summary of localization dynamics of RNF212B, RNF212, and HEI10 at designated

crossover sites and noncrossover sites in spermatocytes and oocytes. See text for details.

(I) Models of crossover maturation and defects in *Rnf212b, Rnf212* and *Hei10* mutants. MutSγ

complexes (grey rings) bind nascent Holliday junctions and convert into sliding clamps that

embrace the interacting duplexes [28]. At designated crossover sites, stabilization of MutSγ (and

other ZMMs) by RNF212 and RNF212B stabilize single-end invasions. HEI10 and an

endonuclease-independent function of MutLγ[58] then facilitate second-end capture to form dHJs.

MutLγ-catalyzed DNA incision and the Bloom complex (BLM-TOPIIIα-RMI1-RMI2; BTRR) then

mediate crossover-specific dHJ resolution [29]. MutSγ is destabilized in *Rnf212b*$^{-/-}$,

Rnf212b$^{RING/RING}$ and *Rnf212*$^{-/-}$ mutants exposing nascent intermediates to unwinding by the

Bloom complex resulting in noncrossovers. In the *Hei10*$^{mei4/mei4}$ mutant, RNF212, RNF212B, and

MutSγ persist at all sites throughout pachytene, and crossover sites fail to differentiate. DSBs

are repaired as noncrossovers only as homologs desynapse and RNF212 and RNF212B

dissociate.

SUPPLEMENTAL FIGURES

Figure S1. 5' UTR variants and isoforms of mouse *Rnf212b*, related to Figure 1.

(A) Protein sequence alignments of mouse and human RNF212B and RNF212 full-length

isoforms. Zinc-coordinating cysteine and histidine (pink) residues of RING finger domain and

other identical residues (blue) are highlighted.

(B) Schematic of 5' UTR variants and isoforms of mouse *Rnf212b*. Top, red and black bars

represent 5' untranslated exons (5' UTRs) and coding exons, respectively. Two 5' UTR variants

were identified in the NCBI database, *variants a* and *b*, which differ by a deletion of the second

untranslated exon in *variant b*. A third *variant c* was identified by 5' RACE from testes and

comprises ~1/3 of the third untranslated exon of *variant a* followed by the 14 coding exons.

Bottom, yellow and blue boxes represent 5' UTRs and coding exons, respectively. Four different

coding isoforms were identified in the NCBI database. For simplicity, 5' UTRs of isoforms a-c are

not shown. Isoform a encodes the full-length RNF212B protein of 297 amino acids; isoform b

uses an alternative splice acceptor site, creating a premature stop codon after exon 10 that is

predicted to produce a protein lacking 86 C-terminal amino acids; isoform c uses an alternative

splice donor site to precisely skip exon 10, which corresponds to 16 amino acids of the

disordered serine-rich C-terminal tail, resulting in a predicted protein of 270 amino acids; and

isoform d lacks the first two exons present in other isoforms and uses an extended exon 3 and

most of exon 4 as a 5' UTR, to encode a predicted protein of 189 amino acids that lacks the N-

terminal RING finger domain. Asterisks indicate positions of stop codons. Horizontal bars

represent positions of primers used to detect each isoform by RT-PCR in (C). Annotated or

related variants and isoforms in the NCBI database are also shown. Expression of full-length a

and b isoforms in testis was confirmed by *de novo* cloning from testis mRNA (the sequence of

cloned isoform b was similar but not identical to XR_874633).

(C) Expression of mouse *Rnf212b* 5' UTR variants and isoforms analyzed by RT-PCR. Total

RNA was extracted from indicated tissues of mature male mice and subjected to RT-PCR.

Gapdh is a loading control.

Figure S2. Domain structure of RNF212B and strategy to generate *Rnf212b* mutant mice

by CRISPR/Cas9, related to Figure 1.

(A) Domain structures of mouse RNF212B and RNF212. Amino acid sequences of the RING

finger domains are shown below. Presumptive zinc-coordinating cysteines and histidines and

other identical residues are highlighted in pink and blue, respectively. The domain structure of

the full-length RNF212B protein mirrors that of RNF212 with an N-terminal RING finger domain

(36.7% amino-acid identity with RNF212), a ~70 amino acid region of predicted coiled-coil

(25.7% identical) and a largely disordered serine-rich C-terminal tail (22.8% identical).

(B) Schematic of the mouse *Rnf212b* gene on chromosome 14. Top row: red and black bars

represent 5' untranslated exons (5' UTRs) and coding exons, respectively. Middle row: yellow

and blue boxes represent 5' UTRs and coding exons, respectively, in the full-length isoform a.

The inverted red triangle represents the translation initiation site in testis identified by 5' RACE.

An asterisk represents the stop codon. Horizontal bars represent the positions of primers used

to detect 5' UTR variants a/b/c by RT-PCR in **Figure S1C**. Bottom row: nucleotide and amino-

acid sequences at the CRISPR/Cas9 targeted site in exon 1. A single-nucleotide insertion

(shown in red) created a frameshift and premature stop codon (asterisk). The guide RNA

(gRNA) sequence is highlighted in blue.

Figure S3. Absence of crossover markers in *Rnf212b*^{-/-} mutant mice, related to Figure 1.

Crossover-specific immunostaining foci of PRR19 (left) and CDK2 (right) foci were absent from

chromosome spreads of *Rnf212b*^{-/-} spermatocytes. Mid/late pachytene nuclei from wild-type and

Rnf212b^{-/-} spermatocytes were immunostained for SYCP3 plus PRR19 (left) or CDK2 (right).

The magnified images show representative chromosomes from the CDK2 staining. Note that

CDK2 also localizes to telomeres, which is not affected by *Rnf212b* mutation. An arrowhead

indicates an interstitial CDK2 focus marking a prospective crossover site. Scale bars, 10 μm for

images of full nuclei and 2 μm for magnified panels.

Figure S4. Dynamics of RAD51 and DMC1 throughout zygonema and pachynema, related

to Figure 1.

(A and B) Representative images of RAD51 (A) and DMC1 (B) immunostainings in wild-type

and *Rnf212b*[-/-] spermatocytes nuclei immunostained for SYCP3 plus RAD51 (A) or DMC1 (B) at

the indicated stages. Arrows indicate the X-Y chromosomes. Scale bars, 10 μm.

(C and D) Focus counts of RAD51 (C) and DMC1 (D). Red bars indicate means ± SDs.

ns, not significant ($p > 0.05$); *$p \leq 0.05$; ***$p \leq 0.001$ for two-tailed Mann-Whitney tests. Total

numbers of nuclei analyzed are indicated below the X axes.

Figure S5. Intermediate steps of recombination are perturbed in *Rnf212b*^{-/-} mutant mice,

related to Figure 1.

(A, C and E) Images of wild-type and *Rnf212b*^{-/-} spermatocyte nuclei immunostained for SYCP3

plus RPA2 (A), TEX11 (C) or MER3 (E) at the indicated stages. Scale bars, 10 µm.

(B, D and F) Counts of RPA2 (B), TEX11 (D) and MER3 (F) immunostaining foci. Red bars

indicate means ± SDs. ns, not significant ($p > 0.05$); *$p \leq 0.05$; ***$p \leq 0.001$; ****$p \leq 0.0001$ for

two-tailed Mann-Whitney tests. Total numbers of nuclei analyzed are indicated below the X

axes.

Ito et al., Figure S6 (related to Figure 2)

976 **Figure S6. Specificity of anti-RNF212B antibody, related to Figure 2.**

977 Wild-type and *Rnf212b*^{-/-} spermatocyte nuclei at indicated stages were immunostained for

978 SYCP3 and RNF212B. Scale bars, 10 µm.

Figure S7. Distinct localization dynamics of RNF212B and RNF212 foci, related to Figure

3.

(A) Quantification of focus intensities for RNF212B and RNF212. Each dot represents each

focus of RNF212B and RNF212 from the early, mid and late pachytene nuclei (one nucleus per

stage) analyzed in **Figure 3D**. Bars indicate means ± SDs. All, all foci; COs, crossover-specific

foci that colocalize with HEI10 foci; Others, foci that don't colocalize with HEI10 foci

(noncrossover foci). ND, not detected. *$p \leq 0.05$; **$p \leq 0.01$; ****$p \leq 0.0001$ for two-tailed Mann-

Whitney tests.

(B) Representation of per chromosome analysis of RNF212B and RNF212 foci for a mid-

pachytene chromosome (shown in the images in the right-hand panels). Each focus of

RNF212B or RNF212 is represented by a dot. The dashed line indicates the signal intensity of

the brightest other focus (noncrossover focus) along the same chromosome. Orange vertical

lines indicate the positions of HEI10-positive crossover sites (a) and (b). Intensities of foci at

these sites relative to that of the brightest other (noncrossover) focus are: (a) 3.5 and (b) 2.3 for

RNF212B; and (a) 0.8 and (b) 1.1 for RNF212.

(C) Total signal intensity of RNF212B and RNF212 per nucleus in early, mid and late pachytene

spermatocytes. Black bars indicate means. ns, not significant ($p > 0.05$); *$p \leq 0.05$; ***$p \leq 0.001$

for two-tailed Mann-Whitney tests. Nuclei analyzed were the same as in **Figure 3D**.

Figure S8. Timing of RNF212B focus amplification and appearance of HEI10 foci in

spermatocytes, related to Figure 3.

(A) Amplification of RNF212B foci relative to the emergence of HEI10 foci. Airyscan images of

an early-to-mid pachytene spermatocyte nucleus immunostained for SYCP3, RNF212B,

RNF212 and HEI10. This nucleus contains 14 HEI10 foci indicating that it is in transition to a full

complement of designated crossovers sites (≥20 HEI10 foci). The white arrowhead indicates an

amplified RNF212B focus without an associated HEI10 focus; and the orange arrowhead

indicates an amplified RNF212B focus colocalized with a small, emerging HEI10 focus. These

configurations indicate that growth of RNF212B foci can precede a detectable HEI10 focus.

(B) Amplification of RNF212B at crossover sites. Crossover-specific foci of RNF212B (i.e.

RNF212B foci that colocalized with HEI10 foci) in early-to-mid pachytene nuclei (containing 5-18

HEI10 foci), and mid-pachytene nuclei (≥19 HEI10 foci) were classified based on their degree of

amplification by measuring their intensity relative to the brightest other (noncrossover)

RNF212B focus along the same chromosome (complete means no other foci detected along the

SC; strong, ≥2-fold brighter; weak, 1.5-2-fold brighter; or no amplification, <1.5-fold brighter). 81

crossover-specific RNF212B foci from 7 early-to-mid pachytene and 186 crossover-specific

RNF212B foci from 8 mid-pachytene nuclei were analyzed by Airyscan imaging. In early-mid

pachytene nuclei, the majority of HEI10 foci are associated with detectable amplification of

RNF212B but 28% are not.

(C) Amplification of RNF212B and emergence of HEI10 foci in early-to-mid pachynema.

Numbers of HEI10 foci (sky blue), and RNF212B foci with weak (yellow) and strong

amplification (orange) are shown. Criteria for RNF212B focus amplification are as in (B). 71.6%

(58/81 foci) of sites marked by HEI10 were coincident with weak or strong amplification of

RNF212B, while 78.4% (40/51 foci) of strongly amplified RNF212B foci colocalized with a HEI10

focus. Thus, with respect to the amplification of RNF212B and the emergence of HEI10 at

prospective crossover sites, a sequential order of events is hard to discern, i.e. these events

may be coincident, consistent with the dependence of RNF212B focus differentiation on HEI10.

7 early-to-mid pachytene nuclei with 5-18 HEI10 foci per nucleus were analyzed by Airyscan

imaging.

1028

Ito et al., Figure S9 (related to Figure 4)

1029

Figure S9. Interdependent chromosomal localization of RNF212B and RNF212 in oocytes,

related to Figure 4.

(A and B) RNF212 (A) and RNF212B (B) are interdependent for chromosomal localization.

Pachytene-stage oocytes at embryonic day 18.5 or 16.5 (E18.5 or E16.5) from the indicated

genotypes were immunostained for SYCP3 and RNF212 (A) or RNF212B (B). Scale bars, 10

μm.

Figure S10. The RING finger domain is required for RNF212B function *in vivo*, related to

Figure 4.

(A) Scheme of the *Rnf212b* RING mutation. A three-nucleotides deletion in addition to a single-

nucleotide substitution (shown in red) induced loss of the 9th cysteine residue with no change in

other protein-coding amino acids. White and black boxes indicate 5' UTR and coding exons,

respectively. The guide RNA (gRNA) sequence is highlighted in blue. Cysteine and histidine

residues of RING finger domain are highlighted in pink.

(B) Reduced testis size in *Rnf212b*$^{RING/RING}$ mutants. Red bars indicate means. ns, not significant

($p > 0.05$); ****$p \leq 0.0001$ for two-tailed *t* tests. Numbers of mice analyzed are indicated below

the X axis.

(C) Diminished chiasma numbers in *Rnf212b*$^{RING/RING}$ meiocytes. Left, wild-type and

Rnf212b$^{RING/RING}$ spermatocytes in diakinesis/metaphase-I stages stained with DAPI. Right,

metaphase-I oocytes from ≥2 months old wild-type and *Rnf212b*$^{RING/RING}$ females were stained

with DAPI and immunostained for centromeres. Scale bars, 10 μm.

(D) Chiasma counts in *Rnf212b*$^{-/-}$ and *Rnf212b*$^{RING/RING}$ meiocytes. Red bars indicate means. ns,

not significant ($p > 0.05$) for two-tailed Mann-Whitney tests. Total numbers of nuclei analyzed

are indicated below the X axis.

(E) Absence of MLH1 foci in *Rnf212b^RING/RING^* spermatocytes. Mid-late pachytene nuclei from

wild-type and *Rnf212b^RING/RING^* spermatocytes immunostained for SYCP3 and MLH1. Scale

bars, 10 µm.

(F) High frequency of unpaired X-Y chromosomes in H1t-positive pachytene *Rnf212b^RING/RING^*

cells. ns, not significant ($p > 0.05$); ****$p ≤ 0.0001$ for Fisher's exact tests. Total numbers of cells

analyzed from 3 mice of each genotype are indicated below the X axis.

(G-J) Focus counts of RPA2 (G), MSH4 (H), TEX11 (I), and MER3 (J) in wild-type, *Rnf212b^-/-^*,

and *Rnf212b^RING/RING^* spermatocytes. Red bars indicate means ± SDs. ns, not significant ($p >$

0.05); *$p ≤ 0.05$; **$p ≤ 0.01$; ***$p ≤ 0.001$; ****$p ≤ 0.0001$ for two-tailed Mann-Whitney tests. Total

numbers of nuclei analyzed are indicated below the X axes.

1064

Figure S11. Role of the RNF212B RING-finger domain for self-interaction, chromosomal

localization, and protein stability, related to Figure 4.

(A) Yeast two-hybrid assay shows that a C9A RING-domain mutation in RNF212B diminishes its

self-interaction but not its interaction with RNF212.

(B) Diminished protein levels of RNF212B and RNF212 in $Rnf212b^{RING/RING}$ spermatocytes.

Whole-testis extract from indicated genotypes were subjected to immunoblotting for RNF212

and IP-immunoblotting for RNF212B. SYCP3 is a loading control of meiotic cells.

(C and D) Chromosomal localization of RNF212B (C) and RNF212 (D) is diminished in

Rnf212b$^{RING/RING}$ mutant meiocytes. Early pachytene spermatocyte nuclei from wild-type and

Rnf212b$^{RING/RING}$ testes (left); and pachytene-stage nuclei from wild-type and *Rnf212b*$^{RING/RING}$

oocytes at E16.5 (right) were immunostained for SYCP3 and RNF212B (C) or RNF212 (D).

Scale bars, 10 μm.

Figure S12. Chromosomal localization of RNF212B and RNF212 in mutants defective for

synapsis and crossing over, related to Figure 5.

(A) A mid zygotene-like $Sycp1^{-/-}$ spermatocyte nucleus immunostained for SYCP3, RNF212B,

RNF212, and RPA2. General staining of RNF212B and RNF212 is absent but limited

localization to small regions where homologous axes are closely aligned can be detected

(magnified in the panels on the right).

(B and C) Quantification of RPA2 foci colocalizing with RNF212B and RNF212. (B) Degree of

RNF212B and RNF212 colocalization with RPA2. (C) Quantification of total and colocalizing

RPA2 foci. Black bars indicate means. 7 early-pachytene $Sycp1^{+/+}$ nuclei and 8 late-

zygotene/early-pachytene-like $Sycp1^{-/-}$ nuclei were analyzed.

(D) Quantification of RNF212B-RNF212 colocalization. Black bars indicate means. 7 early

pachytene $Sycp1^{+/+}$ nuclei and 8 late zygotene/early pachytene-like $Sycp1^{-/-}$ nuclei were

analyzed.

(E) Normal chromosomal localization of RNF212B and RNF212 in early pachytene $Mlh3^{-/-}$

mutant spermatocyte nuclei. An early pachytene nucleus immunostained for SYCP3, RNF212B

and RNF212 is shown.

1094 (F) Numbers of RNF212B and RNF212 foci in wild-type and *Mlh3*$^{-/-}$ late-pachytene

1095 spermatocyte nuclei. Left, numbers of large and small RNF212B foci. Right, numbers of

1096 RNF212 foci associated with crossover sites (colocalizing with large RNF212B foci) and other

1097 sites. Black bars indicate means. 20 *Mlh3*$^{+/+}$ and 18 *Mlh3*$^{-/-}$ nuclei were analyzed.

1098 ns, not significant ($p > 0.05$); **$p \le 0.01$; ****p \le 0.001$; *****p \le 0.0001$ for two-tailed Mann-

1099 Whitney tests. Scale bars, 10 µm in images of full nuclei (A and E) and 2 µm in the magnified

1100 panels (A).

Figure S13. Localization of HEI10 and MLH1 in oocytes, related to Figures 6 and 7.

(A) Total signal intensity of RNF212B and RNF212 per nucleus in early/mid and mid/late

pachytene oocytes. Black bars indicate means. ns, not significant ($p > 0.05$) for two-tailed

Mann-Whitney tests. Nuclei analyzed were the same as in **Figure 6H**.

(B) HEI10 localization at successive prophase-I stages of fetal oocyte nuclei from embryonic

day 16.5 (E16.5). Surface-spread prophase-I oocyte nuclei in mid zygonema, early pachynema,

early/mid pachynema, and mid pachynema were immunostained for SYCP3, HEI10, and MLH1.

Arrowheads indicate crossover sites marked by MLH1. The magnified images show

representative chromosomal regions. Scale bars, 10 µm for images of full nuclei and 2 µm for

magnified panels.

(C) Representations of per chromosome analysis of HEI10 foci for the chromosomes highlighted

in (B). Each HEI10 focus is represented by a dot. Magenta dots indicate HEI10 foci at crossover

sites (colocalizing with MLH1 foci). The dashed line indicates the signal intensity of the brightest

other focus (noncrossover focus) along the same chromosome.

(D) Per chromosome analysis showing that HEI10 foci located at crossover sites (colocalizing

with MLH1) increase in relative intensity as pachytene progresses. Mean intensities are

indicated by red lines and values shown below the graph. ns, not significant ($p > 0.05$); ***$p \leq$

0.001 for two-tailed Mann-Whitney tests. 31, 91, and 105 crossover foci from 14 early

pachytene, 9 early/mid pachytene, and 6 mid pachytene nuclei were analyzed, respectively.

1121

MATERIALS AND METHODS

Mice

Mice were maintained and euthanized according to the guidelines of the Institutional Animal

Care and Use Committees of the University of California Davis. All mice were congenic with the

C57BL/6J background except for *Sycp1*[-/-] mutant mice. The *Rnf212*[-/-], *Hei10*[mei4/mei4], *Spo11*[-/-],

Sycp1[-/-] and *Mlh3*[-/-] mutant lines were previously described [15, 25, 49, 51, 52]. Mature adult mice (2-6

months old) were used for experiments unless otherwise noted.

1129

Rnf212b mutant alleles were generated by The Cornell Stem Cell and Transgenic Core Facility.

124 bp of DNA template for sgRNA, comprising the T7 polymerase binding site and sgRNA

target sequence (5'-CCCCATCTTTTCGGAAACAC-3') followed by the remaining sgRNA

sequence, were amplified by PCR as previously described [63] and purified using a GeneJET

PCR Purification Kit (Thermo Scientific, K0702). 600 ng of purified DNA template was *in vitro*

transcribed for 4 h at 37^{o}C using MEGAshortscript T7 Transcription Kit (Invitrogen, AM1354).

sgRNA was purified using MEGAclear Transcription Clean-Up Kit (Invitrogen, AM1908), diluted

to 1.6 µg/µl and stored at -80^{o}C prior to embryo injection. 12.5 ng/µl of purified sgRNA and

either 12.5 or 25 ng/µl of Cas9 mRNA were microinjected into pronuclei of 2-cell stage of

C57BL/6J homozygous embryos. Embryos were transferred to pseudo-pregnant female

recipients 2 days after pronuclear injection at the 4-cell stage. Genomic DNA of founder mice

was isolated from toe clips, PCR-amplified, cloned into the pCR4Blunt-TOPO vector (Invitrogen,

450031), and Sanger-sequenced to identify mutations. Founder mice were backcrossed to

C57BL/6J mice for ≥4 generations and heterozygous mice were bred to homozygosity.

Genotyping was performed by PCR on genomic DNA isolated from mouse tails. Primers used

for cloning and genotyping are listed in **Table S2**.

1146

Total RNA extraction, RT-PCR, and 5' RACE

Tissues were dissected from adult male mice, washed in phosphate-buffered saline (PBS),

frozen in liquid nitrogen and stored at -80^{o}C prior to RNA extraction. Total RNA was extracted

using TRIzol according to manufacturer's instructions (Invitrogen). 1 µg of total RNA was

reverse-transcribed using the SuperScript IV First-Strand Synthesis System (Invitrogen,

18091050) and synthesized cDNA was PCR-amplified. For 5' RACE, 4 µg of total RNA from

testes was reverse-transcribed using 5' RACE System for Rapid Amplification of cDNA Ends

(Invitrogen, 18374058) and synthesized cDNA was PCR-amplified, cloned into the pCR4-TOPO

TA vector (Invitrogen, K457501) and Sanger-sequenced to identify translation initiation site.

Primers used are listed in **Table S2**.

1157

Yeast two-hybrid assays

Full-length mouse *Rnf212b* and *Rnf212* cDNAs were PCR amplified from mouse testis cDNA,

prepared as above, and cloned into pGADT7 and pGBKT7 vectors (Clontech) using a Gibson

Assembly Cloning Kit (NEB, E2611S). Full-length mouse *Hei10* coding sequence was

synthesized by Twist Bioscience and cloned into pGADT7 and pGBKT7. Prey and bait vectors

were co-transformed into the Y2HGold strain. Transformants were suspended in 20 µl sterile

water, and 2 µl spotted onto fresh plates with and without selection: SD/-Trp/-Leu plate, SD/-Trp/-

Leu containing 200 ng/ml of Aureobasidin A (Clontech, 630466) to select for the *AUR1-C* reporter

gene, or SD/-Trp/-Leu /-Ade/-His to select for *ADE2* and *HIS3* reporters. After incubation of the

plates at 30°C for 3 days, single colonies were inoculated into SD/-Trp/-Leu media and grown

overnight. Colonies were resuspended at OD_{600} = 1, four 10-fold serial dilutions were prepared,

and 10 μl drops were spotted onto selection plates and incubated at 30°C for 3 to 5 days.

Primers used for cloning and point mutagenesis are listed in **Table S2**.

1171

Antibody production

Polyclonal antibodies against mouse RNF212B were raised in Guinea pigs. Codon-optimized of

full-length mouse RNF212B was cloned into pET-28b (+) (Addgene) with a C-terminal 6xHis tag.

E. coli Arctic Express (DE3) cells (Agilent Technologies Inc.) transformed with the mouse

RNF212B-6xHis expression vector were grown in 2L of LB at 30°C to an OD_{600} of 0.8, and

protein expression was induced with 0.5 mM IPTG followed by incubation at 11°C in media

supplemented with 0.1 mM $ZnCl_2$. ~5 g of cells were pelleted by centrifugation, suspended in 60

ml of denaturing lysis buffer A (6M Guanidine, 25 mM sodium-phosphate pH 7.4, 500 mM NaCl,

0.1 mM $ZnCl_2$, 1 mM β-mercaptoethanol, 20 mM imidazole, 10% glycerol) and sonicated.

Samples were centrifuged at 35,000 rpm in a Ti-45 rotor (Beckman) for 45 min and soluble

extract was applied to a 5 ml HisTrap FF column (GE healthcare Inc.) using a GE AKTA Avant

25 FPLC system. After equilibrating with 5 column volumes (CV) of lysis buffer A, followed by

washes with 10 CV of wash buffer B (6M urea, 25 mM sodium-phosphate pH 7.4, 500 mM

NaCl, 1 mM β-mercaptoethanol, 10% glycerol) supplemented with 60 mM imidazole, bound

proteins were eluted with a linear gradient of imidazole (60-600 mM) in buffer B. Peak fractions

were pooled and dialyzed extensively in decreasing urea concentrations (6M to 1M urea in 25

mM sodium-phosphate pH 7.4, 500 mM NaCl, 1 mM β-mercaptoethanol, 10% glycerol). Protein

concentrations were determined by Bradford and spectrophotometric (A_{280}) methods. Two

Guinea pigs were immunized with purified RNF212B protein by Antibodies Incorporated (Davis,

CA). The IgG fraction was purified from the resultant sera, using the Montage antibody

purification kit according to manufacturer's instructions (Millipore Sigma), and dialyzed against

PBS containing 10% glycerol and aliquots stored at -80°C.

1194

Histology

Testes from adult male mice were dissected, punctured with a 27-guage needle, and fixed in

10% buffered formalin overnight at room temperature. Ovaries from 18 days postpartum (18

dpp) female mice were dissected and fixed in 10% buffered formalin overnight at room

temperature. Fixed testes and ovaries were washed in PBS twice and stored in 70% ethanol at

4^{o}C prior to embedding. Tissues embedded in paraffin were sectioned (5 μm) onto glass slides

(Fisher Scientific, 12-550-15), deparaffinized, rehydrated, and incubated with antigen retrieval

buffer (10 mM Sodium Citrate, 0.05% Tween-20) for 50 min at 100^{o}C. Slides were then stained

with hematoxylin and eosin (testes), or immunostained with anti-p63 antibody, as described

below, and counterstained with hematoxylin (ovaries), and mounted with Permount (Fisher

Scientific, SP15-100).

1206

Surface spreads of spermatocyte chromosomes

Surface-spread chromosomes of spermatocytes were prepared as described previously [64]with

slight modification. Testes were dissected, the tunica albuginea removed, and adherent

extratubular tissues removed by rinsing and dissociating seminiferous tubules in PBS using a

pair of 25-gauge (adult) or 27-guage (juvenile) needles in a 35 mm petri dish (Corning, 430165).

Dissociated seminiferous tubules were incubated in hypotonic extraction buffer (30 mM Tris-HCl

pH 8.0, 50 mM sucrose, 17 mM trisodium citrate dihydrate, 5 mM EDTA, 0.5 mM dithiothreitol

(DTT) and 0.5 mM phenylmethylsulphonyl fluoride (PMSF), pH 8.2-8.3) for 15-45 min at room

temperature. After quickly rinsing seminiferous tubules in 100 mM sucrose, a small amount of

tubules were placed in 40 µl of 100 mM sucrose on a glass depression slide. Tubules were torn

to pieces using two fine forceps, and large pieces of tubular remnant were removed. The

volume was increased to 40-60 µl by adding 100 mM sucrose and a cell suspension was made

by pipetting. A clean glass slide (Fisher Scientific, 12-544-7) was dipped into freshly made PFA

solution (1% paraformaldehyde adjusted to pH 9.2 using 1.25 M sodium borate, 0.15% Triton X-

100) in a 50 ml Falcon tube, and excess solution was drained onto a paper towel. 20 µl of cell

suspension was placed at the upper right corner of the slide and slowly dispersed in horizontal

and vertical directions to homogeneously cover the slide. Hot tap water was added to a humid

chamber and slides were slowly dried in the closed chamber overnight at room temperature.

Slides were further dried for 3 hr with lid ajar, and then for 1-2 hr with the lid removed. Slides

were washed once for 5 min in deionized water and twice for 5 min in 0.4% Photo-Flo 200

solution (Kodak, 1464510) in a coplin jar before air-drying at room temperature. Slides were

either directly processed for immunostaining, as described below, or stored wrapped in

aluminum foil at -80°C prior to immunostaining. For SIM imaging, 15 µl of cell suspension was

placed at the upper right corner of a coverslip (Fisher Scientific, 12-544-E) coated with 90 µl of

1% PFA, 0.15% Triton X-100 solution and slowly dispersed as above.

1232

Surface spreads of oocyte chromosomes

Surface-spread oocyte chromosomes were prepared as described for spermatocytes with

modifications. Ovaries from fetal females were dissected into PBS and kept on ice prior to

incubation in hypotonic extraction buffer. A pair of ovaries were incubated in hypotonic extraction

buffer for 10 min at room temperature, quickly rinsed in 100 mM sucrose, and placed in 50 µl of

100 mM sucrose on a glass depression slide. Ovaries were torn to pieces using a pair of 25-

guage needles, and large pieces of ovarian remnant were removed. The volume was increased

to 40 µl by adding 100 mM sucrose and a cell suspension was made by pipetting. One half of a

clean glass slide was coated with 50 µl of freshly made 1% PFA 0.15% Triton X-100, and 10 µl

of cells suspension was placed at the upper right corner of the slide slowly dispersed in

horizontal and vertical directions to homogeneously cover the half slide. Slides were then dried,

washed, and stored as described for spermatocyte chromosomes spreads.

1245

Chromosome spreads of diakinesis/metaphase-I spermatocytes

Testes from adult male mice were dissected, tunica albuginea removed, and seminiferous

tubules placed in 2 ml of hypotonic solution (1% trisodium citrate) in a 35 mm petri dish.

Seminiferous tubules were torn to pieces using two fine forceps and, after adding 1 ml of

hypotonic solution, 3 ml of tubule suspension was transferred to a 15 ml Falcon tube using a

plastic transfer pipette (Phenix, PP-137030). Tubule fragments were allowed to settle out for 3

min and the supernatant containing suspended cells was transferred to another 15 ml Falcon

tube. Remnant tubule fragments in the petri dish were suspended in 2 ml of hypotonic solution

and transferred to the first 15 ml Falcon tube containing tubule fragments. The mixture of tubule

fragments was resuspended using a transfer pipette, tubule fragments were settled out, and the

supernatant was transferred to the 15 ml Falcon tube containing the first supernatant. The

mixture of supernatants was filtered through 70 μm and 40 μm Cell Strainers (Corning 352350;

Celltreat, 229481) and suspended cells were pelleted by centrifugation at 900 rpm for 10 min at

room temperature. Cells were fixed by adding 3 ml of freshly made fixative solution 1 (75%

methanol, 25% acetic acid with 0.375% chloroform) drop-by-drop while gentle vortexing. Cells

pelleted by centrifugation at 900 rpm for 10 min at room temperature, resuspended in 3 ml of

freshly made fixative solution 2 (75% methanol, 25% acetic acid, chilled to -20^{o}C), pelleted

again, and resuspended in 0.5 ml of fixative solution 2. Fixed cell suspension was dropped onto

a clean glass slide (Fisher Scientific, 12-544-7) from a height of 2-3 feet height using a glass

Pasteur pipette in a room with >40% humidity. Slides were air-dried for 10 min at room

temperature and either stained immediately with ProLong Gold or Diamond Antifade Mountant

(Thermo Fisher Scientific, P36930 or P36970) containing 1 µg/ml DAPI, or stored at 4^{o}C prior to

staining.

1269

Chromosome spreads of metaphase-I oocytes

Ovaries from adult female mice without prior hormonal stimulation were dissected, follicles were

punctured in pre-warmed (37^{o}C) M2 media (Sigma-Aldrich, M7167) using a 25-gauge needle,

and germinal-vesicle stage oocytes with integral cumulus cell layers were collected.

Surrounding cumulus cells were mechanically removed by pipetting and oocytes were cultured

in M2 media for 7 hr at 37^{o}C. Metaphase-I oocytes were transferred into Tyrode's acidic solution

(Sigma-Aldrich, T1788) to remove the zona pellucida (ZP), and ZP-free oocytes were

transferred and maintained in M2 media prior to spreading. 5-10 µl of 1% PFA solution pH 9.2

containing 0.15% Triton X-100 and 3 mM DTT was placed in each well of a 12-well glass slide

(Electron Microscopy Sciences, 63425-05) and one ZP-free oocyte was transferred to each well.

Slides were air-dried overnight at room temperature, and either directly processed for

immunostaining or stored at 4^{o}C for up to several days prior to immunostaining.

1282

Immunostaining

Slides were rehydrated with Tris-buffered saline (TBS) pH 8.0 containing 0.05% of Triton X-100

(TBST) for 3 min, blocked twice with blocking buffer (1% normal goat or donkey serum, 3%

bovine serum albumin (BSA), 1x TBS pH 8.0, 0.05% Triton X-100, 0.05% sodium azide) for 15

min at room temperature and incubated with primary antibodies in antibody dilution buffer (10%

normal goat or donkey serum, 3% BSA, 1x TBS pH 8.0, 0.05% Triton X-100, 0.05% sodium

azide) in a humid chamber overnight at room temperature. Slides were briefly rinsed with TBST,

washed twice with TBST for 5 min, blocked twice with blocking buffer for 15 min at room

temperature and then incubated with secondary antibodies in antibody dilution buffer in a humid

chamber for 1 hr at 37^{o}C. Slides were rinsed with TBST, washed three times with TBST for 5

min, washed once with Milli-Q water for 2 min, and air-dried prior to mounting with ProLong

Gold or Diamond Antifade Mountant. For SIM imaging, cover slips were blocked three times for

15 min prior to primary antibody incubation.

1296

For metaphase-I oocytes, slides were briefly rehydrated with TBST, washed three times with

TBST for 5 min, blocked twice with blocking buffer for 15 min at room temperature, and

incubated with primary antibodies in antibody dilution buffer in a humid chamber overnight at

room temperature. Slides were rinsed with TBST, briefly washed twice with TBST, blocked twice

with blocking buffer for 15 min at room temperature, and incubated with secondary antibodies in

antibody dilution buffer in a humid chamber for 1 hr at room temperature. Slides were rinsed

with TBST, briefly washed with TBST, and then washed three times with TBST for 5 min. Slides

were then incubated with TBST containing 5 µg/ml DAPI for 10 min at room temperature and

mounted with 50% ProLong Gold or Diamond Antifade Mountant in TBST.

All primary and secondary antibodies used are listed in **Table S3**.

1307

Protein blot analysis

Testes were dissected from 16-18 dpp juvenile male mice and frozen in liquid nitrogen. Five

pairs of testes were homogenized in 1 mL ice-cold RIPA buffer (50 mM Tris-HCl pH 7.5, 150 mM

NaCl, 1 mM EDTA, 1% NP-40, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate, SDS)

supplemented with protease and isopeptidase inhibitors (1x Complete protease inhibitor EDTA-

free, Roche, 04693159001), 1 mM PMSF, 10 mM N-ethylmaleimide, NEM) , incubated for 15

min on ice, sonicated (20% duty cycle, output 2 in burts for 2.5 min), and then incubated for 15

min on ice. Following centrifugation at 14,000 rpm for 15 min at 4^{o}C, supernatants were

collected as whole-testis extracts. Protein concentrations determined by Bradford assay were

normalized, and samples were subjected to electrophoresis and immunoblotting. Membranes

were blocked with TBST plus 2.5% non-fat milk for 1 hr at room temperature and incubated with

primary antibodies overnight at 4^{o}C. Membranes were washed three times with TBST for 10 min

each, then incubated with HRP-conjugated secondary antibodies in TBST for 1 hr at room

temperature. After four washes with TBST for 10 min each, HRP signal was developed using the

SuperSignal West Pico Chemiluminescent Substrate (Thermo Scientific, PI-34080) and

detected using an Amersham Imager 600 (GE Healthcare).

1324

1325   All primary and HRP-conjugated secondary antibodies used are listed in **Table S3**.

1326

Image acquisition

Images of surface-spread prophase chromosomes and metaphase-I chromosome spreads were

acquired using a Zeiss AxioPlan II microscope with a 63 x Plan-Apochromat 1.4 NA objective

and EXFO X-Cite metal halide light source, captured with a Hamamatsu ORCA-ER CCD

camera and processed using Volocity (Perkin Elmer) and Photoshop (Adobe) software. SIM

images were acquired using a Nikon N-SIM super-resolution microscope and processed using

NIS-Elements 2 image processing software. Airyscan images were acquired using a Zeiss

LSM800 with Airyscan microscope with a 60 x 1.4 NA objective and processed using ZEN

imaging software (Carl Zeiss). Images of testis sections were acquired using a Zeiss Axio

Imager M2 microscope with a 20 x Plan-Apochromat 0.8 NA objective, captured with a

Hamamatsu ORCA-Flash 4.0 V3 sCMOS camera and processed using ZEN imaging software

(Carl Zeiss). Images of ovary sections were acquired using a ScanScope digital scanner

(Asperio) and processed using ImageScope software.

1340

Image analysis

Comparisons were made between animals that were either littermates or matched by age.

Numbers and colocalization of foci were determined manually. *Rnf212b*$^{-/-}$, *Rnf212*$^{-/-}$ and

Rnf212b$^{RING/RING}$ mutant analysis was blinded with respect to the genotype of the animals.

Results from more ≥2 independent experiments/animals were pooled for quantification of

number of foci and chiasmata. *Spo11*$^{-/-}$, *Sycp1*$^{-/-}$, *Mlh3*$^{-/-}$ and *Hei10*$^{mei4/mei4}$ mutant analysis was

not blinded to genotype because phenotypes are overt. For focus intensity analyses, RNF212B

and RNF212 foci along homolog axes were automatically defined by thresholding intensity on

SYCP3 staining and manually inspecting images to separate any adjacent focus pairs defined

as one focus. For *Spo11*$^{-/-}$ mutant analysis, synapsed regions marked by SYCP1 were manually

cropped and intensities of RNF212B and RNF212 were measured. Two regions of interest (ROI)

were drawn inside nuclei but adjacent to homolog axes and their average intensity was

subtracted as background for each nucleus. Results from one experiment/animal among ≥2

independent experiments/animals were shown for quantification of intensities.

1355

Prophase-I stages were defined by SYCP3 staining (spermatocytes) and both SYCP3 staining

and fetal age (oocytes) using standard criteria. Leptonema was defined by short SYCP3

stretches without evidence of synapsis determined by thickening of SYCP3 staining. Zygonema

was defined by longer stretches of SYCP3 with various degree of synapsis: early, mid- and late-

zygonema were defined by having <25%, 25-75% and >75% of synapsis, respectively.

Pachynema was defined by full synapsis of all autosomes. For spermatocytes, early pachynema

was defined by extensive synapsis between the X-Y chromosomes, mid pachynema was

defined by limited/end-to-end X-Y synapsis, and late pachynema was defined by

decondensed/elongated X-Y chromosomes with figure-of eight configuration and thickening of

SYCP3 staining at telomeres of autosomes. Diplonema in spermatocytes was defined by

internal desynapsis with various degrees of residual synapsis: early, mid and late diplonema

were defined by having >75%, 75-25% and<25% of synapsis, respectively. For oocytes, nuclei

with unsynapsed/desynapsed chromosomes at embryonic day 15.5 (E15.5) and at 0.5 day

postpartum (0.5 dpp) were defined as zygonema and diplonema, respectively. Dictyate stage

was defined by short, ragged SYCP3 stretches without evidence of synapsis at 0.5 dpp.

1371

1372   For ovary sections, numbers of oocytes were manually counted for every fifth section and

1373   counts were multiplied by five to calculate the total number of oocytes per pair of ovaries, as

1374   described previously [65].

1375

1376   **Statistical analysis**

1377   Statistical analyses were performed using Graphpad Prism software v.8-9 and R version 3.5.2.

1378   Statistical parameters and tests, sample sizes, means, error bars, and standard deviations

1379   (SDs) are described in the figures and/or corresponding figure legends. Sample sizes were not

1380   predetermined using any statistical tests.

1381

Acknowledgements

We thank M.A. Handel, P. Cohen, C. Höög, J. Ward, and R.J. Pezza for antibodies; the UC

Davis MCB Light Microscopy Imaging Facility and M.R. Paddy for help with SIM imaging; the

Cornell Stem Cell and Transgenic Core Facility, supported in part by Empire State Stem Cell

Fund Contract Number C024174, J. Schimenti and R.J. Munroe for generating *Rnf212b* mutant

mice; and members of the Hunter lab for helpful discussions. M.I. was supported by a Japan

Society for the Promotion of Science postdoctoral fellowship for research abroad. S.S. was

supported by A.P. Giannini postdoctoral fellowship. B.N. was supported by the Howard Hughes

Medical Institute EXROP program. N.H. is an Investigator with the Howard Hughes Medical

Institute.

1392

Author contributions

M.I. and N.H. conceived the study and designed the experiments. M.I., S.L., L.H., K.L., N.L.,

R.M.H., R.P., C.H., and I.H. performed the experiments and analyzed the data. Y.Y. performed

oocyte culture experiments and analyzed the data. S.S. and B.N. performed histological

experiments and analyzed the data. D.S.K. generated anti-RNF212B antibodies. M.I. and N.H.

wrote the manuscripts with input from Y.Y., S.L., S.S., and D.S.K. All authors edited the

manuscript.

1400

1401   **Competing financial interests**

1402   The authors declare no competing financial interests.

1403

REFERENCES

1.   Watanabe, Y. Geometry and force behind kinetochore orientation: lessons from meiosis.

Nat Rev Mol Cell Biol **13**, 370-382 (2012).

2.   Hunter, N. Meiotic Recombination: The Essence of Heredity. *Cold Spring Harb Perspect*

Biol **7** (2015).

3.   Zickler, D. & Kleckner, N. Meiosis: Dances Between Homologs. *Annu Rev Genet* (2023).

4.   Faisal, I. & Kauppi, L. Reduced MAD2 levels dampen the apoptotic response to non-

exchange sex chromosomes and lead to sperm aneuploidy. *Development* **144**, 1988-

1996 (2017).

5.   Hassold, T., Hall, H. & Hunt, P. The origin of human aneuploidy: where we have been,

where we are going. *Human molecular genetics* **16 Spec No. 2**, R203-208 (2007).

6.   Hassold, T. & Hunt, P. To err (meiotically) is human: the genesis of human aneuploidy.

Nat Rev Genet **2**, 280-291 (2001).

7.   Faisal, I. & Kauppi, L. Sex chromosome recombination failure, apoptosis, and fertility in

male mice. *Chromosoma* **125**, 227-235 (2016).

8.   Hassold, T. *et al.* Failure to recombine is a common feature of human oogenesis. *Am J*

Hum Genet **108**, 16-24 (2021).

9.   Wang, S. *et al.* Inefficient Crossover Maturation Underlies Elevated Aneuploidy in

Human Female Meiosis. *Cell* **168**, 977-989 e917 (2017).

10.   Wartosch, L. *et al.* Origins and mechanisms leading to aneuploidy in human eggs.

Prenat Diagn **41**, 620-630 (2021).

11.   Cole, F. *et al.* Homeostatic control of recombination is implemented progressively in

mouse meiosis. *Nat Cell Biol* **14**, 424-430 (2012).

12.   Wang, S., Zickler, D., Kleckner, N. & Zhang, L. Meiotic crossover patterns: obligatory

crossover, interference and homeostasis in a single process. *Cell cycle* **14**, 305-314

(2015).

13.   Chelysheva, L. *et al.* The Arabidopsis HEI10 is a new ZMM protein related to Zip3. *PLoS*

Genet **8**, e1002799 (2012).

14.   Cheng, C.H. *et al.* SUMO modifications control assembly of synaptonemal complex and

polycomplex in meiosis of Saccharomyces cerevisiae. *Genes & development* **20**, 2067-

2081 (2006).

15.   Reynolds, A. *et al.* RNF212 is a dosage-sensitive regulator of crossing-over during

mammalian meiosis. *Nat Genet* **45**, 269-278 (2013).

16.   Qiao, H. *et al.* Antagonistic roles of ubiquitin ligase HEI10 and SUMO ligase RNF212

regulate meiotic recombination. *Nat Genet* **46**, 194-199 (2014).

17.   Rao, H.B. *et al.* A SUMO-ubiquitin relay recruits proteasomes to chromosome axes to

regulate meiotic recombination. *Science* **355**, 403-407 (2017).

18.   De Muyt, A. *et al.* E3 ligase Hei10: a multifaceted structure-based signaling molecule

with roles within and beyond meiosis. *Genes Dev* **28**, 1111-1123 (2014).

19.   Zhang, L., Kohler, S., Rillo-Bohn, R. & Dernburg, A.F. A compartmentalized signaling

network mediates crossover control in meiosis. *Elife* **7**, e30789 doi:

30710.37554/eLife.30789 (2018).

20.   Nguyen, H., Labella, S., Silva, N., Jantsch, V. & Zetka, M. C. elegans ZHP-4 is required

at multiple distinct steps in the formation of crossovers and their transition to segregation

competent chiasmata. *PLoS genetics* **14**, e1007776 (2018).

21.   Lake, C.M. *et al.* Narya, a RING finger domain-containing protein, is required for meiotic

DNA double-strand break formation and crossover maturation in Drosophila

melanogaster. *PLoS genetics* **15**, e1007886 (2019).

22.   Wang, K. *et al.* The role of rice HEI10 in the formation of meiotic crossovers. *PLoS*

Genet **8**, e1002809 (2012).

23.   Lake, C.M. *et al.* Vilya, a component of the recombination nodule, is required for meiotic

double-strand break formation in Drosophila. *Elife* **4**, e08287 (2015).

24.   Bhalla, N., Wynne, D.J., Jantsch, V. & Dernburg, A.F. ZHP-3 acts at crossovers to couple

meiotic recombination with synaptonemal complex disassembly and bivalent formation in

C. elegans. *PLoS Genet* **4**, e1000235 (2008).

25.   Ward, J.O. *et al.* Mutation in mouse hei10, an e3 ubiquitin ligase, disrupts meiotic

crossing over. *PLoS Genet* **3**, e139 (2007).

26.   Shinohara, M., Oh, S.D., Hunter, N. & Shinohara, A. Crossover assurance and crossover

interference are distinctly regulated by the ZMM proteins during yeast meiosis. *Nat*

Genet **40**, 299-309 (2008).

27.   Pyatnitskaya, A., Borde, V. & De Muyt, A. Crossing and zipping: molecular duties of the

ZMM proteins in meiosis. *Chromosoma* **128**, 181-198 (2019).

28.   Snowden, T., Acharya, S., Butz, C., Berardini, M. & Fishel, R. hMSH4-hMSH5

recognizes Holliday Junctions and forms a meiosis-specific sliding clamp that embraces

homologous chromosomes. *Molecular cell* **15**, 437-451 (2004).

29.   Kulkarni, D.S. *et al.* PCNA activates the MutLgamma endonuclease to promote meiotic

crossing over. *Nature* (2020).

30.   Cannavo, E. *et al.* Regulation of the MLH1-MLH3 endonuclease in meiosis. *Nature*

(2020).

31.   Jessop, L., Rockmill, B., Roeder, G.S. & Lichten, M. Meiotic chromosome synapsis-

promoting proteins antagonize the anti-crossover activity of sgs1. *PLoS Genet* **2**, e155

(2006).

32.   De Muyt, A. *et al.* BLM helicase ortholog Sgs1 is a central regulator of meiotic

recombination intermediate metabolism. *Mol Cell* **46**, 43-53 (2012).

33.   Zakharyevich, K., Tang, S., Ma, Y. & Hunter, N. Delineation of joint molecule resolution

pathways in meiosis identifies a crossover-specific resolvase. *Cell* **149**, 334-347 (2012).

34.   Tang, S., Wu, M.K.Y., Zhang, R. & Hunter, N. Pervasive and essential roles of the Top3-

Rmi1 decatenase orchestrate recombination and facilitate chromosome segregation in

meiosis. *Mol Cell* **57**, 607-621 (2015).

35.   Morgan, C. *et al.* Diffusion-mediated HEI10 coarsening can explain meiotic crossover

positioning in Arabidopsis. *Nat Commun* **12**, 4674 (2021).

36.   Fozard, J.A., Morgan, C. & Howard, M. Coarsening dynamics can explain meiotic

crossover patterning in both the presence and absence of the synaptonemal complex.

Elife **12** (2023).

37.   Zhang, L., Stauffer, W., Zwicker, D., A.F., D. & Crossover patterning through kinase-

regulated condensation and coarsening of recombination nodules. *bioRxiv*

2021.08.26.457865 (2021).

38.   Kleckner, N. *et al.* A mechanical basis for chromosome function. *Proc Natl Acad Sci U S*

A **101**, 12592-12597 (2004).

39.   Zhang, L., Liang, Z., Hutchinson, J. & Kleckner, N. Crossover patterning by the beam-

film model: analysis and implications. *PLoS Genet* **10**, e1004042 (2014).

40.   Johnston, S.E., Stoffel, M.A. & Pemberton, J.M. Variants at *RNF212* and *RNF212B* are

associated with recombination rate variation in Soay sheep (*Ovis aries*). *bioRxiv*

2020.07.26.217802 (2020).

41.   Gershoni, M. *et al.* A pathogenic variant in the uncharacterized RNF212B gene results in

severe aneuploidy male infertility and repeated IVF failure. *HGG Adv* **4**, 100189 (2023).

42.   Kadri, N.K. *et al.* Coding and noncoding variants in HFM1, MLH3, MSH4, MSH5,

RNF212, and RNF212B affect recombination rate in cattle. *Genome Res* **26**, 1323-1332

(2016).

43.   Johnston, S.E., Huisman, J. & Pemberton, J.M. A Genomic Region Containing REC8

and RNF212B Is Associated with Individual Recombination Rate Variation in a Wild

Population of Red Deer (Cervus elaphus). *G3 (Bethesda)* **8**, 2265-2276 (2018).

44.   Johnston, S.E., Berenos, C., Slate, J. & Pemberton, J.M. Conserved Genetic

Architecture Underlying Individual Recombination Rate Variation in a Wild Population of

Soay Sheep (Ovis aries). *Genetics* **203**, 583-598 (2016).

45.   Condezo, Y.B. *et al.* RNF212B E3 ligase is essential for crossover designation and

maturation during male and female meiosis in the mouse. *Proc Natl Acad Sci U S A* **121**,

e2320995121 (2024).

46.   Bondarieva, A. *et al.* Proline-rich protein PRR19 functions with cyclin-like CNTD1 to

promote meiotic crossing over in mouse. *Nat Commun* **11**, 3101 (2020).

47.   Ashley, T., Walpita, D. & de Rooij, D.G. Localization of two mammalian cyclin dependent

kinases during mammalian meiosis. *Journal of cell science* **114**, 685-693 (2001).

48.   Brown, M.S. & Bishop, D.K. DNA strand exchange and RecA homologs in meiosis. *Cold*

Spring Harb Perspect Biol **7**, a016659 (2014).

49.   Hinch, A.G. *et al.* The Configuration of RPA, RAD51, and DMC1 Binding in Meiosis

Reveals the Nature of Critical Recombination Intermediates. *Mol Cell* **79**, 689-701 e610

(2020).

50.   Baudat, F., Manova, K., Yuen, J.P., Jasin, M. & Keeney, S. Chromosome synapsis

defects and sexually dimorphic meiotic progression in mice lacking Spo11. *Molecular cell*

6, 989-998 (2000).

51.   Romanienko, P.J. & Camerini-Otero, R.D. The mouse Spo11 gene is required for meiotic

chromosome synapsis. *Molecular cell* **6**, 975-987 (2000).

52.   de Vries, F.A. *et al.* Mouse Sycp1 functions in synaptonemal complex assembly, meiotic

recombination, and XY body formation. *Genes Dev* **19**, 1376-1389 (2005).

53.   Lipkin, S.M. *et al.* Meiotic arrest and aneuploidy in MLH3-deficient mice. *Nat Genet* **31**,

385-390 (2002).

54.   McMahill, M.S., Sham, C.W. & Bishop, D.K. Synthesis-dependent strand annealing in

meiosis. *PLoS biology* **5**, e299 (2007).

55.   Palmer, N. *et al.* A novel function for CDK2 activity at meiotic crossover sites. *PLoS*

biology **18**, e3000903 (2020).

56.   Holloway, J.K., Sun, X., Yokoo, R., Villeneuve, A.M. & Cohen, P.E. Mammalian CNTD1 is

1535   critical for meiotic crossover maturation and deselection of excess precrossover sites. *J*

1536   *Cell Biol* **205**, 633-641 (2014).

1537   57.   Qiao, H. *et al.* Interplay between synaptonemal complex, homologous recombination,

1538   and centromeres during mammalian meiosis. *PLoS genetics* **8**, e1002790 (2012).

1539   58.   Serrentino, M.E., Chaplais, E., Sommermeyer, V. & Borde, V. Differential association of

1540   the conserved SUMO ligase Zip3 with meiotic double-strand break sites reveals regional

1541   variations in the outcome of meiotic recombination. *PLoS Genet* **9**, e1003416 (2013).

1542   59.   Premkumar, T. *et al.* Genetic dissection of crossover mutants defines discrete

1543   intermediates in mouse meiosis. *Mol Cell* **83**, 2941-2958 e2947 (2023).

1544   60.   van der Heijden, G.W. *et al.* Chromosome-wide nucleosome replacement and H3.3

1545   incorporation during mammalian meiotic sex chromosome inactivation. *Nat Genet* **39**,

1546   251-258 (2007).

1547   61.   Ghafari, F., Gutierrez, C.G. & Hartshorne, G.M. Apoptosis in mouse fetal and neonatal

1548   oocytes during meiotic prophase one. *BMC Dev Biol* **7**, 87 (2007).

62.   Qiao, H. *et al.* Impeding DNA Break Repair Enables Oocyte Quality Control. *Mol Cell* **72**,

211-221 e213 (2018).

63.   Abe, H. *et al.* The Initiation of Meiotic Sex Chromosome Inactivation Sequesters DNA

Damage Signaling from Autosomes in Mouse Spermatogenesis. *Curr Biol* **30**, 408-420

e405 (2020).

64.   Bassett, A.R., Tibbit, C., Ponting, C.P. & Liu, J.L. Highly efficient targeted mutagenesis of

Drosophila with the CRISPR/Cas9 system. *Cell Rep* **4**, 220-228 (2013).

65.   Peters, A.H., Plug, A.W., van Vugt, M.J. & de Boer, P. A drying-down technique for the

spreading of mammalian meiocytes from the male and female germline. *Chromosome*

Res **5**, 66-68 (1997).

66.   Di Giacomo, M. *et al.* Distinct DNA-damage-dependent and -independent responses

drive the loss of oocytes in recombination-defective mouse mutants. *Proc Natl Acad Sci*

U S A **102**, 737-742 (2005).

1562

1563