Nanoscale cellular organization of viral RNA and proteins in SARS-CoV-2 replication organelles

Leonid Andronov*[1], Mengting Han*[2], Yanyu Zhu[2], Ashwin Balaji[1,5], Anish R. Roy[1], Andrew E. S.

Barentine[1], Puja Patel[3], Jaishree Garhyan[3], Lei S. Qi[2,4,6,#], and W.E. Moerner[1,4,#]

[1]Department of Chemistry; [2]Department of Bioengineering; [3]In Vitro Biosafety Level 3 (BSL-3)

Service Center, School of Medicine; [4]Sarafan ChEM-H; [5]Biophysics PhD Program; Stanford

University, Stanford, CA 94305 U.S.A. [6]Chan Zuckerberg Biohub – San Francisco, San Francisco, CA

94158 U.S.A.

*These authors contributed equally.

#Correspondence to: W. E. Moerner, wmoerner@stanford.edu; Lei S. Qi, sqi@stanford.edu

Abstract:

The SARS-CoV-2 viral infection transforms host cells and produces special organelles in many ways, and we focus on the replication organelle where the replication of viral genomic RNA (vgRNA) occurs. To date, the precise cellular localization of key RNA molecules and replication intermediates has been elusive in electron microscopy studies. We use super-resolution fluorescence microscopy and specific labeling to reveal the nanoscopic organization of replication organelles that contain vgRNA clusters along with viral double-stranded RNA (dsRNA) clusters and the replication enzyme, encapsulated by membranes derived from the host endoplasmic reticulum (ER). We show that the replication organelles are organized differently at early and late stages of infection. Surprisingly, vgRNA accumulates into distinct globular clusters in the cytoplasmic perinuclear region, which grow and accommodate more vgRNA molecules as infection time increases. The localization of ER labels and nsp3 (a component of the double-membrane vesicle, DMV) at the periphery of the vgRNA clusters suggests that replication organelles are enclosed by DMVs at early infection stages which then merge into vesicle packets as infection progresses. Precise co-imaging of the nanoscale cellular organization of vgRNA, dsRNA, and viral proteins in replication organelles of SARS-CoV-2 may inform therapeutic approaches that target viral replication and associated processes.

Keywords: SARS-CoV-2 coronavirus, viral replication, fluorescence imaging, super-resolution fluorescence microscopy, viral RNA localization, COVID-19, viral proteins, cell infection

Introduction

Due to its global health impact, the SARS-CoV-2 betacoronavirus and its infection of mammalian cells have been the subject of a large number of studies across multiple fields. Biochemical methods have allowed researchers to investigate the interactions between the viral RNA and the host proteins *in vitro* and in cellular extracts, leading to much insight[1,2]. There have also been electron microscopy (EM) studies of resin-embedded samples as well as vitrified samples using cryo-electron tomography, all of which have been profiting from the large increase in EM resolution and contrast in recent years. These EM studies can provide very high-resolution structures of protein complexes as well as tomograms of organelles in the cellular context. High-contrast filamentous structures and membranes appear regularly in such images, allowing identification of single- and double-membrane vesicles (DMVs)[3,4,5]. However, the all-important viral RNA and associated proteins are challenging to identify by EM due to a lack of specific contrast. While some researchers have detected RNA-like filaments in vesicles[4,5], further investigations are needed to identify specific viral RNAs in the cellular context.

Fluorescence microscopy offers a highly useful and complementary set of capabilities, most importantly the specific labeling of proteins or RNA sequences. However, conventional diffraction-limited (DL) fluorescence microscopy, with its resolution constrained to ~250 nm, is unable to resolve the tiny structures that are hidden in a blurred DL image. Super-resolution (SR) microscopy based on single molecules (PALM[6], (d)STORM[7,8]) or on structured patterns of molecular depletion (STED[9], SIM[10]), however, offers far better optical resolution down to 10 nm and below. A wealth of important cellular patterns and structures has been identified in recent years, such as the banding patterns of axonal proteins in neuronal cells[11] and many others[12,13,14]. The specificity of SR imaging is useful to apply to the study of viral genomic RNA (vgRNA) and other RNA molecules; moreover, additional nanoscale imaging of critical proteins involved in coronavirus infection of cells provides crucial context for the nearby partners and surroundings of the viral RNA. In a previous proof-of-principle study, we explored the relatively safe human coronavirus 229E (HCoV-229E) from the alphacoronavirus family, which uses the APN receptor and produces only mild cold symptoms[15].

In this work, we apply multicolor confocal microscopy and SR microscopy to explore the
localization patterns of viral RNA, related viral proteins, and altered host cell structures for SARS-
CoV-2 betacoronavirus during the early and late infection of mammalian cells. The SARS-CoV-2
life cycle starts with viral entry into a host cell, facilitated by binding of viral spike protein to its
canonical receptor at the cell surface, the angiotensin-converting enzyme 2 (ACE2)[16], or one of
the alternative receptors[17]. The subsequent fusion of the viral and the host cell membranes
releases the viral genetic material, positive-sense single-stranded viral genomic RNA (vgRNA), into
the cytoplasm, where it is readily translated by host ribosomes. SARS-CoV-2 vgRNA (Fig. 1a)
encodes at least 29 proteins, including structural proteins that make up the virions, and non-
structural (NSPs) and accessory proteins that exist only within host cells and regulate various
processes in the intracellular viral life cycle. All NSPs originate from polyproteins that are
translated directly from vgRNA and are self-cleaved by viral proteases. Structural and accessory
proteins are translated from shorter viral genome fragments called subgenomic RNAs (sgRNAs)
that are transcribed from vgRNA.

Replication and transcription of the viral genome is carried out by the RNA-dependent RNA
polymerase complex (RdRp), which is assembled from nsp12 (RdRp catalytic subunit) along with
nsp7 and nsp8 (accessory subunits)[18]. RdRp first synthesizes either a full-length negative-sense
copy of vgRNA or a subgenomic negative-sense copy of vgRNA, producing double-stranded RNA
(dsRNA) that forms between vgRNA and the negative-sense copy. Next, using this negative-sense
template, a new vgRNA or an sgRNA is generated by the same polymerase enzyme. Additional
NSPs modify newly synthesized viral RNAs to form 5' cap structures[19] that mimic cellular mRNAs
to be translated by host ribosomes. The replication intermediates, such as dsRNA and uncapped
RNAs, might be degraded or trigger innate immune response[20] and therefore need to be
protected from cellular machinery. SARS-CoV-2 transforms host ER into DMVs[21] that are abundant
in the perinuclear region of infected cells[4,5,22] and likely encapsulate dsRNA[3,5] and newly
synthesized viral RNA[4,23]. However, the precise intracellular localization of replicating RdRp
enzymes and therefore of the replication events is not well established to date[3,23,24].

In this work, we focus particularly on vgRNA, dsRNA and key RdRp subunits nsp12, nsp7 and nsp8.
We also co-image a series of molecules, including membrane markers, nucleocapsid protein, spike

protein, and the nsp3 protein (reported to be a major component of a molecular pore spanning

both membranes of DMVs[25]), all to provide context and support for the view that vgRNA, dsRNA,

and RdRp act spatially in replication organelles (ROs) during viral replication. Thus, we provide

key information about where these important players are found in infected cells and how they

change with time during early *vs* late infection. Our results yield a nanoscale optical readout of

viral nucleic acid organization and viral proteins down to 20-40 nm during SARS-CoV-2 infection,

highlight the structural importance of ROs, and could potentially benefit development of future

therapeutic approaches.

Results

Labeling and imaging of SARS-CoV-2 virions

To specifically detect SARS-CoV-2 vgRNA, we applied RNA fluorescence *in situ* hybridization (RNA

FISH) with 48 antisense DNA oligonucleotide probes[26] specifically targeting the open reading

frame 1a (ORF1a) region which is only present in vgRNA and not in subgenomic RNAs (sgRNAs),

ensuring detection of only full-length viral positive-sense vgRNA (Fig. 1a). Each probe was

conjugated with a single blinking fluorophore for (d)STORM (direct Stochastic Optical

Reconstruction Microscopy)[8]. To test this labeling and imaging approach, we first imaged vgRNA

along with SARS-CoV-2 spike protein in purified virions (Fig. S1). While the size of SARS-CoV-2

virions is too small to resolve in conventional DL fluorescence microscopy (Fig. S1a), in SR the

internal concentric organization of the virions can be observed with vgRNA found in their center

and spike at the surface (Fig. S1b). The labeling efficiency with these probes is around 6

dyes/vgRNA in partially Proteinase K-digested virions, which was higher than in intact virions (1.7

dyes/vgRNA) due to poorer accessibility of their vgRNA (Fig. S1c-i).

Next, we imaged SARS-CoV-2 infected Vero E6 cells that were fixed at 24 hours post infection (hpi)

and then labeled for immunofluorescence imaging (Methods). SR microscopy of spike and

nucleocapsid proteins in these cells revealed assembled virions mostly at the cellular periphery,

often at cytoplasmic tubular projections of the plasma membrane, indicating active viral

production (Fig. S1j), similar to previously reported results[5,27]. These studies of virions highlight

the improved resolution of SR microscopy and validate the labeling approach, but much more is

to be learned by imaging viral RNA and proteins in the cellular interior. We now turn to the main

focus of this study, the intracellular replication of viral genomic RNA.

SARS-CoV-2 genomic RNA clusters in cytoplasm of infected cells

Confocal screening demonstrated three patterns of intracellular vgRNA localization (Fig. S2a):

scattered puncta in the cytoplasm (Type 1, Fig. 1b), appearance of bright foci in the perinuclear

region (Type 2, Fig. S2a), and concentration of vgRNA into large dense structures that occupy most

of the perinuclear region (Type 3, Fig. 1f). We find that Type 1 cells were most abundant at 6 hpi,

and Type 3 cells at 24 hpi, indicating that the vgRNA localization progresses from Type 1 to Type

3 as infection advances in time (Fig. S2b). We also find that the cell-integrated vgRNA FISH signal

in infected cells increases 2.2x on average from 6 to 24 hpi (Fig. S2c), representing active viral

replication and accumulation of vgRNA inside the cells.

Fig. 1: **Clustering of vgRNA in the cytoplasm of infected cells.**

a, Scheme of SARS-CoV-2 genome with constructs used for its detection in infected cells. 48 antisense DNA oligonucleotide probes were used to target the ORF1a-coding region of vgRNA that is exclusive to the positive-sense vgRNA and does not occur in the sgRNAs. The RNA FISH probes are conjugated with AF647 or CF568. **b**, Representative confocal images of vgRNA in infected Vero E6 cells at 6 hpi display scattered diffraction-limited (DL) puncta. **c**, Representative SR image of an infected cell at 6 hpi reveals distinct vgRNA clusters in the cytoplasm. **d**, Zoomed-in region of the SR image (green frame in **c**) displays an agglomeration of vgRNA clusters. **e**, Zoomed-in region of the SR image (red frame in **c**) shows nanoscale puncta of individual vgRNA molecules. **f**, Representative confocal images of vgRNA in infected Vero E6 cells at 24 hpi display large DL foci in the perinuclear region of the cytoplasm. **g**, Representative SR image of an infected cell at 24 hpi reveals large perinuclear vgRNA clusters. **h**, Zoomed-in region of the SR image (blue frame in **g**) displays dense vgRNA clusters. **i**, Zoomed-in region of the SR image (yellow frame in **g**) displays nanoscale puncta of vgRNA molecules. **j**, BIC-GMM cluster analysis of the region shown in **d**. **k**, BIC-GMM cluster analysis of the cell shown in **g**. **l**, BIC-GMM cluster analysis of the region shown in **h**. **m**, Histogram of the radii of gyration (Rg) of the vgRNA clusters indicate their size increase between 6 hpi (magenta) and 24 hpi (green). Scale bars, 10 μm (**b**, **f**), 2 μm (**c**, **g**, **k**), 500 nm (**d**, **e**, **h**, **i**, **j**, **l**). Dashed lines in **c** and **g** indicate the boundary of the cell nucleus (large dark region). Localizations that belong to the same cluster in **j**, **k**, **l** are depicted with the same color, but colors are reused. Color bars in **c**, **d**, **e**, **g**, **h**, **i** show the number of single-molecule localizations within each SR pixel (20 x 20 nm^2).

The higher spatial resolution of SR microscopy revealed that at 6 hpi (Type 1 and Type 2 cells), most vgRNA localizes into clusters with an approximately round shape and a diameter of 100-250 nm that scatter in the cytoplasm (Fig. 1c, d). At 24 hpi (Type 2 and Type 3 cells), the vgRNA localization pattern transformed into a fascinating dense perinuclear network of approximately round structures with a diameter of 300-700 nm (Fig. 1g, h). The lower number of localizations in many cluster centers suggests a possibly hollow structure in these 2D images; future 3D imaging can explore more aspects of the cellular localization. To quantify the transformation of vgRNA clusters in infected cells, we performed a clustering analysis using a Bayesian Information Criterion-optimized Gaussian Mixture Model (BIC-GMM) (Fig. 1j-l; see Methods). This analysis showed an increase in the median vgRNA cluster size (Rg, radius of gyration) from 73 nm at 6 hpi to 187 nm at 24 hpi (Fig. 1m), reflecting the drastic change in vgRNA localization pattern.

Besides dense vgRNA clusters, we observe isolated localizations of individual vgRNA molecules

scattered in the cytoplasm at both time points, in line with previously reported results[15,26]. These

appear as a haze in confocal images (Fig. S2a, Type 3) but are resolved as sparse nanoscale puncta

(d < 50 nm) in SR (Fig. 1e, i; Fig. S3a) which we assume to be single vgRNA copies (even though

the puncta are denser at 24 hpi). Using the average number of single-molecule (SM) localizations

per vgRNA punctum as a calibration for the number of localizations per single vgRNA, we

estimated the average number of vgRNA molecules in the vgRNA clusters to be around 26

vgRNA/cluster at 6 hpi, increasing by almost an order of magnitude to 181 vgRNA/cluster at 24

hpi (Fig. S3b-c; procedure detailed in Methods).

dsRNA associates with vgRNA clusters

Next, we proceeded to assess the relation of vgRNA cluster locations to viral replication. For this,

we immunofluorescently labeled an intermediate of coronavirus replication and transcription,

the hybridized dsRNA objects composed of positive-sense vgRNA and negative-sense copy, and

co-imaged dsRNA with vgRNA using two-color confocal and SR microscopy. In confocal

microscopy, dsRNA labeling was present in all cells with detectable vgRNA FISH fluorescence,

including in early infection, demonstrating the high sensitivity of our dsRNA immunofluorescence

detection (Fig. S2d). dsRNA and vgRNA appeared mostly colocalized at both time points at low

resolution (Fig. 2a, b), suggesting that vgRNA clusters are often found close to the replication

centers of SARS-CoV-2. SR microscopy revealed that dsRNA aggregates into clusters of a relatively

compact size (d ≈ 100-200 nm) with distinct patterns of colocalization with vgRNA at 6 or 24 hpi

(Fig. 2c, d).

To quantify the spatial relationship between dsRNA and vgRNA, we conducted pair-pair

correlation analysis[28]. We calculated a bivariate pair-correlation function $g_{12}(r)$, *i.e.*, the

distribution of the pairwise distances between the localizations of the two species[29]. The function

is computed only in perinuclear regions and is normalized in a way that $g_{12}(r) = 1$ for two randomly

and homogeneously distributed species without interaction, signifying complete spatial

randomness (CSR). Closely associated or colocalized species have a prevalence of short pairwise

distances resulting in a peak in $g_{12}(r)$ near r = 0, while anti-correlated species lack short
interparticle distances, which lowers $g_{12}(r)$ at r = 0 followed by peaking at r > 0.

Fig. 2: Association of dsRNA with vgRNA clusters

a-b, Representative confocal images of SARS-CoV-2 infected cells display DL colocalization
between dsRNA (green) and vgRNA (magenta) at both 6 hpi (**a**) and 24 hpi (**b**). **c-d**, Representative
SR images of SARS-CoV-2 infected cells indicate association between dsRNA and vgRNA at 6 hpi

(**c**) and short-range anti-correlation often with concentric localization at 24 hpi (**d**). Bottom panels, zoomed-in images of corresponding colored boxes. **e**, Bivariate pair-correlation functions $g_{12}(r)$ calculated between the localizations of dsRNA and vgRNA indicate their close association at 6 hpi. **f**, Histogram of Rg of dsRNA clusters as determined by the BIC-GMM cluster analysis. **g**, Median Rg of dsRNA clusters per cell significantly decreases between 6 hpi and 24 hpi. p-value = $8 \cdot 10^{-4}$, two-tailed t-test. Error bars represent mean ± standard deviation of median Rg values of dsRNA clusters in individual cells. **h**, Bivariate pair-correlation functions $g_{12}(r)$ reveal nanoscale anti-correlation between dsRNA and vgRNA at 24 hpi. CSR, complete spatial randomness. Thin lines correspond to $g_{12}(r)$ of individual cells and bold lines are the mean values of $g_{12}(r)$ from all analyzed cells. Scale bars, 10 μm (**a-b**), 2 μm (**c-d**), 500 nm (**c-d**, bottom panels). Dashed lines in **c** and **d** indicate the boundary of the cell nucleus.

At early infection stages (6 hpi), dsRNA clusters appear closely associated with or adjacent to vgRNA clusters both visually and by pair-pair correlation analysis (Fig. 2c, e). By contrast, during late infection (24 hpi), dsRNA clusters anticorrelate with vgRNA at short distance scales with an average separation between them around 120 nm as indicated by bivariate pair-correlation functions $g_{12}(r)$ (Fig. 2h). Moreover, at 24 hpi, dsRNA clusters can often be found in the central voids of the large vgRNA structures (Fig. 2d), suggesting their possible concentric localization in the same ROs.

Contrary to vgRNA, the size of dsRNA clusters slightly decreases (Fig. 2f, g) and the total brightness of cellular dsRNA labeling does not significantly change between 6 hpi and 24 hpi (Fig. S2e). Interestingly, at 6 hpi but not at 24 hpi, the dsRNA signal per cell positively correlates with that of vgRNA (Fig. S2f, g). These findings indicate that the amount of dsRNA increases at early infection but reaches saturation by 24 hpi. This may suggest that after the rapid initial production of a dsRNA pool, further generation of negative-sense copies slows down and the replication shifts to the generation of vgRNA from the pool of available negative-sense templates, which is common in other coronaviruses[30].

vgRNA clusters denote the replication centers of SARS-CoV-2 genome

To investigate SARS-CoV-2 replication activity within the vgRNA clusters in more detail, we co-

imaged them with the RdRp complex, the replicating SARS-CoV-2 RNA-dependent RNA

polymerase[18,31], using immunofluorescent labeling of its catalytic subunit nsp12[32]. In confocal

images, nsp12 adopts a similar pattern as vgRNA, colocalizing with it at both 6 hpi and 24 hpi (Fig.

3a-b), which suggests ongoing replication at the vgRNA clusters. In SR images, nsp12 localized in

small sparse puncta (d < 50 nm) that were scattered within and next to the vgRNA clusters at both

time points (Fig. 3c-d). Because nanoscale nsp12 puncta are well separated from each other, and

oligomerization is not expected[18,31,33], each nanoscale punctum is likely to represent a single

replicating enzyme. On average, we detected 2.5 nsp12 puncta per vgRNA cluster at 6 hpi and 7.6

at 24 hpi (Fig. 3h).

From comparison of DL and SR images, one may infer fundamentally different (large-scale) nsp12

structures at 6 hpi and 24 hpi in confocal microscopy (Fig. 3a-b). In DL microscopy, ROs do look

like individual diffraction-limited dots at 6 hpi when they are sparse (Fig. 3a), *i.e.*, the average

distance between them is larger than the diffraction limit (even though the individual RdRp

complexes inside ROs are still not resolved). The same organelles when they are dense at 24 hpi

resemble large irregular blobs because the distance between the individual organelles becomes

smaller than the diffraction limit (Fig. 3b). This filling in with optically overlapping ROs creates a

misleading perception of distinct structures in confocal microscopy. However, SR microscopy,

which sees spatial details on the scale of 20-40 nm, resolves both types of structures much better.

The nsp12 puncta are small in both cases because they arise from individual RdRp enzymes, yet

the vgRNA clusters are smaller at 6 hpi and larger at 24 hpi, which is a better representation of

the size of these assemblies.

Therefore, in contrast to vgRNA but similar to dsRNA, the total cellular amount of nsp12 does not

significantly increase (Fig. S2h) and its nanoscale localization pattern stays the same as infection

progresses from 6 to 24 hpi (Fig. 3c, d). This suggests that the growth of vgRNA clusters arises

from a relatively constant small number of replication components between 6 and 24 hpi

highlighted by the constant amount of dsRNA and RdRp. Bivariate cross-correlation functions

calculated between nsp12 and vgRNA localizations peaked at 0 nm indicating association of these

two targets at both 6 and 24 hpi (Fig. 3i). Since vgRNA clusters colocalize with the catalytic subunit

 of RdRp, we suggest that vgRNA clusters combined with the nearby RdRp enzymes and dsRNA

 highlight ROs that act as centers for replication and transcription of SARS-CoV-2.

Fig. 3: Association of SARS-CoV-2 replication enzyme with vgRNA clusters

a-b, Representative confocal images of SARS-CoV-2 infected cells display DL colocalization between nsp12, the catalytic subunit of RdRp (green) and vgRNA (magenta) at both 6 hpi (**a**) and 24 hpi (**b**). **c-d**, Representative SR images of SARS-CoV-2 infected cells indicate nanoscale association between nsp12 and vgRNA at both 6 hpi (**c**) and 24 hpi (**d**). Insets show magnified images of corresponding regions in the colored boxes. **e-f**, Representative SR images of vgRNA with nsp7 (**e**) or nsp8 (**f**) in the perinuclear regions of SARS-CoV-2 infected cells indicate association of nsp7 and nsp8 with vgRNA clusters. **g**, Representative SR image of vgRNA with

newly synthesized viral RNAs labeled by BrU in a SARS-CoV-2 infected cell indicates localization of
newly synthesized viral RNAs within the perinuclear clusters of vgRNA. **h**, Number of nanoscale
puncta of nsp12 per vgRNA cluster. **i**, Bivariate pair-correlation functions for vgRNA and nsp12,
nsp7, nsp8 and newly transcribed viral RNA labeled with BrU peak at r = 0 nm indicating
association between these target pairs. Scale bars, 10 µm (**a**-**b**), 2 µm (**c**-**d**), 500 nm (**e**-**g** and insets
in **c**-**d**). Dashed lines in **c** and **d** indicate the edge of the cell nucleus.

To verify that nsp12 labeling is a good reporter of assembled replication complexes, we have also

imaged two accessory subunits of RdRp, nsp7 and nsp8. We find close association of these

subunits with vgRNA as shown in Fig. 3e, f, and in the pair-correlation functions of Fig. 3i (see also

Fig. S4 and S5). Nsp12 and nsp8 colocalized with each other on the nanoscale (Fig. S6), indicating

their interaction within ROs, as expected for subunits of assembled RdRp.

Finally, to confirm that the vgRNA clusters we observe contain newly replicated viral RNA, we

provided brominated uridine (BrU) to the infected cells in the form of 5-bromouridine 5'-

triphosphate (BrUTP) for 1 hour before fixation while endogenous transcription was inhibited by

actinomycin D[34,35]. Immunofluorescent labeling of BrU then highlights newly replicated RNA.

Confocal and SR imaging localizes RNA-containing BrU to the perinuclear clusters of vgRNA (Fig.

3g, Fig. S7) and close to nsp12 (Fig. S8), further proving that these structures are the sites of active

replication and transcription of viral RNA.

vgRNA clusters are enclosed in ER-derived membranous organelles

Coronaviruses are known to transform the host ER into replication-permissive structures, such as

convoluted membranes and DMVs[3,22,36]. To investigate the relation of vgRNA clusters with cellular

ER, we immunofluorescently labeled Sec61β, an ER membrane protein[37], in Vero E6 cells stably

expressing Sec61β-GFP[15]. Confocal images of these cells show the appearance of Sec61β spots

that colocalize with vgRNA against the mostly unaltered ER background at 6 hpi (Fig. 4a). At

24 hpi, however, substantial amounts of Sec61β accumulate close to the perinuclear vgRNA

clusters, while the ER tubules outside these regions become poorly visible (Fig. 4a), consistent

with the virus-induced rearrangement of the ER and the inhibition of host gene expression by SARS-CoV-2[38].

Fig. 4: vgRNA clusters are encapsulated in membranes of remodeled ER

a, Representative confocal images of SARS-CoV-2 infected cells indicate an appearance of dense perinuclear foci of Sec61β ER labeling (green) at 24 hpi that colocalizes with vgRNA and nsp3. **b**, SR images reveal concentric organization of Sec61β around vgRNA, dsRNA and nsp12, and colocalization of Sec61β with nsp3. **c**, Bivariate pair-correlation functions indicate anti-correlation of Sec61β with vgRNA and dsRNA and association of Sec61β with nsp3. Scale bars, 20 μm (**a**) and 1 μm (**b**). Dashed lines in **b** indicate the boundary of the cell nucleus.

In SR, we observe encapsulation of the vgRNA clusters by ring-like structures of the altered ER at

6 hpi (Fig. 4b, Fig. S9). As infection progresses, the ER-derived ring- or sphere-like structures grow

to accommodate larger vgRNA clusters at 24 hpi (Fig. 4b, Fig. S10). Pair-correlation functions peak

at the distance of the typical radius of vgRNA clusters indicating nanoscale anti-correlation

compatible with the ER-derived encapsulation of vgRNA (Fig. 4c). dsRNA (Fig. 4b, Fig. S11) and

nsp12 (Fig. 4b) are also found to be encapsulated by the same remodeled ER membranes

suggesting that vgRNA, dsRNA and RdRp are all located within the same ER-derived ROs.

To further confirm that these clusters are surrounded by membranes, we used a (d)STORM-

compatible general membrane marker CellMask Deep Red[39]. This dye broadly stains cellular

membranes, including the nuclear envelope, mitochondrial membranes, and SARS-CoV-2 virions

at the plasma membrane (Fig. S12). The nanoscale image contrast with CellMask Deep Red is

poorer than specific protein labeling of the Sec61β ER label due to background from membranes

of different cellular organelles. Nevertheless, in the perinuclear region of infected cells, we

observed the appearance of a complex membranous network that anti-correlates with vgRNA

and dsRNA, with visible encapsulation of vgRNA and dsRNA clusters (Fig. S12, S13). Taken

together, these findings indicate that each vgRNA-dsRNA-RdRp cluster is located inside a

membrane-bound RO that originates from altered host ER transformed by SARS-CoV-2.

Nsp3, spike and nucleocapsid proteins localize at the surface of SARS-CoV-2 replication organelles

Because the nsp3 protein of betacoronaviruses is essential for the DMV formation[21,40], and nsp3

is a constituent of a DMV molecular pore[25], we proceeded to localize this non-structural protein

to relate the ROs to the SARS-CoV-2-induced DMVs. At DL resolution, nsp3 labeling adopts a

pattern that colocalizes with vgRNA at both 6 and 24 hpi, similar to dsRNA and nsp12 (Fig. 5a, f).

SR imaging of these cells, however, revealed striking nanoscale positioning of nsp3. At 6 hpi,

sparse nsp3 can be found surrounding isolated vgRNA clusters (Fig. 5b, c), while larger nsp3

aggregates are situated amidst bunched vgRNA clusters (Fig. 5d). At 24 hpi, nsp3 localizes at the

borders of the large vgRNA clusters, encircling them in incomplete rings and forming a partial

perinuclear network (Fig. 5g-i). Similar nsp3 arrangements can be observed in relation to dsRNA

(Fig. S14).

Fig. 5: Nsp3 localizes at the surface of vgRNA clusters

a, Representative confocal images of a SARS-CoV-2 infected cell display DL colocalization between

punctate vgRNA (magenta) and nsp3 (green) labeling at 6 hpi. **b,** Representative SR image of a

SARS-CoV-2 infected cell at 6 hpi. **c,** Zoomed-in images of selected vgRNA particles (yellow boxes

in **b**) indicate the localization of nsp3 at the surface of the vgRNA clusters. **d,** Magnified region

with aggregates of vgRNA clusters (blue box in **b**) displays dense nsp3 localization in the core of

these aggregates. **e**, Bivariate pair-correlation functions calculated between the SM localizations

of vgRNA and nsp3 indicate nanoscale anti-correlation of these targets at 6 hpi. **f**, Confocal images

show that vgRNA and nsp3 occupy approximately the same regions in a SARS-CoV-2 infected cell

at 24 hpi. **g**, Representative SR image of a SARS-CoV-2 infected cell at 24 hpi. **h-i**, Magnified regions

of the SR image (colored boxes in **g**) reveal that nsp3 localizes in interstitial regions or

encapsulates vgRNA clusters. **j**, Bivariate pair-correlation functions indicate nanoscale anti-

correlation between vgRNA and nsp3 at 24hpi. Scale bars, 10 µm (**a**, **f**), 2 µm (**b**, **g**), 500 nm (**d**, **h**,

i), 200nm (**c**). Dashed lines in **b**, **g** indicate the boundary of the cell nucleus.

The anti-correlation of vgRNA with nsp3 and dsRNA with nsp3 (Fig. 5, Fig. S14) closely resemble

the pattern observed with vgRNA and dsRNA with Sec61β (Fig. 4, Fig. S11), suggesting that nsp3

may also be localized at the ER-derived membranous surface of the ROs. To further confirm this

hypothesis, we co-imaged nsp3 with Sec61β and CellMask (Fig. 4, Fig. S13, S15). The SR images

and the pair-correlation analysis indicated colocalization between nsp3 and both membrane

markers at both time points (Fig. 4b-c, Fig. S13, S15), confirming that nsp3 localizes on the

membranes encircling the SARS-CoV-2 ROs.

Besides these characteristic localization patterns of nsp3, we observed a few cells with two

different phenotypes at 24 hpi, one with an ER-like network that occupies large regions in the

cytoplasm (Fig. S16a), and another one with nsp3 densely diffused throughout the whole

cytoplasm (Fig. S16b). The ER-like network may represent nsp3 proteins being heavily translated

on ER membranes, while nsp3 proteins found outside the perinuclear region are less likely to be

associated with the SARS-CoV-2 replication process and might represent other nsp3 functions,

such as a papain-like proteolytic function[41] or post-translational modification of host proteins[42],

which can become objects of future SR studies.

The localization of nsp3 at the surface of isolated vgRNA-dsRNA clusters at 6 hpi is consistent with

the localization of molecular pores on the DMV membrane observed by cryo-EM[25]. At late

infection times, DMVs have been observed to merge into vesicle packets (VPs)[5] that are also likely

to contain pores, however molecular pores in the VP membranes have not yet been studied in

detail to our knowledge. Nevertheless, previous studies report that in late infection the

perinuclear region becomes filled with DMVs and VPs[22] that strongly resemble the ROs reported

here. The size of vgRNA clusters at 6 hpi and at 24 hpi from our data is similar to the previously
reported size of DMVs and VPs, correspondingly[5].

To search for a possible role of perinuclear vgRNA clusters in virion assembly, we co-imaged
vgRNA with two SARS-CoV-2 structural proteins, spike and nucleocapsid (Fig. S17, S18). Spike
labeling forms typical ~ 150 nm hollow particles at the cell periphery, and we detect weak vgRNA
signal in the center of some of these particles (Fig. S17b), consistent with the structure of SARS-
CoV-2 virions that contain a single vgRNA molecule. Inside the host cells, spike localizes at the
nuclear envelope and in some cytoplasmic organelles; however, it is mostly excluded from the
perinuclear vgRNA clusters (Fig. S17a, c). Nucleocapsid protein demonstrates rather diffuse
localization throughout the cytoplasm, in accordance with its function in the formation of SARS-
CoV-2 ribonucleocapsid complexes[43], but is also excluded from the RO interior (Fig. S18a).
Nevertheless, in the perinuclear region we detect sparse localizations of both spike and
nucleocapsid proteins next to the vgRNA clusters and between them, likely at the DMV
membranes, as highlighted by anti-correlation of these proteins with vgRNA at r < 200 nm (Fig.
S17c, S18b), similar to the nsp3/vgRNA and Sec61β/vgRNA pairs. The localization of nucleocapsid
protein at the RO membranes has already been reported[44], and spike protein has a
transmembrane domain[45] and tends to localize not only to virion membranes, but also to
intracellular membranes, such as the nuclear envelope (Fig. S17a); therefore, small amounts of
spike can also be present at RO membranes. Our SR data suggests that while the vgRNA clusters
are not directly involved in SARS-CoV-2 virion assembly, it is possible that early stages of virion
assembly start at the RO membrane, once vgRNA molecules leave the ROs.

Taken together, our results provide evidence that vgRNA accumulates in DMVs at 6 hpi and in VPs
at 24 hpi. dsRNA clusters occur within the same vesicles but occupy distinct parts of them
compared to vgRNA. Our data suggests a model (Fig. 6) where SARS-CoV-2 RNA is replicated and
transcribed within these DMVs and VPs as highlighted by the proximal localizations of RdRp.

Discussion

Previous biochemical and EM studies allowed researchers to build models of the intracellular life cycle of SARS-CoV-2[24,46,47]; however, precise localization of specific viral proteins and RNA molecules is challenging due to lack of specific contrast in EM and low resolution in DL fluorescence microscopy. SR fluorescence microscopy is well suited for coronavirus studies in cells as it provides both specific contrast and high resolution (~20 nm and below depending upon photons collected[48]). However, to date few studies have employed this method for coronavirus biology[15], with even less focus on SARS-CoV-2[36,44,49], and none of them addressed the SARS-CoV-2 replication process in detail. Here we apply SR fluorescence microscopy to precisely localize the key players of SARS-CoV-2 replication at different time points in infected cells. Building upon a previously developed method for simultaneous labeling of coronavirus vgRNA with dsRNA and protein immunofluorescence[15], and using improved fixation and multi-color SR imaging protocols (see Methods), we obtain and quantify the appearance and molecular compositions of ROs of SARS-CoV-2 in cells at different stages of infection.

Fig. 6: Proposed model for SARS-CoV-2 replication organelles containing various RNA and protein molecules at early and late stages of infection.

In this study, our results taken together depict a compelling and novel picture of ROs containing various molecules including vgRNA, dsRNA, RdRp, nsp3, and ER membrane (Fig. 6). In this model, we compare the organization of ROs at early and late stages of infection and show how specific RNA and protein molecules are spatially organized in ROs. Compared to the simpler and less

pathogenic HCoV-229E case, SARS-CoV-2 appears to generate more complex clusters of vgRNA,

and with the imaging of viral proteins involved in vgRNA replication and in DMV formation, the

structural importance of ROs is now clear.

The detailed intracellular localization of the central SARS-CoV-2 component, vgRNA, has remained

vague in the literature. Our RNA FISH method[15] targets specific sequences in vgRNA (Fig. 1a) and

detects single vgRNA molecules (Fig. 1e, i; Fig. S1, S3a, S17b), allowing counting of the number of

vgRNA molecules within specific regions (Fig. S3b-c). We find for the first time that most cellular

vgRNA localizes into dense clusters of an approximately round shape that grow and migrate to

the perinuclear region as infection time increases. We show that these clusters appear confined

in membranous vesicles derived from ER as emphasized by the localization of Sec61β and

CellMask at their surface (Fig. 4b, Fig. S9-S13). From comparison with earlier EM images[5,22,25] and

from nsp3 localization at their surface[25] (Fig. 5), we can conclude that these vesicles are most

likely DMVs at an early-mid infection time that grow and merge into VPs as infection progresses.

Previously, metabolic radioactive labeling was used to localize newly synthesized RNA in SARS-

CoV-1 and MERS-CoV-infected cells to DMVs[4]. However, metabolic labeling only localizes a

fraction of vgRNA molecules with little sequence specificity and with a background of viral sgRNA.

Here, we specifically label vgRNA of SARS-CoV-2 for SR microscopy and show that it also localizes

in patterns that suggest confinement in DMVs, consistent with the earlier findings on SARS-CoV-

1 and MERS-CoV[4]. Our metabolic labeling of infected cells with BrUTP also localized newly

synthesized viral RNAs to the perinuclear vgRNA clusters (Fig. 3g, Fig. S7), which agrees with

earlier results[4,23,34] and solidifies our conclusions on the spatial localization of vgRNA and viral

replication machinery inside DMVs.

Previous studies also suggested the presence of dsRNA in DMVs of SARS-CoV-1[3] and SARS-CoV-

2[5]. EM images of DMVs often display a complex filamentous network in their interior that was

attributed to viral RNA molecules[5]. However, the exact type of these RNAs was not determined

due to the absence of specific labeling. As one might expect, single-stranded vgRNA can form a

secondary structure that includes many short dsRNA fragments *e.g.*, in stem loops[50,51]. This makes

it difficult to distinguish between viral dsRNA and vgRNA by measuring the diameter of the

filaments, taking into account that the detection probability of ssRNA might be lower due to a

decreased EM contrast for ssRNA than for dsRNA. Reported abundant branching of filaments in

DMVs[5], however, is typical for ssRNA secondary structures[52]. Indeed, these references present

some evidence about the presence of both dsRNA and vgRNA in DMVs; however, to our

knowledge, there was no simultaneous observation of both vgRNA and dsRNA within the same

DMVs.

Here we use the J2 anti-dsRNA antibody that recognizes only long dsRNA fragments (≥ 40bp) with

no detection of the ssRNA secondary structures[53,54]. The J2 antibody has been reported to

underestimate dsRNA localization[26]; however, using optimized antibody concentrations (Fig. S19,

S20) and optimized staining protocols as detailed in Methods, we achieved excellent sensitivity

to dsRNA with signal present in all infected cells, even in early infection with very low vgRNA

levels (Fig. S2d). Our two-color SR imaging revealed for the first time that most dsRNA and vgRNA

are located within the same DMVs and VPs, occupying distinct regions of these vesicles, and

adopting an anti-correlation pattern at short distances (r < 100 nm) at 24 hpi (Fig. 2). Another

novel observation is the relatively constant amount of dsRNA and a slight decrease in dsRNA

cluster size between 6 and 24 hpi despite the huge change in the vgRNA landscape (Fig. 2, Fig.

S2c, e).

It has been proposed that the RdRp complex of SARS-CoV-1 is located at convoluted membranes

and inside DMVs based on immunogold labeling of nsp8[3]. However, nsp8 has intracellular

functions other than as an RdRp accessory subunit[55,56] that might be exercised at the convoluted

membranes. Here we label the catalytic RdRp subunit, nsp12[18], and find that it mostly localizes

to the vgRNA clusters at both 6 and 24 hpi (Fig. 3a-d, i), suggesting that SARS-CoV-2 replication

and transcription occur preferentially in the vgRNA-filled ROs, where dsRNA resides as well.

Additional experiments revealed that two other RdRp subunits, nsp7 and nsp8, as well as newly

synthesized viral RNA also localize to the vgRNA clusters (Fig. 3e-g, i; Fig. S4-S8), further proving

the role of these clusters as replication organelles.

Nsp3 of betacoronaviruses (SARS-CoV-1, MERS-CoV and MHV) was previously localized to the

convoluted membranes and to the DMV membranes using immuno-EM[3,4,57,58] and cryo-ET[25];

however, these studies were limited to early-mid infection at 8-12 hpi. In our study, we report

two localization patterns of nsp3 of SARS-CoV-2 at 6hpi: 1) sparse nsp3 at the surface of isolated

vgRNA-dsRNA clusters (Fig. 5c, Fig. S14a); and 2) dense nsp3 within the accumulations of vgRNA-

dsRNA clusters (Fig. 5d, Fig. S14a). While the first pattern most likely corresponds to the RO/DMV

membranes considering the role of nsp3 as a DMV pore[25], the second one resembles a pattern

found in other coronaviruses that was attributed to the convoluted membranes[3,4,59]. Convoluted

membranes are typically found within dense groups of DMVs in early-mid infection[3,4] and

localization of nsp3 on them might represent early steps of viral transformation of ER into DMVs.

We found this nsp3 pattern anti-correlated with vgRNA (Fig. 5d, e) and with dsRNA (Fig. S14a, b),

suggesting little to no vgRNA or dsRNA at the convoluted membranes, in line with previous studies

on other coronaviruses[4].

At 24 hpi, we did not observe these early infection patterns of nsp3 localization. Instead, we show

for the first time that at 24 hpi, nsp3 densely localizes at the membranes that separate large

vgRNA clusters and grows into a considerable perinuclear network that contains the ROs (Fig. 5g-

i, Fig. S14c). Since the molecular pores of VPs have not yet been investigated in detail, we can

speculate that this late infection nsp3 pattern corresponds to the pores of VPs that should also

be much denser than those of isolated DMVs, considering the increased density of nsp3 labeling.

Additional rare phenotypes of nsp3 localization that we also report for the first time (Fig. S16)

illustrate the variability of SARS-CoV-2 infection course and should lead to further research on the

other intracellular functions of this viral protein.

Taken together, we investigated several key factors of SARS-CoV-2 replication: vgRNA, dsRNA,

RdRp and nsp3 inside infected cells with SR microscopy for the first time. We discovered and

characterized the nanoscale structure of perinuclear clusters of vgRNA and demonstrated by

RdRp labeling that they associate with SARS-CoV-2 ROs. We found that the ROs also contain

dsRNA and are encapsulated in ER-derived membranes. Using SR data on nsp3, we conclude that

these virus-induced organelles correspond to DMVs.

This study expands the knowledge of the biology of coronaviruses and opens new possibilities for

therapeutics against SARS-CoV-2, considering that clusters of vgRNA have also been reported in

SARS-CoV-2 infected interstitial macrophages of human lungs[17], suggesting their importance in

COVID-19. Careful examination of the organization of ROs may provide new avenues to target the

organelles to disrupt SARS-CoV-2 replication and transcription. Examining localization patterns for

different viral variants or in different host cells will be useful to broaden understanding of the viral

infection. It will also be important to examine how the structures reported in this study change

upon the addition of drug treatments. Our imaging approach may also offer insights into long

COVID by investigating cells that are infected by SARS-CoV-2 that may still contain RO-like

structures after symptoms disappear.

Methods

Antibodies

Primary antibodies and the optimal dilutions and concentrations used are as follows: goat polyclonal anti-spike S2 (Novus Biologicals, AF10774-SP, 1:20, 10 µg/mL), mouse monoclonal anti-dsRNA (SCICONS, 10010200, 1:200, 5 µg/mL), rabbit polyclonal anti-RdRp/nsp12 (Sigma-Aldrich, SAB3501287-100UG, 1:500, 2 µg/mL), mouse monoclonal anti-nucleocapsid (Thermo Fisher, MA5-29981, 1:500, 2 µg/mL), rabbit polyclonal anti-nsp3 (Thermo Fisher, PA5-116947, 1:134, 5 µg/mL), sheep polyclonal anti-GFP (Bio-Rad, 4745-1051, 1:1000, 5 µg/mL), rabbit polyclonal anti-GFP (Novus Biologicals, NB600-308SS, 1:163, 5 µg/mL), rabbit monoclonal anti-nsp7 (GeneTex, GTX636719, 1:200, 2 µg/mL), mouse monoclonal anti-nsp8 (GeneTex, GTX632696, 1:134, 5 µg/mL), mouse monoclonal anti-BrdU (Thermo Fisher, B35128, 1:50, 2 µg/mL). Secondary antibodies and the optimal dilutions and concentrations used are as follows: AF647-conjugated donkey anti-mouse IgG (Thermo Fisher, A-31571, 1:500, 4 µg/mL), AF647-conjugated donkey anti-rabbit IgG (Thermo Fisher, A-31573, 1:500, 4 µg/mL), AF647-conjugated donkey anti-sheep IgG (Thermo Fisher, A-21448, 1:500, 4 µg/mL), CF568-conjugated donkey anti-goat IgG (Sigma-Aldrich, SAB4600074-50UL, 1:500, 4 µg/mL), CF568-conjugated donkey anti-rabbit IgG (Sigma-Aldrich, SAB4600076-50UL, 1:500, 4 µg/mL), CF568-conjugated donkey anti-mouse IgG (Sigma-Aldrich, SAB4600075-50UL, 1:500, 4 µg/mL), CF568-conjugated donkey anti-sheep IgG (Sigma-Aldrich, SAB4600078-50UL, 1:500, 4 µg/mL), CF583R-conjugated donkey anti-mouse IgG (Custom CF Dye, lot 23C1122, Biotium, 1:250, 4 µg/mL), CF583R-conjugated donkey anti-rabbit IgG (Custom CF Dye, lot 23C0811, Biotium, 1:250, 4 µg/mL). To confirm that the fluorophore attached to the secondary antibody does not produce artifacts, in a number of cases we switched the labels by switching the secondary antibodies, and found no difference in the SR structures observed.

Culture of cell lines

The Vero E6 cells (African green monkey kidney epithelial cells, ATCC, CRL-1586), HEK293T cells (human embryonic kidney epithelial cells, ATCC, CRL-3216), and Vero E6-TMPRSS2 cells were cultured in Dulbecco's modified Eagle medium (DMEM) with GlutaMAX, 25 mM D-Glucose, and

1 mM sodium pyruvate (Gibco, 10569010) in 10% FBS (Sigma-Aldrich, F0926) at 37°C and 5% CO2

in a humidified incubator. Cell lines were not authenticated after purchase prior to use. For Vero

E6-TMPRSS2, Geneticin (G418) was added at a final concentration of 1mg/ml.

Lentivirus production for ER labeling with Sec61β

To produce lentivirus, HEK293T cells were cultured in 10-cm dishes and transiently transfected

with 9 µg lentiviral plasmid pLV-ER-GFP (Addgene, 80069, a gift from Pantelis Tsoulfas), 8 µg

pCMV-dR8.91, and 1 µg PMD2.G packaging plasmids using 25 µL TransIT-LT1 Transfection Reagent

(Mirus, MIR 2306). After 72 h of transfection, supernatant was filtered through 0.45 µm filters,

concentrated using Lentivirus Precipitation Solution (ALSTEM, VC100) at 4°C overnight, and

centrifuged at 1,500x g for 30 min at 4°C to collect virus pellets. The virus pellets were

resuspended in cold DMEM for storage at -80°C for transduction of cells.

Generation of stable cell line

To generate a Vero E6 cell line stably expressing Sec61β-GFP, 2×10^5 Vero E6 cells were seeded in

one well of a 6-well plate and infected with one quarter of concentrated lentivirus expressing pLV-

ER-GFP produced from one 10-cm dish of HEK293T cells while seeding. After two days incubation,

monoclonal cells expressing GFP were sorted out using a SONY SH800S sorter. These transduced

cells were only used for ER imaging; all other experiments used wild type (WT) cells.

SARS-CoV-2 viral stocks preparation

The SARS-CoV-2 WA 1, isolate USA-WA1/2020 (NR-52281, BEI Resources) was passaged 3 times

in Vero E6-TMPRSS2 cells as previously described[60,61]. Briefly, a Vero E6-TMPRSS2 monolayer was

infected with virus obtained from BEI; post 72 hours of infection (hpi), P1 virus-containing tissue

culture supernatants were collected and stored at -80°C. Following titration, P1 virus stock was

used to generate a P2 stock by infecting Vero E6 TMPRSS2 monolayers with multiplicity of

infection (MOI) of 0.0001 for 72 hours. P2 virus was passaged again in Vero E6-TMPRSS2 cells to

obtain P3 stock. Viral titers were determined by standard plaque assay on Vero E6 cells.

Infection of cells by SARS-CoV-2

Vero E6 cells previously cultured in 8-well μ-slides were infected in the BSL-3 facility with SARS-

CoV-2 WA 1 (USA212 WA1/2020) in triplicates (MOI=0.5 SARS-CoV-2 WA1 (P3)) at an MOI of 2 for

6 hpi and MOI of 0.2 for 24 hpi. After 6 and 24 hrs of incubation, cells were washed with PBS and

fixed by 4% PFA (Electron Microscopy Sciences #15710) and 0.1% glutaraldehyde (Electron

Microscopy Sciences #16350) in PBS for 1 hour and removed from BSL-3 for further processing.

All work involving viral stock preparation and infection using WT SARS-CoV-2 was conducted at

the high containment BSL-3 facility of Stanford University according to CDC and institutional

guidelines. All the experiments were performed using a P3 SARS-CoV-2 USA-WA1/2020,

containing 100% WT population with no deletion in the spike multi-basic cleavage site.

Synthesis of the RNA FISH probes

vgRNA FISH probes targeting the ORF1a region of SARS-CoV-2[26] were ordered with 5AmMC6

modifications from Integrated DNA Technologies, Inc. in plate format of 25 nmol scale with

standard desalting. Each probe was dissolved in water to a final concentration of 100 μM. The

same set of probes was combined with equal volumes of each probe to get a stock of 100 μM

mixed probes. The mixed probes were further desalted using ethanol precipitation. Briefly, 120

μL 100 μM probes were mixed with 12 μL 3 M sodium acetate (pH 5.5), followed by 400 μL

ethanol. After precipitation at -80C overnight, probes were pelleted through centrifugation at

12,000x g for 10 min at 4°C, washed with precooled 70% (vol./vol.) ethanol three times, air dried,

and dissolved in water to make a 100 μM solution of probes. Then, 18 μL 100 μM probes were

mixed with 2 μL 1 M NaHCO₃ (pH 8.5), followed by 100 μg Alexa Fluor™ 647 succinimidyl ester

(NHS) (Invitrogen, A37573) or CF568 succinimidyl ester (NHS) (Biotium, 92131) dissolved in 2 μL

dry DMSO (Invitrogen, D12345). The mixture was incubated for 3 days at 37C in the dark for

conjugation and purified for 3 rounds using Monarch PCR & DNA Cleanup Kit (5 µg) (NEB, T1030S)

following the manufacturer's instructions. The estimated labeling efficiency of probes was

calculated using the following equation:

$$Modification\ ratio = \frac{20}{(A_{base} \times \varepsilon_{dye}) / (A_{dye} \times \varepsilon_{base})}$$

where ε_{dye} is 239,000 cm^{-1}M^{-1}, ε_{base} is 8,919 cm^{-1}M^{-1}, A_{base} is the absorbance of the nucleic acid at

260 nm, and A_{dye} is the absorbance of the dye at 650 nm. For the probes labeled with CF568, ε_{dye}

is 100,000 cm^{-1}M^{-1}, ε_{base} is 8,919 cm^{-1}M^{-1}, A_{base} is the absorbance of the nucleic acid at 260 nm,

and A_{dye} is the absorbance of the dye at 562 nm.

RNA FISH, immunofluorescence (IF), and CellMask staining

Fixed cells from BLS3 as described above were washed twice with a freshly prepared 0.1% NaBH$_4$

solution at room temperature for 5 min, and washed with PBS three times. For staining without

CellMask (Thermo Fisher, C10046), cells were permeabilized in 70% ethanol at 4°C overnight. For

CellMask staining, cells were permeabilized in 0.1% Triton X-100 at room temperature for 30 min.

For RNA FISH staining, permeabilized cells were washed with 200 µL Wash Buffer A [40 µL Stellaris

RNA FISH Wash Buffer A (LGC Biosearch Technologies, SMF-WA1-60), 20 µL deionized formamide,

140 µL H2O] at room temperature for 5 min, and incubated with 110 µL Hybridization Buffer [99

µL Stellaris RNA FISH Hybridization Buffer (LGC Biosearch Technologies, SMF-HB1-10), 11 µL

deionized formamide] containing 1.1 µL 12.5 µM vgRNA FISH probes for 4 hours at 37°C in the

dark. Then cells were washed with Wash Buffer A for 30 min at 37°C in the dark, washed with

Wash Buffer A containing DAPI for 30 min at 37°C in the dark, and stored in Wash Buffer B (LGC

Biosearch Technologies, SMF-WB1-20) for imaging. DAPI was only added to the samples for

confocal imaging and not added to the samples for SR imaging.

For IF staining with antibodies, permeabilized cells were washed with PBS twice, incubated with

3% BSA in PBS at room temperature for 30 min, and incubated with primary antibodies in PBS at

37°C for 1 hour. After incubation with primary antibodies, cells were washed twice with PBST

buffer (0.1% Tween-20 in PBS) at room temperature for 5 min, washed with PBS once, incubated

with secondary antibodies in PBS at room temperature for 30 min, washed with PBST buffer three

times at room temperature for 5 min, and stored in PBS for imaging.

For simultaneous RNA FISH and IF staining, permeabilized cells were washed with 200 μL Wash

Buffer A at room temperature for 5 min, and incubated with 110 μL Hybridization Buffer (99 μL

Stellaris RNA FISH Hybridization Buffer, 11 μL deionized formamide) containing 1.1 μL 12.5 μM

vgRNA FISH probes, 1 U/μL RNase inhibitor (NxGen, F83923-1), and primary antibodies for 4 hours

at 37°C in the dark. Then cells were washed with 2xSSC buffer once, washed with Wash Buffer A

containing secondary antibodies for 30 min at 37°C in the dark, washed with Wash Buffer A for

30 min at 37°C in the dark, washed with Wash Buffer B once, and stored in Wash Buffer B for

imaging. For CellMask staining, several more steps were performed from here. Cells were washed

with PBS once, stained with 1:20k CellMask and 1 U/μL RNase inhibitor in PBS for 20 min at room

temperature in the dark, and washed with PBS three times before imaging.

RNA FISH and IF staining of purified virions

8-well μ-slides (ibidi, 80827-90) were first treated with poly-D-lysine solution (Thermo Fisher,

A3890401) at 4°C overnight. Then in the BSL-3 facility, the poly-D-lysine solution was removed

and 150 μL SARS-CoV-2 WA1 (P3) virus solution of titer 1.82×10^5 PFU/mL was added into one well

of poly-D-lysine-treated 8-well μ-slides for incubation at 4°C for 24 hours to coat the virions onto

the surface of the well. After incubation, the medium containing virions was removed and the

well was washed with PBS twice. Virions on the surface of the well were fixed with 4% PFA in PBS

for 1 hour at room temperature and the sample was removed from BSL-3. The sample was

washed twice with a freshly prepared 0.1% $NaBH_4$ solution at room temperature for 5 min, and

then washed with PBS three times. The fixed virions were permeabilized in 70% ethanol at 4°C

overnight and washed with PBS twice. For the group with Proteinase K digestion, virions were

incubated with 0.2 mg/mL Proteinase K (NEB #P8107S) in 120 μL PBS at 37°C for 30 min and

washed with PBST buffer three times. Virions were washed with Wash Buffer A once and

incubated with 110 μL Hybridization Buffer (99 μL Stellaris RNA FISH Hybridization Buffer, 11 μL

deionized formamide) containing 1.1 µL 12.5 µM vgRNA FISH probes, 1 U/µL RNase inhibitor, and

primary antibodies for 4 hours at 37°C in the dark. Then virions were washed with 2xSSC buffer

once, washed with Wash Buffer A containing secondary antibodies for 30 min at 37°C in the dark,

washed with Wash Buffer A for 30 min at 37°C in the dark, washed with Wash Buffer B once, and

stored in Wash Buffer B for imaging.

Labeling of newly replicated RNA with BrUTP

Vero E6 cells cultured in 8-well µ-slides were infected with SARS-CoV-2 WA 1 (USA212 WA1/2020)

as described above. After 24 hours of incubation, the culture medium was switched to low

glucose DMEM medium (Thermo Fisher, 10567014) supplemented with 20 mM glucosamine for

30 mins to deplete uridine. Both infected and uninfected groups were further treated with 15 µM

actinomycin D (Sigma, A4262) at 37°C for 30 min to inhibit cellular transcription. To transfect cells

with BrUTP, each well of cells was treated with 10 mM BrUTP (Sigma, B7166) and 12 µL

Lipofectamine 2000 (Thermo Fisher, 11668030) for 1 hour at 37°C. Cells were then washed twice

with PBS, followed by 4% PFA and 0.1% glutaraldehyde fixation for 1 hour, and removed from BSL-

3 following BSL-3 SOP of sample removal. The fixed cells were then washed twice with a freshly

prepared 0.1% $NaBH_4$ solution at room temperature for 5 min, washed with PBS three times,

permeabilized in 70% ethanol at 4°C overnight, and washed twice with PBS.

The co-staining of BrU and nsp12 with antibodies follows the IF staining procedure detailed in the

section "RNA FISH, immunofluorescence (IF), and CellMask staining". For the co-staining of BrU

and vgRNA, cells were first incubated with BrU antibody and 1 U/µL RNase inhibitor in PBS at 37°C

for 30 min. Cells were then washed twice with PBST buffer, washed once with PBS, and incubated

with the secondary antibody and 0.5 U/µL RNase inhibitor in PBS at room temperature for 30

min. Cells were then washed with PBST buffer three times, fixed again with 4% PFA and 0.1%

glutaraldehyde in PBS at room temperature for 10 min, and washed with PBS three times. After

that, cells were washed with Wash Buffer A at room temperature for 5 min, incubated with 110

µL Hybridization Buffer containing 1.1 µL 12.5 µM vgRNA FISH probes and 1 U/µL RNase inhibitor

for 4 hours at 37°C in the dark. Then cells were washed twice with Wash Buffer A for 30 min at

37°C in the dark and stored in Wash Buffer B for imaging.

Spinning disk confocal microscopy

Confocal microscopy was performed at the Stanford University Cell Sciences Imaging Core Facility

with a Nikon TiE inverted spinning disk confocal microscope (SDCM) equipped with a

Photometrics Prime 95B camera, a CSU-X1 confocal scanner unit with microlenses, and 405 nm,

488 nm, 561 nm, and 642 nm lasers, using the 60x/1.27 NA PLAN APO IR water immersion

objective. Images were taken using NIS Elements software version 4.60 with Z stacks at 0.3 μm

steps. The camera pixel size of SDCM is 0.183 μm/pixel and the pinhole size is 50 μm. Only one Z

slice is used for all images shown.

Analysis of confocal data

To extract the intensity of vgRNA, dsRNA and RdRp in each infected cell (Fig. S2c, e-h), the

summation projection of each z stack was created by Fiji[62]. The intensity of each target species in

each cell was measured by Fiji, subtracting the background of the same color channel. The

infected cells were characterized manually into three types based on the morphology of vgRNA.

Type 1 shows scattered dot-like localization of vgRNA. Type 3 shows large clustered vgRNA. Type

2 contains features of both type 1 and type 3.

Optimization of antibody concentrations

We optimized the concentration of antibodies in this study by quantifying their signal-to-

background ratio (SBR), where the signal is the brightness of the IF labeling in the cells that

express the given target (virus-infected sample or cells expressing Sec61β-GFP), and the

background is the brightness in the negative control cells (not-infected or WT cells).

To optimize the concentration of primary antibodies against the viral targets, different

concentrations of the primary antibody were applied to stain Vero E6 cells in SARS-CoV-2-infected

and not-infected samples under a constant secondary antibody concentration (Fig. S19). To

optimize the concentration of secondary antibodies, different concentrations of the secondary

antibody were applied to stain Vero E6 cells in infected (virus+) and not-infected (virus−) samples

under a constant primary antibody concentration (Fig. S20). For each cell, a 11 pixel x 11 pixel box

was drawn in the region with brightest signal in the cell and the mean intensity within that region

was measured to represent the intensity of target antibody in that cell. The SBR was calculated,

after subtraction of the dark signal I_{dark}, using the following equation:

$$SBR = \frac{< I_{virus+} - I_{dark} >}{< I_{virus-} - I_{dark} >}$$

To optimize the concentration of the anti-GFP antibodies, different concentrations of primary

antibody were applied to stain Vero E6 Sec61B-GFP cells and WT Vero E6 cells under a constant

secondary antibody concentration (Fig. S19). For each cell, a 11 pixel x 11 pixel box was drawn in

the region with the brightest signal in the cell and the mean intensities of both the GFP and the

antibody signals within that region were measured after subtraction of the dark signals. To

account for the variable expression levels among different cells, the IF signal I_{IF} was normalized

by the GFP signal I_{GFP} within the given region. The SBR was calculated using the following

equation:

$$SBR = \frac{< I_{IF,Sec61B\text{-}GFP}/I_{GFP,Sec61B\text{-}GFP} >}{< I_{IF,WT}/I_{GFP,WT} >}$$

For the primary antibodies against GFP, nsp3, nucleocapsid, nsp12 and for the secondary antibody

for the dsRNA labeling, we chose the antibody concentration that produces the highest SBR as

the optimal concentration. For the primary antibodies against spike S2 and dsRNA and for the

secondary antibody for the spike S2 labeling, we chose the concentration that yields the second

highest SBR because it provides a significantly lower non-specific background with only a minor

decrease of the estimated SBR.

Optical setup for SR microscopy

(d)STORM SR microscopy was performed on a custom-built system (Fig. S14), consisting of a Nikon Diaphot 200 inverted microscope frame with an oil-immersion objective 60x/1.35 NA (Olympus UPLSAPO60XO) and a Si EMCCD camera (Andor iXon Ultra 897). We used 642 nm and 560 nm 1W continuous-wave (CW) lasers (MPB Communications Inc.) for excitation of AF647 or CellMask and CF568 or CF583R, accordingly. For reactivation of fluorophores from the dark state we used a 405 nm 50 mW CW diode laser (Coherent OBIS). All laser beams were expanded and co-aligned in free space and coupled into a square-core multi-mode fiber with a shaker for speckle reduction (Newport F-DS-ASQR200-FC/PC). The output tip of the fiber (200 x 200 μm^2 core size) was imaged with a 10x/0.25 NA objective and magnified to achieve a square illumination region of 47.6 x 47.6 μm^2 with a constant intensity in the sample image plane of the main objective. The fluorescence was split from the excitation light with a multi-band dichroic mirror (ZT405/488/561/640rpcv2, Chroma) and filtered with dichroic filters (ZET635NF, ZET561NF, T690LPxxr, all Chroma). The fluorescence of AF647 and CellMask was additionally filtered with a band-pass filter (ET685/70M, Chroma) and that of CF568 and CF583R with a combination of 561LP and 607/70BP (Semrock, EdgeBasic and BrightLine). The sample image was focused with a tube lens (f = 400 mm) on the EMCCD camera, providing a pixel size of 117 x 117 nm^2 in sample coordinates.

Axial drift was compensated with a custom Focus Lock system[63]. We used an 808 nm fiber-coupled diode laser (Thorlabs S1FC808) whose output fiber tip was conjugated with the back focal plane of the imaging objective, allowing changing the angle of this beam out of the objective by translating the fiber tip (Fig. S14). This inclined beam was partially reflected from the coverslip-water interface and the reflected beam was focused with a cylindrical lens onto a CMOS sensor (UI-3240CP-NIR, IDS Imaging). The 808 nm beam was aligned such that the image of the reflected beam would shift laterally when the axial position of the sample changes. The sample was mounted on two stacked piezo stages (U-780.DOS for coarse and P-545.3C8S for fine movement, both Physik Instrumente). The position of the reflected beam image was recorded when the sample was set at the desired Z position for imaging. During imaging, the Z-position of the fine

stage was directed to move proportionally to the shift of the reflected beam image from the

recorded position, compensating for Z-drift. The Focus Lock control code was programmed in

Matlab (MathWorks, Inc.).

SR imaging procedure

For (d)STORM, the sample chamber was filled with 300 µl of a photoblinking buffer consisting of

200 U/ml glucose oxidase, 1000 U/ml catalase, 10% w/v glucose, 200 mM Tris-HCl pH 8.0, 15 mM

NaCl and 50 mM cysteamine. The buffer was prepared using the following stock solutions[48]: 1)

4 kU/ml glucose oxidase (G2133, Sigma), 20 kU/ml catalase (C1345, Sigma), 25 mM KCl (P217,

Fisher), 4 mM TCEP (646547, Sigma), 50% v/v glycerol (BP229, Fisher) and 22 mM Tris-HCl pH 7.0

(BP1756, Fisher), stored at −20 °C; 2) 1 M cysteamine-HCl (30080, Sigma), stored at −20 °C; 3) 37%

w/v glucose (49139, Sigma) with 56 mM NaCl (S271, Fisher) and 0.74 M Tris-HCl pH 8.0

(J22638.AE, Fisher), stored at +4 °C. For samples with RNA FISH labeling, the buffer was

supplemented with 1 U/µl of an RNase inhibitor (302811, LGC Biosearch Technologies).

The SR imaging started with a DL image of cells from each fluorophore at a low power (*e.g.*, 2

W/cm^2). For (d)STORM acquisitions, we began with AF647 or CellMask, followed by CF568 or

CF583R[64]. We used an excitation power density of ~20 kW/cm^2 for shelving and blinking of CF568,

~13 kW/cm^2 for CF583R and 6-20 kW/cm^2 for AF647. The power density of the 405 nm

illumination for both dyes was increased from 0 to 50 W/cm^2 throughout an acquisition to keep

the reactivation rate approximately constant. The exposure time was 10.57 ms per frame and the

calibrated EM gain was either 43 or 84. The image recording started after the initial shelving phase

upon observation of clear SM blinking; the blinking movies were acquired for approximately

$6 \cdot 10^4 - 8 \cdot 10^4$ frames for each fluorophore.

SR data analysis

SM movies were processed with the ThunderStorm plugin[65] for Fiji. First, the images were filtered

with a wavelet filter with a b-spline order of 3 and a scale of 2. The coarse localizations were

found as local maxima with an 8-neighborhood connectivity and a threshold of 2·std(Wave.F1).

These localizations were weighted least squares-fitted with the integrated Gaussian model using

a radius of 4 pixels and an initial sigma of 1.1. Then, we performed drift correction estimated by

cross-correlation between successive subsets of localizations in ThunderStorm, or in SharpViSu[66]

when the drift correction in ThunderStorm was unsuccessful. For further processing, we kept only

localizations with fitted sigma between 160 nm and 80 nm. This choice effectively rejects

molecules away from the focal plane, providing an approximate axial sectioning of the images to

roughly 500 nm[67].

For image registration, we imaged 200 nm TetraSpeck beads (T7280, Thermo Fisher Scientific) in

both channels, whose images were processed similarly to the SM movies. The transformation

between the channels was calculated using an affine transformation with help of Matlab function

'fitgeotrans'. The calculated transformation was then applied to the CF568 or CF583R localizations

using a Matlab function 'transformPointsInverse'.

Localizations found within 50 nm on consecutive frames that could originate from multiple

localizations of a single molecule were treated in two ways. For SR images, to improve the

resolution, these localizations were refined by selecting them from a normal distribution with a

mean at the weighted mean of the initial localizations and a standard deviation (SD) that equals

$120 \cdot (N_{ph})^{-1/2}$ nm, where N_{ph} is the total number of photons acquired from all localizations in the

given consecutive series[48]. For data analysis other than SR image reconstruction, to suppress

overcounting, the localizations of the consecutive series were reduced to a single localization at

the weighted mean position. The weights of localizations were proportional to the photon counts

of these individual localizations. After this correction, the SR data of antibody-detected Spike, N,

nsp12, BrU, nsp8, nsp7 was additionally filtered by removing localizations that had 3 or less

neighbors within 30 nm. SR images were reconstucted as 2D histograms with a bin size of 20 x

20 nm^2. However, SR images where one of the channels contained the CellMask labeling had a

bin size of $30 \times 30 \text{ nm}^2$. SR images acquired with CellMask were additionally filtered with a

Gaussian filter with σ = 0.5 pixels.

Cluster analysis with BIC-GMM

Gaussian Mixture Models (GMM) implemented in Python were fitted to vgRNA and dsRNA

localization datasets, yielding a representation of localization densities as a collection of

potentially elliptical and/or rotated 2D Gaussians. The number of components most suitable for

each field of view was determined using an iterative grid search, evaluating 4 candidate GMMs

using the Bayesian Information Criterion (BIC)[68]. The first grid iteration tested [1, 2500]

components with test points $t_i = \{ 1, 834, 1667, 2500 \}$, where i denotes the index in the set such

that $t_0 = 1$. For each iteration of the grid search, the model with the lowest BIC (corresponding to

the best candidate), t_k was selected, and the next iteration of the grid was narrowed, to be

bounded by $[t_{max\ (k-1,\ 0)} + 1, t_{min\ (k+1,\ 3)} - 1]$, until the stride of the grid was 1 component, or the test

point with the best BIC was on a rail ($k = 0$ or 3). To reduce memory requirements, this GMM

optimization was performed on a random subset of up to 200,000 localizations from each data

set, but the optimized GMM was then used to predict a component assignment for all original

localizations. These components were regarded as clusters, and refined by removing localizations

with a log probability of being an event from their assigned Gaussian component of less than −25.

The radius of gyration, Rg, was then calculated for each cluster, and the number of localizations

in each cluster, N_{loc}, was used to approximate a cluster density as $\delta = N_{loc} / (\pi \cdot Rg^2)$. Clusters with

δ below a threshold of 0.008 localizations/nm^2 for dsDNA, or below an ROI-dependent threshold

between 0.005 and 0.013 localizations/nm^2 for vgRNA, were removed from further quantification

as sparse background. This analysis and resulting visualizations were carried out in the PYthon

Microscopy Environment (https://doi.org/10.5281/zenodo.4289803)[69], using a plugin

(github.com/barentine/bic-gmm) and the scikit-learn GMM implementation[70].

Counting of vgRNA molecules in the clusters

The number of vgRNA molecules in a vgRNA cluster was defined as a quotient between the

number of vgRNA-FISH localizations in the cluster and the average number of localizations

produced by a single FISH-labeled vgRNA molecule in the given cell. The average number of

localizations per vgRNA molecule was estimated from isolated nanoscale vgRNA puncta in the

cytoplasm (Fig. 3a). This number was defined as the median of the number of localizations within

50 nm from each localization in the region with vgRNA puncta. The estimated number of vgRNA

molecules was calculated for every cluster determined by the BIC-GMM cluster analysis and the

median value per cell was shown in a chart (Fig. 3b-c).

Counting of nsp12 puncta in the vgRNA clusters

The center of nsp12 puncta is obtained by fitting the SR images in ThunderStorm[65]. The SR

localizations of nsp12 were first converted into a 2D histogram image with a bin size of

20 x 20 nm^2. The approximate localization of the center was found as a centroid of connected

components with a threshold of 5·std(Wave.F1) without filter. These localizations were least

squares-fitted with the integrated Gaussian model using a fitting radius of 2 pixels and an initial

sigma of 0.4. We next removed duplicates among localizations within a 20 nm radius. The puncta

whose sigma were smaller than 5 nm were further filtered out to avoid localizing single-pixel-

sized background localizations. For each vgRNA cluster with its center and the radius of gyration

(Rg) determined using BIC-GMM, we counted the number of nsp12 puncta within a 1.5·Rg

distance of the center of the vgRNA cluster. For nsp12 puncta found within the cutoff distance of

more than one vgRNA cluster, we assigned them to their closest cluster based on the relative

distance d/Rg, with d being the distance between the center of the vgRNA cluster and center of

the nsp12 punctum.

Bivariate pair-correlation functions

For calculation of bivariate pair-correlation functions[29] $g_{12}(r)$, we first manually selected the

cytoplasmic regions with dense vgRNA clusters. The pair-correlation functions were calculated by

counting the number of localizations of the second species within a distance between r and r+dr

from each localization of the first species. These were normalized by dividing the number of

localizations by the area of the corresponding ring of radii r and r+dr and by the average density

of the second species in the region. Finally, the obtained numbers were averaged across the

localizations of the first species. r was scanned over the range between 0 and 500 nm and dr was

set to 1 nm. For the complete spatial randomness (CSR) case, a test CSR dataset was generated

with the same average density as for the experimental case across the same ROI. $g_{12}(r)$ traces

were calculated from these CSR datasets as described above. No edge effect correction was

performed leading to a slight decrease of $g_{12}(r)$ at large r. Plots in the figures display experimental

and CSR $g_{12}(r)$ for each analyzed cell as faint lines as well as the mean $g_{12}(r)$ calculated from all

cells in bold lines.

Estimation of RNA FISH labeling efficiency in virions

Dye molecules inside virions were counted using fluorescence bleaching with SM calibration.

Virions attached to the coverslip were labeled using the RNA-FISH+IF protocol with PFA-only

fixation. The density of virions was around 0.5 μm^{-2} ensuring observation of most virions as single

DL spots without overlap (Fig. S1a, d). vgRNA was FISH-labeled with AF647 and spike protein was

IF-stained with CF568. Glass-bottom chambers with virions were kept in PBS for this experiment.

Samples were illuminated with 642 nm light at 20 W/cm^2 and were imaged with an exposure time

of 200 ms and an EM gain of 43 until bleaching of all AF647 in the imaging region (around 200 s).

A separate DL image of spike was taken with 560 nm excitation. The AF647 bleaching movies were

processed in ThunderStorm using a wavelet filter with a b-spline order of 3 and a scale of 2, a

local maximum approximate localization with a threshold of 1.2·std(Wave.F1) and an 8-

neighborhood connectivity. These localizations were weighted least squares-fitted with the

integrated Gaussian model using a radius of 3 pixels and an initial sigma of 1.1. Then, we kept

only localizations with sigma < 160 nm & sigma > 80 nm and removed duplicates within 300 nm

on each frame.

Further processing was done in Matlab with a custom script. We considered only vgRNA-AF647

localizations that had a spike-CF568 signal within 200 nm to avoid counting AF647 molecules

outside virions. The bleaching time traces (Fig. S1c, f) were found by searching in consecutive

frames within 200 nm of the localization from the first frame and allowing up to 5 empty frames

between frames with detections. The number of bleaching steps was defined as the rounded

quotient between the initial and the final brightness of a spot in a time trace serving as the SM

calibration. For each bleaching trace, the initial brightness (in photons) was defined as the median

value of the brightness in the first 4 localizations and the final brightness as the median brightness

value of the last 4 localizations. If the trace contained only 7-8 detections, the range for the initial

and the final brightness was reduced to 3 frames; for traces with 5-6 detections, this was reduced

to 2; for traces with 3-4 frames – to 1; for traces containing only 1 or 2 detections, the number of

bleaching steps was set to 1. For each analyzed region containing around 200 bleaching traces,

the number of bleaching steps was fitted with a zero-truncated Poisson distribution (Fig. S1g-h).

The expected values ± SD obtained from the fit of 5 regions for each of not-treated and PK-treated

cells are shown in a chart (Fig. S1i).

Data availability

923 Source data for analysis figures are provided with this paper. The code utilized in this study is

924 available at Stanford Digital Repository at https://doi.org/10.25740/td954gx5320. All data

925 generated in this study are available from the Corresponding Authors upon reasonable request.

Author contribution

L.A., M.H., L.S.Q. and W.E.M. conceived the project. L.A. designed the optical set-up, performed

the SR acquisitions and data analysis. M.H. performed cell culture, labeling and confocal imaging.

Y.Z. performed confocal and SR data analysis and helped with sample preparation and confocal

imaging. J.G. and P.P. performed SARS-CoV-2 infection experiments at the BSL-3 facility with staff

listed in the Acknowledgements. A.R.R. contributed to the concept and SR experiments at the

early stages of the project. A.E.S.B. designed the BIC-GMM cluster analysis method and

contributed to the optical set-up design. A.B. performed portions of the BrU SR imaging. L.A. and

W.E.M. wrote the manuscript with input from all authors.

Acknowledgements

We thank Amol Pohane for assistance with the cell culturing in the BSL-3 facility. We thank Leiping

Zeng for sample preparation and discussion about the results and experimental plan with the

other authors. This work was supported in part by the National Institute of General Medical

Sciences Grant Nos. R35GM118067 (to W.E.M.) and the National Institutes of Health Common

Fund 4D Nucleome Program No. U01 DK127405 (to L.S.Q.). We also acknowledge Stanford

University Cell Sciences Imaging Core Facility (RRID:SCR_017787). L.S.Q. is a Chan Zuckerberg

Biohub – San Francisco Investigator, and W.E.M. is a Sarafan ChEM-H Fellow.

An earlier version of this research was published on bioRxiv on November 8, 2023

https://doi.org/10.1101/2023.11.07.566110

References

1.   Schmidt N*, et al.* The SARS-CoV-2 RNA–protein interactome in infected human cells.
Nature Microbiology **6**, 339-353 (2021).

2.   Flynn RA*, et al.* Discovery and functional interrogation of SARS-CoV-2 RNA-host protein
interactions. *Cell* **184**, 2394-2411.e2316 (2021).

3.   Knoops K*, et al.* SARS-coronavirus replication is supported by a reticulovesicular network
of modified endoplasmic reticulum. *PLoS Biol* **6**, e226 (2008).

4.   Snijder EJ*, et al.* A unifying structural and functional model of the coronavirus replication
organelle: Tracking down RNA synthesis. *PLoS Biol* **18**, e3000715 (2020).

5.   Klein S*, et al.* SARS-CoV-2 structure and replication characterized by in situ cryo-electron
tomography. *Nat Commun* **11**, 5885 (2020).

6. Betzig E*, et al.* Imaging intracellular fluorescent proteins at nanometer resolution. *Science*
967   **313**, 1642-1645 (2006).

969 7. Rust MJ, Bates M, Zhuang X. Sub-diffraction-limit imaging by stochastic optical
970   reconstruction microscopy (STORM). *Nat Methods* **3**, 793-796 (2006).

972 8. Heilemann M*, et al.* Subdiffraction-Resolution Fluorescence Imaging with Conventional
973   Fluorescent Probes. *Angew Chem Int Ed* **47**, 6172-6176 (2008).

975 9. Klar TA, Jakobs S, Dyba M, Egner A, Hell SW. Fluorescence microscopy with diffraction
976   resolution barrier broken by stimulated emission. *Proceedings of the National Academy*
977   *of Sciences of the United States of America* **97**, 8206-8210 (2000).

979 10. Gustafsson MGL. Surpassing the lateral resolution limit by a factor of two using structured
980   illumination microscopy. *J Microsc* **198**, 82-87 (2000).

982 11. Xu K, Zhong G, Zhuang X. Actin, Spectrin, and Associated Proteins Form a Periodic
983   Cytoskeletal Structure in Axons. *Science* **339**, 452-456 (2013).

985 12. Andronov L, Ouararhni K, Stoll I, Klaholz BP, Hamiche A. CENP-A nucleosome clusters form
986   rosette-like structures around HJURP during G1. *Nature Communications* **10**, 4436 (2019).

988 13. Reinhardt SCM*, et al.* Ångström-resolution fluorescence microscopy. *Nature* **617**, 711-716
989   (2023).

991 14. Baddeley D, Bewersdorf J. Biological Insight from Super-Resolution Microscopy: What We
992   Can Learn from Localization-Based Images. *Annual Review of Biochemistry* **87**, 965-989
993   (2018).

995 15. Wang J*, et al.* Multi-color super-resolution imaging to study human coronavirus RNA
996   during cellular infection. *Cell Reports Methods* **2**, 100170 (2022).

998 16. Jackson CB, Farzan M, Chen B, Choe H. Mechanisms of SARS-CoV-2 entry into cells. *Nature*
999   *Reviews Molecular Cell Biology* **23**, 3-20 (2022).

1000

1001 17. Timothy Ting-Hsuan W*, et al.* Activated interstitial macrophages are a predominant target
1002   of viral takeover and focus of inflammation in COVID-19 initiation in human lung. *bioRxiv*,
1003   2022.2005.2010.491266 (2023).

18.   Hillen HS, Kokic G, Farnung L, Dienemann C, Tegunov D, Cramer P. Structure of replicating SARS-CoV-2 polymerase. *Nature* **584**, 154-156 (2020).

19.   Park GJ*, et al.* The mechanism of RNA capping by SARS-CoV-2. *Nature* **609**, 793-800 (2022).

20.   Decroly E, Ferron F, Lescar J, Canard B. Conventional and unconventional mechanisms for capping viral mRNA. *Nature Reviews Microbiology* **10**, 51-65 (2012).

21.   Zimmermann L*, et al.* SARS-CoV-2 nsp3 and nsp4 are minimal constituents of a pore spanning replication organelle. *Nature Communications* **14**, 7894 (2023).

22.   Eymieux S*, et al.* Ultrastructural modifications induced by SARS-CoV-2 in Vero cells: a kinetic analysis of viral factory formation, viral particle morphogenesis and virion release. *Cellular and Molecular Life Sciences* **78**, 3565-3576 (2021).

23.   Doyle N, Simpson J, Hawes PC, Maier HJ. Coronavirus RNA Synthesis Takes Place within Membrane-Bound Sites. *Viruses* **13**, 2540 (2021).

24.   V'Kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. Coronavirus biology and replication: implications for SARS-CoV-2. *Nat Rev Microbiol* **19**, 155-170 (2021).

25.   Wolff G*, et al.* A molecular pore spans the double membrane of the coronavirus replication organelle. *Science* **369**, 1395-1398 (2020).

26.   Lee JY*, et al.* Absolute quantitation of individual SARS-CoV-2 RNA molecules provides a new paradigm for infection dynamics and variant differences. *eLife* **11**, e74153 (2022).

27.   Pepe A, Pietropaoli S, Vos M, Barba-Spaeth G, Zurzolo C. Tunneling nanotubes provide a route for SARS-CoV-2 spreading. *Science Advances* **8**, eabo0171 (2022).

28.   Sherman E*, et al.* Functional Nanoscale Organization of Signaling Molecules Downstream of the T Cell Antigen Receptor. *Immunity* **35**, 705-720 (2011).

29.   Razvag Y, Neve-Oz Y, Sajman J, Reches M, Sherman E. Nanoscale kinetic segregation of TCR and CD45 in engaged microvilli facilitates early T cell activation. *Nature Communications* **9**, 732 (2018).

1041

1042   30.   Sawicki SG, Sawicki DL. Coronavirus minus-strand RNA synthesis and effect of
1043   cycloheximide on coronavirus RNA synthesis. *Journal of Virology* **57**, 328-334 (1986).

1044

1045   31.   Gao Y*, et al.* Structure of the RNA-dependent RNA polymerase from COVID-19 virus.
1046   *Science* **368**, 779-782 (2020).

1047

1048   32.   Ahn D-G, Choi J-K, Taylor DR, Oh J-W. Biochemical characterization of a recombinant SARS
1049   coronavirus nsp12 RNA-dependent RNA polymerase capable of copying viral RNA
1050   templates. *Archives of Virology* **157**, 2095-2104 (2012).

1051

1052   33.   Naydenova K*, et al.* Structure of the SARS-CoV-2 RNA-dependent RNA polymerase in the
1053   presence of favipiravir-RTP. *Proceedings of the National Academy of Sciences* **118**,
1054   e2021946118 (2021).

1055

1056   34.   Gosert R, Kanjanahaluethai A, Egger D, Bienz K, Baker Susan C. RNA Replication of Mouse
1057   Hepatitis Virus Takes Place at Double-Membrane Vesicles. *Journal of Virology* **76**, 3697-
1058   3708 (2002).

1059

1060   35.   Heinrich BS, Cureton DK, Rahmeh AA, Whelan SPJ. Protein Expression Redirects Vesicular
1061   Stomatitis Virus RNA Synthesis to Cytoplasmic Inclusions. *PLOS Pathogens* **6**, e1000958
1062   (2010).

1063

1064   36.   Cortese M*, et al.* Integrative Imaging Reveals SARS-CoV-2-Induced Reshaping of
1065   Subcellular Morphologies. *Cell Host & Microbe* **28**, 853-866.e855 (2020).

1066

1067   37.   Lang S*, et al.* An Update on Sec61 Channel Functions, Mechanisms, and Related Diseases.
1068   *Front Physiol* **8**, 887 (2017).

1069

1070   38.   Hsu JC-C, Laurent-Rolle M, Pawlak JB, Wilen CB, Cresswell P. Translational shutdown and
1071   evasion of the innate immune response by SARS-CoV-2 NSP14 protein. *Proceedings of the
1072   National Academy of Sciences* **118**, e2101161118 (2021).

1073

1074   39.   Mönkemöller V, Schüttpelz M, McCourt P, Sørensen K, Smedsrød B, Huser T. Imaging
1075   fenestrations in liver sinusoidal endothelial cells by optical localization microscopy.
1076   *Physical Chemistry Chemical Physics* **16**, 12576-12581 (2014).

1077

1078 40. Oudshoorn D*, et al.* Expression and Cleavage of Middle East Respiratory Syndrome
1079   Coronavirus nsp3-4 Polyprotein Induce the Formation of Double-Membrane Vesicles That
1080   Mimic Those Associated with Coronaviral RNA Replication. *mBio* **8**, 10.1128/mbio.01658-
1081   01617 (2017).

1082

1083 41. Han Y-S*, et al.* Papain-Like Protease 2 (PLP2) from Severe Acute Respiratory Syndrome
1084   Coronavirus (SARS-CoV): Expression, Purification, Characterization, and Inhibition.
1085   *Biochemistry* **44**, 10349-10359 (2005).

1086

1087 42. Shin D*, et al.* Papain-like protease regulates SARS-CoV-2 viral spread and innate immunity.
1088   *Nature* **587**, 657-662 (2020).

1089

1090 43. Bai Z, Cao Y, Liu W, Li J. The SARS-CoV-2 Nucleocapsid Protein and Its Role in Viral Structure,
1091   Biological Functions, and a Potential Target for Drug or Vaccine Mitigation (2021).

1092

1093 44. Scherer KM*, et al.* SARS-CoV-2 nucleocapsid protein adheres to replication organelles
1094   before viral assembly at the Golgi/ERGIC and lysosome-mediated egress. *Science*
1095   *Advances* **8**, eabl4895 (2022).

1096

1097 45. Xia X. Domains and Functions of Spike Protein in SARS-Cov-2 in the Context of Vaccine
1098   Design (2021).

1099

1100 46. Hartenian E, Nandakumar D, Lari A, Ly M, Tucker JM, Glaunsinger BA. The molecular
1101   virology of coronaviruses. *J Biol Chem* **295**, 12910-12934 (2020).

1102

1103 47. Baggen J, Vanstreels E, Jansen S, Daelemans D. Cellular host factors for SARS-CoV-2
1104   infection. *Nature Microbiology* **6**, 1219-1232 (2021).

1105

1106 48. Andronov L, Genthial R, Hentsch D, Klaholz BP. splitSMLM, a spectral demixing method for
1107   high-precision multi-color localization microscopy applied to nuclear pore complexes.
1108   *Communications Biology* **5**, 1100 (2022).

1109

1110 49. Storti B*, et al.* A spatial multi-scale fluorescence microscopy toolbox discloses entry
1111   checkpoints of SARS-CoV-2 variants in Vero E6 cells. *Computational and Structural*
1112   *Biotechnology Journal* **19**, 6140-6156 (2021).

1113

1114 50. Cao C*, et al.* The architecture of the SARS-CoV-2 RNA genome inside virion. *Nature*
1115   *Communications* **12**, 3917 (2021).

1116

1117   51.   Lan TCT*, et al.* Secondary structural ensembles of the SARS-CoV-2 RNA genome in infected
1118   cells. *Nature Communications* **13**, 1128 (2022).

1119

1120   52.   Gopal A, Zhou ZH, Knobler CM, Gelbart WM. Visualizing large RNA molecules in solution.
1121   *RNA* **18**, 284-299 (2012).

1122

1123   53.   Schonborn J, Oberstrass J, Breyel E, Tittgen J, Schumacher J, Lukacs N. Monoclonal
1124   antibodies to double-stranded RNA as probes of RNA structure in crude nucleic acid
1125   extracts. *Nucleic Acids Res* **19**, 2993-3000 (1991).

1126

1127   54.   Ku J*, et al.* Reactive Polymer Targeting dsRNA as Universal Virus Detection Platform with
1128   Enhanced Sensitivity. *Biomacromolecules* **21**, 2440-2454 (2020).

1129

1130   55.   Deng J*, et al.* SARS-CoV-2 NSP8 suppresses type I and III IFN responses by modulating the
1131   RIG-I/MDA5, TRIF, and STING signaling pathways. *Journal of Medical Virology* **95**, e28680
1132   (2023).

1133

1134   56.   Kumar P*, et al.* The nonstructural protein 8 (nsp8) of the SARS coronavirus interacts with
1135   its ORF6 accessory protein. *Virology* **366**, 293-303 (2007).

1136

1137   57.   Snijder EJ*, et al.* Ultrastructure and origin of membrane vesicles associated with the severe
1138   acute respiratory syndrome coronavirus replication complex. *J Virol* **80**, 5927-5940 (2006).

1139

1140   58.   Stertz S*, et al.* The intracellular sites of early replication and budding of SARS-coronavirus.
1141   *Virology* **361**, 304-315 (2007).

1142

1143   59.   Ulasli M, Verheije MH, de Haan CAM, Reggiori F. Qualitative and quantitative
1144   ultrastructural analysis of the membrane rearrangements induced by coronavirus. *Cellular*
1145   *Microbiology* **12**, 844-861 (2010).

1146

1147   60.   Chiem K, Ye C, Martinez-Sobrido L. Generation of Recombinant SARS-CoV-2 Using a
1148   Bacterial Artificial Chromosome. *Current Protocols in Microbiology* **59**, e126 (2020).

1149

1150   61.   Ye C*, et al.* Rescue of SARS-CoV-2 from a Single Bacterial Artificial Chromosome. *mBio* **11**,
1151   10.1128/mbio.02168-02120 (2020).

1152

1153   62.   Schindelin J*, et al.* Fiji: an open-source platform for biological-image analysis. *Nature*
1154   *methods* **9**, 676-682 (2012).

1155

1156   63.   Barentine AES*, et al.* An integrated platform for high-throughput nanoscopy. *Nature*
1157   *Biotechnology*, (2023).

1158

1159   64.   Wang B, Xiong M, Susanto J, Li X, Leung W-Y, Xu K. Transforming Rhodamine Dyes for
1160   (d)STORM Super-Resolution Microscopy via 1,3-Disubstituted Imidazolium Substitution.
1161   *Angewandte Chemie International Edition* **61**, e202113612 (2022).

1162

1163   65.   Ovesny M, Krizek P, Borkovec J, Svindrych Z, Hagen GM. ThunderSTORM: a comprehensive
1164   ImageJ plug-in for PALM and STORM data analysis and super-resolution imaging.
1165   *Bioinformatics (Oxford, England)* **30**, 2389-2390 (2014).

1166

1167   66.   Andronov L, Lutz Y, Vonesch J-L, Klaholz BP. SharpViSu: integrated analysis and
1168   segmentation of super-resolution microscopy data. *Bioinformatics* **32**, 2239-2241 (2016).

1169

1170   67.   Möckl L*, et al.* Quantitative Super-Resolution Microscopy of the Mammalian Glycocalyx.
1171   *Developmental Cell* **50**, 57-72 (2019).

1172

1173   68.   Schwarz G. Estimating the dimension of a model. *The annals of statistics* **6**, 461-464
1174   (1978).

1175

1176   69.   Marin Z*, et al.* PYMEVisualize: an open-source tool for exploring 3D super-resolution data.
1177   *Nature Methods* **18**, 582-584 (2021).

1178

1179   70.   Pedregosa F*, et al.* Scikit-learn: Machine Learning in Python. *J Mach Learn Res* **12**, 2825–
1180   2830 (2011).

1181

1182

1183

1184

1185

1186
Supplementary figures

1187

1188

Fig. S1. Validation of the labeling and imaging approach.

1189

a, DL image of SARS-CoV-2 virions where vgRNA was labeled with AF647 by RNA FISH and the spike

proteins were labeled by primary anti-spike S2 antibody with secondary CF568-conjugated antibody. **b**,

Representative two-color SR images of individual virions reveal concentric localization of spike around

vgRNA. **c**, Bleaching time trace of AF647 emission from a single virion (yellow arrow in **a**) demonstrates

two-step bleaching. **d**, DL image of virions that were treated with Proteinase K (PK) before labeling. **e**, SR

images of PK-treated virions reveal incomplete spike labeling due to digestion of proteins by the PK. **f**,

Bleaching time trace of AF647 emission from a single virion (yellow arrow in **d**) shows 6-step bleaching

suggesting increased vgRNA labeling efficiency in PK-treated virions. **g**-**h**, Histograms of the number of

fluorophores per virion in untreated (**g**) or PK-treated (**h**) samples and their fits with a Poisson distribution.

i, Mean number of AF647 molecules per virion from the fit for 5 different regions in both untreated and

PK-treated samples. p-value = $2 \cdot 10^{-8}$, two-tailed t-test. The error bars indicate mean ± SD value for the

untreated and PK-treated groups. **j**, SR image of a SARS-CoV-2 infected cell with the cell body to the left

reveals assembled virions at its cytoplasmic tubular projections at 24 hpi. Scale bars, 100 nm (**b**, **e**) and 1

μm (**a**, **d**, **j**).

1204

Fig. S3. Estimation of the number of vgRNA molecules in vgRNA clusters.

a, SR localizations of single vgRNA molecules found in the cytoplasm of infected cells outside the dense vgRNA clusters. On a cell-by-cell basis, similar images are used as a calibration for the number of SR detections per one vgRNA molecule. Examples of SR images of single vgRNA molecules are indicated with white circles (r = 50 nm). **b**, Estimated number of vgRNA molecules per cluster at 6 and 24 hpi from all analyzed cells. The histogram counts are normalized by the number of analyzed cells; the histogram counts for 24 hpi were additionally divided by 3 to account for the 3x wider bin size than at 6 hpi. **c**, Median estimated counts of vgRNA molecules per cluster for each analyzed cell (individual yellow points). The error bars represent mean ± SD values of these median vgRNA molecule counts for each time point. P-value = 5·10⁻⁴, two-tailed t-test. Scale bars, 50 x 50 nm².

1229

1230

Fig. S4. Association of nsp7 with perinuclear vgRNA structures.

Representative SR image of a SARS-CoV-2 infected cell at 24 hpi labeled for vgRNA (magenta) and nsp7 (green) with magnified regions shown in the colored boxes. Scale bars, 2 μm and 500 nm (bottom panels).

1235

1236

1237

1238 **Fig. S5. Association of nsp8 with perinuclear vgRNA structures.**

1239 Representative SR image of a SARS-CoV-2 infected cell at 24 hpi labeled for vgRNA (magenta) and

1240 nsp8 (green) with magnified regions shown in the colored boxes. Scale bars, 2 µm and 500 nm

1241 (bottom panels).

1242

1243

Fig. S6. Colocalization of nsp12 with nsp8.

a, Representative confocal images of cells co-labeled for nsp8 and nsp12 demonstrate their DL colocalization in the perinuclear region of infected cells (Virus+, 24 hpi) and low background immunofluorescence signal in non-infected cells (Virus−). **b**. Representative SR image of an infected cell at 24 hpi reveals punctate localization of both nsp12 and nsp8 in the perinuclear region. (bottom panels) Magnified images of the regions in the colored boxes reveal nanoscale colocalization of nsp12 with nsp8. **c**. Bivariate pair-correlation functions calculated in the perinuclear regions of infected cells demonstrate colocalization of nsp12 and nsp8 at r < 100 nm. Scale bars, 10 μm (**a**), 2 μm (**b**) and 500 nm (bottom panels).

Fig. S7. Association of newly synthesized viral RNAs with perinuclear clusters of vgRNA.

a, Representative confocal images of cells co-labeled for vgRNA and BrU demonstrate their DL colocalization in the perinuclear region of infected cells treated with BrUTP for 1 h before fixation (Virus+ BrUTP+); low background BrU signal in infected cells not treated with BrUTP (Virus+ BrUTP−) and low background signal of both targets in non-infected cells treated with BrUTP for 1 h (Virus− BrUTP+). Endogenous transcription was inhibited with Actinomycin D in all conditions (ActD+). Virus+ cells were fixed at 24 hpi. **b**. Representative SR image of an infected cell at 24 hpi treated with BrUTP and Actinomycin D demonstrates association of BrU labeling with vgRNA clusters. Scale bars, 10 μm (**a**), 2 μm (**b**).

1263

Fig. S8. Association of newly synthesized viral RNAs with nsp12.

a, Representative confocal images of cells co-labeled for nsp12 and BrU demonstrate their DL colocalization in the perinuclear region of infected cells treated with BrUTP for 1 h (Virus+ BrUTP+); low background BrU signal in infected cells not treated with BrUTP (Virus+ BrUTP−); and low background signal of both targets in non-infected cells treated with BrUTP for 1 h (Virus− BrUTP+). Endogenous transcription was inhibited with Actinomycin D in all conditions (ActD+). Virus+ cells were fixed at 24 hpi. **b**. SR image of an infected cell (type 1, early infection) treated with BrUTP demonstrates association of BrU labeling with nsp12. **c**. Bivariate pair-correlation functions calculated in the perinuclear regions of infected and BrUTP-treated cells reveal nanoscale association of nsp12 and BrU. Scale bars, 10 μm (**a**), 2 μm (**b**) and 500 nm (bottom zoomed-in panels).

1274

1275

1276

1277 **Fig. S9. Alterations of host cell ER at 6 hpi.**

1278 SR image of vgRNA in a SARS-CoV-2 infected Vero E6 cell, stably expressing Sec61β-GFP. Altered

1279 ER forms ring-like structures that partially encapsulate vgRNA clusters in the perinuclear region.

1280 Left: green (Sec61β) / magenta (vgRNA) coloring; right: color scale of Sec61β localizations. Scale

1281 bar, 2 µm.

1282

1283

1284

1285

1286 **Fig. S10. Alterations of host cell ER at 24 hpi.**

1287 SR image of vgRNA in a SARS-CoV-2 infected Vero E6 cell, stably expressing Sec61β-GFP. Altered

1288 ER forms ring-like structures that encapsulate vgRNA clusters in the perinuclear region, while the

1289 Sec61β signal at the ER tubules decreases compared to 6 hpi (Fig. S9). Left: green (Sec61β) /

1290 magenta (vgRNA) coloring; right: color scale of Sec61β localizations. Scale bar, 2 µm.

1291

1292

1293

1294

1295 **Fig. S11. Encapsulation of dsRNA by altered host ER at 24 hpi.**

1296 SR image of dsRNA in a SARS-CoV-2 infected Vero E6 cell, stably expressing Sec61β-GFP. Ring-like

1297 structures of altered ER encapsulate dsRNA clusters in the perinuclear region. Left: green (dsRNA)

1298 / magenta (Sec61β) coloring; right: color scale of Sec61β localizations. Scale bar, 2 μm.

1299

1300

1301

Fig. S12. Encapsulation of dsRNA into membrane-bound organelles.

a, SR image of dsRNA and membranes in a SARS-CoV-2 infected cell at 24 hpi with membranes labeled by CellMask Deep Red (magenta) and dsRNA labeled with immunofluorescence (green). CellMask-labeled membranes can be observed around dsRNA clusters. Virions at the plasma membrane are seen as bright puncta (right side and lower right corner of the image). **b**, Zoomed-in image that corresponds to the white box in **a**. **c**, Bivariate pair-correlation functions indicate nanoscale anti-correlation between dsRNA and CellMask, consistent with dsRNA encapsulation in membrane-bound organelles at both 6 and 24 hpi. Scale bars, 2 μm (**a**) and 500 nm (**b**).

1310

1311

1312

1313

Fig. S13. Nanoscale co-organization of viral components with host cell membranes.

a, SR images of nsp3, Sec61β, vgRNA and membranes (CellMask) in SARS-CoV-2 infected cells at 6 and 24 hpi. **b**, Bivariate pair-correlation functions indicate nanoscale association between nsp3 and Sec61β, nsp3 and CellMask, and nanoscale anti-correlation between vgRNA and CellMask at both time points. Scale bar, 1 µm.

1319

1320

Fig. S14. Nanoscale anti-correlation of nsp3 with dsRNA.

a, SR image of a SARS-CoV-2 infected cell at 6 hpi with nsp3 and dsRNA labeled by immunofluorescence. Nsp3 can be observed at the surface of isolated dsRNA clusters (white boxes & right panel) or in dense aggregates between dsRNA clusters (blue box & blue insets). **b**, Bivariate pair-correlation functions indicate nanoscale anti-correlation between dsRNA and nsp3 at 6 hpi. **c**, SR image of a SARS-CoV-2 infected cell at 24 hpi. Nsp3 forms a network-like pattern that encapsulates dsRNA clusters. **d**, Bivariate pair-correlation functions indicate nanoscale anti-correlation between dsRNA and nsp3 at 24 hpi. Scale bars, 2 μm (**a**, **c**) and 500 nm (insets in **a**, **c** and right panel in **a**).

1329

1330

Fig. S15. Nanoscale colocalization of nsp3 with membranes at 24 hpi.

SR image of nsp3 (green) and membranes as labeled by CellMask (magenta) in SARS-CoV-2 infected cells at 24 hpi. Nsp3 forms a network-like pattern in the perinuclear region that colocalizes with the CellMask pattern. Scale bar, 2 μm.

1335

1336

1337

1338

1339 **Fig. S16. Less common patterns of nanoscale nsp3 localization at 24 hpi.**

1340 **a**, Nsp3 forms an ER-like network that occupies a large part of the cytoplasm. **b**, Besides the
1341 common perinuclear pattern, Nsp3 is also diffusely localized throughout the whole cytoplasm.
1342 Scale bars, 2 μm.

1343

1344

1345

Fig. S17. Nanoscale localization of spike protein at 24 hpi.

a, SR image of a SARS-CoV-2 infected cell at 24 hpi labeled for spike (green) and vgRNA (magenta). **b**, Examples of assembled virions encapsulated by the spike proteins and with vgRNA in their interior, detected at the cell periphery (yellow arrows in **a**). (bottom panel) Bivariate pair-correlation functions calculated in the plasma membrane regions indicate colocalization of these targets at r < 100 nm. **c**, Magnified image that corresponds to the blue frame in **a** displays spike localizations mostly excluded from the interior of the perinuclear vgRNA clusters with possible localization at their membrane. (bottom panel) Bivariate pair-correlation functions calculated in the perinuclear regions of infected cells indicate nanoscale anti-correlation of spike with SARS-CoV-2 replication organelles. Scale bars, 2 µm (**a**), 100 nm (**b**), 500 nm (**c**).

1356

1357

1358

Fig. S18. Nanoscale anti-correlation of nucleocapsid protein with SARS-CoV-2 replication organelles at 24 hpi.

a, SR image of a SARS-CoV-2 infected cell at 24 hpi labeled for the nucleocapsid protein (green) and vgRNA (magenta). The magnified image in the blue frame displays nucleocapsid protein localizations mostly excluded from the interior of the perinuclear vgRNA clusters with possible localization at their membrane. **b**, Bivariate pair-correlation functions calculated in the perinuclear regions of the infected cells indicate nanoscale anti-correlation of the nucleocapsid protein with vgRNA. Scale bars, 2 μm and 500 nm (bottom panels).

1367

1368

Fig. S19. Optimization of primary antibody concentrations.

The concentration of primary antibodies was optimized to minimize the background or to

maximize the signal to background ratio (SBR) between SARS-CoV-2 infected and non-infected

cells or between cells expressing Sec61β-GFP and WT cells (see Methods). Box plots: center line,

median; box limits, upper and lower quartiles; whiskers, 1.5x interquartile range; dots, values for

individual cells. SBR plots show mean ± SD.

Fig. S20. Optimization of secondary antibody concentrations.

The concentration of secondary antibodies was optimized to minimize the background or to maximize the SBR between SARS-CoV-2 infected and non-infected cells (see Methods). Box plots: center line, median; box limits, upper and lower quartiles; whiskers, 1.5x interquartile range; dots, values for individual cells. SBR plots show mean ± SD.

1385

1386

1387

1388

Fig. S21. Path diagram of SR microscope used in this study.

Black-filled icons: mirrors; thin empty rectangles: dichroic or neutral density filters; dashed rectangles: movable or motorized components; boxes: cameras or lasers; bent lines: optical fiber; icons with blue edges: lenses or a beam splitter cube; QWP: quarter-wave plate; IP: image plane; IIP: intermediate image plane; BS: beam splitter; OD: optical density. Optics are shown for producing a second image on the EMCCD, but the second path was not used in this study. The gray lines denote the 808 nm beam in the focus lock apparatus.

1396

1397