Nanoscale cellular organization of viral RNA and proteins in SARS-CoV-2 replication organelles

Leonid Andronov*[1], Mengting Han*[2], Yanyu Zhu[2], Anish R. Roy[1], Andrew E. S. Barentine[1],

Jaishree Garhyan[3], Lei S. Qi[2,4,5], W.E. Moerner[1,4]

[1]Department of Chemistry; [2]Department of Bioengineering; [3]In Vitro Biosafety Level 3 (BSL-3)

Service Center, School of Medicine; [4]Sarafan ChEM-H; Stanford University, Stanford, CA 94305

U.S.A. [5]Chan Zuckerberg Biohub – San Francisco, San Francisco, CA 94158 U.S.A.

*These authors contributed equally.

#Correspondence to: W. E. Moerner, wmoerner@stanford.edu; Lei S. Qi, sqi@stanford.edu

Abstract:

The SARS-CoV-2 viral infection transforms host cells and produces special organelles in many ways, and we focus on the replication organelle where the replication of viral genomic RNA (vgRNA) occurs. To date, the precise cellular localization of key RNA molecules and replication intermediates has been elusive in electron microscopy studies. We use super-resolution fluorescence microscopy and specific labeling to reveal the nanoscopic organization of replication organelles that contain vgRNA clusters along with viral double-stranded RNA (dsRNA) clusters and the replication enzyme, encapsulated by membranes derived from the host endoplasmic reticulum (ER). We show that the replication organelles are organized differently at early and late stages of infection. Surprisingly, vgRNA accumulates into distinct globular clusters in the cytoplasmic perinuclear region, which grow and accommodate more vgRNA molecules as infection time increases. The localization of ER labels and nsp3 (a component of the double-membrane vesicle, DMV) at the periphery of the vgRNA clusters suggests that replication organelles are enclosed by DMVs at early infection stages which then merge into vesicle packets as infection progresses. Precise co-imaging of the nanoscale cellular organization of vgRNA, dsRNA, and viral proteins in replication organelles of SARS-CoV-2 may inform therapeutic approaches that target viral replication and associated processes.

Keywords: SARS-CoV-2 coronavirus, viral replication, fluorescence imaging, super-resolution fluorescence microscopy, viral RNA localization, COVID-19, viral proteins, cell infection

Introduction

Due to its global health impact, the SARS-CoV-2 coronavirus and its infection of mammalian cells have been the subject of a large number of studies across multiple fields. Biochemical methods have allowed researchers to investigate the interactions between the viral oligonucleotides and the host proteins *in vitro* and in cellular extracts, leading to much insight[1,2]. There have also been electron microscopy (EM) studies of resin-embedded samples as well as vitrified samples using cryo-electro tomography, all of which have been profiting from the large increase in EM resolution and contrast in recent years. These EM studies can provide very high-resolution structures of protein complexes as well as tomograms of organelles in the cellular context. High-contrast filamentous structures and membranes appear regularly in such images, allowing identification of single- and double-membrane vesicles (DMVs)[3-5]. However, the all-important viral RNA and associated proteins are challenging to identify by EM due to a lack of specific contrast. While some researchers have detected RNA-like filaments in vesicles[4,5], further investigations are needed to identify specific viral RNAs in the cellular context.

Fluorescence microscopy offers a highly useful and complementary set of capabilities, most importantly the specific labeling of proteins or RNA sequences. However, conventional diffraction-limited (DL) fluorescence microscopy, with its resolution constrained to ~250 nm, is unable to resolve the tiny structures that are hidden in a blurred DL image. Super-resolution (SR) microscopy based on single molecules (PALM[6], (d)STORM[7,8]) or on structured patterns of molecular depletion (STED[9], SIM[10]), however, offers far better optical resolution down to 10 nm and below. A wealth of important cellular patterns and structures have been identified in recent years, such as the banding patterns of axonal proteins in neuronal cells[11] and many others[12-14]. The specificity of SR imaging is useful to apply to the study of viral genomic RNA (vgRNA) and other RNA molecules as well as protein players involved in coronavirus infection of cells, demonstrated previously for the less pathogenic human coronavirus 229E (HCoV-229E)[15].

In this work, we apply multicolor confocal microscopy and SR microscopy to explore the localization patterns of viral RNA and protein molecules for SARS-CoV-2 during the early and late infection of mammalian cells. We focus particularly on viral genomic RNA (vgRNA) and its relative,

the double-stranded RNA (dsRNA) that forms between the (+) sense vgRNA and the (-) sense copy.

After the initial infection with a few copies of vgRNA, more vgRNA and dsRNA are synthesized by

the RNA-dependent RNA polymerase complex (RdRp), an early essential enzyme formed by

ribosomal polyprotein synthesis and the viral proteases. We also co-image a series of molecules,

including membrane markers, nucleocapsid protein, spike protein, and the nsp3 protein (reported

to be a major component of a molecular pore spanning both membranes of DMVs[16]), all to

provide context and support for the view that vgRNA, dsRNA, and RdRp act spatially in replication

organelles (ROs) during virus replication. Thus, we provide key information about where these

important players are found in infected cells and how they change with time during infection. Our

results yield a nanoscale optical readout of viral nucleic acid organization during SARS-CoV-2

infection, highlight the structural importance of ROs, and could potentially benefit development

of future therapeutic approaches.

Results

Labeling and imaging of SARS-CoV-2 virions

To specifically detect SARS-CoV-2 vgRNA, we applied RNA fluorescence *in situ* hybridization (RNA

FISH) with 48 antisense DNA oligonucleotide probes[17] specifically targeting the open reading

frame 1a (ORF1a) region which is only present in vgRNA and not in sgRNAs, insuring detection of

only full-length viral (+)vgRNA (Fig. 1a). Each probe was conjugated with a single blinking

fluorophore for (d)STORM (direct Stochastic Optical Reconstruction Microscopy)[8]. To test this

labelling and imaging approach, we first imaged vgRNA along with SARS-CoV-2 spike protein in

purified virions (Extended Data Fig. 1). While the size of SARS-CoV-2 virions is too small to resolve

in conventional DL fluorescence microscopy (Extended Data Fig. 1a), in SR the internal concentric

organization of the virions can be observed with vgRNA found in their center and spike at the

surface (Extended Data Fig. 1b). The labelling efficiency with these probes is around 6 dyes/vgRNA

in partially Proteinase K-digested virions, which was higher than in intact virions due to poorer

accessibility of their vgRNA (Extended Data Fig. 1c-i).

Next, we imaged SARS-CoV-2 infected Vero E6 cells that were fixed at 24 hours post infection (hpi)

and then labeled for immunofluorescence imaging (Methods). Spike and nucleocapsid SR

microscopy in these cells revealed assembled virions mostly at the cellular periphery, often at

cytoplasmic tubular projections, indicating active viral production (Extended Data Fig. 1j), similar

to previously reported results[5,18]. We now turn to the main focus of this study, the replication of

viral genomic RNA.

Fig. 1: Clustering of vgRNA in the cytoplasm of infected cells.

a, Scheme of SARS-CoV-2 genome with constructs used for its detection in infected cells. 48 antisense DNA oligonucleotide probes were used to target the ORF1a-coding region of vgRNA that is exclusive to the (+)vgRNA and does not occur in the sgRNAs. The RNA FISH probes are conjugated with AF647 or CF568. **b**, Representative confocal images of vgRNA in infected Vero E6 cells at 6 hpi display scattered DL puncta. **c**, Representative SR image of an infected cell at 6 hpi reveals distinct vgRNA clusters in the cytoplasm. **d**, Histogram of the radii of gyration (Rg) of the

vgRNA clusters indicate their size increase between 6 hpi (magenta) and 24 hpi (green). **e**,
Zoomed-in region of the SR image (green frame in **c**) displays an agglomeration of vgRNA clusters.
f, BIC-GMM cluster analysis of the region shown in **e**. **g**, Representative confocal images of vgRNA
in infected Vero E6 cells at 24 hpi display large DL foci in the perinuclear region of the cytoplasm.
h, Representative SR image of an infected cell at 24 hpi reveals large perinuclear vgRNA clusters.
i, BIC-GMM cluster analysis of the cell shown in **h**. **j**, Zoomed-in region of the SR image (blue frame
in **h**) displays dense vgRNA clusters. **k**, BIC-GMM cluster analysis of the region shown in **j**. Scale
bars, 10 μm (**b**, **g**), 1 μm (**c**, **h**), 200 nm (**e**, **f**, **j**, **k**). Dashed lines in **c** and **h** indicate the position of
the cell nucleus. Localizations that belong to the same cluster in **f**, **i**, **k** are depicted with the same
color. Color bars in **c**, **e**, **h**, **j** show the number of SM localizations within each SR pixel (20 x 20
nm^2).

SARS-CoV-2 genomic RNA clusters in cytoplasm of infected cells

Confocal screening demonstrated three patterns of intracellular vgRNA localization (Extended
Data Fig. 2a): scattered puncta in the cytoplasm (Type 1, Fig. 1b), appearance of bright foci in the
perinuclear region (Type 2, Extended Data Fig. 2a), and concentration of vgRNA into large dense
structures that occupy most of the perinuclear region (Type 3, Fig. 1g). We find that Type 1 cells
were most abundant at 6 hpi, and Type 3 cells at 24 hpi, indicating that the vgRNA localization
progresses from Type 1 to Type 3 as infection advances in time (Extended Data Fig. 2b). We also
find that the cell-integrated vgRNA FISH signal in infected cells increases 2.2x on average from 6
to 24 hpi (Extended Data Fig. 2c), representing active viral replication and accumulation of vgRNA
inside the cells.

The higher spatial resolution of SR microscopy revealed that at 6 hpi (Type 1 and Type 2 cells),
most vgRNA localizes into clusters with an approximately round shape and a diameter of 100-250
nm that scatter in the cytoplasm (Fig. 1c, e). At 24 hpi (Type 2 and Type 3 cells), the vgRNA
localization pattern transformed into a fascinating dense perinuclear network of approximately
round, often hollow structures with a diameter of 300-700 nm (Fig. 1h, j). To quantify the
transformation of vgRNA to clusters in infected cells, we performed a Bayesian Information
Criterion-optimized Gaussian Mixture Model clustering analysis (BIC-GMM) (Fig. 1f, i, k; See

Methods). This analysis showed an increase in the median vgRNA cluster size (radius of gyration)

from 73 nm at 6 hpi to 187 nm at 24 hpi (Fig. 1d, see inset) reflecting the drastic change in vgRNA

localization pattern.

Besides dense vgRNA clusters, we observe isolated localizations of individual vgRNA molecules

scattered in the cytoplasm at both time points, in line with previously reported results on the

229E virus[15,17]. These appear as a haze in confocal images (Extended Data Fig. 2a, Type 3) but are

resolved as sparse nanoscale puncta (d < 50 nm) in SR (Fig. 1h, Extended Data Fig. 3a) which we

assume to be single vgRNA copies (even though the puncta are more dense at 24 hpi). Using the

average number of single-molecule (SM) localizations per vgRNA punctum as a calibration for the

number of localizations per single vgRNA, we estimated the average number of vgRNA molecules

in the vgRNA clusters to be around 26 vgRNA/cluster at 6 hpi, increasing by almost an order of

magnitude to 181 vgRNA/cluster at 24 hpi (Extended Data Fig. 3b-c).

dsRNA associates with vgRNA clusters

Next, we proceeded to assess the relation of vgRNA cluster locations to viral replication. For this,

we immunofluorescently labelled an intermediate of coronavirus replication and transcription,

the hybridized dsRNA objects composed of (+) sense vgRNA and (-) sense copy, and co-imaged

dsRNA with vgRNA using two-color confocal and SR microscopy. These targets appeared mostly

colocalized at both timepoints at low resolution (Fig. 2a-b), suggesting that vgRNA clusters are

often found close to the replication centers of SARS-CoV-2. SR microscopy revealed that dsRNA

aggregates into clusters of a relatively compact size (d ≈ 100-200 nm) with distinct patterns of

colocalization with vgRNA at 6 or 24 hpi (Fig. 2c-e).

To quantify the spatial relationship between dsRNA and vgRNA, we conducted pair-pair

correlation analysis[19]. We calculated a bivariate pair-correlation function $g_{12}(r)$, *i.e.*, the

distribution of the pairwise distances between the localizations of the two species[20]. The function

is computed only in perinuclear regions and is normalized in a way that $g_{12}(r) = 1$ for two randomly

and homogeneously distributed species, signifying complete spatial randomness (CSR). Closely

associated or colocalized species have a prevalence of short pairwise distances resulting in a peak

in $g_{12}(r)$ near r = 0, while anti-correlated species lack short interparticle distances, which lowers

$g_{12}(r)$ at r = 0 followed by peaking at r > 0.

At early infection stages (6 hpi), dsRNA clusters appear closely associated with or adjacent to

vgRNA clusters both visually and by pair-pair correlation analysis (Fig. 2c, f). By contrast, during

late infection (24 hpi), dsRNA clusters anticorrelate with vgRNA at short distance scales with an

average separation between them around 120 nm as indicated by bivariate pair-correlation

functions $g_{12}(r)$ (Fig. 2g). Moreover, at 24 hpi, dsRNA clusters can often be found in the voids of

the large vgRNA structures (Fig. 2d), suggesting their possible concentric localization in the same

ROs.

Contrary to vgRNA, the size of dsRNA clusters slightly decreases and the total brightness of

cellular dsRNA labelling does not significantly change between 6 hpi and 24 hpi (Fig. 2e, Extended

Data Fig. 2d). Interestingly, at 6 hpi but not at 24 hpi, the dsRNA signal per cell positively correlates

with that of vgRNA signal (Extended Data Fig. 2f-g). These findings indicate that the amount of

dsRNA increases at early infection but reaches saturation by 24 hpi. This may suggest that after

the rapid initial production of a dsRNA pool, further generation of (-) sense copies slows down

and the replication shifts to the generation of vgRNA from the pool of available (-) sense copy

templates, which is common in other coronaviruses[21]. Similar anticorrelation between the

localizations of vgRNA and dsRNA, but constant dsRNA cluster size with infection time have been

previously observed for the 229E coronavirus[15], consistent with the related biology of 229E and

SARS-CoV-2.

Fig. 2: Association of dsRNA with vgRNA clusters

a-b, Representative confocal images of SARS-CoV-2 infected cells display DL colocalization between dsRNA (green) and vgRNA (magenta) at both 6 hpi (**a**) and 24 hpi (**b**). **c-d**, Representative SR images of SARS-CoV-2 infected cells indicate association between dsRNA and vgRNA at 6 hpi (**c**) and short-range anti-correlation often with concentric localization at 24 hpi (**d**). Bottom panels, zoomed-in images of corresponding colored boxes. **e**, Histogram of Rg of dsRNA clusters as determined by the BIC-GMM cluster analysis (left). Median Rg of dsRNA clusters significantly decreases between 6 hpi and 24 hpi (right). p-value = $8 \cdot 10^{-4}$, two-tailed t-test. (**f-g**), Bivariate pair-correlation functions $g_{12}(r)$ calculated between the localizations of dsRNA and vgRNA indicate close association at 6 hpi (**f**) and nanoscale anti-correlation at 24 hpi (**g**). CSR, complete spatial randomness. Thin lines correspond to $g_{12}(r)$ of individual cells and bold lines are the mean values of $g_{12}(r)$ from all analyzed cells. Scale bars, 10 μm (**a-b**), 1 μm (**c-d**), 500 nm (**c-d**, bottom panels). Dashed lines in **c** and **d** indicate the position of the cell nucleus.

vgRNA clusters denote the replication centers of SARS-CoV-2 genome

To investigate SARS-CoV-2 replication activity at the vgRNA clusters in more detail, we co-imaged

them with the RdRp complex, the replicating SARS-CoV-2 RNA-dependent RNA polymerase[22,23],

using immunofluorescent labelling of its catalytic subunit nsp12[24]. In confocal images, nsp12

adopts a similar pattern as vgRNA, colocalizing with it at both 6 hpi and 24 hpi (Fig. 3a-b), which

suggests ongoing replication at the vgRNA clusters. In SR images, nsp12 localized in small sparse

puncta (d < 50 nm) that were scattered within and next to the vgRNA clusters at both time points

(Fig. 3c-d). Because nsp12 puncta are well separated from each other, and oligomerization is not

expected[22,23,25], each nanoscale punctum is likely to represent a single replicating enzyme. On

average, we detected 2.5 nsp12 puncta per vgRNA cluster at 6 hpi and 7.6 at 24 hpi (Fig. 3g).

Interestingly, in contrast to vgRNA but similar to dsRNA, the total nsp12 amount does not

significantly increase (Extended Data Fig. 2e) and its nanoscale localization pattern stays the same

as infection progresses from 6 to 24 hpi (Fig. 3c-d). This suggests that the growth of vgRNA clusters

arises from relatively constant small numbers of replication components between 6 and 24 hpi

highlighted by the constant amount of dsRNA and RdRp. Bivariate cross-correlation functions

calculated between nsp12 and vgRNA localizations peaked at 0 nm indicating association of these

two targets at both 6 and 24 hpi (Fig. 3e-f). Since vgRNA clusters colocalize with the catalytic

subunit of RdRp, we conclude that vgRNA clusters combined with the nearby RdRp enzymes and

dsRNA highlight ROs that act as centers for replication and transcription of SARS-CoV-2.

Fig. 3: Association of SARS-CoV-2 replication enzyme with vgRNA clusters

a-b, Representative confocal images of SARS-CoV-2 infected cells display DL colocalization between nsp12, the catalytic subunit of RdRp (green) and vgRNA (magenta) at both 6 hpi (**a**) and 24 hpi (**b**). **c-d**, Representative SR images of SARS-CoV-2 infected cells indicate nanoscale association between nsp12 and vgRNA at both 6 hpi (**c**) and 24 hpi (**d**). Insets show magnified images of corresponding regions in colored boxes. **e-f**, Bivariate pair-correlation functions peak at r = 0 nm indicating association between nsp12 and vgRNA. **g**, Number of nanoscale puncta of nsp12 per vgRNA cluster. **h**, SR image of nsp12 with Sec61β suggests encapsulation of nsp12 within ER-derived membranes. Scale bars, 10 μm (**a-b**), 1 μm (**c-d**), 500 nm (**h** and insets in **c-d**). Dashed lines in **c**, **d** and **h** indicate the edge of the cell nucleus.

vgRNA clusters are enclosed in ER-derived membranous organelles

Coronaviruses are thought to transform the host ER into replication-permissive structures, such as convoluted membranes and DMVs[3,26,27]. To investigate the relation of vgRNA clusters with cellular ER, we immunofluorescently labelled Sec61β, an ER membrane protein[28], in Vero E6 cells

stably expressing Sec61β-GFP[15]. Confocal images of these cells show the appearance of Sec61β

spots that colocalize with vgRNA against the mostly unaltered ER background at 6 hpi (Fig. 4a). At

24 hpi, however, substantial amounts of Sec61β accumulate close to the perinuclear vgRNA

clusters, while the ER tubules outside these regions become poorly visible (Fig. 4a), consistent

with the virus-induced rearrangement of the ER and the inhibition of host gene expression by

SARS-CoV-2[29].

In SR, we observe encapsulation of the vgRNA clusters by ring-like structures of the altered ER at

6 hpi (Fig. 4b, Extended Data Fig. 4). As infection progresses, the ER-derived ring- or sphere-like

structures grow to accommodate larger vgRNA clusters at 24 hpi (Fig. 4b, Extended Data Fig. 5).

Pair-correlation functions peak at the distance of the typical radius of vgRNA clusters indicating

nanoscale anti-correlation compatible with the ER-derived encapsulation of vgRNA (Fig. 4c).

dsRNA (Fig. 4b, Extended Data Fig. 6) and nsp12 (Fig. 3h) are also found to be encapsulated by

the same remodeled ER membranes suggesting that vgRNA, dsRNA and RdRp are all located

within the same ER-derived replication organelles.

To further confirm that these clusters are surrounded by membranes, we used a (d)STORM-

compatible general membrane marker CellMask Deep Red[30]. This dye broadly stains cellular

membranes, including the nuclear envelope, the mitochondrial membranes, and SARS-CoV-2

virions at the plasma membrane (Extended Data Fig. 7). The nanoscale image contrast with

CellMask Deep Red is poorer than specific protein labelling of the Sec61β ER label due to

background from membranes of different cellular organelles. Nevertheless, in the perinuclear

region of infected cells, we observed the appearance of a complex membranous network that

anti-correlates with vgRNA and dsRNA, with visible encapsulation of vgRNA and dsRNA clusters

(Extended Data Fig. 7, 9). Taken together, these findings indicate that the vgRNA-dsRNA-RdRp

clusters are located inside membrane-bound replication organelles (ROs) that originate from

altered host ER transformed by SARS-CoV-2.

Fig. 4: vgRNA clusters are encapsulated in membranes of remodeled ER

a, Representative confocal images of SARS-CoV-2 infected cells indicate an appearance of dense perinuclear foci of Sec61β ER labelling at 24 hpi that colocalizes with vgRNA and nsp3. **b**, SR images reveal concentric organization of Sec61β around vgRNA and dsRNA and colocalization of Sec61β with nsp3. **c**, Bivariate pair-correlation functions indicate anti-correlation of Sec61β with vgRNA and dsRNA and association of Sec61β with nsp3. Scale bars, 20 μm (**a**) and 1 μm (**b**). Dashed lines in **b** indicate the position of the cell nucleus.

Nsp3 localizes at the surface of SARS-CoV-2 replication organelles

Because the nsp3 protein of betacoronaviruses is essential for the DMV formation[31,32], and nsp3 is a constituent of a DMV molecular pore[16], we proceeded to localize this non-structural protein to relate the ROs to the SARS-CoV-2-induced DMVs. At DL resolution, nsp3 labelling adopts a pattern that colocalizes with vgRNA at both 6 and 24 hpi, similarly to dsRNA and nsp12 (Fig. 5a, f). SR imaging of these cells, however, revealed striking nanoscale positioning of nsp3. At 6 hpi, sparse nsp3 can be found surrounding isolated vgRNA clusters (Fig. 5c-d), while larger nsp3 aggregates are situated amidst bunched vgRNA clusters (Fig. 5e). At 24 hpi, nsp3 localizes at the borders of the large vgRNA clusters, encircling them in incomplete rings and forming a partial perinuclear network (Fig. 5g, i, j). Similar nsp3 arrangements can be observed in relation to dsRNA (Extended Data Fig. 8).

The anti-correlation of vgRNA with nsp3 and dsRNA with nsp3 (Fig. 5, Extended Data Fig. 8) closely resemble the pattern observed with vgRNA and dsRNA with Sec61β (Fig. 4, Extended Data Fig. 6), suggesting that nsp3 may also be localized at the ER-derived membranous surface of the ROs. To further confirm this hypothesis, we co-imaged nsp3 with Sec61β and CellMask (Fig. 4, Extended Data Fig. 9, Supplementary Fig. S1). The SR images and the pair-correlation analysis indicated colocalization between nsp3 and both membrane markers at both time points (Fig. 4b-c, Extended Data Fig. 9, Supplementary Fig. S1), confirming that nsp3 localizes on the membranes encircling the SARS-CoV-2 ROs.

Besides these characteristic localization patterns of nsp3, we observed a few cells with two different phenotypes at 24 hpi, one with an ER-like network that occupies large regions in the cytoplasm (Extended Data Fig. 10a), and another one with nsp3 densely diffused throughout the whole cytoplasm (Extended Data Fig. 10b). The ER-like network may represent nsp3 proteins being heavily translated on ER membrane, while nsp3 proteins found outside the perinuclear region are less likely to be associated with the SARS-CoV-2 replication process and might represent other nsp3 functions, such as a papain-like proteolytic function[33] or post-translational modification of host proteins[34], which can become objects of future SR studies.

The localization of nsp3 at the surface of isolated vgRNA-dsRNA clusters at 6 hpi is consistent with

the localization of molecular pores on the DMV membrane observed by cryo-EM[16]. At late

infection times, DMVs have been observed to merge into vesicle packets (VPs)[5] that are also likely

to contain pores, however molecular pores in the VP membranes have not yet been studied in

detail to our knowledge. Nevertheless, previous studies report that in late infection the

perinuclear region becomes filled with DMVs and VPs[26] that strongly resemble the ROs reported

here. The size of vgRNA clusters at 6 hpi and at 24 hpi from our data is similar to the previously

reported size of DMVs and VPs, correspondingly[5]. Taken together, our results provide evidence

that vgRNA accumulates, possibly in DMVs at 6 hpi and in VPs at 24 hpi. dsRNA clusters occur

within the same vesicles but occupy distinct parts of them. Our data are consistent with the

picture that SARS-CoV-2 RNA is replicated and transcribed within these DMVs and VPs as

highlighted by the proximal localizations of RdRp.

321

Fig. 5: Nsp3 localizes at the surface of vgRNA clusters

a, Representative confocal images of a SARS-CoV-2 infected cell display DL colocalization between punctate vgRNA and nsp3 labeling at 6 hpi. **b**, Bivariate pair-correlation functions calculated between the SR localizations of vgRNA and nsp3 indicate nanoscale anti-correlation of these targets at 6 hpi. **c**, Representative SR image of a SARS-CoV-2 infected cell at 6 hpi. **d**, Zoomed-in images of selected vgRNA particles (yellow boxes in **c**) indicate the localization of nsp3 at the surface of the vgRNA clusters. **e**, Magnified region with aggregates of vgRNA clusters (blue box in **c**) displays dense nsp3 localization in the core of these aggregates. **f**, Confocal images indicate that vgRNA and nsp3 occupy approximately the same regions in a SARS-CoV-2 infected cell at 24 hpi. **g**, Representative SR image of a SARS-CoV-2 infected cell at 24 hpi. **h**, Bivariate pair-

correlation functions indicate nanoscale anti-correlation between vgRNA and nsp3 at 24hpi. **i-j**,

Magnified regions of the SR image (colored boxes in **g**) show that nsp3 localizes in interstitial

regions or encapsulates vgRNA clusters. Scale bars, 10 μm (**a**, **f**), 1 μm (**c**, **g**), 500 nm (**i**, **j**), 200nm

(**d**, **e**). Dashed line in **g** indicates the position of the cell nucleus.

Discussion

Previous biochemical and EM studies allowed researchers to build models of the intracellular life

cycle of SARS-CoV-2[35-37]; however, precise localization of specific viral proteins and RNA molecules

is challenging due to lack of specific contrast in EM and low resolution in DL fluorescence

microscopy. SR fluorescence microscopy is well suited for coronavirus studies in cells as it provides

both specific contrast and high resolution (~20 nm and below depending upon photons

collected[38]). However, to date few studies have employed this method for coronavirus biology[15],

even less for SARS-CoV-2[27,39,40], and none of them focused on the SARS-CoV-2 replication process

in detail. Here we apply SR fluorescence microscopy to precisely localize the key players of SARS-

CoV-2 replication at different timepoints in infected cells. Building upon a previously developed

method for simultaneous labelling of coronavirus vgRNA with dsRNA and protein

immunofluorescence[15], and using improved fixation and multi-color SR imaging protocols (see

Methods), we obtain and quantify the appearance and molecular compositions of ROs of SARS-

CoV-2 in cells at different stages of infection.

In this study, our results taken together depict a compelling and novel picture of ROs containing

various molecules including vgRNA, dsRNA, RdRp, nsp3, and ER membrane (Fig. 6). In this model,

we compare organization of ROs at early and late stages of infection and show how specific RNA

and protein molecules are spatially organized in ROs. Compared to the 229E case, SARS-CoV-2

appears to generate more complex clusters of vgRNA, and with the additional imaging of relevant

targets in this study, the structural importance of ROs is now clear.

Fig. 6: Proposed model for SARS-CoV-2 replication organelles containing various RNA and protein molecules at early and late stages of infection.

The detailed intracellular localization of the central SARS-CoV-2 component, vgRNA, has remained vague in the literature. Our RNA FISH method[15] targets specific sequences in vgRNA (Fig. 1a) and detects single vgRNA molecules (Extended Data Fig. 1, 3a), allowing counting of the number of vgRNA molecules within specific regions (Extended Data Fig. 3b-c). We find for the first time that most cellular vgRNA localizes into dense clusters of an approximately round shape that grow and migrate to the perinuclear region as infection time increases. We show that these clusters appear confined in membranous vesicles derived from ER as emphasized by the localization of Sec61β and CellMask at their surface (Fig. 4b, Extended Data Fig. 4, 5, 9). From comparison with earlier EM images[5,16,26] and from nsp3 localization at their surface[16] (Fig. 5), we can conclude that these vesicles are most likely DMVs at an early-mid infection time that grow and merge into VPs as infection progresses.

Previously, metabolic radioactive labelling was used to localize newly synthesized RNA in SARS-CoV-1 and MERS-CoV-infected cells to DMVs[4]. However, metabolic labelling could only localize a fraction of viral RNA molecules with little sequence specificity and with a background of newly transcribed cellular RNA. Here, we specifically label vgRNA of SARS-CoV-2 for SR microscopy and

show that it also localizes in patterns that suggest confinement in DMVs, confirming the earlier

findings on SARS-CoV-1 and MERS-CoV[4].

Previous studies also reported presence of dsRNA in DMVs of SARS-CoV-1[3] and SARS-CoV-2[5]. EM

images of DMVs often display a complex filamentous network in their interior, that was attributed

to viral RNA molecules[5]. However, the exact type of these RNAs was not determined due to the

absence of specific labelling. As one might expect, single-stranded vgRNA can form a secondary

structure that includes many short dsRNA fragments *e.g.*, in stem loops[41,42]. This makes it difficult

to distinguish between viral dsRNA and vgRNA by measuring the diameter of the filaments, taking

into account that the detection probability of ssRNA might be lower due to a decreased EM

contrast for ssRNA than for dsRNA. Reported abundant branching of filaments in DMVs[5], however,

is typical for ssRNA secondary structures[43]. Indeed, there was evidence in the literature about

presence of both dsRNA and vgRNA in DMVs; however, to our knowledge, there was no

simultaneous observation of both vgRNA and dsRNA within the same DMVs.

Here we use the J2 anti-dsRNA antibody that recognizes only long dsRNA fragments (≥ 40bp) with

no detection of the ssRNA secondary structures[44,45]. Two-color SR imaging revealed for the first

time that most dsRNA and vgRNA are located within the same DMVs and VPs, occupying distinct

regions of these vesicles, and adopting an anti-correlation pattern at short distances (r < 100 nm)

at 24 hpi (Fig. 2). A similar nanoscale anti-correlation has been observed in the 229E coronavirus

but the DMVs or ROs were not specifically labelled in that study[15]. Another novel observation is

the relatively constant amount of dsRNA and little change in dsRNA cluster size between 6 and

24 hpi despite the huge change in the vgRNA landscape (Fig. 2).

It has been proposed that the RdRp complex of SARS-CoV-1 is located at convoluted membranes

and inside DMVs based on immunogold labelling of nsp8[3]. However, nsp8 has intracellular

functions other than as an RdRp accessory subunit[46,47] that might be exercised at the convoluted

membranes. Here we label the catalytic RdRp subunit, nsp12[22], and find that it mostly localizes

to the vgRNA clusters at both 6 and 24 hpi (Fig. 3), suggesting that SARS-CoV-2 replication and

transcription occurs preferentially in the vgRNA-filled ROs, where dsRNA resides as well.

Nsp3 of betacoronaviruses (SARS-CoV-1, MERS-CoV and MHV) was previously localized to the

convoluted membranes and to the DMV membranes using immuno-EM[3,4,48,49] and cryo-ET[16];

however, these studies were limited to early-mid infection at 8-12 hpi. In our study, we report

two localization patterns of nsp3 of SARS-CoV-2 at 6hpi: 1) sparse nsp3 at the surface of isolated

vgRNA-dsRNA clusters (Fig. 5d, Extended Data Fig. 8a); 2) dense nsp3 within the accumulations

of vgRNA-dsRNA clusters (Fig. 5e, Extended Data Fig. 8a). While the first pattern most likely

corresponds to the RO/DMV membranes considering the role of nsp3 as a DMV pore[16], the

second one resembles a pattern found in other coronaviruses that was attributed to the

convoluted membranes[3,4,50]. Convoluted membranes are typically found within dense groups of

DMVs in early-mid infection[3,4] and localization of nsp3 on them might represent early steps of

viral transformation of ER into DMVs. We found this nsp3 pattern anti-correlated with vgRNA (Fig.

5c, e) and with dsRNA (Extended Data Fig. 8a-b), suggesting little to no vgRNA or dsRNA at the

convoluted membranes, in line with previous studies on other coronaviruses[4].

At 24 hpi, we did not observe these early infection patterns of nsp3 localization. Instead, we show

for the first time that at 24 hpi, nsp3 densely localizes at the membranes that separate large

vgRNA clusters and grows into a considerable perinuclear network that contains the ROs (Fig. 5g,

i-j, Extended Data Fig. 8c). Since the molecular pores of VPs have not yet been investigated in

detail, we can speculate that this late infection nsp3 pattern corresponds to the pores of VPs that

should also be much denser than those of isolated DMVs, considering the increased density of

nsp3 labelling. Additional rare phenotypes of nsp3 localization that we also report for the first

time (Extended Data Fig. 10) illustrate the variability of SARS-CoV-2 infection course and should

lead to further research on the other intracellular functions of this viral protein.

Taken together, we investigated several key factors of SARS-CoV-2 replication: vgRNA, dsRNA,

RdRp and nsp3 inside infected cells with SR microscopy for the first time. We discovered and

characterized perinuclear clusters of vgRNA and demonstrated by RdRp labelling that they

associate with SARS-CoV-2 ROs. We found that the ROs also contain dsRNA and are encapsulated

in ER-derived membranes. Using SR data on nsp3, we conclude that these virus-induced

organelles correspond to DMVs.

This study expands the knowledge of the biology of coronaviruses and opens new possibilities for

therapeutics against SARS-CoV-2. Careful examination of the organization of replication

organelles may provide new avenues to target the organelles to disrupt SARS-CoV-2 replication

and transcription. Examining localization patterns for different viral variants or in different host

cells will be useful to broaden understanding of the viral infection. It will also be important to

examine how the structures reported in this study change upon the addition of drug treatments.

Our imaging approach may also offer insights into long COVID by investigating cells that are

infected by SARS-CoV-2 that may still contain RO-like structures after symptoms disappear.

Methods

Antibodies

Primary antibodies and the optimal dilutions and concentrations used are as follows: goat polyclonal anti-spike S2 (Novus Biologicals, AF10774-SP, 1:20, 10 µg/mL), mouse monoclonal anti-dsRNA (SCICONS, 10010200, 1:200, 5 µg/mL), rabbit polyclonal anti-RdRp/nsp12 (Sigma-Aldrich, SAB3501287-100UG, 1:500, 2 µg/mL), mouse monoclonal anti-nucleocapsid (Thermo Fisher, MA5-29981, 1:500, 2 µg/mL), rabbit polyclonal anti-nsp3 (Thermo Fisher, PA5-116947, 1:134, 5 µg/mL), sheep polyclonal anti-GFP (Bio-Rad, 4745-1051, 1:1000, 5 µg/mL), and rabbit polyclonal anti-GFP (Novus Biologicals, NB600-308SS, 1:163, 5 µg/mL). Secondary antibodies and the optimal dilutions and concentrations used are as follows: AF647-conjugated donkey anti-mouse IgG (Thermo Fisher, A-31571, 1:500, 4 µg/mL), AF647-conjugated donkey anti-rabbit IgG (Thermo Fisher, A-31573, 1:500, 4 µg/mL), AF647-conjugated donkey anti-sheep IgG (Thermo Fisher, A-21448, 1:500, 4 µg/mL), CF568-conjugated donkey anti-goat IgG (Sigma-Aldrich, SAB4600074-50UL, 1:500, 4 µg/mL), CF568-conjugated donkey anti-rabbit IgG (Sigma-Aldrich, SAB4600076-50UL, 1:500, 4 µg/mL), CF568-conjugated donkey anti-mouse IgG (Sigma-Aldrich, SAB4600075-50UL, 1:500, 4 µg/mL), and CF568-conjugated donkey anti-sheep IgG (Sigma-Aldrich, SAB4600078-50UL, 1:500, 4 µg/mL).

Culture of cell lines

The Vero E6 cells (African green monkey kidney epithelial cells, ATCC, CRL-1586), HEK293T cells (human embryonic kidney epithelial cells, ATCC, CRL-3216), and Vero E6-TMPRSS2 cells were cultured in Dulbecco's modified Eagle medium (DMEM) with GlutaMAX, 25 mM D-Glucose, and 1 mM sodium pyruvate (Gibco, 10569010) in 10% FBS (Sigma-Aldrich, F0926) at 37°C and 5% CO_2 in a humidified incubator. Cell lines were not authenticated after purchase prior to use. For Vero E6-TMPRSS2, Geneticin (G418) was added at a final concentration of 1mg/ml.

Lentivirus production for ER labeling with Sec61β

To produce lentivirus, HEK293T cells were cultured in 10-cm dishes and transiently transfected

with 9 µg lentiviral plasmid pLV-ER-GFP (Addgene, 80069, a gift from Pantelis Tsoulfas), 8 µg

pCMV-dR8.91, and 1 µg PMD2.G packaging plasmids using 25 µL TransIT-LT1 Transfection Reagent

(Mirus, MIR 2306). After 72 h of transfection, supernatant was filtered through 0.45 µm filters,

concentrated using Lentivirus Precipitation Solution (ALSTEM, VC100) at 4°C overnight, and

centrifuged at 1,500x g for 30 min at 4°C to collect virus pellets. The virus pellets were

resuspended in cold DMEM for storage at -80°C for transduction of cells.

Generation of stable cell line

To generate a Vero E6 cell line stably expressing Sec61β-GFP, $2x10^5$ Vero E6 cells were seeded in

one well of a 6-well plate and infected with one quarter of concentrated lentivirus expressing pLV-

ER-GFP produced from one 10-cm dish of HEK293T cells while seeding. After two days incubation,

monoclonal cells expressing GFP were sorted out using a SONY SH800S sorter. These transduced

cells were only used for ER imaging; all other experiments used wild type (WT) cells.

SARS-CoV-2 viral stocks preparation

The SARS-CoV-2 WA 1, isolate USA-WA1/2020 (NR-52281, BEI Resources) was passaged 3 times

in Vero E6-TMPRSS2 cells as previously described[51,52]. Briefly, a Vero E6-TMPRSS2 monolayer was

infected with virus obtained from BEI; post 72 hours of infection (hpi), P1 virus-containing tissue

culture supernatants were collected and stored at -80°C. Following titration, P1 virus stock was

used to generate a P2 stock by infecting Vero E6 TMPRSS2 monolayers with multiplicity of

infection (MOI) of 0.0001 for 72 hours. P2 virus was passaged again in Vero E6-TMPRSS2 cells to

obtain P3 stock. Viral titers were determined by standard plaque assay on Vero E6 cells.

Infection of cells by SARS-CoV-2

Vero E6 cells previously cultured in 8-well μ-slides were infected in the BSL3 facility with SARS-
CoV-2 WA 1 (USA212 WA1/2020) in triplicates (MOI=0.5 SARS-CoV-2 WA1 (P3)) at an MOI of 2 for
6 hpi and MOI of 0.2 for 24 hpi. After 6 and 24 hrs of incubation, cells were washed with PBS and
fixed by 4% PFA (Electron Microscopy Sciences #15710) and 0.1% glutaraldehyde (Electron
Microscopy Sciences #16350) in PBS for 1 hour and removed from BSL3 for further processing. All
work involving viral stock preparation and infection using WT SARS-CoV-2 was conducted at the
high containment BSL3 facility of Stanford University according to CDC and institutional
guidelines. All the experiments were performed using a P3 SARS-CoV-2 USA-WA1/2020,
containing 100% WT population with no deletion in the spike multi-basic cleavage site.

Synthesis of the RNA FISH probes

vgRNA FISH probes targeting the ORF1a region of SARS-CoV-2[17] were ordered with 5AmMC6
modifications from Integrated DNA Technologies, Inc. in plate format of 25 nmol scale with
standard desalting. Each probe was dissolved in water to a final concentration of 100 μM. The
same set of probes was combined with equal volumes of each probe to get a stock of 100 μM
mixed probes. The mixed probes were further desalted using ethanol precipitation. Briefly, 120
μL 100 μM probes were mixed with 12 μL 3 M sodium acetate (pH 5.5), followed by 400 μL
ethanol. After precipitation at -80C overnight, probes were pelleted through centrifugation at
12,000x g for 10 min at 4°C, washed with precooled 70% (vol./vol.) ethanol three times, air dried,
and dissolved in water to make a 100 μM solution of probes. Then, 18 μL 100 μM probes were
mixed with 2 μL 1 M NaHCO$_3$ (pH 8.5), followed by 100 μg Alexa Fluor^{TM} 647 succinimidyl ester
(NHS) (Invitrogen, A37573) or CF568 succinimidyl ester (NHS) (Biotium, 92131) dissolved in 2 μL
dry DMSO (Invitrogen, D12345). The mixture was incubated for 3 days at 37C in the dark for
conjugation and purified for 3 rounds using Monarch PCR & DNA Cleanup Kit (5 μg) (NEB, T1030S)
following the manufacturer's instructions. The estimated labeling efficiency of probes was
calculated using the following equation:

523
$$Modification\ ratio = \frac{20}{(A_{base} \times \varepsilon_{dye}) / (A_{dye} \times \varepsilon_{base})}$$

524 where ε_{dye} is 239,000 cm^{-1}M^{-1}, ε_{base} is 8,919 cm^{-1}M^{-1}, A_{base} is the absorbance of the nucleic acid at

525 260 nm, and A_{dye} is the absorbance of the dye at 650 nm. For the probes labeled with CF568, ε_{dye}

526 is 100,000 cm^{-1}M^{-1}, ε_{base} is 8,919 cm^{-1}M^{-1}, A_{base} is the absorbance of the nucleic acid at 260 nm,

527 and A_{dye} is the absorbance of the dye at 562 nm.

529 *RNA FISH, immunofluorescence (IF), and CellMask staining*

530 Fixed cells from BLS3 as described above were washed twice with a freshly prepared 0.1% NaBH$_4$

531 solution at room temperature for 5 min, and washed with PBS three times. For staining without

532 CellMask (Thermo Fisher, C10046), cells were permeabilized in 70% ethanol at 4°C overnight. For

533 CellMask staining, cells were permeabilized in 0.1% Triton X-100 at room temperature for 30 min.

534 For RNA FISH staining, permeabilized cells were washed with 200 µL Wash Buffer A [40 µL Stellaris

535 RNA FISH Wash Buffer A (LGC Biosearch Technologies, SMF-WA1-60), 20 µL deionized formamide,

536 140 µL H2O] at room temperature for 5 min, and incubated with 110 µL Hybridization Buffer [99

537 µL Stellaris RNA FISH Hybridization Buffer (LGC Biosearch Technologies, SMF-HB1-10), 11 µL

538 deionized formamide] containing 1.1 µL 12.5 µM vgRNA FISH probes for 4 hours at 37°C in the

539 dark. Then cells were washed with Wash Buffer A for 30 min at 37°C in the dark, washed with

540 Wash Buffer A containing DAPI for 30 min at 37°C in the dark, and stored in Wash Buffer B (LGC

541 Biosearch Technologies, SMF-WB1-20) for imaging. DAPI was only added to the samples for

542 confocal imaging and not added to the samples for SR imaging.

543 For IF staining with antibodies, permeabilized cells were washed with PBS twice, incubated with

544 3% BSA in PBS at room temperature for 30 min, and incubated with primary antibodies in PBS at

545 37°C for 1 hour. After incubation with primary antibodies, cells were washed twice with PBST

546 buffer (0.1% Tween-20 in PBS) at room temperature for 5 min, washed with PBS once, incubated

547 with secondary antibodies in PBS at room temperature for 30 min, washed with PBST buffer three

548 times at room temperature for 5 min, and stored in PBS for imaging.

549 For simultaneous RNA FISH and IF staining, permeabilized cells were washed with 200 µL Wash

550 Buffer A at room temperature for 5 min, and incubated with 110 µL Hybridization Buffer (99 µL

Stellaris RNA FISH Hybridization Buffer, 11 µL deionized formamide) containing 1.1 µL 12.5 µM

vgRNA FISH probes, 1 U/µL RNase inhibitor (NxGen, F83923-1), and primary antibodies for 4 hours

at 37°C in the dark. Then cells were washed with 2xSSC buffer once, washed with Wash Buffer A

containing secondary antibodies for 30 min at 37°C in the dark, washed with Wash Buffer A for

30 min at 37°C in the dark, washed with Wash Buffer B once, and stored in Wash Buffer B for

imaging. For CellMask staining, several more steps were performed from here. Cells were washed

with PBS once, stained with 1:20k CellMask and 1 U/µL RNase inhibitor in PBS for 20 min at room

temperature in the dark, and washed with PBS three times before imaging.

RNA FISH and IF staining of purified virions

8-well µ-slides (ibidi, 80827-90) were first treated with poly-D-lysine solution (Thermo Fisher,

A3890401) at 4°C overnight. Then in the BSL3 facility, the poly-D-lysine solution was removed and

150 µL SARS-CoV-2 WA1 (P3) virus solution of titer 1.82×10^5 PFU/mL was added into one well of

poly-D-lysine-treated 8-well µ-slides for incubation at 4°C for 24 hours to coat the virions onto

the surface of the well. After incubation, the medium containing virions was removed and the

well was washed with PBS twice. Virions on the surface of the well were fixed with 4% PFA in PBS

for 1 hour at room temperature and the sample was removed from BSL3. The sample was washed

twice with a freshly prepared 0.1% $NaBH_4$ solution at room temperature for 5 min, and then

washed with PBS three times. The fixed virions were permeabilized in 70% ethanol at 4°C

overnight and washed with PBS twice. For the group with Proteinase K digestion, virions were

incubated with 0.2 mg/mL Proteinase K (NEB #P8107S) in 120 µL PBS at 37°C for 30 min and

washed with PBST buffer three times. Virions were washed with Wash Buffer A once and

incubated with 110 µL Hybridization Buffer (99 µL Stellaris RNA FISH Hybridization Buffer, 11 µL

deionized formamide) containing 1.1 µL 12.5 µM vgRNA FISH probes, 1 U/µL RNase inhibitor, and

primary antibodies for 4 hours at 37°C in the dark. Then virions were washed with 2xSSC buffer

once, washed with Wash Buffer A containing secondary antibodies for 30 min at 37°C in the dark,

washed with Wash Buffer A for 30 min at 37°C in the dark, washed with Wash Buffer B once, and

stored in Wash Buffer B for imaging.

Spinning disk confocal microscopy

Confocal microscopy was performed at the Stanford University Cell Sciences Imaging Core Facility with a Nikon TiE inverted spinning disk confocal microscope (SDCM) equipped with a Photometrics Prime 95B camera, a CSU-X1 confocal scanner unit with microlenses, and 405 nm, 488 nm, 561 nm, and 642 nm lasers, using the 60x/1.27 NA PLAN APO IR water immersion objective. Images were taken using NIS Elements software version 4.60 with Z stacks at 0.3 μm steps. The camera pixel size of SDCM is 0.183 μm/pixel and the pinhole size is 50 μm. Only one Z slice is used for all images shown.

Analysis of confocal data

To extract the intensity of vgRNA, dsRNA and RdRp in each infected cell (Extended Data Fig. 2), the summation projection of each z stack was created by Fiji[53]. The intensity of each target species in each cell was measured by Fiji, subtracting the background of the same color channel. The infected cells were characterized manually into three types based on the morphology of vgRNA. Type 1 shows scattered dot-like localization of vgRNA. Type 3 shows large clustered vgRNA. Type 2 contains features of both type 1 and type 3.

Optimization of antibody concentrations

We optimized the concentration of antibodies in this study by quantifying their signal-to-background ratio (SBR), where the signal is the brightness of the IF labelling in the cells that express the given target (virus-infected sample or cells expressing Sec61β-GFP), and the background is the brightness in the negative control cells (not-infected or WT cells).

To optimize the concentration of primary antibodies against the viral targets, different concentrations of the primary antibody were applied to stain Vero E6 cells in SARS-CoV-2-infected and not-infected samples under a constant secondary antibody concentration (Supplementary

Fig. S2). To optimize the concentration of secondary antibodies, different concentrations of the secondary antibody were applied to stain Vero E6 cells in infected (virus+) and not-infected (virus−) samples under a constant primary antibody concentration (Supplementary Fig. S3). For each cell, a 11 pixel x 11 pixel box was drawn in the region with brightest signal in the cell and the mean intensity within that region was measured to represent the intensity of target antibody in that cell. The SBR was calculated, after subtraction of the dark signal I_{dark}, using the following equation:

$$SBR = \frac{< I_{virus+} - I_{dark} >}{< I_{virus-} - I_{dark} >}$$

To optimize the concentration of the anti-GFP antibodies, different concentrations of primary antibody were applied to stain Vero E6 Sec61B-GFP cells and WT Vero E6 cells under a constant secondary antibody concentration (Supplementary Fig. S2). For each cell, a 11 pixel x 11 pixel box was drawn in the region with the brightest signal in the cell and the mean intensities of both the GFP and the antibody signals within that region were measured after subtraction of the dark signals. To account for the variable expression levels among different cells, the IF signal I_{IF} was normalized by the GFP signal I_{GFP} within the given region. The SBR was calculated using the following equation:

$$SBR = \frac{< I_{IF,Sec61B\text{-}GFP}/I_{GFP,Sec61B\text{-}GFP} >}{< I_{IF,WT}/I_{GFP,WT} >}$$

For the primary antibodies against GFP, nsp3, nucleocapsid, nsp12 and for the secondary antibody for the dsRNA labelling, we chose the antibody concentration that produces the highest SBR as the optimal concentration. For the primary antibodies against spike S2 and dsRNA and for the secondary antibody for the spike S2 labelling, we chose the concentration that yields the second highest SBR because it provides a significantly lower non-specific background with only a minor decrease of the estimated SBR.

Optical setup for SR microscopy

(d)STORM SR microscopy was performed on a custom-built system (Supplementary Fig. S4), consisting of a Nikon Diaphot 200 inverted microscope frame with an oil-immersion objective 60x/1.35 NA (Olympus UPLSAPO60XO) and a Si EMCCD camera (Andor iXon Ultra 897). We used 642 nm and 560 nm 1W continuous-wave (CW) lasers (MPB Communications Inc.) for excitation of AF647 or CellMask and CF568, accordingly. For reactivation of fluorophores from the dark state we used a 405 nm 50 mW CW diode laser (Coherent OBIS). All laser beams were expanded and co-aligned in free space and coupled into a square-core multi-mode fiber with a shaker for speckle reduction (Newport F-DS-ASQR200-FC/PC). The output tip of the fiber (200 x 200 μm^2 core size) was imaged with a 10x/0.25 NA objective and magnified to achieve a square illumination region of 47.6 x 47.6 μm^2 with a constant intensity in the sample image plane of the main objective. The fluorescence was split from the excitation light with a multi-band dichroic mirror (ZT405/488/561/640rpcv2, Chroma) and filtered with dichroic filters (ZET635NF, ZET561NF, T690LPxxr, all Chroma). The fluorescence of AF647 and CellMask was additionally filtered with a band-pass filter (ET685/70M, Chroma) and that of CF568 with a combination of 561LP and 607/70BP (Semrock, EdgeBasic and BrightLine). The sample image was focused with a tube lens (f = 400 mm) on the EMCCD camera, providing a pixel size of 117 x 117 nm^2 in sample coordinates.

Axial drift was compensated with a custom Focus Lock system[54]. We used an 808 nm fiber-coupled diode laser (Thorlabs S1FC808) whose output fiber tip was conjugated with the back focal plane of the imaging objective, allowing changing the angle of this beam out of the objective by translating the fiber tip (Supplementary Fig. S4). This inclined beam was partially reflected from the coverslip-water interface and the reflected beam was focused with a cylindrical lens onto a CMOS sensor (UI-3240CP-NIR, IDS Imaging). The 808 nm beam was aligned such that the image of the reflected beam would shift laterally when the axial position of the sample changes. The sample was mounted on two stacked piezo stages (U-780.DOS for coarse and P-545.3C8S for fine movement, both Physik Instrumente). The position of the reflected beam image was recorded when the sample was set at the desired Z position for imaging. During imaging, the Z-position of the fine stage was directed to move proportionally to the shift of the reflected beam image from

the recorded position, compensating for Z-drift. The Focus Lock control code was programmed in

Matlab (MathWorks, Inc.).

SR imaging procedure

For (d)STORM, the sample chamber was filled with 300 µl of a photoblinking buffer consisting of

200 U/ml glucose oxidase, 1000 U/ml catalase, 10% w/v glucose, 200 mM Tris-HCl pH 8.0, 15 mM

NaCl and 50 mM cysteamine. The buffer was prepared using the following stock solutions[38]: 1)

4 kU/ml glucose oxidase (G2133, Sigma), 20 kU/ml catalase (C1345, Sigma), 25 mM KCl (P217,

Fisher), 4 mM TCEP (646547, Sigma), 50% v/v glycerol (BP229, Fisher) and 22 mM Tris-HCl pH 7.0

(BP1756, Fisher), stored at −20 °C; 2) 1 M cysteamine-HCl (30080, Sigma), stored at −20 °C; 3) 37%

w/v glucose (49139, Sigma) with 56 mM NaCl (S271, Fisher) and 0.74 M Tris-HCl pH 8.0

(J22638.AE, Fisher), stored at +4 °C. For samples with RNA FISH labelling, the buffer was

supplemented with 1 U/µl of an RNase inhibitor (302811, LGC Biosearch Technologies).

The SR imaging started with a DL image of cells from each fluorophore at a low power (*e.g.*, 2

W/cm^2). For (d)STORM acquisitions, we began with AF647 or CellMask, followed by CF568. We

used an excitation power density of ~20 kW/cm^2 for shelving and blinking of CF568 and ~6-20

kW/cm^2 for AF647. The power density of the 405 nm illumination for both dyes was increased

from 0 to 50 W/cm^2 throughout an acquisition to keep the reactivation rate approximately

constant. The exposure time was 10.57 ms per frame and the calibrated EM gain was 43. The

image recording started after the initial shelving phase upon observation of clear SM blinking; the

blinking movies were acquired for approximately 60000 frames for each fluorophore.

SR data analysis

SM movies were processed with the ThunderStorm plugin[55] for Fiji. First, the images were filtered

with a wavelet filter with a b-spline order of 3 and a scale of 2. The coarse localizations were

found as local maxima with an 8-neighborhood connectivity and a threshold of 2·std(Wave.F1).

These localizations were weighted least squares-fitted with the integrated Gaussian model using

a radius of 4 pixels and an initial sigma of 1.1. Then, we performed drift correction estimated by

cross-correlation between successive subsets of localizations. For further processing, we kept

only localizations with fitted sigma between 160 nm and 80 nm.

For image registration, we imaged 200 nm TetraSpeck beads (T7280, Thermo Fisher Scientific) in

both channels, whose images were processed similarly to the SM movies. The transformation

between the channels was calculated using an affine transformation with help of Matlab function

'fitgeotrans'. The calculated transformation was then applied to the CF568 localizations using a

Matlab function 'transformPointsInverse'.

Localizations found within 50 nm on consecutive frames that could originate from multiple

localizations of a single molecule were treated in two ways. For SR images, these localizations

were refined to suppress overcounting by selecting them from a normal distribution with a mean

at the weighted mean of the initial localizations and a standard deviation (SD) that equals

$120 \cdot (N_{ph})^{-1/2}$ nm, where N_{ph} is the total number of photons acquired from all localizations in the

given consecutive series[38]. For data analysis other than SR image reconstruction, the localizations

of the consecutive series were reduced to a single localization at the weighted mean position.

The weights of localizations were proportional to the photon counts of these individual

localizations. SR images were reconstructed as 2D histograms with a bin size of 20 x 20 nm^2.

However, SR images where one of the channels contained the CellMask labelling had a bin size of

30 x 30 nm^2. SR images acquired with CellMask were additionally filtered with a Gaussian filter

with σ = 0.5 pixels.

Cluster analysis with BIC-GMM

Gaussian Mixture Models (GMM) implemented in Python were fitted to vgRNA and dsRNA

localization datasets, yielding a representation of localization densities as a collection of

potentially elliptical and/or rotated 2D Gaussians. The number of components most suitable for

each field of view was determined using an iterative grid search, evaluating 4 candidate GMMs

using the Bayesian Information Criterion (BIC)[56]. The first grid iteration tested [1, 2500]

components with test points $t_i = \{1, 834, 1667, 2500\}$, where i denotes the index in the set such

that $t_0 = 1$. For each iteration of the grid search, the model with the lowest BIC was selected the

best candidate, t_k, and the next iteration of the grid was narrowed, to be bounded by $[t_{max\,(k-1,\,0)}$

$+ 1, t_{min\,(k+1,\,3)} - 1]$, until the stride of the grid was 1 component, or the test point with the best BIC

was on a rail ($k = 0$ or 3). To reduce memory requirements, this GMM optimization was performed

on a random subset of up to 200,000 localizations from each data set, but the optimized GMM

was then used to predict a component assignment for all original localizations. These components

were regarded as clusters, and refined by removing localizations with a log probability of being

an event from their assigned Gaussian component of less than −25. The radius of gyration, Rg,

was then calculated for each cluster, and the number of localizations in each cluster, N_{loc}, was

used to approximate a cluster density as $\delta = N_{loc} / (\pi \cdot Rg^2)$. Clusters with δ below a threshold of

0.008 localizations/nm^2 for dsDNA, or below an ROI-dependent threshold between 0.005 and

0.013 localizations/nm^2 for vgRNA, were removed from further quantification as sparse

background. This analysis and resulting visualizations were carried out in the PYthon Microscopy

Environment (https://doi.org/10.5281/zenodo.4289803)[57], using a plugin

(github.com/barentine/bic-gmm) and the scikit-learn GMM implementation[58].

Counting of vgRNA molecules in the clusters

The number of vgRNA molecules in a vgRNA cluster was defined as a quotient between the

number of vgRNA-FISH localizations in the cluster and the average number of localizations

produced by a single FISH-labelled vgRNA molecule in the given cell. The average number of

localizations per vgRNA molecule was estimated from isolated nanoscale vgRNA puncta in the

cytoplasm (Supplementary Fig. 3a). This number was defined as the median of the number of

localizations within 50 nm from each localization in the region with vgRNA puncta. The estimated

number of vgRNA molecules was calculated for every cluster determined by the BIC-GMM cluster

analysis and the median value per cell was shown in a chart (Supplementary Fig. 3b-c).

Counting of nsp12 puncta in the vgRNA clusters

The center of nsp12 puncta is obtained by fitting the SR images in ThunderStorm[55]. The SR localizations of nsp12 were first converted into a 2D histogram image with a bin size of 20 x 20 nm^2. The approximate localization of the center was found as a centroid of connected components with a threshold of 5·std(Wave.F1) without filter. These localizations were least squares-fitted with the integrated Gaussian model using a fitting radius of 2 pixels and an initial sigma of 0.4. We next removed duplicates among localizations within a 20 nm radius. The puncta whose sigma were smaller than 5 nm were further filtered out to avoid localizing single-pixel-sized background localizations. For each vgRNA cluster with its center and the radius of gyration (Rg) determined using BIC-GMM, we counted the number of nsp12 puncta within a 1.5·Rg distance of the center of the vgRNA cluster. For nsp12 puncta found within the cutoff distance of more than one vgRNA cluster, we assigned them to their closest cluster based on the relative distance d/Rg, with d being the distance between the center of the vgRNA cluster and center of the nsp12 punctum.

Bivariate pair-correlation functions

For calculation of bivariate pair-correlation functions[20] $g_{12}(r)$, we first manually selected the cytoplasmic regions with dense vgRNA clusters. The pair-correlation functions were calculated by counting the number of localizations of the second species within a distance between r and r+dr from each localization of the first species. These were normalized by dividing the number of localizations by the area of the corresponding ring of radii r and r+dr and by the average density of the second species in the region. Finally, the obtained numbers were averaged across the localizations of the first species. r was scanned over the range between 0 and 500 nm and dr was set to 1 nm. For the complete spatial randomness (CSR) case, a test CSR dataset was generated with the same average density as for the experimental case across the same ROI. $g_{12}(r)$ traces were calculated from these CSR datasets as described above. No edge effect correction was performed leading to a slight decrease of $g_{12}(r)$ at large r. Plots in the figures display experimental

and CSR $g_{12}(r)$ for each analyzed cell as faint lines as well as the mean $g_{12}(r)$ calculated from all

cells in bold lines.

Estimation of RNA FISH labelling efficiency in virions

Dye molecules inside virions were counted using fluorescence bleaching with SM calibration.

Virions attached to the coverslip were labelled using the RNA-FISH+IF protocol with PFA-only

fixation. The density of virions was around 0.5 μm^{-2} insuring observation of most virions as single

DL spots without overlap (Extended Data Fig. 1a, d). vgRNA was FISH-labelled with AF647 and

spike protein was IF-stained with CF568. Glass-bottom chambers with virions were kept in PBS for

this experiment. Samples were illuminated with 642 nm light at 20 W/cm² and were imaged with

an exposure time of 200 ms and an EM gain of 43 until bleaching of all AF647 in the imaging

region (around 200 s). A separate DL image of spike was taken with 560 nm excitation. The AF647

bleaching movies were processed in ThunderStorm using a wavelet filter with a b-spline order of

3 and a scale of 2, a local maximum approximate localization with a threshold of 1.2·std(Wave.F1)

and an 8-neighborhood connectivity. These localizations were weighted least squares-fitted with

the integrated Gaussian model using a radius of 3 pixels and an initial sigma of 1.1. Then, we kept

only localizations with sigma < 160 nm & sigma > 80 nm and removed duplicates within 300 nm

on each frame.

Further processing was done in Matlab with a custom script. We considered only vgRNA-AF647

localizations that had a spike-CF568 signal within 200 nm to avoid counting AF647 molecules

outside virions. The bleaching time traces (Extended Data Fig. 1c, f) were found by searching in

consecutive frames within 200 nm of the localization from the first frame and allowing up to 5

empty frames between frames with detections. The number of bleaching steps was defined as

the rounded quotient between the initial and the final brightness of a spot in a time trace serving

as the SM calibration. For each bleaching trace, the initial brightness (in photons) was defined as

the median value of the brightness in the first 4 localizations and the final brightness as the

median brightness value of the last 4 localizations. If the trace contained only 7-8 detections, the

range for the initial and the final brightness was reduced to 3 frames; for traces with 5-6

detections, this was reduced to 2; for traces with 3-4 frames – to 1; for traces containing only 1

or 2 detections, the number of bleaching steps was set to 1. For each analyzed region containing

around 200 bleaching traces, the number of bleaching steps was fitted with a zero-truncated

Poisson distribution (Extended Data Fig. 1g-h). The expected values ± SD obtained from the fit of

5 regions for each of not-treated and PK-treated cells are shown in a chart (Extended Data Fig.

1i).

Author contribution

L.A., M.H., L.S.Q. and W.E.M. conceived the project. L.A. designed the optical set-up, performed

the SR acquisitions and data analysis. M.H. performed cell culture, labelling and confocal imaging.

Y.Z. performed confocal and SR data analysis and helped with sample preparation and confocal

imaging. J.G. performed SARS-CoV-2 infection experiments at the BSL-3 facility with staff listed in

the Acknowledgements. A.R.R. contributed to the concept and SR experiments at the early stages

of the project. A.E.S.B. designed the BIC-GMM cluster analysis method and contributed to the

optical set-up design. L.A and W.E.M. wrote the manuscript with input from all authors.

Acknowledgements

We thank Puja Patel and Amol Pohane for their assistance with the cell culturing in the BSL3

facility. We thank Leiping Zeng for sample preparation and discussion about the results and

experimental plan with the other authors. This work was supported in part by the National

Institute of General Medical Sciences Grant Nos. R35GM118067 (to W.E.M.) and the National

Institutes of Health Common Fund 4D Nucleome Program No. U01 DK127405 (to L.S.Q.). We also

acknowledge Stanford University Cell Sciences Imaging Core Facility (RRID:SCR_017787). L.S.Q. is

a Chan Zuckerberg Biohub Investigator, and W.E.M. is a Sarafan ChEM-H Fellow.

References

1. Schmidt, N. *et al.* The SARS-CoV-2 RNA–protein interactome in infected human cells. *Nature Microbiology* **6**, 339-353 (2021).
2. Flynn, R.A. *et al.* Discovery and functional interrogation of SARS-CoV-2 RNA-host protein interactions. *Cell* **184**, 2394-2411.e2316 (2021).
3. Knoops, K. *et al.* SARS-coronavirus replication is supported by a reticulovesicular network of modified endoplasmic reticulum. *PLoS Biol* **6**, e226 (2008).
4. Snijder, E.J. *et al.* A unifying structural and functional model of the coronavirus replication organelle: Tracking down RNA synthesis. *PLoS Biol* **18**, e3000715 (2020).
5. Klein, S. *et al.* SARS-CoV-2 structure and replication characterized by in situ cryo-electron tomography. *Nat Commun* **11**, 5885 (2020).
6. Betzig, E. *et al.* Imaging intracellular fluorescent proteins at nanometer resolution. *Science* **313**, 1642-1645 (2006).
7. Rust, M.J., Bates, M. & Zhuang, X. Sub-diffraction-limit imaging by stochastic optical reconstruction microscopy (STORM). *Nat. Methods* **3**, 793-796 (2006).
8. Heilemann, M. *et al.* Subdiffraction-Resolution Fluorescence Imaging with Conventional Fluorescent Probes. *Angew. Chem. Int. Ed.* **47**, 6172-6176 (2008).
9. Klar, T.A., Jakobs, S., Dyba, M., Egner, A. & Hell, S.W. Fluorescence microscopy with diffraction resolution barrier broken by stimulated emission. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 8206-8210 (2000).
10. Gustafsson, M.G.L. Surpassing the lateral resolution limit by a factor of two using structured illumination microscopy. *J. Microsc.* **198**, 82-87 (2000).
11. Xu, K., Zhong, G. & Zhuang, X. Actin, Spectrin, and Associated Proteins Form a Periodic Cytoskeletal Structure in Axons. *Science* **339**, 452-456 (2013).
12. Andronov, L., Ouararhni, K., Stoll, I., Klaholz, B.P. & Hamiche, A. CENP-A nucleosome clusters form rosette-like structures around HJURP during G1. *Nature Communications* **10**, 4436 (2019).
13. Reinhardt, S.C.M. *et al.* Ångström-resolution fluorescence microscopy. *Nature* **617**, 711-716 (2023).
14. Baddeley, D. & Bewersdorf, J. Biological Insight from Super-Resolution Microscopy: What We Can Learn from Localization-Based Images. *Annual Review of Biochemistry* **87**, 965-989 (2018).
15. Wang, J. *et al.* Multi-color super-resolution imaging to study human coronavirus RNA during cellular infection. *Cell Reports Methods* **2**, 100170 (2022).
16. Wolff, G. *et al.* A molecular pore spans the double membrane of the coronavirus replication organelle. *Science* **369**, 1395-1398 (2020).
17. Lee, J.Y. *et al.* Absolute quantitation of individual SARS-CoV-2 RNA molecules provides a new paradigm for infection dynamics and variant differences. *eLife* **11**, e74153 (2022).
18. Pepe, A., Pietropaoli, S., Vos, M., Barba-Spaeth, G. & Zurzolo, C. Tunneling nanotubes provide a route for SARS-CoV-2 spreading. *Science Advances* **8**, eabo0171 (2022).
19. Sherman, E. *et al.* Functional Nanoscale Organization of Signaling Molecules Downstream of the T Cell Antigen Receptor. *Immunity* **35**, 705-720 (2011).
20. Razvag, Y., Neve-Oz, Y., Sajman, J., Reches, M. & Sherman, E. Nanoscale kinetic segregation of TCR and CD45 in engaged microvilli facilitates early T cell activation. *Nature Communications* **9**, 732 (2018).

21.   Sawicki, S.G. & Sawicki, D.L. Coronavirus minus-strand RNA synthesis and effect of cycloheximide
on coronavirus RNA synthesis. *Journal of Virology* **57**, 328-334 (1986).

22.   Hillen, H.S. *et al.* Structure of replicating SARS-CoV-2 polymerase. *Nature* **584**, 154-156 (2020).

23.   Gao, Y. *et al.* Structure of the RNA-dependent RNA polymerase from COVID-19 virus. *Science* **368**,
779-782 (2020).

24.   Ahn, D.-G., Choi, J.-K., Taylor, D.R. & Oh, J.-W. Biochemical characterization of a recombinant SARS
coronavirus nsp12 RNA-dependent RNA polymerase capable of copying viral RNA templates.
Archives of Virology **157**, 2095-2104 (2012).

25.   Naydenova, K. *et al.* Structure of the SARS-CoV-2 RNA-dependent RNA polymerase in the presence
of favipiravir-RTP. *Proceedings of the National Academy of Sciences* **118**, e2021946118 (2021).

26.   Eymieux, S. *et al.* Ultrastructural modifications induced by SARS-CoV-2 in Vero cells: a kinetic
analysis of viral factory formation, viral particle morphogenesis and virion release. *Cellular and*
Molecular Life Sciences **78**, 3565-3576 (2021).

27.   Cortese, M. *et al.* Integrative Imaging Reveals SARS-CoV-2-Induced Reshaping of Subcellular
Morphologies. *Cell Host & Microbe* **28**, 853-866.e855 (2020).

28.   Lang, S. *et al.* An Update on Sec61 Channel Functions, Mechanisms, and Related Diseases. *Front*
Physiol **8**, 887 (2017).

29.   Hsu, J.C.-C., Laurent-Rolle, M., Pawlak, J.B., Wilen, C.B. & Cresswell, P. Translational shutdown and
evasion of the innate immune response by SARS-CoV-2 NSP14 protein. *Proceedings of the National*
Academy of Sciences **118**, e2101161118 (2021).

30.   Mönkemöller, V. *et al.* Imaging fenestrations in liver sinusoidal endothelial cells by optical
localization microscopy. *Physical Chemistry Chemical Physics* **16**, 12576-12581 (2014).

31.   Oudshoorn, D. *et al.* Expression and Cleavage of Middle East Respiratory Syndrome Coronavirus
nsp3-4 Polyprotein Induce the Formation of Double-Membrane Vesicles That Mimic Those
Associated with Coronaviral RNA Replication. *mBio* **8**, 10.1128/mbio.01658-01617 (2017).

32.   Liv, Z. *et al.* SARS-CoV-2 nsp3-4 suffice to form a pore shaping replication organelles. *bioRxiv*,
2022.2010.2021.513196 (2022).

33.   Han, Y.-S. *et al.* Papain-Like Protease 2 (PLP2) from Severe Acute Respiratory Syndrome
Coronavirus (SARS-CoV): Expression, Purification, Characterization, and Inhibition. *Biochemistry*
44, 10349-10359 (2005).

34.   Shin, D. *et al.* Papain-like protease regulates SARS-CoV-2 viral spread and innate immunity. *Nature*
587, 657-662 (2020).

35.   Hartenian, E. *et al.* The molecular virology of coronaviruses. *J Biol Chem* **295**, 12910-12934 (2020).

36.   V'Kovski, P., Kratzel, A., Steiner, S., Stalder, H. & Thiel, V. Coronavirus biology and replication:
implications for SARS-CoV-2. *Nat Rev Microbiol* **19**, 155-170 (2021).

37.   Baggen, J., Vanstreels, E., Jansen, S. & Daelemans, D. Cellular host factors for SARS-CoV-2 infection.
Nature Microbiology **6**, 1219-1232 (2021).

38.   Andronov, L., Genthial, R., Hentsch, D. & Klaholz, B.P. splitSMLM, a spectral demixing method for
high-precision multi-color localization microscopy applied to nuclear pore complexes.
Communications Biology **5**, 1100 (2022).

39.   Storti, B. *et al.* A spatial multi-scale fluorescence microscopy toolbox discloses entry checkpoints
of SARS-CoV-2 variants in Vero E6 cells. *Computational and Structural Biotechnology Journal* **19**,
6140-6156 (2021).

40.   Scherer, K.M. *et al.* SARS-CoV-2 nucleocapsid protein adheres to replication organelles before viral
assembly at the Golgi/ERGIC and lysosome-mediated egress. *Science Advances* **8**, eabl4895 (2022).

41.   Cao, C. *et al.* The architecture of the SARS-CoV-2 RNA genome inside virion. *Nature*
Communications **12**, 3917 (2021).

42.   Lan, T.C.T. *et al.* Secondary structural ensembles of the SARS-CoV-2 RNA genome in infected cells.
Nature Communications **13**, 1128 (2022).

43.   Gopal, A., Zhou, Z.H., Knobler, C.M. & Gelbart, W.M. Visualizing large RNA molecules in solution.
RNA **18**, 284-299 (2012).

44.   Schonborn, J. *et al.* Monoclonal antibodies to double-stranded RNA as probes of RNA structure in
crude nucleic acid extracts. *Nucleic Acids Res* **19**, 2993-3000 (1991).

45.   Ku, J. *et al.* Reactive Polymer Targeting dsRNA as Universal Virus Detection Platform with Enhanced
Sensitivity. *Biomacromolecules* **21**, 2440-2454 (2020).

46.   Deng, J. *et al.* SARS-CoV-2 NSP8 suppresses type I and III IFN responses by modulating the RIG-
I/MDA5, TRIF, and STING signaling pathways. *Journal of Medical Virology* **95**, e28680 (2023).

47.   Kumar, P. *et al.* The nonstructural protein 8 (nsp8) of the SARS coronavirus interacts with its ORF6
accessory protein. *Virology* **366**, 293-303 (2007).

48.   Snijder, E.J. *et al.* Ultrastructure and origin of membrane vesicles associated with the severe acute
respiratory syndrome coronavirus replication complex. *J Virol* **80**, 5927-5940 (2006).

49.   Stertz, S. *et al.* The intracellular sites of early replication and budding of SARS-coronavirus. *Virology*
361, 304-315 (2007).

50.   Ulasli, M., Verheije, M.H., de Haan, C.A.M. & Reggiori, F. Qualitative and quantitative
ultrastructural analysis of the membrane rearrangements induced by coronavirus. *Cellular*
Microbiology **12**, 844-861 (2010).

51.   Chiem, K., Ye, C. & Martinez-Sobrido, L. Generation of Recombinant SARS-CoV-2 Using a Bacterial
Artificial Chromosome. *Current Protocols in Microbiology* **59**, e126 (2020).

52.   Ye, C. *et al.* Rescue of SARS-CoV-2 from a Single Bacterial Artificial Chromosome. *mBio* **11**,
10.1128/mbio.02168-02120 (2020).

53.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature methods* **9**,
676-682 (2012).

54.   Barentine, A.E.S. *et al.* An integrated platform for high-throughput nanoscopy. *Nature*
Biotechnology (2023).

55.   Ovesny, M., Krizek, P., Borkovec, J., Svindrych, Z. & Hagen, G.M. ThunderSTORM: a comprehensive
ImageJ plug-in for PALM and STORM data analysis and super-resolution imaging. *Bioinformatics*
(Oxford, England) **30**, 2389-2390 (2014).

56.   Schwarz, G. Estimating the dimension of a model. *The annals of statistics* **6**, 461-464 (1978).

57.   Marin, Z. *et al.* PYMEVisualize: an open-source tool for exploring 3D super-resolution data. *Nature*
Methods **18**, 582-584 (2021).

58.   Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830
(2011).

Extended Data Figures

Extended Data Fig. 1. Validation of the labelling and imaging approach.

a, DL image of SARS-CoV-2 virions where vgRNA was labelled with AF647 by RNA FISH and the spike proteins were labelled by primary anti-spike S2 antibody with secondary CF568-conjugated antibody. **b**, Representative two-color SR images of individual virions reveal concentric localization of spike around vgRNA. **c**, Bleaching time trace of AF647 emission from a single virion (yellow arrow in **a**) demonstrates two-step bleaching. **d**, DL image of virions that were treated with Proteinase K (PK) before labelling. **e**, SR images of PK-treated virions reveal incomplete spike labelling due to digestion of proteins by the PK. **f**, Bleaching time trace of AF647 emission from a single virion (yellow arrow in **d**) shows 6-step bleaching suggesting increased vgRNA labelling efficiency in PK-treated virions. **g-h**, Histograms of the number of fluorophores per virion in

untreated (**g**) or PK-treated (**h**) samples and their fits with a Poisson distribution. **i**, Mean number of AF647 molecules per virion from the fit for 5 different regions in both untreated and PK-treated samples. p-value = $2 \cdot 10^{-8}$, two-tailed t-test. The error bar indicates mean ± SD value for the untreated and PK-treated groups. **j**, SR image of a SARS-CoV-2 infected cell with the cell body to the left reveals assembled virions at its cytoplasmic tubular projections at 24 hpi. Scale bars, 100 nm (**b**, **e**) and 1 μm (**a**, **d**, **j**).

Extended Data Fig. 2. Screening and quantification of vgRNA, dsRNA and nsp12 by confocal microscopy.

a, Representative confocal images show three types of vgRNA distribution in SARS-CoV-2 infected cells. **b**, Number of cells assigned to one of the three types at 6 or 24 hpi. **c**, Cell-integrated vgRNA signal increases significantly from 6 hpi to 24 hpi. p-value = 6·10^{-8}, two-tailed t-test. **d-e**, Cell-integrated signal of immunofluorescently detected dsRNA (**d**) and nsp12 (**e**) in SARS-CoV-2 infected cells does not significantly change from 6 hpi to 24 hpi. p-values = 0.13 (**d**) and 0.23 (**e**), two-tailed t-test. **f**, dsRNA signal correlates with vgRNA signal at 6 hpi (Pearson's r = 0.76). **g**, dsRNA signal does not correlate with vgRNA signal at 24 hpi (Pearson's r = 0.18). Error bars represent mean + SD of the values from individual cells. Scale bars, 10 μm.

1000

Extended Data Fig. 3. Estimation of the number of vgRNA molecules in vgRNA clusters.

a, SR localizations of single vgRNA molecules found in the cytoplasm of infected cells outside the

dense vgRNA clusters. On a cell by cell basis, similar images are used as a calibration for the

number of SR detections per one vgRNA molecule. **b**, The estimated number of vgRNA molecules

per cluster at 6 and 24 hpi from all analyzed cells. The histogram counts are normalized by the

number of analyzed cells; the histogram counts for 24 hpi are additionally divided by 3 to account

for the 3x wider bin size than at 6 hpi. **c**, Median estimated counts of vgRNA molecules per cluster

for each analyzed cell (individual yellow points). The error bars represent mean ± SD values of

these median vgRNA molecule counts for each time point. p-value = $5 \cdot 10^{-4}$, two-tailed t-test. Scale

bars, 50 nm x 50 nm.

1011

1012

1013

1014 **Extended Data Fig. 4. Host ER modification at 6 hpi.**

1015 SR image of vgRNA in a SARS-CoV-2 infected Vero E6 cell, stably expressing Sec61β-GFP. Altered

1016 ER forms ring-like structures that partially encapsulate vgRNA clusters in the perinuclear region.

1017 Left: green/magenta coloring; right: color scale of localizations. Scale bar, 1 μm.

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027 **Extended Data Fig. 5. Host ER modification at 24 hpi.**

1028 SR image of vgRNA in a SARS-CoV-2 infected Vero E6 cell, stably expressing Sec61β-GFP. Altered

1029 ER forms ring-like structures that encapsulate vgRNA clusters in the perinuclear region, while the

1030 Sec61β signal at the ER tubules decreases compared to 6 hpi (Extended Data Fig. 4). Scale bar, 1

1031 μm.

1032

1033

1034

1035

1036

1037 **Extended Data Fig. 6. Encapsulation of dsRNA by altered host ER at 24 hpi.**

1038 SR image of dsRNA in a SARS-CoV-2 infected Vero E6 cell, stably expressing Sec61β-GFP. Ring-like
1039 structures of altered ER encapsulate dsRNA clusters. Scale bar, 1 μm.

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

Extended Data Fig. 7. dsRNA is encapsulated into membrane-bound organelles.

a, SR image of dsRNA in a SARS-CoV-2 infected cell at 24 hpi with membranes labelled by CellMask Deep Red and dsRNA labelled with immunofluorescence. CellMask-labelled membranes can be observed around dsRNA clusters. Virions at the plasma membrane are seen as bright puncta (right side and lower right corner of the image). **b**, Zoomed-in image that corresponds to the white box in **a**. **c**, Bivariate pair-correlation functions indicate nanoscale anti-correlation between dsRNA and CellMask, consistent with dsRNA encapsulation in membrane-bound organelles at both 6 and 24 hpi. Scale bar, 1 μm (**a**) and 500 nm (**b**).

Extended Data Fig. 8. Nsp3 anti-correlates with dsRNA.

a, SR image of a SARS-CoV-2 infected cell at 6 hpi with nsp3 and dsRNA labelled by immunofluorescence. Nsp3 can be observed at the surface of isolated dsRNA clusters (white boxes & white insets) or in dense aggregates between dsRNA clusters (blue boxes & blue insets). **b**, Bivariate pair-correlation functions indicate nanoscale anti-correlation between dsRNA and nsp3 at 6 hpi. **c**, SR image of a SARS-CoV-2 infected cell at 24 hpi with nsp3 and dsRNA labelled by immunofluorescence. Nsp3 forms a network-like pattern that encapsulates dsRNA clusters. **d**, Bivariate pair-correlation functions indicate nanoscale anti-correlation between dsRNA and nsp3 at 24 hpi. Scale bars, 1 μm (**a**, **c**), 500 nm (blue insets in **a** and insets in **c**) and 200 nm (white insets in **a**).

1086

Extended Data Fig. 9. Nanoscale co-organization of viral components with host membranes.

a, SR images of nsp3, Sec61β, vgRNA and membranes (CellMask) in SARS-CoV-2 infected cells at 6 and 24 hpi. **b**, Bivariate pair-correlation functions indicate nanoscale association between nsp3 and Sec61β, nsp3 and CellMask and nanoscale anti-correlation between vgRNA and CellMask at both time points. Scale bar, 1 μm.

1092

1093

1094

Extended Data Fig. 10. Less common patterns of nanoscale nsp3 localization at 24 hpi.

a, Nsp3 forms an ER-like network that occupies a large part of the cytoplasm. **b**, Besides the common perinuclear pattern, Nsp3 is also diffusely localized throughout the whole cytoplasm. Scale bars, 1 µm.