Rediscovering the *Rete Ovarii*: a secreting auxiliary structure to the ovary

Dilara N. Anbarci[1], Jennifer McKey1[1,2], Daniel S. Levic[1], Michel Bagnat[1], and Blanche Capel[1]*

[1]Department of Cell Biology, Duke University Medical Center, Durham NC 27710; [2]Section of Developmental

Biology, Department of Pediatrics, University of Colorado Anschutz Medical Campus, Aurora CO 80045

*Corresponding author: blanche.capel@duke.edu

Keywords: rete ovarii; ovary; proteomics; SNARE complex; fluid secretion

Abstract

The *rete ovarii* (RO) is an appendage of the ovary that has been given little attention. Although the RO appears in drawings of the ovary in early versions of Gray's Anatomy, it disappeared from recent textbooks, and is often dismissed as a functionless vestige in the adult ovary. Using PAX8 immunostaining and confocal microscopy, we characterized the fetal development of the RO in the context of the ovary. The RO consists of three distinct regions that persist in adult life, the intraovarian rete (IOR), the extraovarian rete (EOR), and the connecting rete (CR). While the cells of the IOR appear to form solid cords within the ovary, the EOR rapidly develops into a convoluted tubular epithelium ending in a distal dilated tip. Cells of the EOR are ciliated and exhibit cellular trafficking capabilities. The CR, connecting the EOR to the IOR, gradually acquires tubular epithelial characteristics by birth. Using microinjections into the distal dilated tip of the EOR, we found that luminal contents flow towards the ovary. Mass spectrometry revealed that the EOR lumen contains secreted proteins potentially important for ovarian function. We show that the cells of the EOR are closely associated with vasculature and macrophages, and are contacted by neuronal projections. The direct proximity of the RO to the ovary and its integration with the extraovarian landscape suggest that it plays an important role in ovary development and homeostasis.

Introduction

The *Rete Ovarii* (RO) is an epithelial structure that is directly connected to the ovary, first described over a century ago (Waldeyer, 1870) as a multi-region structure of mesonephric origin. Despite high conservation among mammalian species including guinea pigs, cows, cats, sheep, swine, prairie deer mice, camel, dogs, monkey (Cassali et al., 2000; Kim et al., 2012), cows (Santos et al., 2012), and humans (Khan et al., 1999), previous work on the RO did not arrive at a consensus on the function of this structure. Thus, it has remained a mysterious orphan structure.

Although the RO has been proposed to be the female homologue of the rete testis (Wenzel and Odend'hal, 1985), our recent work revealed that it is more complex(McKey et al., 2022). The RO is composed of three distinct regions: (1) the extraovarian rete (EOR) which consists of columnar epithelial cells that create a single convoluted tubule structure ending in a blind distal dilated tip (DDT), (2) the connecting rete (CR) which consists of pseudo-columnar cells; and (3) the intraovarian rete (IOR) which consists of squamous epithelial cells that branch and form a fine network of thin solid cell cords approximately 1-3 cells thick (Byskov and Lintern-Moore, 1973; McKey et al., 2022). It was previously shown that the RO consists of ciliated and non-ciliated cells rich in apical microvilli and mitochondria (Czernobilsky et al., 1985). Fine granular Periodic acid-Schiff (PAS) positive material was found in the cytoplasm of the EOR and CR cells, indicating the presence of polysaccharides (e.g. glycogen and mucins) (Stein and Anderson, 1979). Because the proportion of PAS-positive material in the RO was found to be influenced by the estrous cycle in cows (Wenzel et al., 1987), researchers concluded that the luminal contents of the RO were under endocrine control (Archbald et al., 1971). Although these results strongly suggested a secretory role for the RO, this function was not experimentally confirmed, and no proteomic or metabolomic investigations were pursued. Previous characterization of the RO relied heavily on imaging serial sections to study the structure. However, these studies lacked contextual information, and the use of sections made it challenging to determine whether tubules were connected to one another or were isolated structures (Wenzel and Odend'hal, 1985; Woolnough et al., 2000). Furthermore, some investigators reported that the whole RO contracted postnatally and that the DDT of the EOR separated from the RO and degenerated (Byskov and Lintern-Moore, 1973). Perhaps for

these reasons, the RO was deemed a functionless vestige, and has been omitted from recent textbook

representations of the female reproductive tract (Girsh, 2021).

The RO was rediscovered and highlighted in our recent study (McKey et al., 2022), where we used confocal

and lightsheet imaging of whole ovaries to study the integration of ovary morphogenesis with the development

of surrounding tissues, including the RO. Recently, the IOR has gained attention as a newly described

progenitor for supporting cells of the murine gonad (Mayère et al., 2022). Recent single cell transcriptomics

studies have also identified cells of the RO within human fetal gonads (Lardenois et al., 2023; Taelman et al.,

2022). We reported that the entire RO expresses high levels of PAX8, and chose to use this as a marker to

visualize and characterize cells of the RO. In the present study, we used the Pax*8-rtTa; Tre-H2B-Gfp* RO

nuclear reporter mouse line, advanced imaging techniques, and secretome analysis to characterize the

development of the intact RO in its native context and to investigate the function and heterogeneity of its cells.

We found that the RO arises from a subset of the mesonephric tubules analogous to the rete testis and

persists into adulthood. Our studies reveal the RO as a continuous structure, surrounded by smooth muscle

actin, a dense vascular network, and several macrophage populations. We also show that the RO is directly

contacted by neurons. The enrichment of secretory machinery in RO cells, our experimental analysis of

directional flow and our analysis of luminal contents, together suggest that the RO sends material to the ovary.

Based on these findings, we suggest that the RO plays a role in ovary function and should be investigated as

an additional organ of the female reproductive tract.

Results

Characterization of the IOR, CR, and EOR

We used immunofluorescence (IF) and confocal imaging to investigate the development and sub-regional

structure of the RO from embryonic day (E) 16.5 to postnatal day (P) 7. Three distinct regions of the RO were

originally defined histologically by cell morphology (Byskov and Lintern-Moore, 1973; Lee et al., 2011). Using

IF and confocal imaging, we found that these three regions are maintained throughout development, but their

relative sizes change. While the entirety of the RO is PAX8+, we took different approaches to identify region

Figure 1. The Rete Ovarii undergoes dynamic changes during development alongside the ovary.

(a,b,c,d top row) Diagrams depicting the development of the three regions (EOR-magenta; CR-cyan; IOR-yellow) of the RO from E16.5-P7 in the whole ovarian complex (ovary-light pink; oviduct-grey). (a,b,c,d bottom rows) Maximum intensity projection from confocal Z-stacks of whole ovary/mesonephros complexes at E16.5 (a), E18.5 (b), P0 (c) and P7 (d) immunostained for PAX8 (yellow), GFRa1 (cyan) and KRT8 (magenta). a,b,c,d second row are composite images, while the third row show separate panels for KRT8 and and the bottom row show panels for GFRa1. *Note that GFRa1 and KRT8 do not co-localize and are specific to the CR and EOR, respectively. All figures are dorsal views of the ovary. Yellow asterisk indicates opening of the infundibulum for reference. Scale bar – 100um

specific markers. First, as GFRa1 was known to be expressed in the rete testis, we hypothesized that it was

expressed in the RO as well. Indeed, immunostaining using antibodies against GFRa1 revealed that part of

the RO is GFRa1+. GFRa1 specifically labeled the CR from E16.5 to P7 (Fig. 1 bottom row). To identify

additional RO markers, we performed bulk and single cell RNA sequencing (ScRNA-seq) of cells from the

ovarian complex of mice at E16.5 and 2 months, and found that *Krt8* was enriched in the RO (Anbarci et al.,

2023). We validated these findings using IF against KRT8, which revealed that KRT8+ cells were specifically

localized to the EOR (Fig. 1 third row).

To characterize RO development, we next analyzed PAX8 and KRT8 expression from E16.5 to P0. At E16.5,

the CR and IOR were the largest of the three regions. In the EOR, several tubules leading from the region of

the CR converged into a single tube of columnar PAX8+ and KRT8+ epithelial cells that lead to a blind end.

Between E16.5 and E18.5, the EOR underwent rapid expansion, and the blind end of the EOR became

dilated. This structure, which we refer to as the distal dilated tip (DDT) was best visualized from the ventral

side of the ovary (SFig. 1). By E18.5, the EOR was the largest of the three regions. At this stage the CR was

still large, but the IOR began to regress. At P0, the EOR remained the largest region, and the IOR had

regressed to the medullary region of the ovary (McKey et al., 2022). Their distinct structure and protein

expression patterns suggest that each region has a different function.

Figure 2. The EOR is highly integrated with its extra-ovarian environment suggesting multifaceted communication.

(a) Top panel is a whole ovarian complex maximum intensity projection of the confocal Z-stack at E18.5 immunostained for ENDOMUCIN (magenta) and PAX8 (cyan). Bottom panel is an optical section showing vasculature tightly surrounding the DDT of the EOR (magenta). (b) Top panel is a whole ovarian complex maximum intensity projection of the confocal Z-stack at E18.5 immunostained for aSMA (magenta) and KRT8 (cyan). Bottom panel is an optical section showing the EOR tightly ensheathed by smooth muscle (magenta). (c) Top panel is a whole ovarian complex maximum intensity projection of the confocal Z-stack at E18.5 immunostained for TUJ1 (magenta) and PAX8 (cyan). Bottom panel is an optical section showing direct contacts between the EOR and neuronal projections (magenta). (d) EOR at P0 immunostained for PAX8 (yellow), F4/80 (magenta) and LYVE-1 (cyan). Top and bottom panel are maximum intensity projection of the confocal Z-stack. Bottom panel shows the absence of LYVE1 macrophages (cyan) proximate to the EOR, in contrast to the closely associated F4/80 macrophages (magenta in top image). Arrowheads show regions devoid of LYVE1 macrophages where F4/80 macrophages are present. Scale bar – 100um

Integration of the EOR with the extra-ovarian environment

During gonadogenesis, the mesonephros is highly vascularized in both XX and XY embryos (Cool et al., 2011). In contrast, the ovary and surrounding tissue are more highly innervated than the testis (McKey et al., 2019). A surprising finding in our transcriptome analysis of the RO was the presence of a high proportion of immune cells and cells with vascular markers that co-isolated with E16.5 RO cells (Anbarci et al., 2023). Therefore, to explore the integration of the RO with its environment, we used IF to investigate the expression of endothelial marker Endomucin (Fig. 2a), mesenchyme marker smooth muscle actin (aSMA) (Fig. 2b), pan-neuronal marker TUJ1 (Fig. 2c), and macrophage markers F4/80 and LYVE1 (Fig. 2d). We found that at E18.5 the EOR was tightly surrounded by vasculature and ensheathed within a layer of aSMA+ mesenchyme. The EOR, and more specifically the DDT, was directly innervated by neurons that contact the PAX8+ epithelial cells. This multifaceted integration with the environment suggests mechanisms through which the RO receives or sends information to other tissues. We also found that the EOR is specifically associated with F4/80+ macrophages (Fig. 4d).

Flow of luminal material within the EOR

Because the EOR lumen is fluid filled, and because the EOR and CR cells are rich in proteoglycans (Byskov, 1975) we hypothesized that the EOR may function in luminal secretions. To investigate this idea,

Figure 3.Fluid moves from the EOR to the ovary.

(a,b) Schematic of dextran injections (syringe) for each group (a- EOR injections, b- ovary injections) (Oviduct-grey, ovary-light pink, IOR-yellow, CR-cyan, EOR-magenta). (Second row) Maximum intensity projection of a confocal Z-stack of whole ovarian complexes at P7, where Dextran was injected into the EOR. Presence of Dextran in the ovary shows that when dextran is injected into the EOR, it diffuses throughout the ovary (dextran-cyan, E-CADHERIN-magenta). (bottom row) Maximum intensity projection from confocal Z-stacks of whole ovary/mesonephros complexes at P7 where Dextran was injected into the ovary. Absence of Dextran signal in the EOR shows that dextran did not travel into the EOR when injected into the ovary near the IOR. Scale bar – 100um

we first chose to determine the direction of flow at P7, when the DDT of the EOR was fully dilated (Fig. 1d).

We first injected fluorescently labeled pH-insensitive Dextran into the DDT of the EOR and found that, within

just 15 min, the fluorescent fluid had readily traveled from the DDT into the ovary, where it then diffused widely

(Fig. 3). By contrast, when we injected fluorescently labeled dextran into the P7 ovary near the IOR, the

dextran remained in the ovary and did not travel to the EOR (Fig. 3). These data indicate that, at least at P7,

the fluid inside the lumen of the EOR travels towards the ovary.

Potential of the EOR and CR for fluid transfer

Figure 4. The EOR and CR acquire the potential for fluid transfer

(a,b,c,d) Cells of the CR acquire E-CADHERIN soon after birth. Maximum intensity projection from confocal Z-stacks of whole ovary/mesonephros complexes at E16.5-P7 immunostained for PAX8 (magenta) and E-CADHERIN (cyan). Arrowhead shows the region of the CR where E-CADHERIN expression is negligible at E16.5 (a), low at E18.5 (b) and fully expressed at P0 (c) and P7 (d). (e) Optical section from confocal Z-stacks of RO complexes at P0 immunostained for E-CADHERIN (yellow), CNN1 (magenta), and aSMA (cyan). Boxed regions show the absence of CNN1 in the aSMA+ sheath around the EOR at P0, and (f) its acquisition by P7, suggesting gain of contractility. (g) The RO expresses cilia marker ARL13b by P0. Maximum intensity projection from confocal Z-stacks of RO complexes at E18.5 immunostained for E-CADHERIN (cyan) and ARL13b (magenta). Outlined boxes show all regions of the EOR have ciliated cells. Scale bar – 100um (h). Airyscan optical section from confocal Z-stacks of RO complexes at E18.5 immunostained for KRT8 (cyan) and ARL13b (magenta). Scale bar – 25um

Without the obvious indication of a lumen within the CR at P7 and taking into consideration the absence of

the epithelial marker KRT8 in those cells, we wondered how fluid could travel through the CR to the ovary.

Using an antibody against E-CADHERIN, a marker of cell junctions, we found that it was specifically present

in the EOR at E16.5, but absent from the CR (Fig. 4a, second row). However, E-Cadherin expression was

gradually gained in the CR and IOR such that, by P7, the entire RO was positive for this epithelial marker

(Fig. 4a-d, second row). These results suggest that tubular epithelial connections between the EOR and IOR

mature gradually between E16.5-P7.

Next, we investigated mechanisms that could facilitate fluid movement through the RO. Previous reports

showed that ciliated and non-ciliated cells are present in the RO (Lee et al., 2011). Using an antibody against

a marker for cilia, ARL13b, we found that primary ciliated cells were abundant in the DDT and throughout the

tubules of the EOR at E18.5 (Fig. 4g-h). Due to the curvature of the DDT, optical slices often include cilia from

neighboring cells. However, an ultrathin (0.6um) optical section suggested that each cell has a single cilium.

Since the EOR was covered in a sheath of aSMA+ mesenchyme (Fig. 2b, Fig. 4e-f), we investigated if, and

when, this layer of smooth muscle became contractile. We used an antibody against the contractile smooth

muscle protein Calponin (CNN1) and found that CNN1 was absent around the EOR during fetal development

and in neonates. However, by P7, the mesenchymal sheath around the EOR gained expression of CNN1,

indicating that it acquired the ability to contract by this stage of development (Fig. 4e-f). These data suggest

that ciliary mechanosensing and/or muscle contraction may aid the directional

movement of the fluid from the DDT to the ovary.

Proteins secreted by the EOR indicate a role for the SNARE-complex

Our data showing fluid flow from the EOR to the ovary prompted us to investigate the nature and identity of

the proteins secreted by the EOR. To address this question, we analyzed the protein contents of the secreted

material using mass spectrometry. Using the *Pax8rtTA; Tre-H2B-GFP* reporter mouse line, we dissected

EORs and isolated luminal fluid by gently homogenizing the tissue with a pestle. Using this method, we

anticipated several problems. First, we expected that some cells within the EOR and surrounding tissue would

be lysed during this procedure, which would release proteins not ordinarily secreted. Second, we expected

Protein	Role
Agt	Pro-peptide for angiotensinogen
Bcam	Glycoprotein
C3	Complement component
Cd200	Glycoprotein
Cfi	Serine proteinase
Clu	**Secreted chaperone***
Cp	Metalloprotein
Cpm	Membrane-bound arginine/lysine carboxypeptidase
Epcam	Antigen
Igfbp2	Binds insulin-like growth factors I and II
Lama5	Laminin
Napsa	Pro-peptide
Sema3c	Secreted glycoprotein
Slit3	Secreted protein
Stxbp2	**Binds syntaxin***

Table 1 Cells of the EOR secrete proteins essential for vesicle transport.

Candidate list of 15 proteins identified by Mass Spectrometry. Bolded proteins were selected for validation due to role in vesicle and protein transport.

158 that some cells closely associated with the 159 EOR would be co-isolated, which could 160 result in contamination with contents from 161 cells that are not part of the EOR. To 162 eliminate intracellular proteins arising from 163 lysed cells, we cross-referenced our 164 proteomic dataset with that of the 165 mammalian secretome (Meinken et al., 166 2015), thereby retaining only known 167 secreted proteins. Next, we compared the 168 resulting list to the E16.5 169 ScRNA-seq data from cells mapping to the 170 RO (Anbarci et al., 2023). This produced a

conservative candidate list of 15 proteins (Tab. 1). Of the candidate proteins, two were selected for validation,

CLU and *STXBP2*, due to their roles in protein and vesicle transport (Argraves and Morales, 2004; Saewu et

al., 2017; Söllner, 2003).

Because no validated antibodies were commercially available to visualize protein expression for CLU and

STXBP2, validations were performed using Hybridization Chain Reaction (HCR), a method for single-

molecule RNA-Fluorescence in situ hybridization (FISH) (https://files.molecularinstruments.com/MI-Protocol-

RNAFISH-Mouse-Rev9.pdf). We found that both *Clu* and *Stxbp2* were expressed in the EOR at P0 (Fig. 5a-

b) and at P7 (Fig. 5c-d). The presence of *Stxbp2* showed that components of the SNARE-complex were

actively transcribed in the EOR, suggesting this may serve as a mechanism for secretion. The presence of

other components of the SNARE-complex were validated using IF. We found that the T-SNARE complex

member STX3, was expressed throughout the EOR and was localized to the apical surface, a cellular position

that is typically associated with active secretion (Fig. 5e) (Söllner, 2003). Ras-associated binding (RAB)

proteins are required in the SNARE complex to tether vesicles to the T-SNARE and allow fusion and secretion

(Takahashi et al., 2012). Using an antibody against the small GTPase found on the surface of vesicles,

RAB11, we found that it was also localized to the sub-apical region of EOR cells (Fig. 5f), which is the typical

docking position for vesicles prior to exocytosis (Söllner, 2003). Taken together, these data suggest that the

EOR actively secretes proteins into the lumen of the structure.

Figure 5. Presence of SNARE-complex members suggests a role for secretion.

(a) Optical section from confocal Z-stacks of PAX8-rtTa; Tre-Cre; Rosa-mTmG (cyan) EOR at E18.5 using HCR for detection of Clu expression (magenta). (b) Optical section from confocal Z-stacks of PAX8-rtTa; Tre-Cre; Rosa-mTmG (cyan) EOR at E18.5 using HCR for Stxbp2 (magenta). (c) Optical section from confocal Z-stacks of PAX8-rtTa; Tre-Cre; Rosa-mTmG (cyan) EOR at P7/8 using HCR for Clu (magenta). (d) Optical section from confocal Z-stacks of PAX8-rtTa; Tre-Cre; Rosa-mTmG (cyan) EOR at P7/8 using HCR for Stxbp2 (magenta). (e) Optical section from confocal Z-stacks of PAX8-rtTa; Tre-Cre; Rosa-mTmG (cyan) EOR at E18.5 immunostained for STX3 (magenta). Outlined higher resolution image acquired with Airyscan (f) Optical section from confocal Z-stacks of PAX8-rtTa; Tre-Cre; Rosa-mTmG (cyan) EOR at E18.5 immunostained for RAB11 (magenta) Outlined higher resolution image acquired with Airyscan . Scale bar – 100um

Discussion

The female reproductive tract is often thought to consist of the vulva, vagina, cervix, uterus, oviduct, and

ovaries (Rendi et al., 2012). We suggest that the RO be added to this list and investigated as an additional

component of female reproductive function. We show that the RO consists of 3 distinct regions that mature

during fetal life and persist into adulthood. The cells of the EOR show secretory characteristics consistent

with the presence of secreted proteins in the lumen identified through mass spectrometry. Importantly, we

show that contents of the EOR are transported to the ovary by P7, suggesting a role in ovary homeostasis.

Historically, the regions of the RO were identified using histological sections, where the EOR was defined as

a columnar epithelium, while the CR was defined as pseudo-columnar. Recently we found that the entire RO

expresses PAX8+, which is usually considered a marker of urogenital epithelial identity. We also showed that

the IOR contained PAX8+/FOXL2+ cells and that the EOR was KRT8+ at E17.5 (McKey et al., 2022). In this

study we further characterized KRT8 and E-Cadherin expression throughout RO development as the CR

acquires epithelial characteristics. We also identified a marker, GFRa1, that is exclusively expressed in the

CR. These molecular distinctions predict that the different regions of the RO are functionally distinct.

While PAS-positive staining within the EOR of sheep indicated the presence of glycogens, glycoproteins, and

proteoglycans (Stein and Anderson, 1979), it remained unclear whether these were secreted by the EOR

cells or were accumulated secretions from the ovary. Using injections of a membrane impermeable dye,

Dextran, into either the RO or ovary, we found that the fluid within the EOR travels towards the ovary. We

used a "milking" process to extrude the contents of EOR, and mass spectrometry, to identify 4,232 proteins.

We expected this group to be a mix of proteins from the luminal fluid, and cytoplasmic proteins from lysed

EOR cells or from cells closely associated with the EOR. We took a very conservative approach to identify

proteins highly likely to be secreted from EOR cells. We compared the proteomics data with the known

mammalian secretome to identify only secreted proteins. We next compared this shortened list with our

available E16.5 Sc-RNA seq data to find proteins that were specific to the EOR at that stage. Although we

recognize that the secretome at P7 likely differs, this candidate list of 15 proteins reveals the first insight into

specific proteins secreted from the RO that may be involved in the secretory process or may affect ovary

function.

Within the protein candidate list, we focused on validating clusterin (CLU) and syntaxin binding protein 2

(STXBP2) due to their roles in protein and vesicle transport. Consistent with our proteomic results, both *Clu*

and *Stxbp2* RNAs were detected in the RO. CLU is a chaperone protein that is present in both reproductive

and non-reproductive tissues (Argraves and Morales, 2004; Saewu et al., 2017). In the epididymis, CLU acts

as a chaperone to direct luminal proteins to the sperm head surface (Saewu et al., 2017). Although the role

of CLU in the RO remains unknown, we hypothesize that it is important for post-secretory protein guidance

to the ovary. STXBP2 binds t-SNARE protein STX and is essential for the vesicle-apical surface membrane

fusion stage of vesicle secretion (Söllner, 2003). The presence of STXBP2 in the candidate list, coupled with

validation of the expression of *Stxbp2* RNA in the cells of the EOR, led us to investigate the presence of other

components of the SNARE complex in the EOR.

The SNARE complex is comprised of a vesicle bound v-SNARE (Vamps), membrane bound t-SNARE (STXs),

small GTPases (RABs), and syntaxin binding proteins (STXBPs) (Söllner, 2003). Using IF staining, we

confirmed the presence of STX3, as well as RAB11. Our ScRNA-seq data indicate that EOR cells also

express *Vamp7* and *Vamp*8, which are v-SNAREs typically associated with STX3 (Dingjan et al., 2018). Small

GTPases, such as RAB11, are essential in the SNARE complex as they mediate the tethering of vesicle

membranes to the apical membrane prior to membrane fusion. Taking into consideration the presence of all

components of the SNARE complex, we hypothesize that the EOR utilizes the SNARE complex to promote

apical secretion into the luminal fluid.

The EOR is highly integrated with the extra-ovarian environment. We found that EOR cells are tightly

surrounded in a web of vasculature. The RO may send or receive information through the vasculature, similar

to other epithelial tissues in the female reproductive tract (Reynolds et al., 2002), raising the possibility that

the RO participates in endocrine signaling. We also found that the EOR is ensheathed within a layer of aSMA+

mesenchyme that expresses the contractile component Calponin (CNN1) (Gao et al., 2014). Our data

suggests that the EOR gains contractility during development, which could facilitate fluid movement within the

EOR lumen in conjunction with cilia within the EOR. We found that the ciliary protein ARL13b was expressed

in the EOR. While the primary cilium is not motile and does not generate flow, we found in our scRNAseq

data that cells in the EOR express *Pkd2*. In renal epithelial cells, nonmotile primary cilia expressing PDK2

sense shear stress during fluid flow and transduce this sensory information (Nauli et al., 2003). Perhaps fluid

movement within the EOR is a two-part process, where the primary cilia sense pressure within the lumen

inducing contraction of the smooth muscle surrounding the EOR to promote fluid movement towards the

ovary. Our E16.5 ScRNA-seq data indicates that the EOR only expresses *Arl13b* and does not express the

multicilia marker *Foxj1*. However, both markers are present in the putative EOR cell clusters in our adult

SnRNA-seq data (Anbarci et al., 2023) suggesting that, after maturation, the EOR likely includes

Arl13b+/Foxj1+ multiciliated cells. It is well documented that adult oviductal multiciliated cells express both

ARL13b and FOXJ1 (Coy et al., 2016). It is possible that the primary cilia give rise to motile multiciliated cells,

as shown in the airway epithelium (Jain et al., 2010) and suggested in the oviduct (Shi et al., 2014).

We also found that the EOR is directly contacted by TUJ1+ neurons, another avenue through which

information may be entering or exiting the EOR. It is unclear whether the innervation contacts the epithelial

cells specifically, similar to the way in which innervation contacts epithelial cells in the gut (Bohórquez et al.,

2015), or whether the innervation contacts the smooth muscle cells that surround the EOR to induce

contraction, similar to the innervation-muscle interaction in the uterus (Morizaki et al., 1989). We also found

that the EOR is specifically associated with F4/80+ LYVE1- macrophages. LYVE1 macrophages are predicted

to have an angiogenic role in the adipose tissue surrounding the epididymis and could possibly be playing a

similar role in the EOR (Cho et al., 2007).

These results demonstrate that the EOR is a secretory epithelium, that its contents move to the ovary at P7,

and that it is highly integrated with the extra-ovarian environment, including vasculature, neurons and

macrophages. This places the EOR in an ideal position to act as an antenna to interpret homeostatic signals

and send information to the ovary. Ongoing investigations will determine whether the EOR responds to

hormones or other physiological signals, and whether proteins secreted by the EOR contribute to the

regulation of ovarian functions.

Methods and Materials

Key Resource Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information	RRID
Strain, strain background (*Mus musculus*)	Crl:CD1(ICR)	Charles River	Strain code: 022		IMSR_CRL:022
Strain, strain background (*M. musculus*)	C57BL/6J	Jackson Laboratory	Stock #:000664		IMSR_JAX:000664
Genetic reagent (*M. musculus*)	Tre-H2B-Gfp (Tg(tetO-HIST1H2BJ/GFP)47Efu/J)	PMID: 14671312	MGI:J:90563		IMSR_JAX:005104
Genetic reagent (*M. musculus*)	Tre-Cre (B6.Cg-Tg(tetO-cre)1Jaw/J)	PMID: 12145322	MGI:J:78365		IMSR_JAX:006234
Genetic reagent (*M. musculus*)	mTmG Gt(ROSA)26Sortm4(ACTB-tdTomato,-EGFP)Luo/J	PMID: 17868096	MGI:J:124702		IMSR_JAX:007576
Genetic reagent (*M. musculus*)	Pax8-rtTA (B6.Cg-Tg(Pax8-rtTA2S*M2)1Koes/J)	PMID: 18724376	MGI:J:140925		IMSR_JAX:007176
Genetic reagent (*M. musculus*)	Lgr5 (B6.129P2-Lgr5tm1(cre/ERT2)Cle/J)	PMID:17934449	MGI:J:127123		IMSR_JAX:008875
Antibody	Smooth muscle alpha action (aSMA) (Cy3-conjugated mouse monoclonal)	Sigma-Aldrich	C6198	1:1000	AB_476856
Antibody	E-Cadherin (rat monoclonal)	Zymed (Thermo Fisher Scientific)	13-1900	1:500	AB_2533005
Antibody	Endomucin (rat monoclonal)	Santa Cruz Biotechnology	sc-65495	1:500	AB_2100037
Antibody	GFP (chicken polyclonal)	Abcam	ab13970	1:1000	AB_300798
Antibody	GFRa1 (goat polyclonal)	R&D Systems	AF560	1:150	AB_2110307
Antibody	KRT8 (rat monoclonal)	DSHB	TROMA-I	1:250	AB_531826
Antibody	PAX8 (rabbit polyclonal)	Proteintech	A10336-1-AP	1:500	AB_2918972
Antibody	CNN1 (rabbit poluclonal)	Proteintech	13938-1-AP	1:200	AB_2082010
Antibody	ARL13b (rabbit polyclonal)	Proteintech	17711-1-AP	1:1000	AB_2060867

Antibody	RAB11 (rabbit monoclonal)	Cell Signaling Technology	5589	1:500	AB_10693925
Antibody	LYVE-1 (goat polyclonal)	R&D Systems	AF2125	1:500	AB_2297188
Antibody	F4/80 (rat monoclonal)	BioRad	MCA497RT	1:2000	AB_1102558
Antibody	STX3 (rabbit monoclonal)	Abcam	ab133750	1:200	
Antibody	TUJ1 (488-conjugated mouse monoclonal)	BioLegend	A488-435L	1:1000	AB_10143904
Antibody	AF647 anti-Rabbit (donkey polyclonal)	Jackson ImmunoResearch	711-605-152	1:500	AB_2492288
Antibody	AF488 anti-Chicken (donkey polyclonal)	Jackson ImmunoResearch	703-545-155	1:500	AB_2340375
Antibody	AF488 anti-Rat (donkey polyclonal)	Life Technologies	A-21208	1:500	AB_2535794
Antibody	Cy3 anti-Goat (donkey polyclonal)	Jackson ImmunoResearch	705-165-147	1:500	AB_2307351
HCR RNA-Fish probe	Stxbp2 (B3 amplifier)	Molecular Instruments	Accession #: XR_001778418.1		
HCR RNA-Fish probe	Clu (B1 amplifier)	Molecular Instruments	Accession #: NM_013492.3		
HCR amplifer	F647 (B3 amplifier)	Molecular Instruments			
HCR amplifer	F546 (B1 amplifier)	Molecular Instruments			
Reagent	Dextran, Alexa FluorTM 568; 10,000 MW	ThermoFisher	D22912		
Chemical compound, drug	Dichloromethane	MilliporeSigma	270997-1L		
Chemical compound, drug	Benzyl Ether	MilliporeSigma	108014-1KG		
Chemical compound, drug	Quadrol = N,N,N′,N′-Tetrakis(2-Hydroxypropyl)ethylene diamine	MilliporeSigma	122262		
Software, algorithm	Zen Black Edition	Carl Zeiss			
Software, algorithm	Imaris v9.6	Bitplane			
Software, algorithm	Adobe Creative Cloud	Adobe		Photoshop, Illustrator, Premier Pro	

279 **Mice and Tissue Collection**

Unless otherwise stated, mice used for experiments were maintained on the CD-1 or mixed CD-1 and C57BL/6J genetic backgrounds. The *Pax8-rtTA* and *Tre-H2BGFP* lines were previously described (Traykova-Brauch et al., 2008; Tumbar et al., 2004) and maintained on a mixed CD-1/C57BL/6J background, and maintained on a mixed CD-1/C57BL/6J background. The *Pax8-rtTa;Tre-Cre;mTmG* line was obtained by crossing the *Pax8-rtTa* line with carriers of the *Tre-Cre* (Perl et al., 2002) and *mTmG* (Muzumdar et al., 2007) alleles, which allowed for visualization of cell membranes of PAX8+ cells. Pregnant and nursing dams were given a doxycycline diet at 625 mg/kg (Teklad Envigo TD.01306) 3 days prior to tissue collection to induce GFP expression in Pax8+ cells. Toe samples were collected from mice for genotyping. Primers used for PCR genotyping are listed in the supplementary table 1. To obtain samples at specific stages of development, males were housed with females for timed matings. Successful mating was determined by the presence of a vaginal plug. Date of plug was considered embryonic day 0.5. Tissue samples were collected in phosphate-buffered saline without calcium or magnesium (PBS -/-), fixed in 4% paraformaldehyde (PFA)/PBS for 30 minutes at room temperature and dehydrated stepwise into 100% methanol, followed by storage at -20°C. All mice were housed in accordance with National Institutes of Health guidelines, and experiments were conducted with the approval of the Duke University Medical Center Institutional Animal Care and Use Committee (protocol #: A078-23-03).

Immunostaining and Confocal Image Acquisition Samples were stepwise rehydrated into 100% PBS, followed by a 30-minute permeabilization wash in PBS 0.1% Triton X-100. Samples were then blocked for 1 hour with PBS, 1% Triton X-100, 10% horse serum, and 3% BSA. Samples were incubated overnight in primary antibodies diluted in blocking solution at 4°C (key resource table). Samples then underwent three 30-minute washes in permeabilization solution and incubated overnight in secondary antibodies (1:500 dilution) and Hoechst vital dye solution diluted in blocking solution. The next day, samples underwent three 20-minute washes in permeabilization solution before being transferred into 100% PBS and stored at 4°C until ready for confocal imaging. Samples were mounted in DABCO mounting solution and stored at -20°C until imaged. Samples were imaged from both the dorsal and ventral sides using 3D-printed reversible slides that utilize two coverslips that allow for flipping. The 3D model can be downloaded on the NIH 3D Print Exchange website at https://3dprint.nih.gov/discover/3DPX-009765. Samples were imaged *in toto* using laser scanning confocal

microscopy captures in the longitudinal plane on Zeiss LSM780 or LSM880 and affiliated Zen software Carl

Zeiss, Inc, Germany) using 10x, 20x, and 63x (also used for Airyscan) objectives.

Hybridization chain reaction

The mouse embryo protocol from Molecular Instruments whole-mount

(https://files.molecularinstruments.com/MI-Protocol-RNAFISH-Mouse-Rev9.pdf) was adjusted for P0 and P7

ovary samples. Samples were stepwise rehydrated into 0.1% Tween 20 / PBS (PBST). Samples were then

subjected to 10 µg/mL proteinase K solution for 10 minutes at room temperature, then washed twice in PBST

for 5 minutes. Samples were then post-fixed in 4% PFA for 10 minutes at room temperature, followed by three

PBST washes for 5 minutes. Samples were then pre-hybridized for 30 minutes in 500ul of hybridization buffer

(Molecular Instruments) at 37°C. Samples were then incubated overnight at 37°C with HCR probes diluted in

500ul hybridization buffer (2pmol in 500ul). Samples were then washed for 15 minutes four times in the

Molecular Instruments probe wash buffer at 37°C, followed by two 5x 0.1% Tween 20/SSC (SSCT) washes

for 5 minutes at room temperature. Fluorescently labeled hairpins were snap cooled and left in a dark drawer

at room temperature for 30 minutes. Following hairpin preparation, samples were incubated with hairpins

diluted in Molecular Instruments amplification buffer overnight in the dark at room temperature. On the third

day, samples were washed in SSCT four times for 15 minutes each before being transferred to PBS -/- and

then mounted in DABCO mounting medium for imaging.

Image Processing

Confocal images were imported into FIJI software for minor image processing (cropping, rotations, maximum

intensity projection/ optical slice montage, and color application). The RO was oriented to the left of the ovary

with the oviduct at the top of the ovary. Images were then imported into Adobe Photoshop CC (Adobe, Inc,

CA) for final processing of channel overlay, brightness, contrast adjustment, and modification of red to

"magenta" (hue adjustment to -30). Channel color accessibility was determined via Photoshop colorblind

proofing.

Dextran Injections

Postnatal day 7 mice were euthanized, and the entire ovarian complex was dissected in PBS +/+. Samples were then placed on agar blocks (1.5%) soaked in PBS +/+. The distal tip of the EOR was identified and punctured using a tungsten needle (0.001mm tip diameter, Fine Science Tools, 10130-05). A microinjection unit (Picospritzer III microinjector) and fine capillary glass needle (5-10 um) loaded with 2-3 nl dextran solution (1.25mg/ml dextran, PBS, 0.05% phenol red) was used to inject dextran into the opening of the EOR. Samples were then left for 15 minutes before fixing with 4% PFA. During the 15 minutes, samples were monitored to visualize movement. Samples were then fixed, stained, and imaged.

Luminal Fluid Collection for Proteomics and Analysis

Postnatal day 7 EORs were collected from *Pax8-rtTa; Tre-H2B-Gfp* mice (as a guide to only collect EOR) and carefully cleaned up to remove as much non-EOR tissue (40 EOR samples). Samples were then placed in a 1.5ml Eppendorf tube and 'pressed' for fluid using a disposable pellet pestle. The sample was then spun down and the supernatant was collected and snap frozen in liquid nitrogen. The sample was then submitted to the Duke University Proteomics Core for mass spectrometry. Proteomic analysis uncovered 4,252 proteins present in the pressed fluid. To exclude proteins that may have contributed due to cell lysis, results were cross-referenced against a database of secreted proteins (Meinken et al., 2015). Overlapping proteins were then cross-referenced to our E16.5 ScRNA-seq data (Anbarci et al, 2023), and candidate proteins were determined by gene expression specific to the EOR cluster.

LC-MS/MS Proteomics Analysis

The sample was subjected to a Bradford (Pierce) protein measurement and 10 ug was removed for downstream processing. The sample was brought to 4% SDS, reduced with 10 mM dithiothreitol for 20 min at 55°C, alkylated with 25mM iodoacetamide for 45 min at room temperature and then subjected to S-trap (Protifi) trypsin digestion using manufacturer recommended protocols. Digested peptides were lyophilized to dryness and resuspended in 50 uL of 0.2% formic acid/2% acetonitrile. The sample was subjected to chromatographic separation on a Waters MClass UPLC equipped with a 1.8 µm Acquity HSS T3 C18 75 µm × 250 mm column (Waters Corp.) with a 90-min linear gradient of 5 to 30% acetonitrile with 0.1% formic acid at a flow rate of 400 nanoliters/minute (nL/min) with a column temperature of 55°C. Data collection on the

Fusion Lumos mass spectrometer with a FAIMS Pro device was performed for three difference compensation voltages (-40v, -60v, -80v). Within each CV, a data-dependent acquisition (DDA) mode of acquisition with a r=120,000 (@ m/z 200) full MS scan from m/z 375 – 1500 with a target AGC value of 4e5 ions was performed. MS/MS scans with HCD settings of 30% were acquired in the linear ion trap in "rapid" mode with a target AGC value of 1e4 and max fill time of 35 ms. The total cycle time for each CV was 0.66s, with total cycle times of 2 sec between like full MS scans. A 20s dynamic exclusion was employed to increase depth of coverage. The total analysis cycle time for each sample injection was approximately 2 hours.

Raw LC-MS/MS data files were processed in Proteome Discoverer 3.0 (Thermo Scientific) and then submitted to independent Sequest database searches against a *Mus musculus* protein database containing both forward and reverse entries of each protein. Search tolerances were 2 ppm for precursor ions and 0.8 Da for product ions using trypsin specificity with up to two missed cleavages. Carbamidomethylation (+57.0214 Da on C) was set as a fixed modification, whereas oxidation (+15.9949 Da on M) was considered a dynamic mass modifications. All searched spectra were imported into Scaffold (v5.3, Proteome Software) and scoring thresholds were set to achieve a peptide false discovery rate of 1% using the PeptideProphet algorithm. Data were output as total spectral matches.

Acknowledgements

The authors would like to thank Benjamin Carlson and Lisa Cameron from the Duke Light Microscopy Core Facility for confocal imaging resources and assistance on high resolution imaging and analysis. The authors would also like to thank Dr. Erik Soderblom from the Duke Center for Genomic and Computational Biology core facility for proteomic analysis and assistance. We are grateful to all members of the Capel and Bagnat laboratories for their constant support, discussion, and suggestions on the work presented in this paper and beyond.

Funding

This project was supported by grants from the National Institutes of Health #1R01HD090050-0 to BC, #1R01DK1321120 to MB, and #K99HD103778 and #R00HD103778 to JM. DNA was supported by #5R01HD039963 to BC and the Duke School of Medicine.

Author contributions

DNA, JM, DSL carried out the experiments; DNA carried out the data curating, validating, and formal analysis; DNA, JM, DSL, and BC designed the experiments; DNA, JM, DSL, MB, and BC drafted, revised, and edited the manuscript; JM, MB, and BC acquired funding for the project.

Competing interests

The authors declare no competing or financial interests.

References

Anbarci, D.N., ORourke, R., Xiang, Y., Peters, D.T., Capel, B., McKey, J., 2023. Transcriptome analysis of the mouse fetal and adult rete ovarii and surrounding tissues. bioRxiv, 2023.2011.2006.565717.

Archbald, L.F., Schultz, R.H., Fahning, M.L., Kurtz, H.J., Zemjanis, R., 1971. Rete ovarii in heifers: a preliminary study. J Reprod Fertil 26, 413-414.

Argraves, W.S., Morales, C.R., 2004. Immunolocalization of cubilin, megalin, apolipoprotein J, and apolipoprotein A-I in the uterus and oviduct. Molecular Reproduction and Development: Incorporating Gamete Research 69, 419-427.

Bohórquez, D.V., Shahid, R.A., Erdmann, A., Kreger, A.M., Wang, Y., Calakos, N., Wang, F., Liddle, R.A., 2015. Neuroepithelial circuit formed by innervation of sensory enteroendocrine cells. The Journal of clinical investigation 125, 782-786.

Byskov, A.G., 1975. The role of the rete ovarii in meiosis and follicle formation in the cat, mink and ferret. Reproduction 45, 201-209.

Byskov, A.G., Lintern-Moore, S., 1973. Follicle formation in the immature mouse ovary: the role of the rete ovarii. J Anat 116, 207-217.

Cassali, G., Nogueira, J., Nascimento, E., Cardoso, J., Ferreira, D., 2000. Morphological and pathological aspects of the rete ovarii in sheep (Ovis aries). Arquivo Brasileiro de Medicina Veterinária e Zootecnia 52, 47-52.

Cho, C.-H., Jun Koh, Y., Han, J., Sung, H.-K., Jong Lee, H., Morisada, T., Schwendener, R.A., Brekken, R.A., Kang, G., Oike, Y., 2007. Angiogenic role of LYVE-1–positive macrophages in adipose tissue. Circulation research 100, e47-e57.

Cool, J., DeFalco, T.J., Capel, B., 2011. Vascular-mesenchymal cross-talk through Vegf and Pdgf drives organ patterning. Proceedings of the National Academy of Sciences 108, 167-172.

Coy, S., Du, Z., Sheu, S.-H., Woo, T., Rodriguez, F.J., Kieran, M.W., Santagata, S., 2016. Distinct patterns of primary and motile cilia in Rathke's cleft cysts and craniopharyngioma subtypes. Modern Pathology 29, 1446-1459.

Czernobilsky, B., Moll, R., Levy, R., Franke, W.W., 1985. Co-expression of cytokeratin and vimentin filaments in mesothelial, granulosa and rete ovarii cells of the human ovary. Eur J Cell Biol 37, 175-190.

Dingjan, I., Linders, P.T., Verboogen, D.R., Revelo, N.H., Ter Beest, M., van den Bogaart, G., 2018. Endosomal and phagosomal SNAREs. Physiological reviews 98, 1465-1492.

Gao, Y., Bayless, K.J., Li, Q., 2014. TGFBR1 is required for mouse myometrial development. Molecular Endocrinology 28, 380-394.

Girsh, E., 2021. Physiology of the Female Reproductive System, A Textbook of Clinical Embryology. Cambridge University Press, pp. 28-55.

Jain, R., Pan, J., Driscoll, J.A., Wisner, J.W., Huang, T., Gunsten, S.P., You, Y., Brody, S.L., 2010.

Temporal relationship between primary and motile ciliogenesis in airway epithelial cells. American journal of

respiratory cell and molecular biology 43, 731-739.

Khan, M., Dodson, A., Heatley, M., 1999. Ki-67, oestrogen receptor, and progesterone receptor proteins in

the human rete ovarii and in endometriosis. Journal of clinical pathology 52, 517-520.

Kim, S.-W., Lee, Y.-H., Lee, S.-R., Kim, K.-M., Lee, Y.-J., Jung, K.-J., Chang, K.-S., Kim, D., Son, H.-Y.,

Reu, D.-S., 2012. Bilateral Ovarian Cysts Originated from Rete Ovarii in an African Green Monkey

(Cercopithecus aethiops). Journal of Veterinary Medical Science, 11-0354.

Lardenois, A., Suglia, A., Moore, C.L., Evrard, B., Noel, L., Rivaud, P., Besson, A., Toupin, M., Leonard, S.,

Lesne, L., 2023. Single-cell transcriptome landscape of developing fetal gonads defines somatic cell lineage

specification in humans. bioRxiv, 2023.2008. 2007.552336.

Lee, S.H., Ichii, O., Otsuka, S., Yaser Hosney, E., Namiki, Y., Hashimoto, Y., Kon, Y., 2011. Ovarian cysts

in MRL/MpJ mice are derived from the extraovarian rete: a developmental study. Journal of Anatomy 219,

743-755.

Mayère, C., Regard, V., Perea-Gomez, A., Bunce, C., Neirijnck, Y., Djari, C., Bellido-Carreras, N., Sararols,

P., Reeves, R., Greenaway, S., 2022. Origin, specification and differentiation of a rare supporting-like

lineage in the developing mouse gonad. Science Advances 8, eabm0972.

McKey, J., Anbarci, D.N., Bunce, C., Ontiveros, A.E., Behringer, R., Capel, B., 2022. Integration of mouse

ovary morphogenesis with developmental dynamics of the oviduct, ovarian ligaments, and rete ovarii. Elife

11, e81088.

McKey, J., Bunce, C., Batchvarov, I.S., Ornitz, D.M., Capel, B., 2019. Neural crest-derived neurons invade

the ovary but not the testis during mouse gonad development. Proceedings of the National Academy of

Sciences 116, 5570-5575.

Meinken, J., Walker, G., Cooper, C.R., Min, X.J., 2015. MetazSecKB: the human and animal secretome and

subcellular proteome knowledgebase. Database 2015.

Morizaki, N., Morizaki, J., Hayashi, R., Garfield, R., 1989. A functional and structural study of the innervation

of the human uterus. American journal of obstetrics and gynecology 160, 218-228.

Muzumdar, M.D., Tasic, B., Miyamichi, K., Li, L., Luo, L., 2007. A global double-fluorescent Cre reporter

mouse. genesis 45, 593-605.

Nauli, S.M., Alenghat, F.J., Luo, Y., Williams, E., Vassilev, P., Li, X., Elia, A.E., Lu, W., Brown, E.M., Quinn,

S.J., 2003. Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. Nature

genetics 33, 129-137.

Perl, A.-K.T., Wert, S.E., Nagy, A., Lobe, C.G., Whitsett, J.A., 2002. Early restriction of peripheral and

proximal cell lineages during formation of the lung. Proceedings of the National Academy of Sciences 99,

10482-10487.

Rendi, M.H., Muehlenbachs, A., Garcia, R.L., Boyd, K.L., 2012. Female reproductive system, Comparative

anatomy and histology. Elsevier, pp. 253-284.

Reynolds, L.P., Grazul-Bilska, A.T., Redmer, D.A., 2002. Angiogenesis in the female reproductive organs:

pathological implications. International journal of experimental pathology 83, 151-164.

Saewu, A., Kadunganattil, S., Raghupathy, R., Kongmanas, K., Diaz-Astudillo, P., Hermo, L., Tanphaichitr,

N., 2017. Clusterin in the mouse epididymis: possible roles in sperm maturation and capacitation.

Reproduction 154, 867-880.

Santos, R.L., Peixoto, D.G.d.M., Turchetti, A.P., Borges, Á.M., do Nascimento, E.F., Paixão, T.A., 2012.

Squamous metaplasia of the rete ovarii in a Zebu cow. BMC Veterinary Research 8, 1-4.

Shi, D., Komatsu, K., Hirao, M., Toyooka, Y., Koyama, H., Tissir, F., Goffinet, A.M., Uemura, T., Fujimori, T.,

2014. Celsr1 is required for the generation of polarity at multiple levels of the mouse oviduct. Development

141, 4558-4568.

Söllner, T.H., 2003. Regulated exocytosis and SNARE function. Molecular membrane biology 20, 209-220.

Stein, L.E., Anderson, C.H., 1979. A qualitative and quantitative study of rete ovarii development in the fetal

rat: correlation with the onset of meiosis and follicle cell appearance. Anat Rec 193, 197-211.

Taelman, J., Czukiewska, S.M., Moustakas, I., Chang, Y.W., Hillenius, S., van der Helm, T., Mei, H., Fan,

X., Chuva de Sousa Lopes, S.M., 2022. Characterization of the human fetal rete region by single cell

transcriptional analysis of gonads and mesonephros/epididymis. bioRxiv, 2022.2007. 2020.500903.

Takahashi, S., Kubo, K., Waguri, S., Yabashi, A., Shin, H.-W., Katoh, Y., Nakayama, K., 2012. Rab11

regulates exocytosis of recycling vesicles at the plasma membrane. Journal of cell science 125, 4049-4057.

Traykova-Brauch, M., Schönig, K., Greiner, O., Miloud, T., Jauch, A., Bode, M., Felsher, D.W., Glick, A.B.,

Kwiatkowski, D.J., Bujard, H., 2008. An efficient and versatile system for acute and chronic modulation of

renal tubular function in transgenic mice. Nature medicine 14, 979-984.

Tumbar, T., Guasch, G., Greco, V., Blanpain, C., Lowry, W.E., Rendl, M., Fuchs, E., 2004. Defining the

epithelial stem cell niche in skin. Science 303, 359-363.

Waldeyer, W., 1870. Eierstock un Ei, Ein Beitrag zur Anatomie und Entwicklungsgeschichte der

Sexualorgane. Рипол Классик.

Wenzel, J.G., Odend'hal, S., 1985. The mammalian rete ovarii: a literature review. Cornell Vet 75, 411-425.

Wenzel, J.G.W., Odend'Hal, S., Player, E., 1987. Histological and histochemical characterization of the

bovine rete ovarii through the estrous cycle and gestation. Anatomia, histologia, embryologia 16, 124-135.

Woolnough, E., Russo, L., Khan, M.S., Heatley, M.K., 2000. An immunohistochemical study of the rete

ovarii and epoophoron. Pathology 32, 77-83.