Bioprocess optimization enables enhanced protopanaxadiol production in yeast

Shangkun Qiu, Mariam Dianat Sabet Gilani, Conrad Müller, Ulf Liebal, Lars M. Blank[*]

iAMB-Institute of Applied Microbiology, ABBt-Aachen Biology and Biotechnology, RWTH Aachen University, Aachen, Germany

[*]Corresponding author: Lars.Blank@rwth-aachen.de

Tel: +49 (0)241 80-26600

Abstract: Ginsenosides are the primary active metabolites of ginseng and have been used in traditional herbal medicine in Asia for more than 4,000 years. Protopanaxadiol (PPD) is the precursor of dammarane-type ginsenosides, exhibiting different pharmacological activities. However, extraction of PPD from plant material is cumbersome because of its low concentration. Precision fermentation using recombinant yeast is a promising alternative strategy to produce PPD. For increasing PPD production, the medium and fermentation conditions were optimized by a Design of Experiment (DoE) approach. PPD production increased by 7.5-fold in the final cultivation condition compared to the reference condition. The PPD titer reached 1.2 g/L during simple 500 mL shake flask cultivations, the highest PPD production in shake flasks to date. The protocol presented facilitates parallel evaluation of recombinant yeast, thereby contributing to the much-needed sustainable synthesis of the versatile molecule class of triterpenoids.

Introduction

Triterpenoids are secondary metabolites ubiquitous in plants with a range of biological properties.[1] The triterpenoid class ginsenosides are mainly found in Ginseng (*Panax* spp.), a plant belonging to the Araliaceae family, which consists of 17 species, of which *Panax quinquefolius*, *Panax ginseng*, and *Panax notoginseng* are commonly industrially exploited. The plant extracts are discussed to possess pharmacological properties, including anti-cancer, anti-diabetic, anti-aging, anti-inflammatory, cardiovascular-protective, and the inhibition of coronavirus disease 2019 (COVID-19).[2,3,4] More than 180 ginsenosides have been identified in the Panax

genus, which can be categorized into two groups based on their aglycones: dammarane-type tetracyclic and oleanane-type pentacyclic ginsenosides.[5,6,7] Dammarane-type tetracyclic ginsenosides can be further divided into three groups, ocotillol, protopanaxadiol (PPD)-type ginsenosides, and protopanaxatriol (PPT)-type ginsenosides, based on the carbohydrate moieties at C3, C6, and C20 positions and the respective hydroxylation.[7]

Ginseng has been used in traditional herbal medicine in Asia for more than 4,000 years.[5,8,3] The term "ginseng" is derived from the Chinese words "人 參 (Renshen)", "the essence of men", indicating its importance.[9] Traditionally, ginsenosides were either consumed using the plant tissue or extracted from the root of ginseng. The root contains a documented total ginsenoside content of 60 mg/g in the dry weight.[3] Changing and low amounts of the single target molecules in plant tissue make commercial production cumbersome.[10,11,8] In addition, Ginseng roots need five to six years of growth until harvesting, with varying influences related to climate, land, soil, and pathogens.[3,5,12] Furthermore, the complex structures of ginsenosides make chemical synthesis economically infeasible.[5,13]

Yeasts are simple eukaryotes that can be engineered for the synthesis of high-value metabolites. With their complete organelle structures, including mitochondria, peroxisomes, Golgi apparatus, endoplasmic reticulum (ER), and vacuoles, which allow post-translational modifications of proteins and operation of compartmented biochemical pathways, they are geared for enzymes from plants.[5,11,14,15] Yeasts also possess redox systems that provide similar biological and physical environments

encountered by cytochrome P450s and uridine diphosphate glycosyltransferase (UGTs)

in plants and mammals.[5,14] Due to these reasons, yeasts are considered excellent cell

factories for producing natural plant products, such as ginsenosides, betulinic acid,

farnesene, amorpha-4,11-diene, and others.[10,11,17,18,13] For example, the production of

α-farnesene was improved by engineered *Yarrowia lipolytica* during fed-batch

fermentation via the expression of the acetyl-CoA acetyltransferase (Erg10p) from

Escherichia coli and HMG-CoA reductase (Hmgp) from *Bordetella petrii*, the final

α-farnesene production reached 26 g/L.[11,19] Also, a highly engineered *Saccharomyces*

cerevisiae expressing the amorpha-4,11-diene synthase (ADS) from *Artemisia annua*

and using an unrestricted ethanol feed allowed the final production of 40 g/L

amorpha-4,11-diene.[13]

Triterpenoids are either synthesized from squalene or 2,3-oxidosqualene, which are

both intermediates of sterol synthesis in yeast.[5,16] The synthesis of ginsenosides by

engineered yeasts from renewable resources is a promising strategy to address the

increasing demand for this interesting class of molecules.[5] The microbial biosynthesis

of ginsenosides has several advantages, including rapid growth, controllable culture

conditions, space efficiency, sustainable production, environmentally friendly, and

well-developed genetic manipulation technology.[5,11,20] At present, only a fraction of

ginsenosides are synthesized by engineered yeast, such as dammarenediol-II (DM),

PPD, PPT, Compound K (CK), ginsenoside Rh1, ginsenoside Rh2, ginsenoside F1,

ginsenoside Rg3, 20S-O-Glc-DM, 3β-O-Glc-DM, ginsenoside Rg1, ginsenoside

NgR1, ginsenoside NgR2, ginsenoside Rf, ginsenoside Glc-Rf, and ginsenoside

Ro.[5,21] Furthermore, there have been some prominent achievements in synthesizing ginsenosides by yeasts. For example, according to a recent study, researchers constructed a PPD-producing *S. cerevisiae* cell factory via overexpression of mevalonate and PPD synthetic pathways and optimization of P450 expression levels; the final PPD production reached 11 g/L in 10 L fed-batch fermentation.[18] Researchers also introduced the ginsenoside CK synthesis pathway into this strain while decreasing UDP-glucose consumption and improving UDP-glucose biosynthesis by engineering strategies; the final CK yield reached almost 6 g/L in fed-batch fermentations.[12]

Research showed that bioprocess optimization, including medium and cultivation conditions, can vastly improve the production of metabolites by cell factories.[22,23] For example, carbon sources (e.g., glucose, glycerol, and glutamate), nitrogen sources, temperature, pH, oxygen, inoculum size, and salts significantly affect the cells' growth and production.[22,24,25] As the parameter space in medium and fermentation parameter optimization is huge, researchers aim for Design of Experiments (DoE) approaches. For example, researchers optimized the concentration of glucose, yeast extract, and peptone via the Box-Behnken design; this optimization increased the geranylgeraniol production by 1.75-fold compared to the traditional YPD medium.[26] Cultured α-farnesene-producing *Y. lipolytica* in a medium optimized by a uniform design (UD) technique reached a final α-farnesene titer of 210 mg/L in a shake flask.[24] In another study, adding citrate (0-10 mM) or acetate (0-10 mM) into the medium improved squalene production. The inhibition of the squalene monooxygenase Erg1p by

terbinafine allowed 730 mg/L of squalene titers.[27] Therefore, improving the

cultivation conditions of engineered yeast is a promising strategy.

This study aims to provide a simple shake flask protocol to evaluate all the different

mutants one might have to generate for further improving triterpenoid synthesis in

yeast. Here, we use PPD synthesis as the main target, as the synthesis requires the

activity of a triterpenoid cyclase and P450 monooxygenase. PPD is one of the

30-carbon dammarane II-type tetracyclic triterpenoids found in ginseng and is the

precursor of high-value ginsenosides.[10] PPD exhibits valuable biological properties,

such as antineoplastic, antioxidant, antidepressant, antitumor, and

anti-inflammation.[28,29] For improved PPD production by yeast in shake flasks, the

following parameters were considered: temperature, glucose and glutamate

concentrations, salts solution, trace element solution, vitamin solution, and oxygen

availability. PPD production reached 1.2 g/L using the best combination of shake

flask set-up and optimized medium, the highest PPD production in shake flasks to

date. This simple set-up facilitates the evaluation of engineered triterpenoid

synthesizing yeast, and the amounts of triterpenoid produced enable advanced

analytics and even first ideas for purification and use of this intriguing class of natural

products.

Figure 1. The PPD synthesis pathway in engineered *S. cerevisiae*. The recombinant enzymes are underlined. G6P,

glucose 6-phosphate; HMG-CoA, 3-hydroxy-3-methylglutaryl-CoA; IPP, isopentenyl pyrophosphate; DMAPP,

dimethylallyl diphosphate; FPP, farnesyl diphosphate; Erg10p, acetyl-CoA acetyltransferase; Erg13p, HMG-CoA

synthase; Hmg1p and Hmg2p, HMG-CoA reductase 1 and 2; Idi1p, isopentenyl-diphosphate delta-isomerase;

Erg20p, farnesyl diphosphate synthase; Erg9p, squalene synthase; Erg1p, squalene monooxygenase; DDS,

dammarenediol-II synthase; PPDS, protopanaxadiol synthase; ER, endoplasmic reticulum. This image was created

by Figdraw.

RESULTS AND DISCUSSION

Effect of medium composition and Oxygen availability on DM and PPD

Production. The DoE experiment included 34 runs designed using Design Expert 11

(Stat-Ease, Inc.). The seven significant factors and the high, medium, and low

categories are shown in Table 1. The system Duetz microtiter plates were used for

cultivation (Enzyscreen). Samples were taken after 72 h, 96 h, and 120 h of

cultivation. DM is the precursor of PPD, which is also a triterpenoid and is the

substrate of the cytochrome P450 monooxygenase protopanaxadiol synthase (PPDS)

for PPD (Figure 1).[30,31] DM and PPD concentrations were highest after 120 h. PPD

production reached 0.3 g/L in four experiments (1, 10, 12, and 33) (Figure 2A).

Among them, runs 1, run 10, and run 12 were all cultured at 20 °C, and run 33 was

cultured at 30 °C. Glucose and glutamate concentrations were three-fold compared to

standard WM8+ medium (100 g/L) for runs 1, 12, and 33. Notably, the salt

concentration of all four runs was 0.33-fold of standard WM8+ medium. Also, higher

oxygen availability was common in these four experiments, using only 1 mL of

culture broth. The PPD concentration reached 0.9 g/L in run 1, the highest PPD

production among all 34 runs in the DoE experiment.

DM production suggests triterpenoid synthesis. However, it is also a potential

limitation in the subsequent P450-catalyzed reaction to PPD. Only seven experiments

showed DM production (Figure 2B). Among the seven experiments, only runs 21 and

27 had DM concentrations above 0.3 g/L. These two runs' glucose, glutamate, and salt

concentrations were three-fold that of standard WM8+ medium. The results suggest

high glucose and glutamate concentrations support high DM and PPD production. At

the same time, low salt concentration, low temperature, and high oxygen

concentration might benefit PPD production, indicating that the P450-catalyzed

hydroxylation is somewhat challenging, with substrate limitation and a slow rate of

catalysis.

Table 1. DoE for improving the standard WM8+ medium and fermentation conditions

Run	Factor A: Glucose	Factor B: Salts	Factor C: Glutamate	Factor D: Trace element	Factor E: Vitamin	Factor F: Temperature [℃]	Factor G: Volume [mL]
1	3	0.33	3	3	0.33	20	1
2	1.67	1.67	1.67	1.67	1.67	25	2
3	3	3	0.33	3	0.33	30	3
4	3	3	0.33	0.33	3	30	3
5	0.33	0.33	0.33	3	0.33	20	3
6	3	3	3	3	3	20	1
7	3	3	0.33	3	0.33	20	1
8	0.33	0.33	3	0.33	0.33	30	1
9	0.33	0.33	0.33	0.33	3	30	1
10	0.33	0.33	3	3	3	20	1
11	0.33	3	3	0.33	3	30	1
12	3	0.33	3	0.33	3	20	1
13	0.33	3	3	3	0.33	20	3
14	0.33	3	3	0.33	3	20	3
15	0.33	0.33	0.33	3	0.33	30	1
16	0.33	3	0.33	0.33	0.33	30	3
17	3	3	0.33	0.33	0.33	30	1
18	0.33	0.33	3	0.33	0.33	20	3
19	0.33	3	0.33	3	3	30	1
20	3	0.33	3	3	3	20	3
21	3	3	3	0.33	0.33	30	3
22	3	0.33	0.33	3	3	20	1
23	1.67	1.67	1.67	1.67	1.67	25	2
24	0.33	3	0.33	0.33	3	20	1
25	1.67	1.67	1.67	1.67	1.67	25	2
26	3	0.33	0.33	0.33	0,33	30	3
27	3	3	3	3	3	30	3
28	3	3	0.33	0.33	0.33	20	3
29	0.33	0.33	3	3	0.33	30	3
30	3	0.33	0.33	0.33	3	20	3
31	0.33	0.33	0.33	3	3	30	3
32	0.33	3	3	0.33	0.33	20	1
33	3	0.33	3	3	3	30	1
34	2	1.67	1.67	1.67	1.67	25	2
CT	1	1	1	1	1	30	2

Every factor has a low (red), middle (yellow), and high (green) value. For factors A-E, the figures in the table

represent fold changes in the actual concentration compared to the standard WM8+ medium. For factor F, the

figures in the table represent the actual temperature. For factor G, figures in the table represent the volume of the

medium in 24-well plates. CT: Control experiment, standard WM8+ medium.

Figure 2. The result of the 34 runs DoE experiment. (A) PPD production by engineered yeast in 24-well plates. (B)

DM production at different fermentation conditions in 24-well plates. CT: Control experiment, standard WM8+

medium.

The ordinary least squares (OLS) Estimation of the DoE Data. DoE data was

analyzed with OLS regression to PPD and DM to identify the predictive factors. We

tested models containing only the individual factor effects (Ind-OLS) as well as

individual factors combined with two-factor interactions (Int-OLS) (Table 2). The

p-value of the F-Statistics identified models with sufficient power to explain the

variance of the observations (PPD) if the p-value is below 0.05. The Ind-OLS and the

more complex Int-OLS explain 41% and 83% of the variance in PPD, respectively

(adj. R²(Ind-OLS)=0.41, adj. R²(Int-OLS)=0.83). The p-values of the individual

factors in each regression model determine whether the factor significantly

contributes to the prediction. There are two significant factors for Ind-OLS (A:

glucose, C: glutamate) and seven for Int-OLS (B: salt, C: glutamate, AxE (vitamins),

AxC, B (salt)xF (temperature), CxD (trace element), BxG (oxygen)) (Figure 3). In the

simple Ind-OLS model with individual factors, only glucose and glutamate are the

most influential predictors that positively contribute to PPD production with

correlation values of 0.4 (Figure 3A). Conversely, salt concentration (B), temperature

(F), and medium volume (G) correlate negatively to the final PPD production, albeit

with low correlation values at -0.2 (Figure 3A). In the more complex Int-OLS model,

glucose is only significant via the two-factor interactions with glutamate (A:C) and

vitamin (A:E) but no longer as an individual factor. The most significant single factors

in the complex Int-OLS model are the positive correlation with glutamate (C) and the

negative correlation with salts (B). While the volume level (G, oxygen) individually is

just above the significance value (p=0.05), it negatively correlates with PPD and is

also negative with the significant interaction with salts (BxG). Thus, less oxygen via

higher volumes decreases PPD (Figure 3A). The yeast increases biomass quickly,

while the oxygen concentration decreases fast in the exponential phase.[25] Furthermore,

NADPH is oxidized to NADP by a P450 reductase in the presence of oxygen, then

transfers electrons to the monooxygenases, such as PPDS.[7] Therefore, controlling the

oxygen concentration will also play a critical role in synthesizing the final product.

Consequently, in the subsequent investigation, we selected glucose, glutamate, salts,

temperature, and oxygen as the main significant factors for further research to check if

these factors will affect the final PPD and DM production.

Table 2. Regression results for OLS with seven factors alone and first-order interactions of two parameters.

Observable	Equation	F-Statistic p-value	Adj. R^2
PPD	Ind-OLS: $\sum c_F F$	0.003	0.41
DM		0.14	0.14
PPD	Int-OLS: $\sum c_F F +$	0.02	0.83
DM	$\sum c_{Fij} F_i F_j$	0.83	-0.53

The F-Statistic shows the p-value for the hypothesis that the independent variables of the OLS model explain the

variability of the dependent variable, R^2 is the percentage of explained variance, Adj. R^2 is a multi-variable

corrected correlation coefficient, Fact. with $p<0.05$, are factors with a p-value below 0.05, along with the estimated

coefficient in the OLS model.

Figure 3. The OLS regression result of the DoE data with prediction of PPD production. (A) The OLS regression

result of the Ind-OLS. (B) The OLS regression result of the Int-OLS. The coefficients with decreasing absolute

values are shown for factors (Ind-OLS) and factor interactions (Int-OLS), significantly explaining the variance

(p-values below 0.05). The larger the coefficient, the greater the influence of this factor or factor interaction on the

final PPD production. A: glucose; B: salts; C: glutamate; D: trace element solution; E: vitamin solution; F:

temperature; G: medium volume.

Effect of Temperature on DM and PPD Production. Sesquiterpenoid production

by engineered *S. cerevisiae* was reported to be better at 25 °C than at 30 °C.[32] We also

investigated temperature as an optimization parameter for PPD production. Here, the

highest PPD production was observed at 20 °C in the 34 runs DoE experiment (Figure

2A). Temperatures above 30 °C were reported to cause stress to baker′s yeast and

decrease cell viability.[25] Therefore, the 15 runs of the DoE experiment cultured at

20 °C were selected to investigate the effect of different temperatures on DM and PPD

production in 24-well plates. To explore the optimum temperature, 18 °C, 20 °C,

22 °C, 25 °C, 28 °C, 30 °C, and 32 °C were selected. The other fermentation and

medium conditions remained constant. In general, the production of PPD increased

with increasing temperature and reached the highest value at 28°C, then decreased

with increasing temperature (Figure 4A, Figure S1). Especially the PPD productions

of runs 1 and 12 are the highest among the 15 runs, reached 1.5 g/L at 28 °C (Figure

4A). But for run 13, run 14, run 18, run 20, run 24, run 30, and run 32, the PPD

production reached the highest value at 32 °C. However, the PPD production for these

seven runs is lower than runs 1 and 12 (Figure S1). So, we can speculate that the

optimum temperature for PPD production is 28 °C.

The DM production of run 1, run 12, and run 24 reached the highest when cultured

at 30 °C. However, the optimum temperature of the run 6 and run 32 is 25 °C, run 10,

run 20, run 22, and run 28 is 28 °C, and run 30 is 32°C (Figure 4B, Figure S1). The

first reason why the optimum temperatures of PPD and DM synthesis in all the runs

are different may be that the different concentrations of compounds in the medium

can affect the DM conversion to PPD at different temperatures. Another reason may

be that the different medium compositions affect the enzyme activities involved in

PPD and DM synthesis at different temperatures. The DM production of run 28 is the

highest when cultured at 28 °C, which reached 1.1 g/L (Figure S1). The DM

production of run 22, run 28, and run 30 are all over 0.8 g/L. The total triterpenoid

production of run 1 is the highest among all the runs, cultured at 30 °C and reached

2.3 g/L (Figure 4C). The total triterpenoid production of run 1, run 12, and run 30 is

over 2 g/L. However, the total triterpenoid production of run 1 and run 12 reached 2

g/L when cultured at 28 °C or 30 °C, while run 30 reached 2 g/L at 32 °C (Figure 4C,

Figure S1). Hence, optimum temperature for DM and total triterpenoid production is

in the range of 28-32 °C.

Figure 4. The effect of different temperatures on DM and PPD production. (A) The comparison of PPD production

between run 1 and run 12 at 18℃, 20℃, 22℃, 25℃, 28℃, 30℃, and 32℃. (B) The comparison of DM

production between run 1 and run 12 at 18℃, 20℃, 22℃, 25℃, 28℃, 30℃, and 32℃. (C) The comparison of

total triterpenoid production between run 1 and run 12 at 18℃, 20℃, 22℃, 25℃, 28℃, 30℃, and 32℃. Other

runs' data is shown in Figure S1.

Effect of Glucose, Glutamate, and Salt Concentration on DM and PPD

Production. According to the results above, runs 1 and 12 have the highest PPD and

total triterpenoid production. The conditions of runs 1 and 12 were chosen for further

investigation (Figure 4A). Glucose, glutamate, and salt were selected to be varied in

concentration, while all other parameters were kept constant. The optimum salt

concentration for PPD production seems to be 0.33, 0.44 or 2 times the WM8+

medium, with the highest PPD production is 2 g/L (Figure 5A). Here, it is indicated

that the optimum concentration of glucose and glutamate for PPD synthesis is two

times that of the standard WM8+ medium. Higher concentrations might cause osmotic

stress and deflect resources from PPD production (Figure 5A).

DM production profits from increased salt concentration, and this is different from

PPD. Again, three times or even higher concentrations of glucose and glutamate

abolish DM production. The highest DM production of all the runs was 1.6 g/L

(Figure 5B). Indeed, under these conditions, doubling the glucose and glutamate

concentrations of WM8+ medium, total triterpenoid production exceeded 3 g/L

(Figure 5C).

Figure 5. The effect of glucose, glutamate, and salt concentration on PPD and DM production (A) PPD production

depends on glucose, glutamate, and salts. (B) DM production is dependent on glucose, glutamate, and salts. (C)

The total triterpenoid production. The glucose and glutamate concentration is two or three times the standard

WM8+ medium. The horizontal coordinate is the salt concentration compared to the standard WM8+ medium.

Runs 1 or 12 means the medium is modified from run 1 or run 12.

For the translation of intracellular products, the yield per biomass is central, as is

the yield on substrate for any biotechnological process. Here, biomass generally

increased with increasing salt concentrations under constant carbon sources (Figure 6).

The highest biomass yield per gram carbon source is 328 mg$_{CDW}$/g$_{glucose\ and\ glutamate}$ in

two-fold salt concentration (Figure 6C). The triterpenoid yield increased with

increasing salt concentrations when glucose and glutamate concentration were twice

that of the standard WM8+ medium. However, the PPD yield first increases and then

decreases with increasing salt concentrations (Figure 6A and C). The highest total

triterpenoid yield is 38 mg/g$_{glucose\ and\ glutamate}$ (Figure 6C). At the same time, the highest

PPD yield is 21 mg/$g_{glucose\ and\ glutamate}$ (Figure 6C).

The total triterpenoid content of ripe ginseng root is 0.1-3 %.[3] We here estimated the CDW content of PPD, DM, and total triterpenoids. The CDW content of the total triterpenoid of three runs is above 120 mg/g_{CDW} (Figure 7A and C). The carbon source concentration of these three runs is two times that of the standard WM8+ medium. The salt concentration is 0.22, 0.33, or 0.44 times the reference. All the data indicates that low salt concentrations favor high CDW content of PPD. In contrast, high salt concentrations can contribute to increased CDW content of DM (Figure 7A and C). The optimum glucose and glutamate concentration for the CDW content of PPD and total triterpenoids is two times of the standard WM8+ medium.

Figure 6. The effect of glucose, glutamate, and salt concentration on triterpenoid yield. (A) The glucose and glutamate concentration is two times the standard WM8+ medium. This medium is modified from run 1. (B) The glucose and glutamate concentration is three times the standard WM8+ medium. This medium is modified from run 1. (C) The glucose and glutamate concentration is twice the standard WM8+ medium. This medium is modified from run 12. (D) The glucose and glutamate concentration is three times the standard WM8+ medium.

This medium is modified from run 12. The horizontal coordinate is the salt concentration compared to the standard

WM8+ medium. All other parameters were used as described in Figure 5.

Figure 7. The effect of glucose, glutamate, and salt concentration on CDW content of triterpenoids. (A) The

glucose and glutamate concentration is two times the standard WM8+ medium. This medium is modified from run

1. (B) The glucose and glutamate concentration is three times the standard WM8+ medium. This medium is

modified from run 1. (C) The glucose and glutamate concentration is twice the standard WM8+ medium. This

medium is modified from run 12. (D) The glucose and glutamate concentration is three times the standard WM8+

medium. This medium is modified from run 12. The horizontal coordinate is the salt concentration compared to the

standard WM8+ medium. All other parameters were used as described in Figure 5.

Triterpenoid Production in 500 mL Shake Flasks. While microtiter plate

cultivation allows strain comparison, shake flask cultivations can be designed to test

product purification ideas and synthesize the first products for use. For this scale increase, we

selected four media from the above experiments, which are medium 1 (3-fold, run 1),

medium 2 (3-fold, run 12), medium 3 (2-fold, run 12), and medium 4 (2-fold, run 12)

(Figure 5A). Yeasts were cultivated in two types of 500 mL shake flasks. The shake

flask was closed by either an aluminum or a membrane lid (Figure 8A). The

membrane lid facilitates better aeration when compared to the aluminum lid. Two

enzymes in PPD synthesis require oxygen: the conversion of squalene to

2,3-oxidosqualene, catalyzed by Erg1p, and the conversion of DM to PPD, catalyzed

by PPDS (Figure 1).[33,34] First, yeasts were cultivated in the shake flask with an

aluminum lid. The smaller the filling volume of the shake flask, the better the aeration.

Two different filling volumes were used, i.e., 25 mL and 50 mL. The yeast was

cultured in the four improved media. The results showed the highest PPD production

of 0.8 g/L occurred in medium 3 with 25 mL filling volume. The PPD production

decreased with the increase in the medium. The PPD production in all four media

tested is higher than that in WM8+ medium, which is 0.16 g/L. It can be speculated

that the higher oxygen concentration benefits the synthesis of PPD, which can

increase the final PPD production in the shake flask (Figure 8B).

Figure 8. Shake flask configuration influences PPD production. (A) Shake flasks with membrane and aluminum lid. (B) PPD production depends on the filling volume using a shake flask with an aluminum lid. (C) Water evaporation rate in the 500 mL shake flask with a membrane lid. (D) Water evaporation rate in the 500 mL shake flask with an aluminum lid. (E) PPD and DM production depends on the filling volume using a shake flask with a membrane lid. UN, unoptimized standard WM8+ medium; M1, Optimum medium 1; M2, Optimum medium 2; M3, Optimum medium 3; M4, Optimum medium 4.

The identical experiments were repeated using the membrane lid for improved aeration. Notably, water evaporation is higher using the membrane lid (Figure 8C and D), although the shaker was equipped with a moisture controller, keeping water content in the air at 70% saturation. To account for evaporation, we calculated the water evaporation in the 500 mL shake flask with both types of lids at 28 °C (Figure

8C and D). We measured the volume after 120 h and calculated the product

concentration without evaporation. The corrections were as follows:

$$\frac{a \times b}{c} = d \qquad (1)$$

where a refers to the PPD or DM production (g/L) measured by HPLC; b refers to the

final volume (mL); c refers to initial volume (mL); d refers to the actual PPD or DM

production (g/L) without evaporation.

The results indicated that the PPD production also decreased with the increase of

medium volume. The PPD productions of optimum media 3 and 4 in this membrane

lid shake flask are higher than the PPD production of the same media with an

aluminum lid, reaching 1.2 g/L and 0.6 g/L when the volume is 25 mL. At the same

time, the DM production increased with medium volume (Figure 8E). This indicated

that the membrane lid and smaller volume allow more oxygen to enter the shake flask,

which supplies more oxygen for improving the conversion of squalene to

2,3-oxidosqualene and the conversion of DM to PPD (Figure 1). The highest PPD

productions of unoptimized standard WM8+ medium in the shake flask with an

aluminum lid or membrane lid are 0.16 g/L or 0.45 g/L, respectively (Figure 8E). This

indicated that the highest PPD production of the optimum medium 3 is 2.6-fold

compared to the unoptimized standard WM8+ medium in the shake flask with a

membrane lid. However, the PPD productions in media 1 and 2 with this membrane

lid shake flask are much lower than the PPD production in medium 3 (data not

shown). This may be due to the higher glucose and glutamate concentration (three

times the standard WM8+ medium) in media 1 and 2, limiting the PPD and DM

synthesis in the shake flask, just like the analysis described in Figure 5.

The fermentation kinetics of PPD and DM in engineered yeast cultured in the

standard WM8+ medium and the M3 medium. Ethanol is a more reduced carbon

source than glucose, which feeds directly into the cytosolic acetyl-CoA pool required

for the mevalonate pathway.[7] Therefore, we also measured the concentration of the

carbon sources and triterpenoid products during the fermentation to investigate the

fermentation kinetics of PPD and DM. As a Crabtree-positive yeast, the engineered *S.*

cerevisiae uses respiro-fermentative glucose metabolism and produces ethanol and

carbon dioxide.[35] After glucose depletion at 24 h, ethanol will be the carbon source in

the presence of oxygen, as the classic diauxic shift of *S. cerevisiae*. The ethanol

concentration also reached the highest value at 24 h, 12.5 g/L, and 35 g/L in the

standard WM8+ and M3 medium (Figure 9A and B). Ethanol has been shown to be a

beneficial carbon source for triterpenoid production.[7] Therefore, PPD and DM

synthesis was very fast when the carbon source became ethanol. The highest PPD

production is 0.6 g/L and 1.8 g/L in the standard WM8+ and M3 medium, respectively,

slightly higher than the PPD production above data (0.45 g/L in the standard WM8+

medium and 1.2 g/L in the M3 medium) (Figure 8E, 9A and B). The reason should be

due to the several samples taken during the fermentation that reduced the volume and

increased the oxygen concentration; this led to higher PPD production. The result

indicated that the M3 medium performs better than the standard WM8+ medium for

PPD synthesis.

Figure 9. Fermentation kinetics of PPD and DM in engineered yeast. (A) The concentration of glucose, ethanol, PPD, and DM during the fermentation in engineered yeast cultured in the standard WM8+ medium. (B) The concentration of glucose, ethanol, PPD, and DM during the fermentation in engineered yeast cultured in the M3 medium.

According to previous studies, it is a promising choice to improve the production of the final metabolites in the cell's factories by optimizing the fermentation bioprocess, including medium compounds and fermentation conditions.[22,23] The PPD production of the M3 medium is nearly 2 g/L in the 24-well plates, which is much more than the PPD production in the shake flask with a membrane lid, 1.2 g/L. The reasons may be due to the cells' sediment, the water evaporation, some cells on the wall, oxygen concentration, and other factors that are different between the two growth formats. We also checked other engineered yeasts that can produce other triterpenoids (betulin and betulinic acid) with the M3 medium. The result showed that the betulin and betulinic acid productions of these two strains in the M3 medium are largely increased compared to the productions in the standard WM8+ medium (Figure 10). This means that the M3 medium also works for other engineered strains. As aimed, the results can be generalized to evaluate engineered yeasts under simple

growth conditions.

Figure 10. Betulin (B) and betulinic acid (BA) produced by engineered yeasts. The aluminum lid was used for

shake flask cover and the media WM8+ and M3 were compared.

In conclusion, we identified the optimum temperature for PPD production at 28 °C.

The M3 medium presented here for producing PPD by yeast was built by stepwise

evaluation and optimization of the WM8+ medium. Furthermore, we used a

membrane lid to improve the final PPD production, which facilitated increased

aeration. Eventually, a remarkable titer of 1.2 g/L PPD was achieved in the 500 mL

shake flask, a 7.5-fold improvement compared to the unoptimized standard WM8+

medium with original fermentation conditions (0.16 g/L). As the medium and growth

conditions are easily established and transferable to other triterpenoid-producing

baker´s yeast, the study might facilitate the simple evaluation of engineered strains in

microtiter plates and shake flask cultivations.

MATERIALS AND METHODS

Strains, Medium, and Reagents. The engineered yeast 102-tm-reerg7-2PP for PPD

synthesis was modified from *S. cerevisiae* CEN.PK 102-5B and originated from a

previous study.[16,36] *S. cerevisiae* BA6Δ (unpublished) is a pah1 deletion mutant of

442 BA6, generated through replacement of PAH1 with a hygromycin resistance cassette

443 (YMR165CΔ:hphMX).[37,38] The engineered yeast BA6Δ for betulin synthesis was

444 modified from *S. cerevisiae* Simo1575.[16] Standard WM8+ medium included 50 g/L

445 glucose, salt solution, 10 g/L glutamate, trace elements solution, vitamins solution,

446 verduyn-trace elements solution, and verduyun-vitamins solution. In addition, the salt

447 solution includes 0.25 g/L $(NH_4)H_2PO_4$, 2.8 g/L NH_4Cl, 0.25 g/L $MgCl_2·6H_2O$, 0.1

448 g/L $CaCl_2·2H_2O$, 2 g/L KH_2PO_4, 0.55 g/L $MgSO_4·7H_2O$, and 0.075 g/L $C_6H_{12}O_6$. The

449 trace elements includes 1.75 mg/L $ZnSO_4·7H_2O$, 0.5 mg/L $FeSO_4·7H_2O$, 0.1 mg/L

450 $CuSO_4·5H_2O$, 0.1 mg/L $MnCl_2·4H_2O$, 0.1 mg/L $Na_2MoO_4·2H_2O$, and EDTA (0.04

451 mM). The vitamins solution includes 10 mg/L $C_6H_5NO_2$ (nicotinic acid), 25 mg/L

452 $C_8H_{11}NO_3$ (pyridoxine), 10 mg/L $C_{12}H_{17}N_4OS$ (thiamine), 2.5 mg/L $C_{10}H_{16}N_2O_3S$

453 (biotin), and 50 mg/L $C_9H_{17}NO_5$ (calcium-d-pantothenate). Verduyn-Trace elements

454 solution includes 15 mg/L Na_2EDTA, 4.5 mg/L $CaCl_2·2H_2O$, 4.5 mg/L $ZnSO_4·7H_2O$,

455 3 mg/L $FeSO_4·7H_2O$, 0.3 mg/L $CuSO_4·5H_2O$, 1 mg/L $MnCl_2·2H_2O$, 0.4 mg/L

456 $Na_2MoO_4·2H_2O$, 0.3 mg/L $CoCl_2·6H_2O$, 1 mg/L H_3BO_3, and 0.1 mg/L KI.

457 Verduyn-vitamins solution includes 25 mg/L $C_6H_{12}O_6$ (myo-inositol), 1 mg/L

458 $C_6H_5NO_2$ (nicotinic acid), 1 mg/L $C_8H_{11}NO_3$ (pyridoxine), 1 mg/L $C_{12}H_{18}Cl_2N_4OS$

459 (thiamin-HCl), 0.05 mg/L $C_{10}H_{16}N_2O_3S$ (biotin), 1 mg/L $C_9H_{17}NO_5$

460 (calcium-pantothenate), and 0.2 mg/L $C_7H_7NO_2$ (p-aminobenzoic acid).[10] The PPD

461 standard was purchased from PhytoLab. The DM standard was purchased from

462 AOBIOUS.

463 **Design of Experiment.** The 34 runs DoE was designed using Design Expert 11

(Stat-Ease, Inc.). Study type: Factorial; Subtype: Randomized; Design type: Minimum

Run Resolution V; 34 runs; Design Model: 2FI; Blocks: No Blocks; Center Points: 4.

The experiments were carried out in 24-well plates (polypropylene square deep well

MTP, re-usable, 24 wells of 11 ml volume, flattened, detoxified, Kuhner shaker,

Germany).

Ordinary least squared (OLS) Estimation. An OLS regression was used with two

sets of linear formulas: 1. containing the individual factor ($\sum c_i F_i$, with c_i as the

coefficient for each factor F_i=[A, B, C, D, E, F, G]) and 2. the individual factor with

first order factor interactions ($\sum c_i F_i + \sum c_{ij} F_i F_j$, with c_{ij} as the coefficient for each

two-factor interaction). A regression was performed with the seven factors A-G

towards either PPD or DM, resulting in four regressions. All data was centered to

have a mean value of 0 and a standard deviation of 1. The evaluation of the OLS

models is based on the F-statistics, which reports whether a combination of

independent variables (the factors in the OLS model) can explain the variance of the

dependent variable (PPD or DM) if the p-value is below 0.05. The factors are selected

by their individual p-values of their contribution to the OLS model. The impact of

each factor is then determined by the respective coefficient. The calculations are

performed with the *ols* package in the *statsmodel* (v0.13.2) library in *Python*, and the

code is provided as Jupyter Notebook in the Supplementaries.

Cultivation of Engineered Yeasts in Shake Flasks. Pre-cultures in the 500 mL shake

flasks with an aluminum lid containing 50 mL WM8+ medium and inoculate each

pre-culture with 500 µL of a thawed cryogenic stock of the engineered yeast (30 °C,

200 rpm, and 50 mm shaking diameter). The pre-culture medium with an OD_{600} of

6-10 was used as the inoculum of the main culture. The start OD_{600} of the main

culture was set to 0.4, measured with a single-use cuvette (polystyrene, semi-micro,

1.6 mL, Carl Roth, Germany) in a photometer (Biochrom Ltd., United Kingdom). All

experiments were carried out in the 500 mL shake flasks with aluminum or membrane

lids. The filling volume differed between 25 and 50 mL (see text for details). The

cultivation temperature differed between 28 °C and 30 °C (200 rpm and 50 mm

shaking diameter). At least duplicates were performed.

Determination of PPD, DM, and Biomass concentrations. 800 µL of cell broth

were sampled, transferred into 2 mL reaction tubes (Eppendorf, Germany), and frozen

at -20 °C until further use. For cell disruption, 250 µL of glass beads (0.5 mm

diameter, Carl Roth, Germany) were added, followed by 800 µL chloroform:

methanol (80:20 ratio) and 80 µL HCl (1 M). The samples were then placed into the

MinibeadBeater (Biospec Products, USA) for 2 minutes twice. For phase separation,

samples were centrifuged for 10 minutes at 4 °C, 13,000 rpm (Eppendorf Centrifuge

5418 R, Eppendorf, Germany). 500 µL of the lower organic phase containing the PPD

and DM were filtered (4 mm syringe filters RC membrane, pore size 0.2 µm,

Phenomenex, Germany) and transferred into HPLC vials (Flat N9 vials, screw neck,

1.5 mL, clear, Macherey-Nagel, Germany) and stored at 4 °C until HPLC analysis.

PPD and DM concentration were determined with an HPLC system (UltiMate 3000

HPLC system, Thermo Fisher Scientific, USA) coupled to a Dionex Corona Veo

Charged Aerosol Detector. The column was an EC150/4.6 NUCLEODUR C18

gravity column (Macherey-Nagel GmbH and Co. KG, Germany), the mobile phase

was Acetonitrile:0.2 % (v/v) formic acid, the column temperature was set to 40 ℃, the

flow rate was programmed to 1.2 mL/min, the injection volume was set to 5 μL, the

runtime was 23 min. To measure the CDW, 2 mL culture was harvested by

centrifugation for 5 minutes at 4 ℃, 13,000 rpm (Eppendorf Centrifuge 5418 R,

Eppendorf, Germany). The collected yeast cells were washed with deionized water

and oven-dried to a constant weight. An OD_{600} to CDW correlation of CDW (g/L)=0.2

$\times OD_{600}$+1.3 was estimated. All values are reported as the average of three replicates.

Water evaporation. Water evaporation was monitored using 500 mL shake flasks

with either the aluminum or membrane lid. Different water volumes (25 mL and 50

mL) were incubated in a shaker for 120 h (28 ℃, 200 rpm, and 50 mm shaking

diameter).

Author Contributions

Shangkun Qiu completed all the experiments, including the DoE design, OLS

estimation analysis, triterpenoids measurement, and data analysis. Shangkun Qiu also

wrote the first draft of the manuscript. Mariam Dianat Sabet Gilani helped with the

yeast fermentation and triterpenoid measurements. Conrad Müller advised on the DoE

experiment design and DoE data analysis. Ulf Liebal advised the OLS estimation

analysis. Lars M. Blank supervised the study, discussed the results, and revised the

manuscript.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

Shangkun Qiu was funded by the China Scholarship Council (CSC). The China

Scholarship Council is the Chinese Ministry of Education's non-profit organization

that provides support for international academic exchange with China. Lars M. Blank

acknowledges funding from the European Union's Horizon 2020 Research and

Innovation Program under Grant Agreement 870294 for the EU/China Project

MIX-UP.

REFERENCES

(1) Li, Z.; Wang, Y.; Wang, L.; Shi, T.; Sun, X.; Huang, H. Advanced strategies for the synthesis of terpenoids in *Yarrowia lipolytica*. *J Agr Food Chem.* **2021**, *69* (8), 2367-2381.

(2) Piao, X.; Zhang, H.; Kang, J.; Yang, D.; Li, Y.; Pang, S.; Jin, Y.; Yang, D.; Wang, Y. Advances in saponin diversity of *Panax ginseng*. *Molecules.* **2020**, *25* (15), 3452.

(3) Hou, M.; Wang, R.; Zhao, S.; Wang, Z. Ginsenosides in Panax genus and their biosynthesis. *Acta Pharm Sin B.* **2021**, *11* (7), 1813-1834.

(4) Jung, E.-M.; Lee, G.-S. Korean red ginseng, a regulator of NLRP3 inflammasome, in the COVID-19 pandemic. *J Ginseng Res.* **2022**, *45* (3), 331-336.

(5) Chu, L.; Montecillo, J.; Bae, H. Recent advances in the metabolic engineering of yeasts for ginsenoside biosynthesis. *Front Bioeng Biotech.* **2020**, *8*, 139.

(6) Li, Y.; Li, J.; Diao, M.; Peng, L.; Huang, S.; Xie, N. Characterization of a group of UDP-glycosyltransferases involved in the biosynthesis of triterpenoid saponins of *Panax notoginseng*. *ACS Synth Biol.* **2022**, *11* (2), 770-779.

(7) Qiu, S.; Blank, L. Recent advances in yeast recombinant biosynthesis of the triterpenoid protopanaxadiol and glycosylated derivatives thereof. *J Agric Food Chem.* **2023**, *71* (5), 2197-2210.

(8) Kim, Y.; Zhang, D.; Yang, D. Biosynthesis and biotechnological production of ginsenosides. *Biotechnol adv.* **2015**, *33*, 717-735.

(9) Xu, J.; Chu, Y.; Liao, B.; Xiao, S.; Yin, Q.; Bai, R.; Su, H.; Dong, L.; Li, X.; Qian, J.; et al. *Panax ginseng* genome examination for ginsenoside biosynthesis. *GigaScience.* **2017**, *6* (11), 1-15.

(10) Czarnotta, E.; Dianat, M.; Korf, M.; Granica, F.; Merz, J.; Maury, J.; Baallal Jacobsen, S.; Förster, J.; Ebert, B.; Blank, L. Fermentation and purification strategies for the production of betulinic acid and its lupane-type precursors in *Saccharomyces cerevisiae*. *Biotechnol Bioeng.* **2017**, *114* (11), 2528-2538.

(11) Muhammad, A.; Feng, X.; Rasool, A.; Sun, W.; Li, C. Production of plant natural products through engineered *Yarrowia lipolytica*. *Biotechnol Adv.* **2020**, *43*, 107555.

(12) Wang, P.; Wang, J.; Zhao, G.; Yan, X.; Zhou, Z. Systematic optimization of the yeast cell factory for sustainable and high efficiency production of bioactive ginsenoside compound K. *Syn Syst Biotechno.* **2021**, *6* (2), 69-76.

(13) Westfall, P. J.; Pitera, D. J.; Lenihan, J. R.; Eng, D.; Woolard, F. X.; Regentin, R.; Horning, T.; Tsuruta, H.; Melis, D. J.; Owens, A.; et al. Production of amorphadiene in yeast, and its conversion to dihydroartemisinic acid, precursor to the antimalarial agent artemisinin. *P Natl A Sci.* **2012**, *109* (3), E111-E118.

(14) Orlean, P. Architecture and biosynthesis of the *Saccharomyces cerevisiae* cell wall. *Genetics.* **2012**, *192* (3), 775-818.

(15) Choi, B. H.; Kang, H. J.; Kim, S. C.; Lee, P. C. Organelle engineering in yeast: enhanced production of protopanaxadiol through manipulation of peroxisome proliferation in *Saccharomyces cerevisiae*. *Microorganisms.* **2022**, *10* (3), 650.

(16) Guo, H.; Jacobsen, S.; Walter, K.; Lewandowski, A.; Czarnotta, E.; Knuf, C.; Polakowski, T.; Maury, J.; Lang, C.; Förster, J.; et al. Triterpenoid production with a minimally engineered *Saccharomyces cerevisiae* chassis. *bioRxiv.* **2022**. DOI: DOI: 10.1101/2022.07.11.499565.

(17) Wang, C.; Liwei, M.; Park, J.; Jeong, S.; Wei, G.; Wang, Y.; Kim, S. Microbial platform for terpenoid production: *Escherichia coli* and Yeast. *Front microbiol.* **2018**, *9*, 2460.

(18) Wang, P.; Wei, W.; Ye, W.; Li, X.; Zhao, W.; Yang, C.; Li, C.; Yan, X.; Zhou, Z. Synthesizing ginsenoside
Rh2 in *Saccharomyces cerevisiae* cell factory at high-efficiency. *Cell discov.* **2019**, *5* (5).

(19) Liu, Y.; Jiang, X.; Cui, Z.; Wang, Z.; Qi, Q.; Hou, J. Engineering the oleaginous yeast *Yarrowia*
lipolytica for production of α-farnesene. *Biotechnol Biofuels.* **2019**, *12*, 296.

(20) Srisawat, P.; Yasumoto, S.; Fukushima, E.; Robertlee, J.; Seki, H.; Muranaka, T. Production of the
bioactive plant-derived triterpenoid morolic acid in engineered *Saccharomyces cerevisiae*. *Biotechnol*
Bioeng. **2020**, *117* (7), 2198-2208.

(21) Zhang, H.; Hua, X.; Zheng, D.; Wu, H.; Li, C.; Rao, P.; Wen, M.; Choi, Y.; Xue, Z.; Wang, Y.; et al. De
novo biosynthesis of oleanane-type ginsenosides in *Saccharomyces cerevisiae* using two types of
glycosyltransferases from *Panax ginseng*. *J Agric Food Chem.* **2022**, *70* (7), 2231-2240.

(22) Zhang, A.; Xie, Y.; He, Y.; Wang, W.; Sen, B.; Wang, G. Bio-based squalene production by
Aurantiochytrium sp. through optimization of culture conditions, and elucidation of the putative
biosynthetic pathway genes. *Bioresour Technol.* **2019**, *287*, 121415.

(23) Zhou, J.; Hu, T.; Gao, L.; Su, P.; Zhang, Y.; Zhao, Y.; Chen, S.; Tu, L.; Song, Y.; Wang, X.; et al.
Friedelane-type triterpene cyclase in celastrol biosynthesis from *Tripterygium wilfordii* and its
application for triterpenes biosynthesis in yeast. *New Phytol.* **2019**, *223* (2), 722-735.

(24) Liu, S.; Liu, Z.; Wei, L.; Hua, Q. Pathway engineering and medium optimization for α-farnesene
biosynthesis in oleaginous yeast *Yarrowia lipolytica*. *J Biotechnol.* **2020**, *319*, 74-81.

(25) Carsanba, E.; Pintado, M.; Oliveira, C. Fermentation strategies for production of pharmaceutical
terpenoids in engineered yeast. *Pharmaceuticals (Basel).* **2021**, *14* (4).

(26) Song, T.; Ding, M.; Zhai, F.; Liu, D.; Liu, H.; Xiao, W.; Yuan, Y. Engineering *Saccharomyces cerevisiae*
for geranylgeraniol overproduction by combinatorial design. *Sci Rep.* **2017**, *7* (1), 14991.

(27) Tang, W.; Wang, D.; Tian, Y.; Fan, X.; Wang, C.; Lu, X.; Li, P.; Ji, X.; Liu, H. Metabolic engineering of
Yarrowia lipolytica for improving squalene production. *Bioresource technol.* **2021**, *323*, 124652.

(28) Zhao, F.; Bai, P.; Nan, W.; Li, D.; Zhang, C.; Lu, C.; Qi, H.; Lu, W. A modular engineering strategy for
high-level production of protopanaxadiol from ethanol by *Saccharomyces cerevisiae*. *Aiche J.* **2018**, *65*
(3), 866-874.

(29) Yang, L.; Zhang, X.; Li, K.; Li, A.; Yang, W.; Yang, R.; Wang, P.; Zhao, Z.; Cui, F.; Qin, Y.; et al.
Protopanaxadiol inhibits epithelial-mesenchymal transition of hepatocellular carcinoma by targeting
STAT3 pathway. *Cell Death Dis.* **2019**, *10* (9), 630.

(30) Gao, X.; Caiyin, Q.; Zhao, F.; Wu, Y.; Lu, W. Engineering *Saccharomyces cerevisiae* for enhanced
production of protopanaxadiol with cofermentation of glucose and xylose. *J Agr Food Chem.* **2018**, *66*
(45), 12009-12016.

(31) Guo, H.; Wang, H.; Chen, T.; Guo, L.; Blank, L. M.; Ebert, B. E.; Huo, Y.-X. Engineering critical amino
acid residues of lanosterol synthase to improve the production of triterpenoids in *Saccharomyces*
cerevisiae. *ACS Synth Biol.* **2022**, *11* (8), 2685-2696.

(32) Ouyang, X.; Cha, Y.; Li, W.; Zhu, C.; Zhu, M.; Li, S.; Zhuo, M.; Huang, S.; Li, J. Stepwise engineering
of *Saccharomyces cerevisiae* to produce (+)-valencene and its related sesquiterpenes. *RSC Adv.* **2019**, *9*
(52), 30171-30181.

(33) Li, T.; Liu, G.; Zhou, W.; Jiang, M.; Ren, Y.; Tao, X.; Liu, M.; Zhao, M.; Wang, F.; Gao, B.; et al.
Metabolic engineering of *Saccharomyces cerevisiae* to overproduce squalene. *J Agr Food chem.* **2020**,
68 (7), 2132-2138.

(34) Istiandari, P.; Yasumoto, S.; Srisawat, P.; Tamura, K.; Chikugo, A.; Suzuki, H.; Seki, H.; Fukushima, E.;
Muranaka, T. Comparative analysis of NADPH-Cytochrome P450 reductases from legumes for

heterologous production of triterpenoids in transgenic *Saccharomyces cerevisiae*. *Front Plant Sci.* **2021**, *12*, 762546.

(35) Anderlei, T.; Zang, W.; Papaspyrou, M.; Büchs, J. Online respiration activity measurement (OTR, CTR, RQ) in shake flasks. *Biochem Eng J.* **2004**, *17* (3), 187-194.

(36) Guo, H. *Optimization of triterpenoid production in engineered Saccharomyces cerevisiae*; RWTH Aachen University, 2019.

(37) Dianat, M.; Münchberg, U.; Blank, L. M.; Freier, E.; Ebert, B. E. Non-invasive monitoring of microbial triterpenoid production using nonlinear microscopy techniques. *Front Bioeng Biotechnol.* **2023**, *11*, 1106566.

(38) Kerstin, W. *Compartment engineering to increase metabolic flux through the mevalonate pathway in Saccharomyces cerevisiae*; RWTH Aachen University, 2018.

A

PPD production (g/L)

Medium 1
Medium 2
Medium 3
Medium 4

0.22-fold 0.33-fold 0.44-fold 0.66-fold 1-fold 2-fold

■ 2-fold, run 1 ■ 3-fold, run 1 ■ 2-fold, run 12 ■ 3-fold, run 12

B

DM production (g/L)

0.22-fold 0.33-fold 0.44-fold 0.66-fold 1-fold 2-fold

■ 2-fold, run 1 ■ 3-fold, run 1 ■ 2-fold, run 12 ■ 3-fold, run 12

C

Total triterpenoids production (g/L)

0.22-fold 0.33-fold 0.44-fold 0.66-fold 1-fold 2-fold

■ 2-fold, run 1 ■ 3-fold, run 1 ■ 2-fold, run 12 ■ 3-fold, run 12