Combinatorial CRISPR screen reveals *FYN* and *KDM4* as targets for synergistic drug combination for

treating triple negative breast cancer

Running title: FYN and KDM4 inhibition synergizes with TKIs

Tackhoon Kim[1,2,3,4,*] Byung-Sun Park[1,2], Soobeen Heo[5], Heeju Jeon[1,2], Jaeyeal Kim[1,2], Donghwa Kim[6], Sang

Kook Lee[6], So-Youn Jung[7], Sun-Young Kong[5,8], Timothy K. Lu[4,*]

[1] Medicinal Materials Research Center, Korea Institute of Science and Technology, 5 Hwarangro-14-gil,

SeongbukGu, Seoul 02792, Republic of Korea

[2] Department of Biological Sciences, Korea University, 145 AnamRo, SeongbukGu, Seoul 02841, Republic of

Korea

[3] Division of Bio-Medical Science and Technology, Korea University of Science and Technology, 217

GajeongRo, YuseongGu, Daejeon 34113, Republic of Korea

[4] Research Lab of Electronics, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge

MA 02139, USA

[5] Targeted Therapy Branch, Division of Rare and Refractory Cancer, Research Institute, National Cancer Center,

Goyang 10408, Korea

[6] College of Pharmacy, Natural Products Research Institute, Seoul National University, 1 Gwanakro, Gwanakgu,

Seoul 08826, Republic of Korea.

[7] Division of Breast Surgery, Department of Surgery, National Cancer Center, Goyang 10408, Korea

[8] Department of Laboratory Medicine, National Cancer Center, Goyang 10408, Republic of Korea

* Correspondence: Tackhoon Kim, 5 Hwarangro-14-gil, SeongbukGu, Seoul 02792, Republic of Korea;

Timothy K. Lu, 500 Technology Square, Cambridge MA 02139, USA.

Email: tackhoon@kist.re.kr; tim@lugroup.org

Competing Interest Statement

The authors declare no conflict of interest

Abstract

Tyrosine kinases play a crucial role in cell proliferation and survival and are extensively investigated as targets for cancer treatment. However, the efficacy of most tyrosine kinase inhibitors (TKIs) in cancer therapy is limited due to resistance. In this study, we identify a synergistic combination therapy involving TKIs for the treatment of triple negative breast cancer. By employing pairwise tyrosine kinase knockout CRISPR screens, we identify *FYN* and *KDM4* as critical targets whose inhibition enhances the effectiveness of TKIs, such as NVP-ADW742 (IGF-1R inhibitor), gefitinib (EGFR inhibitor), and imatinib (ABL inhibitor) both *in vitro* and *in vivo*. Mechanistically, treatment with TKIs upregulates the transcription of *KDM4*, which in turn demethylates H3K9me3 at *FYN* enhancer for *FYN* transcription. This compensatory activation of *FYN* and *KDM4* contributes to the resistance against TKIs. *FYN* expression is associated with therapy resistance and persistence by demonstrating its upregulation in various experimental models of drug-tolerant persisters and residual disease following targeted therapy, chemotherapy, and radiotherapy. Collectively, our study provides novel targets and mechanistic insights that can guide the development of effective combinatorial targeted therapies, thus maximizing the therapeutic benefits of TKIs.

Introduction

Tyrosine kinases have emerged as important drug targets in cancer therapy due to their druggability and pivotal roles in cell proliferation and survival(1). They are implicated in various aspects of cancer development(2), such as cell survival, proliferation, angiogenesis, and invasion, making them attractive targets for drug intervention. Consequently, tyrosine kinase inhibitors (TKIs) have gained considerable attention as primary agents for cancer treatment.

Triple negative breast cancer (TNBC) treatment has limited options for targeted therapy. TNBC, characterized by the absence of estrogen receptor, progesterone receptor, and HER2 expression, exhibits elevated activity of tyrosine kinases, including EGFR and IGF1R(3, 4). However, several clinical trials investigating TKIs, such as VEGFR inhibitors, EGFR inhibitors, and FGFR inhibitors, in TNBC treatment have yielded disappointing results due to inadequate efficacy. Therefore, it is crucial to comprehend the mechanisms underlying TNBC's suboptimal response to TKIs to enable the development of more effective targeted therapies against TNBC.

The therapeutic efficacy of TKIs is compromised by intrinsic and acquired resistance(5). For instance, EGFR inhibitor gefitinib extended the median progression-free survival by only five months compared to conventional

chemotherapy in non-small cell lung cancer (NSCLC) patients with EGFR mutation(6). Significant subset of drug resistance is driven by gene interactions that enable compensatory changes in signal transduction upon drug treatment. Compensatory activation of mitogenic signals, such as MET, PIK3CA amplification, and MAPK/ERK signaling activation, counterbalances the inhibition of EGFR by TKI osimertinib in a significant portion of NSCLC patients(7). Simultaneous inhibition of multiple signaling molecules that compensate for each other's loss is proposed as an effective strategy to overcome resistance to kinase inhibitor therapy, emphasizing the importance of combinatorial therapy(8).

Until recently, a highly scalable method for screening combinatorial therapy has been lacking. Combinatorial CRISPR screens have emerged as efficient tools to identify synergistic targets for combinatorial therapy. We and others recently developed combinatorial CRISPR screens to elucidate pairwise gene interactions(9-12). Our combinatorial genetic screen platform, combinatorial genetics *en Masse* (combiGEM), was successfully implemented to identify combinations of epigenetic regulators causing synthetic lethality in ovarian cancer cells (9).

In this study, we utilize CombiGEM-CRISPR technology to identify tyrosine kinase inhibitor combinations with synergistic effect in TNBC cell line and xenograft models for potential combinatorial therapy against TNBC. We highlight FYN as a key therapeutic target that, when inhibited, enhances the cytotoxic effect of inhibition of other tyrosine kinases (IGF1R, EGFR, and ABL2). Mechanistic studies reveal KDM4 as a crucial epigenetic regulator that demethylates H3K9me3 and transcriptionally upregulates *FYN* upon TKI treatment. *In vitro* and *in vivo* validation demonstrates the synergistic TNBC-shrinking effects of combining PP2, saracatinib (Src family kinase / FYN inhibitor) or QC6352 (KDM4 inhibitor) with TKIs. Additionally, we demonstrate the clinical significance of our findings by observing upregulation of *FYN* in various models of drug tolerant persisters and residual tumors after chemo-, radio-, or targeted therapy. Therefore, simultaneous targeting of *FYN-KDM4* and tyrosine kinase pathways through combinatorial therapy holds promise for effective therapy against TNBC.

Results

Pairwise tyrosine kinase knockout CRISPR screen reveals synergistic tyrosine kinase inhibition combination

For efficient translation of CRISPR screening data to drug combination, we selected 76 tyrosine kinases that

could be inhibited by at least one drug from the drug repurposing hub database (table S1) (13). For pairwise

CombiGEM library construction, we chose three guide RNAs from the optimized Brunello single guide RNA

(sgRNA) list(14) employing the iterative cloning method as previously described(9, 15). The resulting library

enabled screens of pairwise knockouts of the 76 tyrosine kinase genes, encompassing 54,289 sgRNA pairs

representing 3,003 pairwise gene disruptions (Fig. 1A). To validate our library, we performed next-generation

sequencing and confirmed that 99.5% (2,989/3,003) of gene pairs were represented by at least 6 pairs of

sgRNAs, with the $\log_{10}$ reads per million of 0.5 (Fig. S1A).

Triple negative breast cancer cell line MDA-MB-231 cells stably expressing Cas9 were transduced with the

lentiviral library at low multiplicity of infection (MOI) of 0.3. Genomic DNA was harvested 3 days after

transduction (designated as day 0[D0]), and 23 days after transduction (D20) (Fig. 1A) to perform PCR

amplification of sgRNA pairs for subsequent next-generation sequencing (NGS) analysis. sgRNAs, instead of

barcodes in our previous CombiGEM screens, were directly sequenced using paired-end sequencing to rule out

uncoupling of barcodes of sgRNAs and barcodes (Fig. S1B-C). We counted the occurrences of each sgRNA pair

in the NGS data and calculated the normalized log2 fold change in counts between the day 20 and day 0 samples

as the growth phenotype score Z (see Methods). The Z scores for the two permutations of an sgRNA pair

(r=0.50 between sgRNA-A + sgRNA-B and sgRNA-B + sgRNA-A pairs), the biological replicates, (r = 0.74

between replicates #2 and #3), and independent sgRNA pairs targeting the same set of genes were positively

correlated (r=0.3-0.72) (Fig. S1D-G).

Gene pairs that synergistically kill cells were identified by calculating gene interaction scores (GI). The GI

scores were derived by comparing the growth phenotype score Z resulting from the disruption of a gene pair

(Z_{A+B}, observed Z score) to the sum of Z scores obtained from the disruption of each gene individually within

the pair ($Z_{A+Con} + Z_{B+Con}$, expected Z score) (Fig. 1B). The expected and observed Z scores for each gene (or

sgRNA) pair exhibited a strong positive correlation (r=0.97 gene level, r=0.88 sgRNA level), suggesting that

most random pairwise combinations of tyrosine kinase perturbations show additive effects (Fig. 1B and S1H).

GI scores were calculated by quantifying each gene pair's normalized deviation from the quadratic fit of the

expected-observed Z score plot(10) (Fig.1C and Fig. S1I, see methods).

We selected thirty synthetic lethal gene pairs using cutoffs for gene level GI score <-2 and p<0.01 for GI scores

determined by RIGER analysis(16) and Z<-5 (Fig. 1D and dataset S1). Among these, the SRC-YES pair was

one of the strongest synthetic lethal gene pairings (GI= -3.95). Notably, SRC-YES belong to the same tyrosine

kinase family and are known to be functionally redundant and are expected to be synthetic lethal(17). These

findings provide evidence for the effectiveness of our screening approach in identifying synthetic lethal gene

pairs.

Synthetic lethal gene pairs are next validated by expressing the pair of sgRNAs targeting them. To achieve this,

we introduced lentiviral vectors carrying two distinct sgRNAs targeting the candidate synthetic lethal pairs, each

tagged with a different fluorescent protein (GFP and mCherry). MDA-MB-231 Cas9 cells were transduced with

the lentivirus at a low titer (MOI ~0.5), resulting in a mixed population of cells expressing either one or both

sgRNAs along with their respective fluorescent proteins (Figs. 2A and S2A). We monitored the decrease in the

number of GFP/mCherry double-positive cells to evludate synthetic lethality (see methods). We validated the

efficacy of the sgRNAs used in Figure 2A through the T7 endonuclease assay, which confirmed efficient gene

editing (Fig. S2B). Consistent with our CRISPR screening results, we observed that the disruption of six out of

eight synergistic target gene combinations led to a reduction in cell viability beyond what was predicted by the

Bliss independence model (Fig. 2B). Moreover, the relative viability of double knockout cells and the rate of

synergistic killing demonstrated a strong correlation with our screening data (r = 0.65 for both viability and

synergistic effect; Fig. S2C). Collectively, our findings provide compelling evidence that our screening approach

successfully identified synthetic lethal gene pairs with a high level of confidence.

FYN inhibition synergizes with IGF1R, EGFR, ABL2 inhibitions in cell killing

We noticed that several validated synergistic target gene pairs included *FYN* (e.g. *FYN*+IGF1R, *FYN*+EGFR,

and *FYN*+ABL2). Notably, network analysis of the 30 candidate synergistic tyrosine kinase pairs revealed that

FYN is one of the key nodes participating in synergistic interactions with multiple genes (Fig. 2C). Expression

of FYN, a member of Src family kinase, has been implicated in cancer malignancy including drug resistance(18-

21). Particularly, recent studies highlighted significant contribution of FYN in TNBC malignancy by promoting

epithelial-to-mesenchymal transition (EMT)(22, 23). Interestingly, we found that *FYN*, but not *SRC*, exhibited

significant upregulation in triple-negative breast cancer (TNBC) compared to other subtypes, as evidenced by

microarray data from primary tumor samples (24) and the cancer cell line encyclopedia (CCLE)(25) (Figs. 2D-

E). In contrast, other key nodes in figure 2C, including FGFR2, FRK and TEK were not expressed at

appreciable levels in MDA-MB-231 (log2(TPM+1) for TEK: 0.0704, FRK:0.124, FGFR2:0.227), and their

expressions were not significantly upregulated in TNBCs compared to other breast cancer subtypes (Fig. S3).

Therefore, we proceeded further in focusing on validating FYN as key candidate synthetic lethal gene. These

findings suggest that FYN could represent an attractive drug target for TNBC treatment. To investigate this

further, we assessed whether simultaneous inhibition of FYN by PP2, which selectively targets the SRC family

kinase inhibitor with the highest potency against FYN, in combination with other kinase inhibitors (TKIs), could

inhibit cancer cell growth(26). PP2 as a single agent significantly downregulated MDA-MB-231 cell viability

(Fig. S4A). Therefore, we focused on synergistic cell death by TKI combinations above additive effects by each

TKI. To this end, analysis using SynergyFinder plus(27)revealed that all TKI combinations involving PP2 and

NVP-ADW742 (IGF1R inhibitor), gefitinib (EGFR inhibitor) or imatinib (ABL inhibitor) synergistically

induced cell death in MDA-MB-231 cells (Fig. 2F). Dose-response curves demonstrated that co-treatment with

PP2 reduced the IC50 of the tested TKIs by 34-61%, indicating that PP2 sensitized cancer cells to TKI treatment

(Fig. 2G). Similar synergy was observed when TKI combinations included saracatinib, a SRC family kinase

inhibitor(28), in place of PP2 (Fig. S4B). Moreover, specific ablation of *FYN*, but not *SRC*, sensitized cells to

TKIs, highlighting the critical role of FYN as a member of SRC kinase family responsible for TKI resistance

(Figs. 2H and S4C). Ablation of FYN itself did not significantly decrease cell viability (Fig. S4D). Importantly,

we observed similar synergy between the same drug combinations in other TNBC cell lines, including Hs578T,

HCC1143, HCC1395, and HCC1937 cells (Figs. S4E-H). Further assessment using live-dead and BrdU assays

revealed that both the PP2+NVP-ADW742 and PP2+gefitinib combinations synergistically induced cell death

while inhibiting cell growth (Fig. 2I).

Persistent activation of MAPK pathway and PI3K-AKT pathway has been associated with TKI resistance in

various cancers(5). Therefore, we investigated which downstream pathways were involved in sensitizing cells to

TKI treatment. Notably, the p38 MAPK was significantly attenuated following treatment with either PP2 or

saracatinib treatment (Fig. 2J). Previous studies with imatinib resistant CML cells identified ERK signaling as

critical downstream of FYN activation(19, 20). However, FYN inhibition failed to significantly downregulate

phosphorylated ERK level upon imatinib treatment, indicating downstream signals of FYN leading to drug

resistance may be context dependent. Genetic ablation of *FYN* similarly reduced p38 activation (Fig. 2K).

Attenuation of p38 activity was also observed in an independent TNBC cell line, Hs578T (Figs. S4I-J).

Importantly, treatment of p38 MAPK pathway inhibitor SB203580 markedly sensitized cells to TKI treatment

(Fig. 2L), while SB203580 as single agent did not significantly change cell viability (Fig. S4K)

FYN mRNA is induced upon TKI treatment in KDM4 dependent manner

Our discovery that inhibition of FYN synergizes with multiple TKIs possessing distinct target profiles suggests

that FYN may play a role in general resistance mechanisms against TKI therapy. Consistently, we observed an

increase in both protein and mRNA levels of *FYN* following TKI treatment, indicating that upregulation of FYN

confers compensatory survival signal in TKI-treated cells (Fig. 3A-B). The phosphorylation levels of FYN was

increased proportional to FYN protein level, indicating specific kinase activity of FYN did not change (Fig.

S5A). Previous study suggested that increased expression of EGR1 transcription factor is responsible for FYN

mRNA accumulation in imatinib resistant CML(18). Consistently, EGR1 expression was increased upon TKI

treatment in MDA-MB-231 cells. However, EGR1 expression was not increased in TKI treated MDA-MB-231

cells, nor did its knockout significantly downregulated FYN mRNA levels (Fig. S5B). To elucidate the

mechanisms underlying the accumulation of *FYN*, we treated MDA-MB-231 cells with inhibitors targeting key

epigenetic modifiers and assessed their synergistic effects with NVP-ADW742 in cell killing, as well as their

impact on *FYN* mRNA accumulation. Multiple drugs, including pinometostat (DOT1L inhibitor(29)),

tazemetostat (EZH2 inhibitor(30)), A366 (G9a inhibitor(31)) and GSK-J4 (KDM6 inhibitor(32)) strongly

decreased cell viability upon TKI treatment (Fig. S5C). As the increase in *FYN* mRNA is responsible for TKI

resistance, we reasoned that the drug that directly affect FYN mRNA level and hence cell viability should be an

inhibitor of epigenetic regulator that enhances transcription. To this end, we focused on pinometostat and GSK-

J4 for further validations. Intriguingly, while GSK-J4 decreased FYN mRNA upon NVP-ADW742 treatment,

pinometostat failed to decrease it (Fig. S5D).. Consistent with this, treatment with NVP-ADW742 increased the

expression of most members of the jumonji domain histone demethylase family (Fig. 3C). This observation is

consistent with a previous study on taxane-resistant H1299 lung cancer cells(33), suggesting that histone

demethylases may play critical roles in activating a drug resistance gene program. However, the ablation of

KDM6, the primary targets of GSK-J4, failed to significantly decrease *FYN* expression (Fig. S5E). GSK-J4 is

known to inhibit other jumonji domain histone demethylase family proteins including KDM4 and KDM5(34).

Therefore, we tested the possibility that other histone demethylase may be involved in regulating *FYN*

expression. Among jumonji domain histone demethylases, *KDM4*, and to a lesser extent *KDM3*, were the only

gene family members whose ablation inhibited *FYN* upregulation and p38 activation upon TKI treatment (Figs.

3D and S5F). Ablation of KDM5, which has been shown to induce drug tolerance in cancer cells(35), did not

significantly alter *FYN* expression (Fig. S5G). Like NVP-ADW742 treatment, gefitinib treatment increased

KDM4 demethylase levels (Fig. S5H). We also analyzed two independent TNBC organoids obtained from

primary tumors and found concurrent upregulation of *KDM4* with *FYN* mRNAs upon NVP-ADW742 and

gefitinib treatment (Fig. S5I). Critically, time course experiment with NVP-ADW742 treated MDA-MB-231

revealed that accumulation of KDM4A protein preceded FYN protein (Fig. 3E), suggesting that KDM4A

accumulation may be responsible for FYN accumulation. Both KDM3 and KDM4 demethylates methylated

H3K9, thereby promoting the opening heterochromatin for transcription(36). Remarkably, expression of

KDM4A, the most abundantly expressed gene among KDM4 demethylases in TNBC cell lines (Fig. S6A) was

enriched in TNBC compared to other breast cancer subtypes (Fig. 3F) and was positively correlated with *FYN*

expression in CCLE database, suggesting that KDM4 regulates *FYN* mRNA levels (Fig. 3G). Genetic ablation

of *KDM3* or *KDM4* (Fig. S6B-C) decreased *FYN* and p38 activity. Also, genetic ablation of KDM3 or KDM4

sensitizing MDA-MB-231 cells to TKIs (Figs. 3H-I). The Ablation of KDM3 or KDM4 only had modest but

statistically insignificant effect on cell viability (Fig. S6D). Likewise, treatment of KDM4 inhibitor QC6352(37)

synergized with TKIs in killing MDA-MB-231 cells (Fig. 3J). QC6352 treatment also significantly attenuated

FYN accumulation upon NVP-ADW742 treatment (Fig. 3K-L). This was consistent with the RNA sequencing

data results in the previous study with breast cancer stem cells treated with QC6352(38). Specifically, *FYN* was

the most significantly downregulated SRC family kinase upon QC6352 treatment (Fig. 3M). Analysis of

chromatin IP (ChIP) sequencing data from the same study revealed KDM4A enrichment near FYN promoter;

and QC6352 treatment increased H3K9me3 enrichment at the same locus (Fig. 3N). Indeed, this FYN promoter

locus exhibited a reduction in H3K9me3 following NVP-ADW742 treatment, while QC6352 treatment restored

H3K9me3 enrichment (Fig. 3O). This finding suggests that KDM4 may directly demethylate H3K9me3 at FYN

promoter to upregulate *FYN* transcription. In contrast, H3K27me3 marks, which is demethylated by KDM6

family demethylases, were not significantly changed at FYN promoter upon NVP-ADW742 treatment (Fig.

S6E). *FYN* accumulation and resistance to TKIs were also confirmed to be attenuated by QC6352 treatment in

other independent TNBC cell lines (Figs. S6F-G).

FYN/KDM4 inhibition synergizes with TKI treatment *in vivo*

We proceeded to investigate the potential clinical application of our synthetic lethal gene pairs as combinatorial

therapy by assessing the *in vivo* efficacy of pharmacological interventions targeting these gene pairs using

MDA-MB-231 xenograft models. Strikingly, co-treatment of saracatinib and NVP-ADW742 synergistically

reduced tumor size, whereas treatment with either agent alone was ineffective in slowing tumor growth (Fig.

4A). All treatment groups exhibited minimal changes in body weight, indicating that the overall health of the

animals was not adversely affected by the combination treatment (Fig. S7A). Saracatinib-gefitinib combination

was not tested as saracatinib can inhibit EGFR(28). Similarly, KDM4 inhibitor QC6352 synergized with

gefitinib in reducing MDA-MB-231 xenograft tumor growth without causing overt changes in animal health

(Figs. 4B and S7B). Additionally, the expression levels of *FYN* and *KDM4A* were found to be correlated with

poor prognosis in a previously reported breast cancer cohort(24), highlighting the potential of targeting these

two genes as therapeutic targets for TNBC (Fig. 4C). Collectively, our results demonstrate that upregulation of

KDM4 upon TKI treatment reduces H3K9me3 mark in *FYN* enhancer, thereby increasing *FYN* expression and

promoting cell survival under TKI treatment (Fig. 4D).

FYN is associated with drug tolerant persister phenotype

The observed epigenetic alterations in regulators conferring resistance to multiple cancer drugs closely resemble

non-genetic changes associated with the generation of drug-tolerant persisters(35). Indeed, prolonged incubation

of MDA-MB-231 cells treated with TKIs or conventional chemotherapy drugs such as doxorubicin or paclitaxel

resulted in increased levels of *FYN* (Fig. 5A). Curiously, *KDM4A* expression was only upregulated upon

treatment with NVP-ADW742 and gefitinib, suggesting that while *FYN* upregulation is a general feature of drug

tolerant cells, the mechanism of *FYN* upregulation may vary depending on the specific drug being used.

Analysis of previously published RNA sequencing data from a series of osimertinib tolerant EGFR mutated lung

cancer cell lines(39) revealed higher expression levels of *FYN* and *KDM4A* in the drug persisters, but not SRC

(Fig. 5B). Consistently, we confirmed upregulation of *FYN* at both the protein and mRNA levels in gefitinib and

osimertinib resistant PC9 and HCC827 cells (Figs. 5C-D). Pharmacological inhibition of FYN or

downregulation of *FYN* expression through inhibition of KDM4 sensitized gefitinib resistant PC9 cells to EGFR

inhibitor, suggesting that *FYN-KDM4* are responsible for gefitinib resistant phenotype in this cell line (Figs.5E-

G).

Importantly, upregulation of *FYN* has been consistently observed in multiple independent studies involving

drug-tolerant cancer cell lines and patient-derived xenografts treated with various drugs that have distinct target

profiles, including TKIs (lapatinib, a HER2 inhibitor, against HER2 positive breast cancer(40)) and

chemotherapy drugs (irinotecan, topoisomerase inhibitor against colorectal cancer(41); gemcitabine against

pancreatic cancer(42); and carboplatin against high grade serous ovarian carcinoma(43)) (Fig. 5H). Moreover,

enrichment of *FYN* has also been observed in residual disease following chemotherapy, including neoadjuvant

chemostherapy plus bevacizumab treated HER2 negative breast cancer(44), neoadjuvant chemoradiotherapy

combined plus atezolizumab treated esophageal cancer(45), and chemoradiotherapy treated colorectal

cancer(46), indicating its potential role in mediating drug tolerance during chemotherapy (Fig. 5I). While the

causal relationship between FYN expression drug tolerance in response to various therapeutic interventions

warrants further study, these evidence suggest that FYN expression is associated with drug tolerance. Notably,

an analogous increase in *KDM4* was not consistently observed across all tumor models tested in Figures 5H-I

(Fig. S8A-B). This suggests that, as previously noted in Figure 5A, while *FYN* serves as a general mediator of

drug tolerance, the specific mechanisms underlying its upregulation may vary depending on the cancer type and

the drug being administered. Taken together, these lines of evidence further support our findings in TNBC cell

lines and suggest that *FYN* acts as a common mediator of drug tolerance.

Discussion

In this study, we employed combinatorial CRISPR screening to identify combinations of TKIs that exhibit

synergistic effects in eliminating triple-negative breast cancer (TNBC). We discovered and validated that

concurrent targeting of FYN, along with other tyrosine kinases such as IGF1R, EGFR or ABL2 can

synergistically eradicate TNBC and impede cancer growth. Our findings also provide evidence that the

transcriptional upregulation of *FYN*, facilitated by the activation of KDM4 histone demethylases, confers

resistance and persistence to TKIs. Upregulation of *FYN* is a general feature of drug tolerant cancer cells,

suggesting the association of *FYN* expression with drug resistance and tumor recurrence after treatment.

This research provides basis for breakthrough combinatorial therapy achieving effective targeted therapy with

minimal risk of developing resistance. Our combinatorial CRISPR screening demonstrates that treatment with

TKIs or histone demethylase inhibitors enhances the sensitivity of cells to other TKIs. Consequently, drug

combinations exhibit a more potent inhibition of cancer growth than the simple sum of the therapeutic effects of

individual drugs. Furthermore, synergistic drug combinations enable a reduction in the dosage of each drug,

with minimal compromise in therapeutic efficacy. Such combinations yield a therapeutic response comparable to

that achieved with significantly higher doses of each individual agent. We anticipate that combinatorial therapy

has the potential to mitigate side effects by minimizing the dosage of each drug, thus widening the therapeutic

window. Further studies should elucidate the downstream mechanisms by which FYN upregulation contributes

to drug tolerance. SRC family kinases are known to upregulate multiple signaling pathways including ERK,

AKT and p38 pathway(47). Although our study showed that, at least in MDA-MB-231 cell line, FYN depends

on p38 activity for TKI resistance (figs. 2J-L), it should further be shown while this downstream mechanism is

generalizable in other cancers. The downstream mechanism of p38 that contributes to drug resistance would also

be of great interest to identify novel therapeutic approaches minimizing drug tolerance. Also, our combinatorial

CRISPR screen results warrant further studies with other synthetic lethal gene combinations other than those

involving FYN that are not deeply investigated in this study. Particularly, FGFR2, TEK, FRK identified as key

nodes in figure 2C may be of particular interest as they are also associated with cancer cell survival(48, 49).

It is intriguing to observe that *FYN* is specifically upregulated in various models of drug resistance and tolerance.

SRC family kinases, which includes FYN, have been linked to drug resistance(47, 50). In line with this,

phosphoproteomic analysis of neoadjuvant chemotherapy resistant TNBC patient derived xenografts showed

upregulated SRC family kinase networks including kinases and their substrates and adaptors(51). Our findings

reveal that *FYN* is specifically upregulated at the mRNA level possibly through epigenetic regulations, providing

further depth to our understanding of drug resistance in cancer therapy. The epigenetic reprogramming of the

drug tolerant cells may be distinct depending on the tumor type or the therapeutic interventions, as KDM4A,

which we show is increased upon TKI treatment in TNBC cell line in figure 3, is not significantly regulated in

other cancers analyzed in figure 5. Therefore, the context dependent mechanisms underlying FYN upregulation,

and its essentiality in constituting drug tolerance remains as subjects for further study.

Furthermore, our work highlights the significance of histone demethylases in TKI resistance. Numerous histone

demethylases have been implicated in drug resistance and tolerance across different cancer drug types. For

instance, the KDM5 family of H3K4 demethylases has been associated with the drug-tolerant persister

phenotype against multiple TKIs(35). In our study, we identify *KDM4* as a critical factor in the generation of

drug-tolerant persisters in breast cancer. *KDM4* is known to be upregulated in various cancers, including breast

cancer, and promotes key malignant traits. Previous studies have demonstrated the essential role of KDM4 in

induced pluripotency through its interaction with pluripotency factors(52). These findings suggest that an

KDM4 inhibitor could be a promising therapeutic target with specific activity against cancer stem cells.

Consistently, specific inhibitors targeting KDM4 have recently been developed and shown to inhibit the

generation of breast cancer stem cells(38). The mechanisms underlying KDM4 upregulation upon drug

treatment is currently unclear and remains as subjects for further study. Nevertheless, given our discoveries

regarding the involvement of *KDM4* in drug resistance in breast and lung cancer, the development of novel

drugs targeting KDM4 holds significant therapeutic potential.

Materials and methods

Cell Culture. HEK293T, MDA-MB-231, Hs578T cells were obtained from American Type Cell Culture (ATCC)

HCC1143, HCC1395, HCC1937 were obtained from Korean cell line bank. HEK293T and MDA-MB-231 were

grown in DMEM supplemented with 10% FBS and penicillin/streptomycin. Hs578T and HCC1143, HCC1395,

HCC1937 were grown in RPMI1640 supplemented with 10% FBS. Parental and gefitinib-, and Osimertinib-

resistant PC9 and HCC827 cells were kind gifts from Jae Cheol Lee (Asan Medical Center, Seoul, Korea) and

were grown in RPMI1640 supplemented with 10% FBS. Gefitinib and osimertinib resistant cells are maintained

in the presence of 1μM gefitnib and 0.5μM osimertinib, respectively. Gefitinib resistant derivatives of PC9 and

HCC827 cells were generated as described previously by treating cells with escalating dose of gefitinib (53). All

cell culture medium and supplements are purchased from Welgene Inc.

Combinatorial library construction. Combinatorial library was constructed as previously described(15). The

sgRNAs used in the screens were cloned in pAWp28 storage vector in two versions: one version containing

human U6 driven sgRNA with wild type scaffold, and another containing mouse U6 driven sgRNA with cr2

variant scaffold. The sgRNA expression cassette consisting of U6 promoter and sgRNA were subject to one-pot,

iterative cloning into lentiviral pTK799 vector using BglII-MfeI restriction sites flanking the sgRNA expression

cassette and BamHI-EcoRI sites in pTK799. pTK799 vector is derived from pAWp12(15) by replacing CMV-

GFP selectable marker to EFS-Puro.

Combinatorial CRISPR screening procedure. Lentivirus was generated in HEK293T cells by transfecting

lentiviral transfer vector, and helper vectors (psPAX2, and pVSV-G) using Fugene HD (Promega). Lentiviral

supernatant was collected 48 hours after transfection, and was frozen and stored in -80C. The appropriate titer

for lentiviral transduction was determined by transducing MDA-MB231 cells with two-fold serial dilution of

lentiviral supernatant, selecting with puromycin 2 days after transduction for 2 days, and determining cell

viability with AQuaeous one cell viability, MTS assay (Promega). After determining the titer of lentiviral

supernatant, 100 million MDA-MB231 cells carrying constitutively expressed Cas9 were transduced with

CombiGEM library at MOI of 0.3. The expected initial coverage is 100 million x 0.3/ (54,289 different sgRNA

combinations)= 553. Three days after transduction, the cells were either harvested as day 0 sample or selected

with 2µg/mL puromycin (Invitrogen). Cells were treated with benzonase before harvesting to minimize

carryover of plasmid DNA in lentiviral supernatant. Cells were grown in the presence of 2µg/mL puromycin for

20 days before harvesting.

The genomic DNA of harvested cells were isolated using Blood & Cell Culture Maxi kit (Qiagen). The PCR

amplicon spanning the two sgRNAs were generated with PCR using Q5 High Fidelity DNA polymerase (New

England Biolabs) and the following primers:

F: CAAGCAGAAGACGGCATACGAGAT CCTAGTAACTATAGAGGCTTAATGTGCG

R: AATGATACGGCGACCACCGAGATCTACAC NNNNNN ACACGAATTCTGCCGTGGATCCAA

The six nucleotides described as "NNNNNN" in reverse primer above represents unique index to identify

biological replicates in multiplexed NGS run.

The PCR protocol involves 60 seconds of initial DNA denaturation at 98C, and 20 cycles of 10 seconds

denaturation at 98C, 10 seconds annealing at 67C, and 120 seconds elongation at 72C. All genomic DNA

isolated were used in PCR reaction at concentration of 40µg/mL. All PCR products were combined and

precipitated with isopropanol at room temperature. The precipitated DNA was resuspended in 400µL EB buffer

(Qiagen) and gel purified. The purified PCR products were sent for NGS by NextSeq500 paired end sequencing

with the following sequencing primers:

Forward read: GGACTAGCCTTATTTGAACTTGCTATGCAGCTTTCTGCTTAGCTCTCAAAC

Forward   index   read:

CGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTAACTTGCTATTTCTAGCTCTAAAAC

Reverse read: GCA CCG AGT CGG TGC TTT TTT GGA TCC ACG GCA GAA TTC GTGT

The raw GI scores calculated as deviation from the quadratic fit of the expected-observed Z score plot. The GI

scores were normalized by dividing the raw GI scores with the standard deviation of the GI scores obtained

from the 200 nearest neighbors in terms of expected Z scores (Fig. S1I)(11).

Data analysis. The sgRNA sequences were identified and their occurrences were counted with C++ script deposited in Github (https://github.com/tackhoonkim/combinatorial-CRISPR-screens-2023).

Validation of screens using sgRNAs. Individual sgRNA was cloned to either pTK1329, and pTK1336 that are both derived from pAWp12 with EFS-GFP and EFS-mCherry, respectively, as selectable markers. Validation of synthetic lethality between gene A and B were analyzed by transducing MDA-MB-231 Cas9 cells with four combinations of lentiviral supernatant pairs (MOI ~ 0.5 each) containing (1) GFP-sgA and mCherry-sgB; (2) GFP-sgA and mCherry-sgCon; (3) GFP-sgCon and mCherry-sgB; (4) GFP-sgCon and mCherry-sgCon. The fraction of GFP/mCherry double positive cells were analyzed using BD Accuri C6 and its accompanying software. The expected fold change in sgA+sgB were calculated as $FC_{sgA+sgCon}$ x $FC_{sgCon+sgB}$, where FC is normalized fold change in fraction of GFP/mCherry double positive cells relative to those transduced with GFP-sgCon and mCherry-sgCon.

MTT Cell viability assay. Cells were seeded at 1000-2000 cells/well in 96 well plate. Tyrosine kinase inhibitors at indicated combination of dose were treated 12 hours after seeding, and cells were grown for 3 days. The relative viability was measured by EzCytox cell viability assay (Dojindo). The absorbance at 450nm wavelength was measured using EnVision multimode plate reader (PerkinElmer).

Cell death and cell proliferation assay. Cells were incubated with tyrosine kinase inhibitors for 48 hours. Cell proliferation was quantified with BrdU assay using FITC conjugated BrdU antibody (Biolegend, 364103) and propidum iodide/RNase A solution (Cell Signaling), analyzed with BD Accuri C6 and accompanied software. Cell death was quantified with Live-Dead cell staining kit (Molecular Probes) by flow cytometry analysis using BD Accuri C6 and accompanied software.

Western blot analysis. Cells were treated with drugs for 48 hours unless otherwise indicated. Western blots were performed as previously reported. Briefly, Cells were lysed in RIPA buffer (50mM Tris-HCl pH 7.5, 150mM sodium chloride, 0.1% sodium dodecyl sulfate, 0.5% sodium deoxycholate, 1% NP-40, 1mM EDTA) supplemented with protease inhibitor (aprotinin, leupeptin, pepstatin A, and phenylmethylsulfonyl fluoride [PMSF]) and phosphatase inhibitor (sodium fluoride and sodium orthovanadate) cocktail. Antibodies used for western blot analysis were: anti-SRC (Santa Cruz biotechnology, mouse sc-), anti-FYN (Cell Signaling Technology, rabbit #4023), anti-phospho-FYN Y530 (Invitrogen, PA5-36644), anti-GAPDH (Cell Signaling Technology, rabbit #5174), anti-KDM4A (Bethyl Laboratories, rabbit A300-861A), anti-phospho-ERK

T202/Y204 (Cell Signaling Technology, rabbit #9101), anti-ERK (Cell Signaling Technology, rabbit #9102),

anti-phospho-STAT3 Y705 (Cell Signaling Technology, rabbit #9145), anti-STAT3 (Cell Signaling Technology,),

anti-phospho-AKT (Cell Signaling Technology, mouse #4051), anti-AKT (Cell Signaling Technology,), anti-

phospho-p38 (Cell Signaling Technology,), anti-p38 (Cell Signaling Technology,).

RT-qPCR analysis. RNA is extracted from cultured cells with Trizol (Invitrogen) according to the

manufacturer's instructions. The precipitated RNA pellet was resuspended in RNase free water (Enzynomics),

and was subject to reverse transcription with M-MLV reverse transcriptase (Enzynomics) at 37C for 2 hours,

followed by heat inactivation at 95C for 5 minutes. The resulting cDNA was used for quantitative real time PCR

using TOPreal SYBR Green qPCR premix (low ROX, Enzynomics) reagent and CFX96 Real-Time PCR

detection system (Bio-Rad).

Xenograft assay. All animal experiments were approved by IACUC of Korea Institute of Science and

Technology (KIST). Six-week-old female nude mice were injected with 5×10^6 MDA-MB-231 cells suspended

in 1:1 (w/w) mixture of PBS and growth factor reduced Matrigel (Corning) in fourth inguinal mammary fat pad.

Starting two weeks after tumor cell injection, saracatinib (50mg/kg mouse body weight, MedChemExpress),

NVP-ADW742 (20mg/kg, Sigma), gefitinib (20mg/kg, MedChemExpress), QC6352 (10mg/kg,

MedChemExpress) in 45% saline+40% polyethyleneglycol 300 (sigma)+5% Tween-80 (sigma)+5% DMSO

(sigma) were injected intraperitoneally every 24 hours for two weeks. Tumor volume was measured by digital

caliper and calculated as $(width)^2 \times length \times 0.5$.

Public database analyses. Gene Expression Omnibus (GEO) data with breast cancer cohort (GSE25066(24))

were analyzed using web based platform Cancer Target Gene Screening (https://ctgs.biohackers.net)(54). Cancer

Cell Line Encyclopedia (CCLE) data were analyzed using depmap R package version 1.14. The list of GEO data

used for analysis are listed in table S2.

Primary TNBC organoid culture and drug treatment. Specimens was obtained from enrolled patients with

TNBC breast cancer with IRB approval. Tumor tissue was collected and transferred using cold RPMI1640

media at the National Cancer Center (Goyang, Republic of Korea). Then the tissues were dissected with a blade

on Petri dish and enzymatically digested with dissociated kit (Multi Tissue Dissociation Kit 1, 130-110-201) by

gentle MACS Dissociators (Miltenyi Biotec, Germany). After cell counting, 1×10^5 cells were embedded in 40

μl of Matrigel (Corning) and seeded into each well of a 24-well cell culture plate. After the matrigel was

solidified, 500 μL organoid medium supplemented with advanced DMEM/F12 medium (Invitrogen, USA), B27

(Invitrogen, USA, 17504), 1.25 mM N-acetylcysteine (Sigma, USA, A9165), 5 ng/mL EGF (PeproTech, USA,

AF-100-15), 20 ng/mL FGF-10 (Peprotech, 100-26), 5ng/ml FGF7 (Peprotech, AF-100-19), 50ng/ml R-spondin

1 (Qkine, Qk006), 5 mM nicotinamide (Sigma, N0636), 500 nM A83-01 (Tocris, 2939), 1X GlutaMAX (Gibco,

35050-061), 100ug/ml primocin (Invivogen, ant-pm), 10mM HEPES (Gibco, 15630-080), 1X Noggin (U-

Protein Express BV, N002), 1X ITS (Gibco, 12585014), 100nM β-estradiol (Sigma, E2758), 1ug/ml

Hydrocortisone (Sigma, H0888), 5nM Heregulin (PeproTech, AF-100-03), 500nM SB202190 (R&D system,

1264), 10uM Y27632 (TOCRIS, 1254) as described by Clevers and colleagues (55), was added to each well and

the cells grown under standard culture conditions (37 ℃, 5% CO2).

Acknowledgments.

We thank BioMicro Center of MIT for sequencing analysis. We thank Jae Cheol Lee (Asan Medical Center,

Korea) for providing gefitinib resistant lung cancer cell lines. This work was supported by Korea Institute of

Science and Technology (KIST) Institutional Programs (2E32331 to T.K.); and National Research Foundation of

Korea, funded by the Korean government (MSIT) (2021R1A2C1093499 to T.K., 2020M3A9A5036362 to

S.Y.K).

Author contributions. T.K., B.-S.P., H.J. and J.K. performed experiments. T.K. and T.K.L. supervised the

research. S.H. S.-Y.J. and S.Y.K. performed experiments with TNBC patient derived organoids. D.K., S.K.L.

generated and provided osimertinib resistant HCC827 cell line.

Conflict of Interest Statement

The authors declare no conflict of interest

Ethics Statement

All experiments with human tumor organoids were conducted in accordance with the requirements of the

National Cancer Center Institutional Review Board (IRB).

Availability of data and materials: The NGS data for CRISPR screening results are available under NCBI

SRA accession code PRJNA976939.

References

1.   Wu P, Nielsen TE, Clausen MH. Small-molecule kinase inhibitors: an analysis of FDA-
approved drugs. Drug Discovery Today. 2016;21(1):5-10.

2.   Hanahan D, Weinberg Robert A. Hallmarks of Cancer: The Next Generation. Cell.
2011;144(5):646-74.

3.   Li Y, Zhan Z, Yin X, Fu S, Deng X. Targeted Therapeutic Strategies for Triple-Negative
Breast Cancer. Frontiers in Oncology. 2021;11.

4.   Litzenburger BC, Creighton CJ, Tsimelzon A, Chan BT, Hilsenbeck SG, Wang T, et al. High
IGF-IR Activity in Triple-Negative Breast Cancer Cell Lines and Tumorgrafts Correlates with
Sensitivity to Anti–IGF-IR Therapy. Clinical Cancer Research. 2011;17(8):2314-27.

5.   Morgillo F, Della Corte CM, Fasano M, Ciardiello F. Mechanisms of resistance to EGFR-
targeted drugs: lung cancer. ESMO Open. 2016;1(3):e000060.

6.   Maemondo M, Inoue A, Kobayashi K, Sugawara S, Oizumi S, Isobe H, et al. Gefitinib or
Chemotherapy for Non–Small-Cell Lung Cancer with Mutated EGFR. New England Journal of
Medicine. 2010;362(25):2380-8.

7.   Leonetti A, Sharma S, Minari R, Perego P, Giovannetti E, Tiseo M. Resistance mechanisms
to osimertinib in EGFR-mutated non-small cell lung cancer. British Journal of Cancer.
2019;121(9):725-37.

8.   Jin H, Wang L, Bernards R. Rational combinations of targeted cancer therapies:
background, advances and challenges. Nature Reviews Drug Discovery. 2023;22(3):213-34.

9.   Wong ASL, Choi GCG, Cui CH, Pregernig G, Milani P, Adam M, et al. Multiplexed barcoded
CRISPR-Cas9 screening enabled by CombiGEM. Proceedings of the National Academy of Sciences.
2016;113(9):2544-9.

10.   Horlbeck MA, Xu A, Wang M, Bennett NK, Park CY, Bogdanoff D, et al. Mapping the
Genetic Landscape of Human Cells. Cell. 2018;174(4):953-67.e22.

11.   Han K, Jeng EE, Hess GT, Morgens DW, Li A, Bassik MC. Synergistic drug combinations for
cancer identified in a CRISPR screen for pairwise genetic interactions. Nature Biotechnology.
2017;35(5):463-74.

12.   Shen JP, Zhao D, Sasik R, Luebeck J, Birmingham A, Bojorquez-Gomez A, et al.
Combinatorial CRISPR–Cas9 screens for de novo mapping of genetic interactions. Nature Methods.
2017;14(6):573-6.

13.   Corsello SM, Bittker JA, Liu Z, Gould J, McCarren P, Hirschman JE, et al. The Drug
Repurposing Hub: a next-generation drug library and information resource. Nature medicine.

2017;23(4):405-8.

14.   Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, et al. Optimized
sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nature
Biotechnology. 2016;34(2):184-91.

15.   Wong ASL, Choi GCG, Cheng AA, Purcell O, Lu TK. Massively parallel high-order
combinatorial genetics in human cells. Nat Biotech. 2015;33(9):952-61.

16.   Luo B, Cheung HW, Subramanian A, Sharifnia T, Okamoto M, Yang X, et al. Highly parallel
identification of essential genes in cancer cells. Proceedings of the National Academy of Sciences.
2008;105(51):20380.

17.   Stein PL, Vogel H, Soriano P. Combined deficiencies of Src, Fyn, and Yes tyrosine kinases
in mutant mice. Genes & Development. 1994;8(17):1999-2007.

18.   Irwin ME, Johnson BP, Manshouri R, Amin HM, Chandra J. A NOX2/Egr-1/Fyn pathway
delineates new targets for TKI-resistant malignancies. Oncotarget. 2015;6(27).

19.   Fenouille N, Puissant A, Dufies M, Robert G, Jacquel A, Ohanna M, et al. Persistent
Activation of the Fyn/ERK Kinase Signaling Axis Mediates Imatinib Resistance in Chronic
Myelogenous Leukemia Cells through Upregulation of Intracellular SPARC. Cancer Research.
2010;70(23):9659-70.

20.   Airiau K, Turcq B, Mahon F-X, Belloc F. A new mechanism of resistance to ABL1 tyrosine
kinase inhibitors in a BCR-ABL1-positive cell line. Leukemia Research. 2017;61:44-52.

21.   Grosso S, Puissant A, Dufies M, Colosetti P, Jacquel A, Lebrigand K, et al. Gene expression
profiling of imatinib and PD166326-resistant CML cell lines identifies Fyn as a gene associated
with resistance to BCR-ABL inhibitors. Molecular Cancer Therapeutics. 2009;8(7):1924-33.

22.   Lee G-H, Yoo K-C, An Y, Lee H-J, Lee M, Uddin N, et al. FYN promotes mesenchymal
phenotypes of basal type breast cancer cells through STAT5/NOTCH2 signaling node. Oncogene.
2018;37(14):1857-68.

23.   Xie Y-G, Yu Y, Hou L-K, Wang X, Zhang B, Cao X-C. FYN promotes breast cancer
progression through epithelial-mesenchymal transition. Oncol Rep. 2016;36(2):1000-6.

24.   Hatzis C, Pusztai L, Valero V, Booser DJ, Esserman L, Lluch A, et al. A Genomic Predictor of
Response and Survival Following Taxane-Anthracycline Chemotherapy for Invasive Breast Cancer.
JAMA. 2011;305(18):1873-81.

25.   Ghandi M, Huang FW, Jané-Valbuena J, Kryukov GV, Lo CC, McDonald ER, et al. Next-
generation characterization of the Cancer Cell Line Encyclopedia. Nature. 2019;569(7757):503-8.

26.   Hanke JH, Gardner JP, Dow RL, Changelian PS, Brissette WH, Weringer EJ, et al. Discovery
of a Novel, Potent, and Src Family-selective Tyrosine Kinase Inhibitor: STUDY OF Lck- AND FynT-
DEPENDENT T CELL ACTIVATION. 1996;271(2):695-701.

27.   Zheng S, Wang W, Aldahdooh J, Malyutina A, Shadbahr T, Tanoli Z, et al. SynergyFinder
Plus: Toward Better Interpretation and Annotation of Drug Combination Screening Datasets.
Genomics, Proteomics & Bioinformatics. 2022;20(3):587-96.

28.   Green TP, Fennell M, Whittaker R, Curwen J, Jacobs V, Allen J, et al. Preclinical anticancer

activity of the potent, oral Src inhibitor AZD0530. Molecular Oncology. 2009;3(3):248-61.

29.   Daigle SR, Olhava EJ, Therkelsen CA, Basavapathruni A, Jin L, Boriack-Sjodin PA, et al.
Potent inhibition of DOT1L as treatment of MLL-fusion leukemia. Blood. 2013;122(6):1017-25.

30.   Knutson SK, Warholic NM, Wigle TJ, Klaus CR, Allain CJ, Raimondi A, et al. Durable tumor
regression in genetically altered malignant rhabdoid tumors by inhibition of methyltransferase
EZH2. Proceedings of the National Academy of Sciences. 2013;110(19):7922-7.

31.   Sweis RF, Pliushchev M, Brown PJ, Guo J, Li F, Maag D, et al. Discovery and Development
of Potent and Selective Inhibitors of Histone Methyltransferase G9a. ACS Medicinal Chemistry
Letters. 2014;5(2):205-9.

32.   Kruidenier L, Chung C-w, Cheng Z, Liddle J, Che K, Joberty G, et al. A selective jumonji
H3K27 demethylase inhibitor modulates the proinflammatory macrophage response. Nature.
2012;488(7411):404-8.

33.   Dalvi MP, Wang L, Zhong R, Kollipara RK, Park H, Bayo J, et al. Taxane-Platin-Resistant
Lung Cancers Co-develop Hypersensitivity to JumonjiC Demethylase Inhibitors. Cell Reports.
2017;19(8):1669-84.

34.   Heinemann B, Nielsen JM, Hudlebusch HR, Lees MJ, Larsen DV, Boesen T, et al. Inhibition
of demethylases by GSK-J1/J4. Nature. 2014;514(7520):E1-E2.

35.   Sharma SV, Lee DY, Li B, Quinlan MP, Takahashi F, Maheswaran S, et al. A Chromatin-
Mediated Reversible Drug-Tolerant State in Cancer Cell Subpopulations. Cell. 2010;141(1):69-80.

36.   Hyun K, Jeon J, Park K, Kim J. Writing, erasing and reading histone lysine methylations.
Experimental & Molecular Medicine. 2017;49(4):e324-e.

37.   Chen YK, Bonaldi T, Cuomo A, Del Rosario JR, Hosfield DJ, Kanouni T, et al. Design of
KDM4 Inhibitors with Antiproliferative Effects in Cancer Models. ACS Medicinal Chemistry Letters.
2017;8(8):869-74.

38.   Metzger E, Stepputtis SS, Strietz J, Preca B-T, Urban S, Willmann D, et al. KDM4 Inhibition
Targets Breast Cancer Stem–like Cells. Cancer Research. 2017;77(21):5900-12.

39.   Gogleva A, Polychronopoulos D, Pfeifer M, Poroshin V, Ughetto M, Martin MJ, et al.
Knowledge graph-based recommendation framework identifies drivers of resistance in EGFR
mutant non-small cell lung cancer. Nature Communications. 2022;13(1):1667.

40.   Chang CA, Jen J, Jiang S, Sayad A, Mer AS, Brown KR, et al. Ontogeny and Vulnerabilities
of Drug-Tolerant Persisters in HER2+ Breast Cancer. Cancer Discovery. 2022;12(4):1022-45.

41.   Rehman SK, Haynes J, Collignon E, Brown KR, Wang Y, Nixon AML, et al. Colorectal
Cancer Cells Enter a Diapause-like DTP State to Survive Chemotherapy. Cell. 2021;184(1):226-
42.e21.

42.   Liu Y, Azizian NG, Sullivan DK, Li Y. mTOR inhibition attenuates chemosensitivity through
the induction of chemotherapy resistant persisters. Nature Communications. 2022;13(1):7047.

43.   du Manoir S, Delpech H, Orsetti B, Jacot W, Pirot N, Noel J, et al. In high-grade ovarian
carcinoma, platinum-sensitive tumor recurrence and acquired-resistance derive from quiescent
residual cancer cells that overexpress CRYAB, CEACAM6, and SOX2. The Journal of Pathology.

2022;257(3):367-78.

44. Kimbung S, Markholm I, Bjöhle J, Lekberg T, von Wachenfeldt A, Azavedo E, et al.
Assessment of early response biomarkers in relation to long-term survival in patients with HER2-
negative breast cancer receiving neoadjuvant chemotherapy plus bevacizumab: Results from the
Phase II PROMIX trial. International Journal of Cancer. 2018;142(3):618-28.

45. van den Ende T, de Clercq NC, van Berge Henegouwen MI, Gisbertz SS, Geijsen ED,
Verhoeven RHA, et al. Neoadjuvant Chemoradiotherapy Combined with Atezolizumab for
Resectable Esophageal Adenocarcinoma: A Single-arm Phase II Feasibility Trial (PERFECT). Clinical
Cancer Research. 2021;27(12):3351-9.

46. Snipstad K, Fenton CG, Kjæve J, Cui G, Anderssen E, Paulssen RH. New specific molecular
targets for radio-chemotherapy of rectal cancer. Molecular Oncology. 2010;4(1):52-64.

47. Zhang S, Yu D. Targeting Src family kinases in anti-cancer therapies: turning promise into
triumph. Trends in Pharmacological Sciences. 2012;33(3):122-8.

48. Formisano L, Lu Y, Servetto A, Hanker AB, Jansen VM, Bauer JA, et al. Aberrant FGFR
signaling mediates resistance to CDK4/6 inhibitors in ER+ breast cancer. Nature Communications.
2019;10(1):1373.

49. Zhang L, Yang Y, Chai L, Bu H, Yang Y, Huang H, et al. FRK plays an oncogenic role in
non-small cell lung cancer by enhancing the stemness phenotype via induction of metabolic
reprogramming. International Journal of Cancer. 2020;146(1):208-22.

50. Girotti MR, Pedersen M, Sanchez-Laorden B, Viros A, Turajlic S, Niculescu-Duvaz D, et al.
Inhibiting EGF Receptor or SRC Family Kinase Signaling Overcomes BRAF Inhibitor Resistance in
Melanoma. Cancer Discovery. 2013;3(2):158-67.

51. Kohale IN, Yu J, Zhuang Y, Fan X, Reddy RJ, Sinnwell J, et al. Identification of Src Family
Kinases as Potential Therapeutic Targets for Chemotherapy-Resistant Triple Negative Breast Cancer.
Cancers [Internet]. 2022; 14(17).

52. Loh Y-H, Zhang W, Chen X, George J, Ng H-H. Jmjd1a and Jmjd2c histone H3 Lys 9
demethylases regulate self-renewal in embryonic stem cells. Genes & Development.
2007;21(20):2545-57.

53. Zhang Z, Lee JC, Lin L, Olivas V, Au V, LaFramboise T, et al. Activation of the AXL kinase
causes resistance to EGFR-targeted therapy in lung cancer. Nature Genetics. 2012;44(8):852-60.

54. Kim H-Y, Choi H-J, Lee J-Y, Kong G. Cancer Target Gene Screening: a web application for
breast cancer target gene screening using multi-omics data analysis. Briefings in Bioinformatics.
2020;21(2):663-75.

55. Sachs N, de Ligt J, Kopper O, Gogola E, Bounova G, Weeber F, et al. A Living Biobank of
Breast Cancer Organoids Captures Disease Heterogeneity. Cell. 2018;172(1):373-86.e10.

Figure 1. Pairwise CRISPR screen reveals combinations of synthetic lethal tyrosine kinase ablations. (A) Schematic diagram of combinatorial screens performed in TNBC cell line MDA-MB-231. (B) Scatter plot of expected growth phenotype Z score and observed growth phenotype Z score of each gene combination. Green dots indicate gene combinations where the identical gene is targeted by the two sgRNAs. Red dots indicate candidate synthetic lethal gene pairs listed in table S1. (C) Scatter plot of growth phenotype Z score and normalized GI score of each gene combination. (D) Scatter plot of gene level GI score and RIGER p value calculated with GI scores of each sgRNA pairs that target the given gene pair.

Figure 2. FYN is critical mediator of TKI resistance. (A) Schematic diagram of *in vitro* validation of

synthetic lethal gene pairs using sgRNAs. (B) Summary of synergistic killing by sgRNAs targeting indicated

gene pairs (n=3). (C) Network analysis of the 30 candidate synthetic lethal gene pairs highlighted in figure 1D.

The size of each node is manually drawn to be proportional to the number of connections the gene has. (D-E)

FYN and SRC mRNA expressions in (D) microarray data of primary breast cancers of GSE25066 cohort, and (E)

in cancer cell line encyclopedia, for indicated subtypes. (F) Summary of MTT assay with MDA-MB-231 cells

treated with the TKI combinations at indicated concentrations (n=2). Synergistic killing is calculated using

SynergyFinder with Bliss independence model. (G) Dose response curve of the indicated TKI in the presence

and absence of 5μM PP2 treated for 72 hours (n=3). (H) MTT assay with MDA-MB-231 Cas9 cells expressing

indicated sgRNAs treated with indicated TKIs for 72 hours (n=3). (I) Cell death and cell proliferation in MDA-

MB-231 cells treated with NVP-ADW742, gefitinib and PP2 either as single agent or as combination for 72

hours (n=3). (J) western blot analysis of MDA-MB-231 cells treated with indicated drugs for 48 hours (K)

western blot analysis of MDA-MB-231 Cas9 cells expressing indicated sgRNA and treated with indicated drugs

for 48 hours. (L) MTT assay of MDA-MB-231 cells treated with indicated drugs for 72 hours (SB203580: 10μ

M, NVP-ADW742: 4μM, gefitinib: 10μM, imatinib: 10μM) (n=3). PP2, Saracatinib, NVP-ADW742, gefitinib

and imatinib were treated at 5μM, 5μM, 4μM, 12μM, 12μM unless otherwise indicated. All data are plotted as

mean±s.d. One sample t-test for B, and unpaired two-sided Student's t-test in D,E,H and L. *, $p<0.05$; **,

$p<0.01$; ***, $p<0.001$; n.s., $p>0.05$. All replicates are biological replicates.

Figure 3. Activation of *KDM4* upregulates *FYN*, conferring drug resistance. (A) Western blot analysis of

MDA-MB-231 cells treated with indicated drugs. Numbers below blots indicate quantification of average±s.d.

expression level normalized to GAPDH in three independent experiments. (B) RT-qPCR analysis of *FYN*

expression levels in MDA-MB-231 cells treated with indicated drugs for 48 hours (n=3). (C) RT-qPCR analysis

of indicated jumonji family histone demethylase expression levels after 48 hours treatment of NVP-ADW742

(n=3). (D) Changes in *FYN* mRNA levels upon 48 hours of NVP-ADW742 treatment in MDA-MB-231 Cas9

cells expressing indicated sgRNAs (n=4). (E) *KDM4A* mRNA levels in primary tumor tissues of indicated

subtypes in GSE25066 cohort. (F) Positive correlation of *FYN* and *KDM4A* mRNA levels in CCLE database. (G)

western blot analysis of MDA-MB-231 Cas9 cells expressing indicated sgRNA and treated with indicated drugs

for 48 hours. (H) MTT assay of MDA-MB-231 Cas9 cells expressing indicated sgRNAs and treated with

indicated drugs for 72 hours (n=3). (I) SynergyFinder analysis of MDA-MB-231 cells treated with indicated

drug combinations (n=2). (J-K) western blot analysis (J) and RT-qPCR analysis (K) of MDA-MB-231 cells

treated with indicated drugs for 48 hours (n=3). (L) mRNA expression levels of SRC family kinases in breast

cancer stem cells treated with QC6352 in RNA sequencing data described in Metzger et. al.(38) (M) H3K9me3

and KDM4A enrichment at genomic locus encoding *FYN* promoter in ChIP sequencing data described in the

same study as (L). (N) H3K9me3 Chromatin immunoprecipitation-qPCR analysis of MDA-MB-231 cells

treated with indicated drug for 48 hours at specified genomic loci (n=3). QC6352, PP2, NVP-ADW742,

gefitinib and imatinib were treated at 10µM, 5µM, 4µM, 12µM, 12µM , respectively, unless otherwise indicated.

All data are plotted as mean±s.d. Unpaired two-sided Student's t-test in B,C,D,E,H and K. Paired two-sided

Student's t-test in N. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; n.s., $p>0.05$. All replicates are biological replicates.

Figure 4. Combination therapy targeting FYN+IGF1R and KDM4+EGFR synergistically eliminates

tumor *in vivo*. (A-B) Tumor volume for MDA-MB-231 xenografts treated with indicated drugs. Additive effects

were calculated by Bliss independence model (n=5). (C) Distant relapse free survival of GSE25066 patient

cohort classified by *FYN* (left) and *KDM4A* (right) mRNA expression. (D) Schematics diagram of the

mechanism of KDM4-FYN conferring TKI resistance. All data are plotted as mean±s.d. *, p<0.05; **, p<0.01;

***, p<0.001; n.s., p>0.05. All replicates are biological replicates.

Figure 5. *FYN* and *KDM4* are associated with drug tolerance. (A) MDA-MB-231 cells treated with indicated

drugs for short (acute: 2 days) and long (DTP: 10 days) time periods. A: 1μM NVP-ADW742, G: 5μM gefitinib,

I: 5μM imatinib, D: 100nM doxorubicin, P: 5nM paclitaxel. Numbers below blots indicate quantification of

average±s.d. expression level normalized to GAPDH in three independent experiments. (B) Summary of

mRNA expressions of indicated genes in EGFR mutant lung cancer cells (parental) and their derivative

osimertinib tolerant persisters (DTP). (C-D) western blots (C) and RT-qPCR (D) analyses of indicated parental

and EGFR inhibitor resistant lung cancer cells. (E-F) MTT assay with PC9 parental (par.) and gefitinib resistant

(GR) cells treated with indicated drug combinations (gefi: 2μM gefitinib, PP2: 5μM PP2, Sara: 5μM saracatinib,

QC6352: 10μM QC6352) for 72 hours. (G) western blot analysis with PC9 cells treated with 10μM QC6352 for

48 hours. (H) *FYN* mRNA expression levels of parental and DTP populations in various cancers treated with

indicated drugs.. (I) *FYN* mRNA expression levels of residual disease after indicated treatments. All data are

660 plotted as mean±s.d. Paired two-sided Student's t-test in B, H (HER2+ BRCA set and HGSOC carboplatin set),

661 and I (BRCA set and ESCA set), and unpaired two-sided Student's t-test in H (COAD and PAAD sets) and I

662 (COAD set). The expression data in B,H,I are obtained from NCBI gene expression omnibus. The accession

663 numbers of the expression data analyzed are listed in Supplementary table S2. *, $p < 0.05$; **, $p < 0.01$; ***,

664 $p < 0.001$; n.s., $p > 0.05$. All replicates are biological replicates.

665