Amoebozoan testate amoebae illuminate the diversity of heterotrophs and the

complexity of ecosystems throughout geological time

Alfredo L. Porfirio-Sousa[1,2]*, Alexander K. Tice[2,3]*, Luana Morais[4,5], Giulia M. Ribeiro[1],
Quentin Blandenier[2], Kenneth Dumack[6], Yana Eglit[7,8,9], Nicholas W. Fry[2], Maria Beatriz
Gomes E Souza[1], Tristan C. Henderson[2], Felicity Kleitz-Singleton[2], David Singer[10], Matthew
W. Brown[2,11]**, Daniel J. G. Lahr[1]**

[1]Department of Zoology, Institute of Biosciences, University of São Paulo, São Paulo, Brazil
[2]Department of Biological Sciences, Mississippi State University, Mississippi State,
Mississippi, USA
[3]Department of Biological Sciences, Texas Tech University, Lubbock 79409, Texas
[4]Department of Geophysics, Institute of Astronomy, Geophysics and Atmospheric Sciences,
University of São Paulo, São Paulo, Brazil
[5]Institute of Geosciences, São Paulo State University, Rio Claro, Brazil
[6]Terrestrial Ecology, Institute of Zoology, University of Cologne, Germany
[7]Department of Biology, Dalhousie University, Halifax, NS, Canada
[8]Institute for Comparative Genomics, Dalhousie University, Halifax, NS, Canada
[9]Department of Biology, University of Victoria, Victoria, BC, Canada
[10]Soil Science and Environment Group, Changins, HES-SO University of Applied Sciences
and Arts Western Switzerland, Nyon, Switzerland
[11]Institute for Genomics, Biocomputing & Biotechnology, Mississippi State University,
Mississippi State, Mississippi, USA
* Contributed equally

** Corresponding authors

Email: dlahr@ib.usp.br and matthew.brown@msstate.edu

Author Contributions: Conceptualization, Supervision, Project Administration,
Visualization, Methodology, Writing – Original Draft, A.L.P.-S, A.K.T., M.W.B., and D.J.G.L.;
Funding Acquisition and Resources, D.J.G.L. and M.W.B.; Sampling and Culture
Maintenance, A.L.P.-S, A.K.T., Q.B., K.D., Y.E., M.B.G.S, T.H., F.K.-S, M.W.B., and D.J.G.L;
Data Generation, A.L.P.-S, A.K.T.; L.M.; G.M.R., Q.B., K.D., N.F., T.H., D.S., M.W.B., and
D.J.G.L.; Phylogenetic Matrices Construction, A.L.P.-S, A.K.T., Q.B., N.F., T.H.; F.K.-S,
M.W.B., and D.J.G.L.; Formal Analysis and Investigation, A.L.P.-S, A.K.T.; L.M.; M.W.B.,
and D.J.G.L.; Review Editing, all authors.

Competing Interest Statement: The authors declare no competing interest.

Deposited as a preprint: bioRxiv, CC BY

Classification: Biological Sciences, Evolution

Keywords: Arcellinida; vase-shaped microfossils; phylogenomics; ancestral state
reconstruction; Arcellinida classification; eukaryotic evolution;

Abstract

Heterotrophic protists are vital in Earth's ecosystems, influencing carbon and nutrient cycles and occupying key positions in food webs as microbial predators. Fossils and molecular data suggest the emergence of predatory microeukaryotes and the transition to a eukaryote-rich marine environment by 800 million years ago (Ma). Neoproterozoic vase-shaped microfossils (VSMs) linked to Arcellinida testate amoebae represent the oldest evidence of heterotrophic microeukaryotes. This study explores the phylogenetic relationship and divergence times of modern Arcellinida and related taxa using a relaxed molecular clock approach. We estimate the origin of nodes leading to extant members of the Arcellinida Order to have happened during the latest Mesoproterozoic and Neoproterozoic (1054 - 661 Ma), while the divergence of extant infraorders postdates the Silurian. Our results demonstrate that at least one major heterotrophic eukaryote lineage originated during the Neoproterozoic. A putative radiation of eukaryotic groups (e.g. Arcellinida) during the early-Neoproterozoic sustained by favorable ecological and environmental conditions may have contributed to eukaryotic life endurance during the Cryogenian severe ice ages. Moreover, we infer that Arcellinida most likely already inhabited terrestrial habitats during the Neoproterozoic, coexisting with terrestrial Fungi and green algae, before land plant radiation. The most recent extant Arcellinida groups diverged during the Silurian Period, alongside other taxa within Fungi and flowering plants. These findings shed light on heterotrophic microeukaryotes' evolutionary history and ecological significance in Earth's ecosystems, using testate amoebae as a proxy.

Significance Statement

Arcellinida shelled amoebae are heterotrophic microbial eukaryotes with an extensive Neoproterozoic fossil record represented by the vase-shaped microfossils (VSMs), a diverse group that is abundant and widespread in late Tonian rocks (VSMs). Here we combined phylogenomic sampling and the fossil record to generate time-calibrated trees. Our results illuminate key events in the history of life, including: i) the Tonian origin of extant microbial eukaryote lineages; ii) a speculative proposed radiation of eukaryotes before the Cryogenian, "Tonian revolution"; iii) the establishment of complex terrestrial habitats before the Cryogenian; iv) a post-

Silurian divergence of modern Arcellinida sub-clades in terrestrial (including freshwater) habitats. Our results provide valuable insights into the evolution of life throughout geological time and are congruent with recent discoveries regarding the early diversification of eukaryotes, including the Precambrian history of eukaryotic protosteroids.

Introduction

Heterotrophic microbial eukaryotes play a crucial ecosystem role by contributing to the carbon and nutrient cycles (1, 2). These organisms, capable of phagocytosis, act as predators on bacterial and eukaryotic communities, playing a significant role in complex food webs supported by primary producers (1). Additionally, predation is an evolutionary innovation that likely contributed to the diversification of eukaryotes (3). The Last Eukaryotic Common Ancestor (LECA) was heterotrophic and capable of phagocytosis. However, the timing and specific conditions under which diverse lineages of heterotrophic microeukaryotes have proliferated in Earth's ecosystems remain unclear (4-6).

Evidence from fossils, biomarkers, geochemical proxies, genomic data, and molecular clocks indicate that eukaryotes first originated during the Stenian (1200-1000 Ma) and Tonian periods (1000-720 Ma) (7-11). This led to a transition from a prokaryotic- to a eukaryotic-rich marine environment (6, 12-14), by 800 Ma, likely triggered by increased phosphorus, nitrate, and silica availability (14-17). From around this time, Neoproterozoic vase-shaped microfossils (VSMs) represent the remnants of an early eukaryotic divergence event. Organisms represented by VSMs are generally thought to have lived in marine environments, although a terrestrial habitat for these organisms is also plausible (18-20). The well-preserved nature of VSMs has allowed for detailed investigation and comparisons of their morphology to modern eukaryotic groups. These investigations support the current interpretation of a large fraction of VSMs being members of the stem or crown groups of testate amoeba order Arcellinida, due to both morphological affinities and congruence with molecular phylogenetic reconstructions (19, 21, 22). Other VSMs, such as *Melicerion poikilon*, had suggested affinities to Euglyphida, a distantly-related, convergent rhizarian clade of testate amoebae (18, 19). However, morphological evidence, in this case, is tentative, and the suggestion for a Euglyphida affinity is currently incongruent with molecular phylogenetic reconstructions, as Euglyphida is

part of a clade of cercozoan filose amoebae, which appears to be much younger, around 292 Ma (23). Arcellinida is a diverse lineage of extant heterotrophic microeukaryotes within the Amoebozoa, found in terrestrial and freshwater environments (19-22). Since many VSMs have been recognized as Arcellinida, they are accepted to represent the oldest and most diverse fossil evidence of heterotrophic microeukaryotes (19-20, 24,25). Elucidating the origin and evolutionary history of Arcellinida (and derived lineages) in light of their microfossil record is pivotal to illuminate the early evolution and possible radiation of heterotrophic microbial eukaryotes, and serve as a proxy to infer the complexity of Earth's ecosystems over geological time (19-20, 25, 26).

Recent efforts of sampling diverse amoebozoan testate amoebae in a phylogenomic framework have resolved their deep phylogenetic relationships (21, 27). Amoebozoa is home to at least two testate amoebae groups: Arcellinida and Corycidia. Arcellinida is a diverse order represented by lineages that build hard extracellular shells, with the potential to generate exceptionally preserved fossilized remains (19, 21). Corycidia is a recently established subclade of Amoebozoa represented by the lineages of testate amoebae that produce flexible shells and do not branch within Arcellinida (27). Despite recent advances, many lineages still remain unsampled (21).

In addition to expanding the diversity of sampled Arcellinida at the genome level, the potential of a highly resolved phylogeny to provide insight into timing the Arcellinida origin and divergence of sub-clades has not been explored (22, 25). VSMs can be interpreted either as stem or crown Arcellinida. In either case, these fossils can be used to calibrate a phylogenetic tree and estimate the divergence time of lineages both within Arcellinida, as well as closely related amoebozoans. These diverse VSMs found in sedimentary deposits around the world have been continuously investigated, and their stratigraphy refined over time, enabling us to constrain these fossils' ages (19-20, 28-36). In this context, combining the VSMs and phylogenetic tree calibration opens up avenues to time the evolution of Arcellinida and closely related groups.

Here, we investigate the origin and divergence times of Arcellinida and closely related amoebozoan taxa using phylogenomics and a relaxed molecular clock approach. We expanded the taxonomic sampling for amoebozoan testate amoebae, including 14 taxa that lacked precise placement, to produce a new phylogenomic

dataset (utilizing 226 genes). We considered the diverse record of VSMs and
Metazoa fossils to time calibrate this well-resolved deep phylogenomic tree. Different
calibration strategies and molecular clock models support the divergence of extant
Arcellinida lineages during the latest Mesoproterozoic and early to mid-
Neoproterozoic, between 1060 and 661 Ma. We thus corroborate the origin of a
major eukaryotic group by the Neoproterozoic, including a recorded establishment of
heterotrophy predating the Cryogenian Period. Overall, using amoebozoan testate
amoebae as a proxy, we provide insights into the evolution of microbial eukaryotes
and Earth's early ecosystems.
149
150　**Results**
151
A resolved tree of amoebozoan testate amoebae
We constructed a concatenated supermatrix using 57 taxa and 226 genes (70,428
amino acid sites) using the PhyloFisher v. 1.2.11 package (37, 38). We include novel
data for 14 testate amoebae taxa, obtained through single-cell or whole-culture
transcriptomics (**Fig. 1; Dataset S01, Tables S1 - S4**). The remaining testate
amoebae and sister-group taxa were sampled based on previously available
genomes and transcriptomes (**Dataset S01, Table S1**). The resulting phylogenomic
tree recovers a monophyletic Arcellinida, with three well-defined suborders
(Phryganellina, Organoconcha, and Glutinoconcha) and five infraorders within
Glutinoconcha (**Fig. 2; Appendix S01, Figs. S1 and S2**). The Corycidia clade is
also recovered with full support, with two families, Trichosidae and Amphizonelliidae
fam. nov. (**Fig. 2**). Nearly all nodes of the tree are fully (=100%) or highly (>92%)
supported by Maximum Likelihood nonparametric Real Bootstraps (MLRB), except
for a single node within Sphaerothecina clade that has lower support (MLRB = 76%;
Fig. 2). We have also produced single-gene reconstructions using SSU rDNA and
Cytochrome Oxidase subunit I (COI) **(Dataset S01, Tables S5-S6),** which are the
genes traditionally used to reconstruct relationships in Arcellinida. These analyses
present broader taxon sampling but failed to recover most of the deeper
relationships in Arcellinida **(Appendix S01, Figs. S3-S4**)
171
Time-calibrated tree of amoebozoan testate amoebae

For estimating divergence times in testate amoebae evolution and closely related
taxa, we expanded our phylogenomic supermatrix to consider a representative
sampling of Amorphea, including Amoebozoa, Fungi, Metazoa, and their protistan
relatives **(Dataset S01, Table S7)**. For a comprehensive approach, taking into
account the alternative interpretations of the VSM record, we implemented four
different calibration strategies: i. calibration of nodes within Metazoa, excluding the
VSM record to calibrate amoebozoan nodes; ii. calibration of nodes within Metazoa
and calibration of Glutinoconcha+Organoconcha and Glutinoconcha nodes,
considering VSMs as derived crown Arcellinida; iii. calibration of nodes within
Metazoa and calibration of the Arcellinida node, considering VSMs as basal crown
Arcellinida; iv. calibration of nodes within Metazoa and calibration of
Arcellinida+Euamoebida node, considering VSMs as stem Arcellinida (**Dataset S01,**
Table S8 - S10). To implement these four fossil calibration strategies, we performed
a total of 36 experiments considering three different distributions (i.e., Uniform,
Skew-Normal, or Truncated-Cauchy short-tail) under an uncorrelated or
autocorrelated relaxed clock model with either a drift parameter of $\alpha = 2$ and $\beta = 2$ or
$\alpha = 1$ and $\beta = 10$. We ran each experiment in two independent MCMC chains to
check for convergence, which was achieved for all analyses (**Appendix S01, Fig.**
S5).

Comparing all the calibration strategies and experiments, we observed overall
similar inferred times with the uncorrelated clock model (median = 591 - 531 Ma and
mean = 627 - 549 Ma; **Dataset S01, Table S11 and Appendix S01, Figs. S6 - S43**)
and the autocorrelated clock model (median = 592 - 531 Ma and mean = 671 - 557
Ma; **Dataset S01, Table S11**). Regarding implemented distributions, overall the
uniform distribution inferred the youngest ages, while Truncated-Cauchy inferred
slightly older ages (**Dataset S01, Table S11**). The drift parameter (i.e., $\alpha = 2$ and $\beta =$
2 vs. $\alpha = 1$ and $\beta = 10$) had virtually no impact independent of the clock model,
distribution, and calibration strategy (**Dataset S01, Table S11**). The estimated times
for the Arcellinida node by excluding VSMs from the calibration (mean = 911 - 734
Ma; 95% highest probability density confidence interval (HPD CI) = 1060 - 605) and
by including VSMs in the calibration (mean = 930 - 746 Ma; 95% HPD CI = 1054 -
661 Ma) are highly congruent. This agrees with the current interpretation that VSMs
represent fossil remains of Arcellinida, supporting their use to calibrate amoebozoan
nodes. Aiming for a comprehensive approach, the time estimation results shown and

discussed hereafter focus on the full range of times estimated based on the calibration strategies that included the VSMs, the three distributions, and the drift parameter of α = 2 and β = 2 under the uncorrelated or autocorrelated relaxed clock model.

The molecular clock analyses inferred a mean time for the root of Amorphea to be between 1640 and 1393 Ma (95% HPD CI = 1843 - 1088 Ma; **Fig. 3, Dataset S01, Table S11, and Appendix S01, Fig. S7 - S43**). For Metazoa, the mean time estimated ranged between 835 and 734 Ma (95% HPD CI = 872 - 673 Ma; **Fig. 3, Dataset S01, Table S11, and Appendix S01, Fig. S7 - S43**). The mean estimated for the origin of Amoebozoa ranged from 1607 - 1298 Ma (95% HPD CI = 1795 - 1045 Ma; **Fig. 3, Dataset S01, Table S11, and Appendix S01, Fig. S7 - S43**). The Arcellinida node is constrained within the mean 930 and 746 Ma (95% HPD CI = 1054 - 661 Ma; **Fig. 3, Dataset S01, Table S11, and Appendix S01, Fig. S7 - S43**), estimating an origin for Arcellinida during the latest Mesoproterozoic and Neoproterozoic (**Fig. 4A**). For the early divergence time of Arcellinida subclades, the estimated times suggest that the split between the Organoconcha and Glutinoconcha branches occurred during the Neoproterozoic (mean = 855 - 679 Ma; 95% HPD CI = 969 - 600 Ma; **Figs. 3 - 4B**). Regarding the suborders of Arcellinida, we inferred a mean origin for Phryganellida between 534 - 265 Ma (95% HPD CI = 661 - 175 Ma; **Figs. 3 - 4C**), for Organoconcha 735 - 550 Ma (95% HPD CI = 839 - 463 Ma; **Figs. 3 - 4D**), and for Glutinoconcha between 790 - 621 Ma (95% HPD CI = 897 - 539 Ma; **Figs. 3 - 4E**). For the deeper nodes of Glutinoconcha, the most sampled suborder of Arcellinida, the estimated divergence times ranged between Cryogenian and Carboniferous (mean = 643 - 393 Ma; 95% HPD CI = 705 - 335; **Figs. 3 - 4F - H**). The inferred ages for Glutinoconcha infraorders are relatively more widespread depending on the calibration strategy when compared to other nodes. Treating VSMs as derived crown taxa constrains the infraorders origin between Ediacaran and early Cretaceous (mean = 422 - 221 Ma; 95% HPD CI = 575 - 123 Ma; **Figs. 3 - 4I - M**) while considering VSMs as basal crown or stem Arcellinida estimate their origin mostly between Silurian and early Cretaceous (mean = 339 - 172 Ma; 95% HPD CI = 444 - 122 Ma; **Figs. 3 - 4I - M**). It is worth noting that only the calibration using a uniform autocorrelated clock model and VSMs as derived crown arcellinids inferred ages as old as the Ediacaran for the Glutinoconcha infraorders. All other distribution-clock models consistently led to ages constrained

within the Paleozoic. Most inferred ages for the nodes representing the origin of the modern extant genera and species of Arcellinida postdate the Silurian (mean = 416 - 125 Ma; **Dataset S01, Table S11**). Besides the testate amoebae, the inferred mean times for the other orders and major groups of Amoebozoa we sampled ranged from 585 Ma to 1288 Ma, placing their origin mostly during the Neoproterozoic (**Dataset S01, Table S11**). The results and time-calibrated trees for all experiments are present in the supplemental material (**Dataset S01, Table S11 and Appendix S01, Fig. S6 - S43**).

Ancestral habitat reconstruction of Arcellinida

We performed statistical analyses on the ancestral habitat of key hypothetical ancestors within Arcellinida considering alternative scenarios of a terrestrial or marine origin for the crown group (**Dataset S01, Table S12 and Appendix S01, Fig. S44**). The unrestrained reconstruction inferred a terrestrial habitat (100% probability) for all nodes within Arcellinida. The ancestral reconstruction that sets the fixed value of a marine state on the last common ancestor of modern Arcellinida inferred a high probability of terrestrial habitat for all nodes within Arcellinida (>93%), implying at least two independent transition events (2TE) from marine to terrestrial habitats (**Fig. 5**). The ancestral reconstruction that sets the fixed value of a marine state on the last common ancestor of both the Arcellinida and the Organoconcha+Glutinoconcha clades inferred with high probability (>88%) a terrestrial habitat for the hypothetical ancestors of Phryganellina, Organoconcha, and Glutinoconcha, implying at least three independent transition events (3TE) from marine to terrestrial habitats (**Fig. 5**).

Discussion

A resolved tree of amoebozoan testate amoebae

The phylogenomic dataset constructed in this study improves several aspects of the previously available amoebozoan testate amoebae dataset (21, 27). Through the PhyloFisher pipeline, we were able to construct a curated phylogenomic matrix that is free of paralogs and contamination (**Appendix S01, SI1**). Moreover, the matrix constructed is an accessible and easy-to-update dataset, since newly sequenced transcriptomes can be easily added through PhyloFisher to further expand the taxonomic sampling of amoebozoan testate amoebae in a phylogenomic approach.

In general terms, the phylogenomic tree obtained here is consistent with the
previously published phylogenomic tree for amoebozoan testate amoebae (**Fig. 2**;
21, 27). By recovering all major groups of the Arcellinida and Corycidia clades with
full support, in accordance with previous reconstructions, we corroborate the
robustness of the backbone of the amoebozoan testate amoebae tree. The
phylogenomic analyses enable us to place previously unsampled taxa within
Arcellinida suborders, supporting the taxonomic actions regarding *Heleopera lucida*
comb. nov. and the genus *Microcorycia* (**Appendix S01, SI2 - SI3**). Moreover, the
phylogenomic tree identifies taxa that will need future taxonomic revision based on
further morphological and molecular studies, those being *Difflugia* cf. *capreolata* and
the genera *Cyclopyxis* and *Phryganella*. Detailed discussion on the placement of
newly sequenced amoebozoan testate amoebae is presented in **Appendix S01, SI2
- SI3.**
288
Robust Amorphea time-calibrated trees using VSMs
Our time-calibrated trees are congruent with several molecular clocks that have
sampled the diversity of eukaryotes (**Fig. 3, Dataset S01, Table S11, and Appendix
S01, Fig. S6 - S43**). The estimated ages for the root of our tree fall within the range
inferred by a recent molecular timescale for eukaryotes (95% confidence interval =
2177 - 1356 Ma for Amorphea; 39). Similarly, the estimated ages for the origin of
other major clades align with previous studies, including Obazoa (95% HPD CI =
2305 - 1526 Ma), opisthokonts (95% HPD CI = 2019 - 1051 Ma), and animals (95%
HPD CI = 833 - 680 Ma) (39-42). The inferred times place the origin of Arcellinida
during the latest Mesoproterozoic and early to mid-Neoproterozoic, most likely during
the Tonian Period and no later than the Cryogenian (**Fig. 4, Dataset S01, Table
S11, and Appendix S01, Fig. S6 - S43**). It is worth noting that by considering
different calibration strategies to account for alternative plausible interpretations of
VSMs and the unavoidable fossil record uncertainties, the times we estimated for
each node have wide ranges. However, independently of the calibration strategy,
distribution, and clock model, the Arcellinida origin is mostly placed within the
Neoproterozoic, and all estimated mean times suggest a Tonian origin. Notably, the
time-calibrated trees we generated without using the VSMs as calibration data
inferred times congruent with our analyses considering the Tonian VSMs. This
demonstrates that inferred ages are not a fossil calibration bias and serves as further

corroboration for the interpretation of VSMs as fossil members of, at the very least,

the stem Arcellinida group (**Fig. 4, Dataset S01, Table S11**). Also, the ages we

inferred for the origin of modern groups (i.e., genera) are highly congruent with the

recent Arcellinida fossil record which postdates the Carboniferous Period and

preserves fossils assigned to genera like *Arcella*, *Difflugia*, and *Centropyxis* (**Dataset**

S01, Table S11; 43-45). Collectively, the consistency of the estimated times, and

their congruence with previous molecular clocks and with the fossil record, support

that including VSMs to calibrate a comprehensive phylogenomic sampling of

Amorphea leads to robust results.

The Neoproterozoic diversification of heterotrophic microbial eukaryotes

Our time-calibrated trees reveal that Arcellinida originated most likely during the

Neoproterozoic, with most inferred times and all of the means falling within the

Tonian Period (**Fig. 4 A - B**). The mean estimated times for other major groups of

Amoebozoa also indicate a Tonian origin. Similarly, previous molecular clock

analyses have indicated the divergence of multiple heterotrophic eukaryotes during

this period (40, 41). However, their diversity has been challenging to examine due to

the lack of a fossil record for these organisms. Nevertheless, the estimated

Neoproterozoic origin for the Arcellinida crown group and the diversity of Tonian

VSMs, currently represented by 14 morphologically diverse genera (19-20, 25, 31),

suggest that by the Tonian Period, Earth's ecosystems had witnessed the origin of

modern heterotrophic microbial eukaryotes.

An inferred Tonian origin for diverse heterotrophic microbial eukaryotes is

congruent with proposals of ecosystem establishment during the Neoproterozoic,

stemming from various disciplines. Previous studies speculate the existence of a

"Tonian revolution", based on evidence from biomarkers, fossil records, and

molecular data that infer a marked transition from prokaryotic- to eukaryotic-rich

ecosystems by 800 Ma (6, 11-14, 46). This transition was likely facilitated by factors

such as increased availability of nitrate, phosphorus, silica, and reduced toxicity,

which provided a favorable condition linked to the documented diversification of

eukaryotic phototrophs at that time (14-17). In turn, the establishment of a

community of phototrophs may have served as a favorable condition for the

diversification of heterotrophs. Fossil and geochemical evidence suggest that

photosynthetic biological mats contributed to the establishment of Oxygen Oases

during the Tonian, likely triggered by a higher capacity of oxygen productivity of
eukaryotic phototrophs (7, 10, 47, 48). These oases probably represented an
increase in aerobic conditions, and food availability, that were permissive to the
survival and proliferation of heterotrophs like Arcellinida. Consequently, a combined
interpretation of the geochemical and fossil records indicates that complex
ecosystems were established by the Neoproterozoic.

VSMs serve as a unique testimony of the putative "Tonian revolution" on
eukaryotic diversification and ecosystems established during the Neoproterozoic.
The VSMs have been found in rocks characterized by organic-rich sediments
corroborating the close association of heterotrophic eukaryotes with microbial mats
(19). Also, the organisms represented by these VSMs likely preyed on both the
bacterial and eukaryotic communities, similar to extant Arcellinida (19, 49, 50).
Culture observations have demonstrated diverse strategies of extant Arcellinida to
prey on various organisms, including diatoms, fungi, and nematodes (51-54).
Moreover, several VSMs exhibit holes on their shells interpreted as predation marks,
suggesting they also served as prey to other heterotrophs (3, 55). Predation has
been interpreted as one of the triggers for eukaryotic diversification, including for
animals, and VSMs are among the oldest records of this evolutionary innovation (3,
56-58). Thus, while most of the microbial eukaryotes did not leave fossils, VSMs
stand out as a robust fossil record highlighting the rise of predation and increase of
food web complexity on Earth's ecosystems no later than the Tonian Period.

Our time-calibrated trees suggest that the divergence of some modern
Arcellinida lineages happened during the Neoproterozoic. The inferred times for the
split and origin of the Organoconcha and Glutinoconcha suborders mostly fall within
the Neoproterozoic between the Tonian and Ediacaran Periods, with only some
estimated times suggesting an early-Paleozoic origin for Organoconcha. Specifically,
the ages predicted for the Organoconcha and Glutinoconcha split either predate or
overlap with the Cryogenian Period and its glaciations. It is worth noting that
Phryganellina is currently the least genomically sampled Arcellinida group,
represented only by two genera, thus it is difficult to assess when this suborder
originated. Altogether, the origin and early divergence of the Arcellinida crown group,
and the diverse VSMs record, imply the capability of Neoproterozoic ecosystems not
only to sustain heterotrophic eukaryote groups but also to allow their diversification.

A possible Pre-Cryogenian Eukaryotic Diversification and Survival during Earth's most severe Ice Ages

The Sturtian and Marinoan glaciation periods witnessed extensive ice coverage across the planet, with glaciers extending into tropical regions (59-61). The survival of life during Cryogenian glaciations can be explained by the presence of refugia and the adoption of dormancy strategies (62-69). The presence of cyst-like structures identified inside VSMs serves as direct evidence that these organisms were already capable of entering dormancy stages, similar to extant Arcellinida (70). Additionally, recent discoveries have revealed that Tonian VSM taxa persisted into the Cryogenian Period, providing fossil records that showcase the diversity of heterotrophic eukaryotes during glaciation periods (69). The inferred times for the split and origin of Glutinoconcha and Organoconcha suggest they possibly originated during Cryogenian, indicating not only survival but also divergence of novel modern eukaryotic taxa during Cryogenian glaciations (**Fig. 4 D - E**). Consequently, the VSM record and the timing of early Arcellinida evolution enable speculation about a possible radiation of heterotrophic eukaryotic life shortly before and during the Cryogenian Period. This, coupled with a capacity for dormancy and the exploitation of habitat refugia, may explain the endurance of life during Earth's most severe ice age.

Timing of terrestrial conquest by Arcellinida

Currently, while it is largely suggested that the organisms represented by Tonian VSMs lived in shallow marine environments, a terrestrial habitat cannot be ruled out. To date, VSMs have been reported from Tonian sedimentary deposits described as fully or partially marine (18-19, 25). However, although scarcely discussed in the literature, it is plausible to hypothesize that the organisms represented by the VSMs may have lived in terrestrial environments and were deposited in marine sediments through a number of possible mechanisms, including: surface runoff, river discharge, wind blowing, or supratidal spillover. In any case, these organisms ultimately fossilized in a marine setting (71). Consequently, considering the alternative interpretations of VSMs' natural habitat and their affinity to Arcellinida (i.e., stem or crown Arcellinida), different scenarios can be explored regarding Arcellinida's conquest of terrestrial habitat.

Our ancestral habitat reconstructions indicate three alternative scenarios, a
terrestrial origin for Arcellinida, a marine origin with two independent transition
events (2TE) from marine to terrestrial habitats, and a marine origin with three
independent transition events (3TE) (**Fig. 5 and Appendix S01, Fig. S44**). Although
the reconstruction of a terrestrial origin is statistically superior to the other scenarios
(Likelihood Ratio Test - LRT), this was already expected since all extant Arcellinida
are terrestrial/freshwater inhabiting, leading to the reconstruction of a terrestrial
ancestor (i.e. a possible systemic bias). However, interpreting VSMs as stem or
crown Arcellinida and enforcing a marine origin for the Arcellinida stem and early
crown groups, 2TE or 3TE are plausible and statistically equivalent based on LRT, in
accordance with previous hypotheses (18-21, 25, 72). Multiple transition events are
biologically plausible: Arcellinida-related amoebozoan lineages are often represented
by both marine and terrestrial species, even within the same genera (e.g., *Vannella*,
Mayorella, and *Trichamoeba*).

Coupling the reconstructed scenarios with the timing of Arcellinida origin and
early divergence of its sub-clades, we infer that many Arcellinida probably inhabited
terrestrial environments already in the Neoproterozoic, no later than the Ediacaran
Period. Even if we consider the latest transition scenario reconstructed (3TE) the
inferred times place the terrestrialization event of Glutinoconcha and Organoconcha
most likely between the Tonian and Ediacaran periods (**Fig. 5**). The time of
diversification of life in terrestrial habitats has been traditionally discussed based on
the time of divergence of land plants (embryophytes), which is constrained within a
Paleozoic diversification (73, 74). However, recent inferences based on
phylogenomic reconstructions and molecular clocks have suggested that modern
eukaryotic lineages, like Fungi and green algae, diverged on land no later than
Cryogenian (73, 74). Congruently, our estimated times for Arcellinida
terrestrialization are constrained within the Neoproterozoic. These findings suggest a
Neoproterozoic establishment of relatively complex terrestrial ecosystems inhabited
by diverse organisms, including phototrophic (green algae), absorptive heterotrophic
(Fungi), and phagotrophic heterotrophic protists.

The inferred divergence of Arcellinida sub-clades in terrestrial habitats, well
represented by Glutinoconcha (currently the best-sampled suborder), is also
congruent to the diversification timing estimated for other eukaryotic groups. The
Glutinoconcha infraorders' split is mostly constrained between the late-

Neoproterozoic and mid-Paleozoic (Devonian Period). The radiation of Fungi and the
diversification of extant land plants are estimated to the same window (ca. 480 Ma)
(75). Subsequently, the origin of extant Arcellinida groups, represented by the origin
of all Glutinoconcha infraorders, is mostly constrained between the Silurian and
Cretaceous. This is contemporaneous with the documented Late Paleozoic
diversification of seed plants and saprotrophic mushrooms (75). Similarly, the
estimated time for the divergence of Arcellinida genera, mostly post-dating early
Mesozoic, is congruent to the radiation of diverse groups of Fungi and land plants,
including pine trees (Pinaceae) and flowering plants (Angiosperm) (75). Altogether,
congruences between the timing of the origin of various eukaryotic groups suggest
an integrated and synchronous diversification of life in terrestrial habitats, enabling
speculation about a possible radiation of Arcellinida in this time period. However, this
claim requires explicit testing and corroboration via well-sampled studies of
background diversification rates using fossils.
458

Conclusions
Timing the origin of modern Arcellinida testate amoebae and the divergence times of
their subclades illuminate the evolution of heterotrophic microbial eukaryotes in
geological time. To estimate this timing, we expanded the phylogenomic sampling of
amoebozoan testate amoebae and generated robust time-calibrated Amorphea trees
based on both Arcellinida and Metazoa fossil records. The estimated times for the
origin of Arcellinida and other amoebozoans, mostly constrained within the Tonian
Period, are congruent with the previously speculated "Tonian revolution" for a
diversification of eukaryotes in this Period. This consistency suggests that Earth's
ecosystems had witnessed the divergence of both phototrophic and heterotrophic
eukaryotic lineages, including Arcellinida, during the Neoproterozoic, no later than
the Tonian Period. A putative radiation of eukaryotes before the Cryogenian Period,
coupled with the exploitation of refugia habitats and dormancy strategy, may have
contributed to their endurance during Earth's most severe ice ages. Although the
ancestral habitat of Arcellinida and the possibility of transition between environments
(marine vs. terrestrial) remain contentious, considering the plausible alternative
scenarios we infer that Arcellinida were most likely already inhabiting terrestrial
habitats between the Tonian and Ediacaran Periods. Together with the previously
suggested diversification of Fungi and green algae on land during the Cryogenian

Period, the inferred time for terrestrial Arcellinida is congruent with a Neoproterozoic
establishment of relatively complex ecosystems composed of phototrophic (green
algae), absorptive heterotrophic (Fungi), and phagotrophic heterotrophic eukaryotes,
preceding the diversification of land plants. Similarly, the estimated post-Silurian
origin of modern Arcellinida (i.e., infraorders) suggests a contemporaneity to the
diversification of other groups, including diverse Fungi and land plants. Ultimately,
we suggest the Arcellinida testate amoebae are a key group to further explore the
diversification of heterotrophic microbial eukaryotes and the establishment of
ecosystems starting in the Neoproterozoic.

Material and Methods

Sampling, RNA extraction, and sequencing
We generated transcriptomes for 14 previously genomically unsampled amoebozoan
testate amoeba species through mRNA extraction from either monoclonal cultures or
single-cells isolated from natural samples (**Dataset S01, Table S1**). We synthesized
cDNA from RNA extractions using an adaptation of the Smart-seq2 protocol (77). We
prepared our cDNA libraries for sequencing on the Illumina platform using a
NexteraXT DNA Library Preparation Kit (Illumina) following the manufacturer's
recommended protocol. Libraries were then pooled and sequenced (**Appendix S01,**
SI1; Dataset S01, Table S1).

Trimming, Transcriptome Assembly, and Quality Assessment
We trimmed primers, adaptors, and low-quality bases from raw Illumina reads using
Trimmomatic v. 0.36 (77). We then assembled the surviving reads with Trinity v.
2.12.0 (78). We predicted amino acid sequences (proteomes) from the assembled
transcriptomes using Transdecoder v. 5.5.0. Finally, we assessed the completeness
of all newly sequenced transcriptomes using BUSCO v. 5.3.2 (79) (**Dataset S01,**
Table S1). Further details on trimming, transcriptome assembly, and quality
assessment are presented in **Appendix S01, SI1.**

Phylogenomic dataset construction
We constructed our amoebozoan phylogenomic dataset using the database and
tools available in PhyloFisher v. 1.2.11 (37) following the steps outlined at

https://thebrownlab.github.io/phylofisher-pages/detailed-example-workflow and in

Jones et al. (38). Our final concatenated matrix used in subsequent phylogenetic

analyses consisted of 226 genes (70,428 amino acid sites) and 57 amoebozoan taxa

(**Dataset S01, Table S4**). From each individual ortholog that was concatenated to

create the aforementioned matrix, we constructed single ortholog trees to be used as

input for coalescent-based phylogenomic analyses. Further details on our approach

for phylogenomic dataset construction are presented in **Appendix S01, SI1.**

Phylogenomic analyses

We performed maximum likelihood phylogenetic reconstruction using our final matrix

with IQ-TREE2 v. 2.0-rc (80). We initially inferred a tree from our matrix under the

LG+C20+G4 model. We used the resulting tree as a guide tree to infer a Posterior

Means Site Frequency model (81) using the ML model LG+C60+G4+PMSF in IQ-

TREE2. We assessed the topological support for the resulting tree by 100 Real

nonparametric Bootstrap replicates in IQ-TREE (IQ-TREE v. 2.1.2 COVID-edition)

using the PMSF model. The topological support values inferred from MLRB were

mapped onto the ML tree using RAxML v. 8.2.12 (82) using the "-f b" option. We

carried out coalescent-based phylogenomic analyses with ASTRAL-III v. 5.7.3 (83).

Statistical support for our ASTRAL-III analysis was assessed via local posterior

probability values.

Bayesian molecular dating

Dataset and Topology

Utilizing previously identified orthologs already present in the publicly available

PhyloFisher database, we expanded our amoebozoan phylogenomic dataset to

include a representative sampling of Amorphea. Amorphea is the eukaryotic clade

composed of Amoebozoa, Fungi, Animals, and some other unicellular lineages.

These ortholog sequences were aligned, trimmed, and concatenated as described

above (*Phylogenomic dataset construction section*). Our final expanded dataset

used in the subsequent phylogenetic reconstruction and for the molecular dating

analysis consisted of 230 genes (73,467 amino acid sites) and 96 taxa (**Dataset**

S01, Table S7). Maximum likelihood phylogenetic analysis was performed as

described above (*Phylogenomic analyses section*).

Fossil calibrations

As external calibration information, we considered fossils to calibrate five internal nodes of the Metazoa clade and explored three different strategies to calibrate amoebozoan nodes (**Dataset S01, Tables S8 - S10**). We strictly derived the fossil calibration for Metazoa from dos Reis et al. (42) and Benton et al., (84), which have carefully evaluated the diversity of fossils available and calibration strategies for the Metazoa lineage. We derived the fossil calibration for Arcellinida based on previous analyses and interpretations of the morphological relationship between VSMs and extant Arcellinida (19, 21, 22). Currently, VSM's can be interpreted either as: i) the fossil record of stem Arcellinida; ii) basal crown Arcellinida closely related to Arcellinida common ancestor, or; iii) derived crown Arcellinida (19, 21, 22). Specifically, interpretations of Bayesian and maximum likelihood ancestral reconstructions of Arcellinida shell morphology suggest a morphological congruence between the VSM *Melanocyrillium* to the Glutinoconcha+Organoconcha hypothetical ancestor and between the VSM *Cycliocyrillum* to Glutinoconcha hypothetical ancestor, thus suggesting they may represent derived crown Arcellinida (21). Consequently, three different calibration strategies can be implemented from these alternative interpretations. VSM's can be considered to calibrate: i) Glutinoconcha+Organoconcha and Glutinoconcha nodes; ii) calibrate the Arcellinida node, or; iii) calibrate the node shared between Arcellinida and its closest sister group, the amoebozoan order Euamoebida. Aiming for a comprehensive approach we considered these three strategies and generated comparable time tree estimations. To constrain the VSMs ages, we considered the literature which has described these microfossils and refined their stratigraphical distribution (**Dataset S01, Tables S9;** 36). Since molecular dating considers the fossil information as statistical distributions, and different distributions may impact the time estimation differently, we followed dos Reis et al. (42) strategy and used a total of three different distributions to represent the calibrations derived from the fossil record: i. uniform; ii. skew-normal; and iii. truncated-cauchy short-tail. Full details on our approach for fossil calibration are presented in **Appendix S01, SI1.**

Divergence time estimation

We performed Bayesian molecular dating with the MCMCTree program, implemented within the PAML package (**Dataset S01, Table S10**; 85). We estimated

a mean substitution rate of 0.03135 replacement site^{-1}10^{-8}Myr^{-1} for our dataset with

IQ-TREE2 v. 2.0.6 (80) phylogenetic dating under the LG+G model. Within the PAML

package, we set the overall substitution rate ('rgene_gamma = α, β' parameter) as a

gamma-Dirichlet prior following dos Reis et al. (42), with α = 2 and β = 63.78. This

substitution rate was implemented for all dating experiments. For the rate drift

parameter ('sigma^2_gamma = α, β'), we independently implemented two alternatives,

α = 2 and β = 2 or α = 1 and β = 10. Similarly, we considered both uncorrelated and

autocorrelated relaxed clock models. For all experiments, we analyzed the data

under the LG+G model as a single partition, constrained the root age between 1.6 -

3.2 Ga, and considered a uniform birth-death tree prior and 100 million years as one

time unit. We performed a total of 36 experiments, considering three distributions

(i.e., uniform, skew-normal, and truncated-Cauchy short-tail), varying the rate drift

parameter (i.e., α = 2 and β = 2 or α = 1 and β = 10) and clock model (i.e.,

uncorrelated and autocorrelated). For each experiment, we ran two independent

MCMC chains to verify convergence, discarding the first 2,000 iterations as burn-in

and considering the following 20 million generations. To check the influence of fossil

calibrations using Neoproterozoic VSMs on the estimated dates, we performed

experiments calibrating only the nodes within the Animal clade, applying Uniform and

Skew-Normal calibration strategies, under an uncorrelated or autocorrelated relaxed

clock model with a drift parameter of α = 2 and β = 2 or α = 1 and β = 10, following

the same approach described above, as detailed in **Appendix S01, SI1.**

Ancestral reconstruction of Arcellinida habitat

We applied a maximum likelihood (ML) method, implemented in BayesTraits v. 4.0.1

(86), to statistically reconstruct the ancestral habitat states of Arcellinida and

compare evolutionary scenarios. Currently, the diverse Tonian VSM record has been

documented from environments described as fully or partially marine, suggesting the

organisms represented by these fossils inhabited marine environments (19, 25).

However, we cannot rule out the possibility of a terrestrial habitat for these

organisms, since their dead remains could have been transported from terrestrial to

marine environments where they fossilized. To explore this issue, we combine the

fossil evidence and the phylogenomic reconstruction with branch lengths to

reconstruct the potential ancestral habitat states (i.e., marine habitat *vs.* terrestrial

habitat) of key Arcellinida clades through the BayesTraits MultiState method (86).

Specifically, we used the 100 topologies with branch lengths obtained for the
phylogenomic Real Bootstrap topological support assessment (see *Phylogenomic*
analyses section) and implemented four different ancestral reconstruction analyses:
i. ancestral reconstruction without fossilizing (assigning a fixed ancestral state value)
nodes; ii. ancestral reconstruction fossilizing Arcellinida node as terrestrial, which
interprets the organisms represented by VSMs as terrestrial; iii. ancestral
reconstruction fossilizing Arcellinida node as marine, which interprets the organisms
represented by VSMs as marine; iv. ancestral reconstruction fossilizing Arcellinida
and Organoconcha+Glutinoconcha nodes as marine, which interprets the organisms
represented by VSMs as derived crown Arcellinida that lived in marine habitat
(**DatasetS01, Table S12 and Appendix S01, Fig. S44**). To compare the
reconstructed scenarios, we applied a likelihood ratio test (LRT), which we
considered as significant a difference of LRT $\geq$ 2 (87).

Data Availability

Raw sequencing files are deposited at the NCBI SRA repository under the Bioproject
PRJNA1032600. Phylogenomic supermatrix, single gene marker datasets, and input
information for the molecular clock and ancestral reconstruction analyses are
presented in DatasetS01. All molecular data associated with this manuscript are
available on FigShare (10.6084/m9.figshare.25749276). This includes transcriptome
assemblies, predicted proteomes, alignments (trimmed and untrimmed), as well as
phylogenetic trees.

Acknowledgments

We are deeply indebted to Susannah Porter for providing thoughtful insights to an
earlier version of this manuscript, as well as a second anonymous reviewer. We
thank Enrique Lara for the discussions and critiques during the construction of this
project. We thank Cori Tice for her help with R and advice on plot aesthetics. This
work was supported in part by the Fundação de Amparo à Pesquisa do Estado de
São Paulo (FAPESP) grants 2019/22692-8 and 2021/09529-0 awarded to A.L.P.S.
and grant 2019/22815-2 awarded to D.J.G.L., and by the United States National
Science Foundation (NSF) Division of Environmental Biology (DEB) grant 2100888

(http://www.nsf.gov) awarded to M.W.B. KD was funded by the German Research Foundation grant 399699069 - DU 1863/1, Y.E. was supported by NSERC grant 298366-2019 to Alastair Simpson (Dalhousie University).

References

1. S. Geisen, *et al.*, Soil protists: a fertile frontier in soil biology research. *FEMS Microbiol. Rev.* 42, 293–323 (2018).

2. D. Singer, *et al.*, Protist taxonomic and functional diversity in soil, freshwater and marine ecosystems. *Environ. Int.* 146, 106262 (2021).

3. P. A. Cohen, L. A. Riedman, It's a protist-eat-protist world: recalcitrance, predation, and evolution in the Tonian–Cryogenian ocean. *Emerg. Top. Life Sci.* 2, 173–180 (2018).

4. I. Zachar, E. Szathmáry, Breath-giving cooperation: Critical review of origin of mitochondria hypotheses: Major unanswered questions point to the importance of early ecology. *Biol. Direct* 12, 1–26 (2017).

5. L. Eme, T. J. Ettema, The eukaryotic ancestor shapes up. *Nature* 562, 352-353 (2018).

6. S. M. Porter, Insights into eukaryogenesis from the fossil record. *Interface Focus.* 10, 20190105 (2020).

7. S. Xiao, Q. Tang, After the boring billion and before the freezing millions: evolutionary patterns and innovations in the Tonian Period. *Emerg. Top. Life Sci.* 2, 161–171 (2018).

8. G. Li, *et al.*, An assemblage of macroscopic and diversified carbonaceous compression fossils from the Tonian Shiwangzhuang Formation in western Shandong, North China. *Precambrian Res.* 346, 105801 (2020).

9. Q. Tang, K. Pang, X. Yuan, S. Xiao, A one-billion-year-old multicellular chlorophyte. *Nat. Ecol. Evol.* 4, 543–549 (2020).

10. A. Del Cortona, *et al.*, Neoproterozoic origin and multiple transitions to
macroscopic growth in green seaweeds. *Proc. Natl. Acad. Sci. U.S.A.* 117, 2551–
2559 (2020).

11. J. J. Brocks, *et al.*, Lost world of complex life and the late rise of the eukaryotic
crown. *Nature* 618, 767–773 (2023).

12. P. A. Cohen, R. B. Kodner, The earliest history of eukaryotic life: uncovering an
evolutionary story through the integration of biological and geological data. *Trends*
Ecol. Evol. 37, 246–256 (2022).

13. D. B Mills, *et al.*, Eukaryogenesis and oxygen in earth history. *Nat. Ecol. Evol.* 6,
520–532 (2022).

14. J. Kang, B. Gill, R. Reid, F. Zhang, S. Xiao, Nitrate limitation in early
Neoproterozoic oceans delayed the ecological rise of eukaryotes. *Sci. Adv.* 9,
eade9647 (2023).

15. R. Siever, The silica cycle in the Precambrian. *Geochim. Cosmochim. Acta* 56,
3265–3272 (1992).

16. C. T. Reinhard, *et al.*, Evolution of the global phosphorus cycle. *Nature* 541,
386–389 (2017).

17. C. T. Reinhard, *et al.*, The impact of marine nutrient abundance on early
eukaryotic ecosystems. *Geobiology* 18, 139–151 (2020).

18. S. M. Porter, A. H. Knoll, Testate amoebae in the Neoproterozoic Era: evidence
from vase-shaped microfossils in the Chuar Group, Grand Canyon. *Paleobiology* 26,
360–385 (2000).

19. S. M. Porter, R. Meisterfeld, A. H. Knoll, Vase-shaped microfossils from the
Neoproterozoic Chuar Group, Grand Canyon: a classification guided by modern
testate amoebae. *J. Paleontol.* 77, 409–429 (2003).

20. L. Morais, *et al.*, Insights into vase-shaped microfossil diversity and
Neoproterozoic biostratigraphy in light of recent Brazilian discoveries. *J. Paleontol.*
93, 612–627 (2019).

21. D. J. Lahr, *et al.*, Phylogenomics and morphological reconstruction of Arcellinida
testate amoebae highlight diversity of microbial eukaryotes in the Neoproterozoic.
Curr. Biol. 29, 991–1001 (2019).

22. S. M. Porter, L. A. Riedman, Evolution: ancient fossilized amoebae find their
home in the tree. *Curr. Biol.* 29, R212–R215 (2019).

23. C. Berney, J. Pawlowski, A molecular time-scale for eukaryote evolution
recalibrated with the continuous microfossil record. *Proc. Royal Soc. B: Biol. Sci.*
273, 1867–1872 (2006).

24. M. M. Mus, M. Moczydłowska, A. H. Knoll, Morphologically diverse vase-shaped
microfossils from the Russøya Member, Elbobreen Formation, in Spitsbergen.
Precambrian Res. 350, 105899 (2020).

25. D. J Lahr, An emerging paradigm for the origin and evolution of shelled
amoebae, integrating advances from molecular phylogenetics, morphology and
paleontology. *Mem. Inst. Oswaldo Cruz* 116, e200620 (2021).

26. C. Strullu-Derrien, P. Kenrick, T. Goral, A. H. Knoll, Testate amoebae in the 407-
million-year-old Rhynie Chert. *Curr. Biol.* 29, 461–467 (2019).

27. S. Kang, *et al.*, Between a pod and a hard test: the deep evolution of amoebae.
Mol. Biol. Evol. 34, 2258–2270 (2017).

28. C. M. Dehler, C. M. Fanning, P. K. Link, E. M. Kingsbury, D. Rybczynski,
Maximum depositional age and provenance of the Uinta Mountain Group and Big
Cottonwood Formation, northern Utah: Paleogeography of rifting western Laurentia.
Bulletin 122, 1686–1699 (2010).

29. V. N. Sergeev, J. W. Schopf, Taxonomy, paleoecology and biostratigraphy of the
late Neoproterozoic Chichkan microbiota of South Kazakhstan: the marine biosphere
on the eve of metazoan radiation. *J. Paleontol.* 84, 363–401 (2010).

30. J. V. Strauss, A. D. Rooney, F. A. Macdonald, A. D. Brandon, A. H. Knoll, 740
Ma vase-shaped microfossils from Yukon, Canada: Implications for Neoproterozoic
chronology and biostratigraphy. *Geology* 42, 659–662 (2014).

31. L. A. Riedman, S. M. Porter, C. R. Calver, Vase-shaped microfossil
biostratigraphy with new data from Tasmania, Svalbard, Greenland, Sweden and the
Yukon. *Precambrian Res.* 319, 19–36 (2018).

32. P. A. Cohen, S. W. Irvine, J. V. Strauss, Vase-shaped microfossils from the
Tonian Callison Lake Formation of Yukon, Canada: taxonomy, taphonomy and
stratigraphic palaeobiology. *Palaeontology* 60, 683–701 (2017).

33. M. Moczydłowska, V. Pease, S. Willman, L. Wickström, H. Agić, A Tonian age for
the Visingsö Group in Sweden constrained by detrital zircon dating and
biochronology: implications for evolutionary events. *Geol. Mag.* 155, 1175–1189
(2018).

34. B. Freitas, *et al.*, Cryogenian glaciostatic and eustatic fluctuations and massive
Marinoan-related deposition of Fe and Mn in the Urucum District, Brazil. *Geology* 49,
1478–1483 (2021).

35. A. D. Rooney, *et al.*, Coupled Re-Os and U-Pb geochronology of the Tonian
Chuar Group, Grand Canyon. *GSA Bull.* 130, 1085–1098 (2018).

36. C. Dehler, *et al.*, Precise U-Pb age models refine Neoproterozoic western
Laurentian rift initiation, correlation, and Earth system changes. *Precambrian Res.*
396, 107156 (2023).

37. A. K. Tice, *et al.*, Phylofisher: a phylogenomic package for resolving eukaryotic
relationships. *PLoS Biol.* 19, e3001365 (2021).

38. R. E. Jones, *et al.*, Create, Analyze, and Visualize Phylogenomic Datasets Using
PhyloFisher. *Curr. Protoc.* 4, e969 (2024).

39. J. F. Strassert, I. Irisarri, T. A. Williams, F. Burki, A molecular timescale for
eukaryote evolution with implications for the origin of red algal-derived plastids. *Nat.*
Commun. 12, 1879 (2021).

40. L. W. Parfrey, D. J. Lahr, A. H. Knoll, L. A. Katz, Estimating the timing of early
eukaryotic diversification with multigene molecular clocks. *Proc. Natl. Acad. Sci.*
U.S.A. 108, 13624–13629 (2011).

41. L. Eme, S. C. Sharpe, M. W. Brown, A. J. Roger, On the age of eukaryotes:
evaluating evidence from fossils and molecular clocks. *Cold Spring Harb. Perspect.*
Biol. 6, a016139 (2014).

42. M. Dos Reis, *et al.*, Uncertainty in the timing of origin of animals and the limits of
precision in molecular timescales. *Curr. Biol.* 25, 2939–2950 (2015).

43. A. R. Schmidt, W. Schönborn, U. Schäfer, Diverse fossil amoebae in German
Mesozoic amber. *Palaeontology* 47, 185–197 (2004).

44. A. Farooqui, A. Kumar, N. Jha, A. Pande, D Bhattacharya, A thecamoebian
assemblage from the Manjir formation (early Permian) of northwest Himalaya, India.
Earth Sci. India 3 (2010).

45. V. Girard, D. Néraudeau, S. M. Adl, G. Breton, Protist-like inclusions in amber, as
evidenced by Charentes amber. *Eur. J. Protistol.* 47, 59–66 (2011).

46. L. A. Riedman, P. M. Sadler, Global species richness record and biostratigraphic
potential of early to middle Neoproterozoic eukaryote fossils. *Precambrian Res.* 319,
6–18 (2018).

47. R. Guilbaud, S. W. Poulton, N. J. Butterfield, M. Zhu, G. A. Shields-Zhou, A
global transition to ferruginous conditions in the early Neoproterozoic oceans. *Nat.*
Geosci. 8, 466–470 (2015).

48. H. Wang, *et al.*, A benthic oxygen oasis in the early Neoproterozoic ocean.
Precambrian Res. 355, 106085 (2021).

49. S. Porter, The rise of predators. *Geology* 39, 607–608 (2011).

50. A. H. Knoll, Paleobiological perspectives on early eukaryotic evolution. *Cold*
Spring Harb. Perspect. Biol. 6, a016121 (2014).

51. S. Geisen, *et al.*, Pack hunting by a common soil amoeba on nematodes.
Environ. Microbiol. 17, 4538–4546 (2015).

52. K. Dumack, C. Kahlich, D. J. Lahr, M. Bonkowski, Reinvestigation of *Phryganella*
paradoxa (Arcellinida, Amoebozoa) Penard 1902. *J. Eukaryot. Microbiol.* 66, 232–
243 (2019).

53. A. H. Estermann, *et al.*, Fungivorous protists in the rhizosphere of *Arabidopsis*
thaliana–diversity, functions, and publicly available cultures for experimental
exploration. *Soil Biol. Biochem.* 187, 109206 (2023).

54. K. Dumack, *et al.*, It's time to consider the Arcellinida shell as a weapon. *Eur. J.*
Protistol. 92, 126051 (2024).

55. S. M. Porter, Tiny vampires in ancient seas: evidence for predation via
perforation in fossils from the 780–740 million-year-old Chuar Group, Grand Canyon,
USA. *Proc. Royal Soc. B: Biol. Sci.* 283, 20160221 (2016).

56. C. Cousyn, *et al.*, Rapid, local adaptation of zooplankton behavior to changes in
predation pressure in the absence of neutral genetic changes. *Proc. Natl. Acad. Sci.*
U.S.A. 98, 6256–6260 (2001).

57. H. D. Rundle, S. M. Vamosi, D. Schluter, Experimental test of predation's effect
on divergent selection during character displacement in sticklebacks. *Proc. Natl.*
Acad. Sci. U.S.A. 100, 14943–14948 (2003).

58. J. R. Meyer, R. Kassen, The effects of competition and predation on
diversification in a model adaptive radiation. *Nature* 446, 432–435 (2007).

59. P. F. Hoffman, Z. X. Li, A palaeogeographic context for Neoproterozoic
glaciation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 277, 158–172 (2009).

60. A. D. Rooney, J. V. Strauss, A. D. Brandon, F. A. Macdonald, A Cryogenian
chronology: Two long-lasting synchronous Neoproterozoic glaciations. *Geology* 43,
459–462 (2015).

61. J. P. Pu, *et al.*, Dodging snowballs: Geochronology of the Gaskiers glaciation
and the first appearance of the Ediacaran biota. *Geology* 44, 955–958 (2016).

62. B. Runnegar, Loophole for snowball Earth. *Nature* 405, 403–404 (2000).

63. W. Vincent, *et al.*, Ice shelf microbial ecosystems in the high arctic and
implications for life on snowball Earth. *Naturwissenschaften* 87, 137–141 (2000).

64. P. F. Hoffman, D. P. Schrag, The snowball Earth hypothesis: testing the limits of
global change. *Terra nova* 14, 129–155 (2002).

65. R. M. Morgan-Kiss, J. C. Priscu, T. Pocock, L. Gudynaite-Savitch, N. P. Huner,
Adaptation and acclimation of photosynthetic microorganisms to permanently cold
environments. *Microbiol. Mol. Biol. Rev.* 70, 222–252 (2006).

66. A. J. Campbell, E. D. Waddington, S. G. Warren, Refugium for surface life on
Snowball Earth in a nearly-enclosed sea? A first simple model for sea-glacier
invasion. *Geophys. Res. Lett.* 38 (2011).

67. A. J. Campbell, E. D. Waddington, S. G. Warren, Refugium for surface life on
Snowball Earth in a nearly enclosed sea? A numerical solution for sea-glacier
invasion through a narrow strait. *J. Geophys. Res. Oceans.* 119, 2679–2690 (2014).

68. H. M. Lawal, *et al.*, Cold climate adaptation is a plausible cause for evolution of
multicellular sporulation in Dictyostelia. *Sci. Rep.* 10, 8797 (2020).

69. L. Morais, *et al.*, Diverse vase-shaped microfossils within a Cryogenian glacial
setting in the Urucum Formation (Brazil). *Precambrian Res.* 367, 106470 (2021).

70. M. M. Mus, M. Moczydowska, Internal morphology and taphonomic history of the
Neoproterozoic vase-shaped microfossils from the Visings Group, Sweden. *Nor.*
Geol. Tidsskr. 80, 213–228 (2000).

71. G. J. Retallack, Why was there a Neoproterozoic Snowball Earth? *Precambrian*
Res. 385, 106952 (2023).

72. K. Wang, *et al.*, Shallow-marine testate amoebae with internal structures from
the Lower Devonian of China. *Iscience* 26 (2023).

73. C. H. Wellman, P. K. Strother, The terrestrial biota prior to the origin of land
plants (embryophytes): a review of the evidence. *Palaeontology* 58, 601–627 (2015).

74. J. Žárský, V. Žárský, M. Hanáček, V. Žárský, Cryogenian glacial habitats as a
plant terrestrialisation cradle–the origin of the anydrophytes and Zygnematophyceae
split. *Front. Plant Sci.* 12, 735020 (2022).

75. F. Lutzoni, *et al.*, Contemporaneous radiations of fungi and plants linked to
symbiosis. *Nat. Commun.* 9, 5451 (2018).

76. S. Picelli, *et al.*, Full-length RNA-seq from single cells using Smart-seq2. *Nat.*
Protoc. 9, 171–181 (2014).

77. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30, 2114–2120 (2014).

78. B. J. Haas, *et al.*, *De novo* transcript sequence reconstruction from RNA-seq
using the Trinity platform for reference generation and analysis. *Nat. Protoc.* 8,
1494–1512 (2013).

79. M. Manni, M. R. Berkeley, M. Seppey, E. M. Zdobnov, BUSCO: assessing
genomic data quality and beyond. *Curr. Protoc.* 1, e323 (2021).

80. B. Q. Minh, *et al.*, IQ-TREE 2: new models and efficient methods for phylogenetic
inference in the genomic era. *Mol. Biol. Evol.* 37, 1530–1534 (2020).

81. H. C. Wang, B. Q. Minh, E. Susko, A. J. Roger, Modeling site heterogeneity with
posterior mean site frequency profiles accelerates accurate phylogenomic
estimation. *Syst. Biol.* 67, 216–235 (2018).

82. A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. *Bioinformatics* 30, 1312–1313 (2014).

83. C. Zhang, M. Rabiee, E. Sayyari, S. Mirarab, ASTRAL-III: polynomial time
species tree reconstruction from partially resolved gene trees. *BMC Bioinform.* 19,
15–30 (2018).

84. M. J. Benton, *et al.*, Constraints on the timescale of animal evolutionary history.
Palaeontol. Electron. 18.1.1FC; 1-106 (2015).

85. Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.*
24, 1586–1591 (2007).

86. A. Meade, M. Pagel, "Ancestral state reconstruction using BayesTraits" in
Environmental Microbial Evolution: Methods and Protocols, H. Luo, Eds. (Springer,
2022), pp. 255–266.

87. M. Pagel, The maximum likelihood approach to reconstructing ancestral
character states of discrete characters on phylogenies. *Syst. Biol.* 48, 612–622
(1999).

Figures

Figure 1. Sampled testate amoebae — Arcellinida and Corycidia. The pictured
organisms were photodocumented prior to molecular processing and represent
individuals or cultures from which the transcriptomic data was obtained. The scale
bars represent 20 μm, except when specified. **A.** *Phryganella paradoxa* T, lateral
view. **B.** *Phryganella acropodia* A, apertural view. **C - D.** *Microcorycia aculeata*,
dorsal view (C) and apertural view (D); **E.** *Microcorycia flava*, dorsal view focusing on
the flexible part of the shell; **F.** *Spumochlamys* sp., dorsal view*;* **G.** *Spumochlamys*
bryora, lateral view; **H - K.** *Heleopera lucida* comb. nov. (previously *Difflugia lucida*),
lateral view focusing on the cell within the shell (H), lateral view focusing on the shell
(I - J), and apertural view focusing on the compressed aspect of the shell (K); **L.**
Difflugia cf. *capreolata*, lateral view, scale bar 40 μm; **M.** *Netzelia lobostoma*, lateral
view, white square focusing on details of the shell; **N.** *Cyclopyxis* sp., apertural view,
scale bar 40 μm; **O.** *Galeripora* sp., apertural view, white square focusing on the
pores which surround the shell aperture; **P.** *Trichosphaerium sp.* KSS. Measured

morphometric characteristics of the newly sequenced testate amoebae taxa are
present on **Dataset S01, Table S2**.

Figure 2. The tree of amoebozoan testate amoebae. 226 genes (70,428 amino acid sites) phylogeny of amoebozoan testate amoebae rooted with Evosea (Amoebozoa). The tree was initially built using IQ-TREE2 v. 2.0-rc1 under the LG+C20+G4 model of protein evolution and further used to infer a Posterior Means Site Frequency model using the ML model LG+C60+G4+PMSF. Topological support was assessed by 100 Maximum Likelihood Real Bootstrap Replicates (MLRB) and local posterior probability values (LPP) calculated using ASTRAL-III v. 5.7.3, and are shown in the format (MLRB/LPP).

911

912 Figure 3. Amorphea time-calibrated tree inferred under autocorrelated relaxed 913 clock model, applying a truncated-Cauchy distribution for node calibration and 914 drift parameter of α = 2 and β = 2, considering VSMs as stem Arcellinida to 915 calibrate Arcellinida+Euamoebida node. Bars at nodes are 95% highest 916 probability density confidence intervals (HPD CI). Asterisks indicate the nodes 917 calibrated based on external fossil information. Out of the 36 time-calibrated trees 918 generated, we display here the one representing the analysis that estimated the 919 youngest minimum 95% HPD CI for the two earliest nodes of Arcellinida, thus 920 showing the youngest age estimated for the origin of nodes leading to extant 921 members of the Arcellinida Order. The results and time-calibrated trees for all

experiments are present in the supplemental material (**Dataset S01, Table S11 and**
Appendix S01, Fig. S6 - S43). Abbreviations: Exc.- Excentrotoma; Vonust. -
Volnustoma; Paleo. - Paleoproterozoic; Sta. - Statherian; Cryo. - Cryogenian; Ediac.
- Ediacaran; Ca. - Cambrian; Or. Ordovician; Si - Silurian; De. - Devonian; Car.
Carboniferous; Per. - Permian; Tri. - Triassic; Jur. - Jurassic; Cret. - Cretaceous;
Cen. - Cenozoic; Pal. - Paleogene; N. - Neogene; My - Million Years.

Figure 4. Estimated times for Arcellinida nodes throughout geological time. The figure highlights key carbon anomalies, ice age events, the Arcellinida fossil record (19, 26, 36, 44, 72), and the divergence time of other groups of organisms

suggested by previous studies (horizontal dotted lines; 75). Displayed for each node
are four bars representing, from bottom to top, the calibration strategy not
considering VSM record to calibrate amoebozoan nodes (gray bar), the calibration
strategy considering VSMs as derived crown Arcellinida, calibration strategy
considering VSMs as basal crown Arcellinida, and calibration strategy considering
VSMs as stem Arcellinida. The bars represent the combination of all 95% highest
probability density confidence intervals estimated by each distribution-clock model
considered and the colored dots represent the mean estimated time by each
distribution-clock model. The results and time-calibrated trees for all experiments are
present in the supplemental material (**Dataset S01, Table S11 and Appendix S01,**
Fig. S6 - S43).

Figure 5. Ancestral reconstructions of Arcellinida habitats using BayesTraits. Pie charts at each node indicate the mean probabilities of a hypothetical marine ancestor (blue) or a hypothetical terrestrial ancestor (purple). **A.** Reconstruction considering Arcellinida ancestor as marine, implying at least two independent transition events (2TE). **B.** Reconstruction considering Arcellinida and Organoconcha+Glutinoconcha ancestors as marine, implying at least three independent transition events (3TE). Bars at nodes are 95% highest probability density confidence intervals estimated by the calibration strategy using an autocorrelated relaxed clock model, applying a truncated-Cauchy distribution for node calibration and drift parameter of α = 2 and β = 2, considering VSMs as stem Arcellinida to calibrate Arcellinida+Euamoebida node. The complete results are shown on **Dataset S01, Table S12 and Appendix S01, Fig. S44**.