Divergent patterns of richness and density in the global soil seed bank

Alistair G. Auffret [1⬚*], Emma Ladouceur [2,3,4,5⬚*], Natalie S. Haussmann[6], Petr Keil [7], Eirini Daouti [1,8], Tatiana G. Elumeeva [9], Ineta Kačergytė [1], Jonas Knape [1], Dorota Kotowska[10,11], Matthew Low [1], Vladimir G. Onipchenko[9], Matthieu Paquet [1,12], Diana Rubene[8, 13], Jan Plue[14]

Author Affiliations

[1] Department of Ecology, Swedish University of Agricultural Sciences, 75 007 Uppsala, Sweden

[2] Department of Biology, University of Prince Edward Island, PE, Canada

[3] Canadian Centre for Climate Change and Adaptation, University of Prince Edward Island, St. Peter's Bay, PE, C0A 2A0, Canada

[4] School of Climate Change and Adaptation, University of Prince Edward Island, Charlottetown, PE, C1A 4P3, Canada

[5] German Centre for Integrative Biodiversity Research (iDiv) Leipzig-Halle-Jena, Puschstraße 4, 04103 Leipzig, Germany

[6] Department of Geography, Geoinformatics and Meteorology, University of Pretoria, Lynnwood Rd, Hatfield 0002, South Africa

[7] Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, 165 00 Praha-Suchdol, Czech Republic

[8] Department of Crop Production Ecology, Swedish University of Agricultural Sciences, 75 007 Uppsala, Sweden

[9] Moscow Lomonosov State University, Moscow, 119234, Russia

[10] Institute of Nature Conservation, Polish Academy of Sciences, Kraków, Poland

[11] 'Lendület' Landscape and Conservation Ecology Group, Institute of Ecology and Botany, MTA–HUN-REN Centre for Ecological Research, Vácrátót, Hungary

[12] Theoretical and Experimental Ecology Station (SETE), National Center for Scientific Research (CNRS), Moulis, France

[13] Greensway AB, Gerda Nilssons väg 2, 756 51 Uppsala, Sweden

[14] Swedish Biodiversity Centre, Swedish University of Agricultural Sciences, 75 007 Uppsala, Sweden

⬚Alistair G. Auffret and Emma Ladouceur should be considered joint first authors
* Corresponding authors alistair.auffret@slu.se, erladouceur@upei.ca

Abstract

Soil seed banks are an important component of plant population and community dynamics, buffering temporal heterogeneity and allowing populations to recover following disturbance. At the same time, patterns in soil seed banks are likely to reflect scale-dependent patterns in above-ground vegetation, patterns that differ strongly across world regions. Here, we investigate components of diversity in the soil seed bank across global biomes and ecosystems. We found that patterns of species richness in the soil seed bank diverged from patterns of seed density, although there were high levels of uncertainty, especially at smaller spatial scales. Importantly, habitat degradation consistently led to lower richness, potentially limiting future ecosystem recovery. Among ecosystems, mediterranean and tropical regions had high species richness, but seed density m^{-2} in the soil was highest in wetlands and arable systems. Lower seed densities were found in both high-diversity tropical biomes that are characterised by short-lived seeds, and low-diversity boreal and tundra biomes with more stable established vegetation. Our findings of divergences between species richness and seed density across global biomes, and how they are impacted by habitat degradation, give valuable insights to the understanding of plant biodiversity worldwide.

Introduction

In an era of global environmental change, the persistence of populations in newly-unsuitable conditions and the establishment of new populations following disturbance are key in determining changes in biodiversity over time and in space. For vascular plants, the soil seed bank can play a vital role in both processes. Soil seed banks are the communities of persistent seeds being naturally stored in the soil that contribute to vegetation regeneration after disturbance[1,2]. Plant species that form seed banks do so as part of a life-history

strategy that allows for the buffering of temporal environmental heterogeneity[3–5], and as

such, seed banks are an important component for understanding plant (meta)population and

community dynamics[6,7].

Despite the potential advantage of producing seeds that persist in the soil, not all species

contribute to soil seed banks. Instead, the ability to form seed banks is a functional strategy

for plants to (re)establish and maintain populations, which has evolved in response to both

predictable and unpredictable disturbance regimes[8]. The species that do form seed banks

also vary in terms of the number of seeds produced, and the longevity of seeds in the soil[9].

The disturbances that have contributed to the development of seed banking are not

randomly distributed in space, and as such plant communities in different bioclimatic

environments should rely on seed banking to different extents, with implications for local

seed bank size and composition. For example, in arid systems dominated by drought, it is

advantageous to have an annual life history and form persistent seed banks to buffer annual

changes in water availability[10,11]. A similar pattern exists in temperate Europe, where species

with warmer distributions have been found to have a higher seed bank persistence than

more northerly species[12]. In tropical systems, where climatic conditions are broadly stable

and exposure to freezing temperatures and drought conditions is rare, relatively few species

produce seeds that are able to survive desiccation[13,14] – a trait associated with long-term

persistence – which may result in relatively small and species-poor soil seed banks.

While it can be interesting to broadly compare plant communities from different world

regions, ecological assemblages can differ considerably among habitats within the same

region. The frequency of disturbance within a habitat is generally mirrored by the strategies

of the resulting plant communities[15]. Therefore, different habitats could be expected to

support different soil seed banks, in terms of the number of species and the density of seeds

present. For example, in temperate and boreal regions, forests are dominated by long-lived

trees, with the understorey often covered by species that largely rely on vegetative

reproduction rather than seed banking[16,17]. Species-rich grazed grasslands in the same

regions are characterised by regular, low-intensity and spatially-heterogeneous

disturbances, and as such harbour plants with a wide range of regeneration strategies[18,19].

Intensively-used arable lands on the other hand are well-known to support large

communities of ruderal species with annual life-histories and high seed production[20,21]. As

such, habitat is an important component in understanding how different functional traits are

represented in different regions, while differences between broad habitat types can differ in

different world regions[22]. Moreover, communities in the same broad habitat categories can

differ in their responses to environmental degradation in different regions[23]. Therefore, when

studying plant communities at global scales, it is important to consider the potential roles of

different habitats and environmental degradation driving any variation within or between

regions. This could be especially relevant in the case of soil seed banks, which exist partially

as a strategy to respond to disturbance.

Studying patterns of soil seed bank diversity across large spatial scales and geographic

gradients can be useful for understanding the biogeography and macroecology of their role

in biodiversity maintenance and recovery[24–26]. However, biodiversity is known to be scale-

dependent, and comparisons of soil seed banks across ecoregions and habitats are

therefore likely to be influenced by the scale at which diversity is estimated. It has been

established that the number of species in soil seed banks increases with area[24,25], but while

it is well-known that this accumulation of species varies according to habitat and region in

the established vegetation as well as in other taxa[27–30], such patterns are still largely

unknown in the soil seed bank, at least not at spatial scales that span multiple biomes.

Integrating a scale-explicit approach to understanding diversity in the soil seed bank across

habitats, within and between ecoregions and in response to habitat degradation is an

important step for plant community and macroecology, allowing critical insights for both

broad scale patterns of biodiversity and the context-dependent interpretations of local

dynamics.

While individual seed bank studies are usually (through necessity) small-scale comparisons

across limited environmental gradients within a specific region, global analysis allows the

large-scale comparison of biodiversity of the seed bank. Here, we use a global database of

soil seed banks, harnessing 1442 studies from all seven of the world's continents (Fig. 1), to

reveal how broad differences in climatic conditions across world regions, and differences in

disturbance regimes across habitat types, combine to determine patterns in both the

richness and density of soil seed bank communities at different spatial scales. The global

seed bank database[31] is the result of a comprehensive search of the seed bank literature,

providing 3096 records containing information regarding the richness and/or density of the

soil seed bank at a given location (Methods). To investigate patterns of seed bank diversity

among ecosystems and global biomes, we used both the WWF map of global ecoregions[32]

and the habitat descriptions used by the authors of the component studies to separate

records into 15 ecosystems. These ecosystems are defined as the combination of biome

(e.g. forest, grassland) and ecoregion (e.g. temperate, tropical). Each ecosystem was

assigned to either the terrestrial, aquatic or transitional realm[33]; see Fig. 1 and Table 1.

We used a hierarchical Bayesian modelling approach to investigate different components of

the soil seed bank separately per biome, with ecoregion, habitat degradation (binary:

undisturbed or degraded) and sampling effort (area sampled and number of sites) included

as predictor variables. Specifically, we investigated [1] species richness, finding highest

richness in tropical forests and mediterranean ecosystems, but differences among

ecosystems and biomes, reflecting our broad knowledge of global patterns in the

aboveground vegetation; [2] seed density (m^{-2}), finding highest values in wetlands and

arable ecosystems; and [3] the impact of habitat degradation, finding consistently lower

richness in degraded habitats, with patterns of density. Importantly, we found that patterns of

species richness in the soil seed bank diverged from patterns of density. That is, a high

density of seeds in the soil seed bank does not necessarily translate to a large number of

species, and that this varies among ecosystems, biomes and in response to degradation.

150
151
Figure 1. The global record of the soil seed bank. Points indicate the location of each
record collected from the literature (n=3096; some records have the same location). Records
are first split according to biome (e.g. tundra, forests; panels a-g), with each biome split
according to one or more ecoregions (e.g. boreal, tropical; colours of points). The
combination of biome and ecoregion results in 15 unique ecosystems. Each record is also
designated either as occurring in undisturbed or degraded ecosystems, apart from arable
records which have their own category (shapes of points). We also categorise each biome
as belonging to the **a-e** Terrestrial, **f** Transitional, or **g** Aquatic realm. See Methods for a
more detailed description of all categories.

169 **Table 1. Division of soil seed bank observation into ecosystems.** Biome and ecoregion are defined according to author-reported habitat types and
170 geographical association to global ecoregions (Methods). The combination of biome and ecoregion is what we refer to as an ecosystem. 'Degraded' indicates
171 whether the area samples were taken from was considered to be degraded by authors of the original study. The number of observations (N_{obs}) collected for
172 each ecoregion, and degraded status listed for species richness, and seed density m^{-2}.

Realm	Biome	Ecoregion	Degraded	N_{obs} species richness	N_{obs} seed density
Terrestrial	Tundra	Tundra	No	43	40
Terrestrial	Tundra	Tundra	Yes	3	3
Terrestrial	Forest	Boreal	No	26	22
Terrestrial	Forest	Boreal	Yes	11	11
Terrestrial	Forest	Temperate	No	266	285
Terrestrial	Forest	Temperate	Yes	160	144
Terrestrial	Forest	Tropical	No	155	152
Terrestrial	Forest	Tropical	Yes	136	135
Terrestrial	Grasslands and Savannas	Temperate and Boreal	No	475	456
Terrestrial	Grasslands and Savannas	Temperate and Boreal	Yes	340	265
Terrestrial	Grasslands and Savannas	Tropical	No	51	45
Terrestrial	Grasslands and Savannas	Tropical	Yes	33	31
Terrestrial	Mediterranean and Desert	Deserts and Xeric Shrublands	No	107	83
Terrestrial	Mediterranean and Desert	Deserts and Xeric Shrublands	Yes	33	35
Terrestrial	Mediterranean and Desert	Mediterranean Forests, Woodlands and Scrub	No	129	121
Terrestrial	Mediterranean and Desert	Mediterranean Forests, Woodlands and Scrub	Yes	69	61
Terrestrial	Arable	Mediterranean and Desert	NA	48	43
Terrestrial	Arable	Temperate and Boreal	NA	159	154
Terrestrial	Arable	Tropical	NA	65	59
Transitional	Wetlands and Flooded Grasslands	Mediterranean and Desert	No	40	23
Transitional	Wetlands and Flooded Grasslands	Mediterranean and Desert	Yes	9	7

Transitional Wetlands and Flooded Grasslands Temperate and Boreal			No	276	249
Transitional Wetlands and Flooded Grasslands Temperate and Boreal			Yes	103	90
Transitional Wetlands and Flooded Grasslands Tropical			No	50	42
Transitional Wetlands and Flooded Grasslands Tropical			Yes	11	10
Aquatic	Aquatic	Aquatic	No	74	64
Total number of observations				2872	2630

174 Results

The global seed bank database[31] contains the results of studies carried out in 94 countries

across the world, including observations of more than 35 million seeds using a large range in

sampling effort and extent (for a more in-depth summary of the data, see Methods). In such

a large and heterogeneous dataset, there was generally a large variation in seed bank

richness and density within and between ecosystems. Here, we describe noteworthy results

in terms of overlap in 50% and 90% credible intervals (hereafter CIs) and consistency in the

relative values of model estimates that are rounded to whole numbers (i.e. whole species or

whole seeds m^{-2}). All model results and further supplementary information can be found in

Tables S1-S4, and Figures S2a-g, and full details of statistical models, including the

definitions, interactions, inclusion and transformation of and between predictor variables are

described in the Methods.

186 Seed bank richness and density across ecosystems and biomes

Using models analysing species richness to predict average soil seed bank diversity for each

ecosystem and at multiple spatial scales, we found that predicted species richness generally

increased from the $0.01m^2$ alpha scale to the 15 m^2 gamma scale. Estimated richness was

always higher at the gamma scale, with 50% credible intervals not overlapping in all

undisturbed and 80% of degraded (including arable) systems. Relative values of predicted

richness varied according to scale, with one example being that boreal and temperate

grasslands had a higher predicted richness than tropical forests at the alpha scale (but with

overlapping CIs), while tropical forests had higher richness at the gamma scale (no overlap

of 50% CIs). Generally, differences in estimated richness across ecosystems were more

clear at the gamma scale, so we place most focus there. In undisturbed systems,

mediterranean forests and shrublands were predicted to have the highest richness

compared to all eleven other undisturbed ecosystems (no overlap of 50% CIs, overlap with

six undisturbed ecosystems at 90% CIs; Figure 2, Table S4). Tundra and aquatic systems

generally had the lowest species richness, each having a lower estimate than the other ten

undisturbed ecosystems, and lower than eight of the other ecosystems (50% CIs).

Undisturbed tropical ecosystems were estimated to be broadly more species-rich than their

equivalent temperate and boreal ecosystems. Estimated richness was higher for tropical

forests than both boreal and temperate forests, for tropical grasslands compared to

temperate and boreal grasslands and tropical arable compared to temperate and boreal

arable, while tropical wetlands had similar estimated richness as temperate and boreal

wetlands. Despite this consistency, estimates were only clearly different in forest

ecosystems (50% CIs), although tropical wetlands were more species-rich than

mediterranean and desert wetlands.

The average estimated number of seeds m^{-2} in the soil seed bank was 4 214 m^{-2} (median 3

590) in the (non-arable) terrestrial realm, 9 434 m^{-2} (median 11 001) for arable systems, 10

318 m^{-2} (median 10 802 m^{-2}) in the transitional realm, and 8 124 m^{-2} (median 9 562) in the

aquatic realm (Figure 3, Table S5). Wetland ecosystems exhibited high seed densities, often

higher than other ecosystems within their equivalent ecoregion (90% CIs). For example,

tropical wetlands had higher seed densities than tropical forests, and temperate and boreal

wetlands had higher seed densities than forests and grasslands in the same regions. Within

the terrestrial realm, undisturbed temperate and boreal grasslands had the highest predicted

seed density, higher than boreal forests and temperate forests (90% CIs).

228
229
Figure 2: Scale-dependent richness in soil seed banks across ecosystems. Panels **a-d**
represent realms, with the terrestrial realm separated into arable and non-arable systems.
Each ecosystem is represented by four density curves and point-and-whiskers, showing
species richness at the alpha (α; 0.01 m^2; lower curves, smaller points) and gamma (γ; 15
m^2; upper curves, larger points) scales, and in undisturbed (coloured curves and circular
points) and degraded (grey curves and triangular points) ecosystems. Note that arable
systems are not separated into disturbed and undisturbed and have only two curves per
ecosystem and a square point. Points represent average predicted species, surrounded by
50% (thick) and 90% (thin) credible intervals. Density curves represent 24 000 predictive
draws of the posterior distributions of overall richness at each scale, within each ecosystem
and degradation group.
241

Figure 3: Density of seeds m⁻² in the soil seed bank across ecosystems. Panels **a-d** represent realms, with the terrestrial realm separated into arable and non-arable systems. Each ecosystem is represented by two density curves and point-and-whiskers, showing soil seed bank density in undisturbed (coloured curves and circular points) and degraded (grey curves and triangular points) ecosystems. Note that arable systems are not separated into disturbed and undisturbed and have only two curves per ecosystem and a square point. Large points show the mean densities, surrounded by 50% (thick) and 90% (thin) credible intervals. Density curves represent 24 000 predictive draws of the posterior distributions of seed density, within each ecosystem and degradation group. Background jittered points represent empirical estimates of seeds density within sampled soil banks. Thick vertical lines in each panel represent the mean fixed effect value across **a** Terrestrial (non-arable records), **b** Terrestrial (arable records), **c** Transitional and **d** Aquatic realms.

257 The effect of habitat degradation

In all ecosystems, habitat degradation always resulted in a lower estimated species richness

at the 15 m^2 scale in terms of predicted values, but again, despite this consistency, the

difference was only clear in mediterranean forests, woodlands and scrub according to the

50% credible intervals. At the 0.01 m^2 scale, the predicted values were often the same in

undisturbed and degraded ecosystems. Compared to species richness, habitat degradation

had much more variable effects on seed density m^{-2} (Figure 3, Figure 4). Broadly speaking,

degradation was more likely to result in lower predicted seed densities compared to

undisturbed habitats in open ecosystems, with clear differences (50% CIs) for tropical

grasslands, mediterranean and desert wetlands and aquatic ecosystems. On the other hand,

seed density estimates were generally higher with degradation in forest ecosystems, with

clear differences in both temperate and tropical forests (50% CIs). No ecosystems exhibited

clear differences in richness or density with degradation according to the 90% CIs.

271 Divergent patterns of richness and density

In many cases, relative patterns of richness and density in the soil seed bank differed both

within and among ecosystems, and in response to degradation. Despite relatively high

species richness, seed bank density was generally low in tropical systems. Tropical forests

had the lowest density across undisturbed ecosystems, being lower than all eleven other

undisturbed ecosystems (no overlap of 50% CIs, of which eight had no overlap of 90% CIs).

Undisturbed tropical wetlands, forests and grasslands had a lower density than their

equivalent ecosystems in temperate and boreal regions (50% CI). Disturbed tropical

grasslands were similarly low in seed density, while arable ecosystems in tropical regions

were lower than the equivalent systems in other regions (90% CI). Otherwise, arable

systems combined relatively low richness with high seed density. Mediterranean and desert,

and temperate and boreal arable ecosystems were predicted to have more seeds in the soil

than all other ecosystems in the terrestrial realm (50% CIs), while arable ecosystems often

had lower richness at the gamma scale than other degraded ecosystems within similar

ecoregions (for example, temperate and boreal arable ecosystems had lower richness than

temperate forests and temperate and boreal wetlands; 50% CIs). On the other hand,

mediterranean forests, woodlands and scrub combined high species richness with relatively

high seed density (higher than all other forest ecosystems, 90% CI), while tundra combined

its low species richness with low density, especially when degraded, where density was

lower than all other degraded ecosystems (50% CI).

Figure 4: The joint relationship between predicted species richness m⁻² and seed density m⁻² in the soil seed bank. Panels **a-g** represent biomes. Each ecosystem is represented by two sets of and point-and-whiskers. Points show the mean estimated species richness and seed density m⁻², surrounded by 50% (thick) and 90% (thin) credible intervals. Circular points denote undisturbed and triangles denote degraded ecosystems. Arable systems are not separated into disturbed and undisturbed and have one (square) point with whiskers per ecosystem.

Discussion

Although there was a high uncertainty in our results, we found that across the world's

biomes and ecosystems, patterns in species richness in the seed bank often diverged from

patterns of seed density. Strikingly, we also found that the richness of soil seed banks

appeared higher in undisturbed compared to degraded habitats, suggesting how across the

world's ecosystems, soil seed banks also reflect the ongoing depletion of biodiversity as a

result of anthropogenic activities[23,34]. This indicates that the potential for seed banks to

contribute to the recovery and restoration of degraded ecosystems might be limited. Seed

density, on the other hand, was not always lower in disturbed habitats, showing how different

aspects of biodiversity can show diverging patterns in response to global change

pressures[35]. Our estimated global average of more than 5000 seeds m^{-2} also means that in

the broadest sense, if you put your finger to the earth, you are likely to be pointing at a seed.

The high density of seeds in the soil is a testament to the formation of soil seed banks as an

important component of plant population and community dynamics, and a valuable strategy

for many plant species to facilitate establishment following disturbance[3,6,36].

Our analyses revealed substantial variation in the diversity of the soil seed bank among the

world's ecosystems, identifying divergent patterns between species richness and seed

density. Although the broad relationship between total abundance and species richness is

well-established[35,37], our understanding of plant biogeography and trait-environment

relationships helps us to interpret these results and put them into context. Overall, species

richness in the soil seed bank reflected the latitudinal gradient in biodiversity[29,30,38]. However,

while tropical and mediterranean ecosystems exhibited relatively high richness compared to

temperate, boreal and tundra regions, the former areas differed strongly in terms of seed

density. Mediterranean forests and shrublands exhibited both high richness and density in

the seed bank (Figures 2-4, Table S3). In regions with warm climates and frequent extreme

climatic events, the soil seed bank can play an important role in vegetation recovery and

stability of biodiversity[39,40]. Therefore, the high plant biodiversity in these regions[41], together

with the large proportion of species producing persistent seeds[42], results in the relatively

large and rich seed banks in these regions. On the other hand, tropical ecosystems host

relatively few species that produce seeds that are able to survive desiccation[13,14], which is a

vital aspect of longevity in the soil. Moreover, the warm and often moist conditions combined

with high diversity of other taxa mean that seeds of species that are able to form seed banks

are subject to degradation, predation and attacks from pathogens[43–45]. Together, these

mechanisms result in low soil seed bank density, but the high overall diversity in the

vegetation means that species richness is still high relative to other ecosystems.

The relatively low richness in the soil seed bank in temperate and boreal regions matched

expectations relating to broad gradients in biodiversity, and previous work suggests that this

effect may be compounded by lower proportions of species in these regions producing

persistent seeds compared to lower-latitude species[12]. In these regions, the divergence in

patterns of richness and density was more apparent when comparing ecosystems, with

density clearly higher in grasslands than forests. The vegetation in the temperate and boreal

zones can be considered more stable compared to other regions, and this is especially true

in forests. Because seed banking is expected to provide a competitive advantage in more

dynamic systems[46,47], it follows that these relatively stable systems are instead dominated by

long-lived perennial species, for which regeneration via the soil seed bank is a less important

strategy than for example clonality[48]. In alpine and tundra environments, low seed bank

density and a high proportion of clonal species could also be expected, in this case due to a

low diversity and density of adult individuals, combined with the risks of seed production

during short growing seasons[49,50].

Our results reveal an imprint of anthropogenic activity on soil seed banks worldwide, and

further variation in the relationship between richness and density. While richness was

consistently lower in degraded systems, reflecting patterns in degraded habitats in the

above-ground vegetation[51–53], the effect on density was more variable. Unsurprisingly, arable

soils contained among the highest number of seeds per square metre (Figures 3 & 4). The

high magnitude and frequency of disturbance in arable fields supports relatively species-

poor communities of opportunistic ruderal "weed" species whose traits include both high

seed longevity in the soil and high seed production[20,21], and that can be problematic for

agricultural production[54–56]. In forest ecosystems, lower richness in the seed bank was often

accompanied by higher seed densities. Again, this can be related to plant species'

adaptations to disturbance. While undisturbed forests are relatively stable systems,

degraded forests often retain the imprint of the more dynamic management regimes that

they are experiencing or recovering from, and therefore their seed banks reflect plant

communities that are adapted to such highly-disturbed conditions[57–59]. However, in

grasslands and other habitats, degradation generally appeared to result in both reduced

richness and density in the soil seed bank. Our findings are concerning, matching general

trends of reduced biodiversity in the established vegetation over time[60,61] (but see ref [62]), and

have consequences for the potential of passive restoration to reverse biodiversity losses[63,64].

In addition to detecting differences in species richness in the seed bank among ecosystems,

we also found differences in the accumulation of richness in space. Our findings support

those of Vandvik et al.[25], who observed that species accumulate more rapidly in alpine and

subalpine grasslands than in boreal zones (empirical), and in forests compared to grassland

and heathland (literature), and we extend the results to include other global biomes. We

found that richness estimates across scales (in our case 0.01 m^2 to 15m^2) appeared to show

relatively large differences in mediterranean and tropical ecosystems compared to temperate

ecosystems, demonstrating how species' spatial distributions affect how we interpret relative

differences in biodiversity across regions (Table S3). These findings have implications for

how we understand biodiversity in soil seed banks at both local and regional-global scales.

At local scales, we found that differences in richness across ecosystems, but critically also

between undisturbed and degraded ecosystems, were marginal to non-existent at the 0.01

m^2 scale. This shows that making comparisons at small spatial grains could inhibit our ability

to detect important differences in biodiversity that might be relevant for conservation

management. At larger spatial scales, comparisons among regions can depend on the

spatial grain of predicted richness. This has previously been shown in the aboveground

vegetation, where Sabatini et al.[30] identified a clearer gradient in species richness at large

spatial grains, while Keil & Chase[29] found that areas of relatively high beta diversity in trees

– where richness was relatively low at smaller scales and relatively high at larger scales – to

be concentrated in low latitudes. Our modelled patterns of soil seed bank richness at the

0.01 m^2 scale were broadly similar to those found by Yang et al.[26], although their findings of

low richness in the tropics and (predicted) higher richness in northern regions were not

supported when considering larger spatial scales.

The large number of soil seed bank studies that have been carried out across the world has

allowed our synthesis to cover a greater global extent than many large reviews of

biodiversity, which often suffer from biases relating to the absence or restricted availability of

data from some regions[65–67]. However, our study may still suffer from other biases, which we

discuss here. First, seed banks have not been sampled randomly across the world, but in

places where scientists have been interested in finding and identifying seeds in the seed

bank[68]. This means that our results rest on the assumption that seed banks were not

consistently sampled in areas that are relatively rich or poor in seeds or species, compared

to areas that have not been sampled. Second, the database is made up of studies using

different sampling efforts to sample the soil seed bank. To address variation in sampled

area, we used an approach (ref [29] ,Methods), whereby the putative drivers of variation in

biodiversity are allowed to vary with sample area in our statistical models, meaning that not

only the direct effect of area is controlled for, but also the fact that species-area relationships

can also vary across ecosystems and in response to degradation[28,69]. While we also

controlled for sampling extent (number of sites) in our models, around half of all observations

in the database were represented by a single site. It is well-known that sampling multiple

sites per treatment gives improved confidence in ecological findings (sample replication[70]),

and we echo previous calls for seed bank sampling to improve representativeness[71,72]. Third,

there was a lack of balance in the number of observations across ecosystems, with some

better represented than others. Although we hierarchically modelled each biome separately,

the distribution of observations among ecoregions and between undisturbed and degraded

observations were often uneven. This will have affected our statistical power, increasing

uncertainty in our model estimates, particularly where sample sizes were low.

There is another aspect of the study that will have contributed to the uncertainty in our

results that is not related to the initial data collection. Our binary categorisation of ecosystem

degradation represents a major simplification of the effects of degradation on biodiversity.

While we are convinced that such simple classifications can be useful and informative in

large-scale studies (see e.g. refs [23,63]), our general lack of clear differences between

undisturbed and degraded ecosystems are likely related to the fact that systems undergoing

environmental change are biased towards increasing richness over time[73]. That is, recently

degraded or restored ecosystems are likely to contain both colonising species representing

the new regime, and species representing the previous regime, which have not yet

disappeared. This is especially relevant in the context of seed banks, which exist to buffer

short-term environmental heterogeneity, and whose communities often reflect previous

habitat conditions[74,75]. In more long-term degraded ecosystems, we may expect the negative

effect on seed bank richness to be more clear. It is also important to note that the seed bank

dynamics of – for example – a young post-agricultural forest are likely to be quite different

from those of a mature plantation, both of which would be classed as degraded forests here.

Therefore, while our results may be broadly representative of average effects of degradation

at the global scale, the understanding of specific systems undergoing degradation or

restoration will require more detailed investigation.

What can we learn from studying the soil seed bank? In addition to the continual publication

of studies investigating soil seed banks at local scales, there has been an upswing in the

synthesis of large datasets to understand broad spatial or ecological patterns relating to soil

seed banks. This includes the macro-environmental determinants of small-scale richness

and density[26], and the evolutionary history of seed bank persistence[76]. Here, we complement

such findings with an understanding of the broad biogeographical patterns in soil seed bank

richness and density, as well as the spatial scaling of species richness in the seed bank.

Other large-scale work considers the important role that soil seed banks may play in species

and community responses to global changes, related to the trade-off related to seeds being

able to persist or disperse in space[77], or the role of soil seed banks in facilitating biological

invasions[78]. We believe that the future of soil seed bank synthesis lies beyond simple

measures of community richness and density, or categorisation of species as having or not

having persistent seeds. Collating a large number of datasets including community data from

the soil seed bank and established vegetation – not forgetting to standardise according to

spatial scale – across biomes and systems (e.g. ref [24]), would allow us to delve deeper into a

mechanistic understanding of the role of soil seed banks, biodiversity, restoration and

responses to global change.

486 Methods

487 The global seed bank database

To study the effects of spatial scale, habitat type and quality on soil seed banks, we used a

global database of seed bank richness, abundance and density[31]. Briefly, the database is

based on a Web of Science search using the search term "seed bank*" OR seedbank*,

including papers included in the Web of Science until the end of 2019 (a few papers returned

by the search were subsequently published in journals in 2020). All resulting papers were

checked for being both community seed bank studies (i.e., taking soil samples and counting

and identifying all seeds, rather than focussing on one or more specific species), and

containing information about species richness (number of species), seed abundance

(number of seeds), and/or seed density (number of seeds per unit area or volume) in the

seed bank. Most of the papers were in English, but data were also extracted from papers

published in, for example, Spanish and Portuguese (often South American studies), as well

as French and German. Due to a large data gap, the database was supplemented by an

additional database and literature search featuring studies published in Russian and carried

out in a number of former Soviet states. In total, this resulted in 7224 unique studies to be

checked, and 1442 studies from which records could be extracted or obtained. A record

refers to a row in the database, and represents a known number of soil samples with

specified (by the study authors) geographical coordinates. Depending on the detail in which

study location and design, and seed bank results were reported in a specific study, a record

can represent one or more locations or environments. Each record can also provide one or

more commonly recorded seed bank metrics (mainly total species, total number of seeds,

seeds m^{-2} density), which we term observations. A study can provide multiple observations

by reporting more than one metric, and/or by (for example) reporting metrics across several

distinct records (i.e. different locations or environments). In total, the 1442 studies provided

8087 observations across 3096 records (Figure 1, Table 1, S1). The database includes soil

seed bank data from 94 countries and all seven continents. Full details regarding which data

were extracted and how are available in Auffret et al.[31]. Below, we describe only the

variables in the database that were used for this analysis.

Summary information and data variation in the global seed bank

From the 1185 studies and 2585 observations contributing values of seed density, the total number of seeds or seedlings counted across all studies was 35 932 626. Five studies (or 0.2% of all observations) found no seeds within samples. This included an investigation of aquatic systems in Canada[79], deserts and xeric shrublands in Egypt[80], temperate forests in Germany and the Netherlands[81,82], and tundra in Antarctica[83]. Ten studies observed only one species in their seed bank samples, while the maximum was 246 species in 88 m^2 of soil across 156 arable sites in the midwest of the United States[84]. The minimum (non-zero) observed density of seeds in the seed bank was 3 seeds m^{-2} in a tropical forest in South Africa[85], with the maximum being 501 119 seeds m^{-2} observed in a study from temperate forests in Germany[86]. The minimum number of seeds greater than zero recorded in a soil seed bank observation was two seeds in two studies in temperate forests in the United States, and tropical and subtropical forests in Indonesia with 0.67 seeds m^{-2} and 0.16 m^{-2} of soil sampled, respectively[87,88], with the maximum being 5 907 132 seeds in samples taken in arable systems in 462 m^2 soil from 924 samples across 22 sites in the United Kingdom[89].

Naturally, such raw maximum and minimum values are at least partly an outcome of each investigation's scientific focus and study design. Indeed, there was a large variation in sampling effort across records. The minimum total number of samples was one sample, used in 19 studies, with the maximum being 114 000 samples taken in arable systems representing 224 m^2 of soil sampled across 228 sites in China[90]. The minimum total sampled area was 0.0015 m^2 (5 sites, 15 total samples) in temperate broadleaf and mixed forests from the United States[91], with the maximum being 816 m^2 (20 sites, 20 400 samples) in another large arable study in the UK[92]. The minimum number of sites was one, representing the majority of our samples (1832 observations from 692 studies), with the maximum being 595 sites (total sampled area of 59.5 m^2) in arable fields across Finland[93].

Modelled variables

From the global seed bank database, we extracted the two most common metrics from the soil seed bank literature as response variables: [1] the total number of species recorded in a reported area soil seed bank sampled; [2] the density of seeds m^{-2} in the soil. Soil seed bank studies often involve the collection of multiple samples of soil with a soil corer. These samples are then usually pooled, in order to account for the spatial patchiness of species and seeds in the soil[72,94,95]. Samples can also be collected (and pooled) at several sites in order to capture variation across different locations representing the same habitat or treatment category, often described as 'replicates' within a study. We assume that authors of component papers chose a combination of soil samples and sites that they expected would characterise the soil seed bank in their study system. In our analysis, we used the total sampled area in m^2 or the sample grain as our measure of sampling scale, calculated as the size of the individual soil samples multiplied by the total number of samples taken for a particular observation. We also included the number of sites contributing to each observation as a measure of geographical extent. While this is an imperfect measure, we found it to be a parsimonious covariate, as well as the only such metric that was available in all records. More than half of our observations represent samples collected within a single site (n = 4763).

In order to categorise the global seed bank observations to an ecosystem in a systematic way, we used a combination of authors self-declared habitat descriptions coming from the global seed bank database (aquatic, open grassy systems, forest, arable, or wetland), broad categories of the WWF terrestrial ecoregions based on an observation's geographic coordinates[32], as well as realms in the fashion of the function-based typology for the Earth's ecosystems[33]. Observations with habitat defined as arable or aquatic were placed in an arable and aquatic 'biome', respectively. Habitat and biome were used to categorise all observations into one of 15 categories (Table 1), with the aim to maintain ecologically-

relevant distinctions of ecosystems (the combination of biome and ecoregion) while also

including a reasonable number of observations within each group. Each ecosystem was then

assigned into either the aquatic, terrestrial or transitional realm. Within the terrestrial realm,

we assigned biomes as tundra, forests, grasslands, mediterranean and desert, and arable

systems, while we included all aquatic observations within an aquatic biome within the

aquatic realm. The transitional realm included wetlands in the mediterranean and desert,

temperate and boreal, and tropical ecoregions (Table 1). Finally, the database labelled non-

arable observations as coming from undisturbed or degraded habitats. Habitat degradation

was included as a binary classification within each ecosystem, based on descriptions from

the component studies. During compilation of the database, the contemporary habitat being

assigned to one of the habitat categories listed above, with an additional 'target habitat' was

also added where necessary. Target habitat refers to the ideal natural or semi-natural habitat

of the system, according to the study's authors, interpreted from site descriptions and/or due

to the common practice in the seed bank literature of comparing assemblages across habitat

gradients. If there was no entry for target habitat for a record in the database, we used the

(contemporary) habitat entry and the record was labelled as being undisturbed. When there

was an entry for target habitat, this was used for the habitat definition and the record was

labelled as degraded. Note that the target habitat can have the same category as the

contemporary habitat if it is still – for example – a forest, but in a degraded state. Examples

of degraded habitats include drained wetlands, early-mid successional secondary forests,

abandoned semi-natural grassland or any habitat under restoration, see Auffret et al.[31] for

further details.

Statistical Models

For each response (i.e. species richness and seed density) and biome (e.g. terrestrial

grasslands and transitional wetlands) we fit a univariate mixed effects linear model. This

amounted to 14 models in total (two responses × seven biomes; Table S2), allowing

estimates to vary independently for each biome without well-sampled biomes and realms

(e.g. temperate forests) influencing the estimates of undersampled biomes and realms (e.g.

tundra, aquatic). In general, if there were different ecoregions represented within a biome

then these were treated as categorical fixed effects that interacted with habitat degradation.

If not (e.g. tundra), then degradation was the only categorical fixed effect. In the arable

biome, there was no fixed effect for degradation, because we consider arable to already

represent a degraded system. For each model, intercepts were allowed to vary for every

record, nested within study identity as random effects.

In the species richness models, we quantified richness as a function of sampling area across

the different biomes by fitting total sampled area (m^2; sample grain) as a continuous

predictor, ecoregion as an interacting categorical predictor (where relevant), degradation as

an interacting categorical predictor (a three-way interaction where relevant), and the number

of sites as a continuous covariate as a proxy for study extent. This approach, allowing

drivers of diversity to vary with sampling grain within a single model, has been found to

effectively unify disparate results from previous studies, enabling efficient integration of

heterogeneous biodiversity data[29]. We assumed a Poisson distribution with a log-link

function and we log-transformed and centred the total sampling area and the number of

sites. We set the lower bound of slopes to be zero as informative prior information. This

prevented uncertainty estimates for shallow slopes (aquatic realm) from overlapping with

negative values, which does not logically hold. Seed density models included only

degradation interacting with ecoregion as fixed effects, and assumed a lognormal

distribution. For all model specifications and details see Table S2.

For Bayesian inference and estimates of uncertainty, we fit models using the Hamiltonian

Monte Carlo (HMC) sampler Stan[96] using the R package brms (citations for all software are

given at the end of the Methods). We fit all models with four chains, differing iterations and

differing assumed distributions (Supplementary Information, Model Details Table S2). We

used weakly regularising priors, visually inspected HMC chains for convergence, and

conducted posterior predictive checks to ensure the model represented the distribution of the

data for each (Supplementary Information, Table S2, Figure S1a-g). We selected model

structure and distributions based on this and on model performance (Rhats, sample sizes)

(Supplementary Information, Table S2 Figure S1a-g).

Quantifying scale-dependent richness

Relative differences in local species richness (alpha diversity) between habitats or biomes

do not always match patterns of richness when larger areas are considered (gamma

diversity). This indicates spatial variation in beta diversity (Whittaker's beta = gamma /

alpha)[97]. To investigate this divergence between scales, we first predicted the overall

average species richness at two spatial scales for every biome using the species richness

models to keep richness estimates for every biome consistent. We predicted richness at

0.01 m^2 (alpha diversity), and at a larger scale, gamma diversity at 15 m^2, keeping modelling

parameters constant. These are arbitrary values, but we consider 0.01 m^2 to be a commonly

sampled size in soil seed bank studies, that falls within the interquartile range of seed bank

sampling in the database[31] and has been previously used for pattern prediction of soil seed

bank richness and density[26]. Gamma diversity at 15 m^2 was chosen as the largest scale that

we felt confident predicting across biomes, given the input data and model outputs. In the

main results we report diversity at both alpha and gamma scales assuming each observation

was carried out at one site, as this represents the majority of our values, and acts as our

baseline estimates for diversity.

We estimated the average number of seeds m^{-2} globally for observations within non-

disturbed natural terrestrial, arable, wetland and aquatic realms. To do so, we combined

estimates of the fixed effect (i.e. fitted values) values across all models (i.e. biomes) from

each realm, and predicted the mean seed density m^{-2} across all fitted values from all models

within each broad group. Finally, to look at the joint-relationship between the density of seed

in the soil seed bank and species richness, we predicted species richness at the 1 m^2 scale

used for seed density for direct comparison between the two. We predicted richness at this

third scale 1 m^2, following the same methods at, as for our alpha and gamma diversity. For

all data cleaning, management, wrangling and analyses, we used the R Project for Statistical

Computing version 4.4.0[98] (R Project). To handle data, quantify metrics, summarise results,

and visualise data and estimates we used the brms, tidyverse, bayesplot, and ggplot2

packages[99–102].

659   Data Availability

660   The global seed bank database is published at https://doi.org/10.5878/bvs7-gk47 and is fully

661   described in ref [31].

663   Code Availability

664   The code used for the analysis is available at https://github.com/Global-Seed-Banks/Seed-

665   Banks

Acknowledgements

We thank the Landscape Ecology unit at SLU, and the Biodiversity Synthesis and Physiological Diversity groups at iDiv for feedback on the project idea, data analysis and presentation. AGA is supported by the Swedish research councils VR (2020- 04276) and Formas (2023-02457). EL was supported by the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (FZT 118, 202548816). PK was funded by the European Union (ERC, BEAST, 101044740). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them. VO thanks RSF for financial support (# 25-14-00032). ED was supported by Formas (2023-01507), as were IK (2022-01752, via Biodiversa+ BiodivProtect) and JP (2018-00961 and 2022-01803).

References

1. Grubb, P. J. The Maintenance of Species-Richness in Plant Communities: The Importance of the Regeneration Niche. *Biological Reviews* **52**, 107–145 (1977).

2. Thompson, K. Seeds and Seed Banks. *New Phytologist* **106**, 23–34 (1987).

3. Chesson, P. Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology and Systematics* **31**, 343–366 (2000).

4. HilleRisLambers, J., Clark, J. S. & Lavine, M. Implications of Seed Banking for Recruitment of Southern Appalachian Woody Species. *Ecology* **86**, 85–95 (2005).

5. Honnay, O., Bossuyt, B., Jacquemyn, H., Shimono, A. & Uchiyama, K. Can a seed bank maintain the genetic variation in the above ground plant population? *Oikos* **117**, 1–5 (2008).

6.   Alexander, H. M. *et al.* Metapopulations and metacommunities: combining spatial

and temporal perspectives in plant ecology. *Journal of Ecology* **100**, 88–103 (2012).

7.   Wisnoski, N. I. & Shoemaker, L. G. Seed banks alter metacommunity diversity: The

interactive effects of competition, dispersal and dormancy. *Ecology Letters* **25**, 740–753

(2022).

8.   Saatkamp, A., Poschlod, P. & Venable, D. L. The functional role of soil seed banks in

natural communities. *Seeds: the ecology of regeneration in plant communities* 263–295

(2014) doi:10.1079/9781780641836.0263.

9.   Saatkamp, A., Affre, L., Dutoit, T. & Poschlod, P. The seed bank longevity index

revisited: limited reliability evident from a burial experiment and database analyses. *Ann Bot*

104, 715–724 (2009).

10.   Friedman, J. The evolution of annual and perennial plant life histories: ecological

correlates and genetic mechanisms. *Annual Review of Ecology, Evolution, and Systematics*

51, 461–481 (2020).

11.   DeMalach, N., Kigel, J. & Sternberg, M. The soil seed bank can buffer long-term

compositional changes in annual plant communities. *Journal of Ecology* **109**, 1275–1283

(2021).

12.   Auffret, A. G. *et al.* More warm-adapted species in soil seed banks than in herb layer

plant communities across Europe. *Journal of Ecology* **111**, 1009–1020 (2023).

13.   Tweddle, J. C., Dickie, J. B., Baskin, C. C. & Baskin, J. M. Ecological aspects of seed

desiccation sensitivity. *Journal of Ecology* **91**, 294–304 (2003).

14.   Wyse, S. V. & Dickie, J. B. Predicting the global incidence of seed desiccation

sensitivity. *Journal of Ecology* **105**, 1082–1093 (2017).

15.   Grime, J. P. Evidence for the Existence of Three Primary Strategies in Plants and Its

Relevance to Ecological and Evolutionary Theory. *The American Naturalist* **111**, 1169–1194

(1977).

16.   Klimešová, J. & Herben, T. Clonal and bud bank traits: patterns across temperate

plant communities. *Journal of Vegetation Science* **26**, 243–253 (2015).

17.   Nilsson, M.-C. & Wardle, D. A. Understory vegetation as a forest ecosystem driver:

evidence from the northern Swedish boreal forest. *Frontiers in Ecology and the Environment*

3, 421–428 (2005).

18.   Eriksson, O. & Jakobsson, A. Abundance, distribution and life histories of grassland

plants: a comparative study of 81 species. *Journal of Ecology* **86**, 922–933 (1998).

19.   Ladouceur, E. *et al.* The functional trait spectrum of European temperate grasslands.

Journal of Vegetation Science **30**, 777–788 (2019).

20.   Gaba, S. *et al.* Response and effect traits of arable weeds in agro-ecosystems: a

review of current knowledge. *Weed Research* **57**, 123–147 (2017).

21.   Kuester, A., Conner, J. K., Culley, T. & Baucom, R. S. How weeds emerge: a

taxonomic and trait-based examination using United States data. *New Phytologist* **202**,

1055–1068 (2014).

22.   Kambach, S. *et al.* Climate-trait relationships exhibit strong habitat specificity in plant

communities across Europe. *Nat Commun* **14**, 712 (2023).

23.   Newbold, T., Oppenheimer, P., Etard, A. & Williams, J. J. Tropical and Mediterranean

biodiversity is disproportionately sensitive to land-use and climate change. *Nat Ecol Evol* **4**,

1630–1638 (2020).

24.   Plue, J. *et al.* Buffering effects of soil seed banks on plant community composition in

response to land use and climate. *Global Ecology and Biogeography* **30**, 128–139 (2021).

25.   Vandvik, V., Klanderud, K., Meineri, E., Måren, I. E. & Töpper, J. Seed banks are

biodiversity reservoirs: species–area relationships above versus below ground. *Oikos* **125**,

218–228 (2016).

26.   Yang, X. *et al.* Global patterns of potential future plant diversity hidden in soil seed

banks. *Nat Commun* **12**, 7023 (2021).

27.   Arrhenius, O. Species and area. *Journal of Ecology* **9**, 95–99 (1921).

28.   Drakare, S., Lennon, J. J. & Hillebrand, H. The imprint of the geographical,

evolutionary and ecological context on species-area relationships. *Ecology Letters* **9**, 215–

227 (2006).

29.   Keil, P. & Chase, J. M. Global patterns and drivers of tree diversity integrated across

a continuum of spatial grains. *Nat Ecol Evol* **3**, 390–399 (2019).

30.   Sabatini, F. M. *et al.* Global patterns of vascular plant alpha diversity. *Nat Commun*

13, 4683 (2022).

31.   Auffret, A. G. *et al.* A global database of soil seed bank richness, density, and

abundance. *Ecology* **105**, e4438 (2024).

32.   Olson, D. M. *et al.* Terrestrial Ecoregions of the World: A New Map of Life on Earth A

new global map of terrestrial ecoregions provides an innovative tool for conserving

biodiversity. *BioScience* **51**, 933–938 (2001).

33.   Keith, D. A. *et al.* A function-based typology for Earth's ecosystems. *Nature* **610**,

513–518 (2022).

34.   Eichenberg, D. *et al.* Widespread decline in Central European plant diversity across

six decades. *Global Change Biology* **27**, 1097–1110 (2021).

35.   Blowes, S. A. *et al.* Local biodiversity change reflects interactions among changing

abundance, evenness, and richness. *Ecology* **103**, e3820 (2022).

36.   Plue, J. *et al.* Where does the community start, and where does it end? Including the

seed bank to reassess forest herb layer responses to the environment. *Journal of Vegetation*

Science **28**, 424–435 (2017).

37.   Storch, D., Bohdalková, E. & Okie, J. The more-individuals hypothesis revisited: the

role of community abundance in species richness regulation and the productivity–diversity

relationship. *Ecology Letters* **21**, 920–937 (2018).

38.   Liang, J. *et al.* Co-limitation towards lower latitudes shapes global forest diversity

gradients. *Nat Ecol Evol* **6**, 1423–1437 (2022).

39.   Lloret, F., Escudero, A., Iriondo, J. M., Martínez-Vilalta, J. & Valladares, F. Extreme

climatic events and vegetation: the role of stabilizing processes. *Global Change Biology* **18**,

797–805 (2012).

40.   DeMalach, N., Kigel, J. & Sternberg, M. Contrasting dynamics of seed banks and

standing vegetation of annuals and perennials along a rainfall gradient. *Perspectives in Plant*

Ecology, Evolution and Systematics **58**, 125718 (2023).

41.   Rundel, P. W. *et al.* Mediterranean Biomes: Evolution of Their Vegetation, Floras,

and Climate. *Annual Review of Ecology, Evolution, and Systematics* **47**, 383–407 (2016).

42.   Benedicto-Royuela, J. *et al.* What is the value of biotic seed dispersal in post-fire

forest regeneration? *Conservation Letters* **17**, e12990 (2024).

43.   Skoglund, J. The role of seed banks in vegetation dynamics and restoration of dry

tropical ecosystems. *Journal of Vegetation Science* **3**, 357–360 (1992).

44.   Dalling, J. W., Swaine, M. D. & Garwood, N. C. Dispersal Patterns and Seed Bank

Dynamics of Pioneer Trees in Moist Tropical Forest. *Ecology* **79**, 564–578 (1998).

45.   Baskin, C. C. & Baskin, J. M. Chapter 9 - A Geographical Perspective on

Germination Ecology: Tropical and Subtropical Zones. in *Seeds (Second Edition)* (eds

Baskin, C. C. & Baskin, J. M.) 375–590 (Academic Press, San Diego, 2014).

doi:10.1016/B978-0-12-416677-6.00009-3.

46.   Childs, D. Z., Metcalf, C. J. E. & Rees, M. Evolutionary bet-hedging in the real world:

empirical evidence and challenges revealed by plants. *Proceedings of the Royal Society B:*

Biological Sciences **277**, 3055–3064 (2010).

47.   Snyder, R. E. Multiple risk reduction mechanisms: can dormancy substitute for

dispersal? *Ecology Letters* **9**, 1106–1114 (2006).

48.   Økland, R. H. Persistence of vascular plants in a Norwegian boreal coniferous forest.

Ecography **18**, 3–14 (1995).

49.   Gough, L. Neighbor effects on germination, survival, and growth in two arctic tundra

plant communities. *Ecography* **29**, 44–56 (2006).

50.   Körner, C. *Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems.*

(Springer International Publishing, Cham, 2021). doi:10.1007/978-3-030-59538-8_1.

51.   Moreno-Mateos, D., Meli, P., Vara-Rodríguez, M. I. & Aronson, J. Ecosystem

response to interventions: lessons from restored and created wetland ecosystems. *Journal*

of Applied Ecology **52**, 1528–1537 (2015).

52.   Sullivan, C. A. *et al.* Modelling semi-natural habitat area on lowland farms in western

Ireland. *Biological Conservation* **144**, 1089–1099 (2011).

53.   Vellend, M. Habitat loss inhibits recovery of plant diversity as forests regrow. *Ecology*

84, 1158–1164 (2003).

54.   Ikeda, F. S., Mitja, D., Vilela, L. & Carmona, R. Banco de sementes no solo em

sistemas de cultivo lavoura-pastagem. *Pesq. agropec. bras.* **42**, 1545–1551 (2007).

55.   Rotchés-Ribalta, R., Armengot, L., Mäder, P., Mayer, J. & Sans, F. X. Long-Term

Management Affects the Community Composition of Arable Soil Seedbanks. *Weed Science*

65, 73–82 (2017).

56.   Schwartz, L. M. *et al.* Seedbank and Field Emergence of Weeds in Glyphosate-

Resistant Cropping Systems in the United States. *Weed Science* **63**, 425–439 (2015).

57.   Daïnou, K. *et al.* Soil seed bank characteristics in Cameroonian rainforests and

implications for post-logging forest recovery. *Ecological Engineering* **37**, 1499–1506 (2011).

58.   Dupuy, J. M. & Chazdon, R. L. Long-Term Effects of Forest Regrowth and Selective

Logging on the Seed Bank of Tropical Forests in NE Costa Rica. *Biotropica* **30**, 223–237

(1998).

59.   Plue, J. *et al.* Seed banks of temperate deciduous forests during secondary

succession. *Journal of Vegetation Science* **21**, 965–978 (2010).

60.   Daru, B. H. *et al.* Widespread homogenization of plant communities in the

Anthropocene. *Nat Commun* **12**, 6983 (2021).

61.   Jandt, U. *et al.* More losses than gains during one century of plant biodiversity

change in Germany. *Nature* **611**, 512–518 (2022).

62.   Vellend, M. *et al.* Global meta-analysis reveals no net change in local-scale plant

biodiversity over time. *PNAS* 201312779 (2013) doi:10.1073/pnas.1312779110.

63.   Atkinson, J. *et al.* Terrestrial ecosystem restoration increases biodiversity and

reduces its variability, but not to reference levels: A global meta-analysis. *Ecology Letters*

25, 1725–1737 (2022).

64.   Ladouceur, E. *et al.* The recovery of plant community composition following passive

restoration across spatial scales. *Journal of Ecology* **111**, 814–829 (2023).

65.   Boakes, E. H. *et al.* Distorted Views of Biodiversity: Spatial and Temporal Bias in

Species Occurrence Data. *PLoS Biol* **8**, e1000385 (2010).

66.   Hughes, A. C. *et al.* Sampling biases shape our view of the natural world. *Ecography*

44, 1259–1269 (2021).

67.   Nuñez, M. A. & Amano, T. Monolingual searches can limit and bias results in global

literature reviews. *Nat Ecol Evol* **5**, 264–264 (2021).

68.   Boyd, R. J., Powney, G. D. & Pescott, O. L. We need to talk about nonprobability

samples. *Trends in Ecology & Evolution* **38**, 521–531 (2023).

69.   Krauss, J. *et al.* Habitat fragmentation causes immediate and time-delayed

biodiversity loss at different trophic levels. *Ecology Letters* **13**, 597–605 (2010).

70.   Filazzola, A. & Cahill Jr, J. F. Replication in field ecology: Identifying challenges and

proposing solutions. *Methods in Ecology and Evolution* **12**, 1780–1792 (2021).

71.   Bigwood, D. W. & Inouye, D. W. Spatial Pattern Analysis of Seed Banks: An

Improved Method and Optimized Sampling. *Ecology* **69**, 497–507 (1988).

72.   Plue, J. & Hermy, M. Consistent seed bank spatial structure across semi-natural

habitats determines plot sampling. *Journal of Vegetation Science* **23**, 505–516 (2012).

73.   Kuczynski, L., Ontiveros, V. J. & Hillebrand, H. Biodiversity time series are biased

towards increasing species richness in changing environments. *Nat Ecol Evol* **7**, 994–1001

(2023).

74.   Auffret, A. G. & Cousins, S. A. O. Past and present management influences the seed

bank and seed rain in a rural landscape mosaic. *Journal of Applied Ecology* **48**, 1278–1285

(2011).

75.   Karlík, P. & Poschlod, P. Soil seed-bank composition reveals the land-use history of

calcareous grasslands. *Acta Oecologica* **58**, 22–34 (2014).

76.   Gioria, M., Pyšek, P., Baskin, C. C. & Carta, A. Phylogenetic relatedness mediates

persistence and density of soil seed banks. *Journal of Ecology* **108**, 2121–2131 (2020).

77.   Chen, S.-C., Poschlod, P., Antonelli, A., Liu, U. & Dickie, J. B. Trade-off between

seed dispersal in space and time. *Ecology Letters* **23**, 1635–1642 (2020).

78.   Gioria, M. *et al.* Persistent soil seed banks promote naturalisation and invasiveness

in flowering plants. *Ecology Letters* **24**, 1655–1667 (2021).

79.   Westcott, K., Whillans, T. H. & Fox, M. G. Viability and abundance of seeds of

submerged macrophytes in the sediment of disturbed and reference shoreline marshes in

Lake Ontario. *Can. J. Bot.* **75**, 451–456 (1997).

80.   Zaghloul, M. S. Diversity in soil seed bank of Sinai and implications for conservation

and restoration. *African Journal of Environmental Science and Technology* **2**, 172–184

(2008).

81.   Kirmer, A. & Mahn, E.-G. Spontaneous and initiated succession on unvegetated

slopes in the abandoned lignite-mining area of Goitsche, Germany. *Applied Vegetation*

Science **4**, 19–27 (2001).

82.   Sarneel, J. M., Beltman, B., Buijze, A., Groen, R. & Soons, M. B. The role of wind in

the dispersal of floating seeds in slow-flowing or stagnant water bodies. *Journal of*

Vegetation Science **25**, 262–274 (2014).

83.   Malfasi, F., Convey, P., Zaccara, S. & Cannone, N. Establishment and eradication of

an alien plant species in Antarctica: Poa annua at Signy Island. *Biodivers Conserv* **29**, 173–

186 (2020).

84.   Gibson, D. J. *et al.* Benchmark study on glyphosate-resistant cropping systems in the

United States. Part 7: Effects of weed management strategy (grower practices versus

academic recommendations) on the weed soil seedbank over 6 years. *Pest Management*

Science **72**, 692–700 (2016).

85.   Boyes, L. J., Gunton, R. M., Griffiths, M. E. & Lawes, M. J. Causes of arrested

succession in coastal dune forest. *Plant Ecol* **212**, 21–32 (2011).

86.   Burmeier, S., Eckstein, R. L., Otte, A. & Donath, T. W. Desiccation cracks act as

natural seed traps in flood-meadow systems. *Plant Soil* **333**, 351–364 (2010).

87.   Baptista, T. L. & Shumway, S. W. A Comparison of the Seed Banks of Sand Dunes

with Different Disturbance Histories on Cape Cod National Seashore. *Rhodora* **100**, 298–

313 (1998).

88.   Graham, L. L. B. & Page, S. E. A limited seed bank in both natural and degraded

tropical peat swamp forest: the implications for restoration. *Mires and Peat* 1–13 (2018)

doi:10.19189/MaP.2017.OMB.302.

89.   Hawes, C., Squire, G. R., Hallett, P. D., Watson, C. A. & Young, M. Arable plant

communities as indicators of farming practice. *Agriculture, Ecosystems & Environment* **138**,

17–26 (2010).

90.   He, Y., Gao, P. & Qiang, S. An investigation of weed seed banks reveals similar

potential weed community diversity among three different farmland types in Anhui Province,

China. *Journal of Integrative Agriculture* **18**, 927–937 (2019).

91.   Hausman, C. E., Fraser, L. H., Kershner, M. W. & de Szalay, F. A. Plant community

establishment in a restored wetland: Effects of soil removal. *Applied Vegetation Science* **10**,

383–390 (2007).

92.   Holland, J. M., Smith, B. M., Southway, S. E., Birkett, T. C. & Aebischer, N. J. The

effect of crop, cultivation and seed addition for birds on surface weed seed densities in

arable crops during winter. *Weed Research* **48**, 503–511 (2008).

93.   Salonen, J., Hyvönen, T. & Jalli, H. Composition of weed flora in spring cereals in

Finland - a fourth survey. https://jukuri.luke.fi/handle/10024/478541 (2011).

94.   Thompson, K. Small-Scale Heterogeneity in the Seed Bank of An Acidic Grassland.

Journal of Ecology **74**, 733–738 (1986).

95.   Matlack, G. R. & Good, R. E. Spatial Heterogeneity in the Soil Seed Bank of a

Mature Coastal Plain Forest. *Bulletin of the Torrey Botanical Club* **117**, 143–152 (1990).

96.   Carpenter, B. *et al.* Stan: A Probabilistic Programming Language. *Journal of*

Statistical Software **76**, 1–32 (2017).

97.   Whittaker, R. H. Evolution and Measurement of Species Diversity. *Taxon* **21**, 213–

251 (1972).

98.   R Core Team. *R: A Language and Environment for Statistical Computing.* (R

Foundation for Statistical Computing, Vienna, 2021).

99.   Bürkner, P.-C. Advanced Bayesian Multilevel Modeling with the R Package brms.

The R Journal **10**, 395–411 (2018).

100.   Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. Visualization in

Bayesian Workflow. *Journal of the Royal Statistical Society Series A: Statistics in Society*

182, 389–402 (2019).

101.   Wickham, H. *Ggplot2 - Elegant Graphics for Data Analysis.* (Springer, New York,

2009).

102.   Wickham, H. *et al.* Welcome to the Tidyverse. *Journal of Open Source Software* **4**,

1686 (2019).

924
925