Divergent patterns of richness and density in the global soil seed bank

Alistair G. Auffret [1⍰*], Emma Ladouceur [2,3,4,5⍰*], Natalie S. Haussmann[6], Petr Keil [7], Eirini Daouti [1,8], Tatiana G. Elumeeva [9], Ineta Kačergytė [1], Jonas Knape [1], Dorota Kotowska[10,11], Matthew Low [1], Vladimir G. Onipchenko[9], Matthieu Paquet [1,12], Diana Rubene[8, 13], Jan Plue[14]

Author Affiliations

[1] Department of Ecology, Swedish University of Agricultural Sciences, 75 007 Uppsala, Sweden

[2] Department of Biology, University of Prince Edward Island, PE, Canada

[3] Canadian Centre for Climate Change and Adaptation, University of Prince Edward Island, St. Peter's Bay, PE, C0A 2A0, Canada

[4] School of Climate Change and Adaptation, University of Prince Edward Island, Charlottetown, PE, C1A 4P3, Canada

[5] German Centre for Integrative Biodiversity Research (iDiv) Leipzig-Halle-Jena, Puschstraße 4, 04103 Leipzig, Germany

[6] Department of Geography, Geoinformatics and Meteorology, University of Pretoria, Lynnwood Rd, Hatfield 0002, South Africa

[7] Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, 165 00 Praha-Suchdol, Czech Republic

[8] Department of Crop Production Ecology, Swedish University of Agricultural Sciences, 75 007 Uppsala, Sweden

[9] Moscow Lomonosov State University, Moscow, 119234, Russia

[10] Institute of Nature Conservation, Polish Academy of Sciences, Kraków, Poland

[11] 'Lendület' Landscape and Conservation Ecology Group, Institute of Ecology and Botany, Centre for Ecological Research, Vácrátót, Hungary

[12] Institute of Mathematics of Bordeaux, University of Bordeaux, CNRS, Bordeaux INP, Talence, France

[13] Greensway AB, Gerda Nilssons väg 2, 756 51 Uppsala, Sweden

[14] Swedish Biodiversity Centre, Swedish University of Agricultural Sciences, 75 007 Uppsala, Sweden

⍰Alistair G. Auffret and Emma Ladouceur should be considered joint first authors
* Corresponding authors alistair.auffret@slu.se, erladouceur@upei.ca

Abstract

Soil seed banks are an important component of plant population and community dynamics. As a functional strategy that has evolved in response to abiotic disturbance regimes, they are likely to play a key role in determining biodiversity patterns across the world's biomes and habitats. At the same time, patterns in soil seed banks are likely to reflect scale-dependent patterns observed in above-ground vegetation. Here, we investigate components of diversity in the soil seed bank across global biomes and ecosystems, accounting for sampled soil area and geographic extent. We found that patterns of species richness in the soil seed bank diverged from patterns of density, although there were high levels of uncertainty in the results, especially at smaller spatial scales. Importantly, habitat degradation consistently led to lower richness, potentially limiting future ecosystem recovery. Among ecosystems, Mediterranean and tropical regions were broadly found to have high species richness, but seed density m^{-2} in the soil was highest in wetlands and arable systems. Lower densities were found generally in both high-diversity tropical biomes that are characterised by short-lived seeds, and low-diversity boreal and tundra biomes with more stable established vegetation. We show how different components of the soil seed bank depend on the interplay between local environmental conditions and global biomes, with implications for understanding biodiversity at the global scale.

Introduction

In an era of global environmental change, the persistence of populations in newly-unsuitable conditions and the establishment of new populations following disturbance are key in determining changes in biodiversity over time and in space. For vascular plants, the soil

seed bank can play a vital role in both processes. Soil seed banks are the communities of

persistent seeds being naturally stored in the soil that contribute to vegetation regeneration

after disturbance (Grubb 1977; Thompson 1987). Plant species that form seed banks do so

as part of a life-history strategy that allows for the buffering of temporal environmental

heterogeneity (Chesson 2000; HilleRisLambers *et al.* 2005; Honnay *et al.* 2008), and as

such, seed banks are an important component for understanding plant (meta)population and

community dynamics (Alexander *et al.* 2012; Wisnoski & Shoemaker 2022). By reflecting

previous habitat conditions following land-use change (Auffret & Cousins 2011; Karlík &

Poschlod 2014), seed banks are often studied in relation to the potential for ecological

restoration of target communities (Bossuyt & Honnay 2008; Török *et al.* 2018). As the

climate warms, seed banks are expected to contribute to the maintenance of species

richness in the above-ground vegetation (Plue *et al.* 2021). However, the abundance of

seeds in the seed bank is predicted to decline with increasing soil temperatures due to a

combination of increased rates of germination and seed degradation (Ooi *et al.* 2009).

Despite the potential advantage of producing seeds that persist in the soil, not all species

contribute to soil seed banks. Instead, the ability to form seed banks is a functional strategy

for plants to (re)establish and maintain populations, which has evolved in response to both

predictable and unpredictable disturbance regimes (Saatkamp *et al.* 2014). Such

disturbances are not randomly distributed in space, and as such plant communities in

different bioclimatic environments can be expected to rely on seed banking to different

extents. For example, in arid systems dominated by drought, it is advantageous to have an

annual life history and form persistent seed banks to buffer annual changes in water

availability (DeMalach *et al.* 2021; Friedman 2020). A similar pattern exists in temperate

Europe, where species with warmer distributions have been found to have a higher seed

bank persistence than more northerly species (Auffret *et al.* 2023). In tropical systems,

where climatic conditions are broadly stable and exposure to freezing temperatures and

drought conditions is historically rare, relatively few species produce seeds that are able to

survive dessication – a trait associated with long-term persistence – which may affect the

size and composition of soil seed banks (Tweddle *et al.* 2003; Wyse & Dickie 2017).

While it can be interesting to broadly compare plant communities from different world

regions, ecological assemblages can differ considerably among habitats within the same

biome. The frequency of disturbance within a habitat is generally mirrored by the strategies

of the resulting plant communities (Grime 1977). For example, in temperate and boreal

regions, forests are dominated by long-lived trees, with the understorey often covered by

species that largely rely on vegetative reproduction (Klimešová & Herben 2015; Nilsson &

Wardle 2005). Species-rich grazed grasslands in the same regions are characterised by

regular, low-intensity and spatially-heterogeneous disturbances, and as such harbour plants

with a wide range of regeneration strategies (Eriksson & Jakobsson 1998; Ladouceur *et al.*

2019). Intensively-used arable lands on the other hand are well-known to support abundant

communities of ruderal species with annual life-histories and high seed production (Gaba *et*

al. 2017; Kuester *et al.* 2014). As such, habitat is an important component in understanding

how different functional traits are represented in different regions, while differences between

broad habitat types can differ in different world regions (Kambach *et al.* 2023). Moreover,

communities in the same broad habitat categories can differ in their responses to

environmental degradation in different biomes (Newbold *et al.* 2020). Therefore, when

studying plant communities at global scales, it is important to consider the potential roles of

different habitats and environmental degradation driving any variation within or between

biomes. This could be especially relevant in the case of soil seed banks, which exist partially

as a strategy to respond to disturbance.

Studying patterns of soil seed bank diversity across large spatial scales and geographic

gradients can be useful for understanding the biogeography and macroecology of their role

in biodiversity maintenance and recovery (Plue *et al.* 2021; Vandvik *et al.* 2016; Yang *et al.*

2021). However, biodiversity is known to be scale-dependent, and comparisons of soil seed

banks across biomes and habitats are therefore likely to be influenced by the scale at which

diversity is estimated. It has been established that the number of species in soil seed banks

increases with area (Plue *et al.* 2021; Vandvik *et al.* 2016), but while it is well-known that this

accumulation of species varies according to habitat and biome in the established vegetation

as well as in other taxa (Arrhenius 1921; Drakare *et al.* 2006; Keil & Chase 2019; Sabatini *et*

al. 2022), such patterns are still largely unknown in the soil seed bank, at least not at spatial

scales that span multiple biomes. Integrating a scale-explicit approach to understanding

diversity in the soil seed bank across habitats, within and between biomes and in response

to habitat degradation is an important step for plant community and macroecology, allowing

critical insights for both broad scale patterns of biodiversity and the context-dependent

interpretations of local dynamics.

In this paper, we use a global database to examine spatial patterns of soil seed banks in

terms of species richness and seed density. Seed bank studies are usually (through

necessity) small-scale comparisons across environmental gradients or investigations into the

effects of environmental degradation. Here, we harness 1442 studies from all seven of the

world's continents, with the large number of studies and variation in study grain allowing us

to address questions not usually possible within the same global synthesis. Specifically, we

analyse seed bank biodiversity at two spatial resolutions, comparing diversity within and

between broad habitat categories and biomes, and also analysing the effects of habitat

degradation. Our study contributes to a broader understanding of seed bank ecology, in

terms of responses to environmental change, habitat restoration, and future investigations of

the soil seed bank.

Methods

There are a number of reasons why soil seed banks lend themselves to the study of global patterns and scale-dependence of biodiversity. First, the global distribution of seed bank ecologists means that empirical studies of the soil seed bank are common and far-reaching. Second, they are sampled in a broadly consistent way, through the extraction of a known amount of soil, with species identified directly from seeds or through germination in the greenhouse. Third, studies have used a relatively wide range of sampling efforts or sampling grains (area of ground covered), including variation in both the size of an individual sample and the number of samples taken. Taken together, these advantages show real potential for giving important insights into how biodiversity is distributed in space, and how vegetation regeneration responses to global change are spatially scaled.

The global seed bank database

To study the effects of spatial scale, habitat type and quality on soil seed banks, we used a global database of seed bank richness, abundance and density (Auffret *et al.* 2024). Briefly, the database is based on a Web of Science search using the search term "seed bank*" OR seedbank*, including papers included in the Web of Science until the end of 2019 (a few papers returned by the search were subsequently published in journals in 2020). All resulting papers were checked for being both community seed bank studies (i.e., taking soil samples and counting and identifying all seeds, rather than focussing on one or more specific species), and containing information about species richness (number of species), seed abundance (number of seeds), and/or seed density (number of seeds per unit area or volume) in the seed bank. Most of the papers were in English, but data were also extracted from papers published in, for example, Spanish and Portuguese (often South American

studies), as well as French and German. Due to a large data gap, the database was

supplemented by an additional database and literature search featuring studies published in

Russian and carried out in a number of former Soviet states. In total, this resulted in 7224

unique studies to be checked, and 1442 studies from which records could be extracted or

obtained. A record refers to a row in the database, and represents a known number of soil

samples with specified (by the study authors) geographical coordinates. Depending on the

detail in which study location and design, and seed bank results were reported in a specific

study, a record can represent one or more locations or environments. Each record can also

provide one or more commonly recorded seed bank metrics (mainly total species, total

number of seeds, seeds m^{-2} density), which we term observations. A study can provide

multiple observations by reporting more than one metric, and/or by (for example) reporting

metrics across several distinct records (i.e. different locations or environments). In total, the

1442 studies provided 8087 observations across 3096 records (Figure 1, Table 1, S1). The

database includes soil seed bank data from 94 countries and all seven continents. Full

details regarding which data were extracted and how are available in Auffret *et al*. (2024).

Below, we describe only the variables in the database that were used for this analysis.

182
183
184

Figure 1: The global record of the soil seed bank. Points indicate the location of each
record collected from the literature (n=3096; some records have the same location).
Colours indicate unique ecosystem realm, biome, and biome functional subtype
categories for each record. Biomes Include: a) Tundra, b) Forests, c) Grasslands &
Savannas, d) Mediterranean & Desert (a-d - Natural Terrestrial Realm), e) Arable (e
- Terrestrial Intensive Land Use Realm), f) Wetlands & Flooded Grasslands (f -
Marine-Terrestrial-Freshwater Transitional Realms), g) Aquatic (g - Marine &
Freshwater Realms). Each Biome is divided into biome subtypes ranging from
boreal-tropical depending on context. Shapes to aid in point differentiation indicate
that forests are diamonds, arable are squares, wetland are open circles, aquatic are
triangles, and all other systems are filled circles.

Modelled variables

From the global seed bank database, we extracted the two most common metrics from the

soil seed bank literature as response variables: [1] the total number of species recorded in a

reported area soil seed bank sampled; [2] the density of seeds m^{-2} in the soil. Soil seed bank

studies often involve the collection of multiple samples of soil with a soil corer. These

samples are then usually pooled, in order to account for the spatial patchiness of species

and seeds in the soil (Matlack & Good 1990; Plue & Hermy 2012; Thompson 1986).

Samples can also be collected (and pooled) at several sites in order to capture variation

across different locations representing the same habitat or treatment category, often

described as 'replicates' within a study. We assume that authors of component papers chose

a combination of soil samples and sites that they expected would characterise the soil seed

bank in their study system. In our analysis, we used the total sampled area in m^2 or the

sample grain as our measure of sampling scale, calculated as the size of the individual soil

samples multiplied by the total number of samples taken for a particular observation. We

also included the number of sites contributing to each observation as a measure of

geographical extent. While this is an imperfect measure, we found it to be a parsimonious

covariate, as well as the only such metric that was available in all records. More than half of

our observations represent samples collected within a single site (n = 4763).

In order to categorise the global seed bank observations to an ecosystem in a systematic

way (McGill & Miller, 2022), we used a combination of authors self-declared habitat

descriptions coming from the global seed bank database (aquatic, open grassy systems,

forest, arable, or wetland; Auffret et al. 2024), the WWF terrestrial ecoregions or biomes

based on an observation's geographic coordinates (Olson *et al.* 2001), as well as realms in

the fashion of the function-based typology for the Earth's ecosystems (Keith et al. 2022).

Observations with habitat defined as arable or aquatic were placed in an arable and aquatic

'biome', respectively. Habitat and biome were used to categorise all observations into one of

15 biome subtypes (Table 1), with the aim to maintain ecologically-relevant distinctions of

ecosystems (the combination of biome and biome subtype) while also including a

reasonable number of observations within each group. Each ecosystem was then assigned

into either the aquatic, terrestrial or transitional realm. Within the terrestrial realm, we

assigned biomes as tundra, forests, grasslands, mediterranean and desert, and arable

systems, while we included all aquatic observations within an aquatic biome within the

aquatic realm. The transitional realm included wetlands in the mediterranean and desert,

temperate and boreal, and tropical biomes (Table 1). Finally, the database labelled non-

arable observations as coming from undisturbed or degraded habitats, based on

descriptions from the individual studies. Examples of degraded habitats include drained

wetlands, early-mid successional secondary forests, abandoned semi-natural grassland or

any habitat under restoration. This was included as a binary classification within each

ecosystem.

Table 1: Realm, Biome and Biome subtype categories assigned to each observation. The combination of biome and biome subtype
is what we refer to as an ecosystem. 'Degraded' indicates whether the area samples were taken from was considered to be
degraded by authors of the original study. The number of observations collected for each Biome subtype, and degraded status
listed for species richness, and seed density m^{-2}.

Realm	Biome	Biome subtype	Degraded	N species	N seeds	N density
Terrestrial	Tundra	Tundra	0	43	42	40
Terrestrial	Tundra	Tundra	1	3	3	3
Terrestrial	Forest	Boreal	0	26	22	22
Terrestrial	Forest	Boreal	1	11	11	11
Terrestrial	Forest	Temperate	0	266	268	285
Terrestrial	Forest	Temperate	1	160	139	144
Terrestrial	Forest	Tropical	0	155	151	152
Terrestrial	Forest	Tropical	1	136	136	135
Terrestrial	Grassland	Temperate and Boreal	0	475	451	456
Terrestrial	Grassland	Temperate and Boreal	1	340	263	265
Terrestrial	Grassland	Tropical	0	51	45	45
Terrestrial	Grassland	Tropical	1	33	32	31
Terrestrial	Mediterranean and Desert	Deserts and Xeric Shrublands	0	107	82	83
Terrestrial	Mediterranean and Desert	Deserts and Xeric Shrublands	1	33	29	35
Terrestrial	Mediterranean and Desert	Mediterranean Forests, Woodlands and Scrub	0	129	123	121
Terrestrial	Mediterranean and Desert	Mediterranean Forests, Woodlands and Scrub	1	69	61	61
Terrestrial	Arable	Mediterranean and Desert	0	48	43	43

Terrestrial	Arable	Temperate and Boreal	0	159	140	154
Terrestrial	Arable	Tropical	0	65	58	59
Transitional	Wetland	Mediterranean and Desert	0	40	23	23
Transitional	Wetland	Mediterranean and Desert	1	9	7	7
Transitional	Wetland	Temperate and Boreal	0	276	250	249
Transitional	Wetland	Temperate and Boreal	1	103	89	90
Transitional	Wetland	Tropical	0	50	42	42
Transitional	Wetland	Tropical	1	11	10	10
Aquatic	Aquatic	Aquatic	0	74	65	64
Total number of observations				2872	2585	2630

Statistical Models

For each response (i.e. species richness and seed density) and biome (e.g. terrestrial grasslands and transitional wetlands) we fit a univariate mixed effects linear model. This amounted to 14 models in total (two responses × seven biomes; Table S2). In general, if there were different biome subtypes within a biome then these were treated as categorical fixed effects that interacted with habitat degradation. If not (e.g. tundra), then degradation was the only categorical fixed effect. In the arable biome, there was no fixed effect for degradation, because we consider arable to already represent a degraded system. For each model, intercepts were allowed to vary for every record, nested within study identity as random effects.

In the species richness models, we quantified richness as a function of sampling area across the different biomes by fitting total sampled area (m^2) (sample grain) as a continuous predictor, biome subtype as an interacting categorical predictor (where relevant), degradation as an interacting categorical predictor (a three-way interaction where relevant), and the number of sites as a continuous covariate as a proxy for study extent. This approach, allowing drivers of diversity to vary with sampling grain within a single model, has been found to effectively unify disparate results from previous studies, enabling efficient integration of heterogeneous biodiversity data (Keil & Chase 2019). We assumed a Poisson distribution with a log-link function and we log-transformed and centred the total sampling area and the number of sites. We set the lower bound of slopes to be zero as informative prior information. This prevented uncertainty estimates for shallow slopes (aquatic realm) from overlapping with negative values, which does not logically hold. Seed density models included only degradation interacting with biome subtype as fixed effects, and assume a lognormal distribution. For all model specifications and details see Table S2.

For Bayesian inference and estimates of uncertainty, we fit models using the Hamiltonian

Monte Carlo (HMC) sampler Stan (Carpenter *et al.* 2017) in brms (Burkner 2018), R Project

for Statistical Computing version 4.4.0 (R Project) (see more detail below). We fit all models

with four chains, differing iterations and differing assumed distributions (Supplementary

Information, Model Details Table S2). We used weakly regularising priors, visually inspected

HMC chains for convergence, and conducted posterior predictive checks to ensure the

model represented the distribution of the data for each (Supplementary Information, Table

S2, Figure S1a-g). We selected model structure and distributions based on this and on

model performance (Rhats, sample sizes) (Supplementary Information, Table S2 Figure

S1a-g).

Quantifying scale-dependent richness

Relative differences in local species richness (alpha diversity) between habitats or biomes

do not always match patterns of richness when larger areas are considered (gamma

diversity). This indicates spatial variation in beta diversity (Whittaker's beta = gamma / alpha)

(Whittaker 1972). To investigate this divergence between scales, we first predicted the

overall average species richness at two spatial scales for every biome using the species

richness models to keep richness estimates for every biome consistent. We predicted

richness at 0.01 m^2 (alpha diversity), and at a larger scale, gamma diversity at 15 m^2,

keeping modelling parameters constant. These are arbitrary values, but we consider 0.01 m^2

to be a commonly sampled size in soil seed bank studies, that falls within the interquartile

range of seed bank sampling in the database (Auffret et al. 2024). Additionally, this is a

common soil sample size in soil seed bank studies, that has been previously used for pattern

prediction of soil seed bank richness and density (Yang *et al.* 2021). Gamma diversity at 15

m^2 was chosen as the largest scale that we felt confident predicting across biomes, given

the input data and model outputs. In the main results we report diversity at both alpha and

gamma scales assuming each observation was carried out at one site, as this represents the

majority of our values, and acts as our baseline estimates for diversity.

We estimated the average number of seeds m^{-2} globally for observations within non-

disturbed natural terrestrial, arable, wetland and aquatic realms. To do so, we combined

estimates of the fixed effect (i.e. fitted values) values across all models (i.e. biomes) from

each realm, and predicted the mean seed density m^{-2} across all fitted values from all models

within each broad group. Finally, to look at the joint-relationship between the density of seed

in the soil seed bank and species richness, we predicted species richness at the 1 m^2 scale

used for seed density, following the same methods for this prediction at $1m^2$, as for our alpha

and gamma diversity. For all data cleaning, management, wrangling and analyses, we used

the R Project for Statistical Computing version 4.4.0 (R Project). To handle data, quantify

metrics, summarise results, and visualise data and estimates we used the brms, tidyverse,

bayesplot, and ggplot2 packages (Bürkner 2018; Gabry *et al.* 2019; Wickham 2009;

Wickham *et al.* 2019).

312 Results

313 Soil seed banks of the world

The total number of seeds counted across the 1185 studies and 2585 observations of seed

density was 35 932 626. Five studies (or 0.2% of all observations) found no seeds within

samples. This included an investigation of aquatic systems in Canada (Westcott *et al.* 1997),

deserts and xeric shrublands in Egypt (Zaghloul 2008), temperate forests in Germany and

the Netherlands (Kirmer & Mahn 2001; Sarneel *et al.* 2014), and tundra in Antarctica (Malfasi

et al. 2020). Ten studies observed only one species in their seed bank samples, while the

maximum was 246 species in 88 m^2 of soil across 156 arable sites in the midwest of the

United States (Gibson *et al.* 2016). The minimum (non-zero) observed density of seeds in

the seed bank was 3 seeds m^{-2} in a tropical forest in South Africa (Boyes *et al.* 2011), with

the maximum being 501 119 seeds m^{-2} observed in a study from temperate forests in

Germany (Burmeier *et al.* 2010). The minimum number of seeds greater than zero recorded

in a soil seed bank observation was two seeds in two studies in temperate forests in the

United States, and tropical and subtropical forests in Indonesia with 0.67 seeds m^{-2} and 0.16

m^{-2} of soil sampled, respectively (Baptista & Shumway 1998; Graham & Page 2018), with

the maximum being 5 907 132 seeds in samples taken in arable systems in 462 m^2 soil from

924 samples across 22 sites in the United Kingdom (Hawes *et al.* 2010).

Naturally, such raw maximum and minimum values are at least partly an outcome of each

investigation's scientific focus and study design. Indeed, there was a large variation in

sampling effort across records. The minimum total number of samples was one sample,

used in 19 studies, with the maximum being 114 000 samples taken in arable systems

representing 224 m^2 of soil sampled across 228 sites in China (He *et al.* 2019). The

minimum total sampled area was 0.0015 m^2 (5 sites, 15 total samples) in temperate

broadleaf and mixed forests from the United States (Hausman *et al.* 2007), with the

maximum being 816 m^2 (20 sites, 20 400 samples) in another large arable study in the UK

(Holland *et al.* 2008). The minimum number of sites was one, representing the majority of

our samples (1832 observations from 692 studies), with the maximum being 595 sites (total

sampled area of 59.5 m^2) in arable fields across Finland (Salonen *et al.* 2011).

Local and global patterns in the soil seed bank

We found that patterns of species richness in the soil seed bank diverged from patterns of

density. That is, high relative abundance of seeds in the soil seed bank does not necessarily

translate to a large number of species. We also find that habitat degradation appears to have

impacted the seed bank across the world.In such a large and heterogeneous dataset, there

was generally a large variation in seed bank richness and density within and between

ecosystems. Here, we describe noteworthy results in terms of overlap in 50% and 90%

credible intervals (hereafter CIs) and consistency in the relative values of model estimates

that are rounded to whole numbers (i.e. whole species or whole seeds m^{-2}). All model results

and further supplementary information can be found in Tables S1-S4, and Figures S2a-g.

Richness and density across ecosystems and biomes

Using the species richness models to predict average soil seed bank diversity for each

ecosystem, we found that species richness generally increased from the 0.01m^2 alpha scale

to the 15 m^2 gamma scale, with model estimates always higher at the gamma scale and

50% credible intervals not overlapping in all undisturbed and 80% of degraded (including

arable) systems. Relative values of predicted richness varied according to scale, with one

example being that boreal and temperate grasslands had a higher predicted richness than

tropical forests at the alpha scale (but with overlapping CIs), while tropical forests had higher

richness at the gamma scale (no overlap of 50% CIs). Generally, differences in estimated

richness across ecosystems were more clear at the gamma scale, so we place most focus

there. In undisturbed systems, mediterranean forests and shrublands were predicted to have

the highest richness compared to all eleven other undisturbed ecosystems (no overlap of

50% CIs, overlap with six undisturbed ecosystems at 90% CIs; Figure 2, Table S4). Tundra

and aquatic systems generally had the lowest species richness, each having a lower

estimate than the other ten undisturbed ecosystems, and lower than eight of the other

ecosystems (50% CIs). Undisturbed tropical ecosystems were estimated to be broadly more

species-rich than their equivalent temperate and boreal ecosystems. Estimated values were

higher for tropical forests than both boreal and temperate forests, for tropical grasslands

compared to temperate and boreal grasslands and tropical arable compared to temperate

and boreal arable, while tropical wetlands had similar estimated richness as temperate and

boreal wetlands. Despite this consistency, estimates were only clearly different in forest

ecosystems (50% CIs), although tropical wetlands were more species-rich than

mediterranean and desert wetlands.

The average estimated number of seeds m^{-2} in the soil seed bank was 5 737 m^{-2} (median 6

129) for natural terrestrial and wetland ecosystems, 9 434 m^{-2} (median 11 001) for arable

systems, 8 124 m^{-2} (median 9 562) for aquatic systems (Figure 3, Table S5). Wetlands

exhibited high seed densities, often higher than other ecosystems within their equivalent

biome subtype (90% CIs). For example, tropical wetlands had higher seed densities than

tropical forests, and temperate and boreal wetlands had higher seed densities than forests

and grasslands in the same regions. Within the terrestrial realm, undisturbed temperate and

boreal grasslands had the highest predicted seed density, higher than boreal forests and

temperate forests (90% CIs).

The effects of habitat degradation

In all ecosystems, habitat degradation always resulted in a lower estimated species richness

at the 15 m^2 scale in terms of predicted values, but again, despite this consistency, the

difference was only clear in mediterranean forests, woodlands and scrub according to the

50% credible intervals. At the 0.01 m^2 scale, the predicted values were often the same in

undisturbed and degraded ecosystems. Compared to species richness, habitat degradation

had much more variable effects on seed density m^{-2} (Figure 3, Figure 4). Broadly speaking,

degradation was more likely to result in lower predicted seed densities compared to

undisturbed habitats in open ecosystems, with clear differences (50% CIs) for tropical

grasslands, mediterranean and desert wetlands and aquatic ecosystems. On the other hand,

seed density estimates were generally higher with degradation in forest ecosystems, with

clear differences in both temperate and tropical forests (50% CIs). No ecosystems exhibited

clear differences in richness or density with degradation according to the 90% CIs.

Divergent patterns of richness and density

Despite relatively high species richness, seed bank density was generally low in tropical

systems. Tropical forests had the lowest density across undisturbed ecosystems, being

lower than all eleven other undisturbed ecosystems (no overlap of 50% CIs, of which eight

had no overlap of 90% CIs). Undisturbed tropical wetlands, forests and grasslands had a

lower density than their equivalent ecosystems in temperate and boreal regions (50% CI).

Disturbed tropical grasslands were similarly low in seed density, while arable ecosystems in

tropical regions were lower than the equivalent systems in other regions (90% CI).

Otherwise, arable systems combined relatively low richness with high seed density.

Mediterranean and desert, and temperate and boreal arable ecosystems were predicted to

have more seeds in the soil than all other ecosystems in the terrestrial realm (50% CIs),

while arable ecosystems often had lower richness at the gamma scale than other degraded

ecosystems within similar biome subtypes (for example, temperate and boreal arable

ecosystems had lower richness than temperate forests and temperate and boreal wetlands;

50% CIs). On the other hand, mediterranean forests, woodlands and scrub combined high

species richness with relatively high seed density (higher than all other forest ecosystems,

90% CI), while tundra combined its low species richness with low density, especially when

degraded, where density was lower than all other degraded ecosystems (50% CI).

Figure 2: Scale-dependent diversity in the soil seed banks of the world. Panels
represent ecosystem realm and biomes. Colours represent ecosystem type and
groups. Points represent average predicted species richness at two spatial scales of
soil sampled across realm/biomes. Large points represent the gamma (15 m^2) scale,
small points represent the alpha (0.01 m^2) scale. Densities represent 24 000
predictive draws of the posterior distributions of overall richness at each scale, within
each Biome subtype and degradation group. The black points show the mean, bar
represents 50 (thick) and 90% (thin) credible intervals. Colours indicate unique
ecosystem Biome subtypes. Shapes indicate undisturbed habitat (circle) and
disturbed/degraded habitat (triangle). Y axes vary for clarity.

Figure 3: Density of seeds m^{-2} in the soil seed bank. Panels represent ecosystem realms and biomes. Background jittered points represent empirical estimates of seeds m^{-2} within sampled soil banks. The large points show the mean, bar represents 50 (thick) and 90% (thin) credible intervals. Colours represent ecosystem types and groups, and the dashed reference line at 0 represents a value of 0. Shapes indicate undisturbed habitat (circle) and disturbed/degraded habitat (triangle). Thick colored lines represent the mean fixed effect value across broad realms (Natural Terrestrial, Arable, Wetlands, Aquatic). Y-axes vary for clarity.

Figure 4: The joint relationship between predicted species richness m^{-2} and seed density m^{-2} in the soil seed banks of the world. Panels represent ecosystem realms and biomes. The large points show the mean, bar represents 50 (thick) and 90% (thin) credible intervals. Colours represent ecosystem groups and habitat state, and the dashed reference line at 0 represents a value of 0. Shapes indicate undisturbed habitat (circle) and disturbed/degraded habitat (triangle). X and Y axes vary for clarity.

Discussion

We have used a global database of published soil seed bank studies to explore patterns of biodiversity in the soil seed bank worldwide. Although there was a high uncertainty in our results, we found that across the world's biomes and ecosystems, patterns in species richness in the seed bank often diverged from patterns of seed density. Strikingly, we also found that the richness of soil seed banks appeared higher in undisturbed compared to degraded habitat, suggesting how across the world's ecosystems, soil seed banks also reflect the ongoing depletion of biodiversity as a result of anthropogenic activities

(Eichenberg *et al.* 2021; Newbold *et al.* 2020). This indicates that the potential for seed

banks to contribute to the recovery and restoration of degraded ecosystems might be limited.

Seed density, on the other hand, was not always lower in disturbed habitats, showing how

different aspects of biodiversity can show diverging patterns in response to global change

pressures (Blowes *et al.* 2022). Our estimated global average of more than 5000 seeds m^{-2}

also means that in the broadest sense, if you put your finger to the earth, you are likely to be

pointing at a seed. The high density of seeds in the soil is a testament to the formation of soil

seed banks as an important component of plant population and community dynamics, and a

valuable strategy for many plant species to facilitate establishment following disturbance

(Alexander *et al.* 2012; Chesson 2000; Plue *et al.* 2017).

Scale-dependence of species richness in soil seed banks worldwide

We found that the species richness in terrestrial soil seed banks generally increases with

increasing area of ground sampled, but that the strength of this relationship varies across

biomes, ecosystems and between undisturbed and degraded habitats (Figure 2). Our

findings support those of Vandvik et al. (2016), who found that species accumulate more

rapidly in alpine and subalpine grasslands than in boreal zones (empirical), and in forests

compared to grassland and heathland (literature), and we extend the results to include other

global biomes. We found that richness estimates across small and large scales (in our case

0.01 m^2 and 15m^2) appeared to show relatively large differences in mediterranean and

tropical biomes compared to temperate biomes, demonstrating how species' spatial

distributions affect how we interpret relative differences in biodiversity across regions (Table

S3). Overall, our broad patterns of richness distribution in the global soil seed bank are in

line with other findings of relatively high tropical vs. temperate and boreal diversity, also

documented in global studies of mammals (Marsh *et al.* 2022), soil fungi (Tedersoo *et al.*

2014) and trees (Keil & Chase 2019; Liang *et al.* 2022). However, like others, we also found

some differences depending on the spatial scale of our predictions. For vascular plants,

Sabatini *et al.* (2022) identified a clearer gradient in species richness at large spatial grains,

while Keil & Chase (2019) found that areas of relatively high beta diversity in trees – where

richness was relatively low at smaller scales and relatively high at larger scales – to be

concentrated in low latitudes. Our modelled patterns of soil seed bank richness at the 0.01

m^2 scale were broadly similar to those found by Yang *et al.* (2021), although their findings of

low richness in the tropics and (predicted) higher richness in northern biomes were not

supported when considering larger spatial scales. Our results were characterised by high

levels of uncertainty. However, global maps of biodiversity do not generally map uncertainty

and only plot according to mean predicted values. Our findings highlight the need to be

cautious when drawing conclusions from such maps, with different aspects of biodiversity

varying across space and scale, and with a large degree of uncertainty.

Patterns and determinants of seed bank richness and density

Our analyses revealed substantial variation in the diversity of the soil seed bank among the

world's ecosystems. Importantly, we often found divergent patterns between species

richness and seed density. While mediterranean and tropical ecosystems were broadly

predicted to have higher richness than temperate and boreal ecosystems and tundra, seed

density was generally characterised by high variation within biomes. However,

mediterranean forests and shrublands systems exhibited both high richness and density

(Figures 2-4, Table S3). In regions that are characterised by a warm climate and frequent

extreme climatic events, the soil seed bank can play an important role in vegetation recovery

and stability of biodiversity (DeMalach *et al.* 2023; Lloret *et al.* 2012), and it follows that

many plant species will form persistent seed banks. Previous work from Europe has shown

that species with relatively warm, southerly distributions are more likely to have persistent

seeds than cool-adapted species (Auffret *et al.* 2023), and we also found lower richness in

temperate and boreal grasslands and forests, and lower density in the majority of temperate

and boreal habitats compared to their Mediterranean counterparts. The vegetation in the

temperate and boreal zones can be considered more stable and dominated by long-lived

perennial species, for which regeneration via the soil seed bank is a less important strategy

than for example clonality (Økland 1995). In alpine and tundra environments, low seed bank

density and a high proportion of clonal species could be expected due to a low diversity and

density of adult individuals, combined with the risks of seed production during short growing

seasons (Gough 2006; Körner 2021). In tropical ecosystems, relatively few species produce

persistent seeds (Tweddle *et al.* 2003; Wyse & Dickie 2017), while the warm and often moist

conditions combined with high diversity of other taxa mean that seeds are subject to

degradation, predation and attacks from pathogens (Baskin & Baskin 2014; Dalling *et al.*

1998; Skoglund 1992). This can result in low soil seed bank density, but the high overall

diversity in the vegetation means that species richness is still high relative to other biomes.

Our results show an imprint of anthropogenic activity on soil seed banks worldwide. Perhaps

unsurprisingly, arable soils contained among the highest number of seeds per square metre

(Figure 3,4). Highly and regularly-disturbed arable systems accumulate the persistent seeds

of comparatively few opportunistic ruderal "weed" species that can be problematic for

agricultural production. Indeed, many of the 202 studies contributing to the 769 arable

observations in the global seed bank database were investigations of the effects of

agricultural management on the weed seed bank (Ikeda *et al.* 2007; Rotchés-Ribalta *et al.*

2017; Schwartz *et al.* 2015). However, in non-arable open ecosystems, seed density was

often lower with habitat degradation, while the opposite was true for forested ecosystems.

Many degraded forests are those that are recovering from previous agricultural or

silvicultural management, and therefore their seed banks can still reflect those of highly-

disturbed conditions (Daïnou *et al.* 2011; Dupuy & Chazdon 1998; Plue *et al.* 2010). In other

habitats, degradation generally appeared to result in both reduced richness and density in

the soil seed bank, in line with reports of reduced biodiversity in the established vegetation

over time (Daru *et al.* 2021; Jandt *et al.* 2022; but see Vellend *et al.* 2013), with

consequences for the potential of passive restoration in reversing biodiversity losses

(Atkinson *et al.* 2022; Ladouceur *et al.* 2023).

Global insights into seed bank sampling

The large number of soil seed bank studies that have been carried out across the world has

allowed our synthesis to cover a greater global extent than many large reviews of

biodiversity, which often suffer from biases relating to the absence or restricted availability of

data from some regions (Boakes *et al.* 2010; Hughes *et al.* 2021; Nuñez & Amano 2021).

However, our study may still suffer from other biases, originating from the fact that seed

banks have not been sampled randomly across the world, but in places where scientists

have been interested in finding and identifying seeds in the seed bank (Boyd *et al.* 2023).

Combined with variation in sampling effort, this wealth of soil seed bank data has still

allowed us to gain interesting insights into the spatial scaling and biogeographic patterns of

seed bank diversity. However, some ecosystems were better-represented than others, which

may affect our predictions of richness at different spatial scales. It is also well-known that

sampling multiple sites per treatment gives improved confidence in ecological findings

(sample replication; Filazzola & Cahill Jr 2021), but around half of all observations in the

database represented a single site. There have previously been calls to improve the seed

bank sampling to improve representativeness (Bigwood & Inouye 1988; Plue & Hermy

2012), and we add to this a request to report soil seed bank data at the smallest scale

possible (e.g. replicate, pooled replicates or site level), or at least making it possible to

estimate the geographic spread of sites included in an observation. This would help to create

an even better understanding of seed bank diversity and its spatial scaling.

The growth of soil seed bank synthesis

What can we learn from studying the soil seed bank? In addition to the continual publication

of studies investigating soil seed banks at local scales, there has been an upswing in the

synthesis of large datasets to understand broad spatial or ecological patterns relating to soil

seed banks. This includes the macro-environmental determinants of small-scale richness

and density (Yang *et al.* 2021), and the evolutionary history of seed bank persistence (Gioria

et al. 2020). Here, we complement such findings with an understanding of the spatial scaling

of species richness in the soil seed bank, and broad biogeographical patterns in seed bank

richness and density. Other large-scale work considers the important role that soil seed

banks may play in species and community responses to global changes, related to the trade-

off related to seeds being able to persist or disperse in space (Chen *et al.* 2020), or the role

of soil seed banks in facilitating biological invasions (Gioria *et al.* 2021). We believe that the

future of soil seed bank synthesis lies beyond simple measures of community richness and

density, or categorisation of species as having or not having persistent seeds. Collating a

large number of datasets including community data from the soil seed bank and established

vegetation – not forgetting to standardise according to spatial scale – across biomes and

systems (see Plue *et al.* 2021 for a less-than-comprehensive European study) would allow

us to delve deeper into a mechanistic understanding of the role of soil seed banks,

biodiversity, restoration and responses to global change.

References

Alexander, H.M., Foster, B.L., Ballantyne, F., Collins, C.D., Antonovics, J. & Holt, R.D.

(2012). Metapopulations and metacommunities: combining spatial and temporal

perspectives in plant ecology. *Journal of Ecology*, 100, 88–103.

Arrhenius, O. (1921). Species and area. *Journal of Ecology*, 9, 95–99.

Atkinson, J., Brudvig, L.A., Mallen-Cooper, M., Nakagawa, S., Moles, A.T. & Bonser, S.P.

(2022). Terrestrial ecosystem restoration increases biodiversity and reduces its

variability, but not to reference levels: A global meta-analysis. *Ecology Letters*, 25,

1725–1737.

Auffret, A.G. & Cousins, S.A.O. (2011). Past and present management influences the seed

bank and seed rain in a rural landscape mosaic. *Journal of Applied Ecology*, 48,

1278–1285.

Auffret, A.G., Ladouceur, E., Haussmann, N.S., Daouti, E., Elumeeva, T.G., Kačergytė, I., *et*

al. (2024). A global database of soil seed bank richness, density, and abundance.

Ecology, 105, e4438.

Auffret, A.G., Vangansbeke, P., De Frenne, P., Auestad, I., Basto, S., Grandin, U., *et al.*

(2023). More warm-adapted species in soil seed banks than in herb layer plant

communities across Europe. *Journal of Ecology*, 111, 1009–1020.

Baptista, T.L. & Shumway, S.W. (1998). A Comparison of the Seed Banks of Sand Dunes

with Different Disturbance Histories on Cape Cod National Seashore. *Rhodora*, 100,

298–313.

Baskin, C.C. & Baskin, J.M. (2014). Chapter 9 - A Geographical Perspective on Germination

Ecology: Tropical and Subtropical Zones. In: *Seeds (Second Edition)* (eds. Baskin,

C.C. & Baskin, J.M.). Academic Press, San Diego, pp. 375–590.

Bigwood, D.W. & Inouye, D.W. (1988). Spatial Pattern Analysis of Seed Banks: An Improved

Method and Optimized Sampling. *Ecology*, 69, 497–507.

Blowes, S.A., Daskalova, G.N., Dornelas, M., Engel, T., Gotelli, N.J., Magurran, A.E., *et al.*

(2022). Local biodiversity change reflects interactions among changing abundance,

evenness, and richness. *Ecology*, 103, e3820.

Boakes, E.H., McGowan, P.J.K., Fuller, R.A., Chang-qing, D., Clark, N.E., O'Connor, K., *et*

al. (2010). Distorted Views of Biodiversity: Spatial and Temporal Bias in Species

Occurrence Data. *PLoS Biol*, 8, e1000385.

Bossuyt, B. & Honnay, O. (2008). Can the seed bank be used for ecological restoration? An

overview of seed bank characteristics in European communities. *Journal of*

Vegetation Science, 19, 875–884.

Boyd, R.J., Powney, G.D. & Pescott, O.L. (2023). We need to talk about nonprobability

samples. *Trends in Ecology & Evolution*, 38, 521–531.

Boyes, L.J., Gunton, R.M., Griffiths, M.E. & Lawes, M.J. (2011). Causes of arrested

succession in coastal dune forest. *Plant Ecol*, 212, 21–32.

Bürkner, P.-C. (2018). Advanced Bayesian Multilevel Modeling with the R Package brms.

The R Journal, 10, 395–411.

Burmeier, S., Eckstein, R.L., Otte, A. & Donath, T.W. (2010). Desiccation cracks act as

natural seed traps in flood-meadow systems. *Plant Soil*, 333, 351–364.

Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M., *et al.*

(2017). Stan: A Probabilistic Programming Language. *Journal of Statistical Software*,

76, 1–32.

Chen, S.-C., Poschlod, P., Antonelli, A., Liu, U. & Dickie, J.B. (2020). Trade-off between

seed dispersal in space and time. *Ecology Letters*, 23, 1635–1642.

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of*

Ecology and Systematics, 31, 343–366.

Daïnou, K., Bauduin, A., Bourland, N., Gillet, J.-F., Fétéké, F. & Doucet, J.-L. (2011). Soil

seed bank characteristics in Cameroonian rainforests and implications for post-

logging forest recovery. *Ecological Engineering*, 37, 1499–1506.

Dalling, J.W., Swaine, M.D. & Garwood, N.C. (1998). Dispersal Patterns and Seed Bank

Dynamics of Pioneer Trees in Moist Tropical Forest. *Ecology*, 79, 564–578.

Daru, B.H., Davies, T.J., Willis, C.G., Meineke, E.K., Ronk, A., Zobel, M., *et al.* (2021).

Widespread homogenization of plant communities in the Anthropocene. *Nat*

Commun, 12, 6983.

DeMalach, N., Kigel, J. & Sternberg, M. (2021). The soil seed bank can buffer long-term

compositional changes in annual plant communities. *Journal of Ecology*, 109, 1275–

1283.

DeMalach, N., Kigel, J. & Sternberg, M. (2023). Contrasting dynamics of seed banks and

standing vegetation of annuals and perennials along a rainfall gradient. *Perspectives*

in Plant Ecology, Evolution and Systematics, 58, 125718.

Drakare, S., Lennon, J.J. & Hillebrand, H. (2006). The imprint of the geographical,

evolutionary and ecological context on species-area relationships. *Ecology Letters*, 9,

215–227.

Dupuy, J.M. & Chazdon, R.L. (1998). Long-Term Effects of Forest Regrowth and Selective

Logging on the Seed Bank of Tropical Forests in NE Costa Rica. *Biotropica*, 30, 223–

237.

Eichenberg, D., Bowler, D.E., Bonn, A., Bruelheide, H., Grescho, V., Harter, D., *et al.* (2021).

Widespread decline in Central European plant diversity across six decades. *Global*

Change Biology, 27, 1097–1110.

Eriksson, O. & Jakobsson, A. (1998). Abundance, distribution and life histories of grassland

plants: a comparative study of 81 species. *Journal of Ecology*, 86, 922–933.

Filazzola, A. & Cahill Jr, J.F. (2021). Replication in field ecology: Identifying challenges and

proposing solutions. *Methods in Ecology and Evolution*, 12, 1780–1792.

Friedman, J. (2020). The evolution of annual and perennial plant life histories: ecological

correlates and genetic mechanisms. *Annual Review of Ecology, Evolution, and*

Systematics, 51, 461–481.

Gaba, S., Perronne, R., Fried, G., Gardarin, A., Bretagnolle, F., Biju-Duval, L., *et al.* (2017).

Response and effect traits of arable weeds in agro-ecosystems: a review of current

knowledge. *Weed Research*, 57, 123–147.

Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. (2019). Visualization in

Bayesian Workflow. *Journal of the Royal Statistical Society Series A: Statistics in*

Society, 182, 389–402.

Gibson, D.J., Young, B.G., Owen, M.D., Gage, K.L., Matthews, J.L., Jordan, D.L., *et al.*

(2016). Benchmark study on glyphosate-resistant cropping systems in the United

States. Part 7: Effects of weed management strategy (grower practices versus

academic recommendations) on the weed soil seedbank over 6 years. *Pest*

Management Science, 72, 692–700.

Gioria, M., Carta, A., Baskin, C.C., Dawson, W., Essl, F., Kreft, H., *et al.* (2021). Persistent

soil seed banks promote naturalisation and invasiveness in flowering plants. *Ecology*

Letters, 24, 1655–1667.

Gioria, M., Pyšek, P., Baskin, C.C. & Carta, A. (2020). Phylogenetic relatedness mediates

persistence and density of soil seed banks. *Journal of Ecology*, 108, 2121–2131.

Gough, L. (2006). Neighbor effects on germination, survival, and growth in two arctic tundra

plant communities. *Ecography*, 29, 44–56.

Graham, L.L.B. & Page, S.E. (2018). A limited seed bank in both natural and degraded

tropical peat swamp forest: the implications for restoration. *Mires and Peat*, 1–13.

Grime, J.P. (1977). Evidence for the Existence of Three Primary Strategies in Plants and Its

Relevance to Ecological and Evolutionary Theory. *The American Naturalist*, 111,

1169–1194.

Grubb, P.J. (1977). The Maintenance of Species-Richness in Plant Communities: The

Importance of the Regeneration Niche. *Biological Reviews*, 52, 107–145.

Hausman, C.E., Fraser, L.H., Kershner, M.W. & de Szalay, F.A. (2007). Plant community

establishment in a restored wetland: Effects of soil removal. *Applied Vegetation*

Science, 10, 383–390.

Hawes, C., Squire, G.R., Hallett, P.D., Watson, C.A. & Young, M. (2010). Arable plant

communities as indicators of farming practice. *Agriculture, Ecosystems &*

Environment, 138, 17–26.

He, Y., Gao, P. & Qiang, S. (2019). An investigation of weed seed banks reveals similar

potential weed community diversity among three different farmland types in Anhui

Province, China. *Journal of Integrative Agriculture*, 18, 927–937.

HilleRisLambers, J., Clark, J.S. & Lavine, M. (2005). Implications of Seed Banking for

Recruitment of Southern Appalachian Woody Species. *Ecology*, 86, 85–95.

Holland, J.M., Smith, B.M., Southway, S.E., Birkett, T.C. & Aebischer, N.J. (2008). The effect

of crop, cultivation and seed addition for birds on surface weed seed densities in

arable crops during winter. *Weed Research*, 48, 503–511.

Honnay, O., Bossuyt, B., Jacquemyn, H., Shimono, A. & Uchiyama, K. (2008). Can a seed

bank maintain the genetic variation in the above ground plant population? *Oikos*,

117, 1–5.

Hughes, A.C., Orr, M.C., Ma, K., Costello, M.J., Waller, J., Provoost, P., *et al.* (2021).

Sampling biases shape our view of the natural world. *Ecography*, 44, 1259–1269.

Ikeda, F.S., Mitja, D., Vilela, L. & Carmona, R. (2007). Banco de sementes no solo em

sistemas de cultivo lavoura-pastagem. *Pesq. agropec. bras.*, 42, 1545–1551.

Jandt, U., Bruelheide, H., Jansen, F., Bonn, A., Grescho, V., Klenke, R.A., *et al.* (2022).

More losses than gains during one century of plant biodiversity change in Germany.

Nature, 611, 512–518.

Kambach, S., Sabatini, F.M., Attorre, F., Biurrun, I., Boenisch, G., Bonari, G., *et al.* (2023).

Climate-trait relationships exhibit strong habitat specificity in plant communities

across Europe. *Nat Commun*, 14, 712.

Karlík, P. & Poschlod, P. (2014). Soil seed-bank composition reveals the land-use history of

calcareous grasslands. *Acta Oecologica*, 58, 22–34.

Keil, P. & Chase, J.M. (2019). Global patterns and drivers of tree diversity integrated across

a continuum of spatial grains. *Nat Ecol Evol*, 3, 390–399.

Kirmer, A. & Mahn, E.-G. (2001). Spontaneous and initiated succession on unvegetated

slopes in the abandoned lignite-mining area of Goitsche, Germany. *Applied*

Vegetation Science, 4, 19–27.

Klimešová, J. & Herben, T. (2015). Clonal and bud bank traits: patterns across temperate

plant communities. *Journal of Vegetation Science*, 26, 243–253.

Körner, C. (2021). *Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems*.

Springer International Publishing, Cham.

Kuester, A., Conner, J.K., Culley, T. & Baucom, R.S. (2014). How weeds emerge: a

taxonomic and trait-based examination using United States data. *New Phytologist*,

202, 1055–1068.

Ladouceur, E., Bonomi, C., Bruelheide, H., Klimešová, J., Burrascano, S., Poschlod, P., *et*

al. (2019). The functional trait spectrum of European temperate grasslands. *Journal*

of Vegetation Science, 30, 777–788.

Ladouceur, E., Isbell, F., Clark, A.T., Harpole, W.S., Reich, P.B., Tilman, G.D., *et al.* (2023).

The recovery of plant community composition following passive restoration across

spatial scales. *Journal of Ecology*, 111, 814–829.

Liang, J., Gamarra, J.G.P., Picard, N., Zhou, M., Pijanowski, B., Jacobs, D.F., *et al.* (2022).

Co-limitation towards lower latitudes shapes global forest diversity gradients. *Nat*

Ecol Evol, 6, 1423–1437.

Lloret, F., Escudero, A., Iriondo, J.M., Martínez-Vilalta, J. & Valladares, F. (2012). Extreme

climatic events and vegetation: the role of stabilizing processes. *Global Change*

Biology, 18, 797–805.

Malfasi, F., Convey, P., Zaccara, S. & Cannone, N. (2020). Establishment and eradication of

an alien plant species in Antarctica: Poa annua at Signy Island. *Biodivers Conserv*,

29, 173–186.

Marsh, C.J., Sica, Y.V., Burgin, C.J., Dorman, W.A., Anderson, R.C., del Toro Mijares, I., *et*

al. (2022). Expert range maps of global mammal distributions harmonised to three

taxonomic authorities. *Journal of Biogeography*, 49, 979–992.

Matlack, G.R. & Good, R.E. (1990). Spatial Heterogeneity in the Soil Seed Bank of a Mature

Coastal Plain Forest. *Bulletin of the Torrey Botanical Club*, 117, 143–152.

Newbold, T., Oppenheimer, P., Etard, A. & Williams, J.J. (2020). Tropical and Mediterranean

biodiversity is disproportionately sensitive to land-use and climate change. *Nat Ecol*

Evol, 4, 1630–1638.

Nilsson, M.-C. & Wardle, D.A. (2005). Understory vegetation as a forest ecosystem driver:

evidence from the northern Swedish boreal forest. *Frontiers in Ecology and the*

Environment, 3, 421–428.

Nuñez, M.A. & Amano, T. (2021). Monolingual searches can limit and bias results in global

literature reviews. *Nat Ecol Evol*, 5, 264–264.

Økland, R.H. (1995). Persistence of vascular plants in a Norwegian boreal coniferous forest.

Ecography, 18, 3–14.

Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V.N.,

Underwood, E.C., *et al.* (2001). Terrestrial Ecoregions of the World: A New Map of

Life on Earth A new global map of terrestrial ecoregions provides an innovative tool

for conserving biodiversity. *BioScience*, 51, 933–938.

Ooi, M.K.J., Auld, T.D. & Denham, A.J. (2009). Climate change and bet-hedging:

interactions between increased soil temperatures and seed bank persistence. *Global*

Change Biology, 15, 2375–2386.

Plue, J., Calster, H.V., Auestad, I., Basto, S., Bekker, R.M., Bruun, H.H., *et al.* (2021).

Buffering effects of soil seed banks on plant community composition in response to

land use and climate. *Global Ecology and Biogeography*, 30, 128–139.

Plue, J., Frenne, P.D., Acharya, K., Brunet, J., Chabrerie, O., Decocq, G., *et al.* (2017).

Where does the community start, and where does it end? Including the seed bank to

reassess forest herb layer responses to the environment. *Journal of Vegetation*

Science, 28, 424–435.

Plue, J. & Hermy, M. (2012). Consistent seed bank spatial structure across semi-natural

habitats determines plot sampling. *Journal of Vegetation Science*, 23, 505–516.

Plue, J., Verheyen, K., Van Calster, H., Marage, D., Thompson, K., Kalamees, R., *et al.*

(2010). Seed banks of temperate deciduous forests during secondary succession.

Journal of Vegetation Science, 21, 965–978.

Rotchés-Ribalta, R., Armengot, L., Mäder, P., Mayer, J. & Sans, F.X. (2017). Long-Term

Management Affects the Community Composition of Arable Soil Seedbanks. *Weed*

Science, 65, 73–82.

Saatkamp, A., Poschlod, P. & Venable, D.L. (2014). The functional role of soil seed banks in

natural communities. *Seeds: the ecology of regeneration in plant communities*, CABI

Books, 263–295.

Sabatini, F.M., Jiménez-Alfaro, B., Jandt, U., Chytrý, M., Field, R., Kessler, M., *et al.* (2022).

Global patterns of vascular plant alpha diversity. *Nat Commun*, 13, 4683.

Salonen, J., Hyvönen, T. & Jalli, H. (2011). Composition of weed flora in spring cereals in

Finland - a fourth survey.

Sarneel, J.M., Beltman, B., Buijze, A., Groen, R. & Soons, M.B. (2014). The role of wind in

the dispersal of floating seeds in slow-flowing or stagnant water bodies. *Journal of*

Vegetation Science, 25, 262–274.

Schwartz, L.M., Gibson, D.J., Gage, K.L., Matthews, J.L., Jordan, D.L., Owen, M.D.K., *et al.*

(2015). Seedbank and Field Emergence of Weeds in Glyphosate-Resistant Cropping

Systems in the United States. *Weed Science*, 63, 425–439.

Skoglund, J. (1992). The role of seed banks in vegetation dynamics and restoration of dry

tropical ecosystems. *Journal of Vegetation Science*, 3, 357–360.

Tedersoo, L., Bahram, M., Põlme, S., Kõljalg, U., Yorou, N.S., Wijesundera, R., *et al.* (2014).

Global diversity and geography of soil fungi. *Science*, 346, 1256688.

Thompson, K. (1986). Small-Scale Heterogeneity in the Seed Bank of An Acidic Grassland.

Journal of Ecology, 74, 733–738.

Thompson, K. (1987). Seeds and Seed Banks. *New Phytologist*, 106, 23–34.

Török, P., Helm, A., Kiehl, K., Buisson, E. & Valkó, O. (2018). Beyond the species pool:

modification of species dispersal, establishment, and assembly by habitat restoration.

Restoration Ecology, 26, S65–S72.

Tweddle, J.C., Dickie, J.B., Baskin, C.C. & Baskin, J.M. (2003). Ecological aspects of seed

desiccation sensitivity. *Journal of Ecology*, 91, 294–304.

Vandvik, V., Klanderud, K., Meineri, E., Måren, I.E. & Töpper, J. (2016). Seed banks are

biodiversity reservoirs: species–area relationships above versus below ground.

Oikos, 125, 218–228.

Vellend, M., Baeten, L., Myers-Smith, I.H., Elmendorf, S.C., Beauséjour, R., Brown, C.D., *et*

al. (2013). Global meta-analysis reveals no net change in local-scale plant

biodiversity over time. *PNAS*, 201312779.

Westcott, K., Whillans, T.H. & Fox, M.G. (1997). Viability and abundance of seeds of

submerged macrophytes in the sediment of disturbed and reference shoreline

marshes in Lake Ontario. *Can. J. Bot.*, 75, 451–456.

Wickham, H. (2009). *ggplot2 - Elegant Graphics for Data Analysis*. Springer, New York.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., *et al.* (2019).

Welcome to the Tidyverse. *Journal of Open Source Software*, 4, 1686.

Wisnoski, N.I. & Shoemaker, L.G. (2022). Seed banks alter metacommunity diversity: The

interactive effects of competition, dispersal and dormancy. *Ecology Letters*, 25, 740–

753.

Wyse, S.V. & Dickie, J.B. (2017). Predicting the global incidence of seed desiccation

sensitivity. *Journal of Ecology*, 105, 1082–1093.

Yang, X., Baskin, C.C., Baskin, J.M., Pakeman, R.J., Huang, Z., Gao, R., *et al.* (2021).

Global patterns of potential future plant diversity hidden in soil seed banks. *Nat*

Commun, 12, 7023.

Zaghloul, M.S. (2008). Diversity in soil seed bank of Sinai and implications for conservation

and restoration. *African Journal of Environmental Science and Technology*, 2, 172–

184.