1 **Experimental evolution for the recovery of growth loss due to genome reduction**

3 Kenya Hitomi[1], Yoichiro Ishii[1], Bei-Wen Ying[1,*]

5 [1]School of Life and Environmental Sciences, University of Tsukuba, Tennodai 1-1-1,

6 Tsukuba, Ibaraki 305-8572, Japan

7 [*]Correspondence: ying.beiwen.gf@u.tsukuba.ac.jp

Abstract

As the genome encodes the information crucial for cell growth, a sizeable genomic deficiency often causes a significant decrease in growth fitness. Whether and how the decreased growth fitness caused by genome reduction could be compensated by evolution was investigated here. Experimental evolution with an *Escherichia coli* strain carrying a reduced genome was conducted in multiple lineages for approximately 1,000 generations. The growth rate, which was largely declined due to genome reduction, was considerably recovered, associated with the improved carrying capacity. Genome mutations accumulated during evolution were significantly varied across the evolutionary lineages and were randomly localized on the reduced genome. Transcriptome reorganization showed a common evolutionary direction and conserved the chromosomal periodicity, regardless of highly diversified gene categories, regulons, and pathways enriched in the differentially expressed genes. Genome mutations and transcriptome reorganization caused by evolution, which were found to be dissimilar to those caused by genome reduction, must have followed divergent mechanisms in individual evolutionary lineages. Gene network reconstruction successfully identified three gene modules functionally differentiated, which were responsible for the evolutionary changes of the reduced genome in growth fitness, genome mutation, and gene expression, respectively. The diversity in evolutionary approaches improved the growth fitness associated with the homeostatic transcriptome architecture as if the evolutionary compensation for genome reduction was like all roads leading to Rome.

Keywords: genome reduction, experimental evolution, growth fitness, transcriptome, genome mutation, chromosomal periodicity, gene expression, network analysis

Introduction

The genome encodes the information for cell growth, and its size is likely the evolutionary consequence [1-3]. To determine the essential genomic sequence of modern cells, removing redundant DNA sequences from the wild-type genome in bacteria, so-called genome reduction, has been challenged to a large extent [4-7]. These efforts have been made to discover the minimal genetic requirement for a free-living organism growing under the defined conditions [8-10]. It resulted in the finding of the coordination of genome with cell growth, i.e., genome reduction significantly decreased the growth rate of *Escherichia coli* (*E. coli*) cells independent of culture media or growth forms [11-13]. Slow growth or fitness decline was commonly observed in the genetically reduced [13,14] and chemically synthesized genomes [10,15].

The growth decrease could be recovered by experimental evolution. Growth fitness generally represents the adaptiveness of the living organism to the defined environment [16]. Although the reduced genomes somehow showed differentiated evolvability compared to the wild-type genomes [17-19], their evolutionary adaptation to the environmental changes has been successfully achieved [20-22]. Experimental evolution under the defined culture condition successfully increased the decreased growth rates of the reduced genomes [17,20,21] and fastened the slow-growing synthetic genome [23]. The evolutionary rescue of the growth rate must be associated with the changes in genomic sequence and gene expression benefited for the growth fitness, as what happened in nature adaptive evolution [24-26]. Our previous study showed that the decreased growth rate caused by the absence of a sizeable genomic sequence could be complemented by introducing the mutations elsewhere in the genome [20,21]. Additionally, the changes in growth rate caused by either genome reduction or experimental evolution were dependent on the genome size but not the specific gene function [20].

A genome-wide understanding of the evolutionary compensated fitness increase of the reduced genome is required. The experimental evolution compensated for the genome reduction-mediated growth changes were considered stringently related to transcriptome reorganization. Previous studies have identified conserved features in transcriptome reorganization, despite significant disruption to gene expression patterns resulting from either genome reduction or experimental evolution [27-29]. The findings

indicated that experimental evolution might reinstate growth rates that have been

disrupted by genome reduction to maintain homeostasis in growing cells. Although the

reduced growth rate caused by genome reduction could be recovered by experimental

evolution, it remains unclear whether such an evolutionary improvement in growth

fitness was a general feature of the reduced genome and how the genome-wide changes

occurred to match the growth fitness increase. In the present study, we performed the

experimental evolution with a reduced genome in multiple lineages and analyzed the

evolutionary changes of the genome and transcriptome.

Results

Fitness recovery of the reduced genome by experimental evolution

Experimental evolution of the reduced genome was conducted to regain the growth

fitness, which was decreased due to genome reduction. The *E. coli* strain carrying a

reduced genome, derived from the wild-type genome W3110, showed a significant

decline in its growth rate in the minimal medium compared to the wild-type strain [13]. To

improve the genome reduction-mediated decreased growth rate, the serial transfer of the

genome-reduced strain was performed with multiple dilution rates to keep the bacterial

growth within the exponential phase (Fig. S1), as described [17,20]. Nine evolutionary

lineages were conducted independently (Fig. S2, Table S1). A gradual increase in

growth rate was observed along with the generation passage, which was combined with

a rapid increase in the early evolutionary phase and a slow increase in the later phase

(Fig. 1A). As the growth increases were calculated according to the initial and final

records, the exponential growth rates of the ancestor and evolved populations were

obtained according to the growth curves for a precise evaluation of the evolutionary

changes in growth. The results demonstrated that most evolved populations (Evos)

showed improved growth rates, in which eight out of nine Evos were highly significant

(Fig. 1B, upper). However, the magnitudes of growth improvement were considerably

varied, and the evolutionary dynamics of the nine lineages were somehow divergent

(Fig. S2). It indicated the diversity in the evolutionary approaches for improved fitness.

Eight of nine Evos achieved faster growth than the genome-reduced ancestor (Anc),

whereas all Evos decreased in their saturated population densities (Fig. 1B, bottom). In

comparison to the wild-type strain carrying the full-length genome (WT), the primarily

decreased growth rate caused by genome reduction was significantly improved by

experimental evolution (Fig. 1C, upper), associated with the considerable reduction in

saturated density (Fig. 1C, bottom). It demonstrated that the experimental evolution

could compensate for the genome reduction with a trade-off in population size (carrying

capacity), consistent with the previous findings [20,21].

Intriguingly, a positive correlation was observed between the growth fitness and the

carrying capacity of the Evos (Fig. 1D). It was somehow consistent with the positive

correlations between the colony growth rate and the colony size of a genome-reduced

strain [11] and between the growth rates and the saturated population size of an assortment

of genome reduced strains [13]. Nevertheless, the negative correlation between growth

rate and carrying capacity, known as the r/K selection[30,31] was often observed as the

trade-off relationship between r and K in the evolution and ecology studies [32] [33,34]. As

the r/K trade-off was proposed to balance the cellular metabolism that resulted from the

cost of enzymes involved [34], the deleted genes might play a role in maintaining the

metabolism balance for the r/K correlation. On the other hand, the experimental

evolution (i.e., serial transfer) was strictly performed within the exponential growth

phase; thus, the evolutionary selection was supposed to be driven by the growth rate

without selective pressure to maintain the carrying capacity. The declined carrying

capacity might have been its neutral "drift" but not a trade-off to the growth rate.

Independent and parallel experimental evolution of the reduced genomes selecting

either r or K is required to clarify the actual mechanisms.

Significant variation and random localization of genome mutations

Genome resequencing (Table S2) identified a total of 65 mutations fixed in the nine

Evos (Table 1). The number of mutations largely varied among the nine Evos, from 2 to

13, and no common mutation was detected in all nine Evos (Table S3). A 1,199-bp

deletion of *insH* was frequently found in the Evos (Table S3, highlighted), which well

agreed with its function as a transposable sequence. 51 out of 65 mutations occurred in

the genes, and 45 out of 51 were SNPs. As 36 out of 45 SNPs were nonsynonymous,

which was highly significant compared to random mutations ($p < 0.01$), the mutated

genes might benefit fitness increase. In addition, the abundance of mutations was

unlikely to be related to the magnitude of fitness increase. There was no significant

correlation between the number of mutations and the growth rate in a statistical view (p

> 0.1). Even from an individual close-up viewpoint, the abundance of mutations poorly

explained the fitness increase. For instance, A2 accumulating only two mutations

presented a highly increased growth rate compared to F2 of eight mutations (Table 1),

which poorly improved the growth (Fig. 1B). B2, D2, and E2 all succeeded in

increasing fitness to an equivalent degree (Fig. 1B), whereas they fixed 13, 7, and 3

mutations, respectively (Table 1). The mutated genes were varied in 14 gene categories,

somehow more frequently in the gene categories of Transporter, Enzyme, and Unknown

function (Fig. S3). They seemed highly related to essentiality[7]

(https://shigen.nig.ac.jp/ecoli/pec/genes.jsp), as 11 out of 49 mutated genes were

essential (Table S3). Although the essentiality of genes might differ between the

wild-type and reduced genomes, the experimentally determined 302 essential genes in

the wild-type *E. coli* strain were used for the analysis, of which 286 were annotated in

the reduced genome. The ratio of essential genes in the mutated genes was significantly

higher than in the total genes (286 out of 3290 genes, Chi-square test $p=0.008$). As the

essential genes were determined according to the growth[35] and were known to be more

conserved than nonessential ones [36,37], the high frequency of the mutations fixed in the

essential genes was highly intriguing and reasonable. The large variety of genome

mutations fixed in the independent lineages might result from a highly rugged fitness

landscape [38]. Nevertheless, it was unclear whether and how these mutations were

explicitly responsible for recovering the growth rate of the reduced genome.

Additionally, there were no overlapping genomic positions for the 65 mutations in the

nine Evos (Fig. 2A). Random simulation was performed to verify whether there was any

bias or hotspot in the genomic location for mutation accumulation due to the genome

reduction. A total of 65 mutations were randomly generated on the reduced genome (Fig.

2B), and the genomic distances from the mutations to the nearest genome

reduction-mediated scars were calculated. Welch's t-test was performed to evaluate

whether the genomic distances calculated from random mutations significantly differed

from those from the mutations accumulated in the Evos. As the mean of p values (1,000

times of random simulations) was insignificant (Fig. 2C, $\mu_p > 0.05$), the mutations fixed

on the reduced genome were either closer or farther to the genomic scars, indicating

there was no locational bias for mutation accumulation caused by genome reduction.

Common evolutionary direction and homeostasis in transcriptome reorganization

Since no specificity was detected in the genome mutations, whether these mutations

disturbed the genome-wide gene expression pattern was investigated. Hierarchical

clustering and principal component analysis (PCA) showed that the evolved

transcriptomes were directed to similar patterns but divergent to the WT transcriptome

(Fig. S4). The evolutionary direction of the transcriptomes of the reduced genome was

not approaching the wild-type transcriptome. As the *E. coli* chromosome was structured,

whether the genome reduction caused the changes in its architecture, which led to the

differentiated transcriptome reorganization in the Evos, was investigated. The

chromosomal periodicity of gene expression was analyzed to determine the structural

feature of genome-wide pattern, as previously described [28,39]. The analytical results

showed that the transcriptomes of all Evos presented a common six-period with

statistical significance, equivalent to those of the wild-type and ancestral reduced

genomes (Fig. 3A, Table S4). It demonstrated that the chromosomal architecture of gene

expression patterns remained highly conserved, regardless of the considerably varied

mutations. The homeostatic periodicity was consistent with our previous findings that

the chromosomal periodicity of the transcriptome was independent of genomic or

environmental variation [27,28].

In addition, the genomic locations of the mutations seemed irrelevant to

chromosomal periodicity (Fig. 3A, red lines). No mutagenesis hotspot was observed

even if these mutations were accumulated on a single genome (Fig. 2A). Alternatively,

the expression levels of the mutated genes were somehow higher than those of the

remaining genes (Fig. 3B). As the regulatory mechanisms or the gene functions were

supposed to be disturbed by the mutations, the expression levels of individual genes

might have been either up- or down-regulated. Nevertheless, the overall expression

levels of all mutated genes tended to be increased. One of the reasons was assumed to

be the mutation essentiality, which remained to be experimentally verified. The ratio of

the essential genes in the mutated genes was ~ 22% (Table S3), much higher than the

ratio (~9%) of essential to all genes (286 out of 3,290) in the reduced genome. As the

essential genes showed higher expression levels than the nonessential ones [40], the high

essentiality of the mutated genes might result in a higher mean expression level. On the

other hand, the high frequency of the mutations fixed in the essential genes was

unexpected because the essential genes were generally more conserved than

nonessential ones [36]. The essentiality of the mutations might have participated in

maintaining the homeostatic transcriptome architecture of the reduced genome.

Diversified functions and pathways of the differentially expressed genes

As the evolved transcriptomes were differentiated from those of the WT and reduced

genomes (Fig. S4), the differentially expressed genes (DEGs) were further identified to

discover the gene functions or biological processes contributing to the fitness changes.

Instead of an in-depth survey on the directional changes of the DEGs, the abundance

and functional enrichment of DEGs were investigated to achieve an overview of how

significant the genome-wide fluctuation in gene expression, which ignored the details of

individual genes. The abundance of DEGs among the Evos varied from 333 to 1,130

genes and of few overlaps (Fig. 4A). Most DEGs were unique to each evolutionary

lineage, and the common DEGs across all Evos were only 108 genes (Table S5). The

number of DEGs partially overlapped among the Evos declined along with the

increased lineages of Evos (Fig. 4B). Enrichment analysis showed that only the

histidine-related pathways were significantly enriched in the common DEGs (Fig. 4C).

Functional enrichment of the DEGs in individual Evos showed that the amino acid

metabolism considerably participated and that the enriched pathways were poorly

overlapped (Fig. S5). These findings strongly suggested no universal rule for

evolutionary changes of the reduced genome.

In comparison, 1,226 DEGs were induced by genome reduction. The common DEGs

of genome reduction and evolution varied from 168 to 540, fewer than half of the DEGs

responsible for genome reduction in all Evos (Fig. 5A). The conclusion remained

consistent even if the DEGs were determined with an alternative method, RankProd

(Fig. S6). Functional enrichment of the DEGs observed the specific transcriptional

regulation and metabolic pathways participating in the transcriptome reorganization in response to genome reduction and evolution. Only σ38 was enriched in the genome reduction-mediated DEGs, which was the only regulon that partially overlapped with the Evos (Fig. 5B). No regulon in common was enriched, besides a few partially overlapped regulons, i.e., GadW, GadX, and RcsB (Fig. 5B). In addition, both the number of enriched pathways and their overlaps were significantly differentiated among the nine Evos or between Evos and reduced genome (Fig. 5C). No common pathways were commonly enriched between the genome reduction-mediated DEGs and Evos, no matter annotated the metabolic pathways with GO or KEGG (Fig. S7). The flagellar assembly, the only enriched pathway in genome reduction-mediated DEGs, was absent in all Evos (Fig. 5D). Alternatively, the amino acids-related metabolisms were frequently detected in the Evos, e.g., histidine metabolism, biosynthesis of amino acids, etc. (Fig. 5D, Fig. S7). The variable pathways in the Evos indicated that evolution compensated for the genome reduction in various ways, which differed from how the genome reduction was caused.

Gene modules responsible for the evolutionary changes of the reduced genome

Genome mutation analysis and transcriptome analysis failed to identify the common gene categories or pathways that correlated to the evolution of the reduced genome; thus, the gene modules correlated to evolution were newly evaluated. The weighted gene co-expression network analysis (WGCNA) [41] was performed toward the evolved transcriptomes as tested previously [27]. Reconstruction of the 3,290 genes in the reduced genome led to 21 gene modules, comprising 8 to 320 genes per module (Fig. S8). Hierarchical clustering of these modules showed that roughly three major classes could be primarily divided (Fig. 6A). Functional correlation analysis showed that three of 21 modules (M2, M10, and M16) were highly significantly correlated to the growth fitness, the number of DEGs, and mutation frequency, respectively (Fig. 6A). These modules M2, M10 and M16 might be considered as the hotspots for the genes responsible for growth fitness, transcriptional reorganization, and mutation accumulation of the reduced genome in evolution, respectively. It indicated that the three modules were highly essential and functionally differentiated for growth control, transcriptional change, and

mutagenesis.

Enrichment analyses further identified the gene categories, regulons, and

metabolisms that significantly appeared in the three modules (Fig. 6B). Two gene

categories of nonessential function were enriched in M10, and the module correlated to

the number of DEGs; in contrast, no gene category was enriched in M2 and M16 (Fig.

6B, left). All three modules successfully enriched the regulons without overlaps (Fig.

6B, middle). It indicates that the main regulatory mechanisms participating in the three

modules for growth control, transcriptional change, and mutagenesis were divergent.

GO enrichment resulted in various biological processes in the three modules, roughly

related to transport, transposition, and translation in M2, M10, and M16, respectively

(Fig. 6B, right). Compared to the enriched functions of the genes that disappeared due

to genome reduction (Fig. 6B, bottom), the gene categories of phage and unknown

function and the biological processes related to DNA transposition and integration were

commonly identified in M10. It strongly indicated that M10 was responsible for

genome reduction. The newly constructed gene networks successfully identified three

modules correlated to mutation, DEGs, and growth, revealing the functional

differentiation responsible for evolution to maintain the homeostatic transcriptome

architecture for a growing cell.

Discussion

The evolutionary compensation for genome reduction was directed toward an

identical goal of increased fitness but differentiated in the manner of genomic changes.

Firstly, various genetic functions seemed to trigger the increased growth rates of the

Evos. A few mutations could compensate for the sizeable genomic deficiency. In

particular, mutations in the essential genes, such as RNA polymerases (*rpoA*, *rpoB*,

rpoD) identified in three Evos (Table S3), were supposed to participate in the global

regulation for improved growth. Nevertheless, the considerable variation in the fixed

mutations without overlaps among the nine Evos (Table 1) implied no common

mutagenetic strategy for the evolutionary improvement of growth fitness. It was

supported by the fact that no genomic locational bias for mutations fixed in the

evolution (Fig. 2). Secondly, the transcriptomes presented conserved chromosomal

architectures (Fig. 3) and universal directional changes (Fig. S4), regardless of the significant and differentiated changes in gene expression in response to evolution (Fig. 4, Fig. S5). Although the periodicity of chromosomal architecture was well known [42,43] and coordinated with the bacterial growth rate[28,44], employing its homeostasis as the evolutionary consequence provided a conceptual and unique understanding of growing cells.

It's unclear whether the differentiation in evolutionary paths was particularly significant for the reduced genome used in the present study. Evolution studies often focus on finding the common mutations accumulated in multiple evolutionary lineages to obtain the reasonable mechanism responsible for the adaptation to the defined condition. Common mutations[22,45] or identical genetic functions[46] were reported in the experimental evolution with different reduced genomes, commonly known as parallel evolution (Fig. 7, i). In addition, as not all mutations contribute to the evolved fitness [22,46], another strategy for varied phenotypes was known as divergent evolution (Fig. 7, ii). The present study accentuated the variety of mutations fixed during evolution. Considering the high essentiality of the mutated genes (Table S3), most or all mutations were assumed to benefit the fitness increase, partially demonstrated previously [20]. Nevertheless, the evolved transcriptomes presented a homeostatic architecture, revealing the divergent to convergent evolutionary strategy (Fig. 7, iii). Multiple evolutionary paths for the reduced genome to improve growth fitness were likely all roads leading to Rome.

In addition, the transcriptome reorganization for fitness increase triggered by evolution differed from that for fitness decrease caused by genome reduction. General analyses failed to detect the regulatory network or genetic function mediated by genome reduction and evolution in common. Instead, the newly constructed gene modules successfully enriched the gene categories of mobile elements and unknown functions (Fig. 6B, left) as the evolutionary compensation for genome reduction. The represented mobile elements, flagella, were known to be responsive to environmental stresses such as hypoosmotic pressure or pH [47,48]. Genome reduction and evolution seemed equivalent to the stress response in *E. coli*. These findings were reasonable as enterobactin protected *E. coli* from oxidative stress [49,50], and enterobactin biosynthesis

was upregulated for biofilm formation in genome-reduced *E. coli* [51]. The compensation

of evolution to genome reduction not only verified the known function and mechanism

from a global regulatory viewpoint but also revealed a novel understanding of the

molecular mechanisms and gene functions.

The discriminated functions of gene modules might play a crucial role in response to

genomic and evolutionary changes. WGCNA was conducted to discover the potential

correlation of gene expression to growth fitness. It succeeded in finding the genes

participating in the evolutionary changes of the reduced genome to regain growth

fitness. Three enriched gene modules were assumed separately responsible for

replication, transcription, and population dynamics (Fig. 6B). The growth-correlated

gene module significantly enriched the iron-related biological functions (M2). Although

the translation was commonly reported to be correlated to the growth rate [52,53], it was

enriched in the gene module coordinated to transcriptional changes (M10). Such

functional differentiation of the gene modules might connect with the differentiated

medium components responsible for varied bacterial growth phases, which was

observed using the high-throughput growth assay in hundreds of medium combinations

combined with machine learning [54,55]. We assumed that the various chemicals disturbed

different metabolic fluxes in which different gene modules might have participated. The

biological meaningfulness of the gene modules suggested an alternative genetic

classification besides the commonly used clustering criteria, such as Gene Orthology [56]

and Regulon [57]. In summary, the present study provided a representative example

showing multiple evolutionary paths (i.e., gene mutation and expression) directed the

reduced genome to the improved fitness with the homeostatic transcriptome.

Materials and Methods

E. coli strains and culture medium

The *E. coli* K-12 W3110 wild-type and its genome-reduced strains were initially

distributed by the National BioResource Project of the National Institute of Genetics.

The reduced genome has been constructed by multiple deletions of large genomic

fragments [58], which led to an approximately 21% smaller size than its parent wild-type

genome W3110. The minimal medium M63 was used as described in detail elsewhere

[13,59]. In brief, the medium contains 62 mM dipotassium hydrogen phosphate, 39 mM

potassium dihydrogen phosphate, 15 mM ammonium sulfate, 15 μM thiamine

hydrochloride, 1.8 μM Iron (II) sulfate, 0.2 mM magnesium sulfate, and 22 mM glucose.

Experimental evolution

The genome-reduced *E. coli* strain was evolved in 2 mL of the M63 medium by serial

transfer, as previously described [17,20]. Nine evolutionary lineages were all initiated from

the identical culture stock prepared in advance. The 24-well microplates specific for

microbe culture (IWAKI) were used, and every four wells of four tenfold serial dilutions,

e.g., $10^3 \sim 10^6$, were used for each lineage. Multiple dilutions changing in order promised

at least one of the wells within the exponential growth phase after the overnight culture.

The microplates were incubated in a microplate bioshaker (Deep Well Maximizer,

Taitec) at 37°C, with rotation at 500 rpm. The serial transfer was performed at ~24-h

intervals. Only one of the four wells (dilutions) showing growth in the early exponential

phase ($OD_{600} = 0.01$-0.1) was selected and diluted into four wells of a new microplate

using four dilution ratios. Serial transfer was repeated until all evolutionary lineages

reached approximately 1,000 generations, which required approximately two months

per lineage. A total of nine lineages were conducted, and the daily records of the nine

lineages were summarized in Table S1. The evolutionary generations (G) and the

growth rates (μ) were calculated according to the following equations (Eq. 1 and Eq. 2).

$$G = \log_2 \left(\frac{N_t}{N_0} \right) \qquad \text{(Eq. 1)}$$

$$\mu = \frac{\ln\left(\frac{N_t}{N_0}\right)}{\Delta t} \qquad \text{(Eq. 2)}$$

Growth assay

Both the ancestor and the evolved *E. coli* populations of the reduced genome were

subjected to the growth assay, as previously described [13,59]. In brief, every 200 μl of

culture was dispensed to each well in the 96-well microplate (Coaster). The microplate

was incubated at 37°C in a plate reader (EPOCH2, BioTek), shaking at 567 cpm (cycles

per minute) for 48 h. The temporal changes in OD_{600} were measured at 30-min intervals.

The growth fitness (r) was calculated using the following equation between any two

consecutive points (Eq. 3).

$$r_i = \frac{ln\left(\frac{C_{i+1}}{C_i}\right)}{t_{i+1} - t_i}$$ (Eq. 3)

Here, t_i and t_{i+1} are the culture times at the two consecutive measurement points, and C_i

and C_{i+1} are the OD_{600} at time points t_i and t_{i+1}. The growth rate was determined as the

average of three consecutive r_i, showing the largest mean and minor variance to avoid

the unreliable calculation caused by the occasionally occurring values. The details of the

experimental and analytical processes can be found at https://doi.org/10.3791/56197.

The mean of the biological triplicates was defined as the growth fitness and used in the

analyses.

Genome resequencing and mutation analysis

The *E. coli* cells were collected at the stationary growth phase (i.e., $OD_{600} > 1.0$) and

subjected to genome resequencing, as described previously [20]. In brief, the bacterial

culture was stopped by adding rifampicin at 300 µg/mL. The cell pellet was collected

for genomic DNA purification using a Wizard Genomic DNA Purification Kit

(Promega) under the manufacturer's instructions. The sequencing library was prepared

using the Nextera XT DNA Sample Prep Kit (Illumina), and the paired-end sequencing

(300 bp × 2) was performed using the Illumina HiSeq platform. The raw datasets of

DNA-seq were deposited in the DDBJ Sequence Read Archive under the accession

number DRA013661. The sequencing reads were aligned to the reference genome *E.*

coli W3110 (AP009048.1, GenBank), and the mutation analysis was performed with the

Breseq pipeline [60,61]. The DNA sequencing and mapping statistics are summarized in

Table S2. The mutations fixed in the coding regions due to evolution are shown in Table

S3.

Calculation and simulation of the genomic positions of mutations

The distances from the genomic positions of the genome mutations fixed in the nine

Evos to the nearest genomic scars caused by the genome reduction were calculated as

previously described [13,39]. The mutation simulation was performed with Python in the

following steps. A total of 65 mutations were randomly generated on the reduced

genome, and the distances from the mutated genomic locations to the nearest genomic

scars caused by genome reduction were calculated. Subsequently, Welch's t-test was
performed to evaluate whether the distances calculated from the random mutations were
significantly longer or shorter than those calculated from the mutations that occurred in
Evos. The random simulation, distance calculation, and statistic test were performed
1,000 times, which resulted in 1,000 p values. Finally, the mean of p values (μ_p) was
calculated, and a 95% reliable region was applied. It was used to evaluate whether the
65 mutations in the Evos were significantly close to the genomic scars, i.e., the
locational bias.

RNA sequencing

The *E. coli* cells were collected at the exponential growth phase (i.e., $5 \times 10^7 \sim 2 \times 10^8$
cells/mL) and subjected to RNA sequencing, as described previously [27,44]. In brief, the
bacterial growth was stopped by mixing with the iced 10% phenol ethanol solution. The
cell pellet was collected to purify the total RNAs using the RNeasy Mini Kit (QIAGEN)
and RNase-Free DNase Set (QIAGEN) according to product instructions. The
paired-end sequencing (150 bp $\times$ 2) was performed using the Novaseq6000
next-generation sequencer (Illumina). The rRNAs were removed from the total RNAs
using the Ribo-Zero Plus rRNA Depletion Kit (Illumina), and the mRNA libraries were
prepared using the Ultra Directional RNA Library Prep Kit for Illumina (NEBNext).
Biological replicates were performed for all conditions (N=2~4). The raw datasets were
deposited in the DDBJ Sequence Read Archive under the accession number
DRA013662.

Data processing and normalization

The FASTQ files were mapped to the reference genome W3110 (accession number
AP009048.1, GenBank) using the mapping software Bowtie2 [62], as described
previously [27,44]. The obtained read counts were converted to FPKM values according to
the gene length and total read count values. Global normalization of the FPKM values
was performed to reach an identical mean value in the logarithmic scale in all datasets.
The resulting normalized FPKM values were statistically equivalent to TPM. The gene
expression level was determined as the logarithmic value of FPKM, and the mean

values of biological replicates were used for the following analyses (Table S6).

Computational analysis

The normalized datasets were subjected to the computational analyses performed

with the R statistical analysis software. A total of 3290 genes were used for hierarchical

clustering and principal component analysis (PCA), which were performed with the R

functions of "hclust" and "prcomp", respectively, as described previously[27]. The

corresponding parameters of "method" and "scale" were set as "average" and "F",

respectively. The R package of DESeq2[63] was used to determine the differentially

expressed genes (DEGs), based on the false discovery rate (FDR < 0.05) [64]. The read

counts were used as the input data for DESeq2, in which the data normalization was

performed at each run for pair comparison.

Enrichment analysis

Functional enrichments were performed according to the features of gene category[65],

transcriptional regulation[57], gene ontology[56], and metabolic pathways[66,67]. Twenty-one

gene categories and 46 regulons, which comprised more than 15 genes and 15

regulatees, were subjected to the enrichment, respectively. The statistical significance

was evaluated by the binomial test with Bonferroni correction. The enrichment analysis

of gene ontology (GO terms) [56,68] and metabolic pathway (Kyoto encyclopedia of genes

and genomes, KEGG) [66,67] was performed using DAVID, a web-based tool for

visualizing the characteristics of gene clusters with expression variation [69] [70]. The

statistical significance was according to FDR.

Chromosomal periodicity analysis

Fourier transform was used to evaluate the chromosomal periodicity of the

transcriptome, as previously described [28,44]. The genome was divided into compartments

of 1 kb each, and the mean expression level of the genes within the corresponding

sections was calculated. Gene expression levels were smoothed with a moving average

of 100 kb and subjected to the periodicity analysis using the function "periodogram" in

R. The max peak (periodic wavelength) of the periodogram was fitted to the gene

expression data using the function "nls" in R by the least squares method according to

the following equation (Eq. 4).

$$\text{Exp}(x) = a * \sin\left(x + \frac{b}{T}\right) + c \quad \text{(Eq. 4)}$$

Here, a, b, T, and c represent the periodic amplitude, the periodic phase, the periodic

wavelength indicated by the max peak, and the mean expression level of the whole

transcriptome as a constant, respectively. Note that altering the moving average did not

change the max peak. The statistical significance of the periodicity was assessed with

Fisher's g test (Table S4), as described previously [28,44]. The genomic position of *ori* was

determined according to the previous reports [44,71]. In addition, the function "abline" in R

was used to point out the genomic positions of the mutations.

Gene network analysis

The weighted gene co-expression network analysis (WGCNA) of the nine Evos was

performed with the R package of WGCNA [41], as described previously [27]. A

Step-by-Step method was used to determine the parameters for constructing the gene

networks. The soft threshold was set at 12, where the R^2 of Scale Free Topology Model

Fit was approximately 0.9, as recommended by the developer's instruction. The

resultant gene networks were clustered with the "hclust" function (method=average) and

reconstructed by merging similar modules using the "mergeCloseModules" function

with a height cut of 0.25 in the "dynamic tree cut" method. Finally, a total of 21

modules were determined for 3,290 genes. The correlation coefficients and p-values

between the expression of the gene modules and the other global features (growth rates,

DEGs, and mutations) were evaluated using the functions "cor" and "corPvalueStudent"

in R, respectively. FDR correction was applied to the p-values to account for

multiplicity. Functional enrichment of the gene modules was performed, and the

statistical significance was evaluated by the binomial test with Bonferroni correction as

described above.

Acknowledgments

We thank NBRP for providing the *E. coli* strains carrying the wild-type and reduced

genomes (W3110 and KHK collection). This work was supported by the JSPS

KAKENHI Grant-in-Aid for Scientific Research (B) (grant number 19H03215) and partially by Grant-in-Aid for Challenging Exploratory Research (grant number 21K19815).

Competing interests

The authors declare that there are no competing interests.

References

Koonin, E. V. Evolution of genome architecture. *Int J Biochem Cell Biol* **41**, 298-306 (2009). https://doi.org/S1357-2725(08)00390-7

Lynch, M. Streamlining and simplification of microbial genome architecture. *Annu Rev Microbiol* **60**, 327-349 (2006). https://doi.org/10.1146/annurev.micro.60.080805.142300

Lynch, M. & Conery, J. S. The origins of genome complexity. *Science* **302**, 1401-1404 (2003). https://doi.org/10.1126/science.1089370

Kotaka, Y., Hashimoto, M., Lee, K. I. & Kato, J. I. Mutations identified in engineered Escherichia coli with a reduced genome. *Front Microbiol* **14**, 1189877 (2023). https://doi.org/10.3389/fmicb.2023.1189877

Kato, J. & Hashimoto, M. Construction of consecutive deletions of the Escherichia coli chromosome. *Mol Syst Biol* **3**, 132 (2007). https://doi.org/10.1038/msb4100174

Pósfai, G. *et al.* Emergent properties of reduced-genome Escherichia coli. *Science* **312**, 1044-1046 (2006). https://doi.org/10.1126/science.1126439

Hashimoto, M. *et al.* Cell size and nucleoid organization of engineered Escherichia coli cells with a reduced genome. *Mol Microbiol* **55**, 137-149 (2005). https://doi.org/10.1111/j.1365-2958.2004.04386.x

Aida, H. & Ying, B.-W. Efforts to Minimise the Bacterial Genome as a Free-Living Growing System. *Biology* **12**, 1170 (2023). https://doi.org/10.3390/biology12091170

Breuer, M. *et al.* Essential metabolism for a minimal cell. *Elife* **8** (2019). https://doi.org/10.7554/eLife.36842

Hutchison, C. A., 3rd *et al.* Design and synthesis of a minimal bacterial genome. *Science* **351**, aad6253 (2016). https://doi.org/10.1126/science.aad6253

Hitomi, K., Weng, J. & Ying, B. W. Contribution of the genomic and nutritional differentiation to the spatial distribution of bacterial colonies. *Front Microbiol*

13, 948657 (2022). https://doi.org/10.3389/fmicb.2022.948657

12   Xue, H., Kurokawa, M. & Ying, B. W. Correlation between the spatial

distribution and colony size was common for monogenetic bacteria in laboratory

conditions. *BMC Microbiol* **21**, 114 (2021).

https://doi.org/10.1186/s12866-021-02180-8

13   Kurokawa, M., Seno, S., Matsuda, H. & Ying, B. W. Correlation between

genome reduction and bacterial growth. *DNA Res* **23**, 517-525 (2016).

https://doi.org/10.1093/dnares/dsw035

14   Karcagi, I. *et al.* Indispensability of Horizontally Transferred Genes and Its

Impact on Bacterial Genome Streamlining. *Mol Biol Evol* **33**, 1257-1269 (2016).

https://doi.org/10.1093/molbev/msw009

15   Gibson, D. G. *et al.* Creation of a bacterial cell controlled by a chemically

synthesized genome. *Science* **329**, 52-56 (2010).

https://doi.org/10.1126/science.1190719

16   Orr, H. A. Fitness and its role in evolutionary genetics. *Nat Rev Genet* **10**,

531-539 (2009). https://doi.org/10.1038/nrg2603

17   Nishimura, I., Kurokawa, M., Liu, L. & Ying, B. W. Coordinated Changes in

Mutation and Growth Rates Induced by Genome Reduction. *mBio* **8** (2017).

https://doi.org/10.1128/mBio.00676-17

18   Csorgo, B., Feher, T., Timar, E., Blattner, F. R. & Posfai, G. Low-mutation-rate,

reduced-genome Escherichia coli: an improved host for faithful maintenance of

engineered genetic constructs. *Microbial cell factories* **11**, 11 (2012).

https://doi.org/10.1186/1475-2859-11-11

19   Umenhoffer, K. *et al.* Reduced evolvability of Escherichia coli MDS42, an

IS-less cellular chassis for molecular and synthetic biology applications.

Microbial cell factories **9**, 38 (2010). https://doi.org/10.1186/1475-2859-9-38

20   Kurokawa, M., Nishimura, I. & Ying, B. W. Experimental Evolution Expands

the Breadth of Adaptation to an Environmental Gradient Correlated With

Genome Reduction. *Front Microbiol* **13**, 826894 (2022).

https://doi.org/10.3389/fmicb.2022.826894

21   Choe, D. *et al.* Adaptive laboratory evolution of a genome-reduced Escherichia

coli. *Nature communications* **10**, 935 (2019).

https://doi.org/10.1038/s41467-019-08888-6

22   Suzuki, S., Horinouchi, T. & Furusawa, C. Prediction of antibiotic resistance by

gene expression profiles. *Nat Commun* **5**, 5792 (2014).

https://doi.org/10.1038/ncomms6792

23   Moger-Reischer, R. Z. *et al.* Evolution of a minimal cell. *Nature* **620**, 122-127
(2023). https://doi.org/10.1038/s41586-023-06288-x

24   Ishikawa, A. *et al.* Different contributions of local- and distant-regulatory
changes to transcriptome divergence between stickleback ecotypes. *Evolution* **71**,
565-581 (2017). https://doi.org/10.1111/evo.13175

25   Rozen, D. E., de Visser, J. A. & Gerrish, P. J. Fitness effects of fixed beneficial
mutations in microbial populations. *Curr Biol* **12**, 1040-1045 (2002).
https://doi.org/10.1016/s0960-9822(02)00896-5

26   Lynch, M. Evolution of the mutation rate. *Trends Genet* **26**, 345-352 (2010).
https://doi.org/10.1016/j.tig.2010.05.003

27   Matsui, Y., Nagai, M. & Ying, B. W. Growth rate-associated transcriptome
reorganization in response to genomic, environmental, and evolutionary
interruptions. *Front Microbiol* **14**, 1145673 (2023).
https://doi.org/10.3389/fmicb.2023.1145673

28   Nagai, M., Kurokawa, M. & Ying, B. W. The highly conserved chromosomal
periodicity of transcriptomes and the correlation of its amplitude with the growth
rate in Escherichia coli. *DNA Res* **27** (2020).
https://doi.org/10.1093/dnares/dsaa018

29   Ying, B. W. & Yama, K. Gene Expression Order Attributed to Genome
Reduction and the Steady Cellular State in Escherichia coli. *Front Microbiol* **9**,
2255 (2018). https://doi.org/10.3389/fmicb.2018.02255

30   Engen, S. & Saether, B. E. r- and K-selection in fluctuating populations is
determined by the evolutionary trade-off between two fitness measures: Growth
rate and lifetime reproductive success. *Evolution* **71**, 167-173 (2017).
https://doi.org/10.1111/evo.13104

31   Luckinbill, L. S. r and K Selection in Experimental Populations of Escherichia
coli. *Science* **202**, 1201-1203 (1978).
https://doi.org/10.1126/science.202.4373.1201

32   Marshall, D. J., Cameron, H. E. & Loreau, M. Relationships between intrinsic
population growth rate, carrying capacity and metabolism in microbial
populations. *The ISME Journal* **17**, 2140-2143 (2023).
https://doi.org/10.1038/s41396-023-01543-5

33   Molenaar, D., van Berlo, R., de Ridder, D. & Teusink, B. Shifts in growth
strategies reflect tradeoffs in cellular economics. *Mol Syst Biol* **5**, 323 (2009).
https://doi.org/10.1038/msb.2009.82

34   Wortel, M. T., Noor, E., Ferris, M., Bruggeman, F. J. & Liebermeister, W.

Metabolic enzyme cost explains variable trade-offs between microbial growth

rate and yield. *PLoS Comput Biol* **14**, e1006010 (2018).

https://doi.org/10.1371/journal.pcbi.1006010

35   Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene

knockout mutants: the Keio collection. *Mol Syst Biol* **2**, 2006 0008 (2006).

https://doi.org/10.1038/msb4100050

36   Jordan, I. K., Rogozin, I. B., Wolf, Y. I. & Koonin, E. V. Essential genes are

more evolutionarily conserved than are nonessential genes in bacteria. *Genome*

research **12**, 962-968 (2002). https://doi.org/10.1101/gr.87702

37   Zhang, J. Important genomic regions mutate less often than do other regions.

Nature **602**, 38-39 (2022). https://doi.org/10.1038/d41586-022-00017-6

38   Van Cleve, J. & Weissman, D. B. Measuring ruggedness in fitness landscapes.

Proceedings of the National Academy of Sciences **112**, 7345-7346 (2015).

https://doi.org/10.1073/pnas.1507916112

39   Ying, B. W., Seno, S., Kaneko, F., Matsuda, H. & Yomo, T. Multilevel

comparative analysis of the contributions of genome reduction and heat shock to

the Escherichia coli transcriptome. *BMC Genomics* **14**, 25 (2013).

https://doi.org/10.1186/1471-2164-14-25

40   Ying, B. W., Seno, S., Matsuda, H. & Yomo, T. A simple comparison of the

extrinsic noise in gene expression between native and foreign regulations in

Escherichia coli. *Biochem Biophys Res Commun* **486**, 852-857 (2017).

https://doi.org/10.1016/j.bbrc.2017.03.148

41   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation

network analysis. *BMC Bioinformatics* **9**, 559 (2008).

https://doi.org/10.1186/1471-2105-9-559

42   Mathelier, A. & Carbone, A. Chromosomal periodicity and positional networks

of genes in Escherichia coli. *Mol Syst Biol* **6**, 366 (2010).

https://doi.org/msb201021

43   Krogh, T. J., Moller-Jensen, J. & Kaleta, C. Impact of Chromosomal

Architecture on the Function and Evolution of Bacterial Genomes. *Front*

Microbiol **9**, 2019 (2018). https://doi.org/10.3389/fmicb.2018.02019

44   Liu, L., Kurokawa, M., Nagai, M., Seno, S. & Ying, B. W. Correlated

chromosomal periodicities according to the growth rate and gene expression. *Sci*

Rep **10**, 15531 (2020). https://doi.org/10.1038/s41598-020-72389-6

45   Choe, D. *et al.* Adaptive laboratory evolution of a genome-reduced Escherichia

coli. *Nature Communications* **10** (2019).

https://doi.org/10.1038/s41467-019-08888-6

46   Lu, H. *et al.* Primordial mimicry induces morphological change in Escherichia

coli. *Commun Biol* **5**, 24 (2022). https://doi.org/10.1038/s42003-021-02954-w

47   Ikeda, T. *et al.* Hypoosmotic stress induces flagellar biosynthesis and swimming

motility in Escherichia albertii. *Commun Biol* **3**, 87 (2020).

https://doi.org/10.1038/s42003-020-0816-5

48   Maurer, L. M., Yohannes, E., Bondurant, S. S., Radmacher, M. & Slonczewski, J.

L. pH regulates genes for flagellar motility, catabolism, and oxidative stress in

Escherichia coli K-12. *J Bacteriol* **187**, 304-319 (2005).

https://doi.org/10.1128/jb.187.1.304-319.2005

49   Adler, C. *et al.* The alternative role of enterobactin as an oxidative stress

protector allows Escherichia coli colony development. *PLoS One* **9**, e84734

(2014). https://doi.org/10.1371/journal.pone.0084734

50   Peralta, D. R. *et al.* Enterobactin as Part of the Oxidative Stress Response

Repertoire. *PLoS One* **11**, e0157799 (2016).

https://doi.org/10.1371/journal.pone.0157799

51   May, T. & Okabe, S. Enterobactin is required for biofilm development in

reduced-genome Escherichia coli. *Environ Microbiol* **13**, 3149-3162 (2011).

https://doi.org/10.1111/j.1462-2920.2011.02607.x

52   Scott, M. & Hwa, T. Shaping bacterial gene expression by physiological and

proteome allocation constraints. *Nat Rev Microbiol* (2022).

https://doi.org/10.1038/s41579-022-00818-6

53   Dai, X. *et al.* Reduction of translating ribosomes enables Escherichia coli to

maintain elongation rates during slow growth. *Nat Microbiol* **2**, 16231 (2016).

https://doi.org/10.1038/nmicrobiol.2016.231

54   Aida, H., Hashizume, T., Ashino, K. & Ying, B. W. Machine learning-assisted

discovery of growth decision elements by relating bacterial population dynamics

to environmental diversity. *Elife* **11** (2022). https://doi.org/10.7554/eLife.76846

55   Ashino, K., Sugano, K., Amagasa, T. & Ying, B. W. Predicting the decision

making chemicals used for bacterial growth. *Sci Rep* **9**, 7251 (2019).

https://doi.org/10.1038/s41598-019-43587-8

56   Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nature*

Genetics **25**, 25-29 (2000). https://doi.org/10.1038/75556

57   Salgado, H. *et al.* RegulonDB v8.0: omics data sets, evolutionary conservation,

regulatory phrases, cross-validated gold standards and more. *Nucleic Acids Res*

41, D203-213 (2013). https://doi.org/10.1093/nar/gks1201

58　Mizoguchi, H., Sawano, Y., Kato, J. & Mori, H. Superpositioning of deletions
promotes growth of Escherichia coli with a reduced genome. *DNA Res* **15**,
277-284 (2008). https://doi.org/10.1093/dnares/dsn019

59　Kurokawa, M. & Ying, B. W. Precise, High-throughput Analysis of Bacterial
Growth. *J Vis Exp* (2017). https://doi.org/10.3791/56197

60　Deatherage, D. E., Traverse, C. C., Wolf, L. N. & Barrick, J. E. Detecting rare
structural variation in evolving microbial populations from new sequence
junctions using breseq. *Front Genet* **5**, 468 (2014).
https://doi.org/10.3389/fgene.2014.00468

61　Barrick, J. E. *et al.* Identifying structural variation in haploid microbial genomes
from short-read resequencing data using breseq. *BMC Genomics* **15**, 1039
(2014). https://doi.org/10.1186/1471-2164-15-1039

62　Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat
Methods* **9**, 357-359 (2012). https://doi.org/10.1038/nmeth.1923

63　Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).
https://doi.org/10.1186/s13059-014-0550-8

64　Storey, J. D. A direct approach to false discovery rates. *Journal of the Royal
Statistical Society: Series B (Statistical Methodology)* **64**, 479-498 (2002).
https://doi.org/https://doi.org/10.1111/1467-9868.00346

65　Riley, M. *et al.* Escherichia coli K-12: a cooperatively developed annotation
snapshot--2005. *Nucleic Acids Res* **34**, 1-9 (2006).
https://doi.org/10.1093/nar/gkj405

66　Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M. & Tanabe, M. KEGG as
a reference resource for gene and protein annotation. *Nucleic Acids Res* **44**,
D457-462 (2016). https://doi.org/10.1093/nar/gkv1070

67　Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes.
Nucleic Acids Res **28**, 27-30 (2000).

68　Carbon, S. *et al.* AmiGO: online access to ontology and annotation data.
Bioinformatics **25**, 288-289 (2009).
https://doi.org/10.1093/bioinformatics/btn615

69　Huang, D. W. *et al.* DAVID Bioinformatics Resources: expanded annotation
database and novel algorithms to better extract biology from large gene lists.
Nucleic Acids Res **35**, W169-175 (2007). https://doi.org/10.1093/nar/gkm415

70　Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative
analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**,

44-57 (2009). https://doi.org/10.1038/nprot.2008.211

71   Bryant, J. A., Sellars, L. E., Busby, S. J. & Lee, D. J. Chromosome position

effects on gene expression in Escherichia coli K-12. *Nucleic Acids Res* **42**,

11383-11392 (2014). https://doi.org/10.1093/nar/gku828

Table 1 Overview of fixed genome mutations. The number of mutations in the nine
Evos are shown separately and summed**.** SNP, N and S indicate single nucleotide
substitution, nonsynonymous and synonymous SNP, respectively.
718

Evos	All	Intergenic	Genic		Genic SNP	
			Indel	SNP	N	S
A1	3	0	0	2	1	1
A2	2	1	0	1	1	0
B2	13	0	1	11	9	2
C2	11	1	0	9	8	1
D2	7	0	1	5	4	1
E2	3	1	0	1	1	0
F2	8	1	0	6	5	1
G2	12	2	2	6	4	2
H2	6	0	2	4	3	1
Sum	65	6	6	45	36	9

 Figure legends

 Figure 1 Fitness increase of the reduced genome mediated by experimental

 evolution. A. Temporal changes in growth rate. Color variation indicates the nine

 evolutionary lineages. **B.** Growth rate and maximal population size of the reduced

 genome. Blue and pink indicate the common ancestor and the nine evolved populations,

 respectively. Standard errors are shown according to the biological replications (N=4~6).

 C. Boxplots of growth rate and maximum. Cross and open circles indicate the mean and

 individual values, respectively. Statistic significance evaluated by Mann-Whitney U

 tests is indicated. **D.** Correlation between growth rate and maximum. Spearman's rank

 correlation coefficient and p-value are indicated.

 Figure 2 Genomic localization of mutations. A. Normalized genomic positions of all

 mutations. The vertical lines highlight the total 65 mutations fixed in the nine Evos.

 Color variation indicates the nine Evos. WT and Reduced represent the wild-type and

 reduced genomes used in the present study. **B.** Normalized genomic positions of random

 mutations. The simulation of 65 mutations randomly fixed in the reduced genome was

 performed 1,000 times. As an example of the simulation, the genomic positions of 65

 random mutations are shown. The vertical lines in purple indicate the mutations. **C.**

 Statistic significance of the genome locational bias of mutations. The distance from the

 mutated location to the nearest genomic scar caused by genome reduction was

 calculated. The mutations accumulated in the nine Evos and the 1,000-time random

 simulation were all subjected to the calculation. The significance of genome locational

 bias of the mutations in Evos was evaluated by Welch's t-test. The histogram of 1,000

 tests for 1,000 simulated results is shown. The mean of *p*-values (μ_p) is indicated, which

 is within the 95% confidence interval ($0.07 < \mu_p < 0.09$).

 Figure 3 Chromosomal periodicity of transcriptome and mutated gene expression.

 A. Chromosomal periodicity of transcriptomes. The transcriptomes of the nine Evos are

 shown. Black lines, red curves, and red vertical lines indicate the gene expression levels,

 fitted periods, and locations of mutations, respectively. *Ori* and *dif* are indicated with

 the vertical broken lines. **B.** Boxplot of gene expression levels. Gene expression levels

of the 49 mutated genes in the nine Evos and the remaining 3,225 genes are shown.

Statistic significance evaluated by Welch's t-test is indicated.

Figure 4 Differentially expressed genes (DEGs) and their enriched functions. A.

Commonality of DEGs in the nine Evos. Closed circles represent the combinations of

the Evos. Vertical and horizontal bars indicate the number of the overlapped DEGs in

the combinations and the number of all DEGs in each Evos, respectively. The

combinations with more than 20 DEGs in common are shown. **B.** The number of DEGs

overlapped among the Evos. The numbers of DEGs overlapped across 2 to 9 Evos are

shown. The number of Evos detected in the single Evos is indicated as 1. **C.** Enriched

function in common. The KEGG and GO terms enriched in the common DEGs across

the nine Evos are shown. The statistical significance (FDR) of the enriched pathways

and biological processes is shown on a logarithmic scale represented by color gradation.

Figure 5 Transcriptome comparison between genome reduction and evolution. A.

Venn diagrams of DEGs induced by genome reduction and evolution. The numbers of

individual and overlapped DEGs are indicated. **B.** Heatmap of enriched regulons.

Statistically significant regulons are shown with the FDR values on a logarithmic scale.

C. Number of enriched functions in common. Left and right panels indicate the numbers

of enriched GO terms and KEGG pathways caused by genome reduction and evolution,

respectively. **D.** Enriched functions in common. The overlapped GO terms enriched in

the nine Evos and genome reduction are shown. Blue and pink represent genome

reduction and evolution, respectively.

Figure 6 Reconstructed gene modules. A. Cluster dendrogram of the gene modules

reconstructed by WGCNA. A total of 21 gene modules (M1~M21) were reconstructed.

The significance of the correlation coefficients of the gene modules to growth, mutation,

and expression is represented in purple gradation. From light to dark indicates the

logarithmic p-values from high to low. **B.** Enriched functions of gene modules and

deletion. Enriched gene categories, regulons, and GO terms are shown from left to right.

The numbers of the genes assigned in the three gene modules and the genomic deletion

for genome reduction are indicated in the brackets. Color gradation indicates the

normalized *p* values on a logarithmic scale.

Figure 7 Schematic drawing of evolutionary approaches for the reduced genome.

Three evolutionary strategies are proposed. Pink and blue arrowed lines indicate

experimental evolution and genome reduction, respectively. The size of the open cycles

represents the genome size. Black and grey indicate the ancestor and evolved genomes,

respectively.

Figure 1

A
Reduced

WT

Normalized genomic position

A1
A2
B2
C2
D2
E2
F2
G2
H2

B
Reduced

WT

Normalized genomic position

C

$\mu_p = 0.09$

Frequency

$\log_{10}$ (*p*-value)

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Genome reduction

Experimental evolution

i

Parallel evolution

ii

Divergent evolution

iii

Divergent to convergent evolution

Figure 7